



MATRIX MATHEMATICS Second Edition



DENNIS S. BERNSTEIN

Matrix Mathematics



Matrix Mathematics

Theory, Facts, and Formulas

Dennis S. Bernstein

PRINCETON UNIVERSITY PRESS
PRINCETON AND OXFORD

Copyright ©2009 by Princeton University Press

Published by Princeton University Press,
41 William Street, Princeton, New Jersey 08540

In the United Kingdom: Princeton University Press,
6 Oxford Street, Woodstock, Oxfordshire, OX20 1TW

All Rights Reserved

Library of Congress Cataloging-in-Publication Data

Bernstein, Dennis S., 1954–

Matrix mathematics: theory, facts, and formulas / Dennis S. Bernstein. – 2nd ed.
p. cm.

Includes bibliographical references and index.

ISBN 978-0-691-13287-7 (hardcover : alk. paper)

ISBN 978-0-691-14039-1 (pbk. : alk. paper)

1. Matrices. 2. Linear systems. I. Title.

QA188.B475 2008

512.9'434—dc22

2008036257

British Library Cataloging-in-Publication Data is available

This book has been composed in Computer Modern and Helvetica.

The publisher would like to acknowledge the author of this volume for providing the camera-ready copy from which this book was printed.

Printed on acid-free paper. ∞

www.press.princeton.edu

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

To the memory of my parents

... vessels, unable to contain the great light flowing into them, shatter and break. ... the remains of the broken vessels fall ... into the lowest world, where they remain scattered and hidden

— D. W. Menzi and Z. Padeh,
The Tree of Life, Chayyim Vital's
Introduction to the Kabbalah of
Isaac Luria, Jason Aaronson,
Northvale, 1999

Thor ... placed the horn to his lips ... He drank with all his might and kept drinking as long as ever he was able; when he paused to look, he could see that the level had sunk a little, ... for the other end lay out in the ocean itself.

— P. A. Munch, *Norse Mythology*,
AMS Press, New York, 1970

Contents

<i>Preface to the Second Edition</i>	xv
<i>Preface to the First Edition</i>	xvii
<i>Special Symbols</i>	xxi
<i>Conventions, Notation, and Terminology</i>	xxxiii
1. Preliminaries	1
1.1 Logic and Sets	1
1.2 Functions	3
1.3 Relations	5
1.4 Graphs	8
1.5 Facts on Logic, Sets, Functions, and Relations	10
1.6 Facts on Graphs	13
1.7 Facts on Binomial Identities and Sums	14
1.8 Facts on Convex Functions	21
1.9 Facts on Scalar Identities and Inequalities in One Variable	22
1.10 Facts on Scalar Identities and Inequalities in Two Variables	30
1.11 Facts on Scalar Identities and Inequalities in Three Variables	39
1.12 Facts on Scalar Identities and Inequalities in Four Variables	46
1.13 Facts on Scalar Identities and Inequalities in Six Variables	47
1.14 Facts on Scalar Identities and Inequalities in Eight Variables	47
1.15 Facts on Scalar Identities and Inequalities in n Variables	48
1.16 Facts on Scalar Identities and Inequalities in $2n$ Variables	60
1.17 Facts on Scalar Identities and Inequalities in $3n$ Variables	67
1.18 Facts on Scalar Identities and Inequalities in Complex Variables	68
1.19 Facts on Trigonometric and Hyperbolic Identities	74
1.20 Notes	76
2. Basic Matrix Properties	77
2.1 Matrix Algebra	77
2.2 Transpose and Inner Product	84
2.3 Convex Sets, Cones, and Subspaces	89
2.4 Range and Null Space	93
2.5 Rank and Defect	95
2.6 Invertibility	98

2.7	The Determinant	102
2.8	Partitioned Matrices	106
2.9	Facts on Polars, Cones, Dual Cones, Convex Hulls, and Subspaces	110
2.10	Facts on Range, Null Space, Rank, and Defect	115
2.11	Facts on the Range, Rank, Null Space, and Defect of Partitioned Matrices	120
2.12	Facts on the Inner Product, Outer Product, Trace, and Matrix Powers	126
2.13	Facts on the Determinant	128
2.14	Facts on the Determinant of Partitioned Matrices	132
2.15	Facts on Left and Right Inverses	140
2.16	Facts on the Adjugate and Inverses	141
2.17	Facts on the Inverse of Partitioned Matrices	146
2.18	Facts on Commutators	149
2.19	Facts on Complex Matrices	151
2.20	Facts on Geometry	154
2.21	Facts on Majorization	162
2.22	Notes	164
3.	Matrix Classes and Transformations	165
3.1	Matrix Classes	165
3.2	Matrices Based on Graphs	170
3.3	Lie Algebras and Groups	171
3.4	Matrix Transformations	173
3.5	Projectors, Idempotent Matrices, and Subspaces	175
3.6	Facts on Group-Invertible and Range-Hermitian Matrices	177
3.7	Facts on Normal, Hermitian, and Skew-Hermitian Matrices	178
3.8	Facts on Commutators	184
3.9	Facts on Linear Interpolation	185
3.10	Facts on the Cross Product	186
3.11	Facts on Unitary and Shifted-Unitary Matrices	189
3.12	Facts on Idempotent Matrices	198
3.13	Facts on Projectors	206
3.14	Facts on Reflectors	211
3.15	Facts on Involutory Matrices	212
3.16	Facts on Tripotent Matrices	212
3.17	Facts on Nilpotent Matrices	213
3.18	Facts on Hankel and Toeplitz Matrices	215
3.19	Facts on Hamiltonian and Symplectic Matrices	216
3.20	Facts on Miscellaneous Types of Matrices	217
3.21	Facts on Groups	221
3.22	Facts on Quaternions	225
3.23	Notes	229
4.	Polynomial Matrices and Rational Transfer Functions	231
4.1	Polynomials	231
4.2	Polynomial Matrices	234
4.3	The Smith Decomposition and Similarity Invariants	236

4.4	Eigenvalues	239
4.5	Eigenvectors	245
4.6	The Minimal Polynomial	247
4.7	Rational Transfer Functions and the Smith-McMillan Decomposition	249
4.8	Facts on Polynomials and Rational Functions	253
4.9	Facts on the Characteristic and Minimal Polynomials	260
4.10	Facts on the Spectrum	265
4.11	Facts on Graphs and Nonnegative Matrices	272
4.12	Notes	281
5.	Matrix Decompositions	283
5.1	Smith Form	283
5.2	Multicompanion Form	283
5.3	Hypercompanion Form and Jordan Form	287
5.4	Schur Decomposition	292
5.5	Eigenstructure Properties	295
5.6	Singular Value Decomposition	301
5.7	Pencils and the Kronecker Canonical Form	304
5.8	Facts on the Inertia	307
5.9	Facts on Matrix Transformations for One Matrix	311
5.10	Facts on Matrix Transformations for Two or More Matrices	316
5.11	Facts on Eigenvalues and Singular Values for One Matrix	321
5.12	Facts on Eigenvalues and Singular Values for Two or More Matrices	333
5.13	Facts on Matrix Pencils	338
5.14	Facts on Matrix Eigenstructure	338
5.15	Facts on Matrix Factorizations	345
5.16	Facts on Companion, Vandermonde, and Circulant Matrices	352
5.17	Facts on Simultaneous Transformations	358
5.18	Facts on the Polar Decomposition	359
5.19	Facts on Additive Decompositions	360
5.20	Notes	361
6.	Generalized Inverses	363
6.1	Moore-Penrose Generalized Inverse	363
6.2	Drazin Generalized Inverse	367
6.3	Facts on the Moore-Penrose Generalized Inverse for One Matrix	369
6.4	Facts on the Moore-Penrose Generalized Inverse for Two or More Matrices	377
6.5	Facts on the Moore-Penrose Generalized Inverse for Partitioned Matrices	385
6.6	Facts on the Drazin and Group Generalized Inverses	393
6.7	Notes	398
7.	Kronecker and Schur Algebra	399
7.1	Kronecker Product	399
7.2	Kronecker Sum and Linear Matrix Equations	402

7.3	Schur Product	404
7.4	Facts on the Kronecker Product	405
7.5	Facts on the Kronecker Sum	409
7.6	Facts on the Schur Product	413
7.7	Notes	416
8.	Positive-Semidefinite Matrices	417
8.1	Positive-Semidefinite and Positive-Definite Orderings	417
8.2	Submatrices	419
8.3	Simultaneous Diagonalization	422
8.4	Eigenvalue Inequalities	424
8.5	Exponential, Square Root, and Logarithm of Hermitian Matrices	430
8.6	Matrix Inequalities	431
8.7	Facts on Range and Rank	443
8.8	Facts on Structured Positive-Semidefinite Matrices	444
8.9	Facts on Identities and Inequalities for One Matrix	450
8.10	Facts on Identities and Inequalities for Two or More Matrices	456
8.11	Facts on Identities and Inequalities for Partitioned Matrices	467
8.12	Facts on the Trace	475
8.13	Facts on the Determinant	485
8.14	Facts on Convex Sets and Convex Functions	494
8.15	Facts on Quadratic Forms	500
8.16	Facts on Simultaneous Diagonalization	507
8.17	Facts on Eigenvalues and Singular Values for One Matrix	508
8.18	Facts on Eigenvalues and Singular Values for Two or More Matrices	512
8.19	Facts on Alternative Partial Orderings	522
8.20	Facts on Generalized Inverses	525
8.21	Facts on the Kronecker and Schur Products	531
8.22	Notes	541
9.	Norms	543
9.1	Vector Norms	543
9.2	Matrix Norms	546
9.3	Compatible Norms	549
9.4	Induced Norms	553
9.5	Induced Lower Bound	558
9.6	Singular Value Inequalities	560
9.7	Facts on Vector Norms	563
9.8	Facts on Matrix Norms for One Matrix	571
9.9	Facts on Matrix Norms for Two or More Matrices	580
9.10	Facts on Matrix Norms for Partitioned Matrices	593
9.11	Facts on Matrix Norms and Eigenvalues Involving One Matrix	596
9.12	Facts on Matrix Norms and Eigenvalues Involving Two or More Matrices	599
9.13	Facts on Matrix Norms and Singular Values for One Matrix	602
9.14	Facts on Matrix Norms and Singular Values for Two or More Matrices	607
9.15	Facts on Least Squares	618

9.16	Notes	619
10.	Functions of Matrices and Their Derivatives	621
10.1	Open Sets and Closed Sets	621
10.2	Limits	622
10.3	Continuity	623
10.4	Derivatives	625
10.5	Functions of a Matrix	628
10.6	Matrix Square Root and Matrix Sign Functions	629
10.7	Matrix Derivatives	630
10.8	Facts Involving One Set	632
10.9	Facts Involving Two or More Sets	634
10.10	Facts on Matrix Functions	637
10.11	Facts on Functions and Derivatives	638
10.12	Notes	642
11.	The Matrix Exponential and Stability Theory	643
11.1	Definition of the Matrix Exponential	643
11.2	Structure of the Matrix Exponential	646
11.3	Explicit Expressions	651
11.4	Matrix Logarithms	654
11.5	The Logarithm Function	656
11.6	Lie Groups	658
11.7	Lyapunov Stability Theory	660
11.8	Linear Stability Theory	662
11.9	The Lyapunov Equation	666
11.10	Discrete-Time Stability Theory	669
11.11	Facts on Matrix Exponential Formulas	671
11.12	Facts on the Matrix Sine and Cosine	677
11.13	Facts on the Matrix Exponential for One Matrix	677
11.14	Facts on the Matrix Exponential for Two or More Matrices	681
11.15	Facts on the Matrix Exponential and Eigenvalues, Singular Values, and Norms for One Matrix	689
11.16	Facts on the Matrix Exponential and Eigenvalues, Singular Values, and Norms for Two or More Matrices	692
11.17	Facts on Stable Polynomials	695
11.18	Facts on Stable Matrices	698
11.19	Facts on Almost Nonnegative Matrices	706
11.20	Facts on Discrete-Time-Stable Polynomials	708
11.21	Facts on Discrete-Time-Stable Matrices	712
11.22	Facts on Lie Groups	715
11.23	Facts on Subspace Decomposition	716
11.24	Notes	722
12.	Linear Systems and Control Theory	723
12.1	State Space and Transfer Function Models	723
12.2	Laplace Transform Analysis	726
12.3	The Unobservable Subspace and Observability	727
12.4	Observable Asymptotic Stability	732

12.5	Detectability	734
12.6	The Controllable Subspace and Controllability	735
12.7	Controllable Asymptotic Stability	743
12.8	Stabilizability	747
12.9	Realization Theory	749
12.10	Zeros	757
12.11	H_2 System Norm	765
12.12	Harmonic Steady-State Response	768
12.13	System Interconnections	770
12.14	Standard Control Problem	772
12.15	Linear-Quadratic Control	775
12.16	Solutions of the Riccati Equation	778
12.17	The Stabilizing Solution of the Riccati Equation	782
12.18	The Maximal Solution of the Riccati Equation	787
12.19	Positive-Semidefinite and Positive-Definite Solutions of the Riccati Equation	789
12.20	Facts on Stability, Observability, and Controllability	790
12.21	Facts on the Lyapunov Equation and Inertia	793
12.22	Facts on Realizations and the H_2 System Norm	798
12.23	Facts on the Riccati Equation	802
12.24	Notes	805
Bibliography		807
Author Index		891
Index		903

Preface to the Second Edition

This second edition of *Matrix Mathematics* represents a major expansion of the original work. While the total number of pages is increased 46% from 752 to 1100, the increase is actually greater since this edition is typeset in a smaller font to facilitate a manageable physical size.

The second edition expands on the first edition in several ways. For example, the new version includes material on graphs (developed within the framework of relations and partially ordered sets), as well as alternative partial orderings of matrices, such as rank subtractivity, star, and generalized Löwner. This edition also includes additional material on the Kronecker canonical form and matrix pencils; realizations of finite groups; zeros of multi-input, multi-output transfer functions; identities and inequalities for real and complex numbers; bounds on the roots of polynomials; convex functions; and vector and matrix norms.

The additional material as well as works published subsequent to the first edition increased the number of cited works from 820 to 1503, an increase of 83%. To increase the utility of the bibliography, this edition uses the “back reference” feature of LATEX, which indicates where each reference is cited in the text. As in the first edition, the second edition includes an author index. The expansion of the first edition resulted in an increase in the size of the index from 108 pages to 156 pages.

The first edition included 57 problems, while the current edition has 73. These problems represent various extensions or generalizations of known results, sometimes motivated by gaps in the literature.

In this edition, I have attempted to correct all errors that appeared in the first edition. As with the first edition, readers are encouraged to contact me about errors or omissions in the current edition, which I will periodically update on my home page.

Acknowledgments

I am grateful to many individuals who graciously provided useful advice and material for this edition. Some readers alerted me to errors, while others suggested additional material. In other cases I sought out researchers to help me understand the precise nature of interesting results. At the risk of omitting those who were helpful, I am pleased to acknowledge the following: Mark Balas, Jason Bernstein, Vijay Chellaboina, Sever Dragomir, Harry Dym, Masatoshi Fujii, Rishi Graham, Wasim Haddad, Nicholas Higham, Diederich Hinrichsen, Iman Izadi, Pierre Kabamba,

Marthe Kassouf, Christopher King, Michael Margliot, Roy Mathias, Peter Mercer, Paul Otanez, Bela Palancz, Harish Palanthandalam-Madapusi, Fotios Paliogiannis, Wei Ren, Mario Santillo, Christoph Schmoeger, Wasin So, Robert Sullivan, Yongge Tian, Panagiotis Tsiotras, Götz Trenkler, Chenwei Zhang, and Fuzhen Zhang.

As with the first edition, I am especially indebted to my family, who endured three more years of my consistent absence to make this revision a reality. It is clear that any attempt to fully embrace the enormous body of mathematics known as matrix theory is a neverending task. After committing almost two decades to the project, I remain, like Thor, barely able to perceive a dent in the vast knowledge that resides in the hundreds of thousands of pages devoted to this fascinating and incredibly useful subject. Yet, it my hope, that this book will prove to be valuable to all of those who use matrices, and will inspire interest in a mathematical construction whose secrets and mysteries know no bounds.

Dennis S. Bernstein
Ann Arbor, Michigan
dsbaero@umich.edu
October 2008

Preface to the First Edition

The idea for this book began with the realization that at the heart of the solution to many problems in science, mathematics, and engineering often lies a “matrix fact,” that is, an identity, inequality, or property of matrices that is crucial to the solution of the problem. Although there are numerous excellent books on linear algebra and matrix theory, no one book contains all or even most of the vast number of matrix facts that appear throughout the scientific, mathematical, and engineering literature. This book is an attempt to organize many of these facts into a reference source for users of matrix theory in diverse applications areas.

Viewed as an extension of scalar mathematics, matrix mathematics provides the means to manipulate and analyze multidimensional quantities. Matrix mathematics thus provides powerful tools for a broad range of problems in science and engineering. For example, the matrix-based analysis of systems of ordinary differential equations accounts for interaction among all of the state variables. The discretization of partial differential equations by means of finite differences and finite elements yields linear algebraic or differential equations whose matrix structure reflects the nature of physical solutions [1238]. Multivariate probability theory and statistical analysis use matrix methods to represent probability distributions, to compute moments, and to perform linear regression for data analysis [504, 606, 654, 702, 947, 1181]. The study of linear differential equations [691, 692, 727] depends heavily on matrix analysis, while linear systems and control theory are matrix-intensive areas of engineering [3, 65, 142, 146, 311, 313, 348, 371, 373, 444, 502, 616, 743, 852, 865, 935, 1094, 1145, 1153, 1197, 1201, 1212, 1336, 1368, 1455, 1498]. In addition, matrices are widely used in rigid body dynamics [26, 726, 733, 789, 806, 850, 970, 1026, 1068, 1069, 1185, 1200, 1222, 1351], structural mechanics [863, 990, 1100], computational fluid dynamics [305, 479, 1426], circuit theory [30], queuing and stochastic systems [642, 919, 1034], econometrics [403, 948, 1119], geodesy [1241], game theory [225, 898, 1233], computer graphics [62, 498], computer vision [941], optimization [255, 374, 953], signal processing [702, 1163, 1361], classical and quantum information theory [353, 702, 1042, 1086], communications systems [778, 779], statistics [580, 654, 948, 1119, 1177], statistical mechanics [16, 159, 160, 1372], demography [297, 805], combinatorics, networks, and graph theory [165, 128, 179, 223, 235, 266, 269, 302, 303, 335, 363, 405, 428, 481, 501, 557, 602, 702, 844, 920, 931, 1143, 1387], optics [549, 659, 798], dimensional analysis [641, 1252], and number theory [841].

In all applications involving matrices, computational techniques are essential for obtaining numerical solutions. The development of efficient and reliable algorithms for matrix computations is therefore an important area of research that has been

extensively developed [95, 304, 396, 569, 681, 683, 721, 752, 1224, 1225, 1227, 1229, 1315, 1369, 1427, 1431, 1433, 1478]. To facilitate the solution of matrix problems, entire computer packages have been developed using the language of matrices. However, this book is concerned with the analytical properties of matrices rather than their computational aspects.

This book encompasses a broad range of fundamental questions in matrix theory, which, in many cases can be viewed as extensions of related questions in scalar mathematics. A few such questions follow.

What are the basic properties of matrices? How can matrices be characterized, classified, and quantified?

How can a matrix be decomposed into simpler matrices? A matrix decomposition may involve addition, multiplication, and partition. Decomposing a matrix into its fundamental components provides insight into its algebraic and geometric properties. For example, the polar decomposition states that every square matrix can be written as the product of a rotation and a dilation analogous to the polar representation of a complex number.

Given a pair of matrices having certain properties, what can be inferred about the sum, product, and concatenation of these matrices? In particular, if a matrix has a given property, to what extent does that property change or remain unchanged if the matrix is perturbed by another matrix of a certain type by means of addition, multiplication, or concatenation? For example, if a matrix is nonsingular, how large can an additive perturbation to that matrix be without the sum becoming singular?

How can properties of a matrix be determined by means of simple operations? For example, how can the location of the eigenvalues of a matrix be estimated directly in terms of the entries of the matrix?

To what extent do matrices satisfy the formal properties of the real numbers? For example, while $0 \leq a \leq b$ implies that $a^r \leq b^r$ for real numbers a, b and a positive integer r , when does $0 \leq A \leq B$ imply $A^r \leq B^r$ for positive-semidefinite matrices A and B and with the positive-semidefinite ordering?

Questions of these types have occupied matrix theorists for at least a century, with motivation from diverse applications. The existing scope and depth of knowledge are enormous. Taken together, this body of knowledge provides a powerful framework for developing and analyzing models for scientific and engineering applications.

This book is intended to be useful to at least four groups of readers. Since linear algebra is a standard course in the mathematical sciences and engineering, graduate students in these fields can use this book to expand the scope of their

linear algebra text. For instructors, many of the facts can be used as exercises to augment standard material in matrix courses. For researchers in the mathematical sciences, including statistics, physics, and engineering, this book can be used as a general reference on matrix theory. Finally, for users of matrices in the applied sciences, this book will provide access to a large body of results in matrix theory. By collecting these results in a single source, it is my hope that this book will prove to be convenient and useful for a broad range of applications. The material in this book is thus intended to complement the large number of classical and modern texts and reference works on linear algebra and matrix theory [10, 376, 503, 540, 541, 558, 586, 701, 790, 872, 939, 956, 963, 1008, 1045, 1051, 1098, 1143, 1194, 1238].

After a review of mathematical preliminaries in Chapter 1, fundamental properties of matrices are described in Chapter 2. Chapter 3 summarizes the major classes of matrices and various matrix transformations. In Chapter 4 we turn to polynomial and rational matrices whose basic properties are essential for understanding the structure of constant matrices. Chapter 5 is concerned with various decompositions of matrices including the Jordan, Schur, and singular value decompositions. Chapter 6 provides a brief treatment of generalized inverses, while Chapter 7 describes the Kronecker and Schur product operations. Chapter 8 is concerned with the properties of positive-semidefinite matrices. A detailed treatment of vector and matrix norms is given in Chapter 9, while formulas for matrix derivatives are given in Chapter 10. Next, Chapter 11 focuses on the matrix exponential and stability theory, which are central to the study of linear differential equations. In Chapter 12 we apply matrix theory to the analysis of linear systems, their state space realizations, and their transfer function representation. This chapter also includes a discussion of the matrix Riccati equation of control theory.

Each chapter provides a core of results with, in many cases, complete proofs. Sections at the end of each chapter provide a collection of Facts organized to correspond to the order of topics in the chapter. These Facts include corollaries and special cases of results presented in the chapter, as well as related results that go beyond the results of the chapter. In some cases the Facts include open problems, illuminating remarks, and hints regarding proofs. The Facts are intended to provide the reader with a useful reference collection of matrix results as well as a gateway to the matrix theory literature.

Acknowledgments

The writing of this book spanned more than a decade and a half, during which time numerous individuals contributed both directly and indirectly. I am grateful for the helpful comments of many people who contributed technical material and insightful suggestions, all of which greatly improved the presentation and content of the book. In addition, numerous individuals generously agreed to read sections or chapters of the book for clarity and accuracy. I wish to thank Jasim Ahmed, Suhail Akhtar, David Bayard, Sanjay Bhat, Tony Bloch, Peter Bullen, Steve Campbell, Agostino Capponi, Ramu Chandra, Jaganath Chandrasekhar, Nalin Chaturvedi, Vijay Chellaboina, Jie Chen, David Clements, Dan Davison, Dimitris Dimogianopoulos, Jiu Ding, D. Z. Djokovic, R. Scott Erwin, R. W. Farebrother, Danny Georgiev, Joseph Grcar, Wassim Haddad, Yoram Halevi, Jesse Hoagg, Roger Horn, David Hyland, Iman Izadi, Pierre Kabamba, Vikram Kapila,

Fuad Kittaneh, Seth Lacy, Thomas Laffey, Cedric Langbort, Alan Laub, Alexander Leonessa, Kai-Yew Lum, Pertti Makila, Roy Mathias, N. Harris McClamroch, Boris Mordukhovich, Sergei Nersesov, JinHyoungh Oh, Concetta Pilotto, Harish Palanthandalum-Madapusi, Michael Piovoso, Leiba Rodman, Phil Roe, Carsten Scherer, Wasin So, Andy Sparks, Edward Tate, Yongge Tian, Panagiotis Tsiotras, Feng Tyan, Ravi Venugopal, Jan Willems, Hong Wong, Vera Zeidan, Xingzhi Zhan, and Fuzhen Zhang for their assistance. Nevertheless, I take full responsibility for any remaining errors, and I encourage readers to alert me to any mistakes, corrections of which will be posted on the web. Solutions to the open problems are also welcome.

Portions of the manuscript were typed by Jill Straehla and Linda Smith at Harris Corporation, and by Debbie Laird, Kathy Stolaruk, and Suzanne Smith at the University of Michigan. John Rogosich of Techsetters, Inc., provided invaluable assistance with LATEX issues, and Jennifer Slater carefully copyedited the entire manuscript. I also thank JinHyoungh Oh and Joshua Kang for writing C code to refine the index.

I especially thank Vickie Kearn of Princeton University Press for her wise guidance and constant encouragement. Vickie managed to address all of my concerns and anxieties, and helped me improve the manuscript in many ways.

Finally, I extend my greatest appreciation for the (uncountably) infinite patience of my family, who endured the days, weeks, months, and years that this project consumed. The writing of this book began with toddlers and ended with a teenager and a twenty-year old. We can all be thankful it is finally finished.

Dennis S. Bernstein
Ann Arbor, Michigan
dsbaero@umich.edu
January 2005

Special Symbols

General Notation

π	3.14159 ...
e	2.71828 ...
$\triangleq$	equals by definition
$\lim_{\varepsilon \downarrow 0}$	limit from the right
$\binom{\alpha}{m}$	$\frac{\alpha(\alpha-1)\cdots(\alpha-m+1)}{m!}$
$\binom{n}{m}$	$\frac{n!}{m!(n-m)!}$
$\lfloor a \rfloor$	largest integer less than or equal to a
δ_{ij}	1 if $i = j$, 0 if $i \neq j$ (Kronecker delta)
$\log$	logarithm with base e
$\text{sign } \alpha$	1 if $\alpha > 0$, -1 if $\alpha < 0$, 0 if $\alpha = 0$

Chapter 1

$\{ \}$	set (p. 2)
$\in$	is an element of (p. 2)
$\notin$	is not an element of (p. 2)
$\emptyset$	empty set (p. 2)
$\{ \}_{\text{ms}}$	multiset (p. 2)
card	cardinality (p. 2)
$\cap$	intersection (p. 2)
$\cup$	union (p. 2)
$\mathcal{Y} \setminus \mathcal{X}$	complement of $\mathcal{X}$ relative to $\mathcal{Y}$ (p. 2)
$\mathcal{X}^{\sim}$	complement of $\mathcal{X}$ (p. 2)

$\subseteq$	is a subset of (p. 2)
$\subset$	is a proper subset of (p. 3)
$(x_1, \dots, x_n)$	tuple or n -tuple (p. 3)
$\text{Graph}(f)$	$\{(x, f(x)): x \in \mathcal{X}\}$ (p. 3)
$f: \mathcal{X} \mapsto \mathcal{Y}$	f is a function with domain $\mathcal{X}$ and codomain $\mathcal{Y}$ (p. 3)
$f \bullet g$	composition of functions f and g (p. 3)
$f^{-1}(\mathcal{S})$	inverse image of $\mathcal{S}$ (p. 4)
$\text{rev}(\mathcal{R})$	reversal of the relation $\mathcal{R}$ (p. 5)
$\mathcal{R}^\sim$	complement of the relation $\mathcal{R}$ (p. 5)
$\text{ref}(\mathcal{R})$	reflexive hull of the relation $\mathcal{R}$ (p. 5)
$\text{sym}(\mathcal{R})$	symmetric hull of the relation $\mathcal{R}$ (p. 5)
$\text{trans}(\mathcal{R})$	transitive hull of the relation $\mathcal{R}$ (p. 5)
$\text{equiv}(\mathcal{R})$	equivalence hull of the relation $\mathcal{R}$ (p. 5)
$x \stackrel{\mathcal{R}}{\equiv} y$	(x, y) is an element of the equivalence relation $\mathcal{R}$ (p. 6)
$\text{glb}(\mathcal{S})$	greatest lower bound of $\mathcal{S}$ (p. 7, Definition 1.3.9)
$\text{lub}(\mathcal{S})$	least upper bound of $\mathcal{S}$ (p. 7, Definition 1.3.9)
$\inf(\mathcal{S})$	infimum of $\mathcal{S}$ (p. 7, Definition 1.3.9)
$\sup(\mathcal{S})$	supremum of $\mathcal{S}$ (p. 7, Definition 1.3.9)
$\text{rev}(\mathcal{G})$	reversal of the graph $\mathcal{G}$ (p. 8)
$\mathcal{G}^\sim$	complement of the graph $\mathcal{G}$ (p. 8)
$\text{ref}(\mathcal{G})$	reflexive hull of the graph $\mathcal{G}$ (p. 8)
$\text{sym}(\mathcal{G})$	symmetric hull of the graph $\mathcal{G}$ (p. 8)
$\text{trans}(\mathcal{G})$	transitive hull of the graph $\mathcal{G}$ (p. 8)
$\text{equiv}(\mathcal{G})$	equivalence hull of the graph $\mathcal{G}$ (p. 8)
$\text{indeg}(x)$	indegree of the node x (p. 9)
$\text{outdeg}(x)$	outdegree of the node x (p. 9)
$\text{deg}(x)$	degree of the node x (p. 9)

Chapter 2

$\mathbb{Z}$	integers (p. 77)
$\mathbb{N}$	nonnegative integers (p. 77)

$\mathbb{P}$	positive integers (p. 77)
$\mathbb{R}$	real numbers (p. 77)
$\mathbb{C}$	complex numbers (p. 77)
$\mathbb{F}$	$\mathbb{R}$ or $\mathbb{C}$ (p. 77)
j	$\sqrt{-1}$ (p. 77)
$\bar{z}$	complex conjugate of $z \in \mathbb{C}$ (p. 77)
$\operatorname{Re} z$	real part of $z \in \mathbb{C}$ (p. 77)
$\operatorname{Im} z$	imaginary part of $z \in \mathbb{C}$ (p. 77)
$ z $	absolute value of $z \in \mathbb{C}$ (p. 77)
OLHP	open left half plane in $\mathbb{C}$ (p. 77)
CLHP	closed left half plane in $\mathbb{C}$ (p. 77)
ORHP	open right half plane in $\mathbb{C}$ (p. 77)
CRHP	closed right half plane in $\mathbb{C}$ (p. 77)
$j\mathbb{R}$	imaginary numbers (p. 77)
$\mathbb{R}^n$	$\mathbb{R}^{n \times 1}$ (real column vectors) (p. 78)
$\mathbb{C}^n$	$\mathbb{C}^{n \times 1}$ (complex column vectors) (p. 78)
$\mathbb{F}^n$	$\mathbb{R}^n$ or $\mathbb{C}^n$ (p. 78)
$x_{(i)}$	i th component of $x \in \mathbb{F}^n$ (p. 78)
$x \geq y$	$x_{(i)} \geq y_{(i)}$ for all i ($x - y$ is nonnegative) (p. 79)
$x > y$	$x_{(i)} > y_{(i)}$ for all i ($x - y$ is positive) (p. 79)
$\mathbb{R}^{n \times m}$	$n \times m$ real matrices (p. 79)
$\mathbb{C}^{n \times m}$	$n \times m$ complex matrices (p. 79)
$\mathbb{F}^{n \times m}$	$\mathbb{R}^{n \times m}$ or $\mathbb{C}^{n \times m}$ (p. 79)
$\operatorname{row}_i(A)$	i th row of A (p. 79)
$\operatorname{col}_i(A)$	i th column of A (p. 79)
$A_{(i,j)}$	(i, j) entry of A (p. 79)
$A \stackrel{i}{\leftarrow} b$	matrix obtained from $A \in \mathbb{F}^{n \times m}$ by replacing $\operatorname{col}_i(A)$ with $b \in \mathbb{F}^n$ or $\operatorname{row}_i(A)$ with $b \in \mathbb{F}^{1 \times m}$ (p. 80)
$d_{\max}(A) \triangleq d_1(A)$	largest diagonal entry of $A \in \mathbb{F}^{n \times n}$ having real diagonal entries (p. 80)
$d_i(A)$	i th largest diagonal entry of $A \in \mathbb{F}^{n \times n}$ having real diagonal entries (p. 80)

$d_{\min}(A) \triangleq d_n(A)$	smallest diagonal entry of $A \in \mathbb{F}^{n \times n}$ having real diagonal entries (p. 80)
$A_{(\mathcal{S}_1, \mathcal{S}_2)}$	submatrix of A formed by retaining the rows of A listed in $\mathcal{S}_1$ and the columns of A listed in $\mathcal{S}_2$ (p. 80)
$A_{(\mathcal{S})}$	$A_{(\mathcal{S}, \mathcal{S})}$ (p. 80)
$A \geq \geq B$	$A_{(i,j)} \geq B_{(i,j)}$ for all i, j ($A - B$ is nonnegative) (p. 81)
$A > > B$	$A_{(i,j)} > B_{(i,j)}$ for all i, j ($A - B$ is positive) (p. 81)
$[A, B]$	commutator $AB - BA$ (p. 82)
$\text{ad}_A(X)$	adjoint operator $[A, X]$ (p. 82)
$x \times y$	cross product of vectors $x, y \in \mathbb{R}^3$ (p. 82)
$K(x)$	cross-product matrix for $x \in \mathbb{R}^3$ (p. 82)
$0_{n \times m}, 0$	$n \times m$ zero matrix (p. 83)
I_n, I	$n \times n$ identity matrix (p. 83)
$e_{i,n}, e_i$	$\text{col}_i(I_n)$ (p. 84)
$\hat{I}_n, \hat{I}$	$n \times n$ reverse identity matrix (p. 84) $\begin{bmatrix} 0 & & 1 \\ & \ddots & \\ 1 & & 0 \end{bmatrix}$
$E_{i,j,n \times m}, E_{i,j}$	$e_{i,n} e_{j,m}^T$ (p. 84)
$1_{n \times m}, 1$	$n \times m$ ones matrix (p. 84)
A^T	transpose of A (p. 86)
$\text{tr } A$	trace of A (p. 86)
$\overline{C}$	complex conjugate of $C \in \mathbb{C}^{n \times m}$ (p. 87)
A^*	$\overline{A}^T$ conjugate transpose of A (p. 87)
$\text{Re } A$	real part of $A \in \mathbb{F}^{n \times m}$ (p. 87)
$\text{Im } A$	imaginary part of $A \in \mathbb{F}^{n \times m}$ (p. 87)
$\overline{\mathcal{S}}$	$\{\overline{Z} : Z \in \mathcal{S}\}$ or $\{\overline{Z} : Z \in \mathcal{S}\}_{\text{ms}}$ (p. 87)
$A^{\hat{T}}$	$\hat{I} A^T \hat{I}$ reverse transpose of A (p. 88)
$A^{\hat{*}}$	$\hat{I} A^* \hat{I}$ reverse complex conjugate transpose of A (p. 88)
$ x $	absolute value of $x \in \mathbb{F}^n$ (p. 88)
$ A $	absolute value of $A \in \mathbb{F}^{n \times n}$ (p. 88)
$\text{sign } x$	sign of $x \in \mathbb{R}^n$ (p. 89)
$\text{sign } A$	sign of $A \in \mathbb{R}^{n \times n}$ (p. 89)

$\text{co } \mathcal{S}$	convex hull of $\mathcal{S}$ (p. 89)
$\text{cone } \mathcal{S}$	conical hull of $\mathcal{S}$ (p. 89)
$\text{coco } \mathcal{S}$	convex conical hull of $\mathcal{S}$ (p. 89)
$\text{span } \mathcal{S}$	span of $\mathcal{S}$ (p. 90)
$\text{aff } \mathcal{S}$	affine hull of $\mathcal{S}$ (p. 90)
$\dim \mathcal{S}$	dimension of $\mathcal{S}$ (p. 90)
$\mathcal{S}^\perp$	orthogonal complement of $\mathcal{S}$ (p. 91)
$\text{polar } \mathcal{S}$	polar of $\mathcal{S}$ (p. 91)
$\text{dcone } \mathcal{S}$	dual cone of $\mathcal{S}$ (p. 91)
$\mathcal{R}(A)$	range of A (p. 93)
$\mathcal{N}(A)$	null space of A (p. 94)
$\text{rank } A$	rank of A (p. 95)
$\text{def } A$	defect of A (p. 96)
A^L	left inverse of A (p. 98)
A^R	right inverse of A (p. 98)
A^{-1}	inverse of A (p. 101)
A^{-T}	$(A^T)^{-1}$ (p. 102)
A^{-*}	$(A^*)^{-1}$ (p. 102)
$\det A$	determinant of A (p. 103)
$A_{[i;j]}$	submatrix $A_{(\{i\}^\sim, \{j\}^\sim)}$ of A obtained by deleting $\text{row}_i(A)$ and $\text{col}_j(A)$ (p. 105)
A^A	adjugate of A (p. 105)
$A \stackrel{\text{rs}}{\leq} B$	rank subtractivity partial ordering (p. 119, Fact 2.10.32)
$A \stackrel{*}{\leq} B$	star partial ordering (p. 120, Fact 2.10.35)

Chapter 3

N_n, N	$n \times n$ standard nilpotent matrix (p. 166)
$\text{diag}(a_1, \dots, a_n)$	$\begin{bmatrix} a_1 & & 0 \\ & \ddots & \\ 0 & & a_n \end{bmatrix}$ (p. 167)
$\text{revdiag}(a_1, \dots, a_n)$	$\begin{bmatrix} 0 & & a_1 \\ & \ddots & \\ a_n & & 0 \end{bmatrix}$ (p. 167)

$\text{diag}(A_1, \dots, A_k)$

block-diagonal matrix $\begin{bmatrix} A_1 & & 0 \\ & \ddots & \\ 0 & & A_k \end{bmatrix}$, where
 $A_i \in \mathbb{F}^{n_i \times m_i}$ (p. 167)

J_{2n}, J

$\begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$ (p. 169)

$\mathfrak{gl}_{\mathbb{F}}(n), \mathfrak{pl}_{\mathbb{C}}(n), \mathfrak{sl}_{\mathbb{F}}(n),$
 $\mathfrak{u}(n), \mathfrak{su}(n), \mathfrak{so}(n),$
 $\text{symp}_{\mathbb{F}}(2n), \text{osymp}_{\mathbb{F}}(2n),$
 $\text{aff}_{\mathbb{F}}(n), \text{se}_{\mathbb{F}}(n), \text{trans}_{\mathbb{F}}(n)$

Lie algebras (p. 171)

$\text{GL}_{\mathbb{F}}(n), \text{PL}_{\mathbb{F}}(n), \text{SL}_{\mathbb{F}}(n),$
 $\text{U}(n), \text{O}(n), \text{U}(n, m),$
 $\text{O}(n, m), \text{SU}(n), \text{SO}(n),$
 $\text{Sym}_{\mathbb{F}}(2n), \text{OSym}_{\mathbb{F}}(2n),$
 $\text{Aff}_{\mathbb{F}}(n), \text{SE}_{\mathbb{F}}(n), \text{Trans}_{\mathbb{F}}(n)$

groups (p. 172)

$A_{\perp}$

complementary idempotent matrix or
projector $I - A$ corresponding to the
idempotent matrix or projector A (p. 175)

$\text{ind } A$

index of A (p. 176)

$\mathbb{H}$

quaternions (p. 225, Fact 3.22.1)

Chapter 4

$\mathbb{F}[s]$

polynomials with coefficients in $\mathbb{F}$ (p. 231)

$\deg p$

degree of $p \in \mathbb{F}[s]$ (p. 231)

$\text{mroots}(p)$

multiset of roots of $p \in \mathbb{F}[s]$ (p. 232)

$\text{roots}(p)$

set of roots of $p \in \mathbb{F}[s]$ (p. 232)

$\text{mult}_p(\lambda)$

multiplicity of λ as a root of $p \in \mathbb{F}[s]$ (p. 232)

$\mathbb{F}^{n \times m}[s]$

$n \times m$ matrices with entries in $\mathbb{F}[s]$ ($n \times m$
polynomial matrices with coefficients in $\mathbb{F}$)
(p. 234)

$\text{rank } P$

rank of $P \in \mathbb{F}^{n \times m}[s]$ (p. 235)

$\text{Szeros}(P)$

set of Smith zeros of $P \in \mathbb{F}^{n \times m}[s]$ (p. 237)

$\text{mSzeros}(P)$

multiset of Smith zeros of $P \in \mathbb{F}^{n \times m}[s]$
(p. 237)

χ_A

characteristic polynomial of A (p. 240)

$\lambda_{\max}(A) \triangleq \lambda_1(A)$

largest eigenvalue of $A \in \mathbb{F}^{n \times n}$ having real
eigenvalues (p. 240)

$\lambda_i(A)$	i th largest eigenvalue of $A \in \mathbb{F}^{n \times n}$ having real eigenvalues (p. 240)
$\lambda_{\min}(A) \triangleq \lambda_n(A)$	smallest eigenvalue of $A \in \mathbb{F}^{n \times n}$ having real eigenvalues (p. 240)
$\text{amult}_A(\lambda)$	algebraic multiplicity of $\lambda \in \text{spec}(A)$ (p. 240)
$\text{spec}(A)$	spectrum of A (p. 240)
$\text{mspec}(A)$	multispectrum of A (p. 240)
$\text{gmult}_A(\lambda)$	geometric multiplicity of $\lambda \in \text{spec}(A)$ (p. 245)
$\text{spabs}(A)$	spectral abscissa of A (p. 245)
$\text{sprad}(A)$	spectral radius of A (p. 245)
$\nu_-(A), \nu_0(A), \nu_+(A)$	number of eigenvalues of A counting algebraic multiplicity having negative, zero, and positive real part, respectively (p. 245)
$\text{In } A$	inertia of A , that is, $[\nu_-(A) \ \nu_0(A) \ \nu_+(A)]^T$ (p. 245)
$\text{sig } A$	signature of A , that is, $\nu_+(A) - \nu_-(A)$ (p. 245)
μ_A	minimal polynomial of A (p. 247)
$\mathbb{F}(s)$	rational functions with coefficients in $\mathbb{F}$ (SISO rational transfer functions) (p. 249)
$\mathbb{F}_{\text{prop}}(s)$	proper rational functions with coefficients in $\mathbb{F}$ (SISO proper rational transfer functions) (p. 249)
$\text{reldeg } g$	relative degree of $g \in \mathbb{F}_{\text{prop}}(s)$ (p. 249)
$\mathbb{F}^{n \times m}(s)$	$n \times m$ matrices with entries in $\mathbb{F}(s)$ (MIMO rational transfer functions) (p. 249)
$\mathbb{F}_{\text{prop}}^{n \times m}(s)$	$n \times m$ matrices with entries in $\mathbb{F}_{\text{prop}}(s)$ (MIMO proper rational transfer functions) (p. 249)
$\text{reldeg } G$	relative degree of $G \in \mathbb{F}_{\text{prop}}^{n \times m}(s)$ (p. 249)
$\text{rank } G$	rank of $G \in \mathbb{F}^{n \times m}(s)$ (p. 249)
$\text{poles}(G)$	set of poles of $G \in \mathbb{F}^{n \times m}(s)$ (p. 249)
$\text{bzeros}(G)$	set of blocking zeros of $G \in \mathbb{F}^{n \times m}(s)$ (p. 249)
$\text{Mcdeg } G$	McMillan degree of $G \in \mathbb{F}^{n \times m}(s)$ (p. 251)
$\text{tzeros}(G)$	set of transmission zeros of $G \in \mathbb{F}^{n \times m}(s)$ (p. 251)
$\text{mpoles}(G)$	multiset of poles of $G \in \mathbb{F}^{n \times m}(s)$ (p. 251)
$\text{mtzeros}(G)$	multiset of transmission zeros of $G \in \mathbb{F}^{n \times m}(s)$ (p. 251)

$\text{mbzeros}(G)$	multiset of blocking zeros of $G \in \mathbb{F}^{n \times m}(s)$ (p. 251)
$B(p, q)$	Bezout matrix of $p, q \in \mathbb{F}[s]$ (p. 255, Fact 4.8.6)
$H(g)$	Hankel matrix of $g \in \mathbb{F}(s)$ (p. 257, Fact 4.8.8)

Chapter 5

$C(p)$	companion matrix for monic polynomial p (p. 283)
$\mathcal{H}_l(q)$	$l \times l$ or $2l \times 2l$ hypercompanion matrix (p. 288)
$\mathcal{J}_l(q)$	$l \times l$ or $2l \times 2l$ real Jordan matrix (p. 289)
$\text{ind}_A(\lambda)$	index of λ with respect to A (p. 295)
$\sigma_i(A)$	i th largest singular value of $A \in \mathbb{F}^{n \times m}$ (p. 301)
$\sigma_{\max}(A) \triangleq \sigma_1(A)$	largest singular value of $A \in \mathbb{F}^{n \times m}$ (p. 301)
$\sigma_{\min}(A) \triangleq \sigma_n(A)$	minimum singular value of $A \in \mathbb{F}^{n \times n}$ (p. 301)
$P_{A,B}$	pencil of (A, B) , where $A, B \in \mathbb{F}^{n \times n}$ (p. 304)
$\text{spec}(A, B)$	generalized spectrum of (A, B) , where $A, B \in \mathbb{F}^{n \times n}$ (p. 304)
$\text{mspec}(A, B)$	generalized multispectrum of (A, B) , where $A, B \in \mathbb{F}^{n \times n}$ (p. 304)
$\chi_{A,B}$	characteristic polynomial of (A, B) , where $A, B \in \mathbb{F}^{n \times n}$ (p. 305)
$V(\lambda_1, \dots, \lambda_n)$	Vandermonde matrix (p. 354, Fact 5.16.1)
$\text{circ}(a_0, \dots, a_{n-1})$	circulant matrix of $a_0, \dots, a_{n-1} \in \mathbb{F}$ (p. 355, Fact 5.16.7)

Chapter 6

A^+	(Moore-Penrose) generalized inverse of A (p. 363)
$D \mathcal{A}$	Schur complement of D with respect to $\mathcal{A}$ (p. 367)
A^D	Drazin generalized inverse of A (p. 367)
$A^\#$	group generalized inverse of A (p. 369)

Chapter 7

$\text{vec } A$	vector formed by stacking columns of A (p. 399)
$\otimes$	Kronecker product (p. 400)
$P_{n,m}$	Kronecker permutation matrix (p. 402)
$\oplus$	Kronecker sum (p. 403)
$A \circ B$	Schur product of A and B (p. 404)
$A^{\circ\alpha}$	Schur power of A , $(A^{\circ\alpha})_{(i,j)} = (A_{(i,j)})^\alpha$ (p. 404)

Chapter 8

$\mathbf{H}^n$	$n \times n$ Hermitian matrices (p. 417)
$\mathbf{N}^n$	$n \times n$ positive-semidefinite matrices (p. 417)
$\mathbf{P}^n$	$n \times n$ positive-definite matrices (p. 417)
$A \geq B$	$A - B \in \mathbf{N}^n$ (p. 417)
$A > B$	$A - B \in \mathbf{P}^n$ (p. 417)
$\langle A \rangle$	$(A^*A)^{1/2}$ (p. 431)
$A \# B$	geometric mean of A and B (p. 461, Fact 8.10.43)
$A \#_\alpha B$	generalized geometric mean of A and B (p. 464, Fact 8.10.45)
$A:B$	parallel sum of A and B (p. 528, Fact 8.20.18)
$\text{sh}(A, B)$	shorted operator (p. 530, Fact 8.20.19)

Chapter 9

$\ x\ _p$	Hölder norm $\left[\sum_{i=1}^n x_{(i)} ^p \right]^{1/p}$ (p. 544)
$\ A\ _p$	Hölder norm $\left[\sum_{i,j=1}^{n,m} A_{(i,j)} ^p \right]^{1/p}$ (p. 547)
$\ A\ _F$	Frobenius norm $\sqrt{\text{tr } A^*A}$ (p. 547)
$\ A\ _{\sigma_P}$	Schatten norm $\left[\sum_{i=1}^{\text{rank } A} \sigma_i^p(A) \right]^{1/p}$ (p. 548)
$\ A\ _{q,p}$	Hölder-induced norm (p. 554)

$\ A\ _{\text{col}}$	column norm $\ A\ _{1,1} = \max_{i \in \{1, \dots, m\}} \ \text{col}_i(A)\ _1$ (p. 556)
$\ A\ _{\text{row}}$	row norm $\ A\ _{\infty, \infty} = \max_{i \in \{1, \dots, n\}} \ \text{row}_i(A)\ _1$ (p. 556)
$\ell(A)$	induced lower bound of A (p. 558)
$\ell_{q,p}(A)$	Hölder-induced lower bound of A (p. 559)
$\ \cdot\ _{\text{D}}$	dual norm (p. 570, Fact 9.7.22)

Chapter 10

$\mathbb{B}_{\varepsilon}(x)$	open ball of radius ε centered at x (p. 621)
$\mathbb{S}_{\varepsilon}(x)$	sphere of radius ε centered at x (p. 621)
$\text{int } \mathcal{S}$	interior of $\mathcal{S}$ (p. 621)
$\text{int}_{\mathcal{S}'} \mathcal{S}$	interior of $\mathcal{S}$ relative to $\mathcal{S}'$ (p. 621)
$\text{cl } \mathcal{S}$	closure of $\mathcal{S}$ (p. 621)
$\text{cl}_{\mathcal{S}'} \mathcal{S}$	closure of $\mathcal{S}$ relative to $\mathcal{S}'$ (p. 622)
$\text{bd } \mathcal{S}$	boundary of $\mathcal{S}$ (p. 622)
$\text{bd}_{\mathcal{S}'} \mathcal{S}$	boundary of $\mathcal{S}$ relative to $\mathcal{S}'$ (p. 622)
$(x_i)_{i=1}^{\infty}$	sequence $(x_1, x_2, \dots)$ (p. 622)
$\text{vcone } \mathcal{D}$	variational cone of $\mathcal{D}$ (p. 625)
$\text{D}_+ f(x_0; \xi)$	one-sided directional derivative of f at x_0 in the direction ξ (p. 625)
$\frac{\partial f(x_0)}{\partial x_{(i)}}$	partial derivative of f with respect to $x_{(i)}$ at x_0 (p. 625)
$f'(x)$	Fréchet derivative of f at x (p. 626)
$\frac{df(x_0)}{dx_{(i)}}$	$f'(x_0)$ (p. 626)
$f^{(k)}(x)$	k th Fréchet derivative of f at x (p. 627)
$\frac{d^+ f(x_0)}{dx_{(i)}}$	right one-sided derivative (p. 627)
$\frac{d^- f(x_0)}{dx_{(i)}}$	left one-sided derivative (p. 627)
$\text{Sign}(A)$	matrix sign of $A \in \mathbb{C}^{n \times n}$ (p. 630)

Chapter 11

e^A or $\exp(A)$	matrix exponential (p. 643)

$\mathcal{L}$	Laplace transform (p. 646)
$\mathcal{S}_s(A)$	asymptotically stable subspace of A (p. 665)
$\mathcal{S}_u(A)$	unstable subspace of A (p. 665)
OD	open unit disk in $\mathbb{C}$ (p. 670)
CUD	closed unit disk in $\mathbb{C}$ (p. 670)

Chapter 12

$\mathcal{U}(A, C)$	unobservable subspace of (A, C) (p. 728)
$\mathcal{O}(A, C)$	$\begin{bmatrix} C \\ CA \\ CA^2 \\ \vdots \\ CA^{n-1} \end{bmatrix} \quad (\text{p. 728})$
$\mathcal{C}(A, B)$	controllable subspace of (A, B) (p. 737)
$\mathcal{K}(A, B)$	$\begin{bmatrix} B & AB & A^2B & \cdots & A^{n-1}B \end{bmatrix} \quad (\text{p. 737})$
$G \sim \left[\begin{array}{c c} A & B \\ \hline C & D \end{array} \right]$	state space realization of $G \in \mathbb{F}_{\text{prop}}^{l \times m}[s]$ (p. 749)
$\mathcal{H}_{i,j,k}(G)$	Markov block-Hankel matrix $\mathcal{O}_i(A, C)\mathcal{K}_j(A, B)$ (p. 754)
$\mathcal{H}(G)$	Markov block-Hankel matrix $\mathcal{O}(A, C)\mathcal{K}(A, B)$ (p. 754)
$G \stackrel{\min}{\sim} \left[\begin{array}{c c} A & B \\ \hline C & D \end{array} \right]$	state space realization of $G \in \mathbb{F}_{\text{prop}}^{l \times m}[s]$ (p. 756)
$\mathcal{H}$	Hamiltonian $\begin{bmatrix} A & \Sigma \\ R_1 & -A^T \end{bmatrix} \quad (\text{p. 780})$

Conventions, Notation, and Terminology

When a word is defined, it is italicized.

The definition of a word, phrase, or symbol should always be understood as an “if and only if” statement, although for brevity “only if” is omitted. The symbol $\triangleq$ means equal by definition, where $A \triangleq B$ means that the left-hand expression A is defined to be the right-hand expression B .

Analogous statements are written in parallel using the following style: If n is (even, odd), then $n + 1$ is (odd, even).

The variables i, j, k, l, m, n always denote integers. Hence, $k \geq 0$ denotes a nonnegative integer, $k \geq 1$ denotes a positive integer, and the limit $\lim_{k \rightarrow \infty} A^k$ is taken over positive integers.

The imaginary unit $\sqrt{-1}$ is always denoted by dotless j .

The letter s always represents a complex scalar. The letter z may or may not represent a complex scalar.

The inequalities $c \leq a \leq d$ and $c \leq b \leq d$ are written simultaneously as

$$c \leq \left\{ \begin{array}{c} a \\ b \end{array} \right\} \leq d.$$

The prefix “non” means “not” in the words nonconstant, nonempty, nonintegral, nonnegative, nonreal, nonsingular, nonsquare, nonunique, and nonzero. In some traditional usage, “non” may mean “not necessarily.”

“Increasing” and “decreasing” indicate strict change for a change in the argument. The word “strict” is superfluous, and thus is omitted. Nonincreasing means nowhere increasing, while nondecreasing means nowhere decreasing.

Multisets can have repeated elements. Hence, $\{x\}_{\text{ms}}$ and $\{x, x\}_{\text{ms}}$ are different. The listed elements α, β, γ of the conventional set $\{\alpha, \beta, \gamma\}$ need not be distinct. For example, $\{\alpha, \beta, \alpha\} = \{\alpha, \beta\}$.

The order in which the elements of the set $\{x_1, \dots, x_n\}$ and the elements of the multiset $\{x_1, \dots, x_n\}_{\text{ms}}$ are listed has no significance. The components of the n -tuple $(x_1, \dots, x_n)$ are ordered.

The notation $(x_i)_{i=1}^{\infty}$ denotes the sequence $(x_1, x_2, \dots)$. A sequence can be viewed as an infinite-tuple, where the order of components is relevant and the components need not be distinct.

The composition of functions f and g is denoted by $f \bullet g$. The traditional notation $f \circ g$ is reserved for the Schur product.

$\mathcal{S}_1 \subset \mathcal{S}_2$ means that $\mathcal{S}_1$ is a proper subset of $\mathcal{S}_2$, whereas $\mathcal{S}_1 \subseteq \mathcal{S}_2$ means that $\mathcal{S}_1$ is either a proper subset of $\mathcal{S}_2$ or is equal to $\mathcal{S}_2$. Hence, $\mathcal{S}_1 \subset \mathcal{S}_2$ is equivalent to $\mathcal{S}_1 \subseteq \mathcal{S}_2$ and $\mathcal{S}_1 \neq \mathcal{S}_2$, while $\mathcal{S}_1 \subseteq \mathcal{S}_2$ is equivalent to either $\mathcal{S}_1 \subset \mathcal{S}_2$ or $\mathcal{S}_1 = \mathcal{S}_2$.

The terminology “graph” corresponds to what is commonly called a “simple directed graph,” while the terminology “symmetric graph” corresponds to a “simple undirected graph.”

The range of $\cos^{-1}$ is $[0, \pi]$, the range of $\sin^{-1}$ is $[-\pi/2, \pi/2]$, and the range of $\tan^{-1}$ is $(-\pi/2, \pi/2)$. The *angle between two vectors* is an element of $[0, \pi]$. Therefore, the inner product of two vectors can be used to compute the angle between two vectors.

$$0! \triangleq 1.$$

$$\text{For all } \alpha \in \mathbb{C}, \binom{\alpha}{0} \triangleq 1. \text{ For all } k \in \mathbb{N}, \binom{0}{k} \triangleq 1.$$

$$0/0 = (\sin 0)/0 = (\sinh 0)/0 \triangleq 1.$$

For all square matrices A , $A^0 \triangleq I$. In particular, $0_{n \times n}^0 \triangleq I_n$. With this convention, it is possible to write

$$\sum_{i=0}^{\infty} \alpha^i = \frac{1}{1 - \alpha}$$

for all $-1 < \alpha < 1$. Of course, $\lim_{x \downarrow 0} 0^x = 0$, $\lim_{x \downarrow 0} x^0 = 1$, and $\lim_{x \downarrow 0} x^x = 1$.

Neither ∞ nor $-\infty$ is a real number. However, some operations are defined for these objects as extended real numbers, such as $\infty + \infty = \infty$, $\infty \infty = \infty$, and, for all nonzero real numbers α , $\alpha \infty = \text{sign}(\alpha)\infty$. 0∞ and $\infty - \infty$ are not defined. See [68, pp. 14, 15].

$$1/\infty \triangleq 0.$$

Let a and b be real numbers such that $a < b$. A *finite interval* is of the form (a, b) , $[a, b)$, $(a, b]$, or $[a, b]$, whereas an *infinite interval* is of the form $(-\infty, a)$, $(-\infty, a]$, (a, ∞) , $[a, \infty)$, or $(-\infty, \infty)$. An *interval* is either a finite interval or an infinite interval. An *extended infinite interval* includes either ∞ or $-\infty$. For example, $[-\infty, a)$ and $[-\infty, a]$ include $-\infty$, $(a, \infty]$ and $[a, \infty]$ include ∞ , and $[-\infty, \infty]$ includes $-\infty$ and ∞ .

The symbol $\mathbb{F}$ denotes either $\mathbb{R}$ or $\mathbb{C}$ consistently in each result. For example, in Theorem 5.6.4, the three appearances of “ $\mathbb{F}$ ” can be read as either all “ $\mathbb{C}$ ” or all “ $\mathbb{R}$.”

The imaginary numbers are denoted by $j\mathbb{R}$. Hence, 0 is both a real number and an imaginary number.

The notation $\operatorname{Re} A$ and $\operatorname{Im} A$ represents the real and imaginary parts of A , respectively. Some books use $\operatorname{Re} A$ and $\operatorname{Im} A$ to denote the Hermitian and skew-Hermitian matrices $\frac{1}{2}(A + A^*)$ and $\frac{1}{2j}(A - A^*)$.

For the scalar ordering “ $\leq$,” if $x \leq y$, then $x < y$ if and only if $x \neq y$. For the entrywise vector and matrix orderings, $x \leq y$ and $x \neq y$ do not imply that $x < y$.

Operations denoted by superscripts are applied before operations represented by preceding operators. For example, $\operatorname{tr} (A + B)^2$ means $\operatorname{tr}[(A + B)^2]$ and $\operatorname{cl} S^\sim$ means $\operatorname{cl}(S^\sim)$. This convention simplifies many formulas.

A vector in $\mathbb{F}^n$ is a column vector, which is also a matrix with one column. In mathematics, “vector” generally refers to an abstract vector not resolved in coordinates.

Sets have elements, vectors have components, and matrices have entries. This terminology has no mathematical consequence.

The notation $x_{(i)}$ represents the i th component of the vector x .

The notation $A_{(i,j)}$ represents the scalar (i, j) entry of A . $A_{i,j}$ or A_{ij} denotes a block or submatrix of A .

All matrices have nonnegative integral dimensions. If at least one of the dimensions of a matrix is zero, then the matrix is empty.

The entries of a submatrix $\hat{A}$ of a matrix A are the entries of A lying in specified rows and columns. $\hat{A}$ is a block of A if $\hat{A}$ is a submatrix of A whose entries are entries of adjacent rows and columns of A . Every matrix is both a submatrix and block of itself.

The determinant of a submatrix is a subdeterminant. Some books use “minor.” The determinant of a matrix is also a subdeterminant of the matrix.

The dimension of the null space of a matrix is its defect. Some books use “nullity.”

A block of a square matrix is diagonally located if the block is square and the diagonal entries of the block are also diagonal entries of the matrix; otherwise, the block is off-diagonally located. This terminology avoids confusion with a “diagonal block,” which is a block that is also a square, diagonal submatrix.

For the partitioned matrix $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{F}^{(n+m) \times (k+l)}$, it can be inferred that $A \in \mathbb{F}^{n \times k}$ and similarly for B, C , and D .

The Schur product of matrices A and B is denoted by $A \circ B$. Matrix multiplication is given priority over Schur multiplication, that is, $A \circ BC$ means $A \circ (BC)$.

The adjugate of $A \in \mathbb{F}^{n \times n}$ is denoted by A^A . The traditional notation is $\text{adj } A$, while the notation A^A is used in [1228]. If $A \in \mathbb{F}$ is a scalar then $A^A = 1$. In particular, $0_{1 \times 1}^A = 1$. However, for all $n \geq 2$, $0_{n \times n}^A = 0_{n \times n}$.

If $\mathbb{F} = \mathbb{R}$, then $\overline{A}$ becomes A , A^* becomes A^T , “Hermitian” becomes “symmetric,” “unitary” becomes “orthogonal,” “unitarily” becomes “orthogonally,” and “congruence” becomes “T-congruence.” A square complex matrix A is symmetric if $A^T = A$ and orthogonal if $A^T A = I$.

The diagonal entries of a matrix $A \in \mathbb{F}^{n \times n}$ all of whose diagonal entries are real are ordered as $d_{\max}(A) = d_1(A) \geq d_2(A) \geq \cdots \geq d_n(A) = d_{\min}(A)$.

Every $n \times n$ matrix has n eigenvalues. Hence, eigenvalues are counted in accordance with their algebraic multiplicity. The phrase “distinct eigenvalues” ignores algebraic multiplicity.

The eigenvalues of a matrix $A \in \mathbb{F}^{n \times n}$ all of whose eigenvalues are real are ordered as $\lambda_{\max}(A) = \lambda_1(A) \geq \lambda_2(A) \geq \cdots \geq \lambda_n(A) = \lambda_{\min}(A)$.

The inertia of a matrix is written as

$$\text{In } A \triangleq \begin{bmatrix} \nu_{-}(A) \\ \nu_0(A) \\ \nu_{+}(A) \end{bmatrix}.$$

Some books use the notation $(\nu(A), \delta(A), \pi(A))$.

For $A \in \mathbb{F}^{n \times n}$, $\text{amult}_A(\lambda)$ is the number of copies of λ in the multispectrum of A , $\text{gmult}_A(\lambda)$ is the number of Jordan blocks of A associated with λ , and $\text{ind}_A(\lambda)$ is the order of the largest Jordan block of A associated with λ . The index of A , denoted by $\text{ind } A = \text{ind}_A(0)$, is the order of the largest Jordan block of A associated with the eigenvalue 0.

The matrix $A \in \mathbb{F}^{n \times n}$ is semisimple if the order of every Jordan block of A is 1, and cyclic if A has exactly one Jordan associated with each of its eigenvalues. Defective means not semisimple, while derogatory means not cyclic.

An $n \times m$ matrix has exactly $\min\{n, m\}$ singular values, exactly $\text{rank } A$ of which are positive.

The $\min\{n, m\}$ singular values of a matrix $A \in \mathbb{F}^{n \times m}$ are ordered as $\sigma_{\max}(A) \triangleq \sigma_1(A) \geq \sigma_2(A) \geq \cdots \geq \sigma_{\min\{n, m\}}(A)$. If $n = m$, then $\sigma_{\min}(A) \triangleq \sigma_n(A)$. The notation $\sigma_{\min}(A)$ is defined only for square matrices.

Positive-semidefinite and positive-definite matrices are Hermitian.

A square matrix with entries in $\mathbb{F}$ is diagonalizable over $\mathbb{F}$ if and only if it can be transformed into a diagonal matrix whose entries are in $\mathbb{F}$ by means of a similarity transformation whose entries are in $\mathbb{F}$. Therefore, a complex matrix is diagonalizable over $\mathbb{C}$ if and only if all of its eigenvalues are semisimple, whereas a real matrix is diagonalizable over $\mathbb{R}$ if and only if all of its eigenvalues are semisimple and real. The real matrix $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ is diagonalizable over $\mathbb{C}$, although it is not diagonalizable over $\mathbb{R}$. The Hermitian matrix $\begin{bmatrix} 1 & j \\ -j & 2 \end{bmatrix}$ is diagonalizable over $\mathbb{C}$, and also has real eigenvalues.

An idempotent matrix $A \in \mathbb{F}^{n \times n}$ satisfies $A^2 = A$, while a projector is a Hermitian, idempotent matrix. Some books use “projector” for idempotent and “orthogonal projector” for projector. A reflector is a Hermitian, involutory matrix. A projector is a normal matrix each of whose eigenvalues is 1 or 0, while a reflector is a normal matrix each of whose eigenvalues is 1 or -1 .

An elementary matrix is a nonsingular matrix formed by adding an outer-product matrix to the identity matrix. An elementary reflector is a reflector exactly one of whose eigenvalues is -1 . An elementary projector is a projector exactly one of whose eigenvalues is 0. Elementary reflectors are elementary matrices. However, elementary projectors are not elementary matrices since elementary projectors are singular.

A range-Hermitian matrix is a square matrix whose range is equal to the range of its complex conjugate transpose. These matrices are also called “EP” matrices.

The polynomials 1 and $s^3 + 5s^2 - 4$ are monic. The zero polynomial is not monic.

The rank of a polynomial matrix P is the maximum rank of $P(s)$ over $\mathbb{C}$. This quantity is also called the normal rank. We denote this quantity by $\text{rank } P$ as distinct from $\text{rank } P(s)$, which denotes the rank of the matrix $P(s)$.

The rank of a rational transfer function G is the maximum rank of $G(s)$ over $\mathbb{C}$ excluding poles of the entries of G . This quantity is also called the normal rank. We denote this quantity by $\text{rank } G$ as distinct from $\text{rank } G(s)$, which denotes the rank of the matrix $G(s)$.

The symbol $\oplus$ denotes the Kronecker sum. Some books use $\oplus$ to denote the direct sum of matrices or subspaces.

The notation $|A|$ represents the matrix obtained by replacing every entry of A by its absolute value.

The notation $\langle A \rangle$ represents the matrix $(A^*A)^{1/2}$. Some books use $|A|$ to denote this matrix.

The Hölder norms for vectors and matrices are denoted by $\|\cdot\|_p$. The matrix norm induced by $\|\cdot\|_q$ on the domain and $\|\cdot\|_p$ on the codomain is denoted by $\|\cdot\|_{p,q}$.

The Schatten norms for matrices are denoted by $\|\cdot\|_{\sigma p}$, and the Frobenius norm is denoted by $\|\cdot\|_F$. Hence, $\|\cdot\|_{\sigma\infty} = \|\cdot\|_{2,2} = \sigma_{\max}(\cdot)$, $\|\cdot\|_{\sigma 2} = \|\cdot\|_F$, and $\|\cdot\|_{\sigma 1} = \text{tr } \langle \cdot \rangle$.

Let “ $\leq$ ” be a partial ordering, let X be a set, and consider the inequality

$$f(x) \leq g(x) \text{ for all } x \in X. \quad (1)$$

Inequality (1) is *sharp* if there exists $x_0 \in X$ such that $f(x_0) = g(x_0)$.

The inequality

$$f(x) \leq f(y) \text{ for all } x \leq y \quad (2)$$

is a monotonicity result.

The inequality

$$f(x) \leq p(x) \leq g(x) \text{ for all } x \in X, \quad (3)$$

where p is not identically equal to either f or g on X , is an *interpolation* or *refinement* of (1). The inequality

$$g(x) \leq \alpha f(x) \text{ for all } x \in X, \quad (4)$$

where $\alpha > 1$, is a *reversal* of (1).

Defining $h(x) \triangleq g(x) - f(x)$, it follows that (1) is equivalent to

$$h(x) \geq 0 \text{ for all } x \in X. \quad (5)$$

Now, suppose that h has a global minimizer $x_0 \in X$. Then, (5) implies that

$$0 \leq h(x_0) = \min_{x \in X} h(x) \leq h(y) \text{ for all } y \in X. \quad (6)$$

Consequently, inequalities are often expressed equivalently in terms of optimization problems, and vice versa.

Many inequalities are based on a single function that is either monotonic or convex.

Matrix Mathematics

Chapter One

Preliminaries

In this chapter we review some basic terminology and results concerning logic, sets, functions, and related concepts. This material is used throughout the book.

1.1 Logic and Sets

Let A and B be statements. The *negation* of A is the statement (not A), the *both* of A and B is the statement (A and B), and the *either* of A and B is the statement (A or B). The statement (A or B) does not contradict (A and B), that is, the word “or” is inclusive. Every *statement* is assumed to be either true or false; likewise, no statement can be both true and false.

The statements “ A and B or C ” and “ A or B and C ” are ambiguous. We therefore write “ A and either B or C ” and “either A or both B and C .”

Let A and B be statements. The *implication* statement “if A is satisfied, then B is satisfied” or, equivalently, “ A implies B ” is written as $A \implies B$, while $A \iff B$ is equivalent to $[(A \implies B) \text{ and } (A \impliedby B)]$. Of course, $A \impliedby B$ means $B \implies A$. A *tautology* is a statement that is true regardless of whether the component statements are true or false. For example, the statement “(A and B) implies A ” is a tautology. A *contradiction* is a statement that is false regardless of whether the component statements are true or false.

Suppose that $A \iff B$. Then, A is satisfied *if and only if* B is satisfied. The implication $A \implies B$ (the “only if” part) is *necessity*, while $B \implies A$ (the “if” part) is *sufficiency*. The *converse* statement of $A \implies B$ is $B \implies A$. The statement $A \implies B$ is equivalent to its *contrapositive* statement (not B) $\implies$ (not A).

A *theorem* is a significant statement, while a *proposition* is a theorem of less significance. The primary role of a *lemma* is to support the proof of a theorem or proposition. Furthermore, a *corollary* is a consequence of a theorem or proposition. Finally, a *fact* is either a theorem, proposition, lemma, or corollary. Theorems, propositions, lemmas, corollaries, and facts are provably true statements.

Suppose that $A' \implies A \implies B \implies B'$. Then, $A' \implies B'$ is a corollary of $A \implies B$.

Let A , B , and C be statements, and assume that $A \implies B$. Then, $A \implies B$ is a *strengthening* of the statement $(A \text{ and } C) \implies B$. If, in addition, $A \implies C$, then the statement $(A \text{ and } C) \implies B$ has a *redundant assumption*.

Let $\mathcal{X} \triangleq \{x, y, z\}$ be a *set*. Then,

$$x \in \mathcal{X} \quad (1.1.1)$$

means that x is an *element* of $\mathcal{X}$. If w is not an element of $\mathcal{X}$, then we write

$$w \notin \mathcal{X}. \quad (1.1.2)$$

The set with no elements, denoted by $\emptyset$, is the *empty set*. If $\mathcal{X} \neq \emptyset$, then $\mathcal{X}$ is *nonempty*.

A set cannot have repeated elements. For example, $\{x, x\} = \{x\}$. However, a *multiset* is a collection of elements that allows for repetition. The multiset consisting of two copies of x is written as $\{x, x\}_{\text{ms}}$. However, we do not assume that the listed elements x, y of the conventional set $\{x, y\}$ are distinct. The number of distinct elements of the set $\mathcal{S}$ or not-necessarily-distinct elements of the multiset $\mathcal{S}$ is the *cardinality* of $\mathcal{S}$, which is denoted by $\text{card}(\mathcal{S})$.

There are two basic types of mathematical statements involving quantifiers. An *existential statement* is of the form

$$\text{there exists } x \in \mathcal{X} \text{ such that statement } Z \text{ is satisfied,} \quad (1.1.3)$$

while a *universal statement* has the structure

$$\text{for all } x \in \mathcal{X}, \text{ it follows that statement } Z \text{ is satisfied,} \quad (1.1.4)$$

or, equivalently,

$$\text{statement } Z \text{ is satisfied for all } x \in \mathcal{X}. \quad (1.1.5)$$

Let $\mathcal{X}$ and $\mathcal{Y}$ be sets. The *intersection* of $\mathcal{X}$ and $\mathcal{Y}$ is the set of common elements of $\mathcal{X}$ and $\mathcal{Y}$ given by

$$\mathcal{X} \cap \mathcal{Y} \triangleq \{x: x \in \mathcal{X} \text{ and } x \in \mathcal{Y}\} = \{x \in \mathcal{X}: x \in \mathcal{Y}\} \quad (1.1.6)$$

$$= \{x \in \mathcal{Y}: x \in \mathcal{X}\} = \mathcal{Y} \cap \mathcal{X}, \quad (1.1.7)$$

while the set of elements in either $\mathcal{X}$ or $\mathcal{Y}$ (the *union* of $\mathcal{X}$ and $\mathcal{Y}$) is

$$\mathcal{X} \cup \mathcal{Y} \triangleq \{x: x \in \mathcal{X} \text{ or } x \in \mathcal{Y}\} = \mathcal{Y} \cup \mathcal{X}. \quad (1.1.8)$$

The *complement* of $\mathcal{X}$ relative to $\mathcal{Y}$ is

$$\mathcal{Y} \setminus \mathcal{X} \triangleq \{x \in \mathcal{Y}: x \notin \mathcal{X}\}. \quad (1.1.9)$$

If $\mathcal{Y}$ is specified, then the *complement* of $\mathcal{X}$ is

$$\mathcal{X}^{\sim} \triangleq \mathcal{Y} \setminus \mathcal{X}. \quad (1.1.10)$$

If $x \in \mathcal{X}$ implies that $x \in \mathcal{Y}$, then $\mathcal{X}$ is *contained* in $\mathcal{Y}$ ($\mathcal{X}$ is a *subset* of $\mathcal{Y}$), which is written as

$$\mathcal{X} \subseteq \mathcal{Y}. \quad (1.1.11)$$

The statement $\mathcal{X} = \mathcal{Y}$ is equivalent to the validity of both $\mathcal{X} \subseteq \mathcal{Y}$ and $\mathcal{Y} \subseteq \mathcal{X}$. If $\mathcal{X} \subseteq \mathcal{Y}$ and $\mathcal{X} \neq \mathcal{Y}$, then $\mathcal{X}$ is a *proper subset* of $\mathcal{Y}$ and we write $\mathcal{X} \subset \mathcal{Y}$. The sets $\mathcal{X}$ and $\mathcal{Y}$ are *disjoint* if $\mathcal{X} \cap \mathcal{Y} = \emptyset$. A *partition* of $\mathcal{X}$ is a set of pairwise-disjoint and nonempty subsets of $\mathcal{X}$ whose union is equal to $\mathcal{X}$.

The operations “ $\cap$,” “ $\cup$,” and “ $\setminus$ ” and the relations “ $\subset$ ” and “ $\subseteq$ ” extend directly to multisets. For example,

$$\{x, x\}_{\text{ms}} \cup \{x\}_{\text{ms}} = \{x, x, x\}_{\text{ms}}. \quad (1.1.12)$$

By ignoring repetitions, a multiset can be converted to a set, while a set can be viewed as a multiset with distinct elements.

The *Cartesian product* $\mathcal{X}_1 \times \cdots \times \mathcal{X}_n$ of sets $\mathcal{X}_1, \dots, \mathcal{X}_n$ is the set consisting of *tuples* of the form $(x_1, \dots, x_n)$, where $x_i \in \mathcal{X}_i$ for all $i = 1, \dots, n$. A tuple with n components is an *n-tuple*. Note that the components of an *n-tuple* are ordered but need not be distinct.

By replacing the logical operations “ $\implies$,” “and,” “or,” and “not” by “ $\subseteq$,” “ $\cup$,” “ $\cap$,” and “ $\sim$,” respectively, statements about statements A and B can be transformed into statements about sets $\mathcal{A}$ and $\mathcal{B}$, and vice versa. For example, the identity

$$A \text{ and } (B \text{ or } C) = (A \text{ and } B) \text{ or } (A \text{ and } C)$$

is equivalent to

$$\mathcal{A} \cap (\mathcal{B} \cup \mathcal{C}) = (\mathcal{A} \cap \mathcal{B}) \cup (\mathcal{A} \cap \mathcal{C}).$$

1.2 Functions

Let $\mathcal{X}$ and $\mathcal{Y}$ be sets. Then, a *function* f that maps $\mathcal{X}$ into $\mathcal{Y}$ is a rule $f: \mathcal{X} \mapsto \mathcal{Y}$ that assigns a unique element $f(x)$ (the *image* of x) of $\mathcal{Y}$ to each element x of $\mathcal{X}$. Equivalently, a function $f: \mathcal{X} \mapsto \mathcal{Y}$ can be viewed as a subset $\mathcal{F}$ of $\mathcal{X} \times \mathcal{Y}$ such that, for all $x \in \mathcal{X}$, it follows that there exists $y \in \mathcal{Y}$ such that $(x, y) \in \mathcal{F}$ and such that, if $(x, y_1), (x, y_2) \in \mathcal{F}$, then $y_1 = y_2$. In this case, $\mathcal{F} = \text{Graph}(f) \triangleq \{(x, f(x)): x \in \mathcal{X}\}$. The set $\mathcal{X}$ is the *domain* of f , while the set $\mathcal{Y}$ is the *codomain* of f . If $f: \mathcal{X} \mapsto \mathcal{X}$, then f is a function on $\mathcal{X}$. For $\mathcal{X}_1 \subseteq \mathcal{X}$, it is convenient to define $f(\mathcal{X}_1) \triangleq \{f(x): x \in \mathcal{X}_1\}$. The set $f(\mathcal{X})$, which is denoted by $\mathcal{R}(f)$, is the *range* of f . If, in addition, $\mathcal{Z}$ is a set and $g: \mathcal{Y} \mapsto \mathcal{Z}$, then $g \bullet f: \mathcal{X} \mapsto \mathcal{Z}$ (the *composition* of g and f) is the function $(g \bullet f)(x) \triangleq g[f(x)]$. If $x_1, x_2 \in \mathcal{X}$ and $f(x_1) = f(x_2)$ implies that $x_1 = x_2$, then f is *one-to-one*; if $\mathcal{R}(f) = \mathcal{Y}$, then f is *onto*. The function $I_{\mathcal{X}}: \mathcal{X} \mapsto \mathcal{X}$ defined by $I_{\mathcal{X}}(x) \triangleq x$ for all $x \in \mathcal{X}$ is the *identity* on $\mathcal{X}$. Finally, $x \in \mathcal{X}$ is a *fixed point* of the function $f: \mathcal{X} \mapsto \mathcal{X}$ if $f(x) = x$.

The following result shows that function composition is associative.

Proposition 1.2.1. Let $\mathcal{X}$, $\mathcal{Y}$, $\mathcal{Z}$, and $\mathcal{W}$ be sets, and let $f: \mathcal{X} \mapsto \mathcal{Y}$, $g: \mathcal{Y} \mapsto \mathcal{Z}$, $h: \mathcal{Z} \mapsto \mathcal{W}$. Then,

$$h \bullet (g \bullet f) = (h \bullet g) \bullet f. \quad (1.2.1)$$

Hence, we write $h \bullet g \bullet f$ for $h \bullet (g \bullet f)$ and $(h \bullet g) \bullet f$.

Let $\mathcal{X}$ be a set, and let $\hat{\mathcal{X}}$ be a partition of $\mathcal{X}$. Furthermore, let $f: \hat{\mathcal{X}} \mapsto \mathcal{X}$, where, for all $\mathcal{S} \in \hat{\mathcal{X}}$, it follows that $f(\mathcal{S}) \in \mathcal{S}$. Then, f is a *canonical mapping*, and $f(\mathcal{S})$ is a *canonical form*. That is, for all components $\mathcal{S}$ of the partition $\hat{\mathcal{X}}$ of $\mathcal{X}$, it follows that the function f assigns an element of $\mathcal{S}$ to the set $\mathcal{S}$.

Let $f: \mathcal{X} \mapsto \mathcal{Y}$. Then, f is *left invertible* if there exists a function $g: \mathcal{Y} \mapsto \mathcal{X}$ (a *left inverse* of f) such that $g \bullet f = I_{\mathcal{X}}$, whereas f is *right invertible* if there exists a function $h: \mathcal{Y} \mapsto \mathcal{X}$ (a *right inverse* of f) such that $f \bullet h = I_{\mathcal{Y}}$. In addition, the function $f: \mathcal{X} \mapsto \mathcal{Y}$ is *invertible* if there exists a function $f^{-1}: \mathcal{Y} \mapsto \mathcal{X}$ (the *inverse* of f) such that $f^{-1} \bullet f = I_{\mathcal{X}}$ and $f \bullet f^{-1} = I_{\mathcal{Y}}$. The *inverse image* $f^{-1}(\mathcal{S})$ of $\mathcal{S} \subseteq \mathcal{Y}$ is defined by

$$f^{-1}(\mathcal{S}) \triangleq \{x \in \mathcal{X}: f(x) \in \mathcal{S}\}. \quad (1.2.2)$$

Theorem 1.2.2. Let $\mathcal{X}$ and $\mathcal{Y}$ be sets, and let $f: \mathcal{X} \mapsto \mathcal{Y}$. Then, the following statements hold:

- i) f is left invertible if and only if f is one-to-one.
- ii) f is right invertible if and only if f is onto.

Furthermore, the following statements are equivalent:

- iii) f is invertible.
- iv) f has a unique inverse.
- v) f is one-to-one and onto.
- vi) f is left invertible and right invertible.
- vii) f has a unique left inverse.
- viii) f has a unique right inverse.

Proof. To prove i), suppose that f is left invertible with left inverse $g: \mathcal{Y} \mapsto \mathcal{X}$. Furthermore, suppose that $x_1, x_2 \in \mathcal{X}$ satisfy $f(x_1) = f(x_2)$. Then, $x_1 = g[f(x_1)] = g[f(x_2)] = x_2$, which shows that f is one-to-one. Conversely, suppose that f is one-to-one so that, for all $y \in \mathcal{R}(f)$, there exists a unique $x \in \mathcal{X}$ such that $f(x) = y$. Hence, define the function $g: \mathcal{Y} \mapsto \mathcal{X}$ by $g(y) \triangleq x$ for all $y = f(x) \in \mathcal{R}(f)$ and by $g(y)$ arbitrary for all $y \in \mathcal{Y} \setminus \mathcal{R}(f)$. Consequently, $g[f(x)] = x$ for all $x \in \mathcal{X}$, which shows that g is a left inverse of f .

To prove ii), suppose that f is right invertible with right inverse $g: \mathcal{Y} \mapsto \mathcal{X}$. Then, for all $y \in \mathcal{Y}$, it follows that $f[g(y)] = y$, which shows that f is onto. Conversely, suppose that f is onto so that, for all $y \in \mathcal{Y}$, there exists at least one $x \in \mathcal{X}$ such that $f(x) = y$. Selecting one such x arbitrarily, define $g: \mathcal{Y} \mapsto \mathcal{X}$ by $g(y) \triangleq x$. Consequently, $f[g(y)] = y$ for all $y \in \mathcal{Y}$, which shows that g is a right inverse of f . $\square$

Definition 1.2.3. Let $\mathcal{J} \subset \mathbb{R}$ be a finite or infinite interval, and let $f: \mathcal{J} \mapsto \mathbb{R}$. Then, f is *convex* if, for all $\alpha \in [0, 1]$ and for all $x, y \in \mathcal{J}$, it follows that

$$f[\alpha x + (1 - \alpha)y] \leq \alpha f(x) + (1 - \alpha)f(y). \quad (1.2.3)$$

Furthermore, f is *strictly convex* if, for all $\alpha \in (0, 1)$ and for all distinct $x, y \in \mathcal{J}$, it follows that

$$f[\alpha x + (1 - \alpha)y] < \alpha f(x) + (1 - \alpha)f(y).$$

A more general definition of convexity is given by Definition 8.6.14.

1.3 Relations

Let $\mathcal{X}$, $\mathcal{X}_1$, and $\mathcal{X}_2$ be sets. A *relation* $\mathcal{R}$ on $\mathcal{X}_1 \times \mathcal{X}_2$ is a subset of $\mathcal{X}_1 \times \mathcal{X}_2$. A *relation* $\mathcal{R}$ on $\mathcal{X}$ is a relation on $\mathcal{X} \times \mathcal{X}$. Likewise, a *multirelation* $\mathcal{R}$ on $\mathcal{X}_1 \times \mathcal{X}_2$ is a multisubset of $\mathcal{X}_1 \times \mathcal{X}_2$, while a *multirelation* $\mathcal{R}$ on $\mathcal{X}$ is a multirelation on $\mathcal{X} \times \mathcal{X}$.

Let $\mathcal{X}$ be a set, and let $\mathcal{R}_1$ and $\mathcal{R}_2$ be relations on $\mathcal{X}$. Then, $\mathcal{R}_1 \cap \mathcal{R}_2$, $\mathcal{R}_1 \setminus \mathcal{R}_2$, and $\mathcal{R}_1 \cup \mathcal{R}_2$ are relations on $\mathcal{X}$. Furthermore, if $\mathcal{R}$ is a relation on $\mathcal{X}$ and $\mathcal{X}_0 \subseteq \mathcal{X}$, then we define $\mathcal{R}|_{\mathcal{X}_0} \triangleq \mathcal{R} \cap (\mathcal{X}_0 \times \mathcal{X}_0)$, which is a relation on $\mathcal{X}_0$.

The following result shows that relations can be viewed as generalizations of functions.

Proposition 1.3.1. Let $\mathcal{X}_1$ and $\mathcal{X}_2$ be sets, and let $\mathcal{R}$ be a relation $\mathcal{X}_1 \times \mathcal{X}_2$. Then, there exists a function $f: \mathcal{X}_1 \mapsto \mathcal{X}_2$ such that $\mathcal{R} = \text{Graph}(f)$ if and only if, for all $x \in \mathcal{X}_1$, there exists a unique $y \in \mathcal{X}_2$ such that $(x, y) \in \mathcal{R}$. In this case, $f(x) = y$.

Definition 1.3.2. Let $\mathcal{R}$ be a relation on $\mathcal{X}$. Then, the following terminology is defined:

- i) $\mathcal{R}$ is *reflexive* if, for all $x \in \mathcal{X}$, it follows that $(x, x) \in \mathcal{R}$.
- ii) $\mathcal{R}$ is *symmetric* if, for all $(x_1, x_2) \in \mathcal{R}$, it follows that $(x_2, x_1) \in \mathcal{R}$.
- iii) $\mathcal{R}$ is *transitive* if, for all $(x_1, x_2) \in \mathcal{R}$ and $(x_2, x_3) \in \mathcal{R}$, it follows that $(x_1, x_3) \in \mathcal{R}$.
- iv) $\mathcal{R}$ is an *equivalence relation* if $\mathcal{R}$ is reflexive, symmetric, and transitive.

Proposition 1.3.3. Let $\mathcal{R}_1$ and $\mathcal{R}_2$ be relations on $\mathcal{X}$. If $\mathcal{R}_1$ and $\mathcal{R}_2$ are (reflexive, symmetric) relations, then so are $\mathcal{R}_1 \cap \mathcal{R}_2$ and $\mathcal{R}_1 \cup \mathcal{R}_2$. If $\mathcal{R}_1$ and $\mathcal{R}_2$ are (transitive, equivalence) relations, then so is $\mathcal{R}_1 \cap \mathcal{R}_2$.

Definition 1.3.4. Let $\mathcal{R}$ be a relation on $\mathcal{X}$. Then, the following terminology is defined:

- i) The *complement* $\mathcal{R}^\sim$ of $\mathcal{R}$ is the relation $\mathcal{R}^\sim \triangleq (\mathcal{X} \times \mathcal{X}) \setminus \mathcal{R}$.
- ii) The *support* $\text{supp}(\mathcal{R})$ of $\mathcal{R}$ is the smallest subset $\mathcal{X}_0$ of $\mathcal{X}$ such that $\mathcal{R}$ is a relation on $\mathcal{X}_0$.

- iii) The *reversal* $\text{rev}(\mathcal{R})$ of $\mathcal{R}$ is the relation $\text{rev}(\mathcal{R}) \triangleq \{(y, x) : (x, y) \in \mathcal{R}\}$.
- iv) The *shortcut* $\text{shortcut}(\mathcal{R})$ of $\mathcal{R}$ is the relation $\text{shortcut}(\mathcal{R}) \triangleq \{(x, y) \in \mathcal{X} \times \mathcal{X} : x \text{ and } y \text{ are distinct and there exist } k \geq 1 \text{ and } x_1, \dots, x_k \in \mathcal{X} \text{ such that } (x, x_1), (x_1, x_2), \dots, (x_k, y) \in \mathcal{R}\}$.
- v) The *reflexive hull* $\text{ref}(\mathcal{R})$ of $\mathcal{R}$ is the smallest reflexive relation on $\mathcal{X}$ that contains $\mathcal{R}$.
- vi) The *symmetric hull* $\text{sym}(\mathcal{R})$ of $\mathcal{R}$ is the smallest symmetric relation on $\mathcal{X}$ that contains $\mathcal{R}$.
- vii) The *transitive hull* $\text{trans}(\mathcal{R})$ of $\mathcal{R}$ is the smallest transitive relation on $\mathcal{X}$ that contains $\mathcal{R}$.
- viii) The *equivalence hull* $\text{equiv}(\mathcal{R})$ of $\mathcal{R}$ is the smallest equivalence relation on $\mathcal{X}$ that contains $\mathcal{R}$.

Proposition 1.3.5. Let $\mathcal{R}$ be a relation on $\mathcal{X}$. Then, the following statements hold:

- i) $\text{ref}(\mathcal{R}) = \mathcal{R} \cup \{(x, x) : x \in \mathcal{X}\}$.
- ii) $\text{sym}(\mathcal{R}) = \mathcal{R} \cup \text{rev}(\mathcal{R})$.
- iii) $\text{trans}(\mathcal{R}) = \mathcal{R} \cup \text{shortcut}(\mathcal{R})$.
- iv) $\text{equiv}(\mathcal{R}) = \mathcal{R} \cup \text{ref}(\mathcal{R}) \cup \text{sym}(\mathcal{R}) \cup \text{trans}(\mathcal{R})$.
- v) $\text{equiv}(\mathcal{R}) = \mathcal{R} \cup \text{ref}(\mathcal{R}) \cup \text{rev}(\mathcal{R}) \cup \text{shortcut}(\mathcal{R})$.

Furthermore, the following statements hold:

- vi) $\mathcal{R}$ is reflexive if and only if $\mathcal{R} = \text{ref}(\mathcal{R})$.
- vii) $\mathcal{R}$ is symmetric if and only if $\mathcal{R} = \text{rev}(\mathcal{R})$.
- viii) $\mathcal{R}$ is transitive if and only if $\mathcal{R} = \text{trans}(\mathcal{R})$.
- ix) $\mathcal{R}$ is an equivalence relation if and only if $\mathcal{R} = \text{equiv}(\mathcal{R})$.

For an equivalence relation $\mathcal{R}$ on $\mathcal{X}$, $(x_1, x_2) \in \mathcal{R}$ is denoted by $x_1 \stackrel{\mathcal{R}}{=} x_2$. If $\mathcal{R}$ is an equivalence relation and $x \in \mathcal{X}$, then the subset $\mathcal{E}_x \triangleq \{y \in \mathcal{X} : y \stackrel{\mathcal{R}}{=} x\}$ of $\mathcal{X}$ is the *equivalence class of x induced by $\mathcal{R}$* .

Theorem 1.3.6. Let $\mathcal{R}$ be an equivalence relation on a set $\mathcal{X}$. Then, the set $\{\mathcal{E}_x : x \in \mathcal{X}\}$ of equivalence classes induced by $\mathcal{R}$ is a partition of $\mathcal{X}$.

Proof. Since $\mathcal{X} = \bigcup_{x \in \mathcal{X}} \mathcal{E}_x$, it suffices to show that if $x, y \in \mathcal{X}$, then either $\mathcal{E}_x = \mathcal{E}_y$ or $\mathcal{E}_x \cap \mathcal{E}_y = \emptyset$. Hence, let $x, y \in \mathcal{X}$, and suppose that $\mathcal{E}_x$ and $\mathcal{E}_y$ are not disjoint so that there exists $z \in \mathcal{E}_x \cap \mathcal{E}_y$. Thus, $(x, z) \in \mathcal{R}$ and $(z, y) \in \mathcal{R}$. Now, let $w \in \mathcal{E}_x$. Then, $(w, x) \in \mathcal{R}$, $(x, z) \in \mathcal{R}$, and $(z, y) \in \mathcal{R}$ imply that $(w, y) \in \mathcal{R}$. Hence, $w \in \mathcal{E}_y$, which implies that $\mathcal{E}_x \subseteq \mathcal{E}_y$. By a similar argument, $\mathcal{E}_y \subseteq \mathcal{E}_x$. Consequently, $\mathcal{E}_x = \mathcal{E}_y$. $\square$

The following result, which is the converse of Theorem 1.3.6, shows that a partition of a set $\mathcal{X}$ defines an equivalence relation on $\mathcal{X}$.

Theorem 1.3.7. Let $\mathcal{X}$ be a set, consider a partition of $\mathcal{X}$, and define the relation $\mathcal{R}$ on $\mathcal{X}$ by $(x, y) \in \mathcal{R}$ if and only if x and y belong to the same partition subset of $\mathcal{X}$. Then, $\mathcal{R}$ is an equivalence relation on $\mathcal{X}$.

Definition 1.3.8. Let $\mathcal{R}$ be a relation on $\mathcal{X}$. Then, the following terminology is defined:

- i) $\mathcal{R}$ is *antisymmetric* if $(x_1, x_2) \in \mathcal{R}$ and $(x_2, x_1) \in \mathcal{R}$ imply that $x_1 = x_2$.
- ii) $\mathcal{R}$ is a *partial ordering* on $\mathcal{X}$ if $\mathcal{R}$ is reflexive, antisymmetric, and transitive.

Let $\mathcal{R}$ be a partial ordering on $\mathcal{X}$. Then, $(x_1, x_2) \in \mathcal{R}$ is denoted by $x_1 \stackrel{\mathcal{R}}{\leq} x_2$. If $x_1 \stackrel{\mathcal{R}}{\leq} x_2$ and $x_2 \stackrel{\mathcal{R}}{\leq} x_1$, then, since $\mathcal{R}$ is antisymmetric, it follows that $x_1 = x_2$. Furthermore, if $x_1 \stackrel{\mathcal{R}}{\leq} x_2$ and $x_2 \stackrel{\mathcal{R}}{\leq} x_3$, then, since $\mathcal{R}$ is transitive, it follows that $x_1 \stackrel{\mathcal{R}}{\leq} x_3$.

Definition 1.3.9. Let " $\stackrel{\mathcal{R}}{\leq}$ " be a partial ordering on $\mathcal{X}$. Then, the following terminology is defined:

- i) Let $\mathcal{S} \subseteq \mathcal{X}$. Then, $y \in \mathcal{X}$ is a *lower bound* for $\mathcal{S}$ if, for all $x \in \mathcal{S}$, it follows that $y \stackrel{\mathcal{R}}{\leq} x$.
- ii) Let $\mathcal{S} \subseteq \mathcal{X}$. Then, $y \in \mathcal{X}$ is an *upper bound* for $\mathcal{S}$ if, for all $x \in \mathcal{S}$, it follows that $x \stackrel{\mathcal{R}}{\leq} y$.
- iii) Let $\mathcal{S} \subseteq \mathcal{X}$. Then, $y \in \mathcal{X}$ is the *least upper bound* $\text{lub}(\mathcal{S})$ for $\mathcal{S}$ if y is an upper bound for $\mathcal{S}$ and, for all upper bounds $x \in \mathcal{X}$ for $\mathcal{S}$, it follows that $y \stackrel{\mathcal{R}}{\leq} x$. In this case, we write $y = \text{lub}(\mathcal{S})$.
- iv) Let $\mathcal{S} \subseteq \mathcal{X}$. Then, $y \in \mathcal{X}$ is the *greatest lower bound* for $\mathcal{S}$ if y is a lower bound for $\mathcal{S}$ and, for all lower bounds $x \in \mathcal{X}$ for $\mathcal{S}$, it follows that $x \stackrel{\mathcal{R}}{\leq} y$. In this case, we write $y = \text{glb}(\mathcal{S})$.
- v) $\stackrel{\mathcal{R}}{\leq}$ is a *lattice* on $\mathcal{X}$ if, for all distinct $x, y \in \mathcal{X}$, the set $\{x, y\}$ has a least upper bound and a greatest lower bound.
- vi) $\mathcal{R}$ is a *total ordering* on $\mathcal{X}$ if, for all $x, y \in \mathcal{X}$, it follows that either $(x, y) \in \mathcal{R}$ or $(y, x) \in \mathcal{R}$.

For a subset $\mathcal{S}$ of the real numbers, it is traditional to write $\inf \mathcal{S}$ and $\sup \mathcal{S}$ for $\text{glb}(\mathcal{S})$ and $\text{lub}(\mathcal{S})$, respectively, where "inf" and "sup" denote infimum and supremum, respectively.

1.4 Graphs

Let $\mathcal{X}$ be a finite, nonempty set, and let $\mathcal{R}$ be a relation on $\mathcal{X}$. Then, the pair $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ is a *graph*. The elements of $\mathcal{X}$ are the *nodes* of $\mathcal{G}$, while the elements of $\mathcal{R}$ are the *arcs* of $\mathcal{G}$. If $\mathcal{R}$ is a multirelation on $\mathcal{X}$, then $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ is a *multigraph*.

The graph $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ can be visualized as a set of points in the plane representing the nodes in $\mathcal{X}$ connected by the arcs in $\mathcal{R}$. Specifically, the arc $(x, y) \in \mathcal{R}$ from x to y can be visualized as a directed line segment or curve connecting node x to node y . The direction of an arc can be denoted by an arrow head. For example, consider a graph that represents a city with streets (arcs) connecting houses (nodes). Then, a symmetric relation is a street plan with no one-way streets, whereas an antisymmetric relation is a street plan with no two-way streets.

Definition 1.4.1. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph. Then, the following terminology is defined:

- i) The *reversal* of $\mathcal{G}$ is the graph $\text{rev}(\mathcal{G}) \triangleq (\mathcal{X}, \text{rev}(\mathcal{R}))$.
- ii) The *complement* of $\mathcal{G}$ is the graph $\mathcal{G}^\sim \triangleq (\mathcal{X}, \mathcal{R}^\sim)$.
- iii) The *reflexive hull* of $\mathcal{G}$ is the graph $\text{ref}(\mathcal{G}) \triangleq (\mathcal{X}, \text{ref}(\mathcal{R}))$.
- iv) The *symmetric hull* of $\mathcal{G}$ is the graph $\text{sym}(\mathcal{G}) \triangleq (\mathcal{X}, \text{sym}(\mathcal{R}))$.
- v) The *transitive hull* of $\mathcal{G}$ is the graph $\text{trans}(\mathcal{G}) \triangleq (\mathcal{X}, \text{trans}(\mathcal{R}))$.
- vi) The *equivalence hull* of $\mathcal{G}$ is the graph $\text{equiv}(\mathcal{G}) \triangleq (\mathcal{X}, \text{equiv}(\mathcal{R}))$.
- vii) $\mathcal{G}$ is *reflexive* if $\mathcal{R}$ is reflexive.
- viii) $\mathcal{G}$ is *symmetric* if $\mathcal{R}$ is symmetric. In this case, the arcs (x, y) and (y, x) in $\mathcal{R}$ are denoted by the subset $\{x, y\}$ of $\mathcal{X}$, called an *edge*.
- ix) $\mathcal{G}$ is *transitive* if $\mathcal{R}$ is transitive.
- x) $\mathcal{G}$ is an *equivalence graph* if $\mathcal{R}$ is an equivalence relation.
- xi) $\mathcal{G}$ is *antisymmetric* if $\mathcal{R}$ is antisymmetric.
- xii) $\mathcal{G}$ is *partially ordered* if $\mathcal{R}$ is a partial ordering on $\mathcal{X}$.
- xiii) $\mathcal{G}$ is *totally ordered* if $\mathcal{R}$ is a total ordering on $\mathcal{X}$.
- xiv) $\mathcal{G}$ is a *tournament* if $\mathcal{G}$ has no self-loops, is antisymmetric, and $\text{sym}(\mathcal{R}) = \mathcal{X} \times \mathcal{X}$.

Definition 1.4.2. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph. Then, the following terminology is defined:

- i) The arc $(x, x) \in \mathcal{R}$ is a *self-loop*.
- ii) The *reversal* of $(x, y) \in \mathcal{R}$ is (y, x) .
- iii) If $x, y \in \mathcal{X}$ and $(x, y) \in \mathcal{R}$, then y is the *head* of (x, y) and x is the *tail* of (x, y) .

- iv) If $x, y \in \mathcal{X}$ and $(x, y) \in \mathcal{R}$, then x is a *parent* of y , and y is a *child* of x .
- v) If $x, y \in \mathcal{X}$ and either $(x, y) \in \mathcal{R}$ or $(y, x) \in \mathcal{R}$, then x and y are *adjacent*.
- vi) If $x \in \mathcal{X}$ has no parent, then x is a *root*.
- vii) If $x \in \mathcal{X}$ has no child, then x is a *leaf*.

Suppose that $(x, x) \in \mathcal{R}$. Then, x is both the head and the tail of (x, x) , and thus x is a parent and child of itself. Consequently, x is neither a root nor a leaf. Furthermore, x is adjacent to itself.

Definition 1.4.3. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph. Then, the following terminology is defined:

- i) The graph $\mathcal{G}' = (\mathcal{X}', \mathcal{R}')$ is a *subgraph* of $\mathcal{G}$ if $\mathcal{X}' \subseteq \mathcal{X}$ and $\mathcal{R}' \subseteq \mathcal{R}$.
- ii) The subgraph $\mathcal{G}' = (\mathcal{X}', \mathcal{R}')$ of $\mathcal{G}$ is a *spanning subgraph* of $\mathcal{G}$ if $\text{supp}(\mathcal{R}) = \text{supp}(\mathcal{R}')$.
- iii) For $x, y \in \mathcal{X}$, a *walk* in $\mathcal{G}$ from x to y is an n -tuple of arcs of the form $((x, y)) \in \mathcal{R}$ for $n = 1$ and $((x, x_1), (x_1, x_2), \dots, (x_{n-1}, y)) \in \mathcal{R}^n$ for $n \geq 2$. The *length* of the walk is n . The nodes $x, x_1, \dots, x_{n-1}, y$ are the *nodes* of the walk. Furthermore, if $n \geq 2$, then the nodes $x_1, \dots, x_{n-1}$ are the *intermediate nodes* of the walk.
- iv) $\mathcal{G}$ is *connected* if, for all distinct $x, y \in \mathcal{X}$, there exists a walk in $\mathcal{G}$ from x to y .
- v) For $x, y \in \mathcal{X}$, a *trail* in $\mathcal{G}$ from x to y is a walk in $\mathcal{G}$ from x to y whose arcs are distinct and such that no reversed arc is also an arc of $\mathcal{G}$.
- vi) For $x, y \in \mathcal{X}$, a *path* in $\mathcal{G}$ from x to y is a trail in $\mathcal{G}$ from x to y whose intermediate nodes (if any) are distinct.
- vii) $\mathcal{G}$ is *traceable* if $\mathcal{G}$ has a path such that every node in $\mathcal{X}$ is a node of the path. Such a path is called a *Hamiltonian path*.
- viii) For $x \in \mathcal{X}$, a *cycle* in $\mathcal{G}$ at x is a path in $\mathcal{G}$ from x to x whose length is greater than 1.
- ix) The *period* of $\mathcal{G}$ is the greatest common divisor of the lengths of the cycles in $\mathcal{G}$. Furthermore, $\mathcal{G}$ is *aperiodic* if the period of $\mathcal{G}$ is 1.
- x) $\mathcal{G}$ is *Hamiltonian* if $\mathcal{G}$ has a cycle such that every node in $\mathcal{X}$ is a node of the cycle. Such a cycle is called a *Hamiltonian cycle*.
- xi) $\mathcal{G}$ is a *forest* if $\mathcal{G}$ is symmetric and has no cycles.
- xii) $\mathcal{G}$ is a *tree* if $\mathcal{G}$ is a forest and is connected.
- xiii) The *indegree* of $x \in \mathcal{X}$ is $\text{indeg}(x) \triangleq \text{card}\{y \in \mathcal{X} : y \text{ is a parent of } x\}$.
- xiv) The *outdegree* of $x \in \mathcal{X}$ is $\text{outdeg}(x) \triangleq \text{card}\{y \in \mathcal{X} : y \text{ is a child of } x\}$.
- xv) If $\mathcal{G}$ is symmetric, then the *degree* of $x \in \mathcal{X}$ is $\text{deg}(x) \triangleq \text{indeg}(x) = \text{outdeg}(x)$.

xvi) If $\mathcal{X}_0 \subseteq \mathcal{X}$, then,

$$\mathcal{G}|_{\mathcal{X}_0} \triangleq (\mathcal{X}_0, \mathcal{R}|_{\mathcal{X}_0}).$$

xvii) If $\mathcal{G}' = (\mathcal{X}', \mathcal{R}')$ is a graph, then $\mathcal{G} \cup \mathcal{G}' \triangleq (\mathcal{X} \cup \mathcal{X}', \mathcal{R} \cup \mathcal{R}')$ and $\mathcal{G} \cap \mathcal{G}' \triangleq (\mathcal{X} \cap \mathcal{X}', \mathcal{R} \cap \mathcal{R}')$.

xviii) Let $\mathcal{X} = \mathcal{X}_1 \cup \mathcal{X}_2$, where $\mathcal{X}_1$ and $\mathcal{X}_2$ are nonempty and disjoint, and assume that $\mathcal{X} = \text{supp}(\mathcal{G})$. Then, $(\mathcal{X}_1, \mathcal{X}_2)$ is a *directed cut* of $\mathcal{G}$ if, for all $x_1 \in \mathcal{X}_1$ and $x_2 \in \mathcal{X}_2$, there does not exist a walk from x_1 to x_2 .

Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph, and let $w: \mathcal{X} \times \mathcal{X} \mapsto [0, \infty)$, where $w(x, y) > 0$ if $(x, y) \in \mathcal{R}$ and $w(x, y) = 0$ if $(x, y) \notin \mathcal{R}$. For each arc $(x, y) \in \mathcal{R}$, $w(x, y)$ is the *weight* associated with the arc (x, y) , and the triple $\mathcal{G} = (\mathcal{X}, \mathcal{R}, w)$ is a *weighted graph*. Every graph can be viewed as a weighted graph by defining $w[(x, y)] \triangleq 1$ for all $(x, y) \in \mathcal{R}$ and $w[(x, y)] \triangleq 0$ for all $(x, y) \notin \mathcal{R}$. The graph $\mathcal{G}' = (\mathcal{X}', \mathcal{R}', w')$ is a *weighted subgraph* of $\mathcal{G}$ if $\mathcal{X} \subseteq \mathcal{X}'$, $\mathcal{R}'$ is a relation on $\mathcal{X}'$, $\mathcal{R}' \subseteq \mathcal{R}$, and w' is the restriction of w to $\mathcal{R}'$. Finally, if $\mathcal{G}$ is symmetric, then w is defined on edges $\{x, y\}$ of $\mathcal{G}$.

1.5 Facts on Logic, Sets, Functions, and Relations

Fact 1.5.1. Let A and B be statements. Then, the following statements hold:

- i) $\text{not}(A \text{ or } B) \iff [(\text{not } A) \text{ and } (\text{not } B)]$.
- ii) $\text{not}(A \text{ and } B) \iff (\text{not } A) \text{ or } (\text{not } B)$.
- iii) $(A \text{ or } B) \iff [(\text{not } A) \implies B]$.
- iv) $[(\text{not } A) \text{ or } B] \iff (A \implies B)$.
- v) $[A \text{ and } (\text{not } B)] \iff [\text{not}(A \implies B)]$.

(Remark: Each statement is a tautology.) (Remark: Statements *i*) and *ii*) are *De Morgan's laws*. See [229, p. 24].)

Fact 1.5.2. The following statements are equivalent:

- i) $A \implies (B \text{ or } C)$.
- ii) $[A \text{ and } (\text{not } B)] \implies C$.

(Remark: The statement that *i*) and *ii*) are equivalent is a tautology.)

Fact 1.5.3. The following statements are equivalent:

$$i) A \iff B.$$

$$ii) [A \text{ or } (\text{not } B)] \text{ and } (\text{not } [A \text{ and } (\text{not } B)]).$$

(Remark: The statement that $i)$ and $ii)$ are equivalent is a tautology.)

Fact 1.5.4. The following statements are equivalent:

$$i) \text{ Not } [\text{for all } x, \text{ there exists } y \text{ such that statement } Z \text{ is satisfied}].$$

$$ii) \text{ There exists } x \text{ such that, for all } y, \text{ statement } Z \text{ is not satisfied.}$$

Fact 1.5.5. Let $\mathcal{A}$, $\mathcal{B}$, and $\mathcal{C}$ be sets, and assume that each of these sets has a finite number of elements. Then,

$$\text{card}(\mathcal{A} \cup \mathcal{B}) = \text{card}(\mathcal{A}) + \text{card}(\mathcal{B}) - \text{card}(\mathcal{A} \cap \mathcal{B})$$

and

$$\begin{aligned} \text{card}(\mathcal{A} \cup \mathcal{B} \cup \mathcal{C}) &= \text{card}(\mathcal{A}) + \text{card}(\mathcal{B}) + \text{card}(\mathcal{C}) \\ &\quad - \text{card}(\mathcal{A} \cap \mathcal{B}) - \text{card}(\mathcal{A} \cap \mathcal{C}) - \text{card}(\mathcal{B} \cap \mathcal{C}) \\ &\quad + \text{card}(\mathcal{A} \cap \mathcal{B} \cap \mathcal{C}). \end{aligned}$$

(Remark: This result is the *inclusion-exclusion principle*. See [177, p. 82] or [1218, pp. 64–67].)

Fact 1.5.6. Let $\mathcal{A}, \mathcal{B}, \mathcal{C}$ be subsets of a set $\mathcal{X}$. Then, the following identities hold:

$$i) \mathcal{A} \cap \mathcal{A} = \mathcal{A} \cup \mathcal{A} = \mathcal{A}.$$

$$ii) (\mathcal{A} \cup \mathcal{B})^\sim = \mathcal{A}^\sim \cap \mathcal{B}^\sim.$$

$$iii) (\mathcal{A} \cap \mathcal{B})^\sim = \mathcal{A}^\sim \cup \mathcal{B}^\sim.$$

$$iv) \mathcal{A} = (\mathcal{A} \setminus \mathcal{B}) \cup (\mathcal{A} \cap \mathcal{B}).$$

$$v) [\mathcal{A} \setminus (\mathcal{A} \cap \mathcal{B})] \cup \mathcal{B} = \mathcal{A} \cup \mathcal{B}.$$

$$vi) (\mathcal{A} \cup \mathcal{B}) \setminus (\mathcal{A} \cap \mathcal{B}) = (\mathcal{A} \cap \mathcal{B}^\sim) \cup (\mathcal{A}^\sim \cap \mathcal{B}).$$

$$vii) \mathcal{A} \cap (\mathcal{B} \cup \mathcal{C}) = (\mathcal{A} \cap \mathcal{B}) \cup (\mathcal{A} \cap \mathcal{C}).$$

$$viii) \mathcal{A} \cup (\mathcal{B} \cap \mathcal{C}) = (\mathcal{A} \cup \mathcal{B}) \cap (\mathcal{A} \cup \mathcal{C}).$$

$$ix) (\mathcal{A} \setminus \mathcal{B}) \setminus \mathcal{C} = \mathcal{A} \setminus (\mathcal{B} \cup \mathcal{C}).$$

$$x) (\mathcal{A} \cap \mathcal{B}) \setminus \mathcal{C} = (\mathcal{A} \setminus \mathcal{C}) \cap (\mathcal{B} \setminus \mathcal{C}).$$

$$xi) (\mathcal{A} \cap \mathcal{B}) \setminus (\mathcal{C} \cap \mathcal{B}) = (\mathcal{A} \setminus \mathcal{C}) \cap \mathcal{B}.$$

$$xii) (\mathcal{A} \cup \mathcal{B}) \setminus \mathcal{C} = (\mathcal{A} \setminus \mathcal{C}) \cup (\mathcal{B} \setminus \mathcal{C}) = [\mathcal{A} \setminus (\mathcal{B} \cup \mathcal{C})] \cup (\mathcal{B} \setminus \mathcal{C}).$$

$$xiii) (\mathcal{A} \cup \mathcal{B}) \setminus (\mathcal{C} \cap \mathcal{B}) = (\mathcal{A} \setminus \mathcal{B}) \cup (\mathcal{B} \setminus \mathcal{C}).$$

$$xiv) (\mathcal{A} \cup \mathcal{B}) \cap (\mathcal{A} \cup \mathcal{B}^\sim) = \mathcal{A}.$$

$$xv) (\mathcal{A} \cup \mathcal{B}) \cap (\mathcal{A}^\sim \cup \mathcal{B}) \cap (\mathcal{A} \cup \mathcal{B}^\sim) = \mathcal{A} \cap \mathcal{B}.$$

Fact 1.5.7. Define the relation $\mathcal{R}$ on $\mathbb{R} \times \mathbb{R}$ by

$$\mathcal{R} \triangleq \{((x_1, y_1), (x_2, y_2)) \in (\mathbb{R} \times \mathbb{R}) \times (\mathbb{R} \times \mathbb{R}) : x_1 \leq x_2 \text{ and } y_1 \leq y_2\}.$$

Then, $\mathcal{R}$ is a partial ordering.

Fact 1.5.8. Define the relation $\mathcal{L}$ on $\mathbb{R} \times \mathbb{R}$ by

$$\mathcal{L} \triangleq \{((x_1, y_1), (x_2, y_2)) \in (\mathbb{R} \times \mathbb{R}) \times (\mathbb{R} \times \mathbb{R}) : \\ x_1 \leq x_2 \text{ and, if } x_1 = x_2, \text{ then } y_1 \leq y_2\}.$$

Then, $\mathcal{L}$ is a total ordering on $\mathbb{R} \times \mathbb{R}$. (Remark: Denoting this total ordering by “ $\stackrel{d}{\leq}$,” note that $(1, 4) \stackrel{d}{\leq} (2, 3)$ and $(1, 4) \stackrel{d}{\leq} (1, 5)$.) (Remark: This ordering is the *lexicographic ordering* or *dictionary ordering*, where ‘book’ $\stackrel{d}{\leq}$ ‘box’. Note that the ordering of words in a dictionary is reflexive, antisymmetric, and transitive, and that every pair of words can be ordered.) (Remark: See Fact 2.9.31.)

Fact 1.5.9. Let $f: \mathcal{X} \mapsto \mathcal{Y}$, and assume that f is invertible. Then,

$$(f^{-1})^{-1} = f.$$

Fact 1.5.10. Let $f: \mathcal{X} \mapsto \mathcal{Y}$ and $g: \mathcal{Y} \mapsto \mathcal{Z}$, and assume that f and g are invertible. Then, $g \bullet f$ is invertible and

$$(g \bullet f)^{-1} = f^{-1} \bullet g^{-1}.$$

Fact 1.5.11. Let $f: \mathcal{X} \mapsto \mathcal{Y}$, and let $A, B \subseteq \mathcal{X}$. Then, the following statements hold:

- i) If $A \subseteq B$, then $f(A) \subseteq f(B)$.
- ii) $f(A \cup B) = f(A) \cup f(B)$.
- iii) $f(A \cap B) \subseteq f(A) \cap f(B)$.

Fact 1.5.12. Let $f: \mathcal{X} \mapsto \mathcal{Y}$, and let $A, B \subseteq \mathcal{Y}$. Then, the following statements hold:

- i) $f[f^{-1}(A)] \subseteq A \subseteq f^{-1}[f(A)]$.
- ii) $f^{-1}(A \cup B) = f^{-1}(A) \cup f^{-1}(B)$.
- iii) $f^{-1}(A \cap B) = f^{-1}(A) \cap f^{-1}(B)$.

Fact 1.5.13. Let $\mathcal{X}$ and $\mathcal{Y}$ be finite sets, assume that $\text{card}(\mathcal{X}) = \text{card}(\mathcal{Y})$, and let $f: \mathcal{X} \mapsto \mathcal{Y}$. Then, f is one-to-one if and only if f is onto. (Remark: See Fact 1.6.1.)

Fact 1.5.14. Let $f: \mathcal{X} \mapsto \mathcal{Y}$. Then, the following statements are equivalent:

- i) f is one-to-one.
- ii) For all $A \subseteq \mathcal{X}$ and $B \subseteq \mathcal{Y}$, it follows that $f(A \cap B) = f(A) \cap f(B)$.
- iii) For all $A \subseteq \mathcal{X}$, it follows that $f^{-1}[f(A)] = A$.

- iv) For all disjoint $A \subseteq \mathcal{X}$ and $B \subseteq \mathcal{Y}$, it follows that $f(A)$ and $f(B)$ are disjoint.
- v) For all $A \subseteq \mathcal{X}$ and $B \subseteq \mathcal{Y}$ such that $A \subseteq B$, it follows that $f(A \setminus B) = f(A) \setminus f(B)$.

(Proof: See [68, pp. 44, 45].)

Fact 1.5.15. Let $f: \mathcal{X} \mapsto \mathcal{Y}$. Then, the following statements are equivalent:

- i) f is onto.
- ii) For all $A \subseteq \mathcal{X}$, it follows that $f[f^{-1}(A)] = A$.

Fact 1.5.16. Let $f: \mathcal{X} \mapsto \mathcal{Y}$, and let $g: \mathcal{Y} \mapsto \mathcal{Z}$. Then, the following statements hold:

- i) If f and g are one-to-one, then $f \bullet g$ is one-to-one.
- ii) If f and g are onto, then $f \bullet g$ is onto.

(Remark: A matrix version of this result is given by Fact 2.10.3.)

Fact 1.5.17. Let $\mathcal{X}$ be a set, and let $\mathfrak{X}$ denote the class of subsets of $\mathcal{X}$. Then, “ $\subset$ ” and “ $\subseteq$ ” are transitive relations on $\mathfrak{X}$, and “ $\subseteq$ ” is a partial ordering on $\mathfrak{X}$.

1.6 Facts on Graphs

Fact 1.6.1. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph. Then, the following statements hold:

- i) $\mathcal{R}$ is the graph of a function on $\mathcal{X}$ if and only if every node in $\mathcal{X}$ has exactly one child.

Furthermore, the following statements are equivalent:

- ii) $\mathcal{R}$ is the graph of a one-to-one function on $\mathcal{X}$.
- iii) $\mathcal{R}$ is the graph of an onto function on $\mathcal{X}$.
- iv) $\mathcal{R}$ is the graph of a one-to-one and onto function on $\mathcal{X}$.
- v) Every node in $\mathcal{X}$ has exactly one child and not more than one parent.
- vi) Every node in $\mathcal{X}$ has exactly one child and at least one parent.
- vii) Every node in $\mathcal{X}$ has exactly one child and exactly one parent.

(Remark: See Fact 1.5.13.)

Fact 1.6.2. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph, and assume that $\mathcal{R}$ is the graph of a function $f: \mathcal{X} \mapsto \mathcal{X}$. Then, either f is the identity map or $\mathcal{G}$ has a cycle.

Fact 1.6.3. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph, and assume that $\mathcal{G}$ has a Hamiltonian cycle. Then, $\mathcal{G}$ has no roots and no leaves.

Fact 1.6.4. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph. Then, $\mathcal{G}$ has either a root or a cycle.

Fact 1.6.5. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a symmetric graph. Then, the following statements are equivalent:

- i) $\mathcal{G}$ is a forest.
- ii) $\mathcal{G}$ has no cycles.
- iii) No pair of nodes is connected by more than one path.

Furthermore, the following statements are equivalent:

- iv) $\mathcal{G}$ is a tree.
- v) $\mathcal{G}$ is a connected forest.
- vi) $\mathcal{G}$ is connected and has no cycles.
- vii) $\mathcal{G}$ is connected and has $\text{card}(\mathcal{X}) - 1$ edges.
- viii) $\mathcal{G}$ has no cycles and has $\text{card}(\mathcal{X}) - 1$ edges.
- ix) Every pair of nodes is connected by exactly one path.

Fact 1.6.6. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a tournament. Then, $\mathcal{G}$ has a Hamiltonian path. Furthermore, the Hamiltonian path is a Hamiltonian cycle if and only if $\mathcal{G}$ is connected.

Fact 1.6.7. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a symmetric graph, where $\mathcal{X} \subset \mathbb{R}^2$, assume that $n \triangleq \text{card}(\mathcal{X}) \geq 3$, and assume that the edges in $\mathcal{R}$ can be represented by line segments lying in a plane that are either disjoint or intersect at a node. Furthermore, let m denote the number of edges of $\mathcal{G}$, and let f denote the number of disjoint regions in $\mathbb{R}^2$ whose boundaries are the edges of $\mathcal{G}$. Then,

$$n - m + f = 2.$$

Consequently, if $n \geq 3$, then

$$m \leq 3(n - 2).$$

(Remark: The identity is *Euler's polyhedron formula*.)

1.7 Facts on Binomial Identities and Sums

Fact 1.7.1. The following identities hold:

- i) Let $0 \leq k \leq n$. Then,

$$\binom{n}{k} = \binom{n}{n-k}.$$

- ii) Let $1 \leq k \leq n$. Then,

$$k \binom{n}{k} = n \binom{n-1}{k-1}.$$

iii) Let $2 \leq k \leq n$. Then,

$$k(k-1) \binom{n}{k} = n(n-1) \binom{n-2}{k-2}.$$

iv) Let $0 \leq k < n$. Then,

$$(n-k) \binom{n}{k} = n \binom{n-1}{k}.$$

v) Let $1 \leq k \leq n$. Then,

$$\binom{n}{k} = \binom{n-1}{k} + \binom{n-1}{k-1}.$$

vi) Let $0 \leq m \leq k \leq n$. Then,

$$\binom{n}{k} \binom{k}{m} = \binom{n}{m} \binom{n-m}{k-m}.$$

vii) Let $m, n \geq 0$. Then,

$$\sum_{i=0}^m \binom{n+i}{n} = \binom{n+m+1}{m}.$$

viii) Let $k \geq 0$ and $n \geq 1$. Then,

$$\sum_{i=0}^{n-1} \frac{(k+i)!}{i!} = k! \binom{k+n}{k+1}.$$

ix) Let $0 \leq k \leq n$. Then,

$$\sum_{i=k}^n \binom{i}{k} = \binom{n+1}{k+1}.$$

x) Let $n, m \geq 0$, and let $0 \leq k \leq \min\{n, m\}$. Then,

$$\sum_{i=0}^k \binom{n}{i} \binom{m}{k-i} = \binom{n+m}{k}.$$

xi) Let $n \geq 0$. Then,

$$\sum_{i=1}^n \binom{n}{i} \binom{n}{i-1} = \binom{2n}{n+1}.$$

xii) Let $0 \leq k \leq n$. Then,

$$\sum_{i=0}^{n-k} \binom{n}{i} \binom{n}{k+i} = \frac{(2n)!}{(n-k)!(n+k)!}.$$

xiii) Let $0 \leq k \leq n/2$. Then,

$$\sum_{i=k}^{n-k} \binom{i}{k} \binom{n-i}{k} = \binom{n+1}{2k+1}.$$

xiv) Let $n \geq 0$. Then,

$$\sum_{i=0}^n \binom{n}{i}^2 = \binom{2n}{n}.$$

xv) Let $n \geq 1$. Then,

$$\sum_{i=0}^n i \binom{n}{i}^2 = n \binom{2n-1}{n-1}.$$

xvi) For all $x, y \in \mathbb{C}$ and $n \geq 0$,

$$(x+y)^n = \sum_{i=0}^n \binom{n}{i} x^{n-i} y^i.$$

xvii) Let $n \geq 0$. Then,

$$\sum_{i=0}^n \binom{n}{i} = 2^n.$$

xviii) Let $n \geq 0$. Then,

$$\sum_{i=0}^n \frac{1}{i+1} \binom{n}{i} = \frac{2^{n+1} - 1}{n+1}.$$

xix) Let $n \geq 0$. Then,

$$\sum_{i=0}^n \binom{2n+1}{i} = \sum_{i=0}^{2n} \binom{2n}{i} = 4^n.$$

xx) Let $n > 1$. Then,

$$\sum_{i=0}^{n-1} (n-i)^2 \binom{2n}{i} = 4^{n-1} n.$$

xxi) Let $n \geq 0$. Then,

$$\sum_{i=0}^{\lfloor n/2 \rfloor} \binom{n}{2i} = 2^{n-1}.$$

xxii) Let $n \geq 0$. Then,

$$\sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \binom{n}{2i+1} = 2^{n-1}.$$

xxiii) Let $n \geq 0$. Then,

$$\sum_{i=0}^{\lfloor n/2 \rfloor} (-1)^i \binom{n}{2i} = 2^{n/2} \cos \frac{n\pi}{4}.$$

xxiv) Let $n \geq 0$. Then,

$$\sum_{i=0}^{\lfloor (n-1)/2 \rfloor} (-1)^i \binom{n}{2i+1} = 2^{n/2} \sin \frac{n\pi}{4}.$$

xxv) Let $n \geq 1$. Then,

$$\sum_{i=1}^n i \binom{n}{i} = n 2^{n-1}.$$

xxvi) Let $n \geq 1$. Then,

$$\sum_{i=0}^n \binom{n}{2i} = 2^{n-1}.$$

xxvii) Let $0 \leq k < n$. Then,

$$\sum_{i=0}^k (-1)^i \binom{n}{i} = (-1)^k \binom{n-1}{k}.$$

xxviii) Let $n \geq 1$. Then,

$$\sum_{i=0}^n (-1)^i \binom{n}{i} = 0.$$

xxix) Let $n \geq 1$. Then,

$$\sum_{i=0}^n \frac{2^i}{i+1} = \frac{2^n}{n+1} \sum_{i=0}^n \frac{1}{\binom{n}{i}}.$$

(Proof: See [177, pp. 64–68, 78], [332], [584, pp. 1, 2], and [668, pp. 2–10, 74]. Statement *xxix*) is given in [238, p. 55].) (Remark: Statement *x*) is *Vandermonde's identity*.)

Fact 1.7.2. The following inequalities hold:

i) Let $n \geq 2$. Then,

$$\frac{4^n}{n+1} < \binom{2n}{n} < 4^n.$$

ii) Let $n \geq 7$. Then,

$$\left(\frac{n}{3}\right)^n < n! < \left(\frac{n}{2}\right)^n.$$

iii) Let $1 \leq k \leq n$. Then,

$$\left(\frac{n}{k}\right)^k \leq \binom{n}{k} \leq \min \left\{ \frac{n^k}{k!}, \left(\frac{n e}{k}\right)^k \right\}.$$

iv) Let $0 \leq k \leq n$. Then,

$$(n+1)^k \binom{n}{k} \leq n^k \binom{n+1}{k}.$$

v) Let $1 \leq k \leq n-1$. Then,

$$\sum_{i=1}^k i(i+1) \binom{2n}{k-i} < \frac{2^{2n-2} k(k+1)}{n}.$$

vi) Let $1 \leq k \leq n$. Then,

$$n^k \leq k^{k/2} (k+1)^{(k-1)/2} \binom{n}{k}.$$

(Proof: Statements *i*) and *ii*) are given in [238, p. 210]. Statement *iv*) is given in [668, p. 111]. Statement *vi*) is given in [451].)

Fact 1.7.3. Let n be a positive integer. Then,

$$\sum_{i=1}^n i = \frac{1}{2} n(n+1),$$

$$\begin{aligned}
\sum_{i=1}^n (2i-1) &= n^2, \\
\sum_{i=1}^n i^2 &= \frac{1}{6}n(n+1)(2n+1), \\
\sum_{i=1}^n i^3 &= \frac{1}{4}n^2(n+1)^2 = \left(\sum_{i=1}^n i\right)^2, \\
\sum_{i=1}^n i^4 &= \frac{1}{30}n(n+1)(2n+1)(3n^2+3n-1), \\
\sum_{i=1}^n i^5 &= \frac{1}{12}n^2(n+1)^2(2n^2+2n-1).
\end{aligned}$$

(Remark: See Fact 1.15.9 and [668, p. 11].)

Fact 1.7.4. Let $n \geq 2$. Then,

$$n(\sqrt[n]{n+1} - 1) < \sum_{i=1}^n \frac{1}{i} < 1 + n\left(1 - \frac{1}{\sqrt[n]{n}}\right).$$

(Proof: See [668, pp. 158, 161].)

Fact 1.7.5. Let n be a positive integer. Then,

$$0 < \sum_{i=1}^n \frac{1}{i} < \log n$$

and

$$\lim_{n \rightarrow \infty} \left[\left(\sum_{i=1}^n \frac{1}{i} \right) - \log n \right] = \gamma \approx 0.57721 \dots$$

Hence,

$$\lim_{n \rightarrow \infty} \frac{\sum_{i=1}^n \frac{1}{i}}{\log n} = 1.$$

(Remark: γ is the *Euler constant*.)

Fact 1.7.6. The following statements hold:

$$\sum_{i=1}^{\infty} \frac{1}{i^i} = \int_0^1 \frac{1}{x^x} dx \approx 1.291$$

and

$$\sum_{i=1}^{\infty} (-1)^{i+1} \frac{1}{i^i} = \int_0^1 x^x dx.$$

(Proof: See [238, pp. 4, 44].)

Fact 1.7.7. The following statements hold:

$$\begin{aligned}
\sum_{i=0}^{\infty} \frac{1}{i!} &= e, \\
\sum_{i=1}^{\infty} \frac{1}{i^2} &= \frac{\pi^2}{6}, \\
\sum_{i=1}^{\infty} \frac{1}{i^4} &= \frac{\pi^4}{90}, \\
\sum_{i=1}^{\infty} \frac{1}{i^6} &= \frac{\pi^6}{945}, \\
\sum_{i=1}^{\infty} \frac{1}{(2i-1)^2} &= \frac{\pi^2}{8}, \\
\sum_{i=1}^{\infty} \frac{1}{(2i-1)^4} &= \frac{\pi^4}{96}, \\
\sum_{i=1}^{\infty} \frac{1}{(2i-1)^6} &= \frac{\pi^6}{960}, \\
\sum_{i=1}^{\infty} (-1)^{i+1} \frac{1}{i^2} &= \frac{\pi^2}{12}, \\
\sum_{i=1}^{\infty} (-1)^{i+1} \frac{1}{i^4} &= \frac{7\pi^4}{720}, \\
\sum_{i=1}^{\infty} (-1)^{i+1} \frac{1}{i^6} &= \frac{31\pi^6}{30240}, \\
\sum_{i=1}^{\infty} (-1)^{i+1} \frac{1}{2i-1} &= \frac{\pi}{4}, \\
\sum_{i=1}^{\infty} (-1)^{i+1} \frac{1}{(2i-1)^3} &= \frac{5\pi^5}{1536}, \\
\sum_{i=1}^{\infty} (-1)^{i+1} \frac{1}{(2i-1)^5} &= \frac{61\pi^7}{184320}.
\end{aligned}$$

Fact 1.7.8. For $i = 1, 2, \dots$, let p_i denote the i th prime number, where $p_1 = 2$. Then,

$$\frac{\pi^2}{6} = \prod_{i=1}^{\infty} \frac{1}{1 - p_i^{-2}} \approx 1.6449.$$

(Remark: This identity is the *Euler product formula* for $\zeta(2)$, where ζ is the zeta function.)

Fact 1.7.9. The following statements hold:

$$\sum_{i=1}^{\infty} \frac{1}{\binom{2i}{i}} = \frac{1}{3} + \frac{2\pi}{9\sqrt{3}},$$

$$\sum_{i=1}^{\infty} \frac{i}{\binom{2i}{i}} = \frac{2}{3} + \frac{2\pi}{9\sqrt{3}},$$

$$\sum_{i=1}^{\infty} \frac{i^2}{\binom{2i}{i}} = \frac{4}{3} + \frac{10\pi}{27\sqrt{3}},$$

$$\sum_{i=1}^{\infty} \frac{1}{i \binom{2i}{i}} = \frac{\pi}{3\sqrt{3}},$$

$$\sum_{i=1}^{\infty} \frac{1}{i^2 \binom{2i}{i}} = \frac{\pi^2}{18},$$

$$\sum_{i=1}^{\infty} \frac{2-i}{\binom{2i}{i}} = \frac{2\pi}{9\sqrt{3}},$$

$$\sum_{i=0}^{\infty} \frac{25i-3}{2^{i-1} \binom{3i}{i}} = \pi.$$

(Proof: See [238, pp. 20, 25, 26].)

Fact 1.7.10. The following statements hold:

$$\prod_{i=2}^{\infty} \frac{i^2-1}{i^2+1} = \frac{1}{2} \prod_{i=2}^{\infty} \frac{i^2}{i^2+1} = \frac{\pi}{\sinh \pi} \approx 0.2720,$$

$$\prod_{i=2}^{\infty} \frac{i^2-1}{i^2} = \frac{1}{2},$$

$$\prod_{i=2}^{\infty} \frac{i^3-1}{i^3+1} = \frac{2}{3},$$

$$\prod_{i=2}^{\infty} \frac{i^4-1}{i^4+1} = \frac{\pi \sinh \pi}{\cosh(\sqrt{2}\pi) - \cos(\sqrt{2}\pi)} \approx 0.8480.$$

(Proof: See [238, pp. 4, 5].)

Fact 1.7.11. The following statements hold for all $x \in \mathbb{R}$:

$$\begin{aligned}\sin x &= x \prod_{i=1}^{\infty} \left(1 - \frac{x^2}{i^2 \pi^2}\right), \\ \cos x &= \prod_{i=1}^{\infty} \left(1 - \frac{4x^2}{(2i-1)^2 \pi^2}\right), \\ \sinh x &= x \prod_{i=1}^{\infty} \left(1 + \frac{x^2}{i^2 \pi^2}\right), \\ \cosh x &= \prod_{i=1}^{\infty} \left(1 + \frac{4x^2}{(2i-1)^2 \pi^2}\right), \\ \sin x &= x \prod_{i=1}^{\infty} \cos \frac{x}{2^i}.\end{aligned}$$

1.8 Facts on Convex Functions

Fact 1.8.1. Let $\mathcal{J}$ be a finite or infinite interval, and let $f: \mathcal{J} \mapsto \mathbb{R}$. Then, in each case below, f is convex:

- i) $\mathcal{J} = (0, \infty)$, $f(x) = -\log x$.
 - ii) $\mathcal{J} = (0, \infty)$, $f(x) = x \log x$.
 - iii) $\mathcal{J} = (0, \infty)$, $f(x) = x^p$, where $p < 0$.
 - iv) $\mathcal{J} = [0, \infty)$, $f(x) = -x^p$, where $p \in (0, 1)$.
 - v) $\mathcal{J} = [0, \infty)$, $f(x) = x^p$, where $p \in (1, \infty)$.
 - vi) $\mathcal{J} = [0, \infty)$, $f(x) = (1 + x^p)^{1/p}$, where $p \in (1, \infty)$.
 - vii) $\mathcal{J} = \mathbb{R}$, $f(x) = \frac{a^x - b^x}{c^x - d^x}$, where $0 < d < c < b < a$.
 - viii) $\mathcal{J} = \mathbb{R}$, $f(x) = \log \frac{a^x - b^x}{c^x - d^x}$, where $0 < d < c < b < a$ and $ad \geq bc$.
- (Proof: Statements *vii*) and *viii*) are given in [238, p. 39].)

Fact 1.8.2. Let $\mathcal{J} \subseteq (0, \infty)$ be a finite or infinite interval, let $f: \mathcal{J} \mapsto \mathbb{R}$, and define $g: \mathcal{J} \mapsto \mathbb{R}$ by $g(x) = xf(1/x)$. Then, f is (convex, strictly convex) if and only if g is (convex, strictly convex). (Proof: See [1039, p. 13].)

Fact 1.8.3. Let $f: \mathbb{R} \mapsto \mathbb{R}$, assume that f is convex, and assume that there exists $\alpha \in \mathbb{R}$ such that, for all $x \in \mathbb{R}$, $f(x) \leq \alpha$. Then, f is constant. (Proof: See [1039, p. 35].)

Fact 1.8.4. Let $\mathcal{J} \subseteq \mathbb{R}$ be a finite or infinite interval, let $f: \mathcal{J} \mapsto \mathbb{R}$, and assume that f is continuous. Then, the following statements are equivalent:

- i) f is convex.
- ii) For all $k \in \mathbb{P}$, $x_1, \dots, x_k \in \mathcal{J}$, and $\alpha_1, \dots, \alpha_n \in [0, 1]$ such that $\sum_{i=1}^n \alpha_i = 1$,

it follows that

$$f\left(\sum_{i=1}^n \alpha_i x_i\right) \leq \sum_{i=1}^n \alpha_i f(x_i).$$

(Remark: This result is *Jensen's inequality*.) (Remark: Setting $f(x) = x^p$ yields Fact 1.15.35, whereas setting $f(x) = \log x$ for $x \in (0, \infty)$ yields the arithmetic-mean–geometric-mean inequality given by Fact 1.15.14.) (Remark: See Fact 10.11.7.)

Fact 1.8.5. Let $[a, b] \subset \mathbb{R}$, let $f: [a, b] \mapsto \mathbb{R}$ be convex, and let $x, y \in [a, b]$. Then,

$$\frac{1}{2}[f(x) + f(y)] - f\left[\frac{1}{2}(x + y)\right] \leq \frac{1}{2}[f(a) + f(b)] - f\left[\frac{1}{2}(a + b)\right].$$

(Remark: This result is *Niculescu's inequality*. See [99, p. 13].)

Fact 1.8.6. Let $J \subseteq \mathbb{R}$ be a finite or infinite interval, let $f: J \mapsto \mathbb{R}$. Then, the following statements are equivalent:

- i) f is convex.
- ii) f is continuous, and, for all $x, y \in J$,

$$\frac{2}{3}(f[\tfrac{1}{2}(x+y)] + f[\tfrac{1}{2}(y+z)] + f[\tfrac{1}{2}(x+z)]) \leq \frac{1}{3}[f(x) + f(y) + f(z)] + f[\tfrac{1}{3}(x+y+z)].$$

(Remark: This result is *Popoviciu's inequality*. See [1039, p. 12].) (Remark: For the case of a scalar argument and $f(x) = |x|$, this result implies Hlawka's inequality given by Fact 9.7.4. See Fact 1.18.2 and [1041].) (Problem: Extend this result so that it yields Hlawka's inequality for vector arguments.)

Fact 1.8.7. Let $[a, b] \subset \mathbb{R}$, let $f: [a, b] \mapsto \mathbb{R}$, and assume that f is convex. Then,

$$f\left[\frac{1}{2}(a + b)\right] \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{1}{2}[f(a) + f(b)].$$

(Proof: See [1039, pp. 50–53] and [1156, 1158].) (Remark: This result is the *Hermite-Hadamard inequality*.)

1.9 Facts on Scalar Identities and Inequalities in One Variable

Fact 1.9.1. Let x and α be real numbers, and assume that $x \geq -1$. Then, the following statements hold:

- i) If $\alpha \leq 0$, then

$$1 + \alpha x \leq (1 + x)^\alpha.$$

Furthermore, equality holds if and only if either $x = 0$ or $\alpha = 0$.

- ii) If $\alpha \in [0, 1]$, then

$$(1 + x)^\alpha \leq 1 + \alpha x.$$

Furthermore, equality holds if and only if either $x = 0$, $\alpha = 0$, or $\alpha = 1$.

- iii) If $\alpha \geq 1$, then

$$1 + \alpha x \leq (1 + x)^\alpha.$$

Furthermore, equality holds if and only if either $x = 0$ or $\alpha = 1$.

(Proof: See [34], [274, p. 4], and [1010, p. 65]. Alternatively, the result follows from Fact 1.9.26. See [1447].) (Remark: These results are *Bernoulli's inequality*. An equivalent version is given by Fact 1.9.2.) (Remark: The proof of *i*) and *iii*) in [34] is based on the fact that, for $x \geq -1$, the function $f(x) \triangleq \frac{(1+x)^\alpha - 1}{x}$ for $x \neq 0$ and $f(0) \triangleq \alpha$, is increasing.)

Fact 1.9.2. Let x be a nonnegative number, and let α be a real number. If $\alpha \in [0, 1]$, then

$$\alpha + x^\alpha \leq 1 + \alpha x,$$

whereas, if either $\alpha \leq 0$ or $\alpha \geq 1$, then

$$1 + \alpha x \leq \alpha + x^\alpha.$$

(Proof: Set $y = x + 1$ in Fact 1.9.1. Alternatively, for the case $\alpha \in [0, 1]$, set $y = 1$ in the right-hand inequality in Fact 1.10.21. For the case $\alpha \geq 1$, note that $f(x) \triangleq \alpha + x^\alpha - 1 - \alpha x$ satisfies $f(1) = 0$, $f'(1) = 0$, and, for all $x \geq 0$, $f''(x) = \alpha(\alpha - 1)x^{\alpha-2} > 0$.) (Remark: This result is equivalent to Bernoulli's inequality. See Fact 1.9.1.) (Remark: For $\alpha \in [0, 1]$ a matrix version is given by Fact 8.9.42.) (Problem: Compare the second inequality to Fact 1.10.22 with $y = 1$.)

Fact 1.9.3. Let x and α be real numbers, assume that either $\alpha \leq 0$ or $\alpha \geq 1$, and assume that $x \in [0, 1]$. Then,

$$(1 + x)^\alpha \leq 1 + (2^\alpha - 1)x.$$

Furthermore, equality holds if and only if either $\alpha = 0$, $\alpha = 1$, $x = 0$, or $x = 1$. (Proof: See [34].)

Fact 1.9.4. Let $x \in (0, 1)$, and let k be a positive integer. Then,

$$(1 - x)^k < \frac{1}{1 + kx}.$$

(Proof: See [668, p. 137].)

Fact 1.9.5. Let x be a nonnegative number. Then,

$$\begin{aligned} 8x &< x^4 + 9, \\ 3x^2 &\leq x^3 + 4, \\ 4x^2 &< x^4 + x^3 + x + 1, \\ 8x^2 &< x^4 + x^3 + 4x + 4, \\ 3x^5 &< x^{11} + x^4 + 1. \end{aligned}$$

Now, let n be a positive integer. Then,

$$(2n + 1)x^n \leq \sum_{i=1}^{2n} x^i.$$

(Proof: See [668, pp. 117, 123, 152, 153, 155].)

Fact 1.9.6. Let x be a positive number. Then,

$$1 + \frac{1}{2}x - \frac{1}{8}x^2 < \sqrt{1+x} < 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3.$$

(Proof: See [783, p. 55].)

Fact 1.9.7. Let $x \in (0, 1)$. Then,

$$\frac{1}{2-x} < x^x < x^2 - x + 1.$$

(Proof: See [668, p. 164].)

Fact 1.9.8. Let $x, p \in [1, \infty)$. Then,

$$x^{1/p}(x-1) < px(x^{1/p}-1).$$

Furthermore, equality holds if and only if either $p = 1$ or $x = 1$. (Proof: See [530, p. 194].)

Fact 1.9.9. If $p \in [\sqrt{2}, 2)$, then, for all $x \in (0, 1)$, it follows that

$$\left[\frac{1-x^p}{p(1-x)} \right]^2 \leq \frac{1}{2}(1+x^{p-1}).$$

Furthermore, if $p \in (1, \sqrt{2})$, then there exists $x \in (0, 1)$, such that

$$\frac{1}{2}(1+x^{p-1}) < \left[\frac{1-x^p}{p(1-x)} \right]^2.$$

(Proof: See [206].)

Fact 1.9.10. Let $x, p \in [1, \infty)$. Then,

$$(p-1)^{p-1}(x^p-1)^p \leq p^p(x-1)(x^p-x)^{p-1}x^{p-1}.$$

Furthermore, equality holds if and only if either $p = 1$ or $x = 1$. (Proof: See [530, p. 194].)

Fact 1.9.11. Let $x \in [1, \infty)$, and let $p, q \in (1, \infty)$ satisfy $1/p + 1/q = 1$. Then,

$$px^{1/q} \leq 1 + (p-1)x.$$

Furthermore, equality holds if and only if $x = 1$. (Proof: See [530, p. 194].)

Fact 1.9.12. Let $x \in [1, \infty)$, and let $p, q \in (1, \infty)$ satisfy $1/p + 1/q = 1$. Then,

$$x-1 \leq p^{1/p}q^{1/q}(x^{1/p}-1)^{1/p}(x^{1/q}-1)^{1/q}x^{2/(pq)}.$$

Furthermore, equality holds if and only if $x = 1$. (Proof: See [530, p. 195].)

Fact 1.9.13. Let x be a real number, and let $p, q \in (1, \infty)$ satisfy $1/p + 1/q = 1$. Then,

$$\frac{1}{p}e^{px} + \frac{1}{q}e^{-qx} \leq e^{p^2q^2x^2/8}.$$

(Proof: See [868, p. 260].)

Fact 1.9.14. Let x and y be positive numbers. If $x \in (0, 1]$ and $y \in [0, x]$, then

$$\left(1 + \frac{1}{x}\right)^y \leq 1 + \frac{y}{x}.$$

Equality holds if and only if either $y = 0$ or $x = y = 1$. If $x \in (0, 1)$, then

$$\left(1 + \frac{1}{x}\right)^x < 2.$$

If $x > 1$ and $y \in [1, x]$, then

$$1 + \frac{y}{x} \leq \left(1 + \frac{1}{x}\right)^y < 1 + \frac{y}{x} + \frac{y^2}{x^2}.$$

The left-hand inequality is an equality if and only if $y = 1$. Finally, if $x > 1$, then

$$2 < \left(1 + \frac{1}{x}\right)^x < 3.$$

(Proof: See [668, p. 137].)

Fact 1.9.15. Let x be a nonnegative number, and let p and q be real numbers such that $0 < p \leq q$. Then,

$$e^x \left(1 + \frac{1}{p}\right)^{-x} \leq \left(1 + \frac{x}{p}\right)^p \leq \left(1 + \frac{x}{q}\right)^q \leq e^x.$$

Furthermore, if $p < q$, then equality holds if and only if $x = 0$. Finally,

$$\lim_{q \rightarrow \infty} \left(1 + \frac{x}{q}\right)^q = e^x.$$

(Proof: See [274, pp. 7, 8].) (Remark: For $q \rightarrow \infty$, $(1 + 1/q)^q = e + O(1/q)$, whereas $(1 + 1/q)^q [1 + 1/(2q)] = e + O(1/q^2)$. See [829].)

Fact 1.9.16. Let x be a positive number. Then,

$$\sqrt{\frac{x}{x+1}} e < \left(1 + \frac{1}{x}\right)^x < \frac{2x+1}{2x+2} e$$

and

$$\begin{aligned} \sqrt{1 + \frac{1}{x}} e^{-1/[12x(x+1)]} &< \frac{2x+2}{2x+1} e^{1/[6(2x+1)^2]} \\ &< \frac{e}{\left(1 + \frac{1}{x}\right)^x} \\ &< \sqrt{1 + \frac{1}{x}} e^{-1/[3(2x+1)^2]}. \end{aligned}$$

(Proof: See [1160].)

Fact 1.9.17. Let x be a positive number. Then,

$$\begin{aligned} \left(1 + \frac{1}{x + \frac{1}{5}}\right)^{1/2} &< \left(1 + \frac{2}{3x + 1}\right)^{3/4} \\ &< \left(1 + \frac{1}{\frac{5}{4}x + \frac{1}{3}}\right)^{5/8} \\ &< \frac{e}{\left(1 + \frac{1}{x}\right)^x} \\ &< \left(1 + \frac{1}{x + \frac{1}{6}}\right)^{1/2}. \end{aligned}$$

(Proof: See [921].)

Fact 1.9.18. e is given by

$$\lim_{q \rightarrow \infty} \left(\frac{q+1}{q-1}\right)^{q/2} = e$$

and

$$\lim_{q \rightarrow \infty} \left[\frac{q^q}{(q-1)^{q-1}} - \frac{(q-1)^{q-1}}{(q-2)^{q-2}} \right] = e.$$

(Proof: These expressions are given in [1157] and [829], respectively.)

Fact 1.9.19. Let $n \geq 2$ be a positive integer. Then,

$$e \left(\frac{n}{e}\right)^n < \sqrt{2\pi n} \left(\frac{n}{e}\right)^n < n! < \sqrt{\frac{n}{n-1}} \sqrt{2\pi n} \left(\frac{n}{e}\right)^n < \left(\frac{n+1}{2}\right)^n < \frac{n^{n+1}}{e^{n-1}} < e \left(\frac{n}{2}\right)^n.$$

(Proof: See [1160].) (Remark: The lower bound for $n!$ is *Stirling's formula*.) (Remark: $(e/2)^n < n$ and $\sqrt{2\pi} < e$.)

Fact 1.9.20. Let n be a positive integer. If $n \geq 3$, then

$$n! < 2^{n(n-1)/2}.$$

If $n \geq 6$, then

$$\left(\frac{n}{3}\right)^2 < n! < \left(\frac{n}{2}\right)^2.$$

(Proof: See [668, p. 137].)

Fact 1.9.21. Let x and a be positive numbers. Then,

$$\log x \leq ax - \log a - 1.$$

In particular,

$$\log x \leq \frac{x}{e}.$$

Fact 1.9.22. Let x be a positive number. Then,

$$\frac{x-1}{x} \leq \log x \leq x-1.$$

Furthermore, equality holds if and only if $x = 1$.

Fact 1.9.23. Let x be a positive number such that $x \neq 1$. Then,

$$\frac{1}{x^2 + 1} \leq \frac{\log x}{x^2 - 1} \leq \frac{1}{2x}.$$

Furthermore, equality holds if and only if $x = 1$.

Fact 1.9.24. Let x be a positive number. Then,

$$\frac{2|x-1|}{x+1} \leq |\log x| \leq \frac{|x-1|(1+x^{1/3})}{x+x^{1/3}} \leq \frac{|x-1|}{\sqrt{x}}.$$

Furthermore, equality holds if and only if $x = 1$. (Proof: See [274, p. 8].)

Fact 1.9.25. If $x \in (0, 1]$, then

$$\frac{x-1}{x} \leq \frac{x^2-1}{2x} \leq \frac{x-1}{\sqrt{x}} \leq \frac{(x-1)(1+x^{1/3})}{x+x^{1/3}} \leq \log x \leq \frac{2(x-1)}{x+1} \leq \frac{x^2-1}{x^2+1} \leq x-1.$$

If $x \geq 1$, then

$$\frac{x-1}{x} \leq \frac{x^2-1}{x^2+1} \leq \frac{2(x-1)}{x+1} \leq \log x \leq \frac{(x-1)(1+x^{1/3})}{x+x^{1/3}} \leq \frac{x-1}{\sqrt{x}} \leq \frac{x^2-1}{2x} \leq x-1.$$

Furthermore, equality holds in all cases if and only if $x = 1$. (Proof: See [274, p. 8] and [625].)

Fact 1.9.26. Let x be a positive number, and let p and q be real numbers such that $0 < p \leq q$. Then,

$$\log x \leq \frac{x^p - 1}{p} \leq \frac{x^q - 1}{q} \leq x^q \log x.$$

In particular,

$$\log x \leq 2(\sqrt{x} - 1) \leq x - 1.$$

Furthermore, equality holds in the second inequality if and only if either $p = q$ or $x = 1$. Finally,

$$\lim_{p \downarrow 0} \frac{x^p - 1}{p} = \log x.$$

(Proof: See [34, 1447] and [274, p. 8].) (Remark: See Proposition 8.6.4.) (Remark: See Fact 8.13.1.)

Fact 1.9.27. Let $x > 0$. Then,

$$x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 < \log(1+x) < x - \frac{1}{2}x^2 + \frac{1}{3}x^3.$$

(Proof: See [783, p. 55].)

Fact 1.9.28. Let $x > 1$. Then,

$$\frac{x-1}{\log x} < \left(\frac{x^{1/2} + x^{1/4} + 1}{3} \right)^2 < \left(\frac{x^{1/3} + 1}{2} \right)^3.$$

(Proof: See [756].)

Fact 1.9.29. Let x be a real number. Then, the following statements hold:

i) If $x \in [0, \pi/2]$, then

$$\left. \begin{array}{c} x \cos x \\ \frac{2}{\pi}x \leq \frac{2}{\pi}x + \frac{1}{\pi^3}x(\pi^2 - 4x^2) \\ \frac{x}{\sqrt{(1 - 4/\pi^2)x^2 + 1}} \end{array} \right\} \leq \sin x \leq \left\{ \begin{array}{c} \frac{2}{\pi}x + \frac{\pi-2}{\pi^3}x(\pi^2 - 4x^2) \\ x \leq \tan x \\ 1 \end{array} \right.$$

ii) If $x \in (0, \pi/2]$, then

$$\cot^2 x < \frac{1}{x^2} < 1 + \cot^2 x.$$

iii) If $x \in (0, \pi)$, then

$$\frac{1}{\pi}x(\pi - x) \leq \sin x \leq \frac{4}{\pi^2}x(\pi - x).$$

iv) If $x \in [-4, 4]$, then

$$\cos x \leq \frac{\sin x}{x} \leq 1.$$

v) If $x \in [-\pi/2, \pi/2]$ and $p \in [0, 3]$, then

$$\cos x \leq \left(\frac{\sin x}{x} \right)^p \leq 1.$$

vi) If $x \neq 0$, then

$$x - \frac{1}{6}x^3 < \sin x < x - \frac{1}{6}x^3 + \frac{1}{120}x^5.$$

vii) If $x \neq 0$, then

$$1 - \frac{1}{2}x^2 < \cos x < 1 - \frac{1}{2}x^2 + \frac{1}{24}x^4.$$

viii) If $x \geq \sqrt{3}$, then

$$1 + x \cos \frac{\pi}{x} < (x + 1) \cos \frac{\pi}{x + 1}.$$

ix) If $x \in [0, \pi/2)$,

$$\frac{4x}{\pi - 2x} \leq \pi \tan x.$$

x) If $x \in [0, \pi/2)$, then

$$2 \leq \frac{16}{\pi^4}x^3 \tan x + 2 \leq \left(\frac{\sin x}{x} \right)^2 + \frac{\tan x}{x} \leq \frac{8}{45}x^3 \tan x + 2.$$

xi) If $x \in (0, \pi/2)$, then

$$3x < 2 \sin x + \tan x.$$

xii) For all $x > 0$,

$$3 \sin x < (2 + \cos x)x.$$

xiii) If $x \in [0, \pi/2]$,

$$2 \log \sec x \leq (\sin x) \tan x.$$

xiv) If $x \in (0, 1)$, then

$$\sin^{-1} x < \frac{x}{1 - x^2}.$$

xv) If $x > 0$, then

$$\left. \begin{array}{c} \frac{x}{x^2 + 1} \\ \frac{3x}{1 + 2\sqrt{x^2 + 1}} \end{array} \right\} < \tan^{-1} x.$$

xvi) If $x \in (0, \pi/2)$, then

$$\sinh x < 2 \tan x.$$

xvii) If $x \in \mathbb{R}$, then

$$1 \leq \frac{\sinh x}{x} \leq \cosh x \leq \left(\frac{\sinh x}{x} \right)^3.$$

xviii) If $x > 0$ and $p \geq 3$, then

$$\cosh x < \left(\frac{\sinh x}{x} \right)^p.$$

xix) If $x > 0$, then

$$2 \leq \frac{8}{45}x^3 \tan x + 2 \leq \left(\frac{\sinh x}{x} \right)^2 + \frac{\tanh x}{x}.$$

xx) If $x > 0$, then

$$\frac{\sinh x}{\sqrt{\sinh^2 x + \cosh^2 x}} < \tanh x < x < \sinh x < \frac{1}{2} \sinh 2x.$$

(Proof: Statements *i*), *iv*), *viii*), *ix*), and *xiii*) are given in [273, pp. 250, 251]. For *i*), see also [783, p. 75] and [902]. Statement *ii*) follows from $\sin x < x < \tan x$ in statement *i*). Statement *iii*) is given in [783, p. 72]. Statement *v*) is given in [1500]. Statements *vi*) and *vii*) are given in [783, p. 68]. Statement *x*) is given in [34, 1432]. See also [274, p. 9], [868, pp. 270–271], and [1499, 1500]. Statement *xi*) is *Huygens's inequality*. See [783, p. 71] and [868, p. 266]. Statement *xii*) is given in [783, p. 71] and [868, p. 266]. Statement *xiv*) is given in [868, p. 271]. Statements *xv*) and *xvi*) are given in [783, pp. 70, 75]. Statement *xvii*) is given in [273, pp. 131] and [673, p. 71]. Statements *xviii*) and *xix*) are given in [1500]. Statement *xx*) is given in [783, p. 74].) (Remark: The inequality $2/\pi \leq (\sin x)/x$ is *Jordan's inequality*. See [902].)

Fact 1.9.30. The following statements hold:

i) If $x \in \mathbb{R}$, then

$$\frac{1 - x^2}{1 + x^2} \leq \frac{\sin \pi x}{\pi x}.$$

ii) If $|x| \geq 1$, then

$$\frac{1 - x^2}{1 + x^2} + \frac{(1 - x)^2}{x(1 + x^2)} \leq \frac{\sin \pi x}{\pi x}.$$

iii) If $x \in (0, 1)$, then

$$\frac{(1 - x^2)(4 - x^2)(9 - x^2)}{x^6 - 2x^4 + 13x^2 + 36} \leq \frac{\sin \pi x}{\pi x} \leq \frac{1 - x^2}{\sqrt{1 + 3x^4}}.$$

(Proof: See [902].)

Fact 1.9.31. Let n be a positive integer, and let r be a positive number. Then,

$$\frac{n}{n+1} \leq \left[\frac{(n+1) \sum_{i=1}^n i^r}{n \sum_{i=1}^{n+1} i^r} \right]^{1/r} \leq \frac{\sqrt[n]{n!}}{\sqrt[n+1]{(n+1)!}}.$$

(Proof: See [4].) (Remark: The left-hand inequality is *Alzer's inequality*, while the right-hand inequality is *Martins's inequality*.)

1.10 Facts on Scalar Identities and Inequalities in Two Variables

Fact 1.10.1. Let m and n be positive integers. Then,

$$(m^2 - n^2)^2 + (2mn)^2 = (m^2 + n^2)^2.$$

In particular, if $m = 2$ and $n = 1$, then

$$3^2 + 4^2 = 5^2,$$

while, if $m = 3$ and $n = 2$, then

$$5^2 + 12^2 = 13^2.$$

Furthermore, if $m = 4$ and $n = 1$, then

$$8^2 + 15^2 = 17^2,$$

whereas, if $m = 4$ and $n = 3$, then

$$7^2 + 24^2 = 25^2.$$

(Remark: This result characterizes all *Pythagorean triples* within an integer multiple.)

Fact 1.10.2. The following integer identities hold:

i) $3^3 + 4^3 + 5^3 = 6^3.$

ii) $1^3 + 12^3 = 9^3 + 10^3.$

iii) $10^2 + 11^2 + 12^2 = 13^2 + 14^2.$

iv) $21^2 + 22^2 + 23^2 + 24^2 = 25^2 + 26^2 + 27^2.$

(Remark: The cube of a positive integer cannot be the sum of the cubes of two positive integers. See [477, p. 7].)

Fact 1.10.3. Let $x, y \in \mathbb{R}$. Then,

$$x^2 - y^2 = (x - y)(x + y),$$

$$x^3 - y^3 = (x - y)(x^2 + xy + y^2),$$

$$x^3 + y^3 = (x + y)(x^2 - xy + y^2),$$

$$x^4 - y^4 = (x - y)(x + y)(x^2 + y^2),$$

$$x^4 + x^2y^2 + y^4 = (x^2 + xy + y^2)(x^2 - xy + y^2),$$

$$\begin{aligned}
 x^4 + (x + y)^4 + y^4 &= 2(x^2 + xy + y^2)^2, \\
 x^5 - y^5 &= (x - y)(x^4 + x^3y + x^2y^2 + xy^3 + y^4), \\
 x^5 + y^5 &= (x + y)(x^4 - x^3y + x^2y^2 - xy^3 + y^4), \\
 x^6 - y^6 &= (x - y)(x + y)(x^2 + xy + y^2)(x^2 - xy + y^2).
 \end{aligned}$$

Fact 1.10.4. Let x and y be real numbers. Then,

$$xy \leq \frac{1}{4}(x + y)^2 \leq \frac{1}{2}(x^2 + y^2).$$

If, in addition, x and y are positive, then

$$2 \leq \frac{x}{y} + \frac{y}{x}$$

and

$$\frac{2}{\frac{1}{x} + \frac{1}{y}} \leq \sqrt{xy} \leq \frac{1}{2}(x + y).$$

(Remark: See Fact 8.10.7.)

Fact 1.10.5. Let x and y be positive numbers, and assume that $0 < x < y$. Then,

$$\frac{(x - y)^2}{8y} < \frac{(x - y)^2}{4(x + y)} < \frac{1}{2}(x + y) - \sqrt{xy} < \frac{(x - y)^2}{8x}.$$

(Proof: See [136, p. 231] and [457, p. 183].)

Fact 1.10.6. Let x and y be real numbers, and let $\alpha \in [0, 1]$. Then,

$$\sqrt{\alpha}x + \sqrt{1 - \alpha}y \leq (x^2 + y^2)^{1/2}.$$

Furthermore, equality holds if and only if one of the following conditions holds:

- i) $x = y = 0$.
- ii) $x = 0$, $y > 0$, and $\alpha = 0$.
- iii) $x > 0$, $y = 0$, and $\alpha = 1$.
- iv) $x > 0$, $y > 0$, and $\alpha = \frac{x^2}{x^2 + y^2}$.

Fact 1.10.7. Let α be a real number. Then,

$$0 \leq x^2 + \alpha xy + y^2$$

for all real numbers x, y if and only if $\alpha \in [-2, 2]$.

Fact 1.10.8. Let x and y be nonnegative numbers. Then,

$$\begin{aligned}
 9xy^2 &\leq 3x^3 + 7y^3, \\
 27x^2y &\leq 4(x + y)^3, \\
 6xy^2 &\leq x^3 + y^6 + 8, \\
 x^2y + y^2x &\leq x^3 + y^3, \\
 x^3y + y^3x &\leq x^4 + y^4, \\
 x^4y + y^4x &\leq x^5 + y^5,
 \end{aligned}$$

$$\begin{aligned}
5x^6y^6 &\leq 2x^{15} + 3y^{10}, \\
8(x^3y + y^3x) &\leq (x + y)^4, \\
4x^2y &\leq x^4 + x^3y + y^2 + xy, \\
4x^2y &\leq x^4 + x^3y^2 + y^2 + x, \\
12xy &\leq 4x^2y + 4y^2x + 4x + y, \\
9xy &\leq (x^2 + x + 1)(y^2 + y + 1), \\
6x^2y^2 &\leq x^4 + 2x^3y + 2y^3x + y^4, \\
4(x^2y + y^2x) &\leq 2(x^2 + y^2)^2 + x^2 + y^2, \\
2(x^2y + y^2x + x^2y^2) &\leq 2(x^4 + y^4) + x^2 + y^2.
\end{aligned}$$

(Proof: See Fact 1.15.8, [457, p. 183], [668, pp. 117, 120, 123, 124, 150, 153, 155].)

Fact 1.10.9. Let x and y be real numbers. Then,

$$\begin{aligned}
x^3y + y^3x &\leq x^4 + y^4, \\
4xy(x - y)^2 &\leq (x^2 - y^2)^2, \\
2x + 2xy &\leq x^2y^2 + x^2 + 2, \\
3(x + y + xy) &\leq (x + y + 1)^2.
\end{aligned}$$

(Proof: See [668, p. 117].)

Fact 1.10.10. Let x and y be real numbers. Then,

$$2|(x + y)(1 - xy)| \leq (1 + x^2)(1 + y^2).$$

(Proof: See [457, p. 185].)

Fact 1.10.11. Let x and y be real numbers, and assume that $xy(x + y) \geq 0$. Then,

$$(x^2 + y^2)(x^3 + y^3) \leq (x + y)(x^4 + y^4).$$

(Proof: See [457, p. 183].)

Fact 1.10.12. Let x and y be real numbers. Then,

$$[x^2 + y^2 + (x + y)^2]^2 = 2[x^4 + y^4 + (x + y)^4].$$

Therefore,

$$\frac{1}{2}(x^2 + y^2)^2 \leq x^4 + y^4 + (x + y)^4$$

and

$$x^4 + y^4 \leq \frac{1}{2}[x^2 + y^2 + (x + y)^2]^2.$$

(Remark: This result is *Candido's identity*. See [25].)

Fact 1.10.13. Let x and y be real numbers. Then,

$$54x^2y^2(x + y)^2 \leq [x^2 + y^2 + (x + y)^2]^3.$$

Equivalently,

$$[x^2y^2(x + y)^2]^{1/3} \leq \frac{1}{\sqrt[3]{2}} \frac{1}{3} [x^2 + y^2 + (x + y)^2]^3.$$

(Remark: This result interpolates the arithmetic-mean–geometric-mean inequality due to the factor $1/\sqrt[3]{2}$.) (Remark: This inequality is used in Fact 4.10.1.)

Fact 1.10.14. Let x and y be real numbers, and let $p \in [1, \infty)$. Then,

$$(p-1)(x-y)^2 + [\tfrac{1}{2}(x+y)]^2 \leq [\tfrac{1}{2}(|x|^p + |y|^p)]^{2/p}.$$

(Proof: See [542, p. 148].)

Fact 1.10.15. Let x and y be complex numbers. If $p \in [1, 2]$, then

$$[|x|^2 + (p-1)|y|^2]^{1/2} \leq [\tfrac{1}{2}(|x+y|^p + |x-y|^p)]^{1/p}.$$

If $p \in [2, \infty]$, then

$$[\tfrac{1}{2}(|x+y|^p + |x-y|^p)]^{1/p} \leq [|x|^2 + (p-1)|y|^2]^{1/2}.$$

(Proof: See Fact 9.9.35.)

Fact 1.10.16. Let x and y be real numbers, let p and q be real numbers, and assume that $1 \leq p \leq q$. Then,

$$[\tfrac{1}{2}(|x + \frac{y}{\sqrt{q-1}}|^q + |x - \frac{y}{\sqrt{q-1}}|^q)]^{1/q} \leq [\tfrac{1}{2}(|x + \frac{y}{\sqrt{p-1}}|^p + |x - \frac{y}{\sqrt{p-1}}|^p)]^{1/p}.$$

(Proof: See [542, p. 206].) (Remark: This result is the scalar version of Bonami's inequality. See Fact 9.7.20.)

Fact 1.10.17. Let x and y be positive numbers, and let n be a positive integer. Then,

$$(n+1)(xy^n)^{1/(n+1)} < x + ny.$$

(Proof: See [868, p. 252].)

Fact 1.10.18. Let x and y be positive numbers such that $x < y$, and let n be a positive integer. Then,

$$(n+1)(y-x)x^n < y^{n+1} - x^{n+1} < (n+1)(y-x)y^n.$$

(Proof: See [868, p. 248].)

Fact 1.10.19. Let $[a, b] \subset \mathbb{R}$, and let $x, y \in [a, b]$. Then,

$$|x| + |y| - |x+y| \leq |a| + |b| - |a+b|.$$

(Proof: Use Fact 1.8.5.)

Fact 1.10.20. Let $[a, b] \subset (0, \infty)$, and let $x, y \in [a, b]$. Then,

$$\sqrt{\frac{x}{y}} + \sqrt{\frac{y}{x}} \leq \sqrt{\frac{a}{b}} + \sqrt{\frac{b}{a}}.$$

(Proof: Use Fact 1.8.5.)

Fact 1.10.21. Let x and y be nonnegative numbers, and let $\alpha \in [0, 1]$. Then,

$$[\alpha x^{-1} + (1-\alpha)y^{-1}]^{-1} \leq x^\alpha y^{1-\alpha} \leq \alpha x + (1-\alpha)y.$$

(Remark: The right-hand inequality follows from the concavity of the logarithm function.) (Remark: The left-hand inequality is the scalar *Young inequality*. See Fact 8.10.46, Fact 8.12.26, and Fact 8.12.27.)

Fact 1.10.22. Let x and y be distinct positive numbers, and let $\alpha \in [0, 1]$. Then,

$$\alpha x + (1 - \alpha)y \leq \gamma x^\alpha y^{1-\alpha},$$

where $\gamma > 0$ is defined by

$$\gamma \triangleq \frac{(h-1)h^{1/(h-1)}}{e \log h}$$

and $h \triangleq \max\{y/x, x/y\}$. In particular,

$$\sqrt{xy} \leq \frac{1}{2}(x + y) \leq \gamma \sqrt{xy}.$$

(Remark: This result is the *reverse Young inequality*. See Fact 1.10.21. The case $\alpha = 1/2$ is the *reverse arithmetic-mean-geometric mean inequality*. See Fact 1.15.19.) (Remark: $\gamma = S(1, h)$ is *Specht's ratio*. See Fact 1.15.19 and Fact 11.14.22.) (Remark: This result is due to Tominaga. See [515].)

Fact 1.10.23. Let x and y be positive numbers. Then,

$$1 < x^y + y^x.$$

(Proof: See [457, p. 184] or [783, p. 75].)

Fact 1.10.24. Let x and y be positive numbers. Then,

$$(x + y) \log \left[\frac{1}{2}(x + y) \right] \leq x \log x + y \log y.$$

(Proof: The result follows from the fact that $f(x) = x \log x$ is convex on $(0, \infty)$. See [783, p. 62].)

Fact 1.10.25. Let x be a positive number and let y be a real number. Then,

$$y - \frac{e^{y-1}}{x} \leq \log x.$$

Furthermore, equality holds if $x = y = 1$.

Fact 1.10.26. Let x and y be real numbers, and let $\alpha \in [0, 1]$. Then,

$$[\alpha e^{-x} + (1 - \alpha)e^{-y}]^{-1} \leq e^{\alpha x + (1-\alpha)y} \leq \alpha e^x + (1 - \alpha)e^y.$$

(Proof: Replace x and y by e^x and e^y , respectively, in Fact 1.10.21.) (Remark: The right-hand inequality follows from the convexity of the exponential function.)

Fact 1.10.27. Let x and y be real numbers, and assume that $x \neq y$. Then,

$$e^{(x+y)/2} \leq \frac{e^x - e^y}{x - y} \leq \frac{1}{2}(e^x + e^y).$$

(Proof: See [24].) (Remark: See Fact 1.10.36.)

Fact 1.10.28. Let x and y be real numbers. Then,

$$2 - y - e^{-x-y} \leq 1 + x \leq y + e^{x-y}.$$

Furthermore, equality holds on the left if and only if $x = -y$, and on the right if and only if $x = y$. In particular,

$$2 - e^{-x} \leq 1 + x \leq e^x.$$

Fact 1.10.29. Let x and y be real numbers. Then, the following statements hold:

i) If $0 \leq x \leq y \leq \pi/2$, then

$$\frac{x}{y} \leq \frac{\sin x}{\sin y} \leq \frac{\pi}{2} \left(\frac{x}{y} \right).$$

ii) If either $x, y \in [0, 1]$ or $x, y \in [1, \pi/2]$, then

$$(\tan x) \tan y \leq (\tan 1) \tan xy.$$

iii) If $x, y \in [0, 1]$, then

$$(\sin^{-1} x) \sin^{-1} y \leq \frac{1}{2} \sin^{-1} xy.$$

iv) If $y \in (0, \pi/2]$ and $x \in [0, y]$, then

$$\left(\frac{\sin y}{y} \right) x \leq \sin x \leq \sin \left[y \left(\frac{x}{y} \right)^{y \cot y} \right].$$

v) If $x, y \in [0, \pi]$ are distinct, then

$$\frac{1}{2}(\sin x + \sin y) < \frac{\cos x - \cos y}{y - x} < \sin \left[\frac{1}{2}(x + y) \right].$$

vi) If $0 \leq x < y < \pi/2$, then

$$\frac{1}{\cos^2 x} < \frac{\tan x - \tan y}{x - y} < \frac{1}{\cos^2 y}.$$

vii) If x and y are positive numbers, then

$$(\sinh x) \sinh xy \leq xy \sinh(x + xy).$$

viii) If $0 < y < x < \pi/2$, then

$$\frac{\sin x}{\sin y} < \frac{x}{y} < \frac{\tan x}{\tan y}.$$

(Proof: Statements i)–iii) are given in [273, pp. 250, 251]. Statement iv) is given in [1039, p. 26]. Statement v) is a consequence of the Hermite-Hadamard inequality given by Fact 1.8.6. See [1039, p. 51]. Statement vi) follows from the mean value theorem and monotonicity of the cosine function. See [868, p. 264]. Statement vii) is given in [673, p. 71]. Statement viii) is given in [868, p. 267].) (Remark: $(\sin 0)/0 = (\sinh 0)/0 = 1$.)

Fact 1.10.30. Let x and y be positive numbers. If $p \in [1, \infty)$, then

$$x^p + y^p \leq (x + y)^p.$$

Furthermore, if $p \in [0, 1)$, then

$$(x + y)^p \leq x^p + y^p.$$

(Proof: For the first statement, set $p = 1$ in Fact 1.15.34. For the second statement, set $q = 1$ in Fact 1.15.34.)

Fact 1.10.31. Let x, y, p, q be nonnegative numbers. Then,

$$x^p y^q + x^q y^p \leq x^{p+q} + y^{p+q}.$$

Furthermore, equality holds if and only if either $pq = 0$ or $x = y$. (Proof: See [668, p. 96].)

Fact 1.10.32. Let x and y be nonnegative numbers, and let $p, q \in (1, \infty)$ satisfy $1/p + 1/q = 1$. Then,

$$xy \leq \frac{x^p}{p} + \frac{y^q}{q}.$$

Furthermore, equality holds if and only if $x^p = y^q$. (Proof: See [430, p. 12] or [431, p. 10].) (Remark: This result is *Young's inequality*. An extension is given by Fact 1.15.31. Matrix versions are given by Fact 8.12.12 and Fact 9.14.22.) (Remark: $1/p + 1/q = 1$ is equivalent to $(p - 1)(q - 1) = 1$.)

Fact 1.10.33. Let x and y be positive numbers, and let p and q be real numbers such that $0 \leq p \leq q$. Then,

$$\frac{x^p + y^p}{(xy)^{p/2}} \leq \frac{x^q + y^q}{(xy)^{q/2}}.$$

(Remark: See Fact 8.8.9.)

Fact 1.10.34. Let x and y be positive numbers, and let p and q be nonzero real numbers such that $p \leq q$. Then,

$$\left(\frac{x^p + y^p}{2} \right)^{1/p} \leq \left(\frac{x^q + y^q}{2} \right)^{1/q}.$$

Furthermore, equality holds if and only if either $p = q$ or $x = y$. Finally,

$$\sqrt{xy} = \lim_{p \rightarrow 0} \left(\frac{x^p + y^p}{2} \right)^{1/p}.$$

Hence, if $p < 0 < q$, then

$$\left(\frac{x^p + y^p}{2} \right)^{1/p} \leq \sqrt{xy} \leq \left(\frac{x^q + y^q}{2} \right)^{1/q}$$

where equality holds if and only if $x = y$. (Proof: See [800, pp. 63–65] and [916].) (Remark: This result is a *power mean inequality*. Letting $q = 1$ yields the arithmetic-mean–geometric-mean inequality $\sqrt{xy} \leq \frac{1}{2}(x + y)$.)

Fact 1.10.35. Let x and y be positive numbers, and let p and q be nonzero real numbers such that $p \leq q$. Then,

$$\frac{x^p + y^p}{x^{p-1} + y^{p-1}} \leq \frac{x^q + y^q}{x^{q-1} + y^{q-1}}.$$

Furthermore, equality holds if and only if either $x = y$ or $p = q$. (Proof: See [99, p. 23].) (Remark: The quantity $\frac{x^p + y^p}{x^{p-1} + y^{p-1}}$ is the *Lehmer mean*.)

Fact 1.10.36. Let x and y be positive numbers such that $x < y$, and define

$$G \triangleq \sqrt{xy}, \quad L \triangleq \frac{y-x}{\log y - \log x}, \quad I \triangleq \frac{1}{e} \left(\frac{x^x}{y^y} \right)^{1/(y-x)}, \quad A \triangleq \frac{1}{2}(x+y).$$

Then,

$$x < G < L < I < A < y,$$

$$G < \sqrt{GA} < \sqrt[3]{G^2A} < \sqrt[3]{\frac{1}{4}(G+A)^2G} < L < \left\{ \begin{array}{l} \frac{1}{3}(2G+A) < \frac{1}{3}(G+2A) \\ \sqrt{LA} < \frac{1}{2}(L+A) \end{array} \right\} < I < A,$$

and

$$G + \frac{(x-y)^2(x+3y)(y+3x)}{8(x+y)(x^2+6xy+y^2)} \leq A.$$

Now, let p and q be real numbers such that $1/3 \leq p < 1 < q$. Then,

$$L < \left(\frac{x^p + y^p}{2} \right)^{1/p} < A < \left(\frac{x^q + y^q}{2} \right)^{1/q}.$$

(Proof: See [916, 1155, 1236] and [668, p. 106]. The inequality $L < \frac{1}{3}(2G+A)$ is *Polya's inequality*. See [1039, p. 53]. The inequality $\frac{1}{3}(G+2A) < I$ is due to Sandor. See [99, p. 24].) (Remark: These inequalities refine the arithmetic-mean-geometric-mean inequality Fact 1.15.14.) (Remark: L is the *logarithmic mean*. Note that $L = \int_0^1 x^t y^{1-t} dt$.) (Remark: I is the *identric mean*. See [1236].) (Remark: See Fact 1.15.26.) (Remark: See Fact 1.10.26.)

Fact 1.10.37. Let x and y be positive numbers, and define

$$L \triangleq \frac{y-x}{\log y - \log x}, \quad H_p \triangleq \left(\frac{x^p + (xy)^{p/2} + y^p}{3} \right)^{1/p}, \quad M_p \triangleq \left(\frac{x^p + y^p}{2} \right)^{1/p}.$$

If p, q are positive numbers such that $p < q$, then

$$M_p < M_q$$

and

$$H_p < H_q.$$

Now, let p, q, r be positive numbers such that $0.5283 \approx (\log 3)/(3 \log 2) \leq p \leq 3q/2$ and $1/3 < r < [(\log 2)/\log 3]p \approx 0.6309p$. Then,

$$L < H_{1/2} < M_{1/3} < M_r < H_p < M_q.$$

In particular, if $r \leq (\log 2)/\log 3 \approx 0.6309$ and $q \geq 2/3 \approx 0.6667$, then

$$\left(\frac{x^r + y^r}{2} \right)^{1/r} < \frac{x + \sqrt{xy} + y}{3} < \left(\frac{x^q + y^q}{2} \right)^{1/q}.$$

Finally, if $1/2 \leq p \leq 3q/2$, then

$$\frac{y-x}{\log y - \log x} < \left(\frac{x^p + (xy)^{p/2} + y^p}{3} \right)^{1/p} < \left(\frac{x^q + y^q}{2} \right)^{1/q}.$$

(Proof: See [275, p. 350] and [604, 756].) (Remark: The center term is the *Heron mean*.)

Fact 1.10.38. Let x and y be distinct positive numbers, and let $\alpha \in [0, 1]$. Then,

$$\sqrt{xy} \leq \frac{1}{2}(x^{1-\alpha}y^\alpha + x^\alpha y^{1-\alpha}) \leq \frac{1}{2}(x + y).$$

Furthermore,

$$\frac{1}{2}(x^{1-\alpha}y^\alpha + x^\alpha y^{1-\alpha}) \leq \frac{y - x}{\log y - \log x}$$

if and only if $\alpha \in [\frac{1}{2}(1 - 1/\sqrt{3}), \frac{1}{2}(1 + 1/\sqrt{3})]$, whereas

$$\frac{y - x}{\log y - \log x} \leq \frac{1}{2}(x^{1-\alpha}y^\alpha + x^\alpha y^{1-\alpha})$$

if and only if $\alpha \in [0, \frac{1}{2}(1 - 1/\sqrt{3})] \cup [\frac{1}{2}(1 + 1/\sqrt{3}), 1]$. (Proof: See [437].) (Remark: The first string of inequalities refines the arithmetic-mean–geometric-mean inequality Fact 1.15.14. The center term is the *Heinz mean*. Monotonicity is considered in Fact 1.16.1, while matrix extensions are given by Fact 9.9.49.)

Fact 1.10.39. Let x and y be positive numbers. Then,

$$\left(\frac{x}{y}\right)^y \leq \left(\frac{x+1}{y+1}\right)^{y+1}.$$

Furthermore, equality holds if and only if $x = y$. (Proof: See [868, p. 267].)

Fact 1.10.40. Let x and y be real numbers. If either $0 < x < y < 1$ or $1 < x < y$, then

$$\frac{y^x}{x^y} < \frac{y}{x}$$

and

$$\frac{y^y}{x^x} < \left(\frac{y}{x}\right)^{xy}.$$

If $0 < x < 1 < y$, then both inequalities are reversed. If either $0 < x < 1 < y$ or $0 < x < y < e$, then

$$1 < \left(\frac{y \log x}{x \log y}\right) \left(\frac{y^x - 1}{x^y - 1}\right) < \frac{y^x}{x^y}.$$

If $e < x < y$, then both inequalities are reversed. (Proof: See [1105].)

Fact 1.10.41. Let x and y be real numbers. If $k \geq 1$, then

$$|x - y|^{2k+1} \leq 2^{2k} |x^{2k+1} - y^{2k+1}|.$$

Now, assume that x and y are nonnegative. If $r \geq 1$, then

$$|x - y|^r \leq |x^r - y^r|.$$

(Proof: See [695].) (Remark: Matrix versions of these results are given in [695]. Applications to nonlinear control appear in [1106].) (Problem: Merge these inequalities.)

1.11 Facts on Scalar Identities and Inequalities in Three Variables

Fact 1.11.1. Let x, y, z be real numbers. Then,

$$|x| + |y| + |z| \leq |x + y - z| + |y + z - x| + |z + x - y|$$

and

$$\frac{|x + y|}{1 + |x + y|} \leq \frac{|x|}{1 + |x|} + \frac{|y|}{1 + |y|}.$$

(Proof: See [457, pp. 181, 183].) (Problem: Extend these results to $\mathbb{C}$ and vector arguments.) (Remark: Equality holds in the first result if x, y, z represent the lengths of the sides of a triangle. See Fact 1.11.17.)

Fact 1.11.2. Let x, y, z be real numbers. Then,

$$2[(x - y)(x - z) + (y - z)(y - x) + (z - x)(z - y)] = (x - y)^2 + (y - z)^2 + (z - x)^2.$$

(Proof: See [136, pp. 242, 402].)

Fact 1.11.3. Let x, y, z be real numbers. Then,

$$(x + y)z \leq \frac{1}{2}(x^2 + y^2) + z^2.$$

(Proof: See [136, p. 230].)

Fact 1.11.4. Let x, y, z be real numbers. Then,

$$\left(\frac{1}{2}x + \frac{1}{3}y + \frac{1}{6}z\right)^2 \leq \frac{1}{2}x^2 + \frac{1}{3}y^2 + \frac{1}{6}z^2.$$

(Proof: See [668, p. 129].)

Fact 1.11.5. Let x, y be nonnegative numbers, and let z be a positive number. Then,

$$x + y \leq z^y x + z^{-x} y.$$

(Proof: See [668, p. 163].)

Fact 1.11.6. Let x, y, z be nonnegative numbers. Then,

$$\sqrt[3]{xyz} \leq \frac{1}{3}(\sqrt{xy} + \sqrt{yz} + \sqrt{zx}) \leq \frac{1}{6}(x + y + z) + \frac{1}{2}\sqrt[3]{xyz} \leq \frac{1}{3}(x + y + z).$$

(Proof: The first inequality is given by Fact 1.15.21, while the second inequality is given in [1040].)

Fact 1.11.7. Let x, y, z be nonnegative numbers. Then,

$$\begin{aligned} xy + yz + zx &\leq (\sqrt{xy} + \sqrt{yz} + \sqrt{zx})^2 \\ &\leq 3(xy + yz + zx) \\ &\leq (x + y + z)^2 \\ &\leq 3(x^2 + y^2 + z^2), \end{aligned}$$

$$\begin{aligned}
4(xy + yz) &\leq (x + y + z)^2, \\
2(x + y + z) &\leq x^2 + y^2 + z^2 + 3, \\
2(xy + yz - zx) &\leq x^2 + y^2 + z^2, \\
5xy + 3yz + 7zx &\leq 6x^2 + 4y^2 + 5z^2.
\end{aligned}$$

(Proof: See Fact 1.15.7 and [668, pp. 117, 126].)

Fact 1.11.8. Let x, y, z be nonnegative numbers. Then,

$$\begin{aligned}
12xy + 6xyz &\leq 6x^2 + y^2(z + 2)(2z + 3), \\
(x + y - z)(y + z - x)(z + x - y) &\leq xyz, \\
8xyz &\leq (x + y)(y + z)(z + x), \\
6xyz &\leq x^2y^2 + y^2z^2 + z^2x^2 + x^2 + y^2 + z^2, \\
15xyz &\leq x^3 + y^3 + z^3 + 2(x^2y + y^2z + z^2x + xy^2 + yz^2 + zx^2), \\
15xyz + x^3 + y^3 + z^3 &\leq 2(x + y + z)(x^2 + y^2 + z^2), \\
16xyz &\leq (x + 1)(y + 1)(x + z)(y + z), \\
27xyz &\leq (x^2 + x + 1)(y^2 + y + 1)(z^2 + z + 1), \\
4xyz &\leq x^2y^2z^2 + xy + yz + zx, \\
x^2y + y^2z + z^2x &\leq x^3 + y^3 + z^3, \\
x^2(z + y - x) + y^2(z + x - y) + z^2(x + y - z) \\
&\leq 3xyz \\
&\leq xy^2 + yz^2 + zx^2 \\
&\leq x^3 + y^3 + z^3, \\
27xyz &\leq 3(x + y + z)(xy + yz + zx) \\
&\leq (x + y + z)^3 \\
&\leq 3(x + y + z)(x^2 + y^2 + z^2) \\
&\leq 9(x^3 + y^3 + z^3).
\end{aligned}$$

(Proof: See Fact 1.11.11, [457, pp. 166, 169, 179, 182], [668, pp. 117, 120, 152], and [868, pp. 247, 257].) (Remark: Note the factorization

$$x^3 + y^3 + z^3 - 3xyz = (x + y + z)(x^2 + y^2 + z^2 - xy - yz - zx),$$

where both sides are nonnegative due to the arithmetic-mean-geometric-mean inequality.) (Remark: For positive x, y, z , the inequality $9xyz \leq (x + y + z)(xy + yz + zx)$ is given by Fact 1.15.16.) (Remark: For positive x, y, z , the inequality $3xyz \leq xy^2 + yz^2 + zx^2$ is given by Fact 1.15.17.)

Fact 1.11.9. Let x, y, z be nonnegative numbers. Then,

$$\begin{aligned}
 \left. \begin{array}{l} xyz(x+y+z) \\ 2xyz|x+y-z| \\ 2xyz|x-y+z| \\ 2xyz|-x+y+z| \end{array} \right\} &\leq \left\{ \begin{array}{l} x^2y^2 + y^2z^2 + z^2x^2 \\ 3xyz(x+y+z) \end{array} \right\} \\
 &\leq (xy + yz + zx)^2 \\
 &\leq 3(x^2y^2 + y^2z^2 + z^2x^2) \\
 &\leq (x^2 + y^2 + z^2)^2 \\
 &\leq (x+y+z)(x^3 + y^3 + z^3) \\
 &\leq \left\{ \begin{array}{l} 3(x^4 + y^4 + z^4) \\ (x+y+z)^4 \end{array} \right\} \\
 &\leq 27(x^4 + y^4 + z^4),
 \end{aligned}$$

$$\begin{aligned}
 x^2y^2 + y^2z^2 + z^2x^2 &\leq \frac{1}{2}[x^4 + y^4 + z^4 + xyz(x+y+z)] \\
 &\leq x^4 + y^4 + z^4 \\
 &\leq (x^2 + y^2 + z^2)^2,
 \end{aligned}$$

$$xyz(x+y+z) \leq x^3y + y^3z + z^3x \leq x^4 + y^4 + z^4,$$

$$\begin{aligned}
 \left. \begin{array}{l} 2xyz|x+y-z| \\ 2xyz|x-y+z| \\ 2xyz|-x+y+z| \end{array} \right\} &\leq 3(x^3y + y^3z + z^3x) \leq (x^2 + y^2 + z^2)^2, \\
 (x^2 + y^2 + z^2)(x^3 + y^3 + z^3) &\leq 3(x^5 + y^5 + z^5).
 \end{aligned}$$

Furthermore,

$$\frac{1}{3}(x+y+z) \leq \frac{x^3}{x^2+xy+y^2} + \frac{y^3}{y^2+yz+z^2} + \frac{z^3}{z^2+zx+x^2}.$$

(Proof: See [457, pp. 170, 180], [668, pp. 106, 108, 149], [868, pp. 247, 257], Fact 1.15.2, Fact 1.15.4, and Fact 1.15.22.) (Remark: The inequality $2xyz(x+y-z) \leq x^2y^2 + y^2z^2 + z^2x^2$ follows from $(xy - yz - zx)^2$, and thus is valid for all real x, y, z . See [457, p. 194].) (Remark: The inequality $3xyz(x+y+z) \leq (xy + yz + zx)^2$ follows from Newton's inequality. See Fact 1.15.11.)

Fact 1.11.10. Let x, y, z be nonnegative numbers. Then,

$$9x^2y^2z^2 \leq (x^2y + y^2z + z^2x)(xy^2 + yz^2 + zx^2),$$

$$\begin{aligned}
27x^2y^2z^2 &\leq 3xyz(x+y+z)(xy+yz+zx) \\
&\leq \left\{ \begin{array}{l} xyz(x+y+z)^3 \\ (xy+yz+zx)^3 \end{array} \right\} \\
&\leq \frac{27}{64}(x+y)^2(y+z)^2(z+x)^2 \\
&\leq \frac{9}{64}[(x+y)^6 + (y+z)^6 + (z+x)^6] \\
&\leq \frac{1}{27}(x+y+z)^6 \\
&\leq 9(x^6 + y^6 + z^6), \\
432xy^2z^3 &\leq (x+y+z)^6, \\
3x^2y^2z^2 &\leq \left\{ \begin{array}{l} x^3yz^2 + x^2y^3z + xy^2z^3 \\ xy^3z^2 + x^2yz^3 + x^3y^2z \end{array} \right\} \leq x^2y^4 + y^2z^4 + z^2x^4, \\
9(x^2 + yz)(y^2 + zx)(z^2 + xy) &\leq 8(x^3 + y^3 + z^3)^2, \\
3xyz(x^3 + y^3 + z^3) &\leq (xy + yz + zx)(x^4 + y^4 + z^4), \\
2(x^3y^3 + y^3z^3 + z^3x^3) &\leq x^6 + y^6 + z^6 + 3x^2y^2z^2, \\
xyz(x + y + z)(x^3 + y^3 + z^3) &\leq (xy + yz + zx)(x^5 + y^5 + z^5), \\
(xy + yz + zx)x^2y^2z^2 &\leq x^8 + y^8 + z^8, \\
(xy + yz + zx)^2(xyz^2 + x^2yz + xy^2z) &\leq 3(y^2z^2 + z^2x^2 + x^2y^2)^2, \\
(xyz + 1)^3 &\leq (x^3 + 1)(y^3 + 1)(z^3 + 1).
\end{aligned}$$

Finally, if $\alpha \in [3/7, 7/3]$, then

$$(\alpha + 1)^6(xy + yz + zx)^3 \leq 27(\alpha x + y)^2(\alpha y + z)^2(\alpha z + x)^2.$$

In particular,

$$64(xy + yz + zx)^3 \leq (x + y)^2(y + z)^2(z + x)^2$$

and

$$27(xy + yz + zx)^3 \leq (2x + y)^2(2y + z)^2(2z + x)^2.$$

(Proof: See [136, p. 229], [273, p. 244], [326, p. 114], [457, pp. 179, 182], [668, pp. 105, 134, 150, 155, 169], [868, pp. 247, 252, 257], [1039, p. 14], [1374], Fact 1.11.11, Fact 1.11.21, Fact 1.15.2, Fact 1.15.4, and Fact 1.15.8. For the last inequality, see [63].) (Remark: The inequality $(xy + yz + zx)^2(xyz^2 + x^2yz + xy^2z) \leq 3(y^2z^2 + z^2x^2 + x^2y^2)^2$ is due to Klamkin. See Fact 2.20.11 and [1374].)

Fact 1.11.11. Let x, y, z be positive numbers. Then,

$$6 \leq \frac{9}{2} + \frac{x}{y+z} + \frac{y}{z+x} + \frac{z}{x+y} \leq \frac{x+y}{z} + \frac{y+z}{x} + \frac{z+x}{y}.$$

(Proof: See [99, pp. 33, 34].)

Fact 1.11.12. Let x, y, z be real numbers. Then,

$$2xyz \leq x^2 + y^2z^2$$

and

$$3x^2y^2z^2 \leq x^4y^2 + x^2y^4 + z^6.$$

(Proof: See [668, p. 117] and [153, p. 78].)

Fact 1.11.13. Let x, y, z be positive numbers, and assume that $x < y + z$. Then,

$$\frac{x}{1+x} < \frac{y}{1+y} + \frac{z}{1+z}.$$

(Proof: See [868, p. 44].)

Fact 1.11.14. Let x, y, z be nonnegative numbers. Then,

$$xy(x+y) + yz(y+z) + zx(z+x) \leq x^3 + y^3 + z^3 + 3xyz.$$

(Proof: See [668, p. 98].)

Fact 1.11.15. Let x, y, z be nonnegative numbers, and assume that $x + y < z$. Then,

$$2(x+y)^2z \leq x^3 + y^3 + z^3 + 3xyz.$$

(Proof: See [668, p. 98].)

Fact 1.11.16. Let x, y, z be nonnegative numbers, and assume that $z < x + y$. Then,

$$2(x+y)z^2 \leq x^3 + y^3 + z^3 + 3xyz.$$

(Proof: See [668, p. 100].)

Fact 1.11.17. Let x, y, z be positive numbers. Then, the following statements are equivalent:

- i) x, y, z represent the lengths of the sides of a triangle.
- ii) $z < x + y$, $x < y + z$, and $y < z + x$.
- iii) $(x + y - z)(y + z - x)(z + x - y) > 0$.
- iv) $x > |y - z|$, $y > |z - x|$, and $z > |x - y|$.
- v) $|y - z| < x < y + z$.
- vi) There exist positive numbers a, b, c such that $x = a + b$, $y = b + c$, and $z = c + a$.
- vii) $2(x^4 + y^4 + z^4) < (x^2 + y^2 + z^2)^2$.

In this case, a, b, c in v) are given by

$$a = \frac{1}{2}(z + x - y), \quad b = \frac{1}{2}(x + y - z), \quad c = \frac{1}{2}(y + z - x).$$

(Proof: See [457, p. 164]. Statements v) and vii) are given in [668, p. 125].)
(Remark: See Fact 8.9.5.)

Fact 1.11.18. Let $n \geq 2$, let x, y, z be positive numbers, and assume that $x^n + y^n = z^n$. Then, x, y, z represent the lengths of the sides of a triangle. (Proof: See [668, p. 112].) (Remark: For $n \geq 3$, a lengthy proof shows that the equation $x^n + y^n = z^n$ has no solution in integers.)

Fact 1.11.19. Let x, y, z be positive numbers that represent the lengths of the sides of a triangle. Then, $1/(x+y)$, $1/(y+z)$, and $1/(z+x)$ represent the lengths of the sides of a triangle. (Proof: See [868, p. 44].) (Remark: See Fact 1.11.17 and Fact 1.11.20.)

Fact 1.11.20. Let x, y, z be positive numbers that represent the lengths of the sides of a triangle. Then, $\sqrt{x}$, $\sqrt{y}$, and $\sqrt{z}$, represent the lengths of the sides of a triangle. (Proof: See [668, p. 99].) (Remark: See Fact 1.11.17 and Fact 1.11.19.)

Fact 1.11.21. Let x, y, z be positive numbers that represent the lengths of the sides of a triangle. Then,

$$\begin{aligned}
3(xy + yz + zx) &< (x + y + z)^2 < 4(xy + yz + zx), \\
2(x^2 + y^2 + z^2) &< (x + y + z)^2 < 3(x^2 + y^2 + z^2), \\
\frac{1}{4}(x + y + z)^2 &\leq \left\{ \frac{xy + yz + zx}{\frac{1}{3}(x + y + z)^2} \right\} \leq x^2 + y^2 + z^2 \leq 2(xy + yz + zx), \\
3 &< \frac{2x}{y+z} + \frac{2y}{z+x} + \frac{2z}{x+y} < 4, \\
x(y^2 + z^2) + y(z^2 + x^2) + z(x^2 + y^2) &\leq 3xyz + x^3 + y^3 + z^3, \\
\frac{1}{4}(x + y + z)^3 &\leq (x + y)(y + z)(z + x) \leq \frac{8}{27}(x + y + z)^3, \\
\frac{13}{27}(x + y + z)^3 &\leq (x^2 + y^2 + z^2)(x + y + z) + 4xyz \leq \frac{1}{2}(x + y + z)^3, \\
xyz(x + y + z) &\leq x^2y^2 + y^2z^2 + z^2x^2 \leq x^3y + y^3z + z^3x, \\
xyz &\leq \frac{1}{8}(x + y)(y + z)(z + x).
\end{aligned}$$

If, in addition, the triangle is isosceles, then

$$\begin{aligned}
3(xy + yz + zx) &< (x + y + z)^2 < \frac{16}{5}(xy + yz + zx), \\
\frac{8}{3}(x^2 + y^2 + z^2) &< (x + y + z)^2 < 3(x^2 + y^2 + z^2), \\
\frac{9}{32}(x + y + z)^3 &\leq (x + y)(y + z)(z + x) \leq \frac{8}{27}(x + y + z)^3.
\end{aligned}$$

(Proof: The first string is given in [868, p. 42]. In the second string, the lower bound is given in [457, p. 179], while the upper bound, which holds for all positive x, y, z , is given in Fact 1.11.8. Both the first and second strings are given in [971, p. 199]. In the third string, the upper leftmost inequality follows from Fact 1.11.21; the upper inequality second from the left follows from Fact 1.11.7 whether or not x, y, z represent the lengths of the sides of a triangle; the rightmost inequality is given in [457, p. 179]; the lower leftmost inequality is immediate; and the lower inequality second from the left follows from Fact 1.15.2. The fourth string is given in [868, pp. 267]. The fifth string is given in [457, p. 183]. This result can be

written as [457, p. 186]

$$3 \leq \frac{x}{y+z-x} + \frac{y}{z+x-y} + \frac{z}{x+y-z}.$$

The sixth string is given in [971, p. 199]. The seventh string is given in [1411]. In the eighth string, the left-hand inequality holds for all positive x, y, z . See Fact 1.11.9. The right-hand inequality, which is given in [457, p. 183], orders and interpolates two upper bounds for $xyz(x+y+z)$ given in Fact 1.11.9. The ninth string is given in [971, p. 201]. The inequalities for the case of an obtuse triangle are given in [236] and [971, p. 199].) (Remark: In the fourth string, the lower left inequality is *Nesbitt's inequality*. See [457, p. 163].) (Remark: See Fact 1.11.17 and Fact 2.20.11.)

Fact 1.11.22. Let x, y, z represent the lengths of the sides of a triangle, then

$$\frac{9}{x+y+z} \leq \frac{1}{x} + \frac{1}{y} + \frac{1}{z} \leq \frac{1}{x+y-z} + \frac{1}{x+z-y} + \frac{1}{y+z-x}.$$

(Proof: The lower bound, which holds for all x, y, z , follows from Fact 1.11.21. The upper bound is given in [971, p. 72].) (Remark: The upper bound is *Walker's inequality*.)

Fact 1.11.23. Let x, y, z be positive numbers such that $x+y+z=1$. Then,

$$\frac{25}{1+48xyz} \leq \frac{1}{x} + \frac{1}{y} + \frac{1}{z}.$$

(Proof: See [1469].)

Fact 1.11.24. Let x, y, z be positive numbers that represent the lengths of the sides of a triangle. Then,

$$\left| \frac{x}{y} + \frac{y}{z} + \frac{z}{x} - \left(\frac{y}{x} + \frac{z}{y} + \frac{x}{z} \right) \right| < 1.$$

(Proof: See [457, p. 181].)

Fact 1.11.25. Let x, y, z be positive numbers that represent the lengths of the sides of a triangle. Then,

$$\left| \frac{x-y}{x+y} + \frac{y-z}{y+z} + \frac{z-x}{z+x} \right| < \frac{1}{8}.$$

(Proof: See [457, p. 183].)

Fact 1.11.26. Let x, y, z be real numbers. Then,

$$\frac{|x-z|}{\sqrt{1+x^2}\sqrt{1+z^2}} \leq \frac{|x-y|}{\sqrt{1+x^2}\sqrt{1+y^2}} + \frac{|y-z|}{\sqrt{1+y^2}\sqrt{1+z^2}}.$$

(Proof: See [457, p. 184].)

1.12 Facts on Scalar Identities and Inequalities in Four Variables

Fact 1.12.1. Let w, x, y, z be nonnegative numbers. Then,

$$\sqrt{wx} + \sqrt{yz} \leq \sqrt{(w+y)(x+z)}$$

and

$$6\sqrt[4]{wxyz} \leq \sqrt{(w+x)(y+z)} + \sqrt{(w+y)(x+z)} + \sqrt{(w+z)(x+y)}.$$

(Proof: Use Fact 1.10.4 and see [668, p. 120].)

Fact 1.12.2. Let w, x, y, z be nonnegative numbers. Then,

$$4(wx + xy + yz + zw) \leq (w + x + y + z)^2,$$

$$8(wx + xy + yz + zw + wy + xz) \leq 3(w + x + y + z)^2,$$

$$16(wxy + xyz + yzw + zwx) \leq (w + x + y + z)^3,$$

$$\begin{aligned} 256wxyz &\leq 16(w + x + y + z)(wxy + xyz + yzw + zwx) \\ &\leq (w + x + y + z)^4 \\ &\leq 16(w + x + y + z)(w^3 + x^3 + y^3 + z^3), \end{aligned}$$

$$4wxyz \leq w^2xy + xyz^2 + y^2zw + zw^2x = (wx + yz)(wy + xz),$$

$$4wxyz \leq wx^2z + xy^2w + yz^2x + zw^2y,$$

$$8wxyz \leq (wx + yz)(w + x)(y + z),$$

$$(wx + wy + wz + xy + xz + yz)^2 \leq 6(w^2x^2 + w^2y^2 + w^2z^2 + x^2y^2 + x^2z^2 + y^2z^2),$$

$$4(wxy + xyz + yzw + zwx)^2 \leq (w^2 + x^2 + y^2 + z^2)^3,$$

$$81wxyz \leq (w^2 + w + 1)(x^2 + x + 1)(y^2 + y + 1)(z^2 + z + 1),$$

$$\begin{aligned} w^3x^3y^3 + x^3y^3z^3 + y^3z^3w^3 + z^3w^3x^3 &\leq (wxy + xyz + yzw + zwx)^3 \\ &\leq 16(w^3x^3y^3 + x^3y^3z^3 + y^3z^3w^3 + z^3w^3x^3), \end{aligned}$$

$$\frac{16}{3(w + x + y + z)} \leq \frac{1}{w + x + y} + \frac{1}{x + y + z} + \frac{1}{y + z + w} + \frac{1}{z + w + x}.$$

(Proof: See [457, p. 179], [668, pp. 120, 123, 124, 134, 144, 161], [797], Fact 1.15.22, and Fact 1.15.20.) (Remark: The inequality $(w + x + y + z)^3 \leq 16(w^3 + x^3 + y^3 + z^3)$ is given by Fact 1.15.2.) (Remark: The inequality $16wxyz \leq (w + x + y + z)(wxy + xyz + yzw + zwx)$ is given by Fact 1.15.16.) (Remark: The inequality $4wxyz \leq w^2xy + xyz^2 + y^2zw + zw^2x$ follows from Fact 1.15.17 with $n = 2$.) (Remark: The inequality $4wxyz \leq wx^2z + xy^2w + yz^2x + zw^2y$ is given by Fact 1.15.17.)

Fact 1.12.3. Let w, x, y, z be real numbers. Then,

$$4wxyz \leq w^2x^2 + x^2y^2 + y^2w^2 + z^4$$

and

$$(wxyz + 1)^3 \leq (w^3 + 1)(x^3 + 1)(y^3 + 1)(z^3 + 1).$$

(Proof: See [153, p. 78] and [668, p. 134].)

Fact 1.12.4. Let w, x, y, z be real numbers. Then,

$$\begin{aligned} (w^2 + x^2)(y^2 + z^2) &= (wz + xy)^2 + (wy - xz)^2 \\ &= (wz - xy)^2 + (wy + xz)^2. \end{aligned}$$

Hence,

$$\left. \begin{aligned} (wz + xy)^2 \\ (wy - xz)^2 \\ (wz - xy)^2 \\ (wy + xz)^2 \end{aligned} \right\} \leq (w^2 + x^2)(y^2 + z^2).$$

(Remark: The identity is a statement of the fact that, for complex numbers z_1, z_2 , $|z_1|^2|z_2|^2 = |z_1z_2|^2 = |\operatorname{Re}(z_1z_2)|^2 + |\operatorname{Im}(z_1z_2)|^2$. See [346, p. 77].)

Fact 1.12.5. Let w, x, y, z be real numbers. Then,

$$w^4 + x^4 + y^4 + z^4 - 4wxyz = (w^2 - x^2)^2 + (y^2 + z^2)^2 + 2(wx - yz)^2.$$

(Remark: This result yields the arithmetic-mean–geometric-mean inequality for four variables. See [136, pp. 226, 367].)

1.13 Facts on Scalar Identities and Inequalities in Six Variables

Fact 1.13.1. Let x, y, z, u, v, w be real numbers. Then,

$$\begin{aligned} &x^6 + y^6 + z^6 + u^6 + v^6 + w^6 - 6xyzuvw \\ &= \frac{1}{2}(x^2 + y^2 + z^2)^2[(x^2 - y^2)^2 + (y^2 - z^2)^2 + (z^2 - x^2)^2] \\ &\quad + \frac{1}{2}(u^2 + v^2 + w^2)^2[(u^2 - v^2)^2 + (v^2 - w^2)^2 + (w^2 - u^2)^2] \\ &\quad + 3(xyz - uvw)^2. \end{aligned}$$

(Remark: This result yields the arithmetic-mean–geometric-mean inequality for six variables. See [136, p. 226].)

1.14 Facts on Scalar Identities and Inequalities in Eight Variables

Fact 1.14.1. Let $x_1, x_2, x_3, x_4, y_1, y_2, y_3, y_4$ be real numbers. Then,

$$\begin{aligned} &(x_1^2 + x_2^2 + x_3^2 + x_4^2)(y_1^2 + y_2^2 + y_3^2 + y_4^2) \\ &= (x_1y_1 - x_2y_2 - x_3y_3 - x_4y_4)^2 + (x_1y_2 + x_2y_1 + x_3y_4 - x_4y_3)^2 \\ &\quad + (x_1y_3 - x_2y_4 + x_3y_1 + x_4y_2)^2 + (x_1y_4 + x_2y_3 - x_3y_2 + x_4y_1)^2. \end{aligned}$$

Hence,

$$\left. \begin{aligned} & (x_1y_1 - x_2y_2 - x_3y_3 - x_4y_4)^2 + (x_1y_2 + x_2y_1 + x_3y_4 - x_4y_3)^2 \\ & \quad + (x_1y_3 - x_2y_4 + x_3y_1 + x_4y_2)^2 \\ & (x_1y_1 - x_2y_2 - x_3y_3 - x_4y_4)^2 + (x_1y_2 + x_2y_1 + x_3y_4 - x_4y_3)^2 \\ & \quad + (x_1y_4 + x_2y_3 - x_3y_2 + x_4y_1)^2 \\ & (x_1y_1 - x_2y_2 - x_3y_3 - x_4y_4)^2 + (x_1y_3 - x_2y_4 + x_3y_1 + x_4y_2)^2 \\ & \quad + (x_1y_4 + x_2y_3 - x_3y_2 + x_4y_1)^2 \\ & (x_1y_2 + x_2y_1 + x_3y_4 - x_4y_3)^2 + (x_1y_3 - x_2y_4 + x_3y_1 + x_4y_2)^2 \\ & \quad + (x_1y_4 + x_2y_3 - x_3y_2 + x_4y_1)^2 \end{aligned} \right\} \\ \leq (x_1^2 + x_2^2 + x_3^2 + x_4^2)(y_1^2 + y_2^2 + y_3^2 + y_4^2).$$

(Remark: The identity represents a relationship between a pair of quaternions. An analogous identity holds for two sets of eight variables representing a pair of octonions. See [346, p. 77].)

1.15 Facts on Scalar Identities and Inequalities in n Variables

Fact 1.15.1. Let $x_1, \dots, x_n$ be real numbers, and let k be a positive integer. Then,

$$\left(\sum_{i=1}^n x_i \right)^k = \sum_{i_1 + \dots + i_n = k} \frac{k!}{i_1! \dots i_n!} x_1^{i_1} \dots x_n^{i_n}.$$

(Remark: This result is the *multinomial theorem*.)

Fact 1.15.2. Let $x_1, \dots, x_n$ be nonnegative numbers, and let k be a positive integer. Then,

$$\sum_{i=1}^n x_i^k \leq \left(\sum_{i=1}^n x_i \right)^k \leq n^{k-1} \sum_{i=1}^n x_i^k.$$

Furthermore, equality holds in the second inequality if and only if $x_1 = \dots = x_n$.

(Remark: The case $n = 4$, $k = 3$ is given by the inequality $(w + x + y + z)^3 \leq 16(w^3 + x^3 + y^3 + z^3)$ of Fact 1.12.2.)

Fact 1.15.3. Let $x_1, \dots, x_n$ be nonnegative numbers. Then,

$$\left(\sum_{i=1}^n x_i \right)^2 \leq n \sum_{i=1}^n x_i^2.$$

Furthermore, equality holds if and only if $x_1 = \dots = x_n$. (Remark: This result is equivalent to i) of Fact 9.8.12 with $m = 1$.)

Fact 1.15.4. Let $x_1, \dots, x_n$ be nonnegative numbers, and let k be a positive integer. Then,

$$\sum_{i=1}^n x_i^k \leq \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n x_i^{k-1} \right) \leq n \sum_{i=1}^n x_i^k.$$

(Proof: See [868, pp. 257, 258].)

Fact 1.15.5. Let $x_1, \dots, x_n$ be nonnegative numbers, and let $p, q \in [1, \infty)$, where $p \leq q$. Then,

$$\left(\sum_{i=1}^n x_i^q \right)^{1/q} \leq \left(\sum_{i=1}^n x_i^p \right)^{1/p} \leq n^{1/p-1/q} \left(\sum_{i=1}^n x_i^q \right)^{1/q}.$$

Equivalently,

$$\sum_{i=1}^n x_i^q \leq \left(\sum_{i=1}^n x_i^p \right)^{q/p} \leq n^{q/p-1} \sum_{i=1}^n x_i^q.$$

(Proof: See Fact 9.7.29.) (Remark: Setting $p = 1$ and $q = k$ yields Fact 1.15.2.)

Fact 1.15.6. Let $x_1, \dots, x_n$ be nonnegative numbers. Then,

$$\left(\sum_{i=1}^n x_i^3 \right)^2 \leq \left(\sum_{i=1}^n x_i^2 \right)^3 \leq n \left(\sum_{i=1}^n x_i^3 \right)^2.$$

(Proof: Set $p = 2$ and $q = 3$ in Fact 1.15.5 and square all terms.)

Fact 1.15.7. Let $x_1, \dots, x_n$ be nonnegative numbers. For $n = 2$,

$$2(x_1x_2 + x_2x_1) \leq (x_1 + x_2)^2.$$

For $n = 3$,

$$3(x_1x_2 + x_2x_3 + x_3x_1) \leq (x_1 + x_2 + x_3)^2.$$

If $n \geq 4$, then

$$4(x_1x_2 + x_2x_3 + \dots + x_nx_1) \leq \left(\sum_{i=1}^n x_i \right)^2.$$

(Proof: See [668, p. 144]. The cases $n = 2, 3, 4$ are given by Fact 1.10.4, Fact 1.11.7, and Fact 1.12.2.) (Problem: Is 4 the best constant for $n \geq 5$?)

Fact 1.15.8. Let $x_1, \dots, x_n$ be nonnegative numbers. Then,

$$\left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n x_i^3 \right) \leq \left(\sum_{i=1}^n x_i^5 \right) \left(\sum_{i=1}^n \frac{1}{x_i} \right).$$

(Proof: See [668, p. 150].)

Fact 1.15.9. Let $x_1, \dots, x_n$ be positive numbers, and assume that, for all $i = 1, \dots, n-1$, $x_i < x_{i+1} \leq x_i + 1$. Then,

$$\sum_{i=1}^n x_i^3 \leq \left(\sum_{i=1}^n x_i \right)^2.$$

(Proof: See [457, p. 183].) (Remark: Equality holds in Fact 1.7.3.)

Fact 1.15.10. Let $x_1, \dots, x_n$ be complex numbers, define $E_0 \triangleq 1$, and, for $1 \leq k \leq n$, define

$$E_k \triangleq \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j}.$$

Furthermore, for each positive integer k define

$$\mu_k \triangleq \sum_{i=1}^n x_i^k.$$

Then, for all $k = 1, \dots, n$,

$$kE_k = \sum_{i=1}^k (-1)^{i-1} E_{k-i} \mu_i.$$

In particular,

$$\begin{aligned} E_1 &= \mu_1, \\ 2E_2 &= E_1\mu_1 - \mu_2, \\ 3E_3 &= E_2\mu_2 - E_1\mu_2 + \mu_3. \end{aligned}$$

Furthermore,

$$\begin{aligned} E_1 &= \mu_1, \\ E_2 &= \frac{1}{2}(\mu_1^2 - \mu_2), \\ E_3 &= \frac{1}{6}(\mu_1^3 - 3\mu_1\mu_2 + 2\mu_3) \end{aligned}$$

and

$$\begin{aligned} \mu_1 &= E_1, \\ \mu_2 &= E_1^2 - 2E_2, \\ \mu_3 &= E_1^3 - 3E_1E_2 + 3E_3. \end{aligned}$$

(Remark: This result is *Newton's identity*. An application to roots of polynomials is given by Fact 4.8.2.) (Remark: E_k is the k th *elementary symmetric polynomial*.) (Remark: See Fact 1.15.11.)

Fact 1.15.11. Let $x_1, \dots, x_n$ be complex numbers, let k be a positive integer such that $1 < k < n$, and define

$$S_k \triangleq \binom{n}{k}^{-1} \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j}.$$

Then,

$$S_{k-1}S_{k+1} \leq S_k^2.$$

(Remark: This result is *Newton's inequality*. The case $n = 3, k = 2$ is given by Fact 1.11.9.) (Remark: S_k is the k th *elementary symmetric mean*.) (Remark: See Fact 1.15.10.)

Fact 1.15.12. Let $x_1, \dots, x_n$ be real numbers, and define

$$\bar{x} \triangleq \frac{1}{n} \sum_{j=1}^n x_j$$

and

$$\sigma \triangleq \sqrt{\frac{1}{n} \sum_{j=1}^n (x_j - \bar{x})^2} = \sqrt{\left(\frac{1}{n} \sum_{j=1}^n x_j^2 \right) - \bar{x}^2}.$$

Then, for all $i = 1, \dots, n$,

$$|x_i - \bar{x}| \leq \sqrt{n-1} \sigma.$$

Equality holds if and only if all of the elements of $\{x_1, \dots, x_n\}_{\text{ms}} \setminus \{x_i\}$ are equal. In addition,

$$\frac{\sigma}{\sqrt{n-1}} \leq \max\{x_1, \dots, x_n\} - \bar{x} \leq \sqrt{n-1} \sigma.$$

Equality holds in either the left-hand inequality or the right-hand inequality if and only if all of the elements of $\{x_1, \dots, x_n\}_{\text{ms}} \setminus \max\{x_1, \dots, x_n\}$ are equal. Finally,

$$\frac{\sigma}{\sqrt{n-1}} \leq \bar{x} - \min\{x_1, \dots, x_n\} \leq \sqrt{n-1} \sigma.$$

Equality holds in either the left-hand inequality or the right-hand inequality if and only if all of the elements of $\{x_1, \dots, x_n\}_{\text{ms}} \setminus \min\{x_1, \dots, x_n\}$ are equal. (Proof: The first result is the *Laguerre-Samuelson inequality*. See [574, 732, 754, 1043, 1140, 1332]. The lower bounds in the second and third strings are given in [1448]. See also [1140].) (Remark: A vector extension of the Laguerre-Samuelson inequality is given by Fact 8.9.35. An application to eigenvalue bounds is given by Fact 5.11.45.)

Fact 1.15.13. Let $x_1, \dots, x_n$ be real numbers, and let α, δ , and p be positive numbers. If $p \geq 1$, then

$$\left(\frac{\alpha}{\alpha + n} \right)^{p-1} \delta^p \leq \left| \delta - \sum_{i=1}^n x_i \right|^p + \alpha^{p-1} \sum_{i=1}^n |x_i|^p.$$

In particular,

$$\frac{\alpha \delta^2}{\alpha + n} \leq \left(\delta - \sum_{i=1}^n x_i \right)^2 + \alpha \sum_{i=1}^n x_i^2.$$

Furthermore, if $p \leq 1$, $x_1, \dots, x_n$ are nonnegative, and $\sum_{i=1}^n x_i \leq \delta$, then

$$\left| \delta - \sum_{i=1}^n x_i \right|^p + \alpha^{p-1} \sum_{i=1}^n |x_i|^p \leq \left(\frac{\alpha}{\alpha + n} \right)^{p-1} \delta^p.$$

Finally, equality holds in all cases if and only if $x_1 = \dots = x_n = \delta/(\alpha + n)$. (Proof: See [1253].) (Remark: This result is *Wang's inequality*. The special case $p = 2$ is *Hua's inequality*. Generalizations are given by Fact 9.7.8 and Fact 9.7.9.)

Fact 1.15.14. Let $x_1, \dots, x_n$ be nonnegative numbers. Then,

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \frac{1}{n} \sum_{i=1}^n x_i.$$

Furthermore, equality holds if and only if $x_1 = x_2 = \cdots = x_n$. (Remark: This result is the *arithmetic-mean-geometric-mean inequality*. Several proofs are given in [275]. See also [314]. Bounds for the difference between these quantities are given in [28, 295, 1343].)

Fact 1.15.15. Let $x_1, \dots, x_n$ be positive numbers. Then,

$$\frac{n}{\frac{1}{x_1} + \cdots + \frac{1}{x_n}} \leq \sqrt[n]{x_1 \cdots x_n} \leq \frac{1}{n}(x_1 + \cdots + x_n) \leq \sqrt{\frac{1}{n} \sum_{i=1}^n x_i^2}.$$

Furthermore, equality holds in each inequality if and only if $x_1 = x_2 = \cdots = x_n$. (Remark: The lower bound for the geometric mean is the *harmonic mean*, while the left-hand inequality is the *arithmetic-mean-harmonic-mean inequality*. See Fact 1.15.37.) (Remark: The upper bound for the arithmetic mean is the *quadratic mean*. See [612] and Fact 1.15.32.)

Fact 1.15.16. Let $x_1, \dots, x_n$ be positive numbers. Then,

$$\frac{n^2}{x_1 + \cdots + x_n} \leq \frac{1}{x_1} + \cdots + \frac{1}{x_n}.$$

(Proof: Use Fact 1.15.15. See also [668, p. 130].) (Remark: The case $n = 3$ yields the inequality $9xyz \leq (x+y+z)(xy+yz+zx)$ of Fact 1.11.8.) (Remark: The case $n = 4$ yields the inequality $16wxyz \leq (w+x+y+z)(wxy+xyz+yzw+zwz)$ of Fact 1.12.2.)

Fact 1.15.17. Let $x_1, \dots, x_n$ be positive numbers. Then,

$$n \leq \frac{x_1}{x_2} + \frac{x_2}{x_3} + \cdots + \frac{x_{n-1}}{x_n} + \frac{x_n}{x_1}.$$

(Remark: The case $n = 3$ yields the inequality $3xyz \leq xy^2 + yz^2 + zx^2$ of Fact 1.11.8.) (Remark: The case $n = 4$ yields the inequality $4wxyz \leq wx^2z + xy^2w + yz^2x + zw^2y$ of Fact 1.12.2.)

Fact 1.15.18. Let $x_1, \dots, x_n$ be nonnegative numbers. Then,

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \frac{1}{n} \sum_{i=1}^n x_i \leq \left(\prod_{i=1}^n x_i \right)^{1/n} + \frac{1}{n} \sum_{i < j} |x_i - x_j|.$$

(Proof: See [457, p. 186].)

Fact 1.15.19. Let $x_1, \dots, x_n$ be positive numbers contained in $[a, b]$, where $a > 0$. Then,

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \frac{1}{n} \sum_{i=1}^n x_i \leq \gamma \left(\prod_{i=1}^n x_i \right)^{1/n},$$

where γ is defined by

$$\gamma \triangleq \frac{(h-1)h^{1/(h-1)}}{e \log h}$$

and $h \triangleq b/a$. (Remark: The right-hand inequality is a *reverse arithmetic-mean-*

geometric mean inequality; see [511, 516, 1470]. This result is due to Specht. For the case $n = 2$, see Fact 1.10.22.) (Remark: $\gamma = S(1, h)$ is Specht's ratio. See Fact 1.10.22 and Fact 11.14.22.) (Remark: Matrix extensions are considered in [19, 809].)

Fact 1.15.20. Let $x_1, \dots, x_n$ be positive numbers, and let k satisfy $1 \leq k \leq n$. Then,

$$\left(\binom{n}{k}^{-1} \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j} \right)^{1/k} \leq \frac{1}{n} \sum_{i=1}^n x_i.$$

Equivalently,

$$\sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j} \leq \binom{n}{k} \left(\frac{1}{n} \sum_{i=1}^n x_i \right)^k.$$

(Proof: The result follows from the fact that the k th elementary symmetric function is Schur concave. See [542, p. 102, Exercise 7.11].) (Remark: Equality holds if $k = 1$. The case $n = k$ is the arithmetic-mean–geometric-mean inequality. The case $n = 3, k = 2$ yields the third inequality in Fact 1.11.7. The cases $n = 4, k = 3$ and $n = 4, k = 2$ are given in Fact 1.12.2.)

Fact 1.15.21. Let $x_1, \dots, x_n$ be positive numbers, and let k and k' satisfy $1 \leq k \leq k' \leq n$. Then,

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \binom{n}{k'}^{-1} \sum_{i_1 < \dots < i_{k'}} \prod_{j=1}^{k'} x_{i_j}^{1/k'} \leq \binom{n}{k}^{-1} \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j}^{1/k} \leq \frac{1}{n} \sum_{i=1}^n x_i.$$

(Proof: See [542, p. 23] and [797].) (Remark: This result is an interpolation of the arithmetic-mean–geometric-mean inequality. An alternative interpolation is given by Fact 1.15.25.) (Remark: If $k = 1$, then the right-hand inequality is an equality. If $k = n$, then the left-hand inequality is an equality. The case $n = 3$ and $k = 2$ is given by Fact 1.11.6.)

Fact 1.15.22. Let $x_1, \dots, x_n$ be nonnegative numbers, and let k be a positive integer such that $1 \leq k \leq n$. Then,

$$\left(\sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j} \right)^k \leq \binom{n}{k}^{k-1} \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j}^k.$$

(Remark: Equality holds if $k = 1$ or $k = n$. The case $n = 3, k = 2$ is given by Fact 1.11.9. The cases $n = 4, k = 3$ and $n = 4, k = 2$ are given by Fact 1.12.2.)

Fact 1.15.23. Let $x_1, \dots, x_n$ be positive numbers, and let k satisfy $1 \leq k \leq n$. Then,

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \binom{n}{k}^{-1} \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j}^{1/k} \leq \left(\binom{n}{k}^{-1} \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j} \right)^{1/k} \leq \frac{1}{n} \sum_{i=1}^n x_i.$$

(Proof: Use Fact 1.15.22 to merge Fact 1.15.20 and Fact 1.15.21.)

Fact 1.15.24. Let $x_1, \dots, x_n$ be positive numbers, and let k and k' satisfy $1 \leq k \leq k' \leq n$. Then,

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \left(\binom{n}{k'}^{-1} \sum_{i_1 < \dots < i_{k'}} \prod_{j=1}^{k'} x_{i_j} \right)^{1/k'} \leq \left(\binom{n}{k}^{-1} \sum_{i_1 < \dots < i_k} \prod_{j=1}^k x_{i_j} \right)^{1/k} \leq \frac{1}{n} \sum_{i=1}^n x_i.$$

(Proof: See [797].)

Fact 1.15.25. Let $x_1, \dots, x_n$ be positive numbers, let $\alpha_1, \dots, \alpha_n$ be nonnegative numbers, and assume that $\sum_{i=1}^n \alpha_i = 1$. Then,

$$\left(\prod_{i=1}^n x_i \right)^{1/n} \leq \frac{1}{n!} \sum \prod_{j=1}^n x_{i_j}^{\alpha_j} \leq \frac{1}{n} \sum_{i=1}^n x_i,$$

where the summation is taken over all $n!$ permutations $\{i_1, \dots, i_n\}$ of $\{1, \dots, n\}$. (Proof: See [542, p. 100].) (Remark: This result is a consequence of *Muirhead's theorem*, which states that the middle expression is a Schur convex function of the exponents. See Fact 2.21.5.)

Fact 1.15.26. Let $x_1, \dots, x_n$ be positive numbers. Then,

$$\left(\prod_{i=1}^n x_i \right)^{1/n} < \frac{1}{n} \left(\frac{x_2 - x_1}{\log x_2 - \log x_1} + \frac{x_3 - x_2}{\log x_3 - \log x_2} + \dots + \frac{x_1 - x_n}{\log x_1 - \log x_n} \right) < \frac{1}{n} \sum_{i=1}^n x_i.$$

(Proof: See [99, p. 44].) (Remark: This result is due to Bencze.) (Remark: This result extends Fact 1.10.36 to n variables. See also [1465].)

Fact 1.15.27. Let $x_1, \dots, x_n$ be positive numbers contained in $[a, b]$, where $a > 0$. Then,

$$\frac{a}{2n^2} \sum_{i < j} (\log x_i - \log x_j)^2 \leq \frac{1}{n} \sum_{i=1}^n x_i - \left(\prod_{i=1}^n x_i \right)^{1/n} \leq \frac{b}{2n^2} \sum_{i < j} (\log x_i - \log x_j)^2.$$

(Proof: See [1039, p. 86] or [1040].)

Fact 1.15.28. Let $x_1, \dots, x_n$ be nonnegative numbers contained in $(0, 1/2]$. Furthermore, define

$$A \triangleq \frac{1}{n} \sum_{i=1}^n x_i, \quad G \triangleq \prod_{i=1}^n x_i^{1/n}, \quad H \triangleq \frac{n}{\sum_{i=1}^n \frac{1}{x_i}}$$

and

$$A' \triangleq \frac{1}{n} \sum_{i=1}^n (1 - x_i), \quad G' \triangleq \prod_{i=1}^n (1 - x_i)^{1/n}, \quad H' \triangleq \frac{n}{\sum_{i=1}^n \frac{1}{1 - x_i}}.$$

Then, the following statements hold:

- i) $A'/G' \leq A/G$. Furthermore, equality holds if and only if $x_1 = \dots = x_n$.

ii) $A' - G' \leq A - G$. Furthermore, equality holds if and only if $x_1 = \cdots = x_n$.

iii) $A^n - G^n \leq A'^n - G'^n$. Furthermore, equality holds for $n = 1$ and $n = 2$, and, for $n \geq 3$, if and only if $x_1 = \cdots = x_n$.

iv) $G'/H' \leq G/H$.

(Proof: See [1141]. For a proof of *iv*), see [1159].) (Remark: Result *i*) is due to Fan. See [1159].)

Fact 1.15.29. Let $x_1, \dots, x_n$ be positive numbers, and, for all $k = 1, \dots, n$, define

$$A_k \triangleq \frac{1}{k} \sum_{i=1}^k x_i, \quad G_k \triangleq \prod_{i=1}^k x_i^{1/k}.$$

Then,

$$1 = \left(\frac{A_1}{G_1} \right)^1 \leq \left(\frac{A_2}{G_2} \right)^2 \leq \cdots \leq \left(\frac{A_n}{G_n} \right)^n$$

and

$$0 = 1(A_1 - G_1) \leq 2(A_2 - G_2) \leq \cdots \leq n(A_n - G_n).$$

(Proof: See [1039, p. 13].) (Remark: The first result is due to Popoviciu, while the second result is due to Rado.)

Fact 1.15.30. Let $x_1, \dots, x_n$ be positive numbers, let p be a real number, and define

$$M_p \triangleq \begin{cases} \left(\prod_{i=1}^n x_i \right)^{1/n}, & p = 0, \\ \left(\frac{1}{n} \sum_{i=1}^n x_i^p \right)^{1/p}, & p \neq 0. \end{cases}$$

Now, let p and q be real numbers such that $p \leq q$. Then,

$$M_p \leq M_q$$

and

$$\lim_{r \rightarrow -\infty} M_r = \min\{x_1, \dots, x_n\} \leq \lim_{r \rightarrow 0} M_r = M_0 \leq \lim_{r \rightarrow \infty} M_r = \max\{x_1, \dots, x_n\}.$$

Finally, $p < q$ and at least two of the numbers $x_1, \dots, x_n$ are distinct if and only if

$$M_p < M_q.$$

(Proof: See [273, p. 210] and [963, p. 105].) If p and q are nonzero and $p \leq q$, then

$$\left(\sum_{i=1}^n x_i^p \right)^{1/p} \leq \left(\frac{1}{n} \right)^{1/q-1/p} \left(\sum_{i=1}^n x_i^q \right)^{1/q},$$

which is a reverse form of Fact 1.15.34. (Proof: To verify the limit, take the log of both sides and use l'Hôpital's rule.) (Remark: This result is a *power mean inequality*. $M_0 \leq M_1$ is the arithmetic-mean–geometric-mean inequality given by Fact 1.15.14.) (Remark: A matrix application of this result is given by Fact 8.12.1.)

Fact 1.15.31. Let $x_1, \dots, x_n$ be nonnegative numbers, let $\alpha_1, \dots, \alpha_n$ be nonnegative numbers, and assume that $\sum_{i=1}^n \alpha_i = 1$. Then,

$$\prod_{i=1}^n x_i \leq \sum_{i=1}^n \alpha_i x_i^{1/\alpha_i}.$$

Furthermore, equality holds if and only if $x_1 = x_2 = \dots = x_n$. (Proof: See [447].) (Remark: This result is a generalization of Young's inequality. See Fact 1.10.32. Matrix versions are given by Fact 8.12.12 and Fact 9.14.22.) (Remark: This result is equivalent to Fact 1.15.32.)

Fact 1.15.32. Let $x_1, \dots, x_n$ be positive numbers, let $\alpha_1, \dots, \alpha_n$ be nonnegative numbers, and assume that $\sum_{i=1}^n \alpha_i = 1$. Then,

$$\frac{1}{\sum_{i=1}^n \frac{\alpha_i}{x_i}} \leq \prod_{i=1}^n x_i^{\alpha_i} \leq \sum_{i=1}^n \alpha_i x_i.$$

Now, let r be a real number, define

$$M_r \triangleq \left(\sum_{i=1}^n \alpha_i x_i^r \right)^{1/r}.$$

and let p and q be real numbers such that $p \leq q$. Then,

$$M_p \leq M_q$$

and

$$\lim_{r \rightarrow -\infty} M_r = \min\{x_1, \dots, x_n\} \leq \lim_{r \rightarrow 0} M_r = M_0 \leq \lim_{r \rightarrow \infty} M_r = \max\{x_1, \dots, x_n\}.$$

Furthermore, equality holds if and only if $x_1 = x_2 = \dots = x_n$. (Remark: This result is the *weighted arithmetic-mean-geometric-mean* inequality. Setting $\alpha_1 = \dots = \alpha_n = 1/n$ yields Fact 1.15.14.) (Proof: Since $f(x) = -\log x$ is convex, it follows that

$$\log \prod_{i=1}^n x_i^{\alpha_i} = \sum_{i=1}^n \alpha_i \log x_i \leq \log \sum_{i=1}^n \alpha_i x_i.$$

To prove the second statement, define $f: [0, \infty)^n \mapsto [0, \infty)$ by $f(\mu_1, \dots, \mu_n) \triangleq \sum_{i=1}^n \alpha_i \mu_i - \prod_{i=1}^n \mu_i^{\alpha_i}$. Note that $f(\mu, \dots, \mu) = 0$ for all $\mu \geq 0$. If $x_1, \dots, x_n$ minimizes f , then $\partial f / \partial \mu_i(x_1, \dots, x_n) = 0$ for all $i = 1, \dots, n$, which implies that $x_1 = x_2 = \dots = x_n$.) (Remark: This result is equivalent to Fact 1.15.31.) (Remark: See [1039, p. 11].)

Fact 1.15.33. Let $x_1, \dots, x_n$ be nonnegative numbers. Then,

$$1 + \left(\prod_{i=1}^n x_i \right)^{1/n} \leq \left[\prod_{i=1}^n (1 + x_i) \right]^{1/n}.$$

Furthermore, equality holds if and only if $x_1 = x_2 = \dots = x_n$. (Proof: Use Fact 1.15.14. See [238, p. 210].) (Remark: This inequality is used to prove Corollary 8.4.15.)

Fact 1.15.34. Let $x_1, \dots, x_n$ be nonnegative numbers, and let p, q be positive numbers such that $p \leq q$. Then,

$$\left(\sum_{i=1}^n x_i^q \right)^{1/q} \leq \left(\sum_{i=1}^n x_i^p \right)^{1/p}.$$

Furthermore, the inequality is strict if and only if $p < q$ and at least two of the numbers $x_1, \dots, x_n$ are nonzero. (Proof: See Proposition 9.1.5.) (Remark: This result is the *power-sum inequality*. See [273, p. 213]. This result implies that the Hölder norm is a monotonic function of the exponent.)

Fact 1.15.35. Let $x_1, \dots, x_n$ be positive numbers, and let $\alpha_1, \dots, \alpha_n \in [0, 1]$ be such that $\sum_{i=1}^n \alpha_i = 1$. If $p \leq 0$ or $p \geq 1$, then

$$\left(\sum_{i=1}^n \alpha_i x_i \right)^p \leq \sum_{i=1}^n \alpha_i x_i^p.$$

Alternatively, if $p \in [0, 1]$, then

$$\sum_{i=1}^n \alpha_i x_i^p \leq \left(\sum_{i=1}^n \alpha_i x_i \right)^p.$$

Finally, equality in both cases holds if and only if either $p = 0$ or $p = 1$ or $x_1 = \dots = x_n$. (Remark: This result is a consequence of Jensen's inequality given by Fact 1.8.4.)

Fact 1.15.36. Let $0 < x_1 < \dots < x_n$, and let $\alpha_1, \dots, \alpha_n \geq 0$ satisfy $\sum_{i=1}^n \alpha_i = 1$. Then,

$$1 \leq \left(\sum_{i=1}^n \alpha_i x_i \right) \left(\sum_{i=1}^n \frac{\alpha_i}{x_i} \right) \leq \frac{(x_1 + x_n)^2}{4x_1 x_n}.$$

(Remark: This result is the *Kantorovich inequality*. See Fact 8.15.9 and [927].) (Remark: See Fact 1.15.37.)

Fact 1.15.37. Let $x_1, \dots, x_n$ be positive numbers, and define $\alpha \triangleq \min_{i=1, \dots, n} x_i$ and $\beta \triangleq \max_{i=1, \dots, n} x_i$. Then,

$$1 \leq \left(\frac{1}{n} \sum_{i=1}^n x_i \right) \left(\frac{1}{n} \sum_{i=1}^n \frac{1}{x_i} \right) \leq \frac{(\alpha + \beta)^2}{4\alpha\beta}.$$

(Proof: Use Fact 1.15.36 or Fact 1.16.21. See [430, p. 94] or [431, p. 119].) (Remark: The left-hand inequality is the arithmetic-mean-harmonic-mean inequality. See Fact 1.15.12. The right-hand inequality is *Schweitzer's inequality*. See [1394, 1409] for historical details.) (Remark: A matrix extension is given by Fact 8.10.29.)

Fact 1.15.38. Let $x_1, \dots, x_n$ be positive numbers, and let p and q be positive numbers. Then,

$$\left(\frac{1}{n} \sum_{i=1}^n x_i^p \right) \left(\frac{1}{n} \sum_{i=1}^n x_i^q \right) \leq \frac{1}{n} \sum_{i=1}^n x_i^{p+q}.$$

In particular, if $p \in [0, 1]$, Then,

$$\left(\frac{1}{n} \sum_{i=1}^n x_i^p \right) \left(\frac{1}{n} \sum_{i=1}^n x_i^{1-p} \right) \leq \frac{1}{n} \sum_{i=1}^n x_i^p.$$

(Proof: See [1398].) (Remark: These inequalities are interpolated in [1398].)

Fact 1.15.39. Let $x_1, \dots, x_n$ be positive numbers. Then,

$$\frac{1}{n} \sum_{k=1}^n \left(\prod_{i=1}^k x_i \right)^{1/k} \leq \left[\prod_{k=1}^n \left(\frac{1}{k} \sum_{i=1}^k x_i \right) \right]^{1/k}.$$

Furthermore, equality holds if and only if $x_1 = \dots = x_n$. (Remark: The result can be expressed as $\frac{1}{n}(z_1 + \dots + z_n) \leq \sqrt[n]{y_1 \cdots y_n}$, where $z_k \triangleq \sqrt[k]{x_1 \cdots x_k} \leq y_k \triangleq \frac{1}{k}(x_1 + \dots + x_k)$.) (Remark: This result is the *mixed arithmetic-geometric mean inequality*. This result is due to Nanjundiah. See [336, 983].)

Fact 1.15.40. Let $x_1, \dots, x_n$ be positive numbers, where $n \geq 2$. Then,

$$\sum_{k=1}^n \left(\prod_{i=1}^k x_i \right)^{1/k} \leq \frac{n}{\sqrt[n]{n!}} \sum_{k=1}^n x_k \leq e^{(n-1)/n} \sum_{k=1}^n x_k \leq e \sum_{k=1}^n x_k.$$

Furthermore, equality holds in all of these inequalities if and only if $x_1 = \dots = x_n = 0$. (Remark: The inequality $\frac{n}{\sqrt[n]{n!}} < e^{(n-1)/n}$, which is equivalent to $e(n/e)^n < n!$, follows from Fact 1.9.19.) (Remark: This result is a finite version of *Carleman's inequality*. See [336] and [542, p. 22].)

Fact 1.15.41. Let $x_1, \dots, x_n$ be positive numbers, not all of which are zero. Then,

$$\left(\sum_{i=1}^n x_i \right)^4 < (2 \tan^{-1} n)^2 \left(\sum_{i=1}^n x_i^2 \right) \sum_{i=1}^n i^2 x_i^2 < \pi^2 \left(\sum_{i=1}^n x_i^2 \right) \sum_{i=1}^n i^2 x_i^2.$$

Furthermore,

$$\left(\sum_{i=1}^n x_i \right)^2 < \frac{\pi^2}{6} \sum_{i=1}^n i^2 x_i^2.$$

(Proof: See [154] or [869, p. 18].) (Remark: The first and third terms in the first inequality constitute a finite version of the *Carlson inequality*. The last inequality follows from the Cauchy-Schwarz inequality. See [457, p. 175].)

Fact 1.15.42. Let $x_1, \dots, x_n$ be nonnegative numbers, and let $p > 1$. Then,

$$\sum_{k=1}^n \left(\frac{1}{k} \sum_{i=1}^k x_i \right)^p \leq \left(\frac{p}{p-1} \right)^p \sum_{k=1}^n x_k^p.$$

(Proof: See [849].) (Remark: This result is the *Hardy inequality*. See [336, 849].)

Fact 1.15.43. Let $x_1, \dots, x_n$ be nonnegative numbers, and let $p > 1$. Then,

$$\sum_{k=1}^n \left(\sum_{i=k}^n \frac{x_i}{i} \right)^p \leq p^p \sum_{k=1}^n x_k^p.$$

(Proof: See [849].) (Remark: This result is the *Copson inequality*.)

Fact 1.15.44. Let $x_1, \dots, x_n$, α , and β be positive numbers, let p and q be real numbers, and assume that one of the following conditions is satisfied:

i) $p \in (-\infty, 1] \setminus \{0\}$ and $(n-1)\alpha \leq \beta$.

ii) $p \geq 1$ and $(n^p - 1)\alpha \leq \beta$.

Then,

$$\frac{n}{(\alpha + \beta)^{1/p}} \leq \sum_{i=1}^n \left(\frac{x_i^q}{\alpha x_i^q + \beta \prod_{k=1}^n x_k^{q/n}} \right)^{1/p}.$$

(Proof: See [1461].)

Fact 1.15.45. Let $x_1, \dots, x_n$ be nonnegative numbers, and assume that $\sum_{i=1}^n x_i = 1$. Then,

$$0 \leq \log n - \sum_{i=1}^n x_i \log \frac{1}{x_i} \leq \frac{1}{2}(n^2 - n) \max_{i,j=1,\dots,n} |x_i - x_j|^2.$$

Furthermore, $\sum_{i=1}^n x_i \log \frac{1}{x_i} = 0$ if and only if $x_i = 1$ for some i , while $\sum_{i=1}^n x_i \log \frac{1}{x_i} = \log n$ if and only if $x_1 = \dots = x_n = 1/n$. (Proof: See [433].) (Remark: Define $0 \log \frac{1}{0} \triangleq 0$.) (Remark: Alternative entropy bounds involving $\max_{i,j=1,\dots,n} x_i/x_j$ are given in [434].)

Fact 1.15.46. Let $x_1, \dots, x_n$ be positive numbers, and assume that $\sum_{i=1}^n x_i = 1$. Then,

$$0 \leq \log n - \sum_{i=1}^n x_i \log \frac{1}{x_i} \leq \left(n \sum_{i=1}^n x_i^2 \right) - 1 \leq \left(\sum_{i=1}^n x_i^3 \right)^{1/2} \left[\left(\sum_{i=1}^n \frac{1}{x_i} \right) - n^2 \right]^{1/2}.$$

Consequently,

$$\log n + 1 - n \sum_{i=1}^n x_i^2 \leq \sum_{i=1}^n x_i \log \frac{1}{x_i} \leq \log n.$$

(Proof: See [433, 982].) (Remark: It follows from Fact 1.15.37 that $n^2 \leq \sum_{i=1}^n \frac{1}{x_i}$.)

Fact 1.15.47. Let $x_1, \dots, x_n$ be positive numbers, assume that $\sum_{i=1}^n x_i = 1$, and define $a \triangleq \min_{i=1,\dots,n} x_i$ and $b \triangleq \max_{i=1,\dots,n} x_i$. Then,

$$0 \leq \log n - \sum_{i=1}^n x_i \log \frac{1}{x_i} \leq \frac{1}{n} \lfloor \frac{n^2}{4} \rfloor (b-a) \log \frac{b}{a} \leq \frac{1}{n} \lfloor \frac{n^2}{4} \rfloor \frac{(b-a)^2}{\sqrt{ab}}.$$

(Proof: See [435].) (Remark: This result is based on Fact 1.16.18.) (Remark: See Fact 2.21.6.)

Fact 1.15.48. Let $x_1, \dots, x_n$ be nonnegative numbers. Then,

$$\frac{e^2}{4} \sum_{i=1}^n x_i^2 \leq \prod_{i=1}^n e^{x_i}.$$

Furthermore, equality holds for $n = 1$ and $x_1 = 2$. (Proof: See [1104].)

1.16 Facts on Scalar Identities and Inequalities in $2n$ Variables

Fact 1.16.1. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, let $\alpha, \beta \in \mathbb{R}$, and assume that either $0 \leq \beta \leq \alpha \leq \frac{1}{2}$ or $\frac{1}{2} \leq \alpha \leq \beta \leq 1$. Then,

$$\sum_{i=1}^n x_i^{1-\alpha} y_i^\alpha \sum_{i=1}^n x_i^\alpha y_i^{1-\alpha} \leq \sum_{i=1}^n x_i^{1-\beta} y_i^\beta \sum_{i=1}^n x_i^\beta y_i^{1-\beta}.$$

Furthermore, if x and y are nonnegative numbers, then

$$x^{1-\alpha} y^\alpha + x^\alpha y^{1-\alpha} \leq x^{1-\beta} y^\beta + x^\beta y^{1-\beta}.$$

(Remark: This monotonicity inequality is due to Callebaut. See [1386].)

Fact 1.16.2. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be real numbers. Furthermore, let $x_{[1]}, \dots, x_{[n]}$ denote a rearrangement of $x_1, \dots, x_n$ such that $x_{[1]} \geq \dots \geq x_{[n]}$. Then,

$$\sum_{i=1}^n (x_{[i]} - y_{[i]})^2 \leq \sum_{i=1}^n (x_{[i]} - y_i)^2.$$

(Proof: See [457, p. 180].)

Fact 1.16.3. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be real numbers, and assume that $x_1 \leq \dots \leq x_n$ and $y_1 \leq \dots \leq y_n$. Furthermore, let $x_{[1]}, \dots, x_{[n]}$ denote a rearrangement of $x_1, \dots, x_n$ such that $x_{[1]} \geq \dots \geq x_{[n]}$. Then,

$$n \sum_{i=1}^n x_{[i]} y_{[n-i+1]} \leq \left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n y_i \right) \leq n \sum_{i=1}^n x_{[i]} y_{[i]}.$$

Furthermore, each inequality is an equality if and only if either $x_1 = \dots = x_n$ or $y_1 = \dots = y_n$. (Proof: See [668, pp. 148, 149].) (Remark: This result is *Chebyshev's inequality*.)

Fact 1.16.4. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be real numbers. Furthermore, let $x_{[1]}, \dots, x_{[n]}$ denote a rearrangement of $x_1, \dots, x_n$ such that $x_{[1]} \geq \dots \geq x_{[n]}$. Then,

$$\sum_{i=1}^n x_{[i]} y_{[n-i+1]} \leq \sum_{i=1}^n x_i y_i \leq \sum_{i=1}^n x_{[i]} y_{[i]}.$$

(Proof: See [236, p. 127] and [971, p. 141].) (Remark: This result is the *Hardy-Littlewood rearrangement inequality*.) (Remark: See Fact 8.18.18.)

Fact 1.16.5. Let $x_1, \dots, x_n$ be nonnegative numbers, and let $y_1, \dots, y_n$ be real numbers. Furthermore, let $y_{[1]}, \dots, y_{[n]}$ denote a rearrangement of $y_1, \dots, y_n$

such that $y_{[1]} \geq \cdots \geq y_{[n]}$. Then, for all $k = 1, \dots, n$, it follows that

$$\sum_{i=1}^k x_{[i]} y_i \leq \sum_{i=1}^k x_{[i]} y_{[i]}$$

and

$$\sum_{i=1}^k x_{[i]} y_{[n-i+1]} \leq \sum_{i=1}^k x_i y_i.$$

Now, assume that $y_1, \dots, y_n$ are nonnegative numbers. Then, for all $k = 1, \dots, n$, it follows that

$$\sum_{i=1}^k x_{[i]} y_{[n-i+1]} \leq \sum_{i=1}^k x_i y_i \leq \sum_{i=1}^k x_{[i]} y_i \leq \sum_{i=1}^k x_{[i]} y_{[i]}.$$

(Proof: See [381, 838] and [971, p. 141].) (Remark: This result is an extension of the *Hardy-Littlewood rearrangement inequality*.)

Fact 1.16.6. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be positive numbers, and let p, q be positive numbers such that, for all $i = 1, \dots, n$,

$$q \leq \frac{x_i}{y_i} \leq p.$$

Furthermore, let $x_{[1]}, \dots, x_{[n]}$ denote a rearrangement of $x_1, \dots, x_n$ such that $x_{[1]} \geq \cdots \geq x_{[n]}$. Then,

$$\sum_{i=1}^n x_{[i]} y_{[i]} \leq \frac{p+q}{2\sqrt{pq}} \sum_{i=1}^n x_i y_i.$$

(Remark: This result is a reverse rearrangement inequality.) (Remark: Equality holds for $x_1 = 2, x_2 = 1, y_1 = 1/2, y_2 = 2, q = 1$, and $p = 4$. Consequently, if $q = \min_{i=1, \dots, n} x_i/y_i$ and $p = \max_{i=1, \dots, n} x_i/y_i$, then the coefficient $\frac{p+q}{2\sqrt{pq}}$ is the best possible.) (Proof: See [251].)

Fact 1.16.7. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and assume that $x_1 \geq \cdots \geq x_n$ and $y_1 \geq \cdots \geq y_n$. Then,

$$\prod_{i=1}^n (x_i^2 + y_i^2) \leq \prod_{i=1}^n (x_i^2 + y_{n-i+1}^2).$$

(Remark: See Fact 8.13.11.)

Fact 1.16.8. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be complex numbers. Then,

$$\left| \sum_{i=1}^n x_i y_i \right|^2 = \sum_{i=1}^n |x_i|^2 \sum_{i=1}^n |y_i|^2 - \sum_{i < j} |\bar{x}_i y_j - \bar{x}_j y_i|^2.$$

(Remark: This result is the *Lagrange identity*. For the complex case, see [430, p. 6] or [431, p. 3]. For the real case, see [1322, 314].)

Fact 1.16.9. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be real numbers. Then,

$$\sum_{i=1}^n x_i y_i \leq \left(\sum_{i=1}^n x_i^2 \right)^{1/2} \left(\sum_{i=1}^n y_i^2 \right)^{1/2}.$$

Furthermore, equality holds if and only if $[x_1 \cdots x_n]^T$ and $[y_1 \cdots y_n]^T$ are linearly dependent. (Remark: This result is the *Cauchy-Schwarz inequality*.)

Fact 1.16.10. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be real numbers, and assume that $x_1 \leq \cdots \leq x_n$ and $y_1 \leq \cdots \leq y_n$. Then,

$$\left(\sum_{i=1}^n x_i \right) \left(\sum_{i=1}^n y_i \right) \leq n \sum_{i=1}^n x_i y_i.$$

(Proof: See [68, p. 27].)

Fact 1.16.11. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and let $\alpha \in [0, 1]$. Then,

$$\sum_{i=1}^n x_i^\alpha y_i^{1-\alpha} \leq \left(\sum_{i=1}^n x_i \right)^\alpha \left(\sum_{i=1}^n y_i \right)^{1-\alpha}.$$

Now, let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$. Then, equivalently,

$$\sum_{i=1}^n x_i y_i \leq \left(\sum_{i=1}^n x_i^p \right)^{1/p} \left(\sum_{i=1}^n y_i^q \right)^{1/q}.$$

Furthermore, equality holds if and only if $[x_1^p \cdots x_n^p]^T$ and $[y_1^q \cdots y_n^q]^T$ are linearly dependent. (Remark: This result is *Hölder's inequality*.) (Remark: Note the relationship between the *conjugate parameters* p, q and the *barycentric coordinates* $\alpha, 1 - \alpha$. See Fact 8.21.50.) (Remark: See Fact 9.7.34.)

Fact 1.16.12. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be complex numbers, let p, q, r be positive numbers, and assume that $1/p + 1/q = 1/r$. If $p \in (0, 1)$, $q < 0$, and $r = 1$, then

$$\left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \left(\sum_{i=1}^n |y_i|^q \right)^{1/q} \leq \sum_{i=1}^n |x_i y_i|.$$

Furthermore, if $p, q, r > 0$, then

$$\left(\sum_{i=1}^n |x_i y_i|^r \right)^{1/r} \leq \left(\sum_{i=1}^n |x_i|^p \right)^{1/p} \left(\sum_{i=1}^n |y_i|^q \right)^{1/q}.$$

(Proof: See [1039, p. 19].) (Remark: This result is the *Rogers-Hölder inequality*.) (Remark: Extensions of this result involving negative values of p, q , and r are considered in [1039, p. 19].) (Remark: See Proposition 9.1.6.)

Fact 1.16.13. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$. Then,

$$\sum_{i=1}^n \sum_{j=1}^n \frac{x_i y_j}{i+j-1} \leq \frac{\pi}{\sin(\pi/p)} \left(\sum_{i=1}^n x_i^p \right)^{1/p} \left(\sum_{i=1}^n y_i^q \right)^{1/q}.$$

In particular,

$$\sum_{i=1}^n \sum_{j=1}^n \frac{x_i y_j}{i+j-1} \leq \pi \left(\sum_{i=1}^n x_i^2 \right)^{1/2} \left(\sum_{i=1}^n y_i^2 \right)^{1/2}.$$

(Proof: See [542, p. 66] or [849].) (Remark: This result is the *Hardy-Hilbert inequality*.) (Remark: It follows from Fact 1.16.11 that

$$\sum_{i=1}^n \sum_{j=1}^n x_i y_j \leq n \left(\sum_{i=1}^n x_i^p \right)^{1/p} \left(\sum_{i=1}^n y_i^q \right)^{1/q}.)$$

Fact 1.16.14. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$. Then,

$$\sum_{i=1}^n \sum_{j=1}^n \frac{x_i y_j}{\max\{i, j\}} \leq pq \left(\sum_{i=1}^n x_i^p \right)^{1/p} \left(\sum_{i=1}^n y_i^q \right)^{1/q}.$$

Furthermore,

$$\sum_{i=2}^n \sum_{j=2}^n \frac{x_i y_j}{\log ij} \leq \frac{\pi}{\sin(\pi/p)} \left(\sum_{i=2}^n i^{p-1} x_i^p \right)^{1/p} \left(\sum_{i=2}^n i^{q-1} y_i^q \right)^{1/q}.$$

In particular,

$$\sum_{i=2}^n \sum_{j=2}^n \frac{x_i y_j}{\log ij} \leq \pi \left(\sum_{i=2}^n i x_i^2 \right)^{1/2} \left(\sum_{i=2}^n i y_i^2 \right)^{1/2}.$$

(Proof: For the first result, see [96]. For the second result see [1472].) (Remark: Related inequalities are given in [1473].)

Fact 1.16.15. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and assume that, for all $i = 1, \dots, n$, $x_i + y_i > 0$. Then,

$$\left(\sum_{i=1}^n x_i y_i \right)^2 \leq \sum_{i=1}^n (x_i^2 + y_i^2) \sum_{i=1}^n \frac{x_i^2 y_i^2}{x_i^2 + y_i^2} \leq \sum_{i=1}^n x_i^2 \sum_{i=1}^n y_i^2.$$

(Proof: See [430, p. 37], [431, p. 51], or [1386].) (Remark: This interpolation of the Cauchy-Schwarz inequality is *Milne's inequality*.)

Fact 1.16.16. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and let $\alpha \in [0, 1]$. Then,

$$\left(\sum_{i=1}^n x_i y_i \right)^2 \leq \sum_{i=1}^n x_i^{1+\alpha} y_i^{1-\alpha} \sum_{i=1}^n x_i^{1-\alpha} y_i^{1+\alpha} \leq \sum_{i=1}^n x_i^2 \sum_{i=1}^n y_i^2.$$

(Proof: See [430, p. 43], [431, p. 51], or [1386].) (Remark: This interpolation of the Cauchy-Schwarz inequality is *Callebaut's inequality*.)

Fact 1.16.17. Let $x_1, \dots, x_{2n}$ and $y_1, \dots, y_{2n}$ be real numbers. Then,

$$\left(\sum_{i=1}^{2n} x_i y_i \right)^2 \leq \left(\sum_{i=1}^{2n} x_i y_i \right)^2 + \left[\sum_{i=1}^n (x_i y_{n+i} - x_{n+i} y_i) \right]^2 \leq \sum_{i=1}^{2n} x_i^2 \sum_{i=1}^{2n} y_i^2.$$

(Proof: See [430, p. 41] or [431, p. 49].) (Remark: This interpolation of the Cauchy-Schwarz inequality is *McLaughlin's inequality*.)

Fact 1.16.18. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and define $a \triangleq \min_{i=1, \dots, n} x_i$, and $b \triangleq \max_{i=1, \dots, n} x_i$, $c \triangleq \min_{i=1, \dots, n} y_i$, and $d \triangleq \max_{i=1, \dots, n} y_i$. Then,

$$\left| \sum_{i=1}^n x_i y_i - \frac{1}{n} \sum_{i=1}^n x_i \sum_{i=1}^n y_i \right| \leq \lfloor \frac{n}{2} \rfloor (1 - \frac{1}{n} \lfloor \frac{n}{2} \rfloor) (b - a)(d - c).$$

(Proof: See [435].) (Remark: This result is used in Fact 1.15.45.)

Fact 1.16.19. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be positive numbers, and assume that $\sum_{i=2}^n x_i^2 < x_1^2$. Then,

$$\left(x_1^2 - \sum_{i=2}^n x_i^2 \right) \left(y_1^2 - \sum_{i=2}^n y_i^2 \right) \leq \left(x_1 y_1 - \sum_{i=2}^n x_i y_i \right)^2.$$

(Remark: This result is *Aczels's inequality*. See [273, p. 16]. Extensions are given in [1462] and Fact 9.7.4.)

Fact 1.16.20. Let $x_1, \dots, x_n$ be real numbers, and let $z_1, \dots, z_n$ be complex numbers. Then,

$$\left| \sum_{i=1}^n x_i z_i \right|^2 \leq \frac{1}{2} \sum_{i=1}^n x_i^2 \left(\sum_{i=1}^n |z_i|^2 + \left| \sum_{i=1}^n z_i^2 \right| \right) \leq \sum_{i=1}^n x_i^2 \sum_{i=1}^n |z_i|^2.$$

(Proof: See [430, p. 40] or [431, p. 48].) (Remark: Conditions for equality in the left-hand inequality are given in [430, p. 40] or [431, p. 48].) (Remark: This interpolation of the Cauchy-Schwarz inequality is *De Bruijn's inequality*.)

Fact 1.16.21. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be positive numbers, and define $\alpha \triangleq \min_{i=1, \dots, n} x_i / y_i$ and $\beta \triangleq \max_{i=1, \dots, n} x_i / y_i$. Then,

$$\left(\sum_{i=1}^n x_i y_i \right)^2 \leq \sum_{i=1}^n x_i^2 \sum_{i=1}^n y_i^2 \leq \frac{(\alpha + \beta)^2}{4\alpha\beta} \left(\sum_{i=1}^n x_i y_i \right)^2.$$

Equivalently, let $a \triangleq \min_{i=1, \dots, n} x_i$, $A \triangleq \max_{i=1, \dots, n} x_i$, $b \triangleq \min_{i=1, \dots, n} y_i$, and $B \triangleq \max_{i=1, \dots, n} y_i$. Then,

$$\left(\sum_{i=1}^n x_i y_i \right)^2 \leq \sum_{i=1}^n x_i^2 \sum_{i=1}^n y_i^2 \leq \frac{(ab + AB)^2}{4abAB} \left(\sum_{i=1}^n x_i y_i \right)^2.$$

(Proof: See [430, p. 73] or [431, p. 92].) (Remark: This reversal of the Cauchy-Schwarz inequality is the *Polya-Szego inequality*.)

Fact 1.16.22. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be positive numbers, let $a \triangleq \min_{i=1, \dots, n} x_i$, $A \triangleq \max_{i=1, \dots, n} x_i$, $b \triangleq \min_{i=1, \dots, n} y_i$, and $B \triangleq \max_{i=1, \dots, n} y_i$, let p, q be positive numbers, and assume that $1/p + 1/q = 1$. Then,

$$\sum_{i=1}^n x_i y_i \leq \left(\sum_{i=1}^n x_i^p \right)^{1/p} \left(\sum_{i=1}^n y_i^q \right)^{1/q} \leq \gamma \sum_{i=1}^n x_i y_i,$$

where

$$\gamma \triangleq \frac{A^p B^q - a^p b^q}{[p(AB B^q - a B b^q)]^{1/p} [q(a B A^p - A b a^p)]^{1/q}}.$$

(Proof: See [1394].) (Remark: The left-hand inequality, which is a reversal of Hölder's inequality, is the *Diaz-Goldman-Metcalf inequality*.) (Remark: Setting $p = q = 1/2$ yields Fact 1.16.21.) (Remark: The case in which $1/p + 1/q = 1/r$ is discussed in [1394].)

Fact 1.16.23. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and define $m_x \triangleq \min_{i=1, \dots, n} x_i$, $m_y \triangleq \min_{i=1, \dots, n} y_i$, $M_x \triangleq \max_{i=1, \dots, n} x_i$, and $M_y \triangleq \max_{i=1, \dots, n} y_i$. Then,

$$\left(\sum_{i=1}^n x_i y_i \right)^2 \leq \sum_{i=1}^n x_i^2 \sum_{i=1}^n y_i^2 \leq \left(\sum_{i=1}^n x_i y_i \right)^2 + \frac{n^2}{3} (M_x M_y - m_x m_y)^2.$$

(Proof: See [748].) (Remark: This reversal of the Cauchy-Schwarz inequality is *Ozeki's inequality*.)

Fact 1.16.24. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and assume that, for all $i = 1, \dots, n$, $x_i + y_i > 0$. Then,

$$\sum_{i=1}^n \frac{x_i y_i}{x_i + y_i} \sum_{i=1}^n (x_i + y_i) \leq \sum_{i=1}^n x_i \sum_{i=1}^n y_i.$$

(Proof: See [430, p. 36] or [431, p. 42].) (Remark: For positive numbers x and y , define the *harmonic mean* $H(x, y)$ of x and y by

$$H(x, y) \triangleq \frac{2}{\frac{1}{x} + \frac{1}{y}}.$$

Then, this result is equivalent to

$$\sum_{i=1}^n H(x_i, y_i) \leq H\left(\sum_{i=1}^n x_i, \sum_{i=1}^n y_i\right).$$

See [430, p. 37] or [431, p. 43]. The factor of 2 appearing on the right-hand side in [430, 431] is not needed.) (Remark: This result is *Dragomir's inequality*.) (Remark: Letting α, β be positive numbers and defining the arithmetic mean $A(\alpha, \beta) \triangleq \frac{1}{2}(\alpha + \beta)$, it follows that

$$\frac{(\alpha + \beta)^2}{4\alpha\beta} = \frac{A(\alpha, \beta)}{H(\alpha, \beta)}.$$

For details, see [1409].)

Fact 1.16.25. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers. If $p \in (0, 1]$, then

$$\left[\sum_{i=1}^n (x_i + y_i)^p \right]^{1/p} \geq \left(\sum_{i=1}^n x_i^p \right)^{1/p} + \left(\sum_{i=1}^n y_i^p \right)^{1/p}.$$

If $p \geq 1$, then

$$\left[\sum_{i=1}^n (x_i + y_i)^p \right]^{1/p} \leq \left(\sum_{i=1}^n x_i^p \right)^{1/p} + \left(\sum_{i=1}^n y_i^p \right)^{1/p}.$$

Furthermore, equality holds if and only if either $p = 1$ or $[x_1 \cdots x_n]^T$ and $[y_1 \cdots y_n]^T$ are linearly dependent. (Remark: This result is *Minkowski's inequality*.) (Proof: See [263].)

Fact 1.16.26. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, let $\alpha_1, \dots, \alpha_n$ be nonnegative numbers, and assume that $\sum_{i=1}^n \alpha_i = 1$. Then,

$$x_1^{\alpha_1} \cdots x_n^{\alpha_n} + y_1^{\alpha_1} \cdots y_n^{\alpha_n} \leq (x_1 + y_1)^{\alpha_1} \cdots (x_n + y_n)^{\alpha_n}.$$

(Proof: See [783, p. 64].)

Fact 1.16.27. Let $x_1, \dots, x_n, y_1, \dots, y_n \in (-1, 1)$, and let m be a positive integer. Then,

$$\left[\sum_{i=1}^n \frac{1}{(1 - x_i y_i)^m} \right]^2 \leq \left[\sum_{i=1}^n \frac{1}{(1 - x_i^2)^m} \right] \left[\sum_{i=1}^n \frac{1}{(1 - y_i^2)^m} \right].$$

(Proof: See [430, p. 19] or [431, p. 19].)

Fact 1.16.28. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and assume that $\sum_{i=1}^n x_i$ and $\sum_{i=1}^n y_i$ are nonzero. Then,

$$\left(\frac{\sum_{i=1}^n x_i}{\sum_{i=1}^n y_i} \right)^{\sum_{i=1}^n x_i} \prod_{i=1}^n y_i^{x_i} \leq \prod_{i=1}^n x_i^{x_i}.$$

Furthermore, equality holds if and only if there exists $\alpha > 0$ such that, for all $i = 1, \dots, n$, $x_i = \alpha y_i$. (Proof: See [130].)

Fact 1.16.29. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be nonnegative numbers, and assume that $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i$. Then,

$$\prod_{i=1}^n y_i^{x_i} \leq \prod_{i=1}^n x_i^{x_i}.$$

In particular,

$$\left(\frac{1}{n} \sum_{i=1}^n x_i \right)^{\sum_{i=1}^n x_i} \leq \prod_{i=1}^n x_i^{x_i}.$$

(Proof: See Fact 1.16.28 and [1160].)

Fact 1.16.30. Let $x_1, \dots, x_n$ and $y_1, \dots, y_n$ be positive numbers. Then,

$$\sum_{i=1}^n x_i \log \frac{\sum_{j=1}^n x_j}{\sum_{j=1}^n y_j} \leq \sum_{i=1}^n x_i \log \frac{x_i}{y_i}.$$

If $\sum_{i=1}^n x_i = 1$, then

$$\sum_{i=1}^n x_i \log \frac{1}{x_i} \leq \sum_{i=1}^n x_i \log \frac{1}{y_i} + \log \sum_{i=1}^n y_i.$$

On the other hand, if $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i$, then

$$0 \leq \sum_{i=1}^n x_i \log \frac{1}{y_i} + \log \sum_{i=1}^n y_i.$$

Finally, if $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i = 1$, then

$$\sum_{i=1}^n x_i \log \frac{1}{x_i} \leq \sum_{i=1}^n x_i \log \frac{1}{y_i},$$

or, equivalently,

$$0 \leq \sum_{i=1}^n x_i \log \frac{x_i}{y_i}.$$

(Proof: See [982].) (Remark: $\sum_{i=1}^n x_i \log \frac{1}{x_i}$ is the *entropy*.) (Remark: A refined upper bound and positive lower bound for $\sum_{i=1}^n x_i \log \frac{x_i}{y_i}$ are given in [625].) (Remark: See Fact 2.21.6.) (Remark: Related results are given in [1184, p. 276].)

1.17 Facts on Scalar Identities and Inequalities in $3n$ Variables

Fact 1.17.1. Let $x_1, \dots, x_n, y_1, \dots, y_n, z_1, \dots, z_n$ be real numbers. Then,

$$\left(\sum_{i=1}^n x_i y_i z_i \right)^4 \leq \left(\sum_{i=1}^n x_i^4 \right) \left(\sum_{i=1}^n y_i^2 \right)^2 \left(\sum_{i=1}^n z_i^4 \right).$$

(Proof: See [68, p. 27].)

Fact 1.17.2. Let $x_1, \dots, x_n, y_1, \dots, y_n, z_1, \dots, z_n$ be complex numbers. Then,

$$\left| \sum_{i=1}^n x_i \overline{z_i} \sum_{i=1}^n z_i \overline{y_i} \right| \leq \frac{1}{2} \left(\sqrt{\sum_{i=1}^n |x_i|^2 \sum_{i=1}^n |y_i|^2} + \left| \sum_{i=1}^n x_i \overline{y_i} \right| \right) \sum_{i=1}^n |z_i|^2.$$

(Proof: See [514].) (Remark: This extension of the Cauchy-Schwarz inequality is *Buzano's inequality*.) (Remark: See *xv*) of Fact 9.7.4.)

1.18 Facts on Scalar Identities and Inequalities in Complex Variables

Fact 1.18.1. Let z be a complex number with complex conjugate $\bar{z}$, real part $\operatorname{Re} z$, and imaginary part $\operatorname{Im} z$. Then, the following statements hold:

- i) $-|z| \leq \operatorname{Re} z \leq |\operatorname{Re} z| \leq |z|$.
- ii) $-|z| \leq \operatorname{Im} z \leq |\operatorname{Im} z| \leq |z|$.
- iii) $0 \leq |z| = |-z| = |\bar{z}|$.
- iv) $\operatorname{Re} z = |\operatorname{Re} z| = |z|$ if and only if $\operatorname{Re} z \geq 0$ and $\operatorname{Im} z = 0$.
- v) $\operatorname{Im} z = |\operatorname{Im} z| = |z|$ if and only if $\operatorname{Im} z \geq 0$ and $\operatorname{Re} z = 0$.
- vi) If $z \neq 0$, then $\overline{z^{-1}} = \bar{z}^{-1}$.
- vii) If $z \neq 0$, then $z^{-1} = \bar{z}/|z|^2$.
- viii) If $z \neq 0$, then $|z^{-1}| = 1/|z|$.
- ix) If $|z| = 1$, then $z^{-1} = \bar{z}$.
- x) If $z \neq 0$, then $\operatorname{Re} z^{-1} = (\operatorname{Re} z)/|z|^2$.
- xi) $\operatorname{Re} z \neq 0$ if and only if $\operatorname{Re} z^{-1} \neq 0$.
- xii) If $\operatorname{Re} z \neq 0$, then $|z| = \sqrt{(\operatorname{Re} z)/(\operatorname{Re} z^{-1})}$.
- xiii) $|z^2| = |z|^2 = z\bar{z}$.
- xiv) $z^2 \geq 0$ if and only if $\operatorname{Im} z = 0$.
- xv) $z^2 \leq 0$ if and only if $\operatorname{Re} z = 0$.
- xvi) $z^2 + \bar{z}^2 + 4(\operatorname{Im} z)^2 = 2|z|^2$.
- xvii) $z^2 + \bar{z}^2 + 2|z|^2 = 4(\operatorname{Re} z)^2$.
- xviii) $z^2 + \bar{z}^2 + 2(\operatorname{Im} z)^2 = 2(\operatorname{Re} z)^2$.
- xix) $z^2 + \bar{z}^2 \leq \left\{ \begin{array}{l} |z^2 + \bar{z}^2| \\ (\operatorname{Re} z)^2 \end{array} \right\} \leq 2|z|^2$.
- xx) $z^2 + \bar{z}^2 = |z^2 + \bar{z}^2| = (\operatorname{Re} z)^2 = 2|z|^2$ if and only if $\operatorname{Im} z = 0$.
- xxi) Let n be a positive integer. If $z \neq 1$, then

$$\frac{1 - z^n}{1 - z} = \sum_{i=0}^{n-1} z^i = 1 + z + \cdots + z^{n-1}.$$

Furthermore,

$$\lim_{z \rightarrow 1} \frac{1 - z^n}{1 - z} = n.$$

(Remark: A matrix version of i) is given in [1271].)

Fact 1.18.2. Let z_1 and z_2 be complex numbers. Then, the following statements hold:

- i) $|z_1 z_2| = |z_1| |z_2|$.
- ii) If $z_2 \neq 0$, then $|z_1/z_2| = |z_1|/|z_2|$.
- iii) $||z_1| - |z_2|| \leq |z_1 + z_2| \leq |z_1| + |z_2|$.
- iv) $|z_1 + z_2| = |z_1| + |z_2|$ if and only if $\operatorname{Re}(z_1 \overline{z_2}) = |z_1| |z_2|$.
- v) $|z_1 + z_2| = |z_1| + |z_2|$ if and only if there exists $\alpha \geq 0$ such that either $z_1 = \alpha z_2$ or $z_2 = \alpha z_1$, that is, if and only if z_1 and z_2 have the same phase angle.
- vi) $||z_1| - |z_2|| \leq |z_1 - z_2|$.
- vii) $||z_1| - |z_2|| = |z_1 - z_2|$ if and only if there exists $\alpha \geq 0$ such that either $z_1 = \alpha z_2$ or $z_2 = \alpha z_1$, that is, if and only if z_1 and z_2 have the same phase angle.
- viii) $|1 + \overline{z_1} z_2|^2 = (1 - |z_1|)^2 (1 - |z_2|)^2 + |z_1 + z_2|^2 = (1 + |z_1|^2)(1 + |z_2|^2) - |z_1 - z_2|^2$.
- ix) $|z_1 - z_2|^2 \leq (1 + |z_1|^2)(1 + |z_2|^2)$.
- x) $\frac{1}{2}|z_1 - z_2| + \left| \frac{z_2}{z_1} z_1 - \frac{z_1}{z_2} z_2 \right| = \frac{1}{2}(|z_1| + |z_2|) \left| \frac{z_1}{|z_1|} - \frac{z_2}{|z_2|} \right| \leq |z_1 - z_2|$.
- xi) $2 \operatorname{Re}(z_1 z_2) \leq |z_1|^2 + |z_2|^2$.
- xii) $2 \operatorname{Re}(z_1 z_2) = |z_1|^2 + |z_2|^2$ if and only if $z_1 = \overline{z_2}$.
- xiii) $\frac{1}{2}(|z_1 + z_2|^2 + |z_1 - z_2|^2) = |z_1|^2 + |z_2|^2$.
- xiv) $z_1 \overline{z_2} = \frac{1}{4}(|z_1 + z_2|^2 - |z_1 - z_2|^2 + j|z_1 + jz_2|^2 - j|z_1 - jz_2|^2)$.
- xv) If $a, b \in \mathbb{C}$, $|a| \neq |b|$, and $z_2 = az_1 + b\overline{z_1}$, then

$$z_1 = \frac{\overline{a}z_2 - b\overline{z_2}}{|a|^2 - |b|^2}.$$
- xvi) If $p \geq 1$, then

$$|z_1 + z_2|^p \leq 2^{p-1}(|z_1|^p + |z_2|^p).$$
- xvii) If $p \geq 2$, then

$$2(|z_1|^p + |z_2|^p) \leq |z_1 + z_2|^p + |z_1 - z_2|^p \leq 2^{p-1}(|z_1|^p + |z_2|^p).$$
- xviii) If $p \geq 2$, $q > 0$, and $1/p + 1/q = 1$, then

$$2(|z_1|^p + |z_2|^p)^{q-1} \leq |z_1 + z_2|^q + |z_1 - z_2|^q.$$
- xix) If $p \in (1, 2]$, $q > 0$, and $1/p + 1/q = 1$, then

$$|z_1 + z_2|^q + |z_1 - z_2|^q \leq 2(|z_1|^p + |z_2|^p)^{q-1}.$$
- xx) Let n be a positive integer. If $z_1 \neq z_2$, then

$$\frac{z_1^n - z_2^n}{z_1 - z_2} = z_1^{n-1} + z_2 z_1^{n-2} + \cdots + z_2^{n-1}.$$

Furthermore,

$$\lim_{z_2 \rightarrow z_1} \frac{z_1^n - z_2^n}{z_1 - z_2} = n z_1^{n-1}.$$

Now, let z_1, z_2 , and z_3 be complex numbers. Then, the following statements hold:

$$xxi) |z_1 + z_2|^2 + |z_2 + z_3|^2 + |z_3 + z_1|^2 = |z_1|^2 + |z_2|^2 + |z_3|^2 + |z_1 + z_2 + z_3|^2.$$

$$xxii) |z_1 + z_2| + |z_2 + z_3| + |z_3 + z_1| \leq |z_1| + |z_2| + |z_3| + |z_1 + z_2 + z_3|.$$

$$xxiii) 4(|z_1|^2 + |z_2|^2 + |z_3|^2) \leq |z_1 + z_2 + z_3|^2 + |z_1 + z_2 - z_3|^2 + |z_1 - z_2 + z_3|^2 + |z_1 - z_2 - z_3|^2.$$

$$xxiv) \text{ If } z_1, z_2, z_3 \text{ are nonzero and } z_1^7 + z_2^7 + z_3^7 = 0, \text{ then } |z_1| = |z_2| = |z_3|.$$

Finally, for $i = 1, \dots, n$, let $z_i = r_i e^{j\phi_i}$ be complex numbers, where $r_i \geq 0$ and $\phi_i \in \mathbb{R}$, and assume that there exist $\theta_1, \theta_2 \in \mathbb{R}$ such that $0 < \theta_2 - \theta_1 < \pi$ and such that, for all $i = 1, \dots, n$, $\theta_1 \leq \phi_i \leq \theta_2$. Then, the following inequality holds:

$$xxv) \cos[\frac{1}{2}(\theta_2 - \theta_1)] \sum_{i=1}^n |z_i| \leq |\sum_{i=1}^n z_i|.$$

(Remark: Matrix versions of *i*), *iii*), *v*)–*vii*) are given in [1271]. Result *viii*) is given in [59, p. 19] and [1467]. Result *x*) is the *Dunkl-Williams inequality*. See [430, p. 43] or [431, p. 52] and *ii*) of Fact 9.7.4. Result *xiii*) is the parallelogram law; see [449] and Fact 9.7.4. Result *xiv*) is the *polarization identity*; see [368, p. 54], [1030, p. 276], and Fact 9.7.4. Result *xv*) is given in [734]. Result *xvi*) is given in [695]. Results *xvii*)–*xix*) are due to Clarkson; see [695], [1010, p. 536], and Fact 9.9.34. Result *xxi*) is given in [59, p. 19]. Result *xxii*) is *Hlawka's inequality*. See Fact 1.8.6 and Fact 9.7.4. Result *xxiii*) is given in [449]. Result *xxiv*) is given in [59, pp. 186, 187]. Result *xxv*) is due to Petrovich; see [432].) (Remark: The absolute value $|z| = |x + jy|$, where x and y are real, is identical to the Euclidean norm $\|[\begin{smallmatrix} x \\ y \end{smallmatrix}]\|_2$. Therefore, each result in Section 9.7 involving the Euclidean norm on $\mathbb{R}^2$ can be recast in terms of complex numbers.) (Problem: Compare the lower bounds for $|z_1 - z_2|$ given by *iv*) and *vii*.)

Fact 1.18.3. Let a, b, c be complex numbers, and assume that $a \neq 0$. Then, $z \in \mathbb{C}$ satisfies

$$az^2 + bz + c = 0$$

if and only if

$$z = \frac{1}{2a}(y - b),$$

where

$$y = \pm \frac{1}{\sqrt{2}}(\sqrt{|\Delta| + \operatorname{Re} \Delta} + j \operatorname{sign}(\operatorname{Im} \Delta) \sqrt{|\Delta| - \operatorname{Re} \Delta})$$

and

$$\Delta \triangleq b^2 - 4ac.$$

If, in addition, a, b, c are real, then $z \in \mathbb{C}$ satisfies

$$az^2 + bz + c = 0$$

if and only if

$$z = \frac{1}{2a}(-b \pm \sqrt{b^2 - 4ac}).$$

(Proof: See [59, pp. 15, 16].)

Fact 1.18.4. Let $z, z_1, \dots, z_n$ be complex numbers. Then,

$$\frac{1}{n} \sum_{i=1}^n |z - z_i|^2 = \left| z - \frac{1}{n} \sum_{i=1}^n z_i \right|^2 + \frac{1}{n} \sum_{1 \leq i < j \leq n} |z_i - z_j|^2.$$

(Proof: See [59, pp. 146].)

Fact 1.18.5. let z_1 and z_2 be complex numbers. Then,

$$\begin{aligned} \frac{|z_1 - z_2| - ||z_1| - |z_2||}{\min\{|z_1|, |z_2|\}} &\leq \left| \frac{z_1}{|z_1|} - \frac{z_2}{|z_2|} \right| \\ &\leq \left\{ \begin{array}{l} \frac{|z_1 - z_2| + ||z_1| - |z_2||}{\max\{|z_1|, |z_2|\}} \\ \frac{2|z_1 - z_2|}{|z_1| + |z_2|} \end{array} \right\} \\ &\leq \left\{ \begin{array}{l} \frac{2|z_1 - z_2|}{\max\{|z_1|, |z_2|\}} \\ \frac{2(|z_1 - z_2| + ||z_1| - |z_2||)}{|z_1| + |z_2|} \end{array} \right\} \\ &\leq \frac{4|z_1 - z_2|}{|z_1| + |z_2|}. \end{aligned}$$

(Proof: See Fact 9.7.10.) (Remark: The second and lower third terms constitute the Dunkl-Williams inequality given by Fact 1.18.2.)

Fact 1.18.6. Let z be a complex number. Then, the following statements hold:

- i) $0 < |e^z| \leq e^{|z|}$.
- ii) $|e^z| = e^{|z|}$ if and only if $\text{Im } z = 0$ and $\text{Re } z \geq 0$.
- iii) $|e^z| = 1$ if and only if $\text{Re } z = 0$.
- iv) $||e^z| - 1| \leq |e^z - 1| \leq e^{|z|} - 1$.
- v) If $|z| < \log 2$, then $|e^z - 1| \leq e^{|z|} - 1 < 1$.
- vi) $e^z = e^{\text{Re } z} [\cos(\text{Im } z) + j \sin(\text{Im } z)]$.
- vii) $\text{Re } e^z = 0$ if and only if $\text{Im } z$ is an odd integer multiple of $\pm\pi/2$.
- viii) $\text{Im } e^z = 0$ if and only if $\text{Im } z$ is an integer multiple of $\pm\pi$.
- ix) If z is nonzero, then $|z^j| < e^\pi$.

Furthermore, let θ_1 and θ_2 be real numbers. Then, the following statements hold:

- x) $|e^{j\theta_1} - e^{j\theta_2}| \leq |\theta_1 - \theta_2|$.
- xi) $|e^{j\theta_1} - e^{j\theta_2}| = |\theta_1 - \theta_2|$ if and only if $\theta_1 = \theta_2$.

Finally, let r_1 and r_2 be nonnegative numbers, at least one of which is positive.

Then, the following statement holds:

$$xii) |e^{j\theta_1} - e^{j\theta_2}| \leq \frac{2|r_1e^{j\theta_1} - r_2e^{j\theta_2}|}{r_1 + r_2}.$$

(Proof: Statement *xii*) is given in [683, p. 218].) (Remark: A matrix version of *x*) is given by Fact 11.16.13.)

Fact 1.18.7. Let z be a complex number. Then, for all nonzero $z \in \mathbb{C}$, there exist infinitely many $s \in \mathbb{C}$ such that $e^s = z$. Specifically, let $z = re^{j\phi}$, where $r > 0$ and $\phi \in \mathbb{R}$. Then, for all $k \in \mathbb{Z}$, $s = \log r + j(\phi + 2\pi k)$ satisfies $e^s = z$, where $\log r$ is the positive logarithm of r . In particular, for all odd integers k , $e^{\pm j\pi k} = -1$, while, for all even integers k , $e^{\pm j\pi k} = 1$. To obtain a single-valued definition of $\log$, let $z \in \mathbb{C}$ be nonzero, and write z uniquely as $z = re^{j\phi}$, where $r > 0$ and $\phi \in (-\pi, \pi]$. Then, the *principal branch* of the log function $\log z \in \mathbb{C}$ is defined as

$$\log z \triangleq \log r + j\phi.$$

The principal branch of the log function

$$\log: \mathbb{C} \setminus \{0\} \mapsto \{z: \operatorname{Re} z \neq 0 \text{ and } -\pi < \operatorname{Im} z \leq \pi\}$$

has the following properties:

i) If $z \in \mathbb{C}$ is nonzero, then

$$e^{\log z} = z.$$

ii) Let $z = re^{j\phi} \in \mathbb{C}$, where $r \geq 0$ and $\phi \in (-\pi, \pi]$, and assume that $r \sin \phi \in (-\pi, \pi]$. Then,

$$\log e^z = z.$$

iii) Let $z_1 = r_1e^{j\phi_1}$ and $z_2 = r_2e^{j\phi_2}$, where $r_1, r_2 > 0$ and $\phi_1, \phi_2 \in (-\pi, \pi]$, and assume that $\phi_1 + \phi_2 \in (-\pi, \pi]$. Then,

$$\log z_1 z_2 = \log z_1 + \log z_2.$$

Now, define $\mathcal{D} \triangleq \{z \in \mathbb{C}: |z - 1| < 1\}$. Then, the following statements hold:

iv) For all $z \in \mathcal{D}$, $\log z$ is given by the convergent series

$$\log z = \sum_{i=1}^{\infty} \frac{(-1)^{i+1}}{i} (z - 1)^i.$$

v) If $z \in \mathcal{D}$, then

$$\log e^z = z.$$

vi) If $z_1, z_2 \in \mathcal{D}$, then

$$\log z_1 z_2 = \log z_1 + \log z_2.$$

vii) If $|z| < 1$, then

$$|\log(1 + z)| \leq -\log(1 - |z|)$$

and

$$\frac{|z|}{1 + |z|} \leq |\log(1 + z)| \leq \frac{|z|(1 + |z|)}{|1 + z|}.$$

(Remark: Let $z = re^{j\theta} \in \mathbb{C}$ satisfy $|z - 1| < 1$. Then, $-\pi/2 < \theta < \pi/2$. Furthermore, $\log z = (\log r) + j\theta$, and thus $-\pi/2 < \operatorname{Im} \log z < \pi/2$. Consequently, the infinite series in *iv*) gives the principal log of z .)

Fact 1.18.8. The following infinite series converge for the given values of the complex argument z :

i) For all $z \in \mathbb{C}$,
$$\sin z = z - \frac{1}{3!}z^3 + \frac{1}{5!}z^5 - \frac{1}{7!}z^7 + \dots$$

ii) For all $z \in \mathbb{C}$,
$$\cos z = 1 - \frac{1}{2!}z^2 + \frac{1}{4!}z^4 - \frac{1}{6!}z^6 + \dots$$

iii) For all $|z| < \pi/2$,
$$\tan z = z + \frac{1}{3}z^3 + \frac{2}{15}z^5 + \frac{17}{315}z^7 + \frac{62}{2835}z^9 + \dots$$

iv) For all $z \in \mathbb{C}$,
$$e^z = 1 + z + \frac{1}{2!}z^2 + \frac{1}{3!}z^3 + \frac{1}{4!}z^4 + \dots$$

v) For all nonzero $z \in \mathbb{C}$ such that $|z - 1| \leq 1$,
$$\log z = -\left[1 - z + \frac{1}{2}(1 - z)^2 + \frac{1}{3}(1 - z)^3 + \frac{1}{4}(1 - z)^4 + \dots\right].$$

vi) For all $z \in \text{CUD} \setminus \{1\}$,
$$\log(1 - z) = -\left(z + \frac{1}{2}z^2 + \frac{1}{3}z^3 + \frac{1}{4}z^4 + \dots\right).$$

vii) For all $z \in \text{CUD} \setminus \{-1\}$,
$$\log(1 + z) = z - \frac{1}{2}z^2 + \frac{1}{3}z^3 - \frac{1}{4}z^4 + \dots$$

viii) For all $z \in \text{CUD} \setminus \{-1, 1\}$,
$$\log \frac{1+z}{1-z} = 2\left(z + \frac{1}{3}z^3 + \frac{1}{5}z^5 + \dots\right).$$

ix) For all $z \in \mathbb{C}$ such that $\text{Re } z > 0$,

$$\log z = \sum_{i=0}^{\infty} \frac{2}{2i+1} \left(\frac{z-1}{z+1} \right)^{2i+1}.$$

x) For all $z \in \mathbb{C}$,
$$\sinh z = \sin jz = z + \frac{1}{3!}z^3 + \frac{1}{5!}z^5 + \frac{1}{7!}z^7 + \dots$$

xi) For all $z \in \mathbb{C}$,
$$\cosh z = \cos jz = 1 + \frac{1}{2!}z^2 + \frac{1}{4!}z^4 + \frac{1}{6!}z^6 + \dots$$

xii) For all $|z| < \pi/2$,
$$\tanh z = \tan jz = z - \frac{1}{3}z^3 + \frac{2}{15}z^5 - \frac{17}{315}z^7 + \frac{62}{2835}z^9 - \dots$$

xiii) For all $\alpha \in \mathbb{C}$ and $|z| \leq 1$ such that either $|z| < 1$ or both $\text{Re } \alpha > -1$ and $|z| \neq -1$,

$$(1+z)^\alpha = 1 + \alpha z + \frac{\alpha(\alpha-1)}{2!}z^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!}z^3 + \frac{\alpha(\alpha-1)(\alpha-2)(\alpha-3)}{4!}z^4 + \dots$$

$$= \binom{\alpha}{0} + \binom{\alpha}{1}z + \binom{\alpha}{2}z^2 + \binom{\alpha}{3}z^3 + \binom{\alpha}{4}z^4 + \dots$$

xiv) For all $\alpha \in \mathbb{C}$ and $|z| < 1$,

$$\frac{1}{(1-z)^{\alpha+1}} = \binom{\alpha}{0} + \binom{1+\alpha}{1}z + \binom{2+\alpha}{2}z^2 + \binom{3+\alpha}{3}z^3 + \binom{4+\alpha}{4}z^4 + \dots$$

xv) For all $|z| < 1$,

$$(1 - z)^{-1} = 1 + z + z^2 + z^3 + z^4 + \cdots.$$

(Proof: See [750, pp. 11, 12]. For $x \in \mathbb{R}$ such that $|x| < 1$, it follows that

$$\frac{d}{dx} \log(1 - x) = \frac{-1}{1 - x} = -(1 + x + x^2 + \cdots).$$

Integrating yields

$$\log(1 - x) = -(x + \frac{1}{2}x^2 + \frac{1}{3}x^3 + \cdots).$$

Using analytic continuation to replace $x \in \mathbb{R}$ satisfying $|x| < 1$ with $z \in \mathbb{C}$ satisfying $|z| < 1$ yields *vii)*. (Remark: *vii)* is *Mercator's series*, while *viii)* and *ix)* are equivalent forms of *Gregory's series*. See [683, p. 273].) (Remark: *xiii)* is the *binomial series*.) (Remark: $\text{CUD} = \{z \in \mathbb{C} : |z| \leq 1\}$.)

1.19 Facts on Trigonometric and Hyperbolic Identities

Fact 1.19.1. Let x be a real number such that the expressions below are defined. Then, the following identities hold:

$$i) \quad \sin x = \frac{1}{2j}(e^{jx} - e^{-jx}).$$

$$ii) \quad \cos x = \frac{1}{2}(e^{jx} + e^{-jx}).$$

$$iii) \quad \sin(x + y) = (\sin x)(\cos y) + (\cos x) \sin y.$$

$$iv) \quad \sin(x - y) = (\sin x)(\cos y) - (\cos x) \sin y.$$

$$v) \quad \cos(x + y) = (\cos x)(\cos y) - (\sin x) \sin y.$$

$$vi) \quad \cos(x - y) = (\cos x)(\cos y) + (\sin x) \sin y.$$

$$vii) \quad (\sin x) \sin y = \frac{1}{2}[\cos(x - y) - \cos(x + y)].$$

$$viii) \quad (\sin x) \cos y = \frac{1}{2}[\sin(x + y) + \sin(x - y)].$$

$$ix) \quad (\cos x) \cos y = \frac{1}{2}[\cos(x + y) + \cos(x - y)].$$

$$x) \quad \sin^2 x - \sin^2 y = [\sin(x + y)] \sin(x - y).$$

$$xi) \quad \cos^2 x - \sin^2 y = [\cos(x + y)] \cos(x - y).$$

$$xii) \quad \cos^2 x - \cos^2 y = [\sin(x + y)] \sin(y - x).$$

$$xiii) \quad \sin x + \sin y = 2[\sin \frac{1}{2}(x + y)] \cos \frac{1}{2}(x - y).$$

$$xiv) \quad \sin x - \sin y = 2[\sin \frac{1}{2}(x - y)] \cos \frac{1}{2}(x + y).$$

$$xv) \quad \cos x + \cos y = 2[\cos \frac{1}{2}(x + y)] \cos \frac{1}{2}(x - y).$$

$$xvi) \quad \cos x - \cos y = 2[\sin \frac{1}{2}(x + y)] \sin \frac{1}{2}(y - x).$$

$$xvii) \quad \tan(x + y) = \frac{(\tan x) + \tan y}{1 - (\tan x) \tan y}.$$

$$xviii) \quad \tan(x - y) = \frac{(\tan x) - \tan y}{1 + (\tan x) \tan y}.$$

$$xix) \quad \tan x + \tan y = \frac{\sin(x + y)}{(\cos x) \cos y}.$$

$$xx) \quad \tan x - \tan y = \frac{\sin(x-y)}{(\cos x) \cos y}.$$

$$xxi) \quad \sin x = 2(\sin \frac{x}{2}) \cos \frac{x}{2}.$$

$$xxii) \quad \cos x = 2(\cos^2 \frac{x}{2}) - 1.$$

$$xxiii) \quad \sin 2x = 2(\sin x) \cos x.$$

$$xxiv) \quad \cos 2x = 2(\cos^2 x) - 1.$$

$$xxv) \quad \tan 2x = \frac{2 \tan x}{1 - \tan^2 x}.$$

$$xxvi) \quad \sin 3x = 3(\sin x) - 4 \sin^3 x.$$

$$xxvii) \quad \cos 3x = 4(\cos^3 x) - 3 \cos x.$$

$$xxviii) \quad \tan 3x = \frac{3(\tan x) - \tan^3 x}{1 - 3 \tan^2 x}.$$

$$xxix) \quad \sin^2 x = \frac{1}{2}(1 - \cos 2x).$$

$$xxx) \quad \cos^2 x = \frac{1}{2}(1 + \cos 2x).$$

$$xxxi) \quad \tan^2 x = \frac{1 - \cos 2x}{1 + \cos 2x}.$$

$$xxxii) \quad \tan x = \frac{\sin 2x}{1 + \cos 2x} = \frac{1 - \cos 2x}{\sin 2x} = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}}.$$

$$xxxiii) \quad \sin^2 \frac{x}{2} = \frac{1}{2}(1 - \cos x).$$

$$xxxiv) \quad \cos^2 \frac{x}{2} = \frac{1}{2}(1 + \cos x).$$

$$xxxv) \quad \tan \frac{1}{2}x = \frac{\sin x}{1 + \cos x} = \frac{1 - \cos x}{\sin x}.$$

$$xxxvi) \quad \text{For all } t \geq 0 \text{ and } \alpha \in (0, 1),$$

$$\int_0^\infty \frac{tx^{\alpha-1}}{t+x} dx = \frac{t^\alpha \pi}{\sin \alpha \pi}.$$

(Remark: See [750, pp. 114–116]. The last result is given in [1503, p. 448, formula 589]. See also [542, p. 69].)

Fact 1.19.2. Let x be a real number such that the expressions below are defined. Then, the following identities hold:

$$i) \quad \sinh x = \frac{1}{2}(e^x - e^{-x}).$$

$$ii) \quad \cosh x = \frac{1}{2}(e^x + e^{-x}).$$

$$iii) \quad \tanh x = \frac{\sinh x}{\cosh x}.$$

$$iv) \quad \sin jx = j \sinh x.$$

$$v) \quad \cos jx = j \cosh x.$$

$$vi) \quad \tan jx = j \tanh x.$$

$$vii) \quad \sinh jx = j \sin x.$$

$$viii) \quad \cosh jx = j \cos x.$$

$$ix) \quad \tanh jx = j \tan x.$$

$$x) \sinh(x + y) = (\sinh x)(\cosh y) + (\cosh x) \sinh y.$$

$$xi) \cosh(x + y) = (\cosh x)(\cosh y) + (\sinh x) \sinh y.$$

$$xii) \tanh(x + y) = \frac{(\tanh x) + \tanh y}{1 + (\tanh x) \tanh y}.$$

(Remark: See [750, pp. 117–119].)

Fact 1.19.3. Let $z = x + jy$, where z is a complex number and x and y are real numbers. Then, the following identities hold:

$$i) \sin z = (\sin x)(\cosh y) + j(\cos x) \sinh y.$$

$$ii) \cos z = (\cos x)(\cosh y) - j(\sin x) \sinh y.$$

$$iii) \tan z = \frac{(\sin 2x) + j \sinh 2y}{(\cos 2x) + \cosh 2y}.$$

1.20 Notes

Much of the preliminary material in this chapter can be found in [1030]. A related treatment of mathematical preliminaries is given in [1129]. An extensive introduction to logic and mathematical fundamentals is given in [229]. In [229], the notation “ $A \rightarrow B$ ” is used to denote an implication, which is called a *disjunction*, while “ $A \implies B$ ” indicates a tautology.

An extensive treatment of partially ordered sets is given in [1179]. Lattices are discussed in [229].

Alternative terminology for “one-to-one” and “onto” is *injective* and *surjective*, respectively, while a function that is injective and surjective is *bijective*.

Reference works on inequalities include [162, 273, 274, 275, 340, 637, 963, 971, 1010, 1221]. Recommended texts on complex variables include [725, 1031, 1066].

Chapter Two

Basic Matrix Properties

In this chapter we provide a detailed treatment of the basic properties of matrices such as range, null space, rank, and invertibility. We also consider properties of convex sets, cones, and subspaces.

2.1 Matrix Algebra

The symbols $\mathbb{Z}$, $\mathbb{N}$, and $\mathbb{P}$ denote the sets of integers, nonnegative integers, and positive integers, respectively. The symbols $\mathbb{R}$ and $\mathbb{C}$ denote the real and complex number fields, respectively, whose elements are *scalars*. Since $\mathbb{R}$ is a proper subset of $\mathbb{C}$, we state many results for $\mathbb{C}$. In other cases, we treat $\mathbb{R}$ and $\mathbb{C}$ separately. To do this efficiently, we use the symbol $\mathbb{F}$ to consistently denote either $\mathbb{R}$ or $\mathbb{C}$.

Let $x \in \mathbb{C}$. Then, $x = y + jz$, where $y, z \in \mathbb{R}$ and $j \triangleq \sqrt{-1}$. Define the *complex conjugate* $\bar{x}$ of x by

$$\bar{x} \triangleq y - jz \quad (2.1.1)$$

and the real part $\operatorname{Re} x$ of x and the imaginary part $\operatorname{Im} x$ of x by

$$\operatorname{Re} x \triangleq \frac{1}{2}(x + \bar{x}) = y \quad (2.1.2)$$

and

$$\operatorname{Im} x \triangleq \frac{1}{2j}(x - \bar{x}) = z. \quad (2.1.3)$$

Furthermore, the *absolute value* $|x|$ of x is defined by

$$|x| \triangleq \sqrt{y^2 + z^2}. \quad (2.1.4)$$

The *closed left half plane* (CLHP), *open left half plane* (OLHP), *closed right half plane* (CRHP), and *open right half plane* (ORHP) are the subsets of $\mathbb{C}$ defined by

$$\text{OLHP} \triangleq \{s \in \mathbb{C}: \operatorname{Re} s < 0\}, \quad (2.1.5)$$

$$\text{CLHP} \triangleq \{s \in \mathbb{C}: \operatorname{Re} s \leq 0\}, \quad (2.1.6)$$

$$\text{ORHP} \triangleq \{s \in \mathbb{C}: \operatorname{Re} s > 0\}, \quad (2.1.7)$$

$$\text{CRHP} \triangleq \{s \in \mathbb{C}: \operatorname{Re} s \geq 0\}. \quad (2.1.8)$$

The imaginary numbers are represented by $j\mathbb{R}$. Note that 0 is both a real number and an imaginary number.

The set $\mathbb{F}^n$ consists of *vectors* x of the form

$$x = \begin{bmatrix} x_{(1)} \\ \vdots \\ x_{(n)} \end{bmatrix}, \quad (2.1.9)$$

where $x_{(1)}, \dots, x_{(n)} \in \mathbb{F}$ are the *components* of x . Hence, the elements of $\mathbb{F}^n$ are *column vectors*. Since $\mathbb{F}^1 = \mathbb{F}$, it follows that every scalar is also a vector. If $x \in \mathbb{R}^n$ and every component of x is nonnegative, then x is *nonnegative*, while, if every component of x is positive, then x is *positive*.

Definition 2.1.1. Let $x, y \in \mathbb{R}^n$, and assume that $x_{(1)} \geq \dots \geq x_{(n)}$ and $y_{(1)} \geq \dots \geq y_{(n)}$. Then, the following terminology is defined:

i) y *weakly majorizes* x if, for all $k = 1, \dots, n$, it follows that

$$\sum_{i=1}^k x_{(i)} \leq \sum_{i=1}^k y_{(i)}. \quad (2.1.10)$$

ii) y *strongly majorizes* x if y weakly majorizes x and

$$\sum_{i=1}^n x_{(i)} = \sum_{i=1}^n y_{(i)}. \quad (2.1.11)$$

Now, assume that x and y are nonnegative. Then, the following terminology is defined:

iii) y *weakly log majorizes* x if, for all $k = 1, \dots, n$, it follows that

$$\prod_{i=1}^k x_{(i)} \leq \prod_{i=1}^k y_{(i)}. \quad (2.1.12)$$

iv) y *strongly log majorizes* x if y weakly log majorizes x and

$$\prod_{i=1}^n x_{(i)} = \prod_{i=1}^n y_{(i)}. \quad (2.1.13)$$

Clearly, if y strongly majorizes x , then y weakly majorizes x , and, if y strongly log majorizes x , then y weakly log majorizes x . Fact 2.21.13 states that, if y weakly log majorizes x , then y weakly majorizes x . Finally, in the notation of Definition 2.1.1, if y majorizes x , then $x_{(1)} \leq y_{(1)}$, while, if y strongly majorizes x , then $y_{(n)} \leq x_{(n)}$.

Definition 2.1.2. Let $\mathcal{S} \subseteq \mathbb{R}^n$, and let $f: \mathcal{S} \mapsto \mathbb{R}$. Then, f is *Schur convex* if, for all $x, y \in \mathcal{S}$ such that y strongly majorizes x , it follows that $f(x) \leq f(y)$. Furthermore, f is *Schur concave* if $-f$ is Schur convex.

If $\alpha \in \mathbb{F}$ and $x \in \mathbb{F}^n$, then $\alpha x \in \mathbb{F}^n$ is given by

$$\alpha x = \begin{bmatrix} \alpha x_{(1)} \\ \vdots \\ \alpha x_{(n)} \end{bmatrix}. \quad (2.1.14)$$

If $x, y \in \mathbb{F}^n$, then x and y are *linearly dependent* if there exists $\alpha \in \mathbb{F}$ such that either $x = \alpha y$ or $y = \alpha x$. Linear dependence for a set of two or more vectors is defined in Section 2.3. Furthermore, vectors add component by component, that is, if $x, y \in \mathbb{F}^n$, then

$$x + y = \begin{bmatrix} x_{(1)} + y_{(1)} \\ \vdots \\ x_{(n)} + y_{(n)} \end{bmatrix}. \quad (2.1.15)$$

Thus, if $\alpha, \beta \in \mathbb{F}$, then the *linear combination* $\alpha x + \beta y$ is given by

$$\alpha x + \beta y = \begin{bmatrix} \alpha x_{(1)} + \beta y_{(1)} \\ \vdots \\ \alpha x_{(n)} + \beta y_{(n)} \end{bmatrix}. \quad (2.1.16)$$

If $x \in \mathbb{R}^n$ and x is nonnegative, then we write $x \geq 0$, while, if x is positive, then we write $x > 0$. If $x, y \in \mathbb{R}^n$, then $x \geq y$ means that $x - y \geq 0$, while $x > y$ means that $x - y > 0$.

The vectors $x_1, \dots, x_m \in \mathbb{F}^n$ placed side by side form the *matrix*

$$A \triangleq [x_1 \quad \cdots \quad x_m], \quad (2.1.17)$$

which has n rows and m columns. The components of the vectors $x_1, \dots, x_m$ are the *entries* of A . We write $A \in \mathbb{F}^{n \times m}$ and say that A has *size* $n \times m$. Since $\mathbb{F}^n = \mathbb{F}^{n \times 1}$, it follows that every vector is also a matrix. Note that $\mathbb{F}^{1 \times 1} = \mathbb{F}^1 = \mathbb{F}$. If $n = m$, then n is the *order* of A , and A is *square*. The i th row of A and the j th column of A are denoted by $\text{row}_i(A)$ and $\text{col}_j(A)$, respectively. Hence,

$$A = \begin{bmatrix} \text{row}_1(A) \\ \vdots \\ \text{row}_n(A) \end{bmatrix} = [\text{col}_1(A) \quad \cdots \quad \text{col}_m(A)]. \quad (2.1.18)$$

The entry $x_{j(i)}$ of A in both the i th row of A and the j th column of A is denoted by $A_{(i,j)}$. Therefore, $x \in \mathbb{F}^n$ can be written as

$$x = \begin{bmatrix} x_{(1)} \\ \vdots \\ x_{(n)} \end{bmatrix} = \begin{bmatrix} x_{(1,1)} \\ \vdots \\ x_{(n,1)} \end{bmatrix}. \quad (2.1.19)$$

Let $A \in \mathbb{F}^{n \times m}$. For $b \in \mathbb{F}^n$, the matrix obtained from A by replacing $\text{col}_i(A)$ with b is denoted by

$$A \stackrel{i}{\leftarrow} b. \quad (2.1.20)$$

Likewise, for $b \in \mathbb{F}^{1 \times m}$, the matrix obtained from A by replacing $\text{row}_i(A)$ with b is denoted by (2.1.20).

Let $A \in \mathbb{F}^{n \times m}$, and let $l \triangleq \min\{n, m\}$. Then, the entries $A_{(i,i)}$ for all $i = 1, \dots, l$ and $A_{(i,j)}$ for all $i \neq j$ are the *diagonal entries* and *off-diagonal entries* of A , respectively. Moreover, for all $i = 1, \dots, l-1$, the entries $A_{(i,i+1)}$ and $A_{(i+1,i)}$ are the *superdiagonal entries* and *subdiagonal entries* of A , respectively. In addition, the entries $A_{(i,l+1-i)}$ for all $i = 1, \dots, l$ are the *reverse-diagonal entries* of A . If the diagonal entries $A_{(1,1)}, \dots, A_{(l,l)}$ of A are real, then the diagonal entries of A are labeled from largest to smallest as

$$d_1(A) \geq \dots \geq d_l(A), \quad (2.1.21)$$

and we define

$$d_{\max}(A) \triangleq d_1(A), \quad d_{\min}(A) \triangleq d_l(A). \quad (2.1.22)$$

Partitioned matrices are of the form

$$\begin{bmatrix} A_{11} & \cdots & A_{1l} \\ \vdots & \ddots & \vdots \\ A_{k1} & \cdots & A_{kl} \end{bmatrix}, \quad (2.1.23)$$

where, for all $i = 1, \dots, k$ and $j = 1, \dots, l$, the *block* A_{ij} of A is a matrix of size $n_i \times m_j$. If $n_i = m_j$ and the diagonal entries of A_{ij} lie on the diagonal of A , then the square matrix A_{ij} is a *diagonally located block*; otherwise, A_{ij} is an *off-diagonally located block*.

Let $A \in \mathbb{F}^{n \times m}$. Then, a *submatrix* of A is formed by deleting rows and columns of A . By convention, A is a submatrix of A . If A is a partitioned matrix, then every block of A is a submatrix of A . A block is thus a submatrix whose entries are entries of adjacent rows and adjacent columns. A submatrix can be specified in terms of the rows and columns that are retained. If like-numbered rows and columns of A are retained, then the resulting square submatrix of A is a *principal submatrix* of A . Every diagonally located block is a principal submatrix. Finally, if rows and columns $1, \dots, j$ of A are retained, then the resulting $j \times j$ submatrix of A is a *leading principal submatrix* of A .

Let $A \in \mathbb{F}^{n \times m}$, and let $\mathcal{S}_1$ and $\mathcal{S}_2$ be subsets of $\{1, \dots, n\}$ and $\{1, \dots, m\}$, respectively. Then, $A_{(\mathcal{S}_1, \mathcal{S}_2)}$ is the $\text{card}(\mathcal{S}_1) \times \text{card}(\mathcal{S}_2)$ submatrix of A formed by retaining the rows of A listed in $\mathcal{S}_1$ and the columns of A listed in $\mathcal{S}_2$. Therefore, $A_{(\mathcal{S}_1^c, \mathcal{S}_2^c)}$ is the $[n - \text{card}(\mathcal{S}_1)] \times [m - \text{card}(\mathcal{S}_2)]$ submatrix of A formed by deleting the rows of A listed in $\mathcal{S}_1$ and the columns of A listed in $\mathcal{S}_2$. If $\mathcal{S} \subseteq \{1, \dots, \min\{n, m\}\}$, then we define $A_{(\mathcal{S})} \triangleq A_{(\mathcal{S}, \mathcal{S})}$, which is a principal submatrix of A .

Matrices of the same size add entry by entry, that is, if $A, B \in \mathbb{F}^{n \times m}$, then, for all $i = 1, \dots, n$ and $j = 1, \dots, m$, $(A + B)_{(i,j)} = A_{(i,j)} + B_{(i,j)}$. Furthermore, for all $i = 1, \dots, n$ and $j = 1, \dots, m$, $(\alpha A)_{(i,j)} = \alpha A_{(i,j)}$ for all $\alpha \in \mathbb{F}$ so that $(\alpha A + \beta B)_{(i,j)} = \alpha A_{(i,j)} + \beta B_{(i,j)}$ for all $\alpha, \beta \in \mathbb{F}$. If $A, B \in \mathbb{F}^{n \times m}$, then A and B are *linearly dependent* if there exists $\alpha \in \mathbb{F}$ such that either $A = \alpha B$ or $B = \alpha A$.

Let $A \in \mathbb{R}^{n \times m}$. If every entry of A is nonnegative, then A is *nonnegative*, which is written as $A \geq 0$. If every entry of A is positive, then A is *positive*, which is written as $A > 0$. If $A, B \in \mathbb{R}^{n \times m}$, then $A \geq B$ means that $A - B \geq 0$, while $A > B$ means that $A - B > 0$.

Let $z \in \mathbb{F}^{1 \times n}$ and $y \in \mathbb{F}^n = \mathbb{F}^{n \times 1}$. Then, the scalar $zy \in \mathbb{F}$ is defined by

$$zy \triangleq \sum_{i=1}^n z_{(1,i)} y_{(i)}. \quad (2.1.24)$$

Now, let $A \in \mathbb{F}^{n \times m}$ and $x \in \mathbb{F}^m$. Then, the matrix-vector product Ax is defined by

$$Ax \triangleq \begin{bmatrix} \text{row}_1(A)x \\ \vdots \\ \text{row}_n(A)x \end{bmatrix}. \quad (2.1.25)$$

It can be seen that Ax is a linear combination of the columns of A , that is,

$$Ax = \sum_{i=1}^m x_{(i)} \text{col}_i(A). \quad (2.1.26)$$

The matrix A can be associated with the function $f: \mathbb{F}^m \mapsto \mathbb{F}^n$ defined by $f(x) \triangleq Ax$ for all $x \in \mathbb{F}^m$. The function $f: \mathbb{F}^m \mapsto \mathbb{F}^n$ is *linear* since, for all $\alpha, \beta \in \mathbb{F}$ and $x, y \in \mathbb{F}^m$, it follows that

$$f(\alpha x + \beta y) = \alpha Ax + \beta Ay. \quad (2.1.27)$$

The function $f: \mathbb{F}^m \mapsto \mathbb{F}^n$ defined by

$$f(x) \triangleq Ax + z, \quad (2.1.28)$$

where $z \in \mathbb{F}^n$, is *affine*.

Theorem 2.1.3. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, and define $f: \mathbb{F}^m \mapsto \mathbb{F}^n$ and $g: \mathbb{F}^l \mapsto \mathbb{F}^m$ by $f(x) \triangleq Ax$ and $g(y) \triangleq By$. Furthermore, define the composition $h \triangleq f \bullet g: \mathbb{F}^l \mapsto \mathbb{F}^n$. Then, for all $y \in \mathbb{F}^l$,

$$h(y) = f[g(y)] = A(By) = (AB)y, \quad (2.1.29)$$

where, for all $i = 1, \dots, n$ and $j = 1, \dots, l$, $AB \in \mathbb{F}^{n \times l}$ is defined by

$$(AB)_{(i,j)} \triangleq \sum_{k=1}^m A_{(i,k)} B_{(k,j)}. \quad (2.1.30)$$

Hence, we write $AB y$ for $(AB)y$ and $A(By)$.

Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, $AB \in \mathbb{F}^{n \times l}$ is the *product* of A and B . The matrices A and B are *conformable*, and the product (2.1.30) defines *matrix multiplication*.

Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, AB can be written as

$$AB = \begin{bmatrix} A \text{col}_1(B) & \cdots & A \text{col}_l(B) \end{bmatrix} = \begin{bmatrix} \text{row}_1(A)B \\ \vdots \\ \text{row}_n(A)B \end{bmatrix}. \quad (2.1.31)$$

Thus, for all $i = 1, \dots, n$ and $j = 1, \dots, l$,

$$(AB)_{(i,j)} = \text{row}_i(A) \text{col}_j(B), \quad (2.1.32)$$

$$\text{col}_j(AB) = A \text{col}_j(B), \quad (2.1.33)$$

$$\text{row}_i(AB) = \text{row}_i(A)B. \quad (2.1.34)$$

For conformable matrices A, B, C , the associative and distributive identities

$$(AB)C = A(BC), \quad (2.1.35)$$

$$A(B + C) = AB + AC, \quad (2.1.36)$$

$$(A + B)C = AC + BC \quad (2.1.37)$$

are valid. Hence, we write ABC for $(AB)C$ and $A(BC)$. Note that (2.1.35) is a special case of (1.2.1).

Let $A, B \in \mathbb{F}^{n \times n}$. Then, the *commutator* $[A, B] \in \mathbb{F}^{n \times n}$ of A and B is the matrix

$$[A, B] \triangleq AB - BA. \quad (2.1.38)$$

The *adjoint operator* $\text{ad}_A: \mathbb{F}^{n \times n} \mapsto \mathbb{F}^{n \times n}$ is defined by

$$\text{ad}_A(X) \triangleq [A, X]. \quad (2.1.39)$$

Let $x, y \in \mathbb{R}^3$. Then, the *cross product* $x \times y \in \mathbb{R}^3$ of x and y is defined by

$$x \times y \triangleq \begin{bmatrix} x_{(2)}y_{(3)} - x_{(3)}y_{(2)} \\ x_{(3)}y_{(1)} - x_{(1)}y_{(3)} \\ x_{(1)}y_{(2)} - x_{(2)}y_{(1)} \end{bmatrix}. \quad (2.1.40)$$

Furthermore, the 3×3 *cross-product matrix* is defined by

$$K(x) \triangleq \begin{bmatrix} 0 & -x_{(3)} & x_{(2)} \\ x_{(3)} & 0 & -x_{(1)} \\ -x_{(2)} & x_{(1)} & 0 \end{bmatrix}. \quad (2.1.41)$$

Note that

$$x \times y = K(x)y. \quad (2.1.42)$$

Multiplication of partitioned matrices is analogous to matrix multiplication with scalar entries. For example, for matrices with conformable blocks,

$$\begin{bmatrix} A & B \end{bmatrix} \begin{bmatrix} C \\ D \end{bmatrix} = AC + BD, \quad (2.1.43)$$

$$\begin{bmatrix} A \\ B \end{bmatrix} C = \begin{bmatrix} AC \\ BC \end{bmatrix}, \quad (2.1.44)$$

$$\begin{bmatrix} A \\ B \end{bmatrix} \begin{bmatrix} C & D \end{bmatrix} = \begin{bmatrix} AC & AD \\ BC & BD \end{bmatrix}, \quad (2.1.45)$$

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} E & F \\ G & H \end{bmatrix} = \begin{bmatrix} AE + BG & AF + BH \\ CE + DG & CF + DH \end{bmatrix}. \quad (2.1.46)$$

The $n \times m$ *zero matrix*, all of whose entries are zero, is written as $0_{n \times m}$. If the dimensions are unambiguous, then we write just 0. Let $x \in \mathbb{F}^m$ and $A \in \mathbb{F}^{n \times m}$. Then, the zero matrix satisfies

$$0_{k \times m} x = 0_k, \quad (2.1.47)$$

$$A 0_{m \times l} = 0_{n \times l}, \quad (2.1.48)$$

$$0_{k \times n} A = 0_{k \times m}. \quad (2.1.49)$$

Another special matrix is the *empty matrix*. For $n \in \mathbb{N}$, the $0 \times n$ empty matrix, which is written as $0_{0 \times n}$, has zero rows and n columns, while the $n \times 0$ empty matrix, which is written as $0_{n \times 0}$, has n rows and zero columns. For $A \in \mathbb{F}^{n \times m}$, where $n, m \in \mathbb{N}$, the empty matrix satisfies the multiplication rules

$$0_{0 \times n} A = 0_{0 \times m} \quad (2.1.50)$$

and

$$A 0_{m \times 0} = 0_{n \times 0}. \quad (2.1.51)$$

Although empty matrices have no entries, it is useful to define the product

$$0_{n \times 0} 0_{0 \times m} \triangleq 0_{n \times m}. \quad (2.1.52)$$

Also, we define

$$I_0 \triangleq \hat{I}_0 \triangleq 0_{0 \times 0}. \quad (2.1.53)$$

For $n, m \in \mathbb{N}$, we define $\mathbb{F}^{0 \times m} \triangleq \{0_{0 \times m}\}$, $\mathbb{F}^{n \times 0} \triangleq \{0_{n \times 0}\}$, and $\mathbb{F}^0 \triangleq \mathbb{F}^{0 \times 1}$. Note that

$$\begin{bmatrix} 0_{n \times 0} & 0_{n \times m} \\ 0_{0 \times 0} & 0_{0 \times m} \end{bmatrix} = 0_{n \times m}. \quad (2.1.54)$$

The empty matrix can be viewed as a useful device for matrices just as 0 is for real numbers and $\emptyset$ is for sets.

The $n \times n$ *identity matrix*, which has 1's on the diagonal and 0's elsewhere, is denoted by I_n or just I . Let $x \in \mathbb{F}^n$ and $A \in \mathbb{F}^{n \times m}$. Then, the identity matrix satisfies

$$I_n x = x \quad (2.1.55)$$

and

$$A I_m = I_n A = A. \quad (2.1.56)$$

Let $A \in \mathbb{F}^{n \times n}$. Then, $A^2 \triangleq AA$ and, for all $k \geq 1$, $A^k \triangleq AA^{k-1}$. We use the convention $A^0 \triangleq I$ even if A is the zero matrix.

The $n \times n$ *reverse identity matrix*, which has 1's on the reverse diagonal and 0's elsewhere, is denoted by $\hat{I}_n$ or just $\hat{I}$. Left multiplication of $A \in \mathbb{F}^{n \times m}$ by $\hat{I}_n$ reverses the rows of A , while right multiplication of A by $\hat{I}_m$ reverses the columns of A . Note that

$$\hat{I}_n^2 = I_n. \quad (2.1.57)$$

2.2 Transpose and Inner Product

A fundamental vector and matrix operation is the transpose. If $x \in \mathbb{F}^n$, then the *transpose* x^T of x is defined to be the row vector

$$x^T \triangleq [x_{(1)} \quad \cdots \quad x_{(n)}] \in \mathbb{F}^{1 \times n}. \quad (2.2.1)$$

Similarly, if $x = [x_{(1,1)} \quad \cdots \quad x_{(1,n)}] \in \mathbb{F}^{1 \times n}$, then

$$x^T = \begin{bmatrix} x_{(1,1)} \\ \vdots \\ x_{(1,n)} \end{bmatrix} \in \mathbb{F}^{n \times 1}. \quad (2.2.2)$$

Let $x, y \in \mathbb{F}^n$. Then, $x^T y \in \mathbb{F}$ is a scalar, and

$$x^T y = y^T x = \sum_{i=1}^n x_{(i)} y_{(i)}. \quad (2.2.3)$$

Note that

$$x^T x = \sum_{i=1}^n x_{(i)}^2. \quad (2.2.4)$$

The vector $e_{i,n} \in \mathbb{R}^n$, or just e_i , has 1 as its i th component and 0's elsewhere. Thus,

$$e_{i,n} = \text{col}_i(I_n). \quad (2.2.5)$$

Let $A \in \mathbb{F}^{n \times m}$. Then, $e_i^T A = \text{row}_i(A)$ and $A e_i = \text{col}_i(A)$. Furthermore, the (i, j) entry of A can be written as

$$A_{(i,j)} = e_i^T A e_j. \quad (2.2.6)$$

The $n \times m$ matrix $E_{i,j,n \times m} \in \mathbb{R}^{n \times m}$, or just $E_{i,j}$, has 1 as its (i, j) entry and 0's elsewhere. Thus,

$$E_{i,j,n \times m} = e_{i,n} e_{j,m}^T. \quad (2.2.7)$$

Note that $E_{i,1,n \times 1} = e_{i,n}$ and

$$I_n = E_{1,1} + \cdots + E_{n,n} = \sum_{i=1}^n e_i e_i^T. \quad (2.2.8)$$

Finally, the $n \times m$ *ones matrix*, all of whose entries are 1, is written as $1_{n \times m}$ or just 1. Thus,

$$1_{n \times m} = \sum_{i,j=1}^{n,m} E_{i,j,n \times m}. \quad (2.2.9)$$

Note that

$$1_{n \times 1} = \sum_{i=1}^n e_{i,n} = \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix} \quad (2.2.10)$$

and

$$1_{n \times m} = 1_{n \times 1} 1_{1 \times m}. \quad (2.2.11)$$

Lemma 2.2.1. Let $x \in \mathbb{R}$. Then, $x^T x = 0$ if and only if $x = 0$.

Let $x, y \in \mathbb{R}^n$. Then, $x^T y \in \mathbb{R}$ is the *inner product* of x and y . Furthermore, x and y are *orthogonal* if $x^T y = 0$. If x and y are nonzero, then the *angle* $\theta \in [0, \pi]$ between x and y is defined by

$$\theta \triangleq \cos^{-1} \frac{x^T y}{\sqrt{x^T x y^T y}}. \quad (2.2.12)$$

Note that x and y are orthogonal if and only if $\theta = \pi/2$.

Let $x \in \mathbb{C}^n$. Then, $x = y + jz$, where $y, z \in \mathbb{R}^n$. Therefore, the transpose x^T of x is given by

$$x^T = y^T + jz^T. \quad (2.2.13)$$

The *complex conjugate* $\bar{x}$ of x is defined by

$$\bar{x} \triangleq y - jz, \quad (2.2.14)$$

while the *complex conjugate transpose* x^* of x is defined by

$$x^* \triangleq \bar{x}^T = y^T - jz^T. \quad (2.2.15)$$

The vectors y and z are the *real* and *imaginary* parts $\operatorname{Re} x$ and $\operatorname{Im} x$ of x , respectively, which are defined by

$$\operatorname{Re} x \triangleq \frac{1}{2}(x + \bar{x}) = y \quad (2.2.16)$$

and

$$\operatorname{Im} x \triangleq \frac{1}{2j}(x - \bar{x}) = z. \quad (2.2.17)$$

Note that

$$x^* x = \sum_{i=1}^n \bar{x}_{(i)} x_{(i)} = \sum_{i=1}^n |x_{(i)}|^2 = \sum_{i=1}^n [y_{(i)}^2 + z_{(i)}^2]. \quad (2.2.18)$$

If $w, x \in \mathbb{C}^n$, then $w^T x = x^T w$.

Lemma 2.2.2. Let $x \in \mathbb{C}^n$. Then, $x^* x = 0$ if and only if $x = 0$.

Let $x, y \in \mathbb{C}^n$. Then, $x^* y \in \mathbb{C}$ is the *inner product* of x and y , which is given by

$$x^* y = \sum_{i=1}^n \bar{x}_{(i)} y_{(i)}. \quad (2.2.19)$$

Furthermore, x and y are *orthogonal* if $x^* y = 0$.

Let $A \in \mathbb{F}^{n \times m}$. Then, the *transpose* $A^T \in \mathbb{F}^{m \times n}$ of A is defined by

$$A^T \triangleq \begin{bmatrix} [\text{row}_1(A)]^T & \cdots & [\text{row}_n(A)]^T \end{bmatrix} = \begin{bmatrix} [\text{col}_1(A)]^T \\ \vdots \\ [\text{col}_m(A)]^T \end{bmatrix}, \quad (2.2.20)$$

that is, $\text{col}_i(A^T) = [\text{row}_i(A)]^T$ for all $i = 1, \dots, n$ and $\text{row}_i(A^T) = [\text{col}_i(A)]^T$ for all $i = 1, \dots, m$. Hence, $(A^T)_{(i,j)} = A_{(j,i)}$ and $(A^T)^T = A$. If $B \in \mathbb{F}^{m \times l}$, then

$$(AB)^T = B^T A^T. \quad (2.2.21)$$

In particular, if $x \in \mathbb{F}^m$, then

$$(Ax)^T = x^T A^T, \quad (2.2.22)$$

while, if, in addition, $y \in \mathbb{F}^n$, then $y^T A x$ is a scalar and

$$y^T A x = (y^T A x)^T = x^T A^T y. \quad (2.2.23)$$

If $B \in \mathbb{F}^{n \times m}$, then, for all $\alpha, \beta \in \mathbb{F}$,

$$(\alpha A + \beta B)^T = \alpha A^T + \beta B^T. \quad (2.2.24)$$

Let $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$. Then, the matrix $xy^T \in \mathbb{F}^{n \times m}$ is the *outer product* of x and y . The outer product xy^T is nonzero if and only if both x and y are nonzero.

The *trace* of a square matrix $A \in \mathbb{F}^{n \times n}$, denoted by $\text{tr } A$, is defined to be the sum of its diagonal entries, that is,

$$\text{tr } A \triangleq \sum_{i=1}^n A_{(i,i)}. \quad (2.2.25)$$

Note that

$$\text{tr } A = \text{tr } A^T. \quad (2.2.26)$$

Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then, AB and BA are square,

$$\text{tr } AB = \text{tr } BA = \text{tr } A^T B^T = \text{tr } B^T A^T = \sum_{i,j=1}^{n,m} A_{(i,j)} B_{(j,i)}, \quad (2.2.27)$$

and

$$\text{tr } AA^T = \text{tr } A^T A = \sum_{i,j=1}^{n,m} A_{(i,j)}^2. \quad (2.2.28)$$

Furthermore, if $n = m$, then, for all $\alpha, \beta \in \mathbb{F}$,

$$\text{tr}(\alpha A + \beta B) = \alpha \text{tr } A + \beta \text{tr } B. \quad (2.2.29)$$

Lemma 2.2.3. Let $A \in \mathbb{R}^{n \times m}$. Then, $\text{tr } A^T A = 0$ if and only if $A = 0$.

Let $A, B \in \mathbb{R}^{n \times m}$. Then, the *inner product* of A and B is $\text{tr } A^T B$. Furthermore, A is *orthogonal* to B if $\text{tr } A^T B = 0$.

Let $C \in \mathbb{C}^{n \times m}$. Then, $C = A + jB$, where $A, B \in \mathbb{R}^{n \times m}$. Therefore, the transpose C^T of C is given by

$$C^T = A^T + jB^T. \quad (2.2.30)$$

The *complex conjugate* $\overline{C}$ of C is

$$\overline{C} \triangleq A - jB, \quad (2.2.31)$$

while the *complex conjugate transpose* C^* of C is

$$C^* \triangleq \overline{C}^T = A^T - jB^T. \quad (2.2.32)$$

Note that $\overline{C} = C$ if and only if $B = 0$, and that

$$(C^T)^T = \overline{\overline{C}} = (C^*)^* = C. \quad (2.2.33)$$

The matrices A and B are the real and imaginary parts $\operatorname{Re} C$ and $\operatorname{Im} C$ of C , respectively, which are denoted by

$$\operatorname{Re} C \triangleq \frac{1}{2}(C + \overline{C}) = A \quad (2.2.34)$$

and

$$\operatorname{Im} C \triangleq \frac{1}{2j}(C - \overline{C}) = B. \quad (2.2.35)$$

If C is square, then

$$\operatorname{tr} C = \operatorname{tr} A + j \operatorname{tr} B \quad (2.2.36)$$

and

$$\operatorname{tr} C = \operatorname{tr} C^T = \overline{\operatorname{tr} \overline{C}} = \overline{\operatorname{tr} C^*}. \quad (2.2.37)$$

If $\mathcal{S} \subseteq \mathbb{C}^{n \times m}$, then

$$\overline{\mathcal{S}} \triangleq \{\overline{A}: A \in \mathcal{S}\}. \quad (2.2.38)$$

If $\mathcal{S}$ is a multiset with elements in $\mathbb{C}^{n \times m}$, then

$$\overline{\mathcal{S}} = \{\overline{A}: A \in \mathcal{S}\}_{\text{ms}}. \quad (2.2.39)$$

Let $A \in \mathbb{F}^{n \times n}$. Then, for all $k \in \mathbb{N}$,

$$A^{kT} \triangleq (A^k)^T = (A^T)^k, \quad (2.2.40)$$

$$\overline{A^k} = \overline{A}^k, \quad (2.2.41)$$

and

$$A^{k*} \triangleq (A^k)^* = (A^*)^k. \quad (2.2.42)$$

Lemma 2.2.4. Let $A \in \mathbb{C}^{n \times m}$. Then, $\operatorname{tr} A^*A = 0$ if and only if $A = 0$.

Let $A, B \in \mathbb{C}^{n \times m}$. Then, the *inner product* of A and B is $\operatorname{tr} A^*B$. Furthermore, A is *orthogonal* to B if $\operatorname{tr} A^*B = 0$.

If $A, B \in \mathbb{C}^{n \times m}$, then, for all $\alpha, \beta \in \mathbb{C}$,

$$(\alpha A + \beta B)^* = \overline{\alpha} A^* + \overline{\beta} B^*, \quad (2.2.43)$$

while, if $A \in \mathbb{C}^{n \times m}$ and $B \in \mathbb{C}^{m \times l}$, then

$$\overline{AB} = \overline{A}\overline{B} \quad (2.2.44)$$

and

$$(AB)^* = B^*A^*. \quad (2.2.45)$$

In particular, if $A \in \mathbb{C}^{n \times m}$ and $x \in \mathbb{C}^m$, then

$$(Ax)^* = x^*A^*, \quad (2.2.46)$$

while, if, in addition, $y \in \mathbb{C}^n$, then

$$y^*Ax = (y^*Ax)^T = x^T A^T \overline{y} \quad (2.2.47)$$

and

$$(y^*Ax)^* = (\overline{y^*Ax})^T = (y^T \overline{Ax})^T = x^*A^*y. \quad (2.2.48)$$

For $A \in \mathbb{F}^{n \times m}$, define the *reverse transpose* of A by

$$A^{\hat{T}} \triangleq \hat{I}_m A^T \hat{I}_n \quad (2.2.49)$$

and the *reverse complex conjugate transpose* of A by

$$A^* \triangleq \hat{I}_m A^* \hat{I}_n. \quad (2.2.50)$$

For example,

$$\begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}^{\hat{T}} = \begin{bmatrix} 6 & 3 \\ 5 & 2 \\ 4 & 1 \end{bmatrix}. \quad (2.2.51)$$

In general,

$$(A^*)^{\hat{*}} = (A^{\hat{*}})^* = (A^T)^{\hat{T}} = (A^{\hat{T}})^T = \hat{I}_n A \hat{I}_m \quad (2.2.52)$$

and

$$(A^{\hat{*}})^{\hat{*}} = (A^{\hat{T}})^{\hat{T}} = A. \quad (2.2.53)$$

Note that, if $B \in \mathbb{F}^{m \times l}$, then

$$(AB)^{\hat{*}} = B^{\hat{*}}A^{\hat{*}} \quad (2.2.54)$$

and

$$(AB)^{\hat{T}} = B^{\hat{T}}A^{\hat{T}}. \quad (2.2.55)$$

For $x \in \mathbb{F}^m$ and $A \in \mathbb{F}^{n \times m}$, every component of x and every entry of A can be replaced by its absolute value to obtain $|x| \in \mathbb{R}^m$ and $|A| \in \mathbb{R}^{n \times m}$ defined by

$$|x|_{(i)} \triangleq |x_{(i)}| \quad (2.2.56)$$

for all $i = 1, \dots, n$ and

$$|A|_{(i,j)} \triangleq |A_{(i,j)}| \quad (2.2.57)$$

for all $i = 1, \dots, n$ and $j = 1, \dots, m$. Note that

$$|Ax| \leq |A||x|. \quad (2.2.58)$$

Furthermore, if $B \in \mathbb{F}^{m \times l}$, then

$$|AB| \leq |A||B|. \quad (2.2.59)$$

For $x \in \mathbb{R}^n$ and $A \in \mathbb{R}^{n \times m}$, every component of x and every entry of A can be replaced by its sign to obtain $\text{sign } x \in \mathbb{R}^n$ and $\text{sign } A \in \mathbb{R}^{n \times m}$ defined by

$$(\text{sign } x)_{(i)} \triangleq \text{sign } x_{(i)} \quad (2.2.60)$$

for all $i = 1, \dots, n$, and

$$(\text{sign } A)_{(i,j)} \triangleq \text{sign } A_{(i,j)} \quad (2.2.61)$$

for all $i = 1, \dots, n$ and $j = 1, \dots, m$.

2.3 Convex Sets, Cones, and Subspaces

The definitions in this section are stated for subsets of $\mathbb{F}^n$. All of these definitions apply to subsets of $\mathbb{F}^{n \times m}$.

Let $\mathcal{S} \subseteq \mathbb{F}^n$. If $\alpha \in \mathbb{F}$, then $\alpha\mathcal{S} \triangleq \{\alpha x: x \in \mathcal{S}\}$ and, if $y \in \mathbb{F}^n$, then $y + \mathcal{S} = \mathcal{S} + y \triangleq \{y + x: x \in \mathcal{S}\}$. We write $-\mathcal{S}$ for $(-1)\mathcal{S}$. The set $\mathcal{S}$ is *symmetric* if $\mathcal{S} = -\mathcal{S}$, that is, $x \in \mathcal{S}$ if and only if $-x \in \mathcal{S}$. For $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ define $\mathcal{S}_1 + \mathcal{S}_2 \triangleq \{x + y: x \in \mathcal{S}_1 \text{ and } y \in \mathcal{S}_2\}$. Note that, for all $\alpha_1, \alpha_2 \in \mathbb{F}$, $(\alpha + \beta)\mathcal{S} \subseteq \alpha\mathcal{S} + \beta\mathcal{S}$. Trivially, $\mathcal{S} + \emptyset = \emptyset$.

If $x, y \in \mathbb{F}^n$ and $\alpha \in [0, 1]$, then $\alpha x + (1 - \alpha)y$ is a *convex combination* of x and y with *barycentric coordinates* α and $1 - \alpha$. The set $\mathcal{S} \subseteq \mathbb{F}^n$ is *convex* if, for all $x, y \in \mathcal{S}$, every convex combination of x and y is an element of $\mathcal{S}$. Trivially, the empty set is convex.

Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then, $\mathcal{S}$ is a *cone* if, for all $x \in \mathcal{S}$ and all $\alpha > 0$, the vector αx is an element of $\mathcal{S}$. Now, assume that $\mathcal{S}$ is a cone. Then, $\mathcal{S}$ is *pointed* if $0 \in \mathcal{S}$, while $\mathcal{S}$ is *blunt* if $0 \notin \mathcal{S}$. Furthermore, $\mathcal{S}$ is *one-sided* if $x, -x \in \mathcal{S}$ implies that $x = 0$. Hence, $\mathcal{S}$ is one-sided if and only if $\mathcal{S} \cap -\mathcal{S} \subseteq \{0\}$. Furthermore, $\mathcal{S}$ is a *convex cone* if it is convex. Trivially, the empty set is a convex cone.

Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then, $\mathcal{S}$ is a *subspace* if, for all $x, y \in \mathcal{S}$ and $\alpha, \beta \in \mathbb{F}$, the vector $\alpha x + \beta y$ is an element of $\mathcal{S}$. Note that, if $\{x_1, \dots, x_r\} \subset \mathbb{F}^n$, then the set $\{\sum_{i=1}^r \alpha_i x_i: \alpha_1, \dots, \alpha_r \in \mathbb{F}\}$ is a subspace. In addition, $\mathcal{S}$ is an *affine subspace* if there exists a vector $z \in \mathbb{F}^n$ such that $\mathcal{S} + z$ is a subspace. Affine subspaces $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ are *parallel* if there exists a vector $z \in \mathbb{F}^n$ such that $\mathcal{S}_1 + z = \mathcal{S}_2$. If $\mathcal{S}$ is an affine subspace, then there exists a unique subspace parallel to $\mathcal{S}$. Trivially, the empty set is a subspace and an affine subspace.

Let $\mathcal{S} \subseteq \mathbb{F}^n$. The *convex hull* of $\mathcal{S}$, denoted by $\text{co } \mathcal{S}$, is the smallest convex set containing $\mathcal{S}$. Hence, $\text{co } \mathcal{S}$ is the intersection of all convex subsets of $\mathbb{F}^n$ that contain $\mathcal{S}$. The *conical hull* of $\mathcal{S}$, denoted by $\text{cone } \mathcal{S}$, is the smallest cone in $\mathbb{F}^n$ containing $\mathcal{S}$, while the *convex conical hull* of $\mathcal{S}$, denoted by $\text{coco } \mathcal{S}$, is the smallest convex cone in $\mathbb{F}^n$ containing $\mathcal{S}$. If $\mathcal{S}$ has a finite number of elements, then $\text{co } \mathcal{S}$ is a *polytope*

and $\text{coco } \mathcal{S}$ is a *polyhedral convex cone*. The *span* of $\mathcal{S}$, denoted by $\text{span } \mathcal{S}$, is the smallest subspace in $\mathbb{F}^n$ containing $\mathcal{S}$, while, if $\mathcal{S}$ is nonempty, then the *affine hull* of $\mathcal{S}$, denoted by $\text{aff } \mathcal{S}$, is the smallest affine subspace in $\mathbb{F}^n$ containing $\mathcal{S}$. Note that $\mathcal{S}$ is convex if and only if $\mathcal{S} = \text{co } \mathcal{S}$, while similar statements hold for $\text{cone } \mathcal{S}$, $\text{coco } \mathcal{S}$, $\text{span } \mathcal{S}$, and $\text{aff } \mathcal{S}$. Trivially, $\text{co } \emptyset = \text{cone } \emptyset = \text{coco } \emptyset = \text{span } \emptyset = \text{aff } \emptyset = \emptyset$.

Let $x_1, \dots, x_r \in \mathbb{F}^n$. Then, $x_1, \dots, x_r$ are *linearly independent* if $\alpha_1, \dots, \alpha_r \in \mathbb{F}$ and

$$\sum_{i=1}^r \alpha_i x_i = 0 \quad (2.3.1)$$

imply that $\alpha_1 = \alpha_2 = \dots = \alpha_r = 0$. Clearly, $x_1, \dots, x_r$ is linearly independent if and only if $\overline{x_1}, \dots, \overline{x_r}$ are linearly independent. If $x_1, \dots, x_r$ are not linearly independent, then $x_1, \dots, x_r$ are *linearly dependent*. Note that $0_{n \times 1}$ is linearly dependent.

Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is not empty. If $\mathcal{S}$ is not equal to $\{0_{n \times 1}\}$, then there exist $r \geq 1$ vectors $x_1, \dots, x_r \in \mathbb{F}^n$ such that $x_1, \dots, x_r$ are linearly independent over $\mathbb{F}$ and such that $\text{span}\{x_1, \dots, x_r\} = \mathcal{S}$. The set of vectors $\{x_1, \dots, x_r\}$ is a *basis* for $\mathcal{S}$. The positive integer r , which is the *dimension* $\dim \mathcal{S}$ of $\mathcal{S}$, is uniquely defined. We define $\dim\{0_{n \times 1}\} = 0$. If $\mathcal{S}$ is an affine subspace, then the *dimension* $\dim \mathcal{S}$ of $\mathcal{S}$ is the dimension of the subspace parallel to $\mathcal{S}$. If $\mathcal{S}$ is not an affine subspace, then the *dimension* $\dim \mathcal{S}$ of $\mathcal{S}$ is the dimension of $\text{aff } \mathcal{S}$. We define $\dim \emptyset \triangleq -\infty$.

Let $x_1, \dots, x_{n+1} \in \mathbb{R}^n$, and define $\mathcal{S} \triangleq \text{co}\{x_1, \dots, x_{n+1}\}$. The set $\mathcal{S}$ is a *simplex* if $\dim \mathcal{S} = n$.

The following result is the *subspace dimension theorem*.

Theorem 2.3.1. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then,

$$\dim(\mathcal{S}_1 + \mathcal{S}_2) + \dim(\mathcal{S}_1 \cap \mathcal{S}_2) = \dim \mathcal{S}_1 + \dim \mathcal{S}_2. \quad (2.3.2)$$

Proof. See [630, p. 227]. $\square$

Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then, $\mathcal{S}_1$ and $\mathcal{S}_2$ are *complementary* if $\mathcal{S}_1 + \mathcal{S}_2 = \mathbb{F}^n$ and $\mathcal{S}_1 \cap \mathcal{S}_2 = \{0\}$. In this case, we say that $\mathcal{S}_1$ is complementary to $\mathcal{S}_2$, or vice versa.

Corollary 2.3.2. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces, and consider the following conditions:

- i) $\dim(\mathcal{S}_1 + \mathcal{S}_2) = n$.
- ii) $\mathcal{S}_1 \cap \mathcal{S}_2 = \{0\}$.
- iii) $\dim \mathcal{S}_1 + \dim \mathcal{S}_2 = n$.
- iv) $\mathcal{S}_1$ and $\mathcal{S}_2$ are complementary subspaces.

Then,

$$[i), ii)] \iff [i), iii)] \iff [ii), iii)] \iff [i), ii), iii)] \iff [iv)].$$

Let $\mathcal{S} \subseteq \mathbb{F}^n$ be nonempty. Then, the *orthogonal complement* $\mathcal{S}^\perp$ of $\mathcal{S}$ is defined by

$$\mathcal{S}^\perp \triangleq \{x \in \mathbb{F}^n: x^*y = 0 \text{ for all } y \in \mathcal{S}\}. \quad (2.3.3)$$

The orthogonal complement $\mathcal{S}^\perp$ of $\mathcal{S}$ is a subspace even if $\mathcal{S}$ is not.

Let $y \in \mathbb{F}^n$ be nonzero. Then, the subspace $\{y\}^\perp$, whose dimension is $n-1$, is a *hyperplane*. Furthermore, $\mathcal{S}$ is an *affine hyperplane* if there exists a vector $z \in \mathbb{F}^n$ such that $\mathcal{S} + z$ is a hyperplane. The set $\{x \in \mathbb{F}^n: \operatorname{Re} x^*y \leq 0\}$ is a *closed half space*, while the set $\{x \in \mathbb{F}^n: \operatorname{Re} x^*y < 0\}$ is an *open half space*. Finally, $\mathcal{S}$ is an *affine (closed, open) half space* if there exists a vector $z \in \mathbb{F}^n$ such that $\mathcal{S} + z$ is a (closed, open) half space.

Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then,

$$\operatorname{polar} \mathcal{S} \triangleq \{x \in \mathbb{F}^n: \operatorname{Re} x^*y \leq 1 \text{ for all } y \in \mathcal{S}\} \quad (2.3.4)$$

is the *polar* of $\mathcal{S}$. Note that $\operatorname{polar} \mathcal{S}$ is a convex set. Furthermore,

$$\operatorname{polar} \mathcal{S} = \operatorname{polar} \operatorname{co} \mathcal{S}. \quad (2.3.5)$$

Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then,

$$\operatorname{dcone} \mathcal{S} \triangleq \{x \in \mathbb{F}^n: \operatorname{Re} x^*y \leq 0 \text{ for all } y \in \mathcal{S}\} \quad (2.3.6)$$

is the *dual cone* of $\mathcal{S}$. Note that $\operatorname{dcone} \mathcal{S}$ is a pointed convex cone. Furthermore,

$$\operatorname{dcone} \mathcal{S} = \operatorname{dcone} \operatorname{cone} \mathcal{S} = \operatorname{dcone} \operatorname{coco} \mathcal{S}. \quad (2.3.7)$$

Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then, $\mathcal{S}_1$ and $\mathcal{S}_2$ are *orthogonally complementary* if $\mathcal{S}_1$ and $\mathcal{S}_2$ are complementary and $x^*y = 0$ for all $x \in \mathcal{S}_1$ and $y \in \mathcal{S}_2$.

Proposition 2.3.3. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then, $\mathcal{S}_1$ and $\mathcal{S}_2$ are orthogonally complementary if and only if $\mathcal{S}_1 = \mathcal{S}_2^\perp$.

For the next result, note that “ $\subset$ ” indicates proper inclusion.

Lemma 2.3.4. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces such that $\mathcal{S}_1 \subseteq \mathcal{S}_2$. Then, $\mathcal{S}_1 \subset \mathcal{S}_2$ if and only if $\dim \mathcal{S}_1 < \dim \mathcal{S}_2$. Equivalently, $\mathcal{S}_1 = \mathcal{S}_2$ if and only if $\dim \mathcal{S}_1 = \dim \mathcal{S}_2$.

The following result provides constructive characterizations of $\operatorname{co} \mathcal{S}$, $\operatorname{cone} \mathcal{S}$, $\operatorname{coco} \mathcal{S}$, $\operatorname{span} \mathcal{S}$, and $\operatorname{aff} \mathcal{S}$.

Theorem 2.3.5. Let $\mathcal{S} \subseteq \mathbb{R}^n$ be nonempty. Then,

$$\operatorname{co} \mathcal{S} = \bigcup_{k \in \mathbb{P}} \left\{ \sum_{i=1}^k \alpha_i x_i: \alpha_i \geq 0, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^k \alpha_i = 1 \right\} \quad (2.3.8)$$

$$= \left\{ \sum_{i=1}^{n+1} \alpha_i x_i: \alpha_i \geq 0, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^{n+1} \alpha_i = 1 \right\}, \quad (2.3.9)$$

$$\text{cone } \mathcal{S} = \{\alpha x: x \in \mathcal{S} \text{ and } \alpha > 0\}, \quad (2.3.10)$$

$$\text{coco } \mathcal{S} = \bigcup_{k \in \mathbb{P}} \left\{ \sum_{i=1}^k \alpha_i x_i: \alpha_i \geq 0, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^k \alpha_i > 0 \right\} \quad (2.3.11)$$

$$= \left\{ \sum_{i=1}^{n+1} \alpha_i x_i: \alpha_i \geq 0, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^n \alpha_i > 0 \right\}, \quad (2.3.12)$$

$$\text{span } \mathcal{S} = \bigcup_{k \in \mathbb{P}} \left\{ \sum_{i=1}^k \alpha_i x_i: \alpha_i \in \mathbb{R} \text{ and } x_i \in \mathcal{S} \right\} \quad (2.3.13)$$

$$= \left\{ \sum_{i=1}^n \alpha_i x_i: \alpha_i \in \mathbb{R} \text{ and } x_i \in \mathcal{S} \right\}, \quad (2.3.14)$$

$$\text{aff } \mathcal{S} = \bigcup_{k \in \mathbb{P}} \left\{ \sum_{i=1}^k \alpha_i x_i: \alpha_i \in \mathbb{R}, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^k \alpha_i = 1 \right\} \quad (2.3.15)$$

$$= \left\{ \sum_{i=1}^{n+1} \alpha_i x_i: \alpha_i \in \mathbb{R}, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^{n+1} \alpha_i = 1 \right\}. \quad (2.3.16)$$

Now, let $\mathcal{S} \subseteq \mathbb{C}^n$. Then,

$$\text{co } \mathcal{S} = \bigcup_{k \in \mathbb{P}} \left\{ \sum_{i=1}^k \alpha_i x_i: \alpha_i \geq 0, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^k \alpha_i = 1 \right\} \quad (2.3.17)$$

$$= \left\{ \sum_{i=1}^{2n+1} \alpha_i x_i: \alpha_i \geq 0, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^{2n+1} \alpha_i = 1 \right\}, \quad (2.3.18)$$

$$\text{cone } \mathcal{S} = \{\alpha x: x \in \mathcal{S} \text{ and } \alpha > 0\}, \quad (2.3.19)$$

$$\text{coco } \mathcal{S} = \bigcup_{k \in \mathbb{P}} \left\{ \sum_{i=1}^k \alpha_i x_i: \alpha_i \geq 0, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^k \alpha_i > 0 \right\} \quad (2.3.20)$$

$$= \left\{ \sum_{i=1}^{2n+1} \alpha_i x_i: \alpha_i \geq 0, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^{2n} \alpha_i > 0 \right\}, \quad (2.3.21)$$

$$\text{span } \mathcal{S} = \bigcup_{k \in \mathbb{P}} \left\{ \sum_{i=1}^k \alpha_i x_i: \alpha_i \in \mathbb{C} \text{ and } x_i \in \mathcal{S} \right\} \quad (2.3.22)$$

$$= \left\{ \sum_{i=1}^n \alpha_i x_i: \alpha_i \in \mathbb{C} \text{ and } x_i \in \mathcal{S} \right\}, \quad (2.3.23)$$

$$\text{aff } \mathcal{S} = \bigcup_{k \in \mathbb{P}} \left\{ \sum_{i=1}^k \alpha_i x_i : \alpha_i \in \mathbb{C}, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^k \alpha_i = 1 \right\} \quad (2.3.24)$$

$$= \left\{ \sum_{i=1}^{n+1} \alpha_i x_i : \alpha_i \in \mathbb{C}, x_i \in \mathcal{S}, \text{ and } \sum_{i=1}^{n+1} \alpha_i = 1 \right\}. \quad (2.3.25)$$

Proof. Result (2.3.8) is immediate, while (2.3.9) is proved in [879, p. 17]. Furthermore, (2.3.10) is immediate. Next, note that, since $\text{coco } \mathcal{S} = \text{co cone } \mathcal{S}$, it follows that (2.3.8) and (2.3.10) imply (2.3.12) with n replaced by $n + 1$. However, every element of $\text{coco } \mathcal{S}$ lies in the convex hull of $n + 1$ points one of which is the origin. It thus follows that we can set $x_{n+1} = 0$, which yields (2.3.12). Similar arguments yield (2.3.14). Finally, note that all vectors of the form $x_1 + \beta(x_2 - x_1)$, where $x_1, x_2 \in \mathcal{S}$ and $\beta \in \mathbb{R}$, are elements of $\text{aff } \mathcal{S}$. Forming the convex hull of these vectors yields (2.3.16). $\square$

The following result shows that cones can be used to induce relations on $\mathbb{F}^n$.

Proposition 2.3.6. Let $\mathcal{S} \subseteq \mathbb{F}^n$ be a cone and, for $x, y \in \mathbb{F}^n$, let $x \leq y$ denote the relation $y - x \in \mathcal{S}$. Then, the following statements hold:

- i) “ $\leq$ ” is reflexive if and only if $\mathcal{S}$ is a pointed cone.
- ii) “ $\leq$ ” is antisymmetric if and only if $\mathcal{S}$ is a one-sided cone.
- iii) “ $\leq$ ” is symmetric if and only if $\mathcal{S}$ is a symmetric cone.
- iv) “ $\leq$ ” is transitive if and only if $\mathcal{S}$ is a convex cone.

Proof. The proofs of i), ii), and iii) are immediate. To prove iv), suppose that “ $\leq$ ” is transitive, and let $x, y \in \mathcal{S}$ so that $0 \leq \alpha x \leq \alpha x + (1 - \alpha)y$ for all $\alpha \in (0, 1]$. Hence, $\alpha x + (1 - \alpha)y \in \mathcal{S}$ for all $\alpha \in (0, 1]$, and thus $\mathcal{S}$ is convex. Conversely, suppose that $\mathcal{S}$ is a convex cone, and assume that $x \leq y$ and $y \leq z$. Then, $y - x \in \mathcal{S}$ and $z - y \in \mathcal{S}$ imply that $z - x = 2[\frac{1}{2}(y - x) + \frac{1}{2}(z - y)] \in \mathcal{S}$. Hence, $x \leq z$, and thus “ $\leq$ ” is transitive. $\square$

2.4 Range and Null Space

Two key features of a matrix $A \in \mathbb{F}^{n \times m}$ are its range and null space, denoted by $\mathcal{R}(A)$ and $\mathcal{N}(A)$, respectively. The *range* of A is defined by

$$\mathcal{R}(A) \triangleq \{Ax : x \in \mathbb{F}^m\}. \quad (2.4.1)$$

Note that $\mathcal{R}(0_{n \times 0}) = \{0_{n \times 1}\}$ and $\mathcal{R}(0_{0 \times m}) = \{0_{0 \times 1}\}$. Letting α_i denote $x_{(i)}$, it can be seen that

$$\mathcal{R}(A) = \left\{ \sum_{i=1}^m \alpha_i \text{col}_i(A) : \alpha_1, \dots, \alpha_m \in \mathbb{F} \right\}, \quad (2.4.2)$$

which shows that $\mathcal{R}(A)$ is a subspace of $\mathbb{F}^n$. It thus follows from Theorem 2.3.5 that

$$\mathcal{R}(A) = \text{span} \{ \text{col}_1(A), \dots, \text{col}_m(A) \}. \quad (2.4.3)$$

By viewing A as a function from $\mathbb{F}^m$ into $\mathbb{F}^n$, we can write $\mathcal{R}(A) = A\mathbb{F}^m$.

The *null space* of $A \in \mathbb{F}^{n \times m}$ is defined by

$$\mathcal{N}(A) \triangleq \{x \in \mathbb{F}^m: Ax = 0\}. \quad (2.4.4)$$

Note that $\mathcal{N}(0_{n \times 0}) = \mathbb{F}^0 = \{0_{0 \times 1}\}$ and $\mathcal{N}(0_{0 \times m}) = \mathbb{F}^m$. Equivalently,

$$\mathcal{N}(A) = \left\{x \in \mathbb{F}^m: x^T [\text{row}_i(A)]^T = 0 \text{ for all } i = 1, \dots, n\right\} \quad (2.4.5)$$

$$= \left\{[\text{row}_1(A)]^T, \dots, [\text{row}_n(A)]^T\right\}^\perp, \quad (2.4.6)$$

which shows that $\mathcal{N}(A)$ is a subspace of $\mathbb{F}^m$. Note that, if $\alpha \in \mathbb{F}$ is nonzero, then $\mathcal{R}(\alpha A) = \mathcal{R}(A)$ and $\mathcal{N}(\alpha A) = \mathcal{N}(A)$. Finally, if $\mathbb{F} = \mathbb{C}$, then $\mathcal{R}(A)$ and $\mathcal{R}(\overline{A})$ are not necessarily identical. For example, let $A \triangleq \begin{bmatrix} 1 \\ j \end{bmatrix}$.

Let $A \in \mathbb{F}^{n \times n}$, and let $\mathcal{S} \subseteq \mathbb{F}^n$ be a subspace. Then, $\mathcal{S}$ is an *invariant subspace* of A if $A\mathcal{S} \subseteq \mathcal{S}$. Note that $A\mathcal{R}(A) \subseteq A\mathbb{F}^n = \mathcal{R}(A)$ and $A\mathcal{N}(A) = \{0_n\} \subseteq \mathcal{N}(A)$. Hence, $\mathcal{R}(A)$ and $\mathcal{N}(A)$ are invariant subspaces of A .

If $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, then it is easy to see that

$$\mathcal{R}(AB) = A\mathcal{R}(B). \quad (2.4.7)$$

Hence, the following result is not surprising.

Lemma 2.4.1. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $C \in \mathbb{F}^{k \times n}$. Then,

$$\mathcal{R}(AB) \subseteq \mathcal{R}(A) \quad (2.4.8)$$

and

$$\mathcal{N}(A) \subseteq \mathcal{N}(CA). \quad (2.4.9)$$

Proof. Since $\mathcal{R}(B) \subseteq \mathbb{F}^m$, it follows that $\mathcal{R}(AB) = A\mathcal{R}(B) \subseteq A\mathbb{F}^m = \mathcal{R}(A)$. Furthermore, $y \in \mathcal{N}(A)$ implies that $Ay = 0$, and thus $CAy = 0$. $\square$

Corollary 2.4.2. Let $A \in \mathbb{F}^{n \times n}$, and let $k \geq 1$. Then,

$$\mathcal{R}(A^k) \subseteq \mathcal{R}(A) \quad (2.4.10)$$

and

$$\mathcal{N}(A) \subseteq \mathcal{N}(A^k). \quad (2.4.11)$$

Although $\mathcal{R}(AB) \subseteq \mathcal{R}(A)$ for arbitrary conformable matrices A, B , we now show that equality holds in the special case $B = A^*$. This result, along with others, is the subject of the following basic theorem.

Theorem 2.4.3. Let $A \in \mathbb{F}^{n \times m}$. Then, the following identities hold:

- i) $\mathcal{R}(A)^\perp = \mathcal{N}(A^*)$.
- ii) $\mathcal{R}(A) = \mathcal{R}(AA^*)$.
- iii) $\mathcal{N}(A) = \mathcal{N}(A^*A)$.

Proof. To prove i), we first show that $\mathcal{R}(A)^\perp \subseteq \mathcal{N}(A^*)$. Let $x \in \mathcal{R}(A)^\perp$. Then, $x^*z = 0$ for all $z \in \mathcal{R}(A)$. Hence, $x^*Ay = 0$ for all $y \in \mathbb{F}^m$. Equivalently,

$y^*A^*x = 0$ for all $y \in \mathbb{R}^m$. Letting $y = A^*x$, it follows that $x^*AA^*x = 0$. Now, Lemma 2.2.2 implies that $A^*x = 0$. Thus, $x \in \mathcal{N}(A^*)$. Conversely, let us show that $\mathcal{N}(A^*) \subseteq \mathcal{R}(A)^\perp$. Letting $x \in \mathcal{N}(A^*)$, it follows that $A^*x = 0$, and, hence, $y^*A^*x = 0$ for all $y \in \mathbb{R}^m$. Equivalently, $x^*Ay = 0$ for all $y \in \mathbb{R}^m$. Hence, $x^*z = 0$ for all $z \in \mathcal{R}(A)$. Thus, $x \in \mathcal{R}(A)^\perp$, which proves *i*).

To prove *ii*), note that Lemma 2.4.1 with $B = A^*$ implies that $\mathcal{R}(AA^*) \subseteq \mathcal{R}(A)$. To show that $\mathcal{R}(A) \subseteq \mathcal{R}(AA^*)$, let $x \in \mathcal{R}(A)$, and suppose that $x \notin \mathcal{R}(AA^*)$. Then, it follows from Proposition 2.3.3 that $x = x_1 + x_2$, where $x_1 \in \mathcal{R}(AA^*)$ and $x_2 \in \mathcal{R}(AA^*)^\perp$ with $x_2 \neq 0$. Thus, $x_2^*AA^*y = 0$ for all $y \in \mathbb{R}^n$, and setting $y = x_2$ yields $x_2^*AA^*x_2 = 0$. Hence, Lemma 2.2.2 implies that $A^*x_2 = 0$, so that, by *i*), $x_2 \in \mathcal{N}(A^*) = \mathcal{R}(A)^\perp$. Since $x \in \mathcal{R}(A)$, it follows that $0 = x_2^*x = x_2^*x_1 + x_2^*x_2$. However, $x_2^*x_1 = 0$ so that $x_2^*x_2 = 0$ and $x_2 = 0$, which is a contradiction. This proves *ii*).

To prove *iii*), note that *ii*) with A replaced by A^* implies that $\mathcal{R}(A^*A)^\perp = \mathcal{R}(A^*)^\perp$. Furthermore, replacing A by A^* in *i*) yields $\mathcal{R}(A^*)^\perp = \mathcal{N}(A)$. Hence, $\mathcal{N}(A) = \mathcal{R}(A^*A)^\perp$. Now, *i*) with A replaced by A^*A implies that $\mathcal{R}(A^*A)^\perp = \mathcal{N}(A^*A)$. Hence, $\mathcal{N}(A) = \mathcal{N}(A^*A)$, which proves *iii*). $\square$

Result *i*) of Theorem 2.4.3 can be written equivalently as

$$\mathcal{N}(A)^\perp = \mathcal{R}(A^*), \quad (2.4.12)$$

$$\mathcal{N}(A) = \mathcal{R}(A^*)^\perp, \quad (2.4.13)$$

$$\mathcal{N}(A^*)^\perp = \mathcal{R}(A), \quad (2.4.14)$$

while replacing A by A^* in *ii*) and *iii*) of Theorem 2.4.3 yields

$$\mathcal{R}(A^*) = \mathcal{R}(A^*A), \quad (2.4.15)$$

$$\mathcal{N}(A^*) = \mathcal{N}(AA^*). \quad (2.4.16)$$

Using *ii*) of Theorem 2.4.3 and (2.4.15), it follows that

$$\mathcal{R}(AA^*A) = A\mathcal{R}(A^*A) = A\mathcal{R}(A^*) = \mathcal{R}(AA^*) = \mathcal{R}(A). \quad (2.4.17)$$

Letting $A \triangleq \begin{bmatrix} 1 & j \end{bmatrix}$ shows that $\mathcal{R}(A)$ and $\mathcal{R}(AA^T)$ may be different.

2.5 Rank and Defect

The *rank* of $A \in \mathbb{F}^{n \times m}$ is defined by

$$\text{rank } A \triangleq \dim \mathcal{R}(A). \quad (2.5.1)$$

It can be seen that the rank of A is equal to the number of linearly independent columns of A over $\mathbb{F}$. For example, if $\mathbb{F} = \mathbb{C}$, then $\text{rank} \begin{bmatrix} 1 & j \end{bmatrix} = 1$, while, if either $\mathbb{F} = \mathbb{R}$ or $\mathbb{F} = \mathbb{C}$, then $\text{rank} \begin{bmatrix} 1 & 1 \end{bmatrix} = 1$. Furthermore, $\text{rank } A = \text{rank } \overline{A}$, $\text{rank } A^T = \text{rank } A^*$, $\text{rank } A \leq m$, and $\text{rank } A^T \leq n$. If $\text{rank } A = m$, then A has *full column rank*, while, if $\text{rank } A^T = n$, then A has *full row rank*. If A has either full

column rank or full row rank, then A has *full rank*. Finally, the *defect* of A is

$$\text{def } A \triangleq \dim \mathcal{N}(A). \quad (2.5.2)$$

The following result follows from Theorem 2.4.3.

Corollary 2.5.1. Let $A \in \mathbb{F}^{n \times m}$. Then, the following identities hold:

- i) $\text{rank } A^* + \text{def } A = m$.
- ii) $\text{rank } A = \text{rank } AA^*$.
- iii) $\text{def } A = \text{def } A^*A$.

Proof. It follows from (2.4.12) and Proposition 2.3.2 that $\text{rank } A^* = \dim \mathcal{R}(A^*) = \dim \mathcal{N}(A)^\perp = m - \dim \mathcal{N}(A) = m - \text{def } A$, which proves i). Results ii) and iii) follow from ii) and iii) of Theorem 2.4.3. $\square$

Replacing A by A^* in Corollary 2.5.1 yields

$$\text{rank } A + \text{def } A^* = n, \quad (2.5.3)$$

$$\text{rank } A^* = \text{rank } A^*A, \quad (2.5.4)$$

$$\text{def } A^* = \text{def } AA^*. \quad (2.5.5)$$

Furthermore, note that

$$\text{def } A = \text{def } \overline{A} \quad (2.5.6)$$

and

$$\text{def } A^T = \text{def } A^*. \quad (2.5.7)$$

Lemma 2.5.2. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\text{rank } AB \leq \min\{\text{rank } A, \text{rank } B\}. \quad (2.5.8)$$

Proof. Since, by Lemma 2.4.1, $\mathcal{R}(AB) \subseteq \mathcal{R}(A)$, it follows that $\text{rank } AB \leq \text{rank } A$. Next, suppose that $\text{rank } B < \text{rank } AB$. Let $\{y_1, \dots, y_r\} \subset \mathbb{F}^n$ be a basis for $\mathcal{R}(AB)$, where $r \triangleq \text{rank } AB$, and, since $y_i \in A\mathcal{R}(B)$ for all $i = 1, \dots, r$, let $x_i \in \mathcal{R}(B)$ be such that $y_i = Ax_i$ for all $i = 1, \dots, r$. Since $\text{rank } B < r$, it follows that $x_1, \dots, x_r$ are linearly dependent. Hence, there exist $\alpha_1, \dots, \alpha_r \in \mathbb{F}$, not all zero, such that $\sum_{i=1}^r \alpha_i x_i = 0$, which implies that $\sum_{i=1}^r \alpha_i Ax_i = \sum_{i=1}^r \alpha_i y_i = 0$. Thus, $y_1, \dots, y_r$ are linearly dependent, which is a contradiction. $\square$

Corollary 2.5.3. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\text{rank } A = \text{rank } A^* \quad (2.5.9)$$

and

$$\text{def } A = \text{def } A^* + m - n. \quad (2.5.10)$$

Therefore,

$$\text{rank } A = \text{rank } A^*A.$$

If, in addition, $n = m$, then

$$\text{def } A = \text{def } A^*. \quad (2.5.11)$$

Proof. It follows from (2.5.8) with $B = A^*$ that $\text{rank } AA^* \leq \text{rank } A^*$. Furthermore, *ii*) of Corollary 2.5.1 implies that $\text{rank } A = \text{rank } AA^*$. Hence, $\text{rank } A \leq \text{rank } A^*$. Interchanging A and A^* and repeating this argument yields $\text{rank } A^* \leq \text{rank } A$. Hence, $\text{rank } A = \text{rank } A^*$. Next, using *i*) of Corollary 2.5.1, (2.5.9), and (2.5.3) it follows that $\text{def } A = m - \text{rank } A^* = m - \text{rank } A = m - (n - \text{def } A^*)$, which proves (2.5.10). $\square$

Corollary 2.5.4. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\text{rank } A \leq \min\{m, n\}. \quad (2.5.12)$$

Proof. By definition, $\text{rank } A \leq m$, while it follows from (2.5.9) that $\text{rank } A = \text{rank } A^* \leq n$. $\square$

The *dimension theorem* is given by (2.5.13) in the following result.

Corollary 2.5.5. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\text{rank } A + \text{def } A = m \quad (2.5.13)$$

and

$$\text{rank } A = \text{rank } A^*A. \quad (2.5.14)$$

Proof. The result (2.5.13) follows from *i*) of Corollary 2.5.1 and (2.5.9), while (2.5.14) follows from (2.5.4) and (2.5.9). $\square$

The following result follows from the subspace dimension theorem and the dimension theorem.

Corollary 2.5.6. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\dim[\mathcal{R}(A) + \mathcal{N}(A)] + \dim[\mathcal{R}(A) \cap \mathcal{N}(A)] = m. \quad (2.5.15)$$

Corollary 2.5.7. Let $A \in \mathbb{F}^{n \times n}$ and $k \geq 1$. Then,

$$\text{rank } A^k \leq \text{rank } A \quad (2.5.16)$$

and

$$\text{def } A \leq \text{def } A^k. \quad (2.5.17)$$

Proposition 2.5.8. Let $A \in \mathbb{F}^{n \times n}$. If $\text{rank } A^2 = \text{rank } A$, then $\text{rank } A^k = \text{rank } A$ for all $k \geq 1$. Equivalently, if $\text{def } A^2 = \text{def } A$, then $\text{def } A^k = \text{def } A$ for all $k \in \mathbb{P}$.

Proof. Since $\text{rank } A^2 = \text{rank } A$ and $\mathcal{R}(A^2) \subseteq \mathcal{R}(A)$, it follows from Lemma 2.3.4 that $\mathcal{R}(A^2) = \mathcal{R}(A)$. Hence, $\mathcal{R}(A^3) = A\mathcal{R}(A^2) = A\mathcal{R}(A) = \mathcal{R}(A^2)$. Thus, $\text{rank } A^3 = \text{rank } A$. Similar arguments yield $\text{rank } A^k = \text{rank } A$ for all $k \geq 1$. $\square$

We now prove *Sylvester's inequality*, which provides a lower bound for the rank of the product of two matrices.

Proposition 2.5.9. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\text{rank } A + \text{rank } B \leq m + \text{rank } AB. \quad (2.5.18)$$

Proof. Using (2.5.8) to obtain the second inequality below, it follows that

$$\begin{aligned} \text{rank } A + \text{rank } B &= \text{rank} \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \\ &\leq \text{rank} \begin{bmatrix} 0 & A \\ B & I \end{bmatrix} \\ &= \text{rank} \begin{bmatrix} I & A \\ 0 & I \end{bmatrix} \begin{bmatrix} -AB & 0 \\ B & I \end{bmatrix} \\ &\leq \text{rank} \begin{bmatrix} -AB & 0 \\ B & I \end{bmatrix} \\ &\leq \text{rank} \begin{bmatrix} -AB & 0 \end{bmatrix} + \text{rank} \begin{bmatrix} B & I \end{bmatrix} \\ &= \text{rank } AB + m. \quad \square \end{aligned}$$

Combining (2.5.8) with (2.5.18) yields the following result.

Corollary 2.5.10. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\text{rank } A + \text{rank } B - m \leq \text{rank } AB \leq \min\{\text{rank } A, \text{rank } B\}. \quad (2.5.19)$$

2.6 Invertibility

Let $A \in \mathbb{F}^{n \times m}$. Then, A is *left invertible* if there exists a matrix $A^L \in \mathbb{F}^{m \times n}$ such that $A^L A = I_m$, while A is *right invertible* if there exists a matrix $A^R \in \mathbb{F}^{m \times n}$ such that $AA^R = I_n$. These definitions are consistent with the definitions of left and right invertibility given in Chapter 1 applied to the function $f: \mathbb{F}^m \mapsto \mathbb{F}^n$ given by $f(x) = Ax$. Note that A^L (when it exists) and A^* are the same size, and likewise for A^R .

Theorem 2.6.1. Let $A \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

- i) A is left invertible.
- ii) A is one-to-one.
- iii) $\text{def } A = 0$.
- iv) $\text{rank } A = m$.
- v) A has full column rank.

The following statements are also equivalent:

- vi) A is right invertible.
- vii) A is onto.
- viii) $\text{def } A = m - n$.

- ix) $\text{rank } A = n$.
- x) A has full row rank.

Proposition 2.6.2. Let $A \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

- i) A has a unique left inverse.
- ii) A has a unique right inverse.
- iii) $\text{rank } A = n = m$.

Proof. To prove that i) implies iii), suppose that $\text{rank } A = m < n$ so that A is left invertible but nonsquare. Then, it follows from the dimension theorem Corollary 2.5.5 that $\text{def } A^T = n - m > 0$. Hence, there exist infinitely many matrices $A^L \in \mathbb{F}^{m \times n}$ such that $A^L A = I_m$. Conversely, suppose that $B \in \mathbb{F}^{n \times n}$ and $C \in \mathbb{F}^{n \times n}$ are left inverses of A . Then, $(B - C)A = 0$, and it follows from Sylvester's inequality Proposition 2.5.9 that $B = C$. $\square$

The following result shows that the rank and defect of a matrix are not affected by either left multiplication by a left invertible matrix or right multiplication by a right invertible matrix.

Proposition 2.6.3. Let $A \in \mathbb{F}^{n \times m}$, and let $C \in \mathbb{F}^{k \times n}$ be left invertible and $B \in \mathbb{F}^{m \times l}$ be right invertible. Then,

$$\mathcal{R}(A) = \mathcal{R}(AB) \quad (2.6.1)$$

and

$$\mathcal{N}(A) = \mathcal{N}(CA). \quad (2.6.2)$$

Furthermore,

$$\text{rank } A = \text{rank } CA = \text{rank } AB \quad (2.6.3)$$

and

$$\text{def } A = \text{def } CA = \text{def } AB + m - l. \quad (2.6.4)$$

Proof. Let C^L be a left inverse of C . Using both inequalities in (2.5.19) and the fact that $\text{rank } A \leq n$, it follows that

$$\text{rank } A = \text{rank } A + \text{rank } C^L C - n \leq \text{rank } C^L C A \leq \text{rank } CA \leq \text{rank } A,$$

which implies that $\text{rank } A = \text{rank } CA$. Next, (2.5.13) and (2.6.3) imply that $m - \text{def } A = m - \text{def } CA = l - \text{def } AB$, which yields (2.6.4). $\square$

As shown in Proposition 2.6.2, left and right inverses of nonsquare matrices are not unique. For example, the matrix $A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ is left invertible and has left inverses $\begin{bmatrix} 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 1 & 1 \end{bmatrix}$. In spite of this nonuniqueness, however, left inverses are useful for solving equations of the form $Ax = b$, where $A \in \mathbb{F}^{n \times m}$, $x \in \mathbb{F}^m$, and $b \in \mathbb{F}^n$. If A is left invertible, then one can formally (although not rigorously) solve $Ax = b$ by noting that $x = A^L Ax = A^L b$, where $A^L \in \mathbb{F}^{m \times n}$ is a left inverse of A . However, it is necessary to determine beforehand whether or not there actually

exists a vector x satisfying $Ax = b$. For example, if $A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ and $b = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, then A is left invertible although there does not exist a vector x satisfying $Ax = b$. The following result addresses the various possibilities that can arise. One interesting feature of this result is that, if there exists a solution of $Ax = b$ and A is left invertible, then the solution is unique even if A does not have a unique left inverse. For this result, $\begin{bmatrix} A & b \end{bmatrix}$ denotes the $n \times (m+1)$ partitioned matrix formed from A and b . Note that $\text{rank } A \leq \text{rank } \begin{bmatrix} A & b \end{bmatrix} \leq m+1$, while $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix}$ is equivalent to $b \in \mathcal{R}(A)$.

Theorem 2.6.4. Let $A \in \mathbb{F}^{n \times m}$ and $b \in \mathbb{F}^n$. Then, the following statements hold:

- i) There does not exist a vector $x \in \mathbb{F}^m$ satisfying $Ax = b$ if and only if $\text{rank } A < \text{rank } \begin{bmatrix} A & b \end{bmatrix}$.
- ii) There exists a unique vector $x \in \mathbb{F}^m$ satisfying $Ax = b$ if and only if $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix} = m$. In this case, if $A^L \in \mathbb{F}^{m \times n}$ is a left inverse of A , then the solution is given by $x = A^L b$.
- iii) There exist infinitely many $x \in \mathbb{F}^m$ satisfying $Ax = b$ if and only if $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix} < m$. In this case, let $\hat{x} \in \mathbb{F}^m$ satisfy $A\hat{x} = b$. Then, the set of solutions of $Ax = b$ is given by $\hat{x} + \mathcal{N}(A)$.
- iv) Assume that $\text{rank } A = n$. Then, there exists at least one vector $x \in \mathbb{F}^m$ satisfying $Ax = b$. Furthermore, if $A^R \in \mathbb{F}^{m \times n}$ is a right inverse of A , then $x = A^R b$ satisfies $Ax = b$. If $n = m$, then $x = A^R b$ is the unique solution of $Ax = b$. If $n < m$ and $\hat{x} \in \mathbb{F}^n$ satisfies $A\hat{x} = b$, then the set of solutions of $Ax = b$ is given by $\hat{x} + \mathcal{N}(A)$.

Proof. To prove i), note that $\text{rank } A < \text{rank } \begin{bmatrix} A & b \end{bmatrix}$ is equivalent to the fact that b cannot be represented as a linear combination of columns of A , that is, $Ax = b$ does not have a solution $x \in \mathbb{F}^m$. To prove ii), suppose that $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix} = m$ so that, by i), $Ax = b$ has a solution $x \in \mathbb{F}^m$. If $\hat{x} \in \mathbb{F}^m$ satisfies $A\hat{x} = b$, then $A(x - \hat{x}) = 0$. Since $\text{rank } A = m$, it follows from Theorem 2.6.1 that A has a left inverse $A^L \in \mathbb{F}^{m \times n}$. Hence, $x - \hat{x} = A^L A(x - \hat{x}) = 0$, which proves that $Ax = b$ has a unique solution. Conversely, suppose that $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix} = m$ and there exist vectors $x, \hat{x} \in \mathbb{F}^m$, where $x \neq \hat{x}$, such that $Ax = b$ and $A\hat{x} = b$. Then, $A(x - \hat{x}) = 0$, which implies that $\text{def } A \geq 1$. Therefore, $\text{rank } A = m - \text{def } A \leq m - 1$, which is a contradiction. This proves the first statement of ii). Assuming $Ax = b$ has a unique solution $x \in \mathbb{F}^m$, multiplying by A^L yields $x = A^L b$. To prove iii), note that it follows from i) that $Ax = b$ has at least one solution $\hat{x} \in \mathbb{F}^m$. Hence, $x \in \mathbb{F}^m$ is a solution of $Ax = b$ if and only if $A(x - \hat{x}) = 0$, or, equivalently, $x \in \hat{x} + \mathcal{N}(A)$. To prove iv), note that, since $\text{rank } A = n$, it follows that $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix}$, and thus either ii) or iii) applies. $\square$

The set of solutions $x \in \mathbb{F}^m$ to $Ax = b$ is explicitly characterized by Proposition 6.1.7.

Let $A \in \mathbb{F}^{n \times m}$. Proposition 2.6.2 considers the uniqueness of left and right inverses of A , but does not consider the case in which a matrix is both a left inverse and a right inverse of A . Consequently, we say that A is *nonsingular* if there exists

a matrix $B \in \mathbb{F}^{m \times n}$, the *inverse* of A , such that $BA = I_m$ and $AB = I_n$, that is, B is both a left and right inverse of A .

Proposition 2.6.5. Let $A \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

- i) A is nonsingular
- ii) $\text{rank } A = n = m$.

In this case, A has a unique inverse.

Proof. If A is nonsingular, then, since B is both left and right invertible, it follows from Theorem 2.6.1 that $\text{rank } A = m$ and $\text{rank } A = n$. Hence, ii) holds. Conversely, it follows from Theorem 2.6.1 that A has both a left inverse B and a right inverse C . Then, $B = BI_n = BAC = I_n C = C$. Hence, B is also a right inverse of A . Thus, A is nonsingular. In fact, the same argument shows that A has a unique inverse. $\square$

The following result can be viewed as a specialization of Theorem 1.2.2 to the function $f: \mathbb{F}^n \mapsto \mathbb{F}^n$, where $f(x) = Ax$.

Corollary 2.6.6. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is nonsingular.
- ii) A has a unique inverse.
- iii) A is one-to-one.
- iv) A is onto.
- v) A is left invertible.
- vi) A is right invertible.
- vii) A has a unique left inverse.
- viii) A has a unique right inverse.
- ix) $\text{rank } A = n$.
- x) $\text{def } A = 0$.

Let $A \in \mathbb{F}^{n \times n}$ be nonsingular. Then, the inverse of A , denoted by A^{-1} , is a unique $n \times n$ matrix with entries in $\mathbb{F}$. If A is not nonsingular, then A is *singular*.

The following result is a specialization of Theorem 2.6.4 to the case $n = m$.

Corollary 2.6.7. Let $A \in \mathbb{F}^{n \times n}$ and $b \in \mathbb{F}^n$. Then, the following statements hold:

- i) A is nonsingular if and only if there exists a unique vector $x \in \mathbb{F}^n$ satisfying $Ax = b$. In this case, $x = A^{-1}b$.
- ii) A is singular and $\text{rank } A = \text{rank} \begin{bmatrix} A & b \end{bmatrix}$ if and only if there exist infinitely

many $x \in \mathbb{R}^n$ satisfying $Ax = b$. In this case, let $\hat{x} \in \mathbb{F}^m$ satisfy $A\hat{x} = b$. Then, the set of solutions of $Ax = b$ is given by $\hat{x} + \mathcal{N}(A)$.

Proposition 2.6.8. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is nonsingular.
- ii) $\overline{A}$ is nonsingular.
- iii) A^T is nonsingular.
- iv) A^* is nonsingular.

In this case,

$$(\overline{A})^{-1} = \overline{A^{-1}}, \quad (2.6.5)$$

$$(A^T)^{-1} = (A^{-1})^T, \quad (2.6.6)$$

$$(A^*)^{-1} = (A^{-1})^*. \quad (2.6.7)$$

Proof. Since $AA^{-1} = I$, it follows that $(A^{-1})^*A^* = I$. Hence, $(A^{-1})^* = (A^*)^{-1}$. $\square$

We thus use A^{-T} to denote $(A^T)^{-1}$ or $(A^{-1})^T$ and A^{-*} to denote $(A^*)^{-1}$ or $(A^{-1})^*$.

Proposition 2.6.9. Let $A, B \in \mathbb{F}^{n \times n}$ be nonsingular. Then,

$$(AB)^{-1} = B^{-1}A^{-1}, \quad (2.6.8)$$

$$(AB)^{-T} = A^{-T}B^{-T}, \quad (2.6.9)$$

$$(AB)^{-*} = A^{-*}B^{-*}. \quad (2.6.10)$$

Proof. Note that $ABB^{-1}A^{-1} = AIA^{-1} = I$, which shows that $B^{-1}A^{-1}$ is the inverse of AB . Similarly, $(AB)^*A^{-*}B^{-*} = B^*A^*A^{-*}B^{-*} = B^*IB^{-*} = I$, which shows that $A^{-*}B^{-*}$ is the inverse of $(AB)^*$. $\square$

For a nonsingular matrix $A \in \mathbb{F}^{n \times n}$ and $r \in \mathbb{Z}$ we write

$$A^{-r} \triangleq (A^r)^{-1} = (A^{-1})^r, \quad (2.6.11)$$

$$A^{-rT} \triangleq (A^r)^{-T} = (A^{-T})^r = (A^{-r})^T = (A^T)^{-r}, \quad (2.6.12)$$

$$A^{-r*} \triangleq (A^r)^{-*} = (A^{-*})^r = (A^{-r})^* = (A^*)^{-r}. \quad (2.6.13)$$

For example, $A^{-2*} = (A^{-*})^2$.

2.7 The Determinant

One of the most useful quantities associated with a square matrix is its determinant. In this section we develop some basic results pertaining to the determinant of a matrix.

The *determinant* of $A \in \mathbb{F}^{n \times n}$ is defined by

$$\det A \triangleq \sum_{\sigma} (-1)^{N_{\sigma}} \prod_{i=1}^n A_{(i, \sigma(i))}, \quad (2.7.1)$$

where the sum is taken over all $n!$ permutations $\sigma = (\sigma(1), \dots, \sigma(n))$ of the column indices $1, \dots, n$, and where N_{σ} is the minimal number of pairwise transpositions needed to transform $\sigma(1), \dots, \sigma(n)$ to $1, \dots, n$. The following result is an immediate consequence of this definition.

Proposition 2.7.1. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\det A^T = \det A, \quad (2.7.2)$$

$$\det \overline{A} = \overline{\det A}, \quad (2.7.3)$$

$$\det A^* = \overline{\det A}, \quad (2.7.4)$$

and, for all $\alpha \in \mathbb{F}$,

$$\det \alpha A = \alpha^n \det A. \quad (2.7.5)$$

If, in addition, $B \in \mathbb{F}^{m \times n}$ and $C \in \mathbb{F}^{n \times m}$, then

$$\det \begin{bmatrix} A & 0 \\ B & C \end{bmatrix} = (\det A)(\det C). \quad (2.7.6)$$

The following observations are immediate consequences of the definition of the determinant.

Proposition 2.7.2. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

i) If every off-diagonal entry of A is zero, then

$$\det A = \prod_{i=1}^n A_{(i,i)}. \quad (2.7.7)$$

In particular, $\det I_n = 1$.

ii) If A has a row or column consisting entirely of 0's, then $\det A = 0$.

iii) If A has two identical rows or two identical columns, then $\det A = 0$.

iv) If $x \in \mathbb{F}^n$ and $i \in \{1, \dots, n\}$, then

$$\det(A + xe_i^T) = \det A + \det(A \overset{i}{\leftarrow} x). \quad (2.7.8)$$

v) If $x \in \mathbb{F}^{1 \times n}$ and $i \in \{1, \dots, n\}$, then

$$\det(A + e_i x) = \det A + \det(A \overset{i}{\leftarrow} x). \quad (2.7.9)$$

vi) If B is identical to A except that, for some $i \in \{1, \dots, n\}$ and $\alpha \in \mathbb{F}$, either $\text{col}_i(B) = \alpha \text{col}_i(A)$ or $\text{row}_i(B) = \alpha \text{row}_i(A)$, then $\det B = \alpha \det A$.

vii) If B is formed from A by interchanging two rows or two columns of A , then $\det B = -\det A$.

viii) If B is formed from A by adding a multiple of a (row, column) of A to another (row, column) of A , then $\det B = \det A$.

Statements vi)–viii) correspond, respectively, to multiplying the matrix A on the left or right by matrices of the form

$$I_n + (\alpha - 1)E_{i,i} = \begin{bmatrix} I_{i-1} & 0 & 0 \\ 0 & \alpha & 0 \\ 0 & 0 & I_{n-i} \end{bmatrix}, \quad (2.7.10)$$

$$I_n + E_{i,j} + E_{j,i} - E_{i,i} - E_{j,j} = \begin{bmatrix} I_{i-1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & I_{j-i-1} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & I_{n-j} \end{bmatrix}, \quad (2.7.11)$$

where $i \neq j$, and

$$I_n + \beta E_{i,j} = \begin{bmatrix} I_{i-1} & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & \beta & 0 \\ 0 & 0 & I_{j-i-1} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & I_{n-j} \end{bmatrix}, \quad (2.7.12)$$

where $\beta \in \mathbb{F}$ and $i \neq j$. The matrices in (2.7.11) and (2.7.12) illustrate the case $i < j$. Since $I + (\alpha - 1)E_{i,i} = I + (\alpha - 1)e_i e_i^T$, $I + E_{i,j} + E_{j,i} - E_{i,i} - E_{j,j} = I - (e_i - e_j)(e_i - e_j)^T$, and $I + \beta E_{i,j} = I + \beta e_i e_j^T$, it follows that all of these matrices are of the form $I - xy^T$. In terms of Definition 3.1.1, (2.7.10) is an elementary matrix if and only if $\alpha \neq 0$, (2.7.11) is an elementary matrix, and (2.7.12) is an elementary matrix if and only if either $i \neq j$ or $\beta \neq -1$.

Proposition 2.7.3. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\det AB = \det BA = (\det A)(\det B). \quad (2.7.13)$$

Proof. First note the identity

$$\begin{bmatrix} A & 0 \\ I & B \end{bmatrix} = \begin{bmatrix} I & A \\ 0 & I \end{bmatrix} \begin{bmatrix} -AB & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} I & 0 \\ B & I \end{bmatrix} \begin{bmatrix} 0 & I \\ I & 0 \end{bmatrix}.$$

The first and third matrices on the right-hand side of this identity add multiples of rows and columns of $\begin{bmatrix} -AB & 0 \\ 0 & I \end{bmatrix}$ to other rows and columns of $\begin{bmatrix} -AB & 0 \\ 0 & I \end{bmatrix}$. As already noted, these operations do not affect the determinant of $\begin{bmatrix} -AB & 0 \\ 0 & I \end{bmatrix}$. In addition, the fourth matrix on the right-hand side of this identity interchanges n pairs of columns of $\begin{bmatrix} 0 & I \\ I & 0 \end{bmatrix}$. Using (2.7.5), (2.7.6), and the fact that every interchange of a pair of columns of $\begin{bmatrix} 0 & I \\ I & 0 \end{bmatrix}$ entails a factor of -1 , it thus follows that $(\det A)(\det B) = \det \begin{bmatrix} A & 0 \\ I & B \end{bmatrix} = (-1)^n \det \begin{bmatrix} -AB & 0 \\ 0 & I \end{bmatrix} = (-1)^n \det(-AB) = \det AB$. $\square$

Corollary 2.7.4. Let $A \in \mathbb{F}^{n \times n}$ be nonsingular. Then, $\det A \neq 0$ and

$$\det A^{-1} = (\det A)^{-1}. \quad (2.7.14)$$

Proof. Since $AA^{-1} = I_n$, it follows that $\det AA^{-1} = (\det A)(\det A^{-1}) = 1$. Hence, $\det A \neq 0$. In addition, $\det A^{-1} = 1/\det A$. $\square$

Let $A \in \mathbb{F}^{n \times m}$. The determinant of a square submatrix of A is a *subdeterminant* of A . By convention, the determinant of A is a subdeterminant of A . The determinant of a $j \times j$ (principal, leading principal) submatrix of A is a $j \times j$ (*principal, leading principal*) *subdeterminant* of A .

Let $A \in \mathbb{F}^{n \times n}$. Then, the *cofactor* of $A_{(i,j)}$, denoted by $A_{[i;j]}$, is the $(n-1) \times (n-1)$ submatrix of A obtained by deleting the i th row and j th column of A . In other words,

$$A_{[i;j]} \triangleq A_{(\{i\}^{\sim}, \{j\}^{\sim})}. \quad (2.7.15)$$

The following result provides a cofactor expansion of $\det A$.

Proposition 2.7.5. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $i = 1, \dots, n$,

$$\sum_{k=1}^n (-1)^{i+k} A_{(i,k)} \det A_{[i;k]} = \det A. \quad (2.7.16)$$

Furthermore, for all $i, j = 1, \dots, n$ such that $j \neq i$,

$$\sum_{k=1}^n (-1)^{i+k} A_{(j,k)} \det A_{[i;k]} = 0. \quad (2.7.17)$$

Proof. Identity (2.7.16) is an equivalent recursive form of the definition $\det A$, while the right-hand side of (2.7.17) is equal to $\det B$, where B is obtained from A by replacing $\text{row}_i(A)$ by $\text{row}_j(A)$. As already noted, $\det B = 0$. $\square$

Let $A \in \mathbb{F}^{n \times n}$, where $n \geq 2$. To simplify (2.7.16) and (2.7.17) it is useful to define the *adjugate* of A , denoted by $A^A \in \mathbb{F}^{n \times n}$, where, for all $i, j = 1, \dots, n$,

$$(A^A)_{(i,j)} \triangleq (-1)^{i+j} \det A_{[j;i]} = \det(A \stackrel{i}{\leftarrow} e_j). \quad (2.7.18)$$

Then, (2.7.16) implies that, for all $i = 1, \dots, n$,

$$\sum_{k=1}^n A_{(i,k)} (A^A)_{(k,i)} = (AA^A)_{(i,i)} = (A^A A)_{(i,i)} = \det A, \quad (2.7.19)$$

while (2.7.17) implies that, for all $i, j = 1, \dots, n$ such that $j \neq i$,

$$\sum_{k=1}^n A_{(i,k)} (A^A)_{(k,j)} = (AA^A)_{(i,j)} = (A^A A)_{(i,j)} = 0. \quad (2.7.20)$$

Thus,

$$AA^A = A^A A = (\det A)I. \quad (2.7.21)$$

Consequently, if $\det A \neq 0$, then

$$A^{-1} = \frac{1}{\det A} A^A, \quad (2.7.22)$$

whereas, if $\det A = 0$, then

$$AA^A = A^A A = 0. \quad (2.7.23)$$

For a scalar $A \in \mathbb{F}$, we define $A^A \triangleq 1$.

The following result provides the converse of Corollary 2.7.4 by using (2.7.22) to construct A^{-1} in terms of $(n-1) \times (n-1)$ subdeterminants of A .

Corollary 2.7.6. Let $A \in \mathbb{F}^{n \times n}$. Then, A is nonsingular if and only if $\det A \neq 0$. In this case, for all $i, j = 1, \dots, n$, the (i, j) entry of A^{-1} is given by

$$(A^{-1})_{(i,j)} = (-1)^{i+j} \frac{\det A_{[j;i]}}{\det A}. \quad (2.7.24)$$

Finally, the following result uses the nonsingularity of submatrices to characterize the rank of a matrix.

Proposition 2.7.7. Let $A \in \mathbb{F}^{n \times m}$. Then, $\text{rank } A$ is the largest order of all nonsingular submatrices of A .

2.8 Partitioned Matrices

Partitioned matrices were used to state or prove several results in this chapter including Proposition 2.5.9, Theorem 2.6.4, Proposition 2.7.1, and Proposition 2.7.3. In this section we give several useful identities involving partitioned matrices.

Proposition 2.8.1. Let $A_{ij} \in \mathbb{F}^{n_i \times m_j}$ for all $i = 1, \dots, k$ and $j = 1, \dots, l$. Then,

$$\begin{bmatrix} A_{11} & \cdots & A_{1l} \\ \vdots & \ddots & \vdots \\ A_{k1} & \cdots & A_{kl} \end{bmatrix}^T = \begin{bmatrix} A_{11}^T & \cdots & A_{k1}^T \\ \vdots & \ddots & \vdots \\ A_{1l}^T & \cdots & A_{kl}^T \end{bmatrix} \quad (2.8.1)$$

and

$$\begin{bmatrix} A_{11} & \cdots & A_{1l} \\ \vdots & \ddots & \vdots \\ A_{k1} & \cdots & A_{kl} \end{bmatrix}^* = \begin{bmatrix} A_{11}^* & \cdots & A_{k1}^* \\ \vdots & \ddots & \vdots \\ A_{1l}^* & \cdots & A_{kl}^* \end{bmatrix}. \quad (2.8.2)$$

If, in addition, $k = l$ and $n_i = m_i$ for all $i = 1, \dots, m$, then

$$\text{tr} \begin{bmatrix} A_{11} & \cdots & A_{1k} \\ \vdots & \ddots & \vdots \\ A_{k1} & \cdots & A_{kk} \end{bmatrix} = \sum_{i=1}^k \text{tr } A_{ii} \quad (2.8.3)$$

and

$$\det \begin{bmatrix} A_{11} & A_{12} & \cdots & A_{1k} \\ 0 & A_{22} & \cdots & A_{2k} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & A_{kk} \end{bmatrix} = \prod_{i=1}^k \det A_{ii}. \quad (2.8.4)$$

Lemma 2.8.2. Let $B \in \mathbb{F}^{n \times m}$ and $C \in \mathbb{F}^{m \times n}$. Then,

$$\begin{bmatrix} I & B \\ 0 & I \end{bmatrix}^{-1} = \begin{bmatrix} I & -B \\ 0 & I \end{bmatrix} \quad (2.8.5)$$

and

$$\begin{bmatrix} I & 0 \\ C & I \end{bmatrix}^{-1} = \begin{bmatrix} I & 0 \\ -C & I \end{bmatrix}. \quad (2.8.6)$$

Let $A \in \mathbb{F}^{n \times n}$ and $D \in \mathbb{F}^{m \times m}$ be nonsingular. Then,

$$\begin{bmatrix} A & 0 \\ 0 & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} & 0 \\ 0 & D^{-1} \end{bmatrix}. \quad (2.8.7)$$

Proposition 2.8.3. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{l \times n}$, and $D \in \mathbb{F}^{l \times m}$, and assume that A is nonsingular. Then,

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} I & 0 \\ CA^{-1} & I \end{bmatrix} \begin{bmatrix} A & 0 \\ 0 & D - CA^{-1}B \end{bmatrix} \begin{bmatrix} I & A^{-1}B \\ 0 & I \end{bmatrix} \quad (2.8.8)$$

and

$$\text{rank} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = n + \text{rank}(D - CA^{-1}B). \quad (2.8.9)$$

If, furthermore, $l = m$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = (\det A) \det(D - CA^{-1}B). \quad (2.8.10)$$

Proposition 2.8.4. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{l \times m}$, and $D \in \mathbb{F}^{l \times l}$, and assume that D is nonsingular. Then,

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} I & BD^{-1} \\ 0 & I \end{bmatrix} \begin{bmatrix} A - BD^{-1}C & 0 \\ 0 & D \end{bmatrix} \begin{bmatrix} I & 0 \\ D^{-1}C & I \end{bmatrix} \quad (2.8.11)$$

and

$$\text{rank} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = l + \text{rank}(A - BD^{-1}C). \quad (2.8.12)$$

If, furthermore, $n = m$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = (\det D) \det(A - BD^{-1}C). \quad (2.8.13)$$

Corollary 2.8.5. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then,

$$\begin{aligned} \begin{bmatrix} I_n & A \\ B & I_m \end{bmatrix} &= \begin{bmatrix} I_n & 0 \\ B & I_m \end{bmatrix} \begin{bmatrix} I_n & 0 \\ 0 & I_m - BA \end{bmatrix} \begin{bmatrix} I_n & A \\ 0 & I_m \end{bmatrix} \\ &= \begin{bmatrix} I_n & A \\ 0 & I_m \end{bmatrix} \begin{bmatrix} I_n - AB & 0 \\ 0 & I_m \end{bmatrix} \begin{bmatrix} I_n & 0 \\ B & I_m \end{bmatrix}. \end{aligned}$$

Hence,

$$\text{rank} \begin{bmatrix} I_n & A \\ B & I_m \end{bmatrix} = n + \text{rank}(I_m - BA) = m + \text{rank}(I_n - AB)$$

and

$$\det \begin{bmatrix} I_n & A \\ B & I_m \end{bmatrix} = \det(I_m - BA) = \det(I_n - AB). \quad (2.8.14)$$

Hence, $I_n + AB$ is nonsingular if and only if $I_m + BA$ is nonsingular.

Lemma 2.8.6. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, and $D \in \mathbb{F}^{m \times m}$. If A and D are nonsingular, then

$$(\det A) \det(D - CA^{-1}B) = (\det D) \det(A - BD^{-1}C), \quad (2.8.15)$$

and thus $D - CA^{-1}B$ is nonsingular if and only if $A - BD^{-1}C$ is nonsingular.

Proposition 2.8.7. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, and $D \in \mathbb{F}^{m \times m}$. If A and $D - CA^{-1}B$ are nonsingular, then

$$\begin{aligned} & \begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} \\ &= \begin{bmatrix} A^{-1} + A^{-1}B(D - CA^{-1}B)^{-1}CA^{-1} & -A^{-1}B(D - CA^{-1}B)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}. \end{aligned} \quad (2.8.16)$$

If D and $A - BD^{-1}C$ are nonsingular, then

$$\begin{aligned} & \begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} \\ &= \begin{bmatrix} (A - BD^{-1}C)^{-1} & -(A - BD^{-1}C)^{-1}BD^{-1} \\ -D^{-1}C(A - BD^{-1}C)^{-1} & D^{-1} + D^{-1}C(A - BD^{-1}C)^{-1}BD^{-1} \end{bmatrix}. \end{aligned} \quad (2.8.17)$$

If A , D , and $D - CA^{-1}B$ are nonsingular, then $A - BD^{-1}C$ is nonsingular, and

$$\begin{aligned} & \begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} \\ &= \begin{bmatrix} (A - BD^{-1}C)^{-1} & -(A - BD^{-1}C)^{-1}BD^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}. \end{aligned} \quad (2.8.18)$$

The following result is the *matrix inversion lemma*.

Corollary 2.8.8. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, and $D \in \mathbb{F}^{m \times m}$. If A , $D - CA^{-1}B$, and D are nonsingular, then $A - BD^{-1}C$ is nonsingular,

$$(A - BD^{-1}C)^{-1} = A^{-1} + A^{-1}B(D - CA^{-1}B)^{-1}CA^{-1}, \quad (2.8.19)$$

and

$$C(A - BD^{-1}C)^{-1}A = D(D - CA^{-1}B)^{-1}C. \quad (2.8.20)$$

If A and $I - CA^{-1}B$ are nonsingular, then $A - BC$ is nonsingular, and

$$(A - BC)^{-1} = A^{-1} + A^{-1}B(I - CA^{-1}B)^{-1}CA^{-1}. \quad (2.8.21)$$

If $D - CB$, and D are nonsingular, then $I - BD^{-1}C$ is nonsingular, and

$$(I - BD^{-1}C)^{-1} = I + B(D - CB)^{-1}C. \quad (2.8.22)$$

If $I - CB$ is nonsingular, then $I - BC$ is nonsingular, and

$$(I - BC)^{-1} = I + B(I - CB)^{-1}C. \quad (2.8.23)$$

Corollary 2.8.9. Let $A, B, C, D \in \mathbb{F}^{n \times n}$. If $A, B, C - DB^{-1}A$, and $D - CA^{-1}B$ are nonsingular, then

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} - (C - DB^{-1}A)^{-1}CA^{-1} & (C - DB^{-1}A)^{-1} \\ -(D - CA^{-1}B)^{-1}CA^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}. \quad (2.8.24)$$

If $A, C, B - AC^{-1}D$, and $D - CA^{-1}B$ are nonsingular, then

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} - A^{-1}B(B - AC^{-1}D)^{-1} & -A^{-1}B(D - CA^{-1}B)^{-1} \\ (B - AC^{-1}D)^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}. \quad (2.8.25)$$

If $A, B, C, B - AC^{-1}D$, and $D - CA^{-1}B$ are nonsingular, then $C - DB^{-1}A$ is nonsingular, and

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} - A^{-1}B(B - AC^{-1}D)^{-1} & (C - DB^{-1}A)^{-1} \\ (B - AC^{-1}D)^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}. \quad (2.8.26)$$

If $B, D, A - BD^{-1}C$, and $C - DB^{-1}A$ are nonsingular, then

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} (A - BD^{-1}C)^{-1} & (C - DB^{-1}A)^{-1} \\ -D^{-1}C(A - BD^{-1}C)^{-1} & D^{-1} - D^{-1}C(C - DB^{-1}A)^{-1} \end{bmatrix}. \quad (2.8.27)$$

If $C, D, A - BD^{-1}C$, and $B - AC^{-1}D$ are nonsingular, then

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} (A - BD^{-1}C)^{-1} & -(A - BD^{-1}C)^{-1}BD^{-1} \\ (B - AC^{-1}D)^{-1} & D^{-1} - (B - AC^{-1}D)^{-1}BD^{-1} \end{bmatrix}. \quad (2.8.28)$$

If $B, C, D, A - BD^{-1}C$, and $C - DB^{-1}A$ are nonsingular, then $B - AC^{-1}D$ is nonsingular, and

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} (A - BD^{-1}C)^{-1} & (C - DB^{-1}A)^{-1} \\ (B - AC^{-1}D)^{-1} & D^{-1} - D^{-1}C(C - DB^{-1}A)^{-1} \end{bmatrix}. \quad (2.8.29)$$

Finally, if $A, B, C, D, A - BD^{-1}C$, and $B - AC^{-1}D$, are nonsingular, then $C - DB^{-1}A$ and $D - CA^{-1}B$ are nonsingular, and

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} (A - BD^{-1}C)^{-1} & (C - DB^{-1}A)^{-1} \\ (B - AC^{-1}D)^{-1} & (D - CA^{-1}B)^{-1} \end{bmatrix}. \quad (2.8.30)$$

Corollary 2.8.10. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and $I - A^{-1}B$ are nonsingular. Then, $A - B$ is nonsingular, and

$$(A - B)^{-1} = A^{-1} + A^{-1}B(I - A^{-1}B)^{-1}A^{-1}. \quad (2.8.31)$$

If, in addition, B is nonsingular, then

$$(A - B)^{-1} = A^{-1} + A^{-1}(B^{-1} - A^{-1})^{-1}A^{-1}. \quad (2.8.32)$$

2.9 Facts on Polars, Cones, Dual Cones, Convex Hulls, and Subspaces

Fact 2.9.1. Let $\mathcal{S} \subseteq \mathbb{F}^n$, assume that $\mathcal{S}$ is convex, and let $\alpha \in [0, 1]$. Then,

$$\alpha\mathcal{S} + (1 - \alpha)\mathcal{S} = \mathcal{S}.$$

Fact 2.9.2. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are convex. Then, $\mathcal{S}_1 + \mathcal{S}_2$ is convex.

Fact 2.9.3. Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then, the following statements hold:

- i) $\text{coco } \mathcal{S} = \text{co cone } \mathcal{S} = \text{cone co } \mathcal{S}$.
- ii) $\mathcal{S}^{\perp\perp} = \text{span } \mathcal{S} = \text{coco}(\mathcal{S} \cup -\mathcal{S})$.
- iii) $\mathcal{S} \subseteq \text{co } \mathcal{S} \subseteq (\text{aff } \mathcal{S} \cap \text{coco } \mathcal{S}) \subseteq \left\{ \begin{smallmatrix} \text{aff } \mathcal{S} \\ \text{coco } \mathcal{S} \end{smallmatrix} \right\} \subseteq \text{span } \mathcal{S}$.
- iv) $\mathcal{S} \subseteq (\text{co } \mathcal{S} \cap \text{cone } \mathcal{S}) \subseteq \left\{ \begin{smallmatrix} \text{co } \mathcal{S} \\ \text{cone } \mathcal{S} \end{smallmatrix} \right\} \subseteq \text{coco } \mathcal{S} \subseteq \text{span } \mathcal{S}$.
- v) $\text{dccone } \text{dccone } \mathcal{S} = \text{cl coco } \mathcal{S}$.

(Proof: For v), see [239, p. 54].) (Remark: See [176, p. 52]. Note that “pointed” in [176] means one-sided.)

Fact 2.9.4. Let $\mathcal{S}, \mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$. Then, the following statements hold:

- i) $\text{polar } \mathcal{S}$ is a closed, convex set containing the origin.
- ii) $\text{polar } \mathbb{F}^n = \{0\}$, and $\text{polar } \{0\} = \mathbb{F}^n$.
- iii) If $\alpha > 0$, then $\text{polar } \alpha\mathcal{S} = \frac{1}{\alpha} \text{polar } \mathcal{S}$.
- iv) $\mathcal{S} \subseteq \text{polar polar } \mathcal{S}$.
- v) If $\mathcal{S}$ is nonempty, then $\text{polar polar polar } \mathcal{S} = \text{polar } \mathcal{S}$.
- vi) If $\mathcal{S}$ is nonempty, then $\text{polar polar } \mathcal{S} = \text{cl co}(\mathcal{S} \cup \{0\})$.
- vii) If $0 \in \mathcal{S}$ and $\mathcal{S}$ is closed and convex, then $\text{polar polar } \mathcal{S} = \mathcal{S}$.
- viii) If $\mathcal{S}_1 \subseteq \mathcal{S}_2$, then $\text{polar } \mathcal{S}_2 \subseteq \text{polar } \mathcal{S}_1$.
- ix) $\text{polar}(\mathcal{S}_1 \cup \mathcal{S}_2) = (\text{polar } \mathcal{S}_1) \cap (\text{polar } \mathcal{S}_2)$.
- x) If $\mathcal{S}$ is a convex cone, then $\text{polar } \mathcal{S} = \text{dccone } \mathcal{S}$.

(Proof: See [153, pp. 143–147].)

Fact 2.9.5. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are cones. Then,

$$\text{dcone}(\mathcal{S}_1 + \mathcal{S}_2) = (\text{dcone } \mathcal{S}_1) \cap (\text{dcone } \mathcal{S}_2).$$

If, in addition, $\mathcal{S}_1$ and $\mathcal{S}_2$ are closed and convex, then

$$\text{dcone}(\mathcal{S}_1 \cap \mathcal{S}_2) = \text{cl}[(\text{dcone } \mathcal{S}_1) + (\text{dcone } \mathcal{S}_2)].$$

(Proof: See [239, pp. 58, 59] or [153, p. 147].)

Fact 2.9.6. Let $\mathcal{S} \subset \mathbb{F}^n$. Then, the following statements hold:

- i) $\mathcal{S}$ is an affine hyperplane if and only if there exist a nonzero vector $y \in \mathbb{F}^n$ and $\alpha \in \mathbb{R}$ such that $\mathcal{S} = \{x: \text{Re } x^*y = \alpha\}$.
- ii) $\mathcal{S}$ is an affine closed half space if and only if there exist a nonzero vector $y \in \mathbb{F}^n$ and $\alpha \in \mathbb{R}$ such that $\mathcal{S} = \{x \in \mathbb{F}^n: \text{Re } x^*y \leq \alpha\}$.
- iii) $\mathcal{S}$ is an affine open half space if and only if there exist a nonzero vector $y \in \mathbb{F}^n$ and $\alpha \in \mathbb{R}$ such that $\mathcal{S} = \{x \in \mathbb{F}^n: \text{Re } x^*y < \alpha\}$.

(Proof: Let $z \in \mathbb{F}^n$ satisfy $z^*y = \alpha$. Then, $\{x: x^*y = \alpha\} = \{y\}^\perp + z$.)

Fact 2.9.7. Let $x_1, \dots, x_k \in \mathbb{F}^n$. Then,

$$\text{aff } \{x_1, \dots, x_k\} = x_1 + \text{span } \{x_2 - x_1, \dots, x_k - x_1\}.$$

(Remark: See Fact 10.8.12.)

Fact 2.9.8. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is an affine subspace. Then, $\mathcal{S}$ is a subspace if and only if $0 \in \mathcal{S}$.

Fact 2.9.9. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be (cones, convex sets, convex cones, subspaces). Then, so are $\mathcal{S}_1 \cap \mathcal{S}_2$ and $\mathcal{S}_1 + \mathcal{S}_2$.

Fact 2.9.10. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be pointed convex cones. Then,

$$\text{co}(\mathcal{S}_1 \cup \mathcal{S}_2) = \mathcal{S}_1 + \mathcal{S}_2.$$

Fact 2.9.11. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then, $\mathcal{S}_1 \cup \mathcal{S}_2$ is a subspace if and only if either $\mathcal{S}_1 \subseteq \mathcal{S}_2$ or $\mathcal{S}_2 \subseteq \mathcal{S}_1$.

Fact 2.9.12. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$. Then,

$$(\text{span } \mathcal{S}_1) \cup (\text{span } \mathcal{S}_2) \subseteq \text{span}(\mathcal{S}_1 \cup \mathcal{S}_2)$$

and

$$\text{span}(\mathcal{S}_1 \cap \mathcal{S}_2) \subseteq (\text{span } \mathcal{S}_1) \cap (\text{span } \mathcal{S}_2).$$

(Proof: See [1184, p. 11].)

Fact 2.9.13. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then,

$$\text{span}(\mathcal{S}_1 \cup \mathcal{S}_2) = \mathcal{S}_1 + \mathcal{S}_2.$$

Therefore, $\mathcal{S}_1 + \mathcal{S}_2$ is the smallest subspace that contains $\mathcal{S}_1 \cup \mathcal{S}_2$.

Fact 2.9.14. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then, the following statements are equivalent:

- i) $\mathcal{S}_1 \subseteq \mathcal{S}_2$
- ii) $\mathcal{S}_2^\perp \subseteq \mathcal{S}_1^\perp$.
- iii) For all $x \in \mathcal{S}_1$ and $y \in \mathcal{S}_2^\perp$, $x^*y = 0$.

Furthermore, $\mathcal{S}_1 \subset \mathcal{S}_2$ if and only if $\mathcal{S}_2^\perp \subset \mathcal{S}_1^\perp$.

Fact 2.9.15. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$. Then,

$$\mathcal{S}_1^\perp \cap \mathcal{S}_2^\perp \subseteq (\mathcal{S}_1 + \mathcal{S}_2)^\perp.$$

(Problem: Determine necessary and sufficient conditions under which equality holds.)

Fact 2.9.16. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then,

$$(\mathcal{S}_1 \cap \mathcal{S}_2)^\perp = \mathcal{S}_1^\perp + \mathcal{S}_2^\perp$$

and

$$(\mathcal{S}_1 + \mathcal{S}_2)^\perp = \mathcal{S}_1^\perp \cap \mathcal{S}_2^\perp.$$

Fact 2.9.17. Let $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3 \subseteq \mathbb{F}^n$ be subspaces. Then,

$$\mathcal{S}_1 + (\mathcal{S}_2 \cap \mathcal{S}_3) \subseteq (\mathcal{S}_1 + \mathcal{S}_2) \cap (\mathcal{S}_1 + \mathcal{S}_3)$$

and

$$\mathcal{S}_1 \cap (\mathcal{S}_2 + \mathcal{S}_3) \supseteq (\mathcal{S}_1 \cap \mathcal{S}_2) + (\mathcal{S}_1 \cap \mathcal{S}_3).$$

Fact 2.9.18. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then, $\mathcal{S}_1, \mathcal{S}_2$ are complementary subspaces if and only if $\mathcal{S}_1^\perp, \mathcal{S}_2^\perp$ are complementary subspaces. (Remark: See Fact 3.12.1.)

Fact 2.9.19. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be nonzero subspaces, and define $\theta \in [0, \pi/2]$ by

$$\cos \theta = \max\{|x^*y| : (x, y) \in \mathcal{S}_1 \times \mathcal{S}_2 \text{ and } x^*x = y^*y = 1\}.$$

Then,

$$\cos \theta = \max\{|x^*y| : (x, y) \in \mathcal{S}_1^\perp \times \mathcal{S}_2^\perp \text{ and } x^*x = y^*y = 1\}.$$

Furthermore, $\theta = 0$ if and only if $\mathcal{S}_1 \cap \mathcal{S}_2 = \{0\}$, and $\theta = \pi/2$ if and only if $\mathcal{S}_1 = \mathcal{S}_2^\perp$. (Remark: See [537, 744].) (Remark: θ is a *principal angle*. See Fact 5.9.29, Fact 5.11.39, and Fact 5.12.17.)

Fact 2.9.20. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces, and assume that $\mathcal{S}_1 \cap \mathcal{S}_2 = \{0\}$. Then,

$$\dim \mathcal{S}_1 + \dim \mathcal{S}_2 \leq n.$$

Fact 2.9.21. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then,

$$\begin{aligned} \dim(\mathcal{S}_1 \cap \mathcal{S}_2) &\leq \min\{\dim \mathcal{S}_1, \dim \mathcal{S}_2\} \\ &\leq \begin{Bmatrix} \dim \mathcal{S}_1 \\ \dim \mathcal{S}_2 \end{Bmatrix} \\ &\leq \max\{\dim \mathcal{S}_1, \dim \mathcal{S}_2\} \\ &\leq \dim(\mathcal{S}_1 + \mathcal{S}_2) \\ &\leq \min\{\dim \mathcal{S}_1 + \dim \mathcal{S}_2, n\}. \end{aligned}$$

Fact 2.9.22. Let $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3 \subseteq \mathbb{F}^n$ be subspaces. Then,

$$\begin{aligned} \dim(\mathcal{S}_1 + \mathcal{S}_2 + \mathcal{S}_3) + \max\{\dim(\mathcal{S}_1 \cap \mathcal{S}_2), \dim(\mathcal{S}_1 \cap \mathcal{S}_3), \dim(\mathcal{S}_2 \cap \mathcal{S}_3)\} \\ \leq \dim \mathcal{S}_1 + \dim \mathcal{S}_2 + \dim \mathcal{S}_3. \end{aligned}$$

(Proof: See [392, p. 124].) (Remark: Setting $\mathcal{S}_3 = \{0\}$ yields a weaker version of Theorem 2.3.1.)

Fact 2.9.23. Let $\mathcal{S}_1, \dots, \mathcal{S}_k \subseteq \mathbb{F}^n$ be subspaces having the same dimension. Then, there exists a subspace $\hat{\mathcal{S}} \subseteq \mathbb{F}^n$ such that, for all $i = 1, \dots, k$, $\hat{\mathcal{S}}$ and $\mathcal{S}_i$ are complementary. (Proof: See [629, pp. 78, 79, 259, 260].)

Fact 2.9.24. Let $\mathcal{S} \subseteq \mathbb{F}^n$ be a subspace. Then, for all $m \geq \dim \mathcal{S}$, there exists a matrix $A \in \mathbb{F}^{n \times m}$ such that $\mathcal{S} = \mathcal{R}(A)$.

Fact 2.9.25. Let $A \in \mathbb{F}^{n \times n}$, let $\mathcal{S} \subseteq \mathbb{F}^n$, assume that $\mathcal{S}$ is a subspace, let $k \triangleq \dim \mathcal{S}$, let $S \in \mathbb{F}^{n \times k}$, and assume that $\mathcal{R}(S) = \mathcal{S}$. Then, $\mathcal{S}$ is an invariant subspace of A if and only if there exists a matrix $M \in \mathbb{F}^{k \times k}$ such that $AS = SM$. (Proof: Set $B = I$ in Fact 5.13.1. See [872, p. 99].)

Fact 2.9.26. Let $\mathcal{S} \subseteq \mathbb{F}^m$, and let $A \in \mathbb{F}^{n \times m}$. Then,

$$\begin{aligned} \text{cone } A\mathcal{S} &= A \text{ cone } \mathcal{S}, \\ \text{co } A\mathcal{S} &= A \text{ co } \mathcal{S}, \\ \text{span } A\mathcal{S} &= A \text{ span } \mathcal{S}, \\ \text{aff } A\mathcal{S} &= A \text{ aff } \mathcal{S}. \end{aligned}$$

Hence, if $\mathcal{S}$ is a (cone, convex set, subspace, affine subspace), then so is $A\mathcal{S}$. Now, assume that A is left invertible, and let $A^L \in \mathbb{F}^{m \times n}$ be a left inverse of A . Then,

$$\begin{aligned} \text{cone } \mathcal{S} &= A^L \text{ cone } A\mathcal{S}, \\ \text{co } \mathcal{S} &= A^L \text{ co } A\mathcal{S}, \\ \text{span } \mathcal{S} &= A^L \text{ span } A\mathcal{S}, \\ \text{aff } \mathcal{S} &= A^L \text{ aff } A\mathcal{S}. \end{aligned}$$

Hence, if $A\mathcal{S}$ is a (cone, convex set, subspace, affine subspace), then so is $\mathcal{S}$.

Fact 2.9.27. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and let $A \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

i) If A is right invertible and A^R is a right inverse of A , then

$$(A\mathcal{S})^\perp \subseteq A^{R*}\mathcal{S}^\perp.$$

ii) If A is left invertible and A^L is a left inverse of A , then

$$A\mathcal{S}^\perp \subseteq (A^{L*}\mathcal{S})^\perp.$$

iii) If $n = m$ and A is nonsingular, then

$$(A\mathcal{S})^\perp = A^{-*}\mathcal{S}^\perp.$$

(Proof: The third statement is an immediate consequence of the first two statements.)

Fact 2.9.28. Let $A \in \mathbb{F}^{n \times m}$, and let $\mathcal{S}_1 \subseteq \mathbb{R}^m$ and $\mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces. Then, the following statements are equivalent:

i) $A\mathcal{S}_1 \subseteq \mathcal{S}_2$.

ii) $A^*\mathcal{S}_2^\perp \subseteq \mathcal{S}_1^\perp$.

(Proof: See [311, p. 12].)

Fact 2.9.29. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^m$ be subspaces, and let $A \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

i) $A(\mathcal{S}_1 \cup \mathcal{S}_2) = A\mathcal{S}_1 \cup A\mathcal{S}_2$.

ii) $A(\mathcal{S}_1 \cap \mathcal{S}_2) \subseteq A\mathcal{S}_1 \cap A\mathcal{S}_2$.

iii) $A(\mathcal{S}_1 + \mathcal{S}_2) = A\mathcal{S}_1 + A\mathcal{S}_2$.

If, in addition, A is left invertible, then the following statement holds:

iv) $A(\mathcal{S}_1 \cap \mathcal{S}_2) = A\mathcal{S}_1 \cap A\mathcal{S}_2$.

(Proof: See Fact 1.5.11, Fact 1.5.14, and [311, p. 12].)

Fact 2.9.30. Let $\mathcal{S}, \mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be subspaces, let $A \in \mathbb{F}^{n \times m}$, and define $f: \mathbb{F}^m \mapsto \mathbb{F}^n$ by $f(x) \triangleq Ax$. Then, the following statements hold:

i) $f[f^{-1}(\mathcal{S})] \subseteq \mathcal{S} \subseteq f^{-1}[f(\mathcal{S})]$.

ii) $[f^{-1}(\mathcal{S})]^\perp = A^*\mathcal{S}^\perp$.

iii) $f^{-1}(\mathcal{S}_1 \cup \mathcal{S}_2) = f^{-1}(\mathcal{S}_1) \cup f^{-1}(\mathcal{S}_2)$.

iv) $f^{-1}(\mathcal{S}_1 \cap \mathcal{S}_2) = f^{-1}(\mathcal{S}_1) \cap f^{-1}(\mathcal{S}_2)$.

v) $f^{-1}(\mathcal{S}_1 + \mathcal{S}_2) \supseteq f^{-1}(\mathcal{S}_1) + f^{-1}(\mathcal{S}_2)$.

(Proof: See Fact 1.5.12 and [311, p. 12].) (Problem: For a subspace $\mathcal{S} \subseteq \mathbb{F}^n$, $A \in \mathbb{F}^{n \times m}$, and $f(x) \triangleq Ax$, determine $B \in \mathbb{F}^{m \times n}$ such that $f^{-1}(\mathcal{S}) = B\mathcal{S}$, that is, $AB\mathcal{S} \subseteq \mathcal{S}$ and $B\mathcal{S}$ is maximal.)

Fact 2.9.31. Define the convex pointed cone $\mathcal{S} \subset \mathbb{R}^2$ by

$$\mathcal{S} \triangleq \{(x_1, x_2) \in [0, \infty) \times \mathbb{R} : \text{if } x_1 = 0, \text{ then } x_2 \geq 0\},$$

that is,

$$\mathcal{S} = ([0, \infty) \times \mathbb{R}) \setminus [\{0\} \times (-\infty, 0)].$$

Furthermore, for $x, y \in \mathbb{R}^2$, define $x \stackrel{d}{\leq} y$ if and only if $y - x \in \mathcal{S}$. Then, “ $\stackrel{d}{\leq}$ ” is a total ordering on $\mathbb{R}^2$. (Remark: “ $\stackrel{d}{\leq}$ ” is the lexicographic or dictionary ordering. See Fact 1.5.8.) (Remark: See [153, p. 161].)

2.10 Facts on Range, Null Space, Rank, and Defect

Fact 2.10.1. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\mathcal{N}(A) \subseteq \mathcal{R}(I - A)$$

and

$$\mathcal{N}(I - A) \subseteq \mathcal{R}(A).$$

(Remark: See Fact 3.12.3.)

Fact 2.10.2. Let $A \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

- i) If $B \in \mathbb{F}^{m \times l}$ and $\text{rank } B = m$, then $\mathcal{R}(A) = \mathcal{R}(AB)$.
- ii) If $C \in \mathbb{F}^{k \times n}$ and $\text{rank } C = n$, then $\mathcal{N}(A) = \mathcal{N}(CA)$.
- iii) If $S \in \mathbb{F}^{m \times m}$ and S is nonsingular, then $\mathcal{N}(A) = \mathcal{N}(AS)$.

(Remark: See Lemma 2.4.1.)

Fact 2.10.3. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, the following statements hold:

- i) If A and B are right invertible, then so is AB .
- ii) If A and B are left invertible, then so is AB .
- iii) If $n = m = l$ and A and B are nonsingular, then so is AB .

(Proof: The result follows from either Corollary 2.5.10 or Proposition 2.6.3.) (Remark: See Fact 1.5.16.)

Fact 2.10.4. Let $\mathcal{S} \subseteq \mathbb{F}^m$, assume that $\mathcal{S}$ is an affine subspace, and let $A \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

- i) $\text{rank } A + \dim \mathcal{S} - m \leq \dim AS \leq \min\{\text{rank } A, \dim \mathcal{S}\}$.
- ii) $\dim(AS) + \dim[\mathcal{N}(A) \cap \mathcal{S}] = \dim \mathcal{S}$.
- iii) $\dim AS \leq \dim \mathcal{S}$.
- iv) If A is left invertible, then $\dim AS = \dim \mathcal{S}$.

(Proof: For ii), see [1129, p. 413]. For iii), note that $\dim AS \leq \dim \mathcal{S} = \dim A^L AS \leq \dim AS$.) (Remark: See Fact 2.9.26 and Fact 10.8.17.)

Fact 2.10.5. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{1 \times m}$. Then, $\mathcal{N}(A) \subseteq \mathcal{N}(B)$ if and only if there exists a vector $\lambda \in \mathbb{F}^n$ such that $B = \lambda^* A$.

Fact 2.10.6. Let $A \in \mathbb{F}^{n \times m}$ and $b \in \mathbb{F}^n$. Then, there exists a vector $x \in \mathbb{F}^m$ satisfying $Ax = b$ if and only if $b^* \lambda = 0$ for all $\lambda \in \mathcal{N}(A^*)$. (Proof: Assume that $A^* \lambda = 0$ implies that $b^* \lambda = 0$. Then, $\mathcal{N}(A^*) \subseteq \mathcal{N}(b^*)$. Hence, $b \in \mathcal{R}(b) \subseteq \mathcal{R}(A)$.)

Fact 2.10.7. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times m}$. Then, $\mathcal{N}(B) \subseteq \mathcal{N}(A)$ if and only if there exists a matrix $C \in \mathbb{F}^{n \times l}$ such that $A = CB$. Now, let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then, $\mathcal{R}(A) \subseteq \mathcal{R}(B)$ if and only if there exists a matrix $C \in \mathbb{F}^{l \times m}$ such that $A = BC$.

Fact 2.10.8. Let $A, B \in \mathbb{F}^{n \times m}$, and let $C \in \mathbb{F}^{m \times l}$ be right invertible. Then, $\mathcal{R}(A) \subseteq \mathcal{R}(B)$ if and only if $\mathcal{R}(AC) \subseteq \mathcal{R}(BC)$. Furthermore, $\mathcal{R}(A) = \mathcal{R}(B)$ if and only if $\mathcal{R}(AC) = \mathcal{R}(BC)$. (Proof: Since C is right invertible, it follows that $\mathcal{R}(A) = \mathcal{R}(AC)$.)

Fact 2.10.9. Let $A, B \in \mathbb{F}^{n \times n}$, and assume there exists $\alpha \in \mathbb{F}$ such that $\alpha A + B$ is nonsingular. Then, $\mathcal{N}(A) \cap \mathcal{N}(B) = \{0\}$. (Remark: The converse is not true. Let $A \triangleq \begin{bmatrix} 1 & 0 \\ 2 & 0 \end{bmatrix}$ and $B \triangleq \begin{bmatrix} 0 & 1 \\ 0 & 2 \end{bmatrix}$.)

Fact 2.10.10. Let $A, B \in \mathbb{F}^{n \times m}$, and let $\alpha \in \mathbb{F}$ be nonzero. Then,

$$\mathcal{N}(A) \cap \mathcal{N}(B) = \mathcal{N}(A) \cap \mathcal{N}(A + \alpha B) = \mathcal{N}(\alpha A + B) \cap \mathcal{N}(B).$$

(Remark: See Fact 2.11.3.)

Fact 2.10.11. Let $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$. If either $x = 0$ or $y \neq 0$, then

$$\mathcal{R}(xy^T) = \mathcal{R}(x) = \text{span}\{x\}.$$

Furthermore, if either $x \neq 0$ or $y = 0$, then

$$\mathcal{N}(xy^T) = \mathcal{N}(y^T) = \{\bar{y}\}^\perp.$$

Fact 2.10.12. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, $\text{rank } AB = \text{rank } A$ if and only if $\mathcal{R}(AB) = \mathcal{R}(A)$. (Proof: If $\mathcal{R}(AB) \subset \mathcal{R}(A)$ (note proper inclusion), then Lemma 2.3.4 implies that $\text{rank } AB < \text{rank } A$.)

Fact 2.10.13. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $C \in \mathbb{F}^{l \times k}$. If $\text{rank } AB = \text{rank } B$, then $\text{rank } ABC = \text{rank } BC$. (Proof: $\text{rank } B^T A^T = \text{rank } B^T$ implies that $\mathcal{R}(C^T B^T A^T) = \mathcal{R}(C^T B^T)$.)

Fact 2.10.14. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, the following statements hold:

- i) $\text{rank } AB + \text{def } A = \dim[\mathcal{N}(A) + \mathcal{R}(B)]$.
- ii) $\text{rank } AB + \dim[\mathcal{N}(A) \cap \mathcal{R}(B)] = \text{rank } B$.
- iii) $\text{rank } AB + \dim[\mathcal{N}(A^*) \cap \mathcal{R}(B^*)] = \text{rank } A$.
- iv) $\text{def } AB + \text{rank } A + \dim[\mathcal{N}(A) + \mathcal{R}(B)] = l + m$.

$$v) \text{ def } AB = \text{def } B + \dim[\mathcal{N}(A) \cap \mathcal{R}(B)].$$

$$vi) \text{ def } AB + m = \text{def } A + \dim[\mathcal{N}(A^*) \cap \mathcal{R}(B^*)] + l.$$

(Remark: $\text{rank } B - \text{rank } AB = \dim[\mathcal{N}(A) \cap \mathcal{R}(B)] \leq \dim \mathcal{N}(A) = m - \text{rank } A$ yields (2.5.18).)

Fact 2.10.15. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\max\{\text{def } A + l - m, \text{def } B\} \leq \text{def } AB \leq \text{def } A + \text{def } B.$$

If, in addition, $m = l$, then

$$\max\{\text{def } A, \text{def } B\} \leq \text{def } AB.$$

(Remark: The first inequality is *Sylvester's law of nullity*.)

Fact 2.10.16. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times p}$. Then, there exists a matrix $X \in \mathbb{F}^{m \times p}$ satisfying $AX = B$ and $\text{rank } X = q$ if and only if

$$\text{rank } B \leq q \leq \min\{m + \text{rank } B - \text{rank } A, p\}.$$

(Proof: See [1353].)

Fact 2.10.17. The following statements hold:

$$i) \text{ rank } A \geq 0 \text{ for all } A \in \mathbb{F}^{n \times m}.$$

$$ii) \text{ rank } A = 0 \text{ if and only if } A = 0.$$

$$iii) \text{ rank } \alpha A = (\text{sign } |\alpha|) \text{rank } A \text{ for all } \alpha \in \mathbb{F} \text{ and } A \in \mathbb{F}^{n \times m}.$$

$$iv) \text{ rank}(A + B) \leq \text{rank } A + \text{rank } B \text{ for all } A, B \in \mathbb{F}^{n \times m}.$$

(Remark: Compare these conditions to the properties of a matrix norm given by Definition 9.2.1.)

Fact 2.10.18. Let $n, m, k \in \mathbb{P}$. Then, $\text{rank } 1_{n \times m} = 1$ and $1_{n \times n}^k = n^{k-1} 1_{n \times n}$.

Fact 2.10.19. Let $A \in \mathbb{F}^{n \times m}$. Then, $\text{rank } A = 1$ if and only if there exist vectors $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$ such that $x \neq 0$, $y \neq 0$, and $A = xy^T$. In this case, $\text{tr } A = y^T x$. (Remark: See Fact 5.14.1.)

Fact 2.10.20. Let $A \in \mathbb{F}^{n \times n}$, $k \geq 1$, and $l \in \mathbb{N}$. Then, the following identities hold:

$$i) \mathcal{R}[(AA^*)^k] = \mathcal{R}[(AA^*)^l A].$$

$$ii) \mathcal{N}[(A^*A)^k] = \mathcal{N}[A(A^*A)^l].$$

$$iii) \text{rank } (AA^*)^k = \text{rank } (AA^*)^l A.$$

$$iv) \text{def } (A^*A)^k = \text{def } A(A^*A)^l.$$

Fact 2.10.21. Let $A \in \mathbb{F}^{n \times m}$, and let $B \in \mathbb{F}^{m \times p}$. Then,

$$\text{rank } AB = \text{rank } A^*AB = \text{rank } ABB^*.$$

(Proof: See [1184, p. 37].)

Fact 2.10.22. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$2\text{rank } A^2 \leq \text{rank } A + \text{rank } A^3.$$

(Proof: See [392, p. 126] and consider a Jordan block of A .)

Fact 2.10.23. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{rank } A + \text{rank}(A - A^3) = \text{rank}(A + A^2) + \text{rank}(A - A^2).$$

Consequently,

$$\text{rank } A \leq \text{rank}(A + A^2) + \text{rank}(A - A^2),$$

and A is tripotent if and only if

$$\text{rank } A = \text{rank}(A + A^2) + \text{rank}(A - A^2).$$

(Proof: See [1308].) (Remark: This result is due to Anderson and Styan.)

Fact 2.10.24. Let $x, y \in \mathbb{F}^n$. Then,

$$\mathcal{R}(xy^T + yx^T) = \mathcal{R}\left(\begin{bmatrix} x & y \end{bmatrix}\right),$$

$$\mathcal{N}(xy^T + yx^T) = \{x\}^\perp \cap \{y\}^\perp,$$

$$\text{rank}(xy^T + yx^T) \leq 2.$$

Furthermore, $\text{rank}(xy^T + yx^T) = 1$ if and only if there exists $\alpha \in \mathbb{F}$ such that $x = \alpha y \neq 0$. (Remark: $xy^T + yx^T$ is a *doublet*. See [374, pp. 539, 540].)

Fact 2.10.25. Let $A \in \mathbb{F}^{n \times m}$, $x \in \mathbb{F}^n$, and $y \in \mathbb{F}^m$. Then,

$$(\text{rank } A) - 1 \leq \text{rank}(A + xy^*) \leq (\text{rank } A) + 1.$$

(Remark: See Fact 6.4.2.)

Fact 2.10.26. Let $A \triangleq \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $B \triangleq \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$. Then, $\text{rank } AB = 1$ and $\text{rank } BA = 0$. (Remark: See Fact 3.7.30.)

Fact 2.10.27. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$|\text{rank } A - \text{rank } B| \leq \left\{ \begin{array}{l} \text{rank}(A + B) \\ \text{rank}(A - B) \end{array} \right\} \leq \text{rank } A + \text{rank } B.$$

If, in addition, $\text{rank } B \leq k$, then

$$(\text{rank } A) - k \leq \left\{ \begin{array}{l} \text{rank}(A + B) \\ \text{rank}(A - B) \end{array} \right\} \leq (\text{rank } A) + k.$$

Fact 2.10.28. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

$$i) \text{ rank}(A + B) = \text{rank } A + \text{rank } B.$$

$$ii) \mathcal{R}(A) \cap \mathcal{R}(B) = \{0\} \text{ and } \mathcal{R}(A^T) \cap \mathcal{R}(B^T) = \{0\}.$$

(Proof: See [281].) (Remark: See Fact 2.10.29.)

Fact 2.10.29. Let $A, B \in \mathbb{F}^{n \times m}$, and assume that $A^*B = 0$ and $BA^* = 0$. Then,

$$\text{rank}(A + B) = \text{rank } A + \text{rank } B.$$

(Proof: Use Fact 2.11.4 and Proposition 6.1.6. See [339].) (Remark: See Fact 2.10.28.)

Fact 2.10.30. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

- i) $\text{rank}(B - A) = \text{rank } B - \text{rank } A$.
- ii) There exists $M \in \mathbb{F}^{m \times n}$ such that $A = BMB$ and $M = MBM$.
- iii) There exists $M \in \mathbb{F}^{m \times n}$ such that $B = BMB$, $MA = 0$, and $AM = 0$.
- iv) There exists $M \in \mathbb{F}^{m \times n}$ such that $A = AMA$, $MB = 0$, and $BM = 0$.

(Proof: See [339].)

Fact 2.10.31. Let $A, B, C \in \mathbb{F}^{n \times m}$, and assume that

$$\text{rank}(B - A) = \text{rank } B - \text{rank } A$$

and

$$\text{rank}(C - B) = \text{rank } C - \text{rank } B.$$

Then,

$$\text{rank}(C - A) = \text{rank } C - \text{rank } A.$$

(Proof: $\text{rank}(C - A) \leq \text{rank}(C - B) + \text{rank}(B - A) = \text{rank } C - \text{rank } A$. Furthermore, $\text{rank } C \leq \text{rank}(C - A) + \text{rank } A$, and thus $\text{rank}(C - A) \geq \text{rank } C - \text{rank } A$. Alternatively, use Fact 2.10.30.) (Remark: This result is due to [647].)

Fact 2.10.32. Let $A, B \in \mathbb{F}^{n \times m}$, and define

$$A \overset{\text{rs}}{\leq} B$$

if and only if

$$\text{rank}(B - A) = \text{rank } B - \text{rank } A.$$

Then, " $\overset{\text{rs}}{\leq}$ " is a partial ordering on $\mathbb{F}^{n \times m}$. (Proof: Use Fact 2.10.31.) (Remark: The relation " $\overset{\text{rs}}{\leq}$ " is the *rank subtractivity partial ordering*.) (Remark: See Fact 8.19.5.)

Fact 2.10.33. Let $A, B \in \mathbb{F}^{n \times m}$, and assume that the following conditions hold:

- i) $A^*A = A^*B$.
- ii) $AA^* = BA^*$.
- iii) $B^*B = B^*A$.
- iv) $BB^* = AB^*$.

Then, $A = B$. (Proof: See [652].)

Fact 2.10.34. Let $A, B, C \in \mathbb{F}^{n \times m}$, and assume that the following conditions hold:

- i) $A^*A = A^*B$.
- ii) $AA^* = BA^*$.
- iii) $B^*B = B^*C$.
- iv) $BB^* = CB^*$.

Then, the following conditions hold:

- v) $A^*A = A^*C$.
- vi) $AA^* = CA^*$.

(Proof: See [652].)

Fact 2.10.35. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$A \leq^* B$$

if and only if

$$A^*A = A^*B$$

and

$$AA^* = BA^*.$$

Then, " $\leq^*$ " is a partial ordering on $\mathbb{F}^{n \times m}$. (Proof: Use Fact 2.10.33 and Fact 2.10.34.)

(Remark: The relation " $\leq^*$ " is the *star partial ordering*. See [111, 652].) (Remark: See Fact 8.19.7.)

Fact 2.10.36. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $A \leq^* B$ and $AB = BA$. Then, $A^2 \leq^* B^2$. (Proof: See [106].) (Remark: See Fact 8.19.5.)

2.11 Facts on the Range, Rank, Null Space, and Defect of Partitioned Matrices

Fact 2.11.1. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\mathcal{R}\left[\begin{array}{cc} A & B \end{array}\right] = \mathcal{R}(A) + \mathcal{R}(B).$$

Consequently,

$$\text{rank}\left[\begin{array}{cc} A & B \end{array}\right] = \dim[\mathcal{R}(A) + \mathcal{R}(B)].$$

Furthermore, the followings statements are equivalent:

- i) $\text{rank}\left[\begin{array}{cc} A & B \end{array}\right] = n$.
- ii) $\text{def}\left[\begin{array}{c} A^* \\ B^* \end{array}\right] = 0$.
- iii) $\mathcal{N}(A^*) \cap \mathcal{N}(B^*) = \{0\}$.

Fact 2.11.2. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times m}$. Then,

$$\text{rank} \begin{bmatrix} A \\ B \end{bmatrix} = \dim[\mathcal{R}(A^*) + \mathcal{R}(B^*)].$$

(Proof: Use Fact 2.11.1.)

Fact 2.11.3. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times m}$. Then,

$$\mathcal{N}\left(\begin{bmatrix} A \\ B \end{bmatrix}\right) = \mathcal{N}(A) \cap \mathcal{N}(B).$$

Consequently,

$$\text{def} \begin{bmatrix} A \\ B \end{bmatrix} = \dim[\mathcal{N}(A) \cap \mathcal{N}(B)].$$

Furthermore, the followings statements are equivalent:

$$i) \text{ rank} \begin{bmatrix} A \\ B \end{bmatrix} = m.$$

$$ii) \text{ def} \begin{bmatrix} A \\ B \end{bmatrix} = 0.$$

$$iii) \mathcal{N}(A) \cap \mathcal{N}(B) = \{0\}.$$

(Remark: See Fact 2.10.10.)

Fact 2.11.4. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

$$i) \text{ rank}(A + B) = \text{rank } A + \text{rank } B.$$

$$ii) \text{ rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank} \begin{bmatrix} A \\ B \end{bmatrix} = \text{rank } A + \text{rank } B.$$

$$iii) \dim[\mathcal{R}(A) \cap \mathcal{R}(B)] = \dim[\mathcal{R}(A^*) \cap \mathcal{R}(B^*)] = 0.$$

$$iv) \mathcal{R}(A) \cap \mathcal{R}(B) = \mathcal{R}(A^*) \cap \mathcal{R}(B^*) = \{0\}.$$

$$v) \text{ There exists a matrix } C \in \mathbb{F}^{m \times n} \text{ such that } ACA = A, CB = 0, \text{ and } BC = 0.$$

(Proof: See [339, 968].) (Remark: Additional conditions are given by Fact 6.4.32 under the assumption that $A + B$ is nonsingular.)

Fact 2.11.5. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\mathcal{R}(A) = \mathcal{R}(B)$$

if and only if

$$\text{rank } A = \text{rank } B = \text{rank} \begin{bmatrix} A & B \end{bmatrix}.$$

Fact 2.11.6. Let $A \in \mathbb{F}^{n \times m}$, and let $A_0 \in \mathbb{F}^{k \times l}$ be a submatrix of A . Then,

$$\text{rank } A_0 \leq \text{rank } A.$$

Fact 2.11.7. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{k \times m}$, $C \in \mathbb{F}^{m \times l}$, and $D \in \mathbb{F}^{m \times p}$, and assume that

$$\text{rank} \begin{bmatrix} A \\ B \end{bmatrix} = \text{rank } A$$

and

$$\text{rank} \begin{bmatrix} C & D \end{bmatrix} = \text{rank } C.$$

Then,

$$\text{rank} \begin{bmatrix} A \\ B \end{bmatrix} \begin{bmatrix} C & D \end{bmatrix} = \text{rank } AC.$$

(Proof: Use *i*) of Fact 2.10.14.)

Fact 2.11.8. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\begin{aligned} \max\{\text{rank } A, \text{rank } B\} &\leq \text{rank} \begin{bmatrix} A & B \end{bmatrix} \\ &= \text{rank } A + \text{rank } B - \dim[\mathcal{R}(A) \cap \mathcal{R}(B)] \\ &\leq \text{rank } A + \text{rank } B \end{aligned}$$

and

$$\begin{aligned} \text{def } A + \text{def } B &\leq \text{def} \begin{bmatrix} A & B \end{bmatrix} \\ &= \text{def } A + \text{def } B + \dim[\mathcal{R}(A) \cap \mathcal{R}(B)] \\ &\leq \min\{l + \text{def } A, m + \text{def } B\}. \end{aligned}$$

If, in addition, $A^*B = 0$, then

$$\text{rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank } A + \text{rank } B$$

and

$$\text{def} \begin{bmatrix} A & B \end{bmatrix} = \text{def } A + \text{def } B.$$

(Proof: To prove the first equality, use Theorem 2.3.1 and Fact 2.11.1. For the case $A^*B = 0$, note that

$$\begin{aligned} \text{rank} \begin{bmatrix} A & B \end{bmatrix} &= \text{rank} \begin{bmatrix} A^* \\ B^* \end{bmatrix} \begin{bmatrix} A & B \end{bmatrix} = \begin{bmatrix} A^*A & 0 \\ 0 & B^*B \end{bmatrix} \\ &= \text{rank } A^*A + \text{rank } B^*B = \text{rank } A + \text{rank } B. \end{aligned}$$

(Remark: See Fact 6.5.6 and Fact 6.4.44.)

Fact 2.11.9. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\text{rank} \begin{bmatrix} A & B \end{bmatrix} + \dim[\mathcal{R}(A) \cap \mathcal{R}(B)] = \text{rank } A + \text{rank } B.$$

(Proof: Use Theorem 2.3.1 and Fact 2.11.1.)

Fact 2.11.10. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times m}$. Then,

$$\text{rank} \begin{bmatrix} A \\ B \end{bmatrix} + \dim[\mathcal{R}(A^*) \cap \mathcal{R}(B^*)] = \text{rank } A + \text{rank } B.$$

(Proof: Use Fact 2.11.9.)

Fact 2.11.11. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times m}$. Then,

$$\begin{aligned} \max\{\text{rank } A, \text{rank } B\} &\leq \text{rank} \begin{bmatrix} A \\ B \end{bmatrix} \\ &= \text{rank } A + \text{rank } B - \dim[\mathcal{R}(A^*) \cap \mathcal{R}(B^*)] \\ &\leq \text{rank } A + \text{rank } B \end{aligned}$$

and

$$\begin{aligned} \text{def } A - \text{rank } B &\leq \text{def } A - \text{rank } B + \dim[\mathcal{R}(A^*) \cap \mathcal{R}(B^*)] \\ &= \text{def} \begin{bmatrix} A \\ B \end{bmatrix} \\ &\leq \min\{\text{def } A, \text{def } B\}. \end{aligned}$$

If, in addition, $AB^* = 0$, then

$$\text{rank} \begin{bmatrix} A \\ B \end{bmatrix} = \text{rank } A + \text{rank } B$$

and

$$\text{def} \begin{bmatrix} A \\ B \end{bmatrix} = \text{def } A - \text{rank } B.$$

(Proof: Use Fact 2.11.8 and Fact 2.9.21.) (Remark: See Fact 6.5.6.)

Fact 2.11.12. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\left. \begin{array}{l} \max\{\text{rank } A, \text{rank } B\} \\ \text{rank}(A + B) \end{array} \right\} \leq \left\{ \begin{array}{l} \text{rank} \begin{bmatrix} A & B \end{bmatrix} \\ \text{rank} \begin{bmatrix} A \\ B \end{bmatrix} \end{array} \right\} \leq \text{rank } A + \text{rank } B$$

and

$$\text{def } A - \text{rank } B \leq \left\{ \begin{array}{l} \text{def} \begin{bmatrix} A & B \end{bmatrix} - m \\ \text{def} \begin{bmatrix} A \\ B \end{bmatrix} \end{array} \right\} \leq \left\{ \begin{array}{l} \min\{\text{def } A, \text{def } B\} \\ \text{def}(A + B). \end{array} \right.$$

(Proof: $\text{rank}(A + B) = \text{rank} \begin{bmatrix} A & B \end{bmatrix} \begin{bmatrix} I \\ I \end{bmatrix} \leq \text{rank} \begin{bmatrix} A & B \end{bmatrix}$, and $\text{rank}(A + B) = \text{rank} \begin{bmatrix} I & I \end{bmatrix} \begin{bmatrix} A \\ B \end{bmatrix} \leq \text{rank} \begin{bmatrix} A \\ B \end{bmatrix}$.)

Fact 2.11.13. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{l \times k}$, and $C \in \mathbb{F}^{l \times m}$. Then,

$$\text{rank } A + \text{rank } B = \text{rank} \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \leq \text{rank} \begin{bmatrix} A & 0 \\ C & B \end{bmatrix}$$

and

$$\text{rank } A + \text{rank } B = \text{rank} \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix} \leq \text{rank} \begin{bmatrix} 0 & A \\ B & C \end{bmatrix}.$$

Fact 2.11.14. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $C \in \mathbb{F}^{l \times k}$. Then,

$$\text{rank } AB + \text{rank } BC \leq \text{rank} \begin{bmatrix} 0 & AB \\ BC & B \end{bmatrix} = \text{rank } B + \text{rank } ABC.$$

Consequently,

$$\text{rank } AB + \text{rank } BC - \text{rank } B \leq \text{rank } ABC.$$

Furthermore, the following statements are equivalent:

- i) $\text{rank} \begin{bmatrix} 0 & AB \\ BC & B \end{bmatrix} = \text{rank } AB + \text{rank } BC.$
- ii) $\text{rank } AB + \text{rank } BC - \text{rank } B = \text{rank } ABC.$
- iii) There exist $X \in \mathbb{F}^{k \times l}$ and $Y \in \mathbb{F}^{m \times n}$ such that

$$BCX + YAB = B.$$

(Remark: This result is the *Frobenius inequality*.) (Proof: Use Fact 2.11.13 and $\begin{bmatrix} 0 & AB \\ BC & B \end{bmatrix} = \begin{bmatrix} I & A \\ 0 & I \end{bmatrix} \begin{bmatrix} -ABC & 0 \\ 0 & B \end{bmatrix} \begin{bmatrix} I & 0 \\ C & I \end{bmatrix}$. The last statement follows from Fact 5.10.21. See [1307, 1308].) (Remark: See Fact 6.5.15 for the case of equality.)

Fact 2.11.15. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\begin{aligned} \text{rank} \begin{bmatrix} A & B \end{bmatrix} + \text{rank} \begin{bmatrix} A \\ B \end{bmatrix} &\leq \text{rank} \begin{bmatrix} 0 & A & B \\ A & A & 0 \\ B & 0 & B \end{bmatrix} \\ &= \text{rank } A + \text{rank } B + \text{rank}(A + B). \end{aligned}$$

(Proof: Use the Frobenius inequality with $A \triangleq C^T \triangleq \begin{bmatrix} I & I \end{bmatrix}$ and with B replaced by $\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$.)

Fact 2.11.16. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, and $C \in \mathbb{F}^{n \times k}$. Then,

$$\begin{aligned} \text{rank} \begin{bmatrix} A & B & C \end{bmatrix} &\leq \text{rank} \begin{bmatrix} A & B \end{bmatrix} + \text{rank} \begin{bmatrix} B & C \end{bmatrix} - \text{rank } B \\ &\leq \text{rank} \begin{bmatrix} A & B \end{bmatrix} + \text{rank } C \\ &\leq \text{rank } A + \text{rank } B + \text{rank } C. \end{aligned}$$

(Proof: See [937].)

Fact 2.11.17. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{k \times l}$, and assume that B is a submatrix of A . Then,

$$k + l - \text{rank } B \leq n + m - \text{rank } A.$$

(Proof: See [134].)

Fact 2.11.18. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then,

$$\begin{aligned} \begin{bmatrix} I_n & I_n - AB \\ B & 0 \end{bmatrix} &= \begin{bmatrix} I_n & A \\ 0 & I_m \end{bmatrix} \begin{bmatrix} 0 & I_n - AB \\ B & 0 \end{bmatrix} \begin{bmatrix} I_n & 0 \\ I_n & I_n \end{bmatrix} \\ &= \begin{bmatrix} I_n & 0 \\ B & I_m \end{bmatrix} \begin{bmatrix} I_n & 0 \\ 0 & BAB - B \end{bmatrix} \begin{bmatrix} I_n & I_n - AB \\ 0 & I_m \end{bmatrix}. \end{aligned}$$

Hence,

$$\text{rank} \begin{bmatrix} I_n & I_n - AB \\ B & 0 \end{bmatrix} = \text{rank } B + \text{rank}(I_n - AB) = n + \text{rank}(BAB - B).$$

(Remark: See Fact 2.14.7.)

Fact 2.11.19. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then,

$$\begin{aligned} \begin{bmatrix} A & AB \\ BA & B \end{bmatrix} &= \begin{bmatrix} I_n & 0 \\ B & I_m \end{bmatrix} \begin{bmatrix} A & 0 \\ 0 & B - BAB \end{bmatrix} \begin{bmatrix} I_m & B \\ 0 & I_n \end{bmatrix} \\ &= \begin{bmatrix} I_n & A \\ 0 & I_m \end{bmatrix} \begin{bmatrix} A - ABA & 0 \\ 0 & B \end{bmatrix} \begin{bmatrix} I_m & 0 \\ A & I_n \end{bmatrix}. \end{aligned}$$

Hence,

$$\text{rank} \begin{bmatrix} A & AB \\ BA & B \end{bmatrix} = \text{rank } A + \text{rank}(B - BAB) = \text{rank } B + \text{rank}(A - ABA).$$

(Remark: See Fact 2.14.10.)

Fact 2.11.20. Let $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{F}^{(n_1+n_2) \times (m_1+m_2)}$, assume that $\begin{bmatrix} A & B \\ C & D \end{bmatrix}$ is nonsingular, and define $\begin{bmatrix} E & F \\ G & H \end{bmatrix} \in \mathbb{F}^{(m_1+m_2) \times (n_1+n_2)}$ by

$$\begin{bmatrix} E & F \\ G & H \end{bmatrix} \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix}^{-1}.$$

Then,

$$\begin{aligned} \text{def } A &= \text{def } H, \\ \text{def } B &= \text{def } F, \\ \text{def } C &= \text{def } G, \\ \text{def } D &= \text{def } E. \end{aligned}$$

More generally, if U and V are complementary submatrices of a matrix and its inverse, then $\text{def } U = \text{def } V$. (Proof: See [1242, 1364] and [1365, p. 38].) (Remark: U and V are *complementary submatrices* if the row numbers not used to create U are the column numbers used to create V , and the column numbers not used to create U are the row numbers used to create V .) (Remark: Note the sizes of the matrix blocks, which differs from Fact 2.14.28.) (Remark: This result is the *nullity theorem*. A history of this result is given in [1242]. See Fact 3.20.5.)

Fact 2.11.21. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $S_1, S_2 \subseteq \{1, \dots, n\}$. Then,

$$\text{rank}(A^{-1})_{(S_1, S_2)} = \text{rank } A_{(S_2^{\sim}, S_1^{\sim})} + \text{card}(S_1) + \text{card}(S_2) - n.$$

(Proof: See [1365, p. 40].) (Remark: See Fact 2.11.22 and Fact 2.13.5.)

Fact 2.11.22. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $S \subseteq \{1, \dots, n\}$. Then,

$$\text{rank}(A^{-1})_{(S, S^{\sim})} = \text{rank } A_{(S, S^{\sim})}.$$

(Proof: Apply Fact 2.11.21 with $S_2 = S_1^{\sim}$.)

2.12 Facts on the Inner Product, Outer Product, Trace, and Matrix Powers

Fact 2.12.1. Let $x, y, z \in \mathbb{F}^n$, and assume that $x^*x = y^*y = z^*z = 1$. Then,

$$\sqrt{1 - |x^*y|^2} \leq \sqrt{1 - |x^*z|^2} + \sqrt{1 - |z^*y|^2}.$$

Equality holds if and only if there exists $\alpha \in \mathbb{F}$ such that either $z = \alpha x$ or $z = \alpha y$. (Proof: See [1490, p. 155].) (Remark: See Fact 3.11.32.)

Fact 2.12.2. Let $x, y \in \mathbb{F}^n$. Then, $x^*x = y^*y$ and $\text{Im } x^*y = 0$ if and only if $x - y$ is orthogonal to $x + y$.

Fact 2.12.3. Let $x, y \in \mathbb{R}^n$. Then, $xx^T = yy^T$ if and only if either $x = y$ or $x = -y$.

Fact 2.12.4. Let $x, y \in \mathbb{R}^n$. Then, $xy^T = yx^T$ if and only if x and y are linearly dependent.

Fact 2.12.5. Let $x, y \in \mathbb{R}^n$. Then, $xy^T = -yx^T$ if and only if either $x = 0$ or $y = 0$. (Proof: If $x_{(i)} \neq 0$ and $y_{(j)} \neq 0$, then $x_{(j)} = y_{(i)} = 0$ and $0 \neq x_{(i)}y_{(j)} \neq x_{(j)}y_{(i)} = 0$.)

Fact 2.12.6. Let $x, y \in \mathbb{R}^n$. Then, $yx^T + xy^T = y^T y x x^T$ if and only if either $x = 0$ or $y = \frac{1}{2}y^T y x$.

Fact 2.12.7. Let $x, y \in \mathbb{F}^n$. Then,

$$(xy^*)^r = (y^*x)^{r-1}xy^*.$$

Fact 2.12.8. Let $x_1, \dots, x_k \in \mathbb{F}^n$, and let $y_1, \dots, y_k \in \mathbb{F}^m$. Then, the following statements are equivalent:

i) $x_1, \dots, x_k$ are linearly independent, and $y_1, \dots, y_k$ are linearly independent.

ii) $\mathcal{R}\left(\sum_{i=1}^k x_i y_i^T\right) = k$.

(Proof: See [374, p. 537].)

Fact 2.12.9. Let $A, B, C \in \mathbb{R}^{2 \times 2}$. Then,

$$\begin{aligned} \text{tr}(ABC + ACB) + (\text{tr } A)(\text{tr } B)\text{tr } C \\ = (\text{tr } A)\text{tr } BC + (\text{tr } B)\text{tr } AC + (\text{tr } C)\text{tr } AB. \end{aligned}$$

(Proof: See [269, p. 330].) (Remark: See Fact 4.9.3.)

Fact 2.12.10. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times k}$. Then,

$$AE_{i,j,m \times l}B = \text{col}_i(A)\text{row}_j(B).$$

Fact 2.12.11. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $C \in \mathbb{F}^{l \times n}$. Then,

$$\operatorname{tr} ABC = \sum_{i=1}^n \operatorname{row}_i(A) B \operatorname{col}_i(C).$$

Fact 2.12.12. Let $A \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

- i) $A = 0$.
- ii) $Ax = 0$ for all $x \in \mathbb{F}^m$.
- iii) $\operatorname{tr} AA^* = 0$.

Fact 2.12.13. Let $A \in \mathbb{F}^{n \times n}$ and $k \geq 1$. Then,

$$\operatorname{Re} \operatorname{tr} A^{2k} \leq \operatorname{tr} A^k A^{k*} \leq \operatorname{tr} (AA^*)^k.$$

(Remark: To prove the left-hand inequality, consider $\operatorname{tr} (A^k - A^{k*})(A^{k*} - A^k)$. For the right-hand inequality when $k = 2$, consider $\operatorname{tr} (AA^* - A^*A)^2$.)

Fact 2.12.14. Let $A \in \mathbb{F}^{n \times n}$. Then, $\operatorname{tr} A^k = 0$ for all $k = 1, \dots, n$ if and only if $A^n = 0$. (Proof: For sufficiency, Fact 4.10.6 implies that $\operatorname{spec}(A) = \{0\}$, and thus the Jordan form of A is a block-diagonal matrix each of whose diagonally located blocks is a standard nilpotent matrix. For necessity, see [1490, p. 112].)

Fact 2.12.15. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\operatorname{tr} A = 0$. If $A^2 = A$, then $A = 0$. If $A^k = A$, where $k \geq 4$ and $2 \leq n < p$, where p is the smallest prime divisor of $k - 1$, then $A = 0$. (Proof: See [344].)

Fact 2.12.16. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\operatorname{Re} \operatorname{tr} AB \leq \frac{1}{2} \operatorname{tr} (AA^* + BB^*).$$

(Proof: See [729].) (Remark: See Fact 8.12.18.)

Fact 2.12.17. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $AB = 0$. Then, for all $k \geq 1$,

$$\operatorname{tr} (A + B)^k = \operatorname{tr} A^k + \operatorname{tr} B^k.$$

Fact 2.12.18. Let $A \in \mathbb{R}^{n \times n}$, let $x, y \in \mathbb{R}^n$, and let $k \geq 1$. Then,

$$(A + xy^T)^k = A^k + B \hat{I}_k C^T,$$

where

$$B \triangleq \begin{bmatrix} x & Ax & \cdots & A^{k-1}x \end{bmatrix}$$

and

$$C \triangleq \begin{bmatrix} y & (A^T + yx^T)y & \cdots & (A^T + yx^T)^k y \end{bmatrix}.$$

(Proof: See [192].)

Fact 2.12.19. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) $AB + BA = \frac{1}{2}[(A + B)^2 - (A - B)^2]$.
- ii) $(A + B)(A - B) = A^2 - B^2 - [A, B]$.

$$iii) (A - B)(A + B) = A^2 - B^2 + [A, B].$$

$$iv) A^2 - B^2 = \frac{1}{2}[(A + B)(A - B) + (A - B)(A + B)].$$

Fact 2.12.20. Let $A, B \in \mathbb{F}^{n \times n}$, and let k be a positive integer. Then,

$$A^k - B^k = \sum_{i=0}^{k-1} A^i (A - B) B^{k-1-i} = \sum_{i=1}^k A^{k-i} (A - B) B^{i-1}.$$

Fact 2.12.21. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and let $k \geq 1$. Then,

$$\begin{bmatrix} A & B \\ 0 & C \end{bmatrix}^k = \begin{bmatrix} A^k & \sum_{i=1}^k A^{k-i} B C^{i-1} \\ 0 & C^k \end{bmatrix}.$$

Fact 2.12.22. Let $A, B \in \mathbb{F}^{n \times n}$, and define $\mathcal{A} \triangleq \begin{bmatrix} A & A \\ A & A \end{bmatrix}$ and $\mathcal{B} \triangleq \begin{bmatrix} B & -B \\ -B & B \end{bmatrix}$. Then,

$$\mathcal{A}\mathcal{B} = \mathcal{B}\mathcal{A} = 0.$$

Fact 2.12.23. A cube root of I_2 is given by

$$\begin{bmatrix} -\frac{1}{2} & \frac{\sqrt{3}}{2} \\ \frac{-\sqrt{3}}{2} & -\frac{1}{2} \end{bmatrix}^3 = \begin{bmatrix} -1 & -1 \\ 1 & 0 \end{bmatrix}^3 = I_2.$$

Fact 2.12.24. Let n be an integer, and define

$$\begin{bmatrix} a_n \\ b_n \\ c_n \end{bmatrix} \triangleq \begin{bmatrix} 63 & 104 & -68 \\ 64 & 104 & -67 \\ 80 & 131 & -85 \end{bmatrix}^n \begin{bmatrix} 1 \\ 2 \\ 2 \end{bmatrix}.$$

Then,

$$\begin{aligned} \sum_{n=0}^{\infty} a_n &= \frac{1 + 53x + 9x^2}{1 - 82x - 82x^2 + x^3}, \\ \sum_{n=0}^{\infty} b_n &= \frac{2 - 26x - 12x^2}{1 - 82x - 82x^2 + x^3}, \\ \sum_{n=0}^{\infty} c_n &= \frac{2 + 8x - 10x^2}{1 - 82x - 82x^2 + x^3}, \end{aligned}$$

and

$$a_n^3 + b_n^3 = c_n^3 + (-1)^n.$$

(Remark: This result is an identity of Ramanujan. See [632].) (Remark: The last identity holds for all integers, not necessarily positive.)

2.13 Facts on the Determinant

Fact 2.13.1. $\det \hat{I}_n = (-1)^{\lfloor n/2 \rfloor} = (-1)^{n(n-1)/2}.$

Fact 2.13.2. $\det(I_n + \alpha 1_{n \times n}) = 1 + \alpha n$.

Fact 2.13.3. Let $A \in \mathbb{F}^{n \times m}$, let $B \in \mathbb{F}^{m \times n}$, and assume that $m < n$. Then, $\det AB = 0$.

Fact 2.13.4. Let $A \in \mathbb{F}^{n \times m}$, let $B \in \mathbb{F}^{m \times n}$, and assume that $n \leq m$. Then,

$$\det AB = \sum_{1 \leq i_1 < \dots < i_n \leq m} \det A_{(\{1, \dots, n\}, \{i_1, \dots, i_n\})} \det B_{(\{i_1, \dots, i_n\}, \{1, \dots, n\})}$$

(Proof: See [447, p. 102].) (Remark: $\det AB$ is equal to the sum of all $\binom{m}{n}$ products of pairs of subdeterminants of A and B formed by choosing n columns of A and the corresponding n rows of B .) (Remark: This identity is a special case of the Binet-Cauchy formula given by Fact 7.5.17. The special case $n = m$ is given by Proposition 2.7.1.) (Remark: Determinantal and minor identities are given in [270, 880].) (Remark: See Fact 2.14.8.)

Fact 2.13.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, let $S_1, S_2 \subseteq \{1, \dots, n\}$, and assume that $\text{card}(S_1) = \text{card}(S_2)$. Then,

$$|\det(A^{-1})_{(S_1, S_2)}| = \frac{|\det A_{(S_1^c, S_2^c)}|}{|\det A|}.$$

(Proof: See [1365, p. 38].) (Remark: When $\text{card}(S_1) = \text{card}(S_2) = 1$, this result yields the absolute value of (2.7.24).) (Remark: See Fact 2.11.21.)

Fact 2.13.6. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $b \in \mathbb{F}^n$. Then, the solution $x \in \mathbb{F}^n$ of $Ax = b$ is given by

$$x = \begin{bmatrix} \frac{\det(A \overset{1}{\leftarrow} b)}{\det A} \\ \vdots \\ \frac{\det(A \overset{n}{\leftarrow} b)}{\det A} \end{bmatrix}.$$

(Proof: Note that $A(I \overset{i}{\leftarrow} x) = A \overset{i}{\leftarrow} b$. Since $\det(I \overset{i}{\leftarrow} x) = x_{(i)}$, it follows that $(\det A)x_{(i)} = \det(A \overset{i}{\leftarrow} b)$.) (Remark: This identity is *Cramer's rule*. See Fact 2.13.7 for extensions to nonsquare A .)

Fact 2.13.7. Let $A \in \mathbb{F}^{n \times m}$ be right invertible, and let $b \in \mathbb{F}^n$. Then, a solution $x \in \mathbb{F}^m$ of $Ax = b$ is given by

$$x_{(i)} = \frac{\det[(A \overset{i}{\leftarrow} b)A^*] - \det[(A \overset{i}{\leftarrow} 0)A^*]}{\det(AA^*)},$$

for all $i = 1, \dots, m$. (Proof: See [862].) (Remark: This result is a generalization of Cramer's rule. See Fact 2.13.6. Extensions to generalized inverses are given in [178, 755, 855] and [1396, Chapter 3].)

Fact 2.13.8. Let $A \in \mathbb{F}^{n \times n}$, and assume that either $A_{(i,j)} = 0$ for all i, j such that $i + j < n + 1$ or $A_{(i,j)} = 0$ for all i, j such that $i + j > n + 1$. Then,

$$\det A = (-1)^{\lfloor n/2 \rfloor} \prod_{i=1}^n A_{(i, n+1-i)}.$$

(Remark: A is *lower reverse triangular*.)

Fact 2.13.9. Define $A \in \mathbb{R}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \end{bmatrix}.$$

Then,

$$\det A = (-1)^{n+1}.$$

Fact 2.13.10. Let $a_1, \dots, a_n \in \mathbb{F}$. Then,

$$\det \begin{bmatrix} 1+a_1 & a_2 & \cdots & a_n \\ a_1 & 1+a_2 & \cdots & a_n \\ \vdots & \vdots & \ddots & \vdots \\ a_1 & a_2 & \cdots & 1+a_n \end{bmatrix} = 1 + \sum_{i=1}^n a_i.$$

Fact 2.13.11. Let $a_1, \dots, a_n \in \mathbb{F}$ be nonzero. Then,

$$\det \begin{bmatrix} \frac{1+a_1}{a_1} & 1 & \cdots & 1 \\ 1 & \frac{1+a_2}{a_2} & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & \frac{1+a_n}{a_n} \end{bmatrix} = \frac{1 + \sum_{i=1}^n a_i}{\prod_{i=1}^n a_i}.$$

Fact 2.13.12. Let $a, b, c_1, \dots, c_n \in \mathbb{F}$, define $A \in \mathbb{F}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} c_1 & a & a & \cdots & a \\ b & c_2 & a & \cdots & a \\ b & b & c_3 & \ddots & a \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ b & b & b & \cdots & c_n \end{bmatrix},$$

and let $p(x) = (c_1 - x)(c_2 - x) \cdots (c_n - x)$ and $p_i(x) = p(x)/(c_i - x)$ for all $i = 1, \dots, n$.

Then,

$$\det A = \begin{cases} \frac{bp(a) - ap(b)}{b - a}, & b \neq a, \\ a \sum_{i=1}^{n-1} p_i(a) + c_n p_n(a), & b = a. \end{cases}$$

(Proof: See [1487, p. 10].)

Fact 2.13.13. Let $a, b \in \mathbb{F}$, and define $A, B \in \mathbb{F}^{n \times n}$ by

$$A \triangleq (a - b)I_n + b1_{n \times n} = \begin{bmatrix} a & b & b & \cdots & b \\ b & a & b & \cdots & b \\ b & b & a & \ddots & b \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ b & b & b & \cdots & a \end{bmatrix}$$

and

$$B \triangleq aI_n + b1_{n \times n} = \begin{bmatrix} a + b & b & b & \cdots & b \\ b & a + b & b & \cdots & b \\ b & b & a + b & \ddots & b \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ b & b & b & \cdots & a + b \end{bmatrix}.$$

Then,

$$\det A = (a - b)^{n-1}[a + b(n - 1)]$$

and, if $\det A \neq 0$,

$$A^{-1} = \frac{1}{a - b}I_n + \frac{b}{(b - a)[a + b(n - 1)]}1_{n \times n}.$$

Furthermore,

$$\det B = a^{n-1}(a + nb)$$

and, if $\det B \neq 0$,

$$B^{-1} = \frac{1}{a} \left(I_n - \frac{b}{a + nb} 1_{n \times n} \right).$$

(Remark: See Fact 2.14.26, Fact 4.10.15, and Fact 8.9.34.) (Remark: The matrix $aI_n + b1_{n \times n}$ arises in combinatorics. See [267, 269].)

Fact 2.13.14. Let $A \in \mathbb{F}^{n \times n}$, and define $\gamma \triangleq \max_{i,j=1,\dots,n} |A_{(i,j)}|$. Then,

$$|\det A| \leq \gamma^n n^{n/2}.$$

(Proof: The result is a consequence of the arithmetic-mean–geometric-mean inequality Fact 1.15.14 and Schur's inequality Fact 8.17.5. See [447, p. 200].) (Remark: See Fact 8.13.34.)

Fact 2.13.15. Let $A \in \mathbb{R}^{n \times n}$, and, for $i = 1, \dots, n$, let α_i denote the sum of the positive components in $\text{row}_i(A)$ and let β_i denote the sum of the positive

components in $\text{row}_i(-A)$. Then,

$$|\det A| \leq \prod_{i=1}^n \max\{\alpha_i, \beta_i\} - \prod_{i=1}^n \min\{\alpha_i, \beta_i\}.$$

(Proof: See [767].) (Remark: This result is an extension of a result due to Schinzel.)

Fact 2.13.16. For $i = 1, \dots, 4$, let $A_i, B_i \in \mathbb{F}^{2 \times 2}$, where $\det A_i = \det B_i = 1$. Furthermore, define $\mathcal{A}, \mathcal{B}, \mathcal{C}, \mathcal{D} \in \mathbb{F}^{4 \times 4}$, where, for $i, j = 1, \dots, 4$,

$$\begin{aligned}\mathcal{A}_{(i,j)} &= \text{tr } A_i A_j, \\ \mathcal{B}_{(i,j)} &= \text{tr } B_i B_j, \\ \mathcal{C}_{(i,j)} &= \text{tr } A_i B_j, \\ \mathcal{D}_{(i,j)} &= \text{tr } A_i B_j^{-1}.\end{aligned}$$

Then,

$$\det \mathcal{C} + \det \mathcal{D} = 0$$

and

$$(\det \mathcal{A})(\det \mathcal{B}) = (\det \mathcal{C})^2.$$

(Remark: These identities are due to Magnus. See [735].)

Fact 2.13.17. Let $\mathcal{J} \subseteq \mathbb{R}$ be a finite or infinite interval, and let $f: \mathcal{J} \mapsto \mathbb{R}$. Then, the following statements are equivalent:

- i) f is convex.
- ii) For all distinct $x, y, z \in \mathcal{J}$,

$$\frac{\det \begin{bmatrix} 1 & x & f(x) \\ 1 & y & f(y) \\ 1 & z & f(z) \end{bmatrix}}{\det \begin{bmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{bmatrix}} \geq 0.$$

- iii) For all $x, y, z \in \mathcal{J}$ such that $x < y < z$,

$$\det \begin{bmatrix} 1 & x & f(x) \\ 1 & y & f(y) \\ 1 & z & f(z) \end{bmatrix} \geq 0.$$

(Proof: See [1039, p. 21].)

2.14 Facts on the Determinant of Partitioned Matrices

Fact 2.14.1. Let $A \in \mathbb{F}^{n \times n}$, let A_0 be the $k \times k$ leading principal submatrix of A , and let $B \in \mathbb{F}^{(n-k) \times (n-k)}$, where, for all $i, j = 1, \dots, n-k$, $B_{(i,j)}$ is the determinant of the submatrix of A comprised of rows $1, \dots, k$ and $k+i$ and columns $1, \dots, k$ and $k+j$. Then,

$$\det B = (\det A_0)^{n-k-1} \det A.$$

If, in addition, A_0 is nonsingular, then

$$\det A = \frac{\det B}{(\det A_0)^{n-k-1}}.$$

(Remark: If $k = n - 1$, then $B = \det A$.) (Remark: This result is *Sylvester's identity*.)

Fact 2.14.2. Let $A \in \mathbb{F}^{n \times n}$, $x, y \in \mathbb{F}^n$, and $a \in \mathbb{F}$. Then,

$$\det \begin{bmatrix} A & x \\ y^T & a \end{bmatrix} = a(\det A) - y^T A^A x.$$

Hence,

$$\det \begin{bmatrix} A & x \\ y^T & a \end{bmatrix} = \begin{cases} (\det A)(a - y^T A^{-1}x), & \det A \neq 0, \\ a \det(A - a^{-1}xy^T), & a \neq 0, \\ -y^T A^A x, & a = 0 \text{ or } \det A = 0. \end{cases}$$

In particular,

$$\det \begin{bmatrix} A & Ax \\ y^T A & y^T A x \end{bmatrix} = 0.$$

Finally,

$$\det(A + xy^T) = \det A + y^T A^A x = -\det \begin{bmatrix} A & x \\ y^T & -1 \end{bmatrix}.$$

(Remark: See Fact 2.16.2, Fact 2.14.3, and Fact 2.16.4.)

Fact 2.14.3. Let $A \in \mathbb{F}^{n \times n}$, $b \in \mathbb{F}^n$, and $a \in \mathbb{F}$. Then,

$$\det \begin{bmatrix} A & b \\ b^* & a \end{bmatrix} = a(\det A) - b^* A^A b.$$

In particular,

$$\det \begin{bmatrix} A & b \\ b^* & a \end{bmatrix} = \begin{cases} (\det A)(a - b^* A^{-1}b), & \det A \neq 0, \\ a \det(A - a^{-1}bb^*), & a \neq 0, \\ -b^* A^A b, & a = 0. \end{cases}$$

(Remark: This identity is a specialization of Fact 2.14.2 with $x = b$ and $y = \bar{b}$.)

(Remark: See Fact 8.15.4.)

Fact 2.14.4. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{rank} \begin{bmatrix} A & A \\ A & A \end{bmatrix} = \text{rank} \begin{bmatrix} A & -A \\ -A & A \end{bmatrix} = \text{rank } A,$$

$$\text{rank} \begin{bmatrix} A & A \\ -A & A \end{bmatrix} = 2\text{rank } A,$$

$$\det \begin{bmatrix} A & A \\ A & A \end{bmatrix} = \det \begin{bmatrix} A & -A \\ -A & A \end{bmatrix} = 0,$$

$$\det \begin{bmatrix} A & A \\ -A & A \end{bmatrix} = 2^n (\det A)^2.$$

(Remark: See Fact 2.14.5.)

Fact 2.14.5. Let $a, b, c, d \in \mathbb{F}$, let $A \in \mathbb{F}^{n \times n}$, and define $\mathcal{A} \triangleq \begin{bmatrix} aA & bA \\ cA & dA \end{bmatrix}$. Then,

$$\text{rank } \mathcal{A} = \left(\text{rank} \begin{bmatrix} a & b \\ c & d \end{bmatrix} \right) \text{rank } A$$

and

$$\det \mathcal{A} = (ad - bc)^n (\det A)^2.$$

(Remark: See Fact 2.14.4.) (Proof: See Proposition 7.1.11 and Fact 7.4.23.)

Fact 2.14.6. $\det \begin{bmatrix} 0 & I_n \\ I_m & 0 \end{bmatrix} = (-1)^{nm}$.

Fact 2.14.7. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\det \begin{bmatrix} I_n & I_n - AB \\ B & 0 \end{bmatrix} = \det \begin{bmatrix} 0 & I_n - AB \\ B & 0 \end{bmatrix} = \det(BAB - B).$$

(Remark: See Fact 2.11.18 and Fact 2.14.6.)

Fact 2.14.8. Let $A \in \mathbb{F}^{n \times m}$, let $B \in \mathbb{F}^{m \times n}$, and assume that $n \leq m$. Then,

$$\det AB = (-1)^{(n+1)m} \det \begin{bmatrix} A & 0_{n \times n} \\ -I_m & B \end{bmatrix}.$$

(Proof: See [447].) (Remark: See Fact 2.13.4.)

Fact 2.14.9. Let A, B, C, D be conformable matrices with entries in $\mathbb{F}$. Then,

$$\begin{bmatrix} A & AB \\ C & D \end{bmatrix} = \begin{bmatrix} I & 0 \\ C & I \end{bmatrix} \begin{bmatrix} A & 0 \\ C - CA & D - CB \end{bmatrix} \begin{bmatrix} I & B \\ 0 & I \end{bmatrix},$$

$$\det \begin{bmatrix} A & AB \\ C & D \end{bmatrix} = (\det A) \det(D - CB),$$

$$\begin{bmatrix} A & B \\ CA & D \end{bmatrix} = \begin{bmatrix} I & 0 \\ C & I \end{bmatrix} \begin{bmatrix} A & B - AB \\ 0 & D - CB \end{bmatrix} \begin{bmatrix} I & B \\ 0 & I \end{bmatrix},$$

$$\det \begin{bmatrix} A & B \\ CA & D \end{bmatrix} = (\det A) \det(D - CB),$$

$$\begin{bmatrix} A & BD \\ C & D \end{bmatrix} = \begin{bmatrix} I & B \\ 0 & I \end{bmatrix} \begin{bmatrix} A - BC & 0 \\ C - DC & D \end{bmatrix} \begin{bmatrix} I & 0 \\ C & I \end{bmatrix},$$

$$\det \begin{bmatrix} A & BD \\ C & D \end{bmatrix} = \det(A - BC) \det D,$$

$$\begin{bmatrix} A & B \\ DC & D \end{bmatrix} = \begin{bmatrix} I & B \\ 0 & I \end{bmatrix} \begin{bmatrix} A - BC & B - BD \\ 0 & D \end{bmatrix} \begin{bmatrix} I & 0 \\ C & I \end{bmatrix},$$

$$\det \begin{bmatrix} A & B \\ DC & D \end{bmatrix} = \det(A - BC) \det D.$$

(Remark: See Fact 6.5.25.)

Fact 2.14.10. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\det \begin{bmatrix} A & AB \\ BA & B \end{bmatrix} = (\det A) \det(B - BAB) = (\det B) \det(A - ABA).$$

(Proof: See Fact 2.11.19 and Fact 2.14.7.)

Fact 2.14.11. Let $A_1, A_2, B_1, B_2 \in \mathbb{F}^{n \times m}$, and define $\mathcal{A} \triangleq \begin{bmatrix} A_1 & A_2 \\ A_2 & A_1 \end{bmatrix}$ and $\mathcal{B} \triangleq \begin{bmatrix} B_1 & B_2 \\ B_2 & B_1 \end{bmatrix}$. Then,

$$\text{rank} \begin{bmatrix} \mathcal{A} & \mathcal{B} \\ \mathcal{B} & \mathcal{A} \end{bmatrix} = \sum_{i=1}^4 \text{rank } C_i,$$

where $C_1 \triangleq A_1 + A_2 + B_1 + B_2$, $C_2 \triangleq A_1 + A_2 - B_1 - B_2$, $C_3 \triangleq A_1 - A_2 + B_1 - B_2$, and $C_4 \triangleq A_1 - A_2 - B_1 + B_2$. If, in addition, $n = m$, then

$$\det \begin{bmatrix} \mathcal{A} & \mathcal{B} \\ \mathcal{B} & \mathcal{A} \end{bmatrix} = \prod_{i=1}^4 \det C_i.$$

(Proof: See [1305].) (Remark: See Fact 3.22.8.)

Fact 2.14.12. Let $A, B, C, D \in \mathbb{F}^{n \times n}$, and assume that $\text{rank} \begin{bmatrix} A & B \\ C & D \end{bmatrix} = n$. Then,

$$\det \begin{bmatrix} \det A & \det B \\ \det C & \det D \end{bmatrix} = 0.$$

Fact 2.14.13. Let $A, B, C, D \in \mathbb{F}^{n \times n}$. Then,

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{cases} \det(DA - CB), & AB = BA, \\ \det(AD - CB), & AC = CA, \\ \det(AD - BC), & DC = CD, \\ \det(DA - BC), & DB = BD. \end{cases}$$

(Remark: These identities are *Schur's formulas*. See [146, p. 11].) (Proof: If A is nonsingular, then

$$\begin{aligned} \det \begin{bmatrix} A & B \\ C & D \end{bmatrix} &= (\det A) \det(D - CA^{-1}B) = \det(DA - CA^{-1}BA) \\ &= \det(DA - CB). \end{aligned}$$

Alternatively, note the identity

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} A & 0 \\ C & DA - CB \end{bmatrix} \begin{bmatrix} I & BA^{-1} \\ 0 & A^{-1} \end{bmatrix}.$$

If A is singular, then replace A by $A + \varepsilon I$ and use continuity.) (Problem: Find a direct proof for the case in which A is singular.)

Fact 2.14.14. Let $A, B, C, D \in \mathbb{F}^{n \times n}$. Then,

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{cases} \det(AD^T - B^TC^T), & AB = BA^T, \\ \det(AD^T - BC), & DC = CD^T, \\ \det(A^TD - CB), & A^TC = CA, \\ \det(A^TD - C^TB^T), & D^TB = BD. \end{cases}$$

(Proof: Define the nonsingular matrix $A_\varepsilon \triangleq A + \varepsilon I$, which satisfies $A_\varepsilon B = BA_\varepsilon^T$. Then,

$$\begin{aligned} \det \begin{bmatrix} A_\varepsilon & B \\ C & D \end{bmatrix} &= (\det A_\varepsilon) \det(D - CA_\varepsilon^{-1}B) \\ &= \det(DA_\varepsilon^T - CA_\varepsilon^{-1}BA_\varepsilon^T) = \det(DA_\varepsilon^T - CB). \end{aligned}$$

Fact 2.14.15. Let $A, B, C, D \in \mathbb{F}^{n \times n}$. Then,

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{cases} (-1)^{\text{rank } C} \det(A^TD + C^TB), & A^TC = -C^TA, \\ (-1)^{n+\text{rank } A} \det(A^TD + C^TB), & A^TC = -C^TA, \\ (-1)^{\text{rank } B} \det(A^TD + C^TB), & B^TD = -D^TB, \\ (-1)^{n+\text{rank } D} \det(A^TD + C^TB), & B^TD = -D^TB, \\ (-1)^{\text{rank } B} \det(AD^T + BC^T), & AB^T = -BA^T, \\ (-1)^{n+\text{rank } A} \det(AD^T + BC^T), & AB^T = -BA^T, \\ (-1)^{\text{rank } C} \det(AD^T + BC^T), & CD^T = -DC^T, \\ (-1)^{n+\text{rank } D} \det(AD^T + BC^T), & CD^T = -DC^T. \end{cases}$$

(Proof: See [960, 1405].) (Remark: This result is due to Callan. See [1405].)
(Remark: If $A^TC = -C^TA$ and $\text{rank } A + \text{rank } C + n$ is odd, then $\begin{bmatrix} A & B \\ C & D \end{bmatrix}$ is singular.)

Fact 2.14.16. Let $A, B, C, D \in \mathbb{F}^{n \times n}$. Then,

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{cases} \det(AD^T - BC^T), & AB^T = BA^T, \\ \det(AD^T - BC^T), & DC^T = CD^T, \\ \det(A^TD - C^TB), & A^TC = C^TA, \\ \det(A^TD - C^TB), & D^TB = B^TD. \end{cases}$$

(Proof: See [960].)

Fact 2.14.17. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times l}$, and assume that $n + k = m + l$. If $AC^T + BD^T = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = \det(AA^T + BB^T) \det(CC^T + DD^T).$$

Alternatively, if $A^TB + C^TD = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = \det(A^TA + C^TC) \det(B^TB + D^TD).$$

(Proof: Form $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix}^T$ and $\begin{bmatrix} A & B \\ C & D \end{bmatrix}^T \begin{bmatrix} A & B \\ C & D \end{bmatrix}$.)

Fact 2.14.18. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times m}$, and assume that $n + k = 2m$. If $AD^T + BC^T = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = (-1)^m \det(AB^T + BA^T) \det(CD^T + DC^T).$$

Alternatively, if $AB^T + BA^T = 0$ or $CD^T + DC^T = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = (-1)^{m^2+nk} \det(AD^T + BC^T)^2.$$

(Proof: Form $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} B^T & D^T \\ A^T & C^T \end{bmatrix}$ and $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} D^T & B^T \\ C^T & A^T \end{bmatrix}$. See [1405].)

Fact 2.14.19. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{n \times m}$, and $D \in \mathbb{F}^{n \times l}$, and assume that $m + l = 2n$. If $A^TD + C^TB = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = (-1)^n \det(C^TA + A^TC) \det(D^TB + B^TD).$$

Alternatively, if $B^TD + D^TB = 0$ or $A^TC + C^TA = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = (-1)^{n^2+ml} \det(A^TD + C^TB)^2.$$

(Proof: Form $\begin{bmatrix} C^T & A^T \\ D^T & B^T \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix}$ and $\begin{bmatrix} D^T & B^T \\ C^T & A^T \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix}$.)

Fact 2.14.20. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times k}$, $C \in \mathbb{F}^{k \times n}$, and $D \in \mathbb{F}^{k \times k}$. If $AB + BD = 0$ or $CA + DC = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = \det(A^2 + BC) \det(CB + D^2).$$

Alternatively, if $A^2 + BC = 0$ or $CB + D^2 = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = (-1)^{nk} \det(AB + BD) \det(CA + DC).$$

(Proof: Form $\begin{bmatrix} A & B \\ C & D \end{bmatrix}^2$ and $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} B & A \\ D & C \end{bmatrix}$.)

Fact 2.14.21. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times n}$, $C \in \mathbb{F}^{m \times m}$, and $D \in \mathbb{F}^{m \times n}$. If $AD + B^2 = 0$ or $C^2 + DA = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = (-1)^{nm} \det(AC + BA) \det(CD + DB).$$

Alternatively, if $AC + BA = 0$ or $CD + DB = 0$, then

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix}^2 = \det(AD + B^2)\det(C^2 + DA).$$

(Proof: Form $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} C & D \\ A & B \end{bmatrix}$ and $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} D & C \\ B & A \end{bmatrix}$.)

Fact 2.14.22. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times l}$, and assume that $n + k = m + l$. If $AC^* + BD^* = 0$, then

$$\left| \det \begin{bmatrix} A & B \\ C & D \end{bmatrix} \right|^2 = \det(AA^* + BB^*)\det(CC^* + DD^*).$$

Alternatively, if $A^*B + C^*D = 0$, then

$$\left| \det \begin{bmatrix} A & B \\ C & D \end{bmatrix} \right|^2 = \det(A^*A + C^*C)\det(B^*B + D^*D).$$

(Proof: Form $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix}^*$ and $\begin{bmatrix} A & B \\ C & D \end{bmatrix}^* \begin{bmatrix} A & B \\ C & D \end{bmatrix}$.) (Remark: See Fact 8.13.27.)

Fact 2.14.23. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times m}$, and assume that $n + k = 2m$. If $AD^* + BC^* = 0$, then

$$\left| \det \begin{bmatrix} A & B \\ C & D \end{bmatrix} \right|^2 = (-1)^m \det(AB^* + BA^*)\det(CD^* + DC^*).$$

Alternatively, if $AB^* + BA^* = 0$ or $CD^* + DC^* = 0$, then

$$\left| \det \begin{bmatrix} A & B \\ C & D \end{bmatrix} \right|^2 = (-1)^{m^2+nk} |\det(AD^* + BC^*)|^2.$$

(Proof: Form $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} B^* & D^* \\ A^* & C^* \end{bmatrix}$ and $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} D^* & B^* \\ C^* & A^* \end{bmatrix}$.) (Remark: If $m^2 + nk$ is odd, then $\begin{bmatrix} A & B \\ C & D \end{bmatrix}$ is singular.)

Fact 2.14.24. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{n \times m}$, and $D \in \mathbb{F}^{n \times l}$, and assume that $m + l = 2n$. If $A^*D + C^*B = 0$, then

$$\left| \det \begin{bmatrix} A & B \\ C & D \end{bmatrix} \right|^2 = (-1)^m \det(C^*A + A^*C)\det(D^*B + B^*D).$$

Alternatively, if $D^*B + B^*D = 0$ or $C^*A + A^*C = 0$, then

$$\left| \det \begin{bmatrix} A & B \\ C & D \end{bmatrix} \right|^2 = (-1)^{n^2+ml} |\det(A^*D + C^*B)|^2.$$

(Proof: Form $\begin{bmatrix} C^* & A^* \\ D^* & B^* \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix}$ and $\begin{bmatrix} D^* & B^* \\ C^* & A^* \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix}$.) (Remark: If $n^2 + ml$ is odd, then $\begin{bmatrix} A & B \\ C & D \end{bmatrix}$ is singular.)

Fact 2.14.25. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\det \begin{bmatrix} A^*A & A^*B \\ B^*A & B^*B \end{bmatrix} = \begin{cases} \det(A^*A)\det[B^*B - B^*A(A^*A)^{-1}A^*B], & \text{rank } A = m, \\ \det(B^*B)\det[A^*A - A^*B(B^*B)^{-1}B^*A], & \text{rank } B = l, \\ 0, & n < m + l. \end{cases}$$

If, in addition, $m + l = n$, then

$$\det \begin{bmatrix} A^*A & A^*B \\ B^*A & B^*B \end{bmatrix} = \det(AA^* + BB^*).$$

(Remark: See Fact 6.5.27.)

Fact 2.14.26. Let $A, B \in \mathbb{F}^{n \times n}$, and define $\mathcal{A} \in \mathbb{F}^{kn \times kn}$ by

$$\mathcal{A} \triangleq \begin{bmatrix} A & B & B & \cdots & B \\ B & A & B & \cdots & B \\ B & B & A & \ddots & B \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ B & B & B & \cdots & A \end{bmatrix}.$$

Then,

$$\det \mathcal{A} = [\det(A - B)]^{k-1} \det[A + (k-1)B].$$

If $k = 2$, then

$$\det \begin{bmatrix} A & B \\ B & A \end{bmatrix} = \det[(A + B)(A - B)] = \det(A^2 - B^2 - [A, B]).$$

(Proof: See [573].) (Remark: For $k = 2$, the result follows from Fact 4.10.25.)
(Remark: See Fact 2.13.13.)

Fact 2.14.27. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, and $D \in \mathbb{F}^{m \times m}$, and define $M \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$. Furthermore, let $\begin{bmatrix} A' & B' \\ C' & D' \end{bmatrix} \triangleq M^A$, where $A' \in \mathbb{F}^{n \times n}$ and $D' \in \mathbb{F}^{m \times m}$. Then,

$$\det D' = (\det M)^{m-1} \det A$$

and

$$\det A' = (\det M)^{n-1} \det D.$$

(Proof: See [1184, p. 297].) (Remark: See Fact 2.14.28.)

Fact 2.14.28. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, and $D \in \mathbb{F}^{m \times m}$, define $M \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, and assume that M is nonsingular. Furthermore, let $\begin{bmatrix} A' & B' \\ C' & D' \end{bmatrix} \triangleq M^{-1}$, where $A' \in \mathbb{F}^{n \times n}$ and $D' \in \mathbb{F}^{m \times m}$. Then,

$$\det D' = \frac{\det A}{\det M}$$

and

$$\det A' = \frac{\det D}{\det M}.$$

Consequently, A is nonsingular if and only if D' is nonsingular, and D is nonsingular if and only if A' is nonsingular. (Proof: Use $M \begin{bmatrix} I & B' \\ 0 & D' \end{bmatrix} = \begin{bmatrix} A & 0 \\ C & I \end{bmatrix}$. See [1188].) (Remark: This identity is a special case of *Jacobi's identity*. See [709, p. 21].) (Remark: See Fact 2.14.27 and Fact 3.11.24.)

2.15 Facts on Left and Right Inverses

Fact 2.15.1. Let $A \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

- i) If $A^L \in \mathbb{F}^{m \times n}$ is a left inverse of A , then $\overline{A^L} \in \mathbb{F}^{m \times n}$ is a left inverse of $\overline{A}$.
- ii) If $A^L \in \mathbb{F}^{m \times n}$ is a left inverse of A , then $A^{LT} \in \mathbb{F}^{n \times m}$ is a right inverse of A^T .
- iii) If $A^L \in \mathbb{F}^{m \times n}$ is a left inverse of A , then $A^{L*} \in \mathbb{F}^{n \times m}$ is a right inverse of A^* .
- iv) If $A^R \in \mathbb{F}^{m \times n}$ is a right inverse of A , then $\overline{A^R} \in \mathbb{F}^{m \times n}$ is a right inverse of $\overline{A}$.
- v) If $A^R \in \mathbb{F}^{m \times n}$ is a right inverse of A , then $A^{RT} \in \mathbb{F}^{n \times m}$ is a left inverse of A^T .
- vi) If $A^R \in \mathbb{F}^{m \times n}$ is a right inverse of A , then $A^{R*} \in \mathbb{F}^{n \times m}$ is a left inverse of A^* .

Furthermore, the following statements are equivalent:

- vii) A is left invertible.
- viii) $\overline{A}$ is left invertible.
- ix) A^T is right invertible.
- x) A^* is right invertible.

Finally, the following statements are equivalent:

- xi) A is right invertible.
- xii) $\overline{A}$ is right invertible.
- xiii) A^T is left invertible.
- xiv) A^* is left invertible.

Fact 2.15.2. Let $A \in \mathbb{F}^{n \times m}$. If $\text{rank } A = m$, then $(A^*A)^{-1}A^*$ is a left inverse of A . If $\text{rank } A = n$, then $A^*(AA^*)^{-1}$ is a right inverse of A . (Remark: See Fact 3.7.25, Fact 3.7.26, and Fact 3.13.6.)

Fact 2.15.3. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = m$. Then, $A^L \in \mathbb{F}^{m \times n}$ is a left inverse of A if and only if there exists a matrix $B \in \mathbb{F}^{m \times n}$ such that BA is nonsingular and

$$A^L = (BA)^{-1}B.$$

(Proof: For necessity, let $B = A^L$.)

Fact 2.15.4. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = n$. Then, $A^R \in \mathbb{F}^{m \times n}$ is a right inverse of A if and only if there exists a matrix $B \in \mathbb{F}^{m \times n}$ such that AB is nonsingular and

$$A^R = B(AB)^{-1}.$$

(Proof: For necessity, let $B = A^R$.)

Fact 2.15.5. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, and assume that A and B are left invertible. Then, AB is left invertible. If, in addition, A^L is a left inverse of A and B^L is a left inverse of B , then $B^L A^L$ is a left inverse of AB .

Fact 2.15.6. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, and assume that A and B are right invertible. Then, AB is right invertible. If, in addition, A^R is a right inverse of A and B^R is a right inverse of B , then $B^R A^R$ is a right inverse of AB .

2.16 Facts on the Adjugate and Inverses

Fact 2.16.1. Let $x, y \in \mathbb{F}^n$. Then,

$$(I + xy^T)^A = (1 + y^T x)I - xy^T$$

and

$$\det(I + xy^T) = \det(I + yx^T) = 1 + x^T y = 1 + y^T x.$$

If, in addition, $x^T y \neq -1$, then

$$(I + xy^T)^{-1} = I - (1 + x^T y)^{-1} xy^T.$$

Fact 2.16.2. Let $A \in \mathbb{F}^{n \times n}$, $x, y \in \mathbb{F}^n$, and $a \in \mathbb{F}$. Then,

$$\begin{bmatrix} A & x \\ y^T & a \end{bmatrix} = \begin{cases} \begin{bmatrix} I & 0 \\ y^T A^{-1} & 1 \end{bmatrix} \begin{bmatrix} A & 0 \\ 0 & a - y^T A^{-1} x \end{bmatrix} \begin{bmatrix} I & A^{-1} x \\ 0 & 1 \end{bmatrix}, & \det A \neq 0, \\ \begin{bmatrix} I & a^{-1} x \\ 0 & 1 \end{bmatrix} \begin{bmatrix} A - a^{-1} x y^T & 0 \\ 0 & a \end{bmatrix} \begin{bmatrix} I & 0 \\ a^{-1} y^T & 1 \end{bmatrix}, & a \neq 0. \end{cases}$$

(Remark: See Fact 6.5.25.)

Fact 2.16.3. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $x, y \in \mathbb{F}^n$. Then,

$$\det(A + xy^T) = (1 + y^T A^{-1} x) \det A$$

and

$$(A + xy^T)^A = (1 + y^T A^{-1} x) (\det A) I - A^A xy^T.$$

Furthermore, the following statements are equivalent:

- i) $\det(A + xy^T) \neq 0$
- ii) $y^T A^{-1} x \neq -1$.
- iii) $\begin{bmatrix} A & x \\ y^T & -1 \end{bmatrix}$ is nonsingular.

In this case,

$$(A + xy^T)^{-1} = A^{-1} - (1 + y^T A^{-1} x)^{-1} A^{-1} xy^T A^{-1}.$$

(Remark: See Fact 2.16.2 and Fact 2.14.2.) (Remark: The last identity, which is a special case of the matrix inversion lemma Corollary 2.8.8, is the *Sherman-Morrison-Woodbury formula*.)

Fact 2.16.4. Let $A \in \mathbb{F}^{n \times n}$, let $x, y \in \mathbb{F}^n$, and let $a \in \mathbb{F}$. Then,

$$\begin{bmatrix} A & x \\ y^T & a \end{bmatrix}^A = \begin{bmatrix} (a+1)A^A - (A + xy^T)^A & -A^A x \\ -y^T A^A & \det A \end{bmatrix}.$$

Now, assume that $\begin{bmatrix} A & x \\ y^T & a \end{bmatrix}$ is nonsingular. Then,

$$\begin{aligned} & \begin{bmatrix} A & x \\ y^T & a \end{bmatrix}^{-1} \\ &= \begin{cases} \frac{1}{(\det A)(a - y^T A^{-1}x)} \begin{bmatrix} (a - y^T A^{-1}x)A^{-1} + A^{-1}xy^T A^{-1} & -A^{-1}x \\ -y^T A^{-1} & 1 \end{bmatrix}, & \det A \neq 0, \\ \frac{1}{a \det(A - a^{-1}xy^T)} \begin{bmatrix} (a+1)A^A - (A + xy^T)^A & -A^A x \\ -y^T A^A & \det A \end{bmatrix}, & a \neq 0, \\ \frac{1}{-y^T A^A x} \begin{bmatrix} (a+1)A^A - (A + xy^T)^A & -A^A x \\ -y^T A^A & \det A \end{bmatrix}, & a = 0. \end{cases} \end{aligned}$$

(Proof: Use Fact 2.14.2 and see [455, 686].)

Fact 2.16.5. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) $(\overline{A})^A = \overline{A^A}$.
- ii) $(A^T)^A = (A^A)^T$.
- iii) $(A^*)^A = (A^A)^*$.
- iv) If $\alpha \in \mathbb{F}$, then $(\alpha A)^A = \alpha^{n-1} A^A$.
- v) $\det A^A = (\det A)^{n-1}$.
- vi) $(A^A)^A = (\det A)^{n-2} A$.
- vii) $\det (A^A)^A = (\det A)^{(n-1)^2}$.
- viii) If A is nonsingular, then $(A^{-1})^A = (A^A)^{-1}$.

(Proof: See [686].) (Remark: With $0/0 \triangleq 1$ in vi), all of these results hold in the degenerate case $n = 1$.)

Fact 2.16.6. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\det(A + 1_{n \times n}) - \det A = 1_{1 \times n}^T A^A 1 = \sum_{i=1}^n \det(A \stackrel{i}{\leftarrow} 1_{n \times 1}).$$

(Proof: See [222].) (Remark: See Fact 2.14.2, Fact 2.16.9, and Fact 10.11.21.)

Fact 2.16.7. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is singular. Then,

$$\mathcal{R}(A) \subseteq \mathcal{N}(A^A).$$

Hence,

$$\text{rank } A \leq \text{def } A^A$$

and

$$\text{rank } A + \text{rank } A^A \leq n.$$

Furthermore, $\mathcal{R}(A) = \mathcal{N}(A^A)$ if and only if $\text{rank } A = n - 1$.

Fact 2.16.8. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) $\text{rank } A^A = n$ if and only if $\text{rank } A = n$.
- ii) $\text{rank } A^A = 1$ if and only if $\text{rank } A = n - 1$.
- iii) $A^A = 0$ if and only if $\text{rank } A \leq n - 2$.

(Proof: See [1098, p. 12].) (Remark: See Fact 4.10.7.) (Remark: Fact 6.3.6 provides an expression for A^A in the case $\text{rank } A^A = 1$.)

Fact 2.16.9. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$(A^A B)_{(i,j)} = \det \left[A \overset{i}{\leftarrow} \text{col}_j(B) \right].$$

(Remark: See Fact 10.11.21.)

Fact 2.16.10. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) $(AB)^A = B^A A^A$.
- ii) If B is nonsingular, then $(BAB^{-1})^A = B A^A B^{-1}$.
- iii) If $AB = BA$, then $A^A B = B A^A$, $AB^A = B^A A$, and $A^A B^A = B^A A^A$.

Fact 2.16.11. Let $A, B, C, D \in \mathbb{F}^{n \times n}$ and $ABCD = I$. Then, $ABCD = DABC = CDAB = BCDA$.

Fact 2.16.12. Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbb{F}^{2 \times 2}$, where $ad - bc \neq 0$. Then,

$$A^{-1} = (ad - bc)^{-1} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

Furthermore, if $A = \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} \in \mathbb{F}^{3 \times 3}$ and $\beta = a(ei - fh) - b(di - fg) + c(dh - eg) \neq 0$, then

$$A^{-1} = \beta^{-1} \begin{bmatrix} ei - fh & -(bi - ch) & bf - ce \\ -(di - fg) & ai - cg & -(af - cd) \\ dh - eg & -(ah - bg) & ae - bd \end{bmatrix}.$$

Fact 2.16.13. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $A + B$ is nonsingular. Then,

$$A(A + B)^{-1}B = B(A + B)^{-1}A = A - A(A + B)^{-1}A = B - B(A + B)^{-1}B.$$

Fact 2.16.14. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are nonsingular. Then,

$$A^{-1} + B^{-1} = A^{-1}(A + B)B^{-1}.$$

Furthermore, $A^{-1} + B^{-1}$ is nonsingular if and only if $A + B$ is nonsingular. In this case,

$$\begin{aligned} (A^{-1} + B^{-1})^{-1} &= A(A + B)^{-1}B \\ &= B(A + B)^{-1}A \\ &= A - A(A + B)^{-1}A \\ &= B - B(A + B)^{-1}B. \end{aligned}$$

Fact 2.16.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are nonsingular. Then,

$$A - B = A(B^{-1} - A^{-1})B.$$

Therefore,

$$\text{rank}(A - B) = \text{rank}(A^{-1} - B^{-1}).$$

In particular, $A - B$ is nonsingular if and only if $A^{-1} - B^{-1}$ is nonsingular. In this case,

$$(A^{-1} - B^{-1})^{-1} = A - A(A - B)^{-1}A.$$

Fact 2.16.16. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$, and assume that $I + AB$ is nonsingular. Then, $I + BA$ is nonsingular and

$$(I_n + AB)^{-1}A = A(I_m + BA)^{-1}.$$

(Remark: This result is the *push-through identity*.) Furthermore,

$$(I + AB)^{-1} = I - (I + AB)^{-1}AB.$$

Fact 2.16.17. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $I + BA$ is nonsingular. Then,

$$(I + AB)^{-1} = I - A(I + BA)^{-1}B.$$

Fact 2.16.18. Let $A \in \mathbb{F}^{n \times n}$, and assume that A and $A + I$ are nonsingular. Then,

$$(A + I)^{-1} + (A^{-1} + I)^{-1} = (A + I)^{-1} + (A + I)^{-1}A = I.$$

Fact 2.16.19. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$(I + AA^*)^{-1} = I - A(I + A^*A)^{-1}A^*.$$

Fact 2.16.20. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, let $B \in \mathbb{F}^{n \times m}$, let $C \in \mathbb{F}^{m \times n}$, and assume that $A + BC$ and $I + CA^{-1}B$ are nonsingular. Then,

$$(A + BC)^{-1}B = A^{-1}B(I + CA^{-1}B)^{-1}.$$

In particular, if $A + BB^*$ and $I + B^*A^{-1}B$ are nonsingular, then

$$(A + BB^*)^{-1}B = A^{-1}B(I + B^*A^{-1}B)^{-1}.$$

Fact 2.16.21. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{l \times n}$, and $D \in \mathbb{F}^{m \times l}$, and assume that A and $A + BDC$ are nonsingular. Then,

$$\begin{aligned} (A + BDC)^{-1} &= A^{-1} - (I + A^{-1}BDC)^{-1}A^{-1}BDCA^{-1} \\ &= A^{-1} - A^{-1}(I + BDCA^{-1})^{-1}BDCA^{-1} \\ &= A^{-1} - A^{-1}B(I + DCA^{-1}B)^{-1}DCA^{-1} \\ &= A^{-1} - A^{-1}BD(I + CA^{-1}BD)^{-1}CA^{-1} \\ &= A^{-1} - A^{-1}BDC(I + A^{-1}BDC)^{-1}A^{-1} \\ &= A^{-1} - A^{-1}BDCA^{-1}(I + BDCA^{-1})^{-1}. \end{aligned}$$

(Proof: See [666].) (Remark: The third identity generalizes the matrix inversion lemma Corollary 2.8.8 in the form

$$(A + BDC)^{-1} = A^{-1} - A^{-1}B(D^{-1} + CA^{-1}B)^{-1}CA^{-1}$$

since D need not be square or invertible.)

Fact 2.16.22. Let $A \in \mathbb{F}^{n \times m}$, let $C, D \in \mathbb{F}^{n \times m}$, and assume that $I + DB$ is nonsingular. Then,

$$I + AC - (A + B)(I + DB)^{-1}(D + C) = (I - AD)(I + BD)^{-1}(I - BC).$$

(Proof: See [1467].) (Remark: See Fact 2.16.23 and Fact 8.11.21.)

Fact 2.16.23. Let $A, B, C \in \mathbb{F}^{n \times m}$. Then,

$$I + AC^* - (A + B)(I + B^*B)^{-1}(B + C)^* = (I - AB^*)(I + BB^*)^{-1}(I - BC^*).$$

(Proof: Set $D = B^*$ and replace C by C^* in Fact 2.16.22.)

Fact 2.16.24. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that B is nonsingular. Then,

$$A = B[I + B^{-1}(A - B)].$$

Fact 2.16.25. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and $A + B$ are nonsingular. Then, for all $k \in \mathbb{N}$,

$$\begin{aligned} (A + B)^{-1} &= \sum_{i=0}^k A^{-1}(-BA^{-1})^i + (-A^{-1}B)^{k+1}(A + B)^{-1} \\ &= \sum_{i=0}^k A^{-1}(-BA^{-1})^i + A^{-1}(-BA^{-1})^{k+1}(I + BA^{-1})^{-1}. \end{aligned}$$

Fact 2.16.26. Let $A \in \mathbb{F}^{n \times n}$, assume that A is either upper triangular or lower triangular, let D denote the diagonal part of A , and assume that D is nonsingular. Then,

$$A^{-1} = \sum_{i=0}^n (I - D^{-1}A)^i D^{-1}.$$

(Remark: Using the Schur product notation, $D = I \circ A$.)

Fact 2.16.27. Let $A, B \in \mathbb{F}^{n \times n}$ and $\alpha \in \mathbb{F}$, and assume that A , B , $\alpha A^{-1} + (1 - \alpha)B^{-1}$, and $\alpha B + (1 - \alpha)A$ are nonsingular. Then,

$$\begin{aligned} \alpha A + (1 - \alpha)B - [\alpha A^{-1} + (1 - \alpha)B^{-1}]^{-1} \\ = \alpha(1 - \alpha)(A - B)[\alpha B + (1 - \alpha)A]^{-1}(A - B). \end{aligned}$$

(Remark: This identity is relevant to *iv*) of Proposition 8.6.17.)

Fact 2.16.28. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and define $A_0 \triangleq I_n$. Furthermore, for all $k = 1, \dots, n$, let

$$\alpha_k = \frac{1}{k} \operatorname{tr} AA_{k-1},$$

and, for all $k = 1, \dots, n - 1$, let

$$A_k = AA_{k-1} - \alpha_k I.$$

Then,

$$A^{-1} = \frac{1}{\alpha_n} A_{n-1}.$$

(Remark: This result is due to Frame. See [170, p. 99].)

Fact 2.16.29. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and define $\{B_i\}_{i=1}^{\infty}$ by

$$B_{i+1} \triangleq 2B_i - B_i A B_i,$$

where $B_0 \in \mathbb{F}^{n \times n}$ satisfies $\operatorname{sprad}(I - B_0 A) < 1$. Then,

$$B_i \rightarrow A^{-1}$$

as $i \rightarrow \infty$. (Proof: See [144, p. 167].) (Remark: This sequence is given by a Newton-Raphson algorithm.) (Remark: See Fact 6.3.35 for the case in which A is singular or nonsquare.)

Fact 2.16.30. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then, $A + A^{-*}$ is nonsingular. (Proof: Note that $AA^* + I$ is positive definite.)

2.17 Facts on the Inverse of Partitioned Matrices

Fact 2.17.1. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, and $D \in \mathbb{F}^{m \times m}$, and assume that A and D are nonsingular. Then,

$$\begin{bmatrix} A & B \\ 0 & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} & -A^{-1}BD^{-1} \\ 0 & D^{-1} \end{bmatrix}$$

and

$$\begin{bmatrix} A & 0 \\ C & D \end{bmatrix}^{-1} = \begin{bmatrix} A^{-1} & 0 \\ -D^{-1}CA^{-1} & D^{-1} \end{bmatrix}.$$

Fact 2.17.2. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{m \times n}$. Then,

$$\det \begin{bmatrix} 0 & A \\ B & C \end{bmatrix} = \det \begin{bmatrix} C & B \\ A & 0 \end{bmatrix} = (-1)^{nm} (\det A) (\det B).$$

If, in addition, A and B are nonsingular, then

$$\begin{bmatrix} 0 & A \\ B & C \end{bmatrix}^{-1} = \begin{bmatrix} -B^{-1}CA^{-1} & B^{-1} \\ A^{-1} & 0 \end{bmatrix}$$

and

$$\begin{bmatrix} C & B \\ A & 0 \end{bmatrix}^{-1} = \begin{bmatrix} 0 & A^{-1} \\ B^{-1} & -B^{-1}CA^{-1} \end{bmatrix}.$$

Fact 2.17.3. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and assume that C is nonsingular. Then,

$$\begin{bmatrix} A & B \\ B^T & C \end{bmatrix} = \begin{bmatrix} A - BC^{-1}B^T & B \\ 0 & C \end{bmatrix} \begin{bmatrix} I & 0 \\ C^{-1}B^T & I \end{bmatrix}.$$

If, in addition, $A - BC^{-1}B^T$ is nonsingular, then $\begin{bmatrix} A & B \\ B^T & C \end{bmatrix}$ is nonsingular and

$$\begin{aligned} & \begin{bmatrix} A & B \\ B^T & C \end{bmatrix}^{-1} \\ &= \begin{bmatrix} (A - BC^{-1}B^T)^{-1} & -(A - BC^{-1}B^T)^{-1}BC^{-1} \\ -C^{-1}B^T(A - BC^{-1}B^T)^{-1} & C^{-1}B^T(A - BC^{-1}B^T)^{-1}BC^{-1} + C^{-1} \end{bmatrix}. \end{aligned}$$

Fact 2.17.4. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\det \begin{bmatrix} I & A \\ B & I \end{bmatrix} = \det(I - AB) = \det(I - BA).$$

If $\det(I - BA) \neq 0$, then

$$\begin{aligned} \begin{bmatrix} I & A \\ B & I \end{bmatrix}^{-1} &= \begin{bmatrix} I + A(I - BA)^{-1}B & -A(I - BA)^{-1} \\ -(I - BA)^{-1}B & (I - BA)^{-1} \end{bmatrix} \\ &= \begin{bmatrix} (I - AB)^{-1} & -(I - AB)^{-1}A \\ -B(I - AB)^{-1} & I + B(I - AB)^{-1}A \end{bmatrix}. \end{aligned}$$

Fact 2.17.5. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\begin{bmatrix} A & B \\ B & A \end{bmatrix} = \frac{1}{2} \begin{bmatrix} I & I \\ I & -I \end{bmatrix} \begin{bmatrix} A + B & 0 \\ 0 & A - B \end{bmatrix} \begin{bmatrix} I & I \\ I & -I \end{bmatrix}.$$

Therefore,

$$\text{rank} \begin{bmatrix} A & B \\ B & A \end{bmatrix} = \text{rank}(A + B) + \text{rank}(A - B).$$

Now, assume that $n = m$. Then,

$$\det \begin{bmatrix} A & B \\ B & A \end{bmatrix} = \det[(A + B)(A - B)] = \det(A^2 - B^2 - [A, B]).$$

Hence, $\begin{bmatrix} A & B \\ B & A \end{bmatrix}$ is nonsingular if and only if $A + B$ and $A - B$ are nonsingular. In

this case,

$$\begin{bmatrix} A & B \\ B & A \end{bmatrix}^{-1} = \frac{1}{2} \begin{bmatrix} (A+B)^{-1} + (A-B)^{-1} & (A+B)^{-1} - (A-B)^{-1} \\ (A+B)^{-1} - (A-B)^{-1} & (A+B)^{-1} + (A-B)^{-1} \end{bmatrix},$$

$$(A+B)^{-1} = \frac{1}{2} \begin{bmatrix} I & I \end{bmatrix} \begin{bmatrix} A & B \\ B & A \end{bmatrix}^{-1} \begin{bmatrix} I \\ I \end{bmatrix},$$

and

$$(A-B)^{-1} = \frac{1}{2} \begin{bmatrix} I & -I \end{bmatrix} \begin{bmatrix} A & B \\ B & A \end{bmatrix}^{-1} \begin{bmatrix} I \\ -I \end{bmatrix}.$$

(Remark: See Fact 6.5.1.)

Fact 2.17.6. Let $A_1, \dots, A_k \in \mathbb{F}^{n \times n}$, and assume that the $kn \times kn$ partitioned matrix below is nonsingular. Then, $A_1 + \dots + A_k$ is nonsingular, and

$$(A_1 + \dots + A_k)^{-1} = \frac{1}{k} \begin{bmatrix} I_n & \dots & I_n \end{bmatrix} \begin{bmatrix} A_1 & A_2 & \dots & A_k \\ A_k & A_1 & \dots & A_{k-1} \\ \vdots & \vdots & \ddots & \vdots \\ A_2 & A_3 & \dots & A_1 \end{bmatrix}^{-1} \begin{bmatrix} I_m \\ \vdots \\ I_m \end{bmatrix}.$$

(Proof: See [1282].) (Remark: The partitioned matrix is *block circulant*. See Fact 6.5.2 and Fact 6.6.1.)

Fact 2.17.7. Let $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ 0_{m \times m} & C \end{bmatrix}$, where $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times n}$, and $C \in \mathbb{F}^{m \times n}$, and assume that CA is nonsingular. Furthermore, define $P \triangleq A(CA)^{-1}C$ and $P_\perp \triangleq I - P$. Then, $\mathcal{A}$ is nonsingular if and only if $P + P_\perp B P_\perp$ is nonsingular. In this case,

$$\mathcal{A}^{-1} = \begin{bmatrix} (CA)^{-1}(C - CBD) & -(CA)^{-1}CB(A - DBA)(CA)^{-1} \\ D & (A - DBA)(CA)^{-1} \end{bmatrix},$$

where $D \triangleq (P + P_\perp B P_\perp)^{-1} P_\perp$. (Proof: See [639].)

Fact 2.17.8. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times (n-m)}$, and assume that $\begin{bmatrix} A & B \end{bmatrix}$ is nonsingular and $A^*B = 0$. Then,

$$\begin{bmatrix} A & B \end{bmatrix}^{-1} = \begin{bmatrix} (A^*A)^{-1}A^* \\ (B^*B)^{-1}B^* \end{bmatrix}.$$

(Remark: See Fact 6.5.18.) (Problem: Find an expression for $\begin{bmatrix} A & B \end{bmatrix}^{-1}$ without assuming $A^*B = 0$.)

Fact 2.17.9. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, and $C \in \mathbb{F}^{m \times l}$. Then,

$$\begin{bmatrix} I_n & A & B \\ 0 & I_m & C \\ 0 & 0 & I_l \end{bmatrix}^{-1} = \begin{bmatrix} I_n & -A & AC - B \\ 0 & I_m & -C \\ 0 & 0 & I_l \end{bmatrix}.$$

Fact 2.17.10. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then, $X = A^{-1}$ is the unique matrix satisfying

$$\text{rank} \begin{bmatrix} A & I \\ I & X \end{bmatrix} = \text{rank } A.$$

(Remark: See Fact 6.3.30 and Fact 6.6.2.) (Proof: See [483].)

2.18 Facts on Commutators

Fact 2.18.1. Let $A, B \in \mathbb{F}^{2 \times 2}$. Then,

$$[A, B]^2 = \frac{1}{2}(\text{tr}[A, B]^2)I_2.$$

(Remark: See [499, 500].)

Fact 2.18.2. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\text{tr}[A, B]^3 = 3\text{tr}(A^2B^2AB - B^2A^2BA) = -3\text{tr}(AB^2A[A, B]).$$

Fact 2.18.3. Let $A, B \in \mathbb{F}^{n \times n}$, assume that $[A, B] = 0$, and let $k, l \in \mathbb{N}$. Then, $[A^k, B^l] = 0$.

Fact 2.18.4. Let $A, B, C \in \mathbb{F}^{n \times n}$. Then, the following identities hold:

- i) $[A, A] = 0$.
- ii) $[A, B] = [-A, -B] = -[B, A]$.
- iii) $[A, B + C] = [A, B] + [A, C]$.
- iv) $[\alpha A, B] = [A, \alpha B] = \alpha[A, B]$ for all $\alpha \in \mathbb{F}$.
- v) $[A, [B, C]] + [B, [C, A]] + [C, [A, B]] = 0$.
- vi) $[A, B]^T = [B^T, A^T] = -[A^T, B^T]$.
- vii) $\text{tr}[A, B] = 0$.
- viii) $\text{tr } A^k[A, B] = \text{tr } B^k[A, B] = 0$ for all $k \geq 1$.
- ix) $[[A, B], B - A] = [[B, A], A - B]$.
- x) $[A, [A, B]] = -[A, [B, A]]$.

(Remark: v) is the *Jacobi identity*.)

Fact 2.18.5. Let $A, B \in \mathbb{F}^{n \times n}$. Then, for all $X \in \mathbb{F}^{n \times n}$,

$$\text{ad}_{[A, B]} = [\text{ad}_A, \text{ad}_B],$$

that is,

$$\text{ad}_{[A, B]}(X) = \text{ad}_A[\text{ad}_B(X)] - \text{ad}_B[\text{ad}_A(X)]$$

or, equivalently,

$$[[A, B], X] = [A, [B, X]] - [B, [A, X]].$$

Fact 2.18.6. Let $A \in \mathbb{F}^{n \times n}$ and, for all $X \in \mathbb{F}^{n \times n}$, define

$$\text{ad}_A^k(X) \triangleq \begin{cases} \text{ad}_A(X), & k = 1, \\ \text{ad}_A^{k-1}[\text{ad}_A(X)], & k \geq 2. \end{cases}$$

Then, for all $X \in \mathbb{F}^{n \times n}$ and $k \geq 1$,

$$\text{ad}_A^2(X) = [A, [A, X]] - [[A, X], A]$$

and

$$\text{ad}_A^k(X) = \sum_{i=0}^k (-1)^{k-i} \binom{k}{i} A^i X A^{k-i}.$$

(Remark: The proof of Proposition 11.4.7 is based on $g(e^{t\text{ad}_A} e^{t\text{ad}_B})$, where $g(z) \triangleq (\log z)/(z-1)$. See [1162, p. 35].) (Remark: See Fact 11.14.4.) (Proof: For the last identity, see [1098, pp. 176, 207].)

Fact 2.18.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $[A, B] = A$. Then, A is singular. (Proof: If A is nonsingular, then $\text{tr } B = \text{tr } ABA^{-1} = \text{tr } B + n$.)

Fact 2.18.8. Let $A, B \in \mathbb{R}^{n \times n}$ be such that $AB = BA$. Then, there exists a matrix $C \in \mathbb{R}^{n \times n}$ such that $A^2 + B^2 = C^2$. (Proof: See [415].) (Remark: This result applies to real matrices only.)

Fact 2.18.9. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$n \leq \dim \{X \in \mathbb{F}^{n \times n}: AX = XA\}$$

and

$$\dim \{[A, X]: X \in \mathbb{F}^{n \times n}\} \leq n^2 - n.$$

(Proof: See [392, pp. 125, 142, 493, 537].) (Remark: The first set is the *centralizer* or *commutant* of A . See Fact 7.5.2.) (Remark: These quantities are the defect and rank, respectively, of the operator $f: \mathbb{F}^{n \times n} \mapsto \mathbb{F}^{n \times n}$ defined by $f(X) \triangleq AX - XA$. See Fact 7.5.2.) (Remark: See Fact 5.14.22 and Fact 5.14.24.)

Fact 2.18.10. Let $A \in \mathbb{F}^{n \times n}$. Then, there exists $\alpha \in \mathbb{F}$ such that $A = \alpha I$ if and only if, for all $X \in \mathbb{F}^{n \times n}$, $AX = XA$. (Proof: To prove sufficiency, note that $A^T \oplus -A = 0$. Hence, $\{0\} = \text{spec}(A^T \oplus -A) = \{\lambda - \mu: \lambda, \mu \in \text{spec}(A)\}$. Therefore, $\text{spec}(A) = \{\alpha\}$, and thus $A = \alpha I + N$, where N is nilpotent. Consequently, for all $X \in \mathbb{F}^{n \times n}$, $NX = XN$. Setting $X = N^*$, it follows that N is normal. Hence, $N = 0$.) (Remark: This result determines the center subgroup of $\text{GL}(n)$.)

Fact 2.18.11. Define $\mathcal{S} \subseteq \mathbb{F}^{n \times n}$ by

$$\mathcal{S} \triangleq \{[X, Y]: X, Y \in \mathbb{F}^{n \times n}\}.$$

Then, $\mathcal{S}$ is a subspace. Furthermore,

$$\mathcal{S} = \{Z \in \mathbb{F}^{n \times n}: \text{tr } Z = 0\}$$

and

$$\dim \mathcal{S} = n^2 - 1.$$

(Proof: See [392, pp. 125, 493]. Alternatively, note that $\text{tr}: \mathbb{F}^{n^2} \mapsto \mathbb{F}$ is onto, and use Corollary 2.5.5.)

Fact 2.18.12. Let $A, B, C, D \in \mathbb{F}^{n \times n}$. Then, there exist $E, F \in \mathbb{F}^{n \times n}$ such that

$$[E, F] = [A, B] + [C, D].$$

(Proof: The result follows from Fact 2.18.11.) (Problem: Construct E and F .)

2.19 Facts on Complex Matrices

Fact 2.19.1. Let $a, b \in \mathbb{R}$. Then, $\begin{bmatrix} a & b \\ -b & a \end{bmatrix}$ is a representation of the complex number $a + jb$ that preserves addition, multiplication and inversion of complex numbers. In particular, if $a^2 + b^2 \neq 0$, then

$$\begin{bmatrix} a & b \\ -b & a \end{bmatrix}^{-1} = \begin{bmatrix} \frac{a}{a^2+b^2} & \frac{-b}{a^2+b^2} \\ \frac{b}{a^2+b^2} & \frac{a}{a^2+b^2} \end{bmatrix}$$

and

$$(a + jb)^{-1} = \frac{a}{a^2 + b^2} - j \frac{b}{a^2 + b^2}.$$

(Remark: $\begin{bmatrix} a & b \\ -b & a \end{bmatrix}$ is a *rotation-dilation*. See Fact 3.22.6.)

Fact 2.19.2. Let $\nu, \omega \in \mathbb{R}$. Then,

$$\begin{aligned} \begin{bmatrix} \nu & \omega \\ -\omega & \nu \end{bmatrix} &= \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ j & -j \end{bmatrix} \begin{bmatrix} \nu + j\omega & 0 \\ 0 & \nu - j\omega \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ j & -j \end{bmatrix}^* \\ &= \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & j \\ j & 1 \end{bmatrix} \begin{bmatrix} \nu + j\omega & 0 \\ 0 & \nu - j\omega \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & j \\ j & 1 \end{bmatrix}^* \\ &= \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -j \\ j & -1 \end{bmatrix} \begin{bmatrix} \nu + j\omega & 0 \\ 0 & \nu - j\omega \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -j \\ j & -1 \end{bmatrix} \end{aligned}$$

and

$$\begin{bmatrix} \nu & \omega \\ -\omega & \nu \end{bmatrix}^{-1} = \frac{1}{\nu^2 + \omega^2} \begin{bmatrix} \nu & -\omega \\ \omega & \nu \end{bmatrix}.$$

(Remark: See Fact 2.19.1.) (Remark: All three transformations are unitary. The third transformation is also Hermitian.)

Fact 2.19.3. Let $A, B \in \mathbb{R}^{n \times m}$. Then,

$$\begin{aligned} \begin{bmatrix} A & B \\ -B & A \end{bmatrix} &= \frac{1}{2} \begin{bmatrix} I & I \\ jI & -jI \end{bmatrix} \begin{bmatrix} A + jB & 0 \\ 0 & A - jB \end{bmatrix} \begin{bmatrix} I & -jI \\ I & jI \end{bmatrix} \\ &= \frac{1}{2} \begin{bmatrix} I & jI \\ -jI & -I \end{bmatrix} \begin{bmatrix} A - jB & 0 \\ 0 & A + jB \end{bmatrix} \begin{bmatrix} I & jI \\ -jI & -I \end{bmatrix} \\ &= \begin{bmatrix} I & 0 \\ jI & I \end{bmatrix} \begin{bmatrix} A + jB & B \\ 0 & A - jB \end{bmatrix} \begin{bmatrix} I & 0 \\ -jI & I \end{bmatrix}. \end{aligned}$$

Consequently,

$$\begin{bmatrix} A + jB & 0 \\ 0 & A - jB \end{bmatrix} = \frac{1}{2} \begin{bmatrix} I & -jI \\ I & jI \end{bmatrix} \begin{bmatrix} A & B \\ -B & A \end{bmatrix} \begin{bmatrix} I & I \\ jI & -jI \end{bmatrix},$$

and thus

$$A + jB = \frac{1}{2} \begin{bmatrix} I & -jI \end{bmatrix} \begin{bmatrix} A & B \\ -B & A \end{bmatrix} \begin{bmatrix} I \\ jI \end{bmatrix}.$$

Furthermore,

$$\text{rank}(A + jB) = \text{rank}(A - jB) = \frac{1}{2} \text{rank} \begin{bmatrix} A & B \\ -B & A \end{bmatrix}.$$

Now, assume that $n = m$. Then,

$$\begin{aligned} \det \begin{bmatrix} A & B \\ -B & A \end{bmatrix} &= \det(A + jB) \det(A - jB) \\ &= |\det(A + jB)|^2 \\ &= \det[A^2 + B^2 + j(AB - BA)] \\ &\geq 0. \end{aligned}$$

Hence, $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$ is nonsingular if and only if $A + jB$ is nonsingular. If A is nonsingular, then

$$\det \begin{bmatrix} A & B \\ -B & A \end{bmatrix} = \det(A^2 + ABA^{-1}B).$$

If $AB = BA$, then

$$\det \begin{bmatrix} A & B \\ -B & A \end{bmatrix} = \det(A^2 + B^2).$$

(Proof: If A is nonsingular, then use

$$\begin{bmatrix} A & B \\ -B & A \end{bmatrix} = \begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix} \begin{bmatrix} I & A^{-1}B \\ -A^{-1}B & I \end{bmatrix}$$

and

$$\det \begin{bmatrix} I & A^{-1}B \\ -A^{-1}B & I \end{bmatrix} = \det[I + (A^{-1}B)^2].)$$

(Remark: See Fact 4.10.26 and [79, 1281].)

Fact 2.19.4. Let $A, B \in \mathbb{R}^{n \times m}$. Then, $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$ and $\begin{bmatrix} A & -B \\ B & A \end{bmatrix}$ are representations of the complex matrices $A + jB$ and $\overline{A + jB}$, respectively. Furthermore, $\begin{bmatrix} A^T & B^T \\ -B^T & A^T \end{bmatrix}$ and $\begin{bmatrix} A^T & -B^T \\ B^T & A^T \end{bmatrix}$ are representations of the complex matrices $(A + jB)^T$ and $(A + jB)^*$, respectively.

Fact 2.19.5. Let $A, B \in \mathbb{R}^{n \times m}$ and $C, D \in \mathbb{R}^{m \times l}$. Then, for all $\alpha, \beta \in \mathbb{R}$, $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$, $\begin{bmatrix} C & D \\ -D & C \end{bmatrix}$, and $\begin{bmatrix} \alpha A + \beta C & \alpha B + \beta D \\ -(\alpha B + \beta D) & \alpha A + \beta C \end{bmatrix} = \alpha \begin{bmatrix} A & B \\ -B & A \end{bmatrix} + \beta \begin{bmatrix} C & D \\ -D & C \end{bmatrix}$ are representations of the complex matrices $A + jB$, $C + jD$, and $\alpha(A + jB) + \beta(C + jD)$, respectively.

Fact 2.19.6. Let $A, B \in \mathbb{R}^{n \times m}$ and $C, D \in \mathbb{R}^{m \times l}$. Then, $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$, $\begin{bmatrix} C & D \\ -D & C \end{bmatrix}$, and $\begin{bmatrix} AC - BD & AD + BC \\ -(AD + BC) & AC - BD \end{bmatrix} = \begin{bmatrix} A & B \\ -B & A \end{bmatrix} \begin{bmatrix} C & D \\ -D & C \end{bmatrix}$ are representations of the complex matrices

$A + jB$, $C + jD$, and $(A + jB)(C + jD)$, respectively.

Fact 2.19.7. Let $A, B \in \mathbb{R}^{n \times n}$. Then, $A + jB$ is nonsingular if and only if $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$ is nonsingular. In this case,

$$(A + jB)^{-1} = \frac{1}{2} \begin{bmatrix} I & -jI \end{bmatrix} \begin{bmatrix} A & B \\ -B & A \end{bmatrix}^{-1} \begin{bmatrix} I \\ jI \end{bmatrix}.$$

If A is nonsingular, then $A + jB$ is nonsingular if and only if $A + BA^{-1}B$ is nonsingular. In this case,

$$\begin{bmatrix} A & B \\ -B & A \end{bmatrix}^{-1} = \begin{bmatrix} (A + BA^{-1}B)^{-1} & -A^{-1}B(A + BA^{-1}B)^{-1} \\ A^{-1}B(A + BA^{-1}B)^{-1} & (A + BA^{-1}B)^{-1} \end{bmatrix}$$

and

$$(A + jB)^{-1} = (A + BA^{-1}B)^{-1} - jA^{-1}B(A + BA^{-1}B)^{-1}.$$

Alternatively, if B is nonsingular. Then, $A + jB$ is nonsingular if and only if $B + AB^{-1}A$ is nonsingular. In this case,

$$\begin{bmatrix} A & B \\ -B & A \end{bmatrix}^{-1} = \begin{bmatrix} B^{-1}A(B + AB^{-1}A)^{-1} & -(B + AB^{-1}A)^{-1} \\ (B + AB^{-1}A)^{-1} & B^{-1}A(B + AB^{-1}A)^{-1} \end{bmatrix}$$

and

$$(A + jB)^{-1} = B^{-1}A(B + AB^{-1}A)^{-1} - j(B + AB^{-1}A)^{-1}.$$

(Remark: See Fact 3.11.27, Fact 6.5.1, and [1282].)

Fact 2.19.8. Let $A \in \mathbb{R}^{n \times n}$. Then,

$$\det(I + A\overline{A}) \geq 0.$$

(Proof: See [416].)

Fact 2.19.9. Let $A, B \in \mathbb{R}^{n \times n}$. Then,

$$\det \begin{bmatrix} A & B \\ -\overline{B} & \overline{A} \end{bmatrix} \geq 0.$$

If, in addition, A is nonsingular, then

$$\det \begin{bmatrix} A & B \\ -\overline{B} & \overline{A} \end{bmatrix} = |\det A|^2 \det(I + \overline{A^{-1}B}A^{-1}B).$$

(Proof: See [1489].) (Remark: Fact 2.19.8 implies that $\det(I + \overline{A^{-1}B}A^{-1}B) \geq 0$.)

Fact 2.19.10. Let $A, B \in \mathbb{R}^{n \times n}$, and define $C \in \mathbb{R}^{2n \times 2n}$ by $C \triangleq \begin{bmatrix} C_{11} & C_{12} & \cdots \\ C_{21} & \cdots & \end{bmatrix}$, where $C_{ij} \triangleq \begin{bmatrix} A_{(i,j)} & B_{(i,j)} \\ -B_{(i,j)} & A_{(i,j)} \end{bmatrix} \in \mathbb{R}^{2 \times 2}$ for all $i, j = 1, \dots, n$. Then,

$$\det C = |\det(A + jB)|^2.$$

(Proof: Note that

$$C = A \otimes I_2 + B \otimes J_2 = P_{2,n}(I_2 \otimes A + J_2 \otimes B)P_{2,n} = P_{2,n} \begin{bmatrix} A & B \\ -B & A \end{bmatrix} P_{2,n}.$$

See [257].)

2.20 Facts on Geometry

Fact 2.20.1. The points $x, y, z \in \mathbb{R}^2$ lie on one line if and only if

$$\det \begin{bmatrix} x & y & z \\ 1 & 1 & 1 \end{bmatrix} = 0.$$

Fact 2.20.2. The points $w, x, y, z \in \mathbb{R}^3$ lie in one plane if and only if

$$\det \begin{bmatrix} w & x & y & z \\ 1 & 1 & 1 & 1 \end{bmatrix} = 0.$$

Fact 2.20.3. Let $x_1, \dots, x_n \in \mathbb{R}^n$. Then,

$$\text{rank} \begin{bmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_n \end{bmatrix} = \text{rank} \begin{bmatrix} 1 & 0 & \cdots & 0 \\ x_1 & x_2 - x_1 & \cdots & x_n - x_1 \end{bmatrix}.$$

Hence,

$$\text{rank} \begin{bmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_n \end{bmatrix} = n$$

if and only if

$$\text{rank} \begin{bmatrix} x_2 - x_1 & \cdots & x_n - x_1 \end{bmatrix} = n - 1.$$

In this case,

$$\text{aff} \{x_1, \dots, x_n\} = x_1 + \text{span} \{x_2 - x_1, \dots, x_n - x_1\},$$

and thus $\text{aff} \{x_1, \dots, x_n\}$ is an affine hyperplane. Finally,

$$\text{aff} \{x_1, \dots, x_n\} = \{x \in \mathbb{R}^n : \det \begin{bmatrix} 1 & 1 & \cdots & 1 \\ x & x_1 & \cdots & x_n \end{bmatrix} = 0\}.$$

(Proof: See [1184, p. 31].) (Remark: See Fact 2.20.4.)

Fact 2.20.4. Let $x_1, \dots, x_{n+1} \in \mathbb{R}^n$. Then, the following statements are equivalent:

- i) $\text{co} \{x_1, \dots, x_{n+1}\}$ is a simplex.
- ii) $\text{co} \{x_1, \dots, x_{n+1}\}$ has nonempty interior.
- iii) $\text{aff} \{x_1, \dots, x_{n+1}\} = \mathbb{R}^n$.
- iv) $\text{span} \{x_2 - x_1, \dots, x_{n+1} - x_1\} = \mathbb{R}^n$.
- v) $\begin{bmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_{n+1} \end{bmatrix}$ is nonsingular.

(Proof: The equivalence of i) and ii) follows from Fact 10.8.9. The equivalence of i) and iv) follows from Fact 2.9.7. Finally, the equivalence of iv) and v) follows from

$$\begin{bmatrix} 1 & \cdots & 1 \\ x_1 & \cdots & x_{n+1} \end{bmatrix} = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ x_1 & x_2 - x_1 & \cdots & x_{n+1} - x_1 \end{bmatrix} \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & \cdots & 1 \end{bmatrix}.)$$

(Remark: See Fact 2.20.3 and Fact 10.8.12.)

Fact 2.20.5. Let z_1, z_2, z be complex numbers, and assume that $z_1 \neq z_2$. Then, the following statements are equivalent:

i) z lies on the line passing through z_1 and z_2 .

ii) $\frac{z-z_1}{z_2-z_1}$ is real.

iii) $\det \begin{bmatrix} z - z_1 & \bar{z} - \bar{z}_1 \\ z_2 - z_1 & \bar{z}_2 - \bar{z}_1 \end{bmatrix} = 0$.

iv) $\det \begin{bmatrix} z & \bar{z} & 1 \\ z_1 & \bar{z}_1 & 1 \\ z_2 & \bar{z}_2 & 1 \end{bmatrix} = 0$.

Furthermore, the following statements are equivalent:

v) z lies on the line segment connecting z_1 and z_2 .

vi) $\frac{z-z_1}{z_2-z_1}$ is a positive number.

vii) There exists $\phi \in (-\pi, \pi]$ such that $|z - z_1|e^{j\phi} = |z_2 - z_1|e^{j\phi}$.

(Proof: See [59, pp. 54–56].)

Fact 2.20.6. Let z_1, z_2, z_3 be distinct complex numbers. Then, the following statements are equivalent:

i) z_1, z_2, z_3 are the vertices of an equilateral triangle.

ii) $|z_1 - z_2| = |z_2 - z_3| = |z_3 - z_1|$.

iii) $z_1^2 + z_2^2 + z_3^2 = z_1z_2 + z_2z_3 + z_3z_1$.

iv) $\frac{z_2-z_1}{z_3-z_2} = \frac{z_3-z_2}{z_1-z_3}$.

(Proof: See [59, pp. 70, 71] and [868, p. 316].)

Fact 2.20.7. Let $\mathcal{S} \subset \mathbb{R}^2$ denote the triangle with vertices $\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix} \in \mathbb{R}^2$. Then,

$$\text{area}(\mathcal{S}) = \frac{1}{2} \left| \det \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix} \right|.$$

Fact 2.20.8. Let $\mathcal{S} \subset \mathbb{R}^2$ denote the triangle with vertices $\begin{bmatrix} x_1 \\ y_1 \end{bmatrix}, \begin{bmatrix} x_2 \\ y_2 \end{bmatrix}, \begin{bmatrix} x_3 \\ y_3 \end{bmatrix} \in \mathbb{R}^2$. Then,

$$\text{area}(\mathcal{S}) = \frac{1}{2} \left| \det \begin{bmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{bmatrix} \right|.$$

(Proof: See [1184, p. 32].)

Fact 2.20.9. Let z_1, z_2, z_3 be complex numbers. Then, the area of the triangle $\mathcal{S}$ formed by z_1, z_2, z_3 is given by

$$\text{area}(\mathcal{S}) = \frac{1}{4} \left| \det \begin{bmatrix} z_1 & \overline{z_1} & 1 \\ z_2 & \overline{z_2} & 1 \\ z_3 & \overline{z_3} & 1 \end{bmatrix} \right|.$$

(Proof: See [59, p. 79].)

Fact 2.20.10. Let $\mathcal{S} \subset \mathbb{R}^3$ denote the triangle with vertices $x, y, z \in \mathbb{R}^3$. Then,

$$\text{area}(\mathcal{S}) = \frac{1}{2} \sqrt{[(y-x) \times (z-x)]^T [(y-x) \times (z-x)]}.$$

Fact 2.20.11. Let $\mathcal{S} \subset \mathbb{R}^2$ denote a triangle whose sides have lengths a, b , and c , let A, B, C denote the angles of the triangle opposite the sides having lengths a, b , and c , respectively, define the semiperimeter $s \triangleq \frac{1}{2}(a+b+c)$, let r denote the radius of the largest inscribed circle, and let R denote the radius of the smallest circumscribed circle. Then, the following identities hold:

- i) $A + B + C = \pi$.
- ii) $a^2 + b^2 = c^2 + 2ab \cos C$.
- iii) $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$.
- iv) $\text{area}(\mathcal{S}) = \frac{1}{2}ab \sin C = \frac{c^2}{2} \frac{(\sin A) \sin B}{\sin C}$.
- v) $\text{area}(\mathcal{S}) = \sqrt{s(s-a)(s-b)(s-c)} = rs = \frac{abc}{4R}$.
- vi) $\text{area}(\mathcal{S}) \leq \frac{\sqrt{3}}{12}(a^2 + b^2 + c^2)$.
- vii) If $\mathcal{S}$ is equilateral, then $\text{area}(\mathcal{S}) = \frac{\sqrt{3}}{4}a^2$ and $R = 2r = \frac{\sqrt{3}}{3}a$.
- viii) a, b, c are the roots of the cubic equation

$$x^3 - 2sx^2 + (s^2 + r^2 + 4rR)x - 4srR = 0.$$

That is,

$$a + b + c = 2s, \quad ab + bc + ca = s^2 + r^2 + 4rR, \quad abc = 4rRs.$$

ix) a, b, c satisfy

$$a^2 + b^2 + c^2 = 2(s^2 - r^2 - 4rR)$$

and

$$a^3 + b^3 + c^3 = 2s(s^2 - 3r^2 - 6rR).$$

x) If r_1, r_2, r_3 denote the altitudes of the triangle, then

$$\frac{1}{r} = \frac{1}{r_1} + \frac{1}{r_2} + \frac{1}{r_3}.$$

xi) $r \leq \frac{1}{2} \left(\frac{2}{1+\sqrt{5}} \right)^{5/2} (a+b) \approx 0.15(a+b)$. If, in addition, $\mathcal{S}$ is equilateral, then
 $r = \frac{\sqrt{3}}{12} (a+b) \approx 0.14(a+b)$.

Furthermore, the following statements hold:

xii) $2 \leq \frac{a}{b} + \frac{b}{a} \leq \frac{R}{r}.$

xiii) $2 \leq \frac{2}{3} \left(\frac{a}{b} + \frac{b}{c} + \frac{c}{a} \right) \leq \frac{a}{b} + \frac{b}{c} + \frac{c}{a} - 1 \leq \frac{1}{2} \left(1 + \frac{a^2}{bc} + \frac{b^2}{ca} + \frac{c^2}{ab} \right) \leq \frac{R}{r}.$

xiv) $1 \leq \frac{2a^2}{2a^2-(b-c)^2} \frac{2b^2}{2b^2-(c-a)^2} \frac{2c^2}{2c^2-(a-b)^2} \leq \frac{R}{2r}.$

xv) $\frac{a}{2} \frac{4r-R}{R} \leq \sqrt{(s-b)(s-c)} \leq \frac{a}{2}.$

xvi) A triangle $\mathcal{S}$ with values $\text{area}(\mathcal{S})$, r , and R exists if and only if

$$\begin{aligned} r\sqrt{2R^2 + 10rR - r^2 - 2(R-2r)\sqrt{R(R-2r)}} \\ \leq \text{area}(\mathcal{S}) \leq r\sqrt{2R^2 + 10rR - r^2 + 2(R-2r)\sqrt{R(R-2r)}}. \end{aligned}$$

xvii) Let $\theta \triangleq \min\{|A-B|, |A-C|, |B-C|\}_{\text{ms}}$. Then,

$$\begin{aligned} r\sqrt{2R^2 + 10rR - r^2 - 2(R-2r)\sqrt{R(R-2r)}} \cos \theta \\ \leq \text{area}(\mathcal{S}) \leq r\sqrt{2R^2 + 10rR - r^2 + 2(R-2r)\sqrt{R(R-2r)}} \cos \theta. \end{aligned}$$

xviii) $\text{area}(\mathcal{S}) \leq (R + \frac{1}{2}r)^2.$

xix) $\text{area}(\mathcal{S}) \leq \frac{1}{\sqrt{3}}(R+r)^2.$

xx) $\text{area}(\mathcal{S}) \leq \frac{3\sqrt{3}}{25}(R+3r)^2.$

xxi) $3\sqrt{3}r^2 \leq \text{area}(\mathcal{S}) \leq 2rR + (3\sqrt{3} - 4)r^2.$

xxii) $r\sqrt{16rR - 5r^2} \leq \text{area}(\mathcal{S}) \leq r\sqrt{4R^2 + 4rR + 3r^2}.$

xxiii) For all $n \geq 0$, $a^n + b^n + c^n \leq 2^{n+1}R^n + 2^n(3^{1+n/2} - 2^{1+n})r^n.$

xxiv) A triangle $\mathcal{S}$ with values $u = \cos A$, $v = \cos B$, and $w = \cos C$ exists if and only if $u + v + w \geq 1$, $uvw \geq -1$, and $u^2 + v^2 + w^2 + 2uvw = 1$.

xxv) If P is a point inside $\mathcal{S}$ and d_1, d_2, d_3 are the distances from P to each of the sides, then

$$\sqrt{d_1} + \sqrt{d_2} + \sqrt{d_3} \leq \sqrt{\frac{a^2+b^2+c^2}{2R}}.$$

In particular,

$$18R^2 \leq a^2 + b^2 + c^2.$$

xxvi) $4r^2[8R^2 - (a^2 + b^2 + c^2)] \leq R^2(R^2 - 4r^2).$

xxvii) $abc \leq 3\sqrt{3}R^3.$

xxviii) The triangle $\mathcal{S}$ is similar to the triangle $\mathcal{S}'$ with sides of length a', b', c' if and only if

$$\sqrt{aa'} + \sqrt{bb'} + \sqrt{cc'} = \sqrt{(a+b+c)(a'+b'+c')}.$$

$$xxix) (\sin \tfrac{1}{2}A)(\sin \tfrac{1}{2}B)(\sin \tfrac{1}{2}C) < (\sin \tfrac{1}{2}\sqrt[3]{ABC})^3 < \tfrac{1}{8}.$$

$$xxx) (\cos \tfrac{1}{2}A)(\cos \tfrac{1}{2}B)(\cos \tfrac{1}{2}C) < [\sin \tfrac{1}{2}\sqrt[3]{(\pi-A)(\pi-B)(\pi-C)}]^3.$$

$$xxxi) (\tan \tfrac{1}{2}\sqrt[3]{ABC})^3 < (\tan \tfrac{1}{2}A)(\tan \tfrac{1}{2}B)(\tan \tfrac{1}{2}C).$$

$$xxxi) 1 \leq \tan^2(\tfrac{1}{2}A) + \tan^2(\tfrac{1}{2}B) + \tan^2(\tfrac{1}{2}C).$$

$$xxxi) \tfrac{\pi}{3}(a+b+c) \leq Aa + Bb + Cc \leq \tfrac{\pi - \min\{A, B, C\}}{2}(a+b+c).$$

xxxi) If x, y, z are positive numbers, then

$$\begin{aligned} x \sin A + y \sin B + z \sin C &\leq \tfrac{1}{2}(xy + yz + zx) \sqrt{\tfrac{1}{xy} + \tfrac{1}{yz} + \tfrac{1}{zx}} \\ &\leq \tfrac{\sqrt{3}}{2} \left(\tfrac{yz}{x} + \tfrac{zx}{y} + \tfrac{xy}{z} \right). \end{aligned}$$

$$xxxi) \sin A + \sin B + \sin C \leq \tfrac{3\sqrt{3}}{2}.$$

(Proof: Results i)– v) are classical. The first expression for $\text{area}(\mathcal{S})$ in v) is *Heron's formula*. Statements ii) and iii) are the *cosine rule* and *sine rule*, respectively. See [1503, p. 319]. Statement vi) is due to Weitzenbock. See [59, p. 145] and [457, p. 170]. The expression for $\text{area}(\mathcal{S})$ in vii) follows from v) and provides the case of equality in vi). Statements $viii$) and ix) are given in [59, pp. 110, 111]. Statement xi) is given in [102]. Statements xii) and $xiii$) are given in [1374]. Statement xiv) is due to [1097]. See [457, p. 174]. Statement xv) is given in [1146]. Statement xvi), which is due to Ramus, is the *fundamental triangle inequality*. See [1011]. The interpolation of xvi) given by $xvii$) is given in [1463]. The bounds $xviii$)– xx) are given in [1464]. The bounds xxi) and $xxii$) are due to Blundon. See [1161]. Statement $xxiii$) is given in [1161]. Statement $xxiv$) is given in [622]. Statement xxv) is given in [868, pp. 255, 256]. Statement $xxvi$) follows from [59, p. 189]. Statement $xxvii$) follows from [59, p. 144]. Statement $xxviii$) is given in [457, p. 183]. Necessity is immediate. Statements $xxix$)– $xxxi$) are given in [1040]. Statement $xxxi$) is given in [136, p. 231]. Statement $xxxi$) is given in [971, p. 203]. The first inequality in statement $xxxi$) is *Klamkin's inequality*. The first and third terms comprise it Vasic's inequality. See [1374]. Statement $xxxi$) follows from statement $xxxi$) with $x = y = z = 1$.) (Remark: $2r \leq R$ in xii) is *Euler's inequality*. The interpolation is *Bandila's inequality*. The inequality involving the second and fifth terms in $xiii$) is due to Zhang and Song. See [1374].) (Remark: The bound xxi) is *Mircea's inequality*, while $xxii$) is due to Carliz and Leuenberger. See [1464].) (Remark: Additional inequalities involving the sides and angles of a triangle are given in Fact 1.11.21, [244], and [971, pp. 192–203].) (Remark: The second inequality in $xxxi$) is given in Fact 1.11.10.)

Fact 2.20.12. Let a be a complex number, let $b \in (0, |a|^2)$, and define

$$\mathcal{S} \triangleq \{z \in \mathbb{C} : |z|^2 - \bar{a}z - a\bar{z} + b = 0\}.$$

Then, $\mathcal{S}$ is the circle with center at a and radius $\sqrt{|a|^2 - b}$. That is,

$$\mathcal{S} = \{z \in \mathbb{C} : |z - a| = \sqrt{|a|^2 - b}\}.$$

(Proof: See [59, p. 84, 85].)

Fact 2.20.13. Let $\mathcal{S} \subset \mathbb{R}^2$ be a convex quadrilateral whose sides have lengths a, b, c, d , define the semiperimeter $s \triangleq \frac{1}{2}(a + b + c + d)$, let A, B, C, D denote the angles of $\mathcal{S}$ labeled consecutively, and define $\theta \triangleq \frac{1}{2}(A + C) = \pi - \frac{1}{2}(B + D)$. Then,

$$\text{area}(\mathcal{S}) = \sqrt{(s - a)(s - b)(s - c)(s - d) - abcd \cos^2 \theta}.$$

Now, let p, q be the lengths of the diagonals of $\mathcal{S}$. Then,

$$pq \leq ac + bc$$

and

$$\text{area}(\mathcal{S}) = \sqrt{(s - a)(s - b)(s - c)(s - d) - \frac{1}{4}(ac + bd + pq)(ac + bd - pq)}.$$

If the quadrilateral has an inscribed circle that contacts all four sides of the quadrilateral, then

$$\text{area}(\mathcal{S}) = \sqrt{abcd} = \sqrt{p^2 q^2 - (ac - bd)^2}.$$

Finally, all of the vertices of $\mathcal{S}$ lie on a circle if and only if

$$pq = ac + bc.$$

In this case,

$$\text{area}(\mathcal{S}) = \sqrt{(s - a)(s - b)(s - c)(s - d)}$$

and

$$\text{area}(\mathcal{S}) = \frac{1}{4R} \sqrt{(ad + bc)(ac + bd)(ab + cd)},$$

where R is the radius of the circumscribed circle. (Proof: See [60, pp. 37, 38], Wikipedia, PlanetMath, and MathWorld.) (Remark: $pq \leq ac + bc$ is *Ptolemy's inequality*, which holds for nonconvex quadrilaterals. The equality case is *Ptolemy's theorem*. See [59, p. 130].) (Remark: The fourth expression for $\text{area}(\mathcal{S})$ is *Brahmagupta's formula*. The limiting case $d = 0$ yields Heron's formula. See Fact 2.20.11.) (Remark: For each quadrilateral, there exists a quadrilateral with the same side lengths and whose vertices lie on a circle. The area of the latter quadrilateral is maximum over all quadrilaterals with the same side lengths. See [1082].) (Problem: For which quadrilaterals does there exist a quadrilateral with the same side lengths and whose sides are tangent to an inscribed circle?) (Remark: See Fact 9.7.5.)

Fact 2.20.14. Let $\mathcal{S} \subset \mathbb{R}^2$ denote the polygon with vertices $\begin{bmatrix} x_1 \\ y_1 \end{bmatrix}, \dots, \begin{bmatrix} x_n \\ y_n \end{bmatrix} \in \mathbb{R}^2$ arranged in counterclockwise order, and assume that the interior of the polygon is either empty or simply connected. Then,

$$\begin{aligned} \text{area}(\mathcal{S}) = & \frac{1}{2} \det \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix} + \frac{1}{2} \det \begin{bmatrix} x_2 & x_3 \\ y_2 & y_3 \end{bmatrix} + \dots \\ & + \frac{1}{2} \det \begin{bmatrix} x_{n-1} & x_n \\ y_{n-1} & y_n \end{bmatrix} + \frac{1}{2} \det \begin{bmatrix} x_n & x_1 \\ y_n & y_1 \end{bmatrix}. \end{aligned}$$

(Remark: The polygon need not be convex, while “counterclockwise” is determined with respect to a point in the interior of the polygon. *Simply connected* means that the polygon has no holes. See [1237].) (Remark: See [59, p. 100].) (Remark: See Fact 9.7.5.)

Fact 2.20.15. Let $\mathcal{S} \subset \mathbb{R}^3$ denote the tetrahedron with vertices $x, y, z, w \in \mathbb{R}^3$. Then,

$$\text{volume}(\mathcal{S}) = \frac{1}{6} |(x - w)^T[(y - w) \times (z - w)]|.$$

(Proof: The volume of the unit simplex $\mathcal{S} \subset \mathbb{R}^3$ with vertices $(0, 0, 0), (1, 0, 0), (0, 1, 0), (0, 0, 1)$ is $1/6$. Now, Fact 2.20.18 implies that the volume of $A\mathcal{S}$ is $(1/6)|\det A|$.) (Remark: The connection between the *signed volume* of a simplex and the determinant is discussed in [878, pp. 32, 33].)

Fact 2.20.16. Let $\mathcal{S} \subset \mathbb{R}^3$ denote the parallelepiped with vertices $x, y, z, x + y, x + z, y + z, x + y + z \in \mathbb{R}^3$. Then,

$$\text{volume}(\mathcal{S}) = |\det \begin{bmatrix} x & y & z \end{bmatrix}|.$$

Fact 2.20.17. Let $A \in \mathbb{R}^{n \times m}$, assume that $\text{rank } A = m$, and let $\mathcal{S} \subset \mathbb{R}^n$ denote the parallelepiped in $\mathbb{R}^n$ with a vertex at 0 and generated by the m columns of A , that is,

$$\mathcal{S} = \left\{ \sum_{i=1}^m \alpha_i \text{col}_i(A) : 0 \leq \alpha_i \leq 1 \text{ for all } i = 1, \dots, m \right\}.$$

Then,

$$\text{volume}(\mathcal{S}) = [\det(A^T A)]^{1/2}.$$

If, in addition, $m = n$, then

$$\text{volume}(\mathcal{S}) = |\det A|.$$

(Remark: $\text{volume}(\mathcal{S})$ denotes the m -dimensional volume of $\mathcal{S}$. If $m = 2$, then $\text{volume}(\mathcal{S})$ is the area of a parallelogram. See [447, p. 202].)

Fact 2.20.18. Let $\mathcal{S} \subset \mathbb{R}^n$ and $A \in \mathbb{R}^{n \times n}$. Then,

$$\text{volume}(A\mathcal{S}) = |\det A| \text{volume}(\mathcal{S}).$$

(Remark: See [998, p. 468].)

Fact 2.20.19. Let $\mathcal{S} \subset \mathbb{R}^n$ be a simplex, and assume that $\mathcal{S}$ is inscribed in a sphere of radius R . Then,

$$\text{volume}(\mathcal{S}) \leq \sqrt{\frac{(n+1)^{n+1}}{n^n} \frac{R^n}{n!}}.$$

Furthermore, equality holds if and only if $\mathcal{S}$ is a regular polytope. (Proof: See [1373].) (Remark: See [482, p. 66-13].)

Fact 2.20.20. Let $x_1, \dots, x_{n+1} \in \mathbb{R}^n$, define

$$\mathcal{S} \triangleq \text{co}\{x_1, \dots, x_{n+1}\},$$

and define $A \in \mathbb{R}^{(n+2) \times (n+2)}$ by

$$A \triangleq \begin{bmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & 0 & \|x_1 - x_2\|_2^2 & \cdots & \|x_1 - x_{n+1}\|_2^2 \\ 1 & \|x_2 - x_1\|_2^2 & 0 & \cdots & \|x_2 - x_{n+1}\|_2^2 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \|x_{n+1} - x_1\|_2^2 & \|x_{n+1} - x_2\|_2^2 & \cdots & 0 \end{bmatrix}.$$

Then, the n -dimensional volume of $\mathcal{S}$ is given by

$$\text{vol}(\mathcal{S}) = \frac{\sqrt{|\det A|}}{2^{n-1}n!}.$$

(Proof: See [232, pp. 97–99] and [238, pp. 234, 235].) (Remark: $\det A$ is the *Cayley-Menger determinant*.) (Remark: In the case $n = 2$, this result yields Heron's formula for the area of a triangle. See Fact 2.20.11.)

Fact 2.20.21. Let $\mathcal{S}$ denote the spherical triangle on the surface of the unit sphere whose vertices are $x, y, z \in \mathbb{R}^3$, and let A, B, C denote the angles of $\mathcal{S}$ located at the points x, y, z , respectively. Furthermore, let a, b, c denote the planar angles subtended by the pairs (y, z) , (x, z) , (x, y) , respectively, or, equivalently, a, b, c denote the sides of the spherical triangle opposite A, B, C , respectively. Finally, define the solid angle Ω to be the area of $\mathcal{S}$. Then,

$$\Omega = A + B + C - \pi.$$

Furthermore,

$$\tan \frac{\Omega}{2} = \frac{|\begin{bmatrix} x & y & z \end{bmatrix}|}{1 + x^T y + x^T z + y^T z}.$$

Equivalently,

$$\tan \frac{\Omega}{2} = \frac{\sqrt{1 - \cos^2 a - \cos^2 b - \cos^2 c + 2(\cos a)(\cos b)\cos c}}{1 + \cos a + \cos b + \cos c}.$$

Finally,

$$\tan \frac{\Omega}{4} = \sqrt{(\tan \frac{s}{2})(\tan \frac{s-a}{2})(\tan \frac{s-b}{2}) \tan \frac{s-c}{2}}.$$

(Proof: See [461] and [1503, pp. 368–371].) (Remark: Spherical triangles are discussed in [477, pp. 253–260], [753, Chapter 2], [1425, pp. 904–907], and [1436, pp. 26–29]. A linear algebraic approach is given in [127].)

Fact 2.20.22. Let $\mathcal{S}$ denote a circular cap on the surface of the unit sphere, where the angle subtended by cross sections of the cone with apex at the center of the sphere is 2θ . Furthermore, define the solid angle Ω to be the area of $\mathcal{S}$. Then,

$$\Omega = 2\pi(1 - \cos \theta).$$

Fact 2.20.23. Let $\mathcal{S}$ denote a region on the surface of the unit sphere subtended by the sides of a right rectangular pyramid with apex at the center of the sphere, where the subtended planar angles of the edges of the pyramid are θ and

ϕ . Furthermore, define the solid angle Ω to be the area of $\mathcal{S}$. Then,

$$\Omega = 4 \sin^{-1} \left[\left(\sin \frac{\theta}{2} \right) \sin \frac{\phi}{2} \right].$$

2.21 Facts on Majorization

Fact 2.21.1. Let $x \in \mathbb{R}^n$, where $x_{(1)} \geq \cdots \geq x_{(n)} \geq 0$, and assume that $\sum_{i=1}^n x_{(i)} = 1$. Then, $e_{1,n}$ strongly majorizes x , and x strongly majorizes $\frac{1}{n} \mathbf{1}_{n \times 1}$. (Proof: See [971, p. 95].) (Remark: See Fact 2.21.2.)

Fact 2.21.2. Let $x, y, z \in \mathbb{R}^n$, assume that $x_{(1)} \geq \cdots \geq x_{(n)}$, $y_{(1)} \geq \cdots \geq y_{(n)}$, and $z_{(1)} \geq \cdots \geq z_{(n)} \geq 0$, and assume that y weakly majorizes x . Then,

$$x^T z \leq y^T z.$$

(Proof: See [971, p. 95].) (Remark: See Fact 2.21.3.)

Fact 2.21.3. Let $x, y, z \in \mathbb{R}^n$, assume that $x_{(1)} \geq \cdots \geq x_{(n)}$, $y_{(1)} \geq \cdots \geq y_{(n)}$, and $z_{(1)} \geq \cdots \geq z_{(n)}$, and assume that y strongly majorizes x . Then,

$$x^T z \leq y^T z.$$

(Proof: See [971, p. 92].)

Fact 2.21.4. Let $a < b$, let $f: (a, b)^n \mapsto \mathbb{R}$, and assume that f is C^1 . Then, f is Schur convex if and only if f is symmetric and, for all $x \in (a, b)^n$,

$$(x_{(1)} - x_{(2)}) \left(\frac{\partial f(x)}{\partial x_{(1)}} - \frac{\partial f(x)}{\partial x_{(2)}} \right) \geq 0.$$

(Proof: See [971, p. 57].) (Remark: f is symmetric means that $f(Ax) = f(x)$ for all $x \in (a, b)^n$ and for every permutation matrix $A \in \mathbb{R}^{n \times n}$. (Remark: See [779].)

Fact 2.21.5. Let $x, y \in \mathbb{R}^n$, assume that $x_{(1)} \geq \cdots \geq x_{(n)} \geq 0$ and $y_{(1)} \geq \cdots \geq y_{(n)} \geq 0$, assume that y strongly majorizes x , and let $p_1, \dots, p_n$ be nonnegative numbers. Then,

$$\sum_{j=1}^n \prod_{i_j}^{x_{(j)}} p_{i_j} \leq \frac{1}{n!} \sum_{j=1}^n \prod_{i_j}^{y_{(j)}} p_{i_j}$$

where the summation is taken over all $n!$ permutations $\{i_1, \dots, i_n\}$ of $\{1, \dots, n\}$. (Proof: See [542, p. 99] and [971, p. 88].) (Remark: This result is *Muirhead's theorem*, which is based on a function that is *Schur convex*. An immediate consequence is an interpolated version of the arithmetic-mean-geometric-mean inequality. See Fact 1.15.25.)

Fact 2.21.6. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \cdots \geq x_{(n)} \geq 0$ and $y_{(1)} \geq \cdots \geq y_{(n)} \geq 0$, assume that y strongly majorizes x , and assume that $\sum_{i=1}^n x_{(i)} = 1$. Then,

$$\sum_{i=1}^n y_i \log \frac{1}{y_{(i)}} \leq \sum_{i=1}^n x_i \log \frac{1}{x_{(i)}} \leq \log n.$$

(Proof: See [542, p. 102] and [971, pp. 71, 405].) (Remark: For $x_{(1)}, x_{(2)} > 0$, note that $(x_{(1)} - x_{(2)}) \log(x_{(1)}/x_{(2)}) \geq 0$. Hence, it follows from Fact 2.21.4 that the entropy function is *Schur concave*.) (Remark: Entropy bounds are given in Fact 1.15.45, Fact 1.15.46, and Fact 1.15.47.)

Fact 2.21.7. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \cdots \geq x_{(n)}$ and $y_{(1)} \geq \cdots \geq y_{(n)}$. Then, the following statements are equivalent:

- i) y strongly majorizes x .
- ii) x is an element of the convex hull of the vectors $y_1, \dots, y_n! \in \mathbb{R}^n$, where each of these $n!$ vectors is formed by permuting the components of y .
- iii) There exists a doubly stochastic matrix $A \in \mathbb{R}^{n \times n}$ such that $y = Ax$.

(Proof: The equivalence of i) and ii) is due to Rado. See [971, p. 113]. The equivalence of i) and iii) is the *Hardy-Littlewood-Polya theorem*. See [197, p. 33], [709, p. 197], and [971, p. 22].) (Remark: See Fact 8.17.8.) (Remark: The matrix A is *doubly stochastic* if it is nonnegative, $1_{1 \times n}A = 1_{1 \times n}$, and $A1_{n \times 1} = 1_{n \times 1}$.)

Fact 2.21.8. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \cdots \geq x_{(n)}$ and $y_{(1)} \geq \cdots \geq y_{(n)}$, assume that y strongly majorizes x , let $f: [\min\{x_{(n)}, y_{(n)}\}, y_{(1)}] \mapsto \mathbb{R}$, assume that f is convex, and let $\{i_1, \dots, i_n\} = \{j_1, \dots, j_n\} = \{1, \dots, n\}$ be such that $f(x_{(i_1)}) \geq \cdots \geq f(x_{(i_n)})$ and $f(y_{(i_1)}) \geq \cdots \geq f(y_{(i_n)})$. Then, $\begin{bmatrix} f(y_{(j_1)}) & \cdots & f(y_{(j_n)}) \end{bmatrix}^T$ weakly majorizes $\begin{bmatrix} f(x_{(i_1)}) & \cdots & f(x_{(i_n)}) \end{bmatrix}^T$. (Proof: See [197, p. 42], [711, p. 173], or [971, p. 116].)

Fact 2.21.9. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \cdots \geq x_{(n)} \geq 0$ and $y_{(1)} \geq \cdots \geq y_{(n)} \geq 0$, assume that y strongly log majorizes x , let $f: [0, \infty) \mapsto \mathbb{R}$, assume that $g: \mathbb{R} \mapsto \mathbb{R}$ defined by $g(z) \triangleq f(e^z)$ is convex, and let $\{i_1, \dots, i_n\} = \{j_1, \dots, j_n\} = \{1, \dots, n\}$ be such that $f(x_{(i_1)}) \geq \cdots \geq f(x_{(i_n)})$ and $f(y_{(j_1)}) \geq \cdots \geq f(y_{(j_n)})$. Then, $\begin{bmatrix} f(y_{(j_1)}) & \cdots & f(y_{(j_n)}) \end{bmatrix}^T$ weakly majorizes $\begin{bmatrix} f(x_{(i_1)}) & \cdots & f(x_{(i_n)}) \end{bmatrix}^T$. (Proof: Apply Fact 2.21.8.)

Fact 2.21.10. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \cdots \geq x_{(n)}$ and $y_{(1)} \geq \cdots \geq y_{(n)}$, assume that y weakly majorizes x , let $f: [\min\{x_{(n)}, y_{(n)}\}, y_{(1)}] \mapsto \mathbb{R}$, assume that f is convex and increasing, and let $\{i_1, \dots, i_n\} = \{j_1, \dots, j_n\} = \{1, \dots, n\}$ be such that $f(x_{(i_1)}) \geq \cdots \geq f(x_{(i_n)})$ and $f(y_{(j_1)}) \geq \cdots \geq f(y_{(j_n)})$. Then, $\begin{bmatrix} f(y_{(j_1)}) & \cdots & f(y_{(j_n)}) \end{bmatrix}^T$ weakly majorizes $\begin{bmatrix} f(x_{(i_1)}) & \cdots & f(x_{(i_n)}) \end{bmatrix}^T$. (Proof: See [197, p. 42], [711, p. 173], or [971, p. 116].) (Remark: See Fact 2.21.11.)

Fact 2.21.11. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \cdots \geq x_{(n)} \geq 0$ and $y_{(1)} \geq \cdots \geq y_{(n)} \geq 0$, assume that y strongly majorizes x , and let $r \geq 1$. Then, $\begin{bmatrix} y_{(1)}^r & \cdots & y_{(n)}^r \end{bmatrix}^T$ weakly majorizes $\begin{bmatrix} x_{(1)}^r & \cdots & x_{(n)}^r \end{bmatrix}^T$. (Proof: Use Fact 2.21.11.) (Remark: Using the Schur power (see Section 7.3), the conclusion can be stated as the fact that y^{or} weakly majorizes x^{or} .)

Fact 2.21.12. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \cdots \geq x_{(n)} \geq 0$ and $y_{(1)} \geq \cdots \geq y_{(n)} \geq 0$, assume that y weakly log majorizes x , let $f: [0, \infty) \mapsto \mathbb{R}$, as-

sume that $g: \mathbb{R} \mapsto \mathbb{R}$ defined by $g(z) \triangleq f(e^z)$ is convex and increasing, and let $\{i_1, \dots, i_n\} = \{j_1, \dots, j_n\} = \{1, \dots, n\}$ be such that $f(x_{(i_1)}) \geq \dots \geq f(x_{(i_n)})$ and $f(y_{(j_1)}) \geq \dots \geq f(y_{(j_n)})$. Then, $[f(y_{(j_1)}) \ \dots \ f(y_{(j_n)})]^T$ weakly majorizes $[f(x_{(i_1)}) \ \dots \ f(x_{(i_n)})]^T$. (Proof: Use Fact 2.21.10.)

Fact 2.21.13. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \dots \geq x_{(n)} \geq 0$ and $y_{(1)} \geq \dots \geq y_{(n)} \geq 0$, and assume that y weakly log majorizes x . Then, y weakly majorizes x . (Proof: Use Fact 2.21.12 with $f(t) = t$. See [1485, p. 19].)

Fact 2.21.14. Let $x, y \in \mathbb{R}^n$, where $x_{(1)} \geq \dots \geq x_{(n)} \geq 0$ and $y_{(1)} \geq \dots \geq y_{(n)} \geq 0$, assume that y weakly majorizes x , and let $p \in [1, \infty)$. Then, for all $k = 1, \dots, n$,

$$\left(\sum_{i=1}^k x_{(i)}^p \right)^{1/p} \leq \left(\sum_{i=1}^k y_{(i)}^p \right)^{1/p}.$$

(Proof: Use Fact 2.21.10. See [971, p. 96].) (Remark: $\phi(x) \triangleq \left(\sum_{i=1}^k x_{(i)}^p \right)^{1/p}$ is a *symmetric gauge function*. See Fact 9.8.42.)

2.22 Notes

The theory of determinants is discussed in [1023, 1346]. Applications to physics are described in [1371, 1372]. Contributors to the development of this subject are highlighted in [581]. The empty matrix is discussed in [382, 1032], [1129, pp. 462–464], and [1235, p. 3]. Recent versions of Matlab follow the properties of the empty matrix given in this chapter [676, pp. 305, 306]. Convexity is the subject of [180, 239, 255, 450, 879, 1133, 1235, 1355, 1412]. Convex optimization theory is developed in [176, 255]. In [239] the dual cone is called the *polar cone*.

The development of rank properties is based on [968]. Theorem 2.6.4 is based on [1045]. The term “subdeterminant” is used in [1081] and is equivalent to *minor*. The notation A^A for adjugate is used in [1228]. Numerous papers on basic topics in matrix theory and linear algebra are collected in [292, 293]. A geometric interpretation of $\mathcal{N}(A)$, $\mathcal{R}(A)$, $\mathcal{N}(A^*)$, and $\mathcal{R}(A^T)$ is given in [1239]. Some reflections on matrix theory are given in [1259, 1276]. Applications of the matrix inversion lemma are discussed in [619]. Some historical notes on the determinant and inverse of partitioned matrices as well as the matrix inversion lemma are given in [666].

The implications of majorization are extensively developed in [971, 973].

Chapter Three

Matrix Classes and Transformations

This chapter presents definitions of various types of matrices as well as transformations for analyzing matrices.

3.1 Matrix Classes

In this section we categorize various types of matrices based on their algebraic and structural properties.

The following definition introduces various types of square matrices.

Definition 3.1.1. For $A \in \mathbb{F}^{n \times n}$ define the following types of matrices:

- i) A is *group invertible* if $\mathcal{R}(A) = \mathcal{R}(A^2)$.
- ii) A is *involutory* if $A^2 = I$.
- iii) A is *skew involutory* if $A^2 = -I$.
- iv) A is *idempotent* if $A^2 = A$.
- v) A is *skew idempotent* if $A^2 = -A$.
- vi) A is *tripotent* if $A^3 = A$.
- vii) A is *nilpotent* if there exists $k \in \mathbb{P}$ such that $A^k = 0$.
- viii) A is *unipotent* if $A - I$ is nilpotent.
- ix) A is *range Hermitian* if $\mathcal{R}(A) = \mathcal{R}(A^*)$.
- x) A is *range symmetric* if $\mathcal{R}(A) = \mathcal{R}(A^T)$.
- xi) A is *Hermitian* if $A = A^*$.
- xii) A is *symmetric* if $A = A^T$.
- xiii) A is *skew Hermitian* if $A = -A^*$.
- xiv) A is *skew symmetric* if $A = -A^T$.
- xv) A is *normal* if $AA^* = A^*A$.
- xvi) A is *positive semidefinite* ($A \geq 0$) if A is Hermitian and $x^*Ax \geq 0$ for all

$x \in \mathbb{F}^n$.

- xvii) A is *negative semidefinite* ($A \leq 0$) if $-A$ is positive semidefinite.
- xviii) A is *positive definite* ($A > 0$) if A is Hermitian and $x^*Ax > 0$ for all $x \in \mathbb{F}^n$ such that $x \neq 0$.
- xix) A is *negative definite* ($A < 0$) if $-A$ is positive definite.
- xx) A is *semidissipative* if $A + A^*$ is negative semidefinite.
- xxi) A is *dissipative* if $A + A^*$ is negative definite.
- xxii) A is *unitary* if $A^*A = I$.
- xxiii) A is *shifted unitary* if $A + A^* = 2A^*A$.
- xxiv) A is *orthogonal* if $A^T A = I$.
- xxv) A is *shifted orthogonal* if $A + A^T = 2A^T A$.
- xxvi) A is a *projector* if A is Hermitian and idempotent.
- xxvii) A is a *reflector* if A is Hermitian and unitary.
- xxviii) A is a *skew reflector* if A is skew Hermitian and unitary.
- xxix) A is an *elementary projector* if there exists a nonzero vector $x \in \mathbb{F}^n$ such that $A = I - (x^*x)^{-1}xx^*$.
- xxx) A is an *elementary reflector* if there exists a nonzero vector $x \in \mathbb{F}^n$ such that $A = I - 2(x^*x)^{-1}xx^*$.
- xxxi) A is an *elementary matrix* if there exist vectors $x, y \in \mathbb{F}^n$ such that $A = I - xy^T$ and $x^T y \neq 1$.
- xxxi) A is *reverse Hermitian* if $A = A^*$.
- xxxi) A is *reverse symmetric* if $A = A^{\hat{T}}$.
- xxxi) A is a *permutation matrix* if each row of A and each column of A possesses one 1 and zeros otherwise.
- xxxi) A is *reducible* if either $n = 1$ and $A = 0$ or $n \geq 2$ and there exist $k \geq 1$ and a permutation matrix $S \in \mathbb{R}^{n \times n}$ such that $SAS^T = \begin{bmatrix} B & C \\ 0_{k \times (n-k)} & D \end{bmatrix}$, where $B \in \mathbb{F}^{(n-k) \times (n-k)}$, $C \in \mathbb{F}^{(n-k) \times k}$, and $D \in \mathbb{F}^{k \times k}$.
- xxxi) A is *irreducible* if A is not reducible.

Let $A \in \mathbb{F}^{n \times n}$ be Hermitian. Then, the function $f: \mathbb{F}^n \mapsto \mathbb{R}$ defined by

$$f(x) \triangleq x^*Ax \quad (3.1.1)$$

is a *quadratic form*.

The $n \times n$ *standard nilpotent matrix*, which has 1's on the superdiagonal and 0's elsewhere, is denoted by N_n or just N . We define $N_1 \triangleq 0$ and $N_0 \triangleq 0_{0 \times 0}$.

The following definition considers matrices that are not necessarily square.

Definition 3.1.2. For $A \in \mathbb{F}^{n \times m}$ define the following types of matrices:

- i) A is *semicontractive* if $I_n - AA^*$ is positive semidefinite.
- ii) A is *contractive* if $I_n - AA^*$ is positive definite.
- iii) A is *left inner* if $A^*A = I_m$.
- iv) A is *right inner* if $AA^* = I_n$.
- v) A is *centrohermitian* if $A = \hat{I}_n \bar{A} \hat{I}_m$.
- vi) A is *centrosymmetric* if $A = \hat{I}_n A \hat{I}_m$.
- vii) A is an *outer-product matrix* if there exist $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$ such that $A = xy^T$.

The following definition introduces various types of structured matrices.

Definition 3.1.3. For $A \in \mathbb{F}^{n \times m}$ define the following types of matrices:

- i) A is *diagonal* if $A_{(i,j)} = 0$ for all $i \neq j$. If $n = m$, then

$$A = \text{diag}(A_{(1,1)}, \dots, A_{(n,n)}).$$
- ii) A is *tridiagonal* if $A_{(i,j)} = 0$ for all $|i - j| > 1$.
- iii) A is *reverse diagonal* if $A_{(i,j)} = 0$ for all $i + j \neq \min\{n, m\} + 1$. If $n = m$, then

$$A = \text{revdiag}(A_{(1,n)}, \dots, A_{(n,1)}).$$
- iv) A is (*upper triangular, strictly upper triangular*) if $A_{(i,j)} = 0$ for all $(i \geq j, i > j)$.
- v) A is (*lower triangular, strictly lower triangular*) if $A_{(i,j)} = 0$ for all $(i \leq j, i < j)$.
- vi) A is (*upper Hessenberg, lower Hessenberg*) if $A_{(i,j)} = 0$ for all $(i > j + 1, i < j + 1)$.
- vii) A is *Toeplitz* if $A_{(i,j)} = A_{(k,l)}$ for all $k - i = l - j$, that is,

$$A = \begin{bmatrix} a & b & c & \cdots \\ d & a & b & \ddots \\ e & d & a & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{bmatrix}.$$

- viii) A is *Hankel* if $A_{(i,j)} = A_{(k,l)}$ for all $i + j = k + l$, that is,

$$A = \begin{bmatrix} a & b & c & \cdots \\ b & c & d & \ddots \\ c & d & e & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{bmatrix}.$$

ix) A is *block diagonal* if

$$A = \begin{bmatrix} A_1 & & 0 \\ & \ddots & \\ 0 & & A_k \end{bmatrix} = \text{diag}(A_1, \dots, A_k),$$

where $A_i \in \mathbb{F}^{n_i \times m_i}$ for all $i = 1, \dots, k$.

x) A is *upper block triangular* if

$$A = \begin{bmatrix} A_{11} & A_{12} & \cdots & A_{1k} \\ 0 & A_{22} & \cdots & A_{2k} \\ \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & A_{kk} \end{bmatrix},$$

where $A_{ij} \in \mathbb{F}^{n_i \times n_j}$ for all $i, j = 1, \dots, k$.

xi) A is *lower block triangular* if

$$A = \begin{bmatrix} A_{11} & 0 & \cdots & 0 \\ A_{21} & A_{22} & \ddots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ A_{k1} & A_{k2} & \cdots & A_{kk} \end{bmatrix},$$

where $A_{ij} \in \mathbb{F}^{n_i \times n_j}$ for all $i, j = 1, \dots, k$.

xii) A is *block Toeplitz* if $A_{(i,j)} = A_{(k,l)}$ for all $k - i = l - j$, that is,

$$A = \begin{bmatrix} A_1 & A_2 & A_3 & \cdots \\ A_4 & A_1 & A_2 & \ddots \\ A_5 & A_4 & A_1 & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{bmatrix},$$

where $A_i \in \mathbb{F}^{n_i \times m_i}$.

xiii) A is *block Hankel* if $A_{(i,j)} = A_{(k,l)}$ for all $i + j = k + l$, that is,

$$A = \begin{bmatrix} A_1 & A_2 & A_3 & \cdots \\ A_2 & A_3 & A_4 & \ddots \\ A_3 & A_4 & A_5 & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{bmatrix},$$

where $A_i \in \mathbb{F}^{n_i \times m_i}$.

Definition 3.1.4. For $A \in \mathbb{R}^{n \times m}$ define the following types of matrices:

i) A is *nonnegative* ($A \geq 0$) if $A_{(i,j)} \geq 0$ for all $i = 1, \dots, n$ and $j = 1, \dots, m$.

ii) A is *positive* ($A >> 0$) if $A_{(i,j)} > 0$ for all $i = 1, \dots, n$ and $j = 1, \dots, m$.

Now, assume that $n = m$. Then, define the following types of matrices:

iii) A is *almost nonnegative* if $A_{(i,j)} \geq 0$ for all $i, j = 1, \dots, n$ such that $i \neq j$.

iv) A is a Z -matrix if $-A$ is almost nonnegative.

Define the *unit imaginary matrix* $J_{2n} \in \mathbb{R}^{2n \times 2n}$ (or just J) by

$$J_{2n} \triangleq \begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}. \quad (3.1.2)$$

In particular,

$$J_2 = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}. \quad (3.1.3)$$

Note that J_{2n} is skew symmetric and orthogonal, that is,

$$J_{2n}^T = -J_{2n} = J_{2n}^{-1}. \quad (3.1.4)$$

Hence, J_{2n} is skew involutory and a skew reflector.

The following definition introduces structured matrices of even order. Note that $\mathbb{F}$ can represent either $\mathbb{R}$ or $\mathbb{C}$, although A^T does not become A^* in the latter case.

Definition 3.1.5. For $A \in \mathbb{F}^{2n \times 2n}$ define the following types of matrices:

- i) A is *Hamiltonian* if $J^{-1}A^T J = -A$.
- ii) A is *symplectic* if A is nonsingular and $J^{-1}A^T J = A^{-1}$.

Proposition 3.1.6. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) If A is Hermitian, skew Hermitian, or unitary, then A is normal.
- ii) If A is nonsingular or normal, then A is range Hermitian.
- iii) If A is range Hermitian, idempotent, or tripotent, then A is group invertible.
- iv) If A is a reflector, then A is tripotent.
- v) If A is a permutation matrix, then A is orthogonal.

Proof. i) is immediate. To prove ii), note that, if A is nonsingular, then $\mathcal{R}(A) = \mathcal{R}(A^*) = \mathbb{F}^n$, and thus A is range Hermitian. If A is normal, then it follows from Theorem 2.4.3 that $\mathcal{R}(A) = \mathcal{R}(AA^*) = \mathcal{R}(A^*A) = \mathcal{R}(A^*)$, which proves that A is range Hermitian. To prove iii), note that, if A is range Hermitian, then $\mathcal{R}(A) = \mathcal{R}(AA^*) = A\mathcal{R}(A^*) = A\mathcal{R}(A) = \mathcal{R}(A^2)$, while, if A is idempotent, then $\mathcal{R}(A) = \mathcal{R}(A^2)$. If A is tripotent, then $\mathcal{R}(A) = \mathcal{R}(A^3) = A^2\mathcal{R}(A) \subseteq \mathcal{R}(A^2) = A\mathcal{R}(A) \subseteq \mathcal{R}(A)$. Hence, $\mathcal{R}(A) = \mathcal{R}(A^2)$. $\square$

Proposition 3.1.7. Let $\mathcal{A} \in \mathbb{F}^{2n \times 2n}$. Then, $\mathcal{A}$ is Hamiltonian if and only if there exist matrices $A, B, C \in \mathbb{F}^{n \times n}$ such that B and C are symmetric and

$$\mathcal{A} = \begin{bmatrix} A & B \\ C & -A^T \end{bmatrix}. \quad (3.1.5)$$

3.2 Matrices Based on Graphs

Definition 3.2.1. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph, where $\mathcal{X} = \{x_1, \dots, x_n\}$. Then, the following terminology is defined:

- i) The *adjacency matrix* $A \in \mathbb{R}^{n \times n}$ of $\mathcal{G}$ is given by $A_{(i,j)} = 1$ if $(x_j, x_i) \in \mathcal{R}$ and $A_{(i,j)} = 0$ if $(x_j, x_i) \notin \mathcal{R}$, for all $i, j = 1, \dots, n$.
- ii) The *inbound Laplacian matrix* $L_{\text{in}} \in \mathbb{R}^{n \times n}$ of $\mathcal{G}$ is given by $L_{\text{in}(i,i)} = \sum_{j=1, j \neq i}^n A_{(i,j)}$, for all $i = 1, \dots, n$, and $L_{\text{in}(i,j)} = -A_{(i,j)}$, for all distinct $i, j = 1, \dots, n$.
- iii) The *outbound Laplacian matrix* $L_{\text{out}} \in \mathbb{R}^{n \times n}$ of $\mathcal{G}$ is given by $L_{\text{out}(i,i)} = \sum_{j=1, j \neq i}^n A_{(j,i)}$, for all $i = 1, \dots, n$, and $L_{\text{out}(i,j)} = -A_{(i,j)}$, for all distinct $i, j = 1, \dots, n$.
- iv) The *indegree matrix* $D_{\text{in}} \in \mathbb{R}^{n \times n}$ is the diagonal matrix such that $D_{\text{in}(i,i)} = \text{indeg}(x_i)$, for all $i = 1, \dots, n$.
- v) The *outdegree matrix* $D_{\text{out}} \in \mathbb{R}^{n \times n}$ is the diagonal matrix such that $D_{\text{out}(i,i)} = \text{outdeg}(x_i)$, for all $i = 1, \dots, n$.
- vi) Assume that $\mathcal{G}$ has no self-loops, and let $\mathcal{R} = \{a_1, \dots, a_m\}$. Then, the *incidence matrix* $B \in \mathbb{R}^{n \times m}$ of $\mathcal{G}$ is given by $B_{(i,j)} = 1$ if i is the tail of a_j , $B_{(i,j)} = -1$ if i is the head of a_j , and $B_{(i,j)} = 0$ otherwise, for all $i = 1, \dots, n$ and $j = 1, \dots, m$.
- vii) If $\mathcal{G}$ is symmetric, then the *Laplacian matrix* of $\mathcal{G}$ is given by $L \triangleq L_{\text{in}} = L_{\text{out}}$.
- viii) If $\mathcal{G}$ is symmetric, then the *degree matrix* $D \in \mathbb{R}^{n \times n}$ of $\mathcal{G}$ is given by $D \triangleq D_{\text{in}} = D_{\text{out}}$.
- ix) If $\mathcal{G} = (\mathcal{X}, \mathcal{R}, w)$ is a weighted graph, then the *adjacency matrix* $A \in \mathbb{R}^{n \times n}$ of $\mathcal{G}$ is given by $A_{(i,j)} = w[(x_j, x_i)]$ if $(x_j, x_i) \in \mathcal{R}$ and $A_{(i,j)} = 0$ if $(x_j, x_i) \notin \mathcal{R}$, for all $i, j = 1, \dots, n$.

Note that the adjacency matrix is nonnegative, while the inbound Laplacian, outbound Laplacian, and Laplacian matrices are Z-matrices. Furthermore, note that the inbound Laplacian, outbound Laplacian, and Laplacian matrices are unaffected by the presence of self-loops. However, the indegree and outdegree matrices account for self-loops. It can be seen that, for the arc a_i given by (x_k, x_l) , the i th column of B is given by $\text{col}_i(B) = e_l - e_k$. Finally, if $\mathcal{G}$ is a symmetric graph, then A and L are symmetric.

Theorem 3.2.2. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph, where $\mathcal{X} = \{x_1, \dots, x_n\}$, and let L_{in} , L_{out} , D_{in} , D_{out} , and A denote the inbound Laplacian, outbound Laplacian, indegree, outdegree, and adjacency matrices of $\mathcal{G}$, respectively. Then,

$$L_{\text{in}} = D_{\text{in}} - A \quad (3.2.1)$$

and

$$L_{\text{out}} = D_{\text{out}} - A. \quad (3.2.2)$$

Theorem 3.2.3. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a symmetric graph, where $\mathcal{X} = \{x_1, \dots, x_n\}$, and let A , L , D , and B denote the adjacency, Laplacian, degree, and incidence matrices of $\mathcal{G}$, respectively. Then,

$$L = D - A. \quad (3.2.3)$$

Now, assume that $\mathcal{G}$ has no self-loops. Then,

$$L = \frac{1}{2}BB^T. \quad (3.2.4)$$

Definition 3.2.4. Let $M \in \mathbb{F}^{n \times n}$, and let $\mathcal{X} = \{x_1, \dots, x_n\}$. Then, the *graph of M* is $\mathcal{G}(M) \triangleq (\mathcal{X}, \mathcal{R})$, where, for all $i, j = 1, \dots, n$, $(x_j, x_i) \in \mathcal{R}$ if and only if $M_{(i,j)} \neq 0$.

Proposition 3.2.5. Let $M \in \mathbb{F}^{n \times n}$. Then, the adjacency matrix A of $\mathcal{G}(M)$ is given by

$$A = \text{sign } |M|. \quad (3.2.5)$$

3.3 Lie Algebras and Groups

In this section we introduce Lie algebras and groups. Lie groups are discussed in Section 11.5. In the following definition, note that the coefficients α and β are required to be real when $\mathbb{F} = \mathbb{C}$.

Definition 3.3.1. Let $\mathcal{S} \subseteq \mathbb{F}^{n \times n}$. Then, $\mathcal{S}$ is a *Lie algebra* if the following conditions are satisfied:

- i) If $A, B \in \mathcal{S}$ and $\alpha, \beta \in \mathbb{R}$, then $\alpha A + \beta B \in \mathcal{S}$.
- ii) If $A, B \in \mathcal{S}$, then $[A, B] \in \mathcal{S}$.

Note that, if $\mathbb{F} = \mathbb{R}$, then statement i) is equivalent to the statement that $\mathcal{S}$ is a subspace. However, if $\mathbb{F} = \mathbb{C}$ and $\mathcal{S}$ contains matrices that are not real, then $\mathcal{S}$ is not a subspace.

Proposition 3.3.2. The following sets are Lie algebras:

- i) $\mathfrak{gl}_{\mathbb{F}}(n) \triangleq \mathbb{F}^{n \times n}$.
- ii) $\mathfrak{pl}_{\mathbb{C}}(n) \triangleq \{A \in \mathbb{C}^{n \times n}: \text{tr } A \in \mathbb{R}\}$.
- iii) $\mathfrak{sl}_{\mathbb{F}}(n) \triangleq \{A \in \mathbb{F}^{n \times n}: \text{tr } A = 0\}$.
- iv) $\mathfrak{u}(n) \triangleq \{A \in \mathbb{C}^{n \times n}: A \text{ is skew Hermitian}\}$.
- v) $\mathfrak{su}(n) \triangleq \{A \in \mathbb{C}^{n \times n}: A \text{ is skew Hermitian and } \text{tr } A = 0\}$.
- vi) $\mathfrak{so}(n) \triangleq \{A \in \mathbb{R}^{n \times n}: A \text{ is skew symmetric}\}$.
- vii) $\mathfrak{su}(n, m) \triangleq \{A \in \mathbb{C}^{(n+m) \times (n+m)}: \text{diag}(I_n, -I_m)A^* \text{diag}(I_n, -I_m) = -A \text{ and } \text{tr } A = 0\}$.
- viii) $\mathfrak{so}(n, m) \triangleq \{A \in \mathbb{R}^{(n+m) \times (n+m)}: \text{diag}(I_n, -I_m)A^T \text{diag}(I_n, -I_m) = -A\}$.

$ix)$ $\text{symp}_{\mathbb{F}}(2n) \triangleq \{A \in \mathbb{F}^{2n \times 2n}: A \text{ is Hamiltonian}\}.$

$x)$ $\text{osymp}_{\mathbb{C}}(2n) \triangleq \text{su}(2n) \cap \text{symp}_{\mathbb{C}}(2n).$

$xi)$ $\text{osymp}_{\mathbb{R}}(2n) \triangleq \text{so}(2n) \cap \text{symp}_{\mathbb{R}}(2n).$

$xii)$ $\text{aff}_{\mathbb{F}}(n) \triangleq \left\{ \begin{bmatrix} A & b \\ 0 & 0 \end{bmatrix} : A \in \text{gl}_{\mathbb{F}}(n), b \in \mathbb{F}^n \right\}.$

$xiii)$ $\text{se}_{\mathbb{C}}(n) \triangleq \left\{ \begin{bmatrix} A & b \\ 0 & 0 \end{bmatrix} : A \in \text{su}(n), b \in \mathbb{C}^n \right\}.$

$xiv)$ $\text{se}_{\mathbb{R}}(n) \triangleq \left\{ \begin{bmatrix} A & b \\ 0 & 0 \end{bmatrix} : A \in \text{so}(n), b \in \mathbb{R}^n \right\}.$

$xv)$ $\text{trans}_{\mathbb{F}}(n) \triangleq \left\{ \begin{bmatrix} 0 & b \\ 0 & 0 \end{bmatrix} : b \in \mathbb{F}^n \right\}.$

Definition 3.3.3. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$. Then, $\mathcal{S}$ is a *group* if the following conditions are satisfied:

$i)$ If $A \in \mathcal{S}$, then A is nonsingular.

$ii)$ If $A \in \mathcal{S}$, then $A^{-1} \in \mathcal{S}$.

$iii)$ If $A, B \in \mathcal{S}$, then $AB \in \mathcal{S}$.

$\mathcal{S}$ is an *Abelian group* if $\mathcal{S}$ is a group and the following condition is also satisfied:

$iv)$ For all $A, B \in \mathcal{S}$, $[A, B] = 0$.

Finally, $\mathcal{S}$ is a *finite group* if $\mathcal{S}$ is a group and has a finite number of elements.

Definition 3.3.4. Let $\mathcal{S}_1 \subset \mathbb{F}^{n_1 \times n_1}$ and $\mathcal{S}_2 \subset \mathbb{F}^{n_1 \times n_1}$ be groups. Then, $\mathcal{S}_1$ and $\mathcal{S}_2$ are *isomorphic* if there exists a one-to-one and onto function $\phi: \mathcal{S}_1 \mapsto \mathcal{S}_2$ such that, for all $A, B \in \mathcal{S}_1$, $\phi(AB) = \phi(A)\phi(B)$. In this case, $\mathcal{S}_1 \approx \mathcal{S}_2$, and ϕ is an *isomorphism*.

Proposition 3.3.5. Let $\mathcal{S}_1 \subset \mathbb{F}^{n_1 \times n_1}$ and $\mathcal{S}_2 \subset \mathbb{F}^{n_1 \times n_1}$ be groups, and assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are isomorphic with isomorphism $\phi: \mathcal{S}_1 \mapsto \mathcal{S}_2$. Then, $\phi(I_{n_1}) = I_{n_2}$, and, for all $A \in \mathcal{S}_1$, $\phi(A^{-1}) = [\phi(A)]^{-1}$.

Note that, if $\mathcal{S} \subset \mathbb{F}^{n \times n}$ is a group, then $I_n \in \mathcal{S}$.

The following result lists classical groups that arise in physics and engineering. For example, $O(1, 3)$ is the *Lorentz group* [1162, p. 16], [1186, p. 126]. The special orthogonal group $SO(n)$ consists of the orthogonal matrices whose determinant is 1. In particular, each matrix in $SO(2)$ and $SO(3)$ is a *rotation matrix*.

Proposition 3.3.6. The following sets are groups:

$i)$ $\text{GL}_{\mathbb{F}}(n) \triangleq \{A \in \mathbb{F}^{n \times n}: \det A \neq 0\}.$

$ii)$ $\text{PL}_{\mathbb{F}}(n) \triangleq \{A \in \mathbb{F}^{n \times n}: \det A > 0\}.$

$iii)$ $\text{SL}_{\mathbb{F}}(n) \triangleq \{A \in \mathbb{F}^{n \times n}: \det A = 1\}.$

- iv) $U(n) \triangleq \{A \in \mathbb{C}^{n \times n}: A \text{ is unitary}\}.$
- v) $O(n) \triangleq \{A \in \mathbb{R}^{n \times n}: A \text{ is orthogonal}\}.$
- vi) $SU(n) \triangleq \{A \in U(n): \det A = 1\}.$
- vii) $SO(n) \triangleq \{A \in O(n): \det A = 1\}.$
- viii) $U(n, m) \triangleq \{A \in \mathbb{C}^{(n+m) \times (n+m)}: A^* \text{diag}(I_n, -I_m) A = \text{diag}(I_n, -I_m)\}.$
- ix) $O(n, m) \triangleq \{A \in \mathbb{R}^{(n+m) \times (n+m)}: A^T \text{diag}(I_n, -I_m) A = \text{diag}(I_n, -I_m)\}.$
- x) $SU(n, m) \triangleq \{A \in U(n, m): \det A = 1\}.$
- xi) $SO(n, m) \triangleq \{A \in O(n, m): \det A = 1\}.$
- xii) $\text{Symp}_{\mathbb{F}}(2n) \triangleq \{A \in \mathbb{F}^{2n \times 2n}: A \text{ is symplectic}\}.$
- xiii) $\text{OSymp}_{\mathbb{C}}(2n) \triangleq U(2n) \cap \text{Symp}_{\mathbb{C}}(2n).$
- xiv) $\text{OSymp}_{\mathbb{R}}(2n) \triangleq O(2n) \cap \text{Symp}_{\mathbb{R}}(2n).$
- xv) $\text{Aff}_{\mathbb{F}}(n) \triangleq \left\{ \begin{bmatrix} A & b \\ 0 & 1 \end{bmatrix} : A \in \text{GL}_{\mathbb{F}}(n), b \in \mathbb{F}^n \right\}.$
- xvi) $\text{SE}_{\mathbb{C}}(n) \triangleq \left\{ \begin{bmatrix} A & b \\ 0 & 1 \end{bmatrix} : A \in SU(n), b \in \mathbb{C}^n \right\}.$
- xvii) $\text{SE}_{\mathbb{R}}(n) \triangleq \left\{ \begin{bmatrix} A & b \\ 0 & 1 \end{bmatrix} : A \in SO(n), b \in \mathbb{R}^n \right\}.$
- xviii) $\text{Trans}_{\mathbb{F}}(n) \triangleq \left\{ \begin{bmatrix} I & b \\ 0 & 1 \end{bmatrix} : b \in \mathbb{F}^n \right\}.$

3.4 Matrix Transformations

The following results use groups to define equivalence relations.

Proposition 3.4.1. Let $\mathcal{S}_1 \subset \mathbb{F}^{n \times n}$ and $\mathcal{S}_2 \subset \mathbb{F}^{m \times m}$ be groups, and let $\mathcal{M} \subseteq \mathbb{F}^{n \times m}$. Then, the subset of $\mathcal{M} \times \mathcal{M}$ defined by

$$\mathcal{R} \triangleq \{(A, B) \in \mathcal{M} \times \mathcal{M}: \\ \text{there exist } S_1 \in \mathcal{S}_1 \text{ and } S_2 \in \mathcal{S}_2 \text{ such that } A = S_1 B S_2\}$$

is an equivalence relation on $\mathcal{M}$.

Proposition 3.4.2. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$ be a group, and let $\mathcal{M} \subseteq \mathbb{F}^{n \times n}$. Then, the following subsets of $\mathcal{M} \times \mathcal{M}$ are equivalence relations:

- i) $\mathcal{R} \triangleq \{(A, B) \in \mathcal{M} \times \mathcal{M}: \text{there exists } S \in \mathcal{S} \text{ such that } A = S B S^{-1}\}.$
- ii) $\mathcal{R} \triangleq \{(A, B) \in \mathcal{M} \times \mathcal{M}: \text{there exists } S \in \mathcal{S} \text{ such that } A = S B S^*\}.$
- iii) $\mathcal{R} \triangleq \{(A, B) \in \mathcal{M} \times \mathcal{M}: \text{there exists } S \in \mathcal{S} \text{ such that } A = S B S^T\}.$

If, in addition, $\mathcal{S}$ is an Abelian group, then the following subset $\mathcal{M} \times \mathcal{M}$ is an

equivalence relation:

$$iv) \mathcal{R} \triangleq \{(A, B) \in \mathcal{M} \times \mathcal{M} : \text{there exists } S \in \mathcal{S} \text{ such that } A = SBS\}.$$

Various transformations can be employed for analyzing matrices. Propositions 3.4.1 and 3.4.2 imply that these transformations define equivalence relations.

Definition 3.4.3. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following terminology is defined:

- i) A and B are *left equivalent* if there exists a nonsingular matrix $S_1 \in \mathbb{F}^{n \times n}$ such that $A = S_1 B$.
- ii) A and B are *right equivalent* if there exists a nonsingular matrix $S_2 \in \mathbb{F}^{m \times m}$ such that $A = B S_2$.
- iii) A and B are *biequivalent* if there exist nonsingular matrices $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$ such that $A = S_1 B S_2$.
- iv) A and B are *unitarily left equivalent* if there exists a unitary matrix $S_1 \in \mathbb{F}^{n \times n}$ such that $A = S_1 B$.
- v) A and B are *unitarily right equivalent* if there exists a unitary matrix $S_2 \in \mathbb{F}^{m \times m}$ such that $A = B S_2$.
- vi) A and B are *unitarily biequivalent* if there exist unitary matrices $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$ such that $A = S_1 B S_2$.

Definition 3.4.4. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following terminology is defined:

- i) A and B are *similar* if there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A = S B S^{-1}$.
- ii) A and B are *congruent* if there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A = S B S^*$.
- iii) A and B are *T-congruent* if there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A = S B S^T$.
- iv) A and B are *unitarily similar* if there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that $A = S B S^* = S B S^{-1}$.

The transformations that appear in Definition 3.4.3 and Definition 3.4.4 are called *left equivalence*, *right equivalence*, *biequivalence*, *unitarily left equivalence*, *unitarily right equivalence*, *unitarily biequivalence*, *similarity*, *congruence*, *T-congruence*, and *unitarily similarity* transformations, respectively. The following results summarize some matrix properties that are preserved under left equivalence, right equivalence, biequivalence, similarity, congruence, and unitary similarity.

Proposition 3.4.5. Let $A, B \in \mathbb{F}^{n \times n}$. If A and B are similar, then the following statements hold:

- i) A and B are biequivalent.
- ii) $\text{tr } A = \text{tr } B$.

- iii) $\det A = \det B$.
- iv) A^k and B^k are similar for all $k \geq 1$.
- v) A^{k*} and B^{k*} are similar for all $k \geq 1$.
- vi) A is nonsingular if and only if B is; in this case, A^{-k} and B^{-k} are similar for all $k \geq 1$.
- vii) A is (group invertible, involutory, skew involutory, idempotent, tripotent, nilpotent) if and only if B is.

If A and B are congruent, then the following statements hold:

- viii) A and B are biequivalent.
- ix) A^* and B^* are congruent.
- x) A is nonsingular if and only if B is; in this case, A^{-1} and B^{-1} are congruent.
- xi) A is (range Hermitian, Hermitian, skew Hermitian, positive semidefinite, positive definite) if and only if B is.

If A and B are unitarily similar, then the following statements hold:

- xii) A and B are similar.
- xiii) A and B are congruent.
- xiv) A is (range Hermitian, group invertible, normal, Hermitian, skew Hermitian, positive semidefinite, positive definite, unitary, involutory, skew involutory, idempotent, tripotent, nilpotent) if and only if B is.

3.5 Projectors, Idempotent Matrices, and Subspaces

The following result shows that a unique projector can be associated with each subspace.

Proposition 3.5.1. Let $\mathcal{S} \subseteq \mathbb{F}^n$ be a subspace. Then, there exists a unique projector $A \in \mathbb{F}^{n \times n}$ such that $\mathcal{S} = \mathcal{R}(A)$. Furthermore, $x \in \mathcal{S}$ if and only if $x = Ax$.

Proof. See [998, p. 386] and Fact 3.13.15. □

For a subspace $\mathcal{S} \subseteq \mathbb{F}^n$, the matrix $A \in \mathbb{F}^{n \times n}$ given by Proposition 3.5.1 is the *projector onto* $\mathcal{S}$.

Let $A \in \mathbb{F}^{n \times n}$ be a projector. Then, the *complementary projector* $A_\perp$ is the projector defined by

$$A_\perp \triangleq I - A. \quad (3.5.1)$$

Proposition 3.5.2. Let $\mathcal{S} \subseteq \mathbb{F}^n$ be a subspace, and let $A \in \mathbb{F}^{n \times n}$ be the projector onto $\mathcal{S}$. Then, $A_\perp$ is the projector onto $\mathcal{S}^\perp$. Furthermore,

$$\mathcal{R}(A)^\perp = \mathcal{N}(A) = \mathcal{R}(A_\perp) = \mathcal{S}^\perp. \quad (3.5.2)$$

The following result shows that a unique idempotent matrix can be associated with each pair of complementary subspaces.

Proposition 3.5.3. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be complementary subspaces. Then, there exists a unique idempotent matrix $A \in \mathbb{F}^{n \times n}$ such that $\mathcal{R}(A) = \mathcal{S}_1$ and $\mathcal{N}(A) = \mathcal{S}_2$.

Proof. See [182, p. 118] or [998, p. 386]. $\square$

For complementary subspaces $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$, the unique idempotent matrix $A \in \mathbb{F}^{n \times n}$ given by Proposition 3.5.3 is the *idempotent matrix onto $\mathcal{S}_1 = \mathcal{R}(A)$ along $\mathcal{S}_2 = \mathcal{N}(A)$* .

For an idempotent matrix $A \in \mathbb{F}^{n \times n}$, the *complementary idempotent matrix $A_\perp$* defined by (3.5.1) is also idempotent.

Proposition 3.5.4. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be complementary subspaces, and let $A \in \mathbb{F}^{n \times n}$ be the idempotent matrix onto $\mathcal{S}_1 = \mathcal{R}(A)$ along $\mathcal{S}_2 = \mathcal{N}(A)$. Then, $\mathcal{R}(A_\perp) = \mathcal{S}_2$ and $\mathcal{N}(A_\perp) = \mathcal{S}_1$, that is, $A_\perp$ is the idempotent matrix onto $\mathcal{S}_2$ along $\mathcal{S}_1$.

Definition 3.5.5. The *index of A* , denoted by $\text{ind } A$, is the smallest nonnegative integer k such that

$$\mathcal{R}(A^k) = \mathcal{R}(A^{k+1}). \quad (3.5.3)$$

Proposition 3.5.6. Let $A \in \mathbb{F}^{n \times n}$. Then, A is nonsingular if and only if $\text{ind } A = 0$. Furthermore, A is group invertible if and only if $\text{ind } A \leq 1$.

Note that $\text{ind } 0_{n \times n} = 1$.

Proposition 3.5.7. Let $A \in \mathbb{F}^{n \times n}$, and let $k \geq 1$. Then, $\text{ind } A \leq k$ if and only if $\mathcal{R}(A^k)$ and $\mathcal{N}(A^k)$ are complementary subspaces.

Fact 3.6.3 states that the null space and range of a range-Hermitian matrix are orthogonally complementary subspaces. Furthermore, Proposition 3.1.6 states that every range-Hermitian matrix is group invertible. Hence, the null space and range of a group-invertible matrix are complementary subspaces. The following corollary of Proposition 3.5.7 shows that the converse is true. Note that every idempotent matrix is group invertible.

Corollary 3.5.8. Let $A \in \mathbb{F}^{n \times n}$. Then, A is group invertible if and only if $\mathcal{R}(A)$ and $\mathcal{N}(A)$ are complementary subspaces.

For a group-invertible matrix $A \in \mathbb{F}^{n \times n}$, the following result shows how to construct the idempotent matrix onto $\mathcal{R}(A)$ along $\mathcal{N}(A)$.

Proposition 3.5.9. Let $A \in \mathbb{F}^{n \times n}$, and let $r \triangleq \text{rank } A$. Then, A is group invertible if and only if there exist matrices $B \in \mathbb{F}^{n \times r}$ and $C \in \mathbb{F}^{r \times n}$ such that $A =$

BC and $\text{rank } B = \text{rank } C = r$. In this case, the idempotent matrix $P \triangleq B(CB)^{-1}C$ is the idempotent matrix onto $\mathcal{R}(A)$ along $\mathcal{N}(A)$.

Proof. See [998, p. 634]. □

An alternative expression for the idempotent matrix onto $\mathcal{R}(A)$ along $\mathcal{N}(A)$ is given by Proposition 6.2.3.

3.6 Facts on Group-Invertible and Range-Hermitian Matrices

Fact 3.6.1. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is group invertible.
- ii) A^* is group invertible.
- iii) A^T is group invertible.
- iv) $\overline{A}$ is group invertible.
- v) $\mathcal{R}(A) = \mathcal{R}(A^2)$.
- vi) $\mathcal{N}(A) = \mathcal{N}(A^2)$.
- vii) $\mathcal{N}(A) \cap \mathcal{R}(A) = \{0\}$.
- viii) $\mathcal{N}(A) + \mathcal{R}(A) = \mathbb{F}^n$.
- ix) A and A^2 are left equivalent.
- x) A and A^2 are right equivalent.
- xi) $\text{ind } A \leq 1$.
- xii) $\text{rank } A = \text{rank } A^2$.
- xiii) $\text{def } A = \text{def } A^2$.
- xiv) $\text{def } A = \text{amult}_A(0)$.

(Remark: See Corollary 3.5.8, Proposition 3.5.9, and Corollary 5.5.9.)

Fact 3.6.2. Let $A \in \mathbb{F}^{n \times n}$. Then, $\text{ind } A \leq k$ if and only if A^k is group invertible.

Fact 3.6.3. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is range Hermitian.
- ii) A^* is range Hermitian.
- iii) $\mathcal{R}(A) = \mathcal{R}(A^*)$.
- iv) $\mathcal{R}(A) \subseteq \mathcal{R}(A^*)$.
- v) $\mathcal{R}(A^*) \subseteq \mathcal{R}(A)$.
- vi) $\mathcal{N}(A) = \mathcal{N}(A^*)$.

vii) A and A^* are right equivalent.

viii) $\mathcal{R}(A)^\perp = \mathcal{N}(A)$.

ix) $\mathcal{N}(A)^\perp = \mathcal{R}(A)$.

x) $\mathcal{R}(A)$ and $\mathcal{N}(A)$ are orthogonally complementary subspaces.

xi) $\text{rank } A = \text{rank} \begin{bmatrix} A & A^* \end{bmatrix}$.

(Proof: See [323, 1277].) (Remark: Using Fact 3.13.15, Proposition 3.5.2, and Proposition 6.1.6, vi) is equivalent to $A^+A = I - (I - A^+A) = AA^+$. See Fact 6.3.9, Fact 6.3.10, and Fact 6.3.11.)

Fact 3.6.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that $A^2 = A^*$. Then, A is range Hermitian. (Proof: See [114].) (Remark: A is a *generalized projector*.)

Fact 3.6.5. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are range Hermitian. Then,

$$\text{rank } AB = \text{rank } BA.$$

(Proof: See [122].)

3.7 Facts on Normal, Hermitian, and Skew-Hermitian Matrices

Fact 3.7.1. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and assume that A is (normal, Hermitian, skew Hermitian, unitary). Then, so is A^{-1} .

Fact 3.7.2. Let $A \in \mathbb{F}^{n \times m}$. Then, $AA^T \in \mathbb{F}^{n \times n}$ and $A^TA \in \mathbb{F}^{m \times m}$ are symmetric.

Fact 3.7.3. Let $\alpha \in \mathbb{R}$ and $A \in \mathbb{R}^{n \times n}$. Then, the matrix equation $\alpha A + A^T = 0$ has a nonzero solution A if and only if $\alpha = 1$ or $\alpha = -1$.

Fact 3.7.4. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and let $k \geq 1$. Then, $\mathcal{R}(A) = \mathcal{R}(A^k)$ and $\mathcal{N}(A) = \mathcal{N}(A^k)$.

Fact 3.7.5. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) $x^TAx = 0$ for all $x \in \mathbb{R}^n$ if and only if A is skew symmetric.
- ii) A is symmetric and $x^TAx = 0$ for all $x \in \mathbb{R}^n$ if and only if $A = 0$.

Fact 3.7.6. Let $A \in \mathbb{C}^{n \times n}$. Then, the following statements hold:

- i) x^*Ax is real for all $x \in \mathbb{C}^n$ if and only if A is Hermitian.
- ii) x^*Ax is imaginary for all $x \in \mathbb{C}^n$ if and only if A is skew Hermitian.
- iii) $x^*Ax = 0$ for all $x \in \mathbb{C}^n$ if and only if $A = 0$.

Fact 3.7.7. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

- i) $x^*Ax > 0$ for all nonzero $x \in \mathbb{C}^n$.
- ii) $x^TAx > 0$ for all nonzero $x \in \mathbb{R}^n$.

Fact 3.7.8. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is block diagonal. Then, A is (normal, Hermitian, skew Hermitian) if and only if every diagonally located block has the same property.

Fact 3.7.9. Let $A \in \mathbb{C}^{n \times n}$. Then, the following statements hold:

- i) A is Hermitian if and only if jA is skew Hermitian.
- ii) A is skew Hermitian if and only if jA is Hermitian.
- iii) A is Hermitian if and only if $\operatorname{Re} A$ is symmetric and $\operatorname{Im} A$ is skew symmetric.
- iv) A is skew Hermitian if and only if $\operatorname{Re} A$ is skew symmetric and $\operatorname{Im} A$ is symmetric.
- v) A is positive semidefinite if and only if $\operatorname{Re} A$ is positive semidefinite.
- vi) A is positive definite if and only if $\operatorname{Re} A$ is positive definite.
- vii) A is symmetric if and only if $\begin{bmatrix} 0 & A \\ A & 0 \end{bmatrix}$ is symmetric.
- viii) A is Hermitian if and only if $\begin{bmatrix} 0 & A \\ A & 0 \end{bmatrix}$ is Hermitian.
- ix) A is symmetric if and only if $\begin{bmatrix} 0 & A \\ -A & 0 \end{bmatrix}$ is skew symmetric.
- x) A is Hermitian if and only if $\begin{bmatrix} 0 & A \\ -A & 0 \end{bmatrix}$ is skew Hermitian.

(Remark: x) is a real analogue of i) since $\begin{bmatrix} 0 & A \\ -A & 0 \end{bmatrix} = I_2 \otimes A$, and I_2 is a real representation of j .)

Fact 3.7.10. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) If A is (normal, unitary, Hermitian, positive semidefinite, positive definite), then so is A^A .
- ii) If A is skew Hermitian and n is odd, then A^A is Hermitian.
- iii) If A is skew Hermitian and n is even, then A^A is skew Hermitian.
- iv) If A is diagonal, then so is A^A , and, for all $i = 1, \dots, n$,

$$(A^A)_{(i,i)} = \prod_{\substack{j=1 \\ j \neq i}}^n A_{(j,j)}.$$

(Proof: Use Fact 2.16.10.) (Remark: See Fact 5.14.5.)

Fact 3.7.11. Let $A \in \mathbb{F}^{n \times n}$, assume that n is even, let $x \in \mathbb{F}^n$, and let $\alpha \in \mathbb{F}$. Then,

$$\det(A + \alpha xx^*) = \det A.$$

(Proof: Use Fact 2.16.3 and Fact 3.7.10.)

Fact 3.7.12. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is normal.
- ii) $A^2 A^* = A A^* A$.
- iii) $A A^* A = A^* A^2$.
- iv) $\operatorname{tr} (A A^*)^2 = \operatorname{tr} A^2 A^{2*}$.
- v) There exists $k \geq 1$ such that

$$\operatorname{tr} (A A^*)^k = \operatorname{tr} A^k A^{k*}.$$

- vi) There exist $k, l \in \mathbb{P}$ such that

$$\operatorname{tr} (A A^*)^{kl} = \operatorname{tr} (A^k A^{k*})^l.$$

- vii) A is range Hermitian, and $A A^* A^2 = A^2 A^* A$.

- viii) $A A^* - A^* A$ is positive semidefinite.

$$ix) [A, A^* A] = 0.$$

$$x) [A, [A, A^*]] = 0.$$

(Proof: See [115, 323, 452, 454, 589, 1208].) (Remark: See Fact 3.11.4, Fact 5.14.15, Fact 5.15.4, Fact 6.3.16, Fact 6.6.10, Fact 8.9.27, Fact 8.12.5, Fact 8.17.5, Fact 11.15.4, and Fact 11.16.14.)

Fact 3.7.13. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is Hermitian.
- ii) $A^2 = A^* A$.
- iii) $A^2 = A A^*$.
- iv) $A^{*2} = A^* A$.
- v) $A^{*2} = A A^*$.
- vi) There exists $\alpha \in \mathbb{F}$ such that $A^2 = \alpha A^* A + (1 - \alpha) A A^*$.
- vii) There exists $\alpha \in \mathbb{F}$ such that $A^{*2} = \alpha A^* A + (1 - \alpha) A A^*$.
- viii) $\operatorname{tr} A^2 = \operatorname{tr} A^* A$.
- ix) $\operatorname{tr} A^2 = \operatorname{tr} A A^*$.
- x) $\operatorname{tr} A^{*2} = \operatorname{tr} A^* A$.
- xi) $\operatorname{tr} A^{*2} = \operatorname{tr} A A^*$.

If, in addition, $\mathbb{F} = \mathbb{R}$, then the following condition is equivalent to i)–xi):

- xii) There exist $\alpha, \beta \in \mathbb{R}$ such that

$$\alpha A^2 + (1 - \alpha) A^{T^2} = \beta A^T A + (1 - \beta) A A^T.$$

(Proof: To prove that viii) implies i), use the Schur decomposition Theorem 5.4.1 to replace A with $D + S$, where D is diagonal and S is strictly upper triangular. Then, $\operatorname{tr} D^* D + \operatorname{tr} S^* S = \operatorname{tr} D^2 \leq \operatorname{tr} D^* D$. Hence, $S = 0$, and thus $\operatorname{tr} D^* D = \operatorname{tr} D^2$,

which implies that D is real. See [115, 856].) (Remark: See Fact 3.13.1.) (Remark: Fact 9.11.3 states that, for all $A \in \mathbb{F}^{n \times n}$, $|\operatorname{tr} A^2| \leq \operatorname{tr} A^*A$.)

Fact 3.7.14. Let $A \in \mathbb{F}^{n \times n}$, let $\alpha, \beta \in \mathbb{F}$, and assume that $\alpha \neq 0$. Then, the following statements are equivalent:

- i) A is normal.
- ii) $\alpha A + \beta I$ is normal.

Now, assume, in addition, that $\alpha, \beta \in \mathbb{R}$. Then, the following statements are equivalent:

- iii) A is Hermitian.
- iv) $\alpha A + \beta I$ is Hermitian.

(Remark: The function $f(A) = \alpha A + \beta I$ is an *affine mapping*.)

Fact 3.7.15. Let $A \in \mathbb{R}^{n \times n}$, assume that A is skew symmetric, and let $\alpha > 0$. Then, $-A^2$ is positive semidefinite, $\det A \geq 0$, and $\det(\alpha I + A) > 0$. If, in addition, n is odd, then $\det A = 0$.

Fact 3.7.16. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is skew Hermitian. If n is even, then $\det A \geq 0$. If n is odd, then $\det A$ is imaginary. (Proof: The first statement follows from Proposition 5.5.21.)

Fact 3.7.17. Let $x, y \in \mathbb{F}^n$, and define

$$A \triangleq \begin{bmatrix} x & y \end{bmatrix}.$$

Then,

$$xy^* - yx^* = AJ_2A^*.$$

Furthermore, $xy^* - yx^*$ is skew Hermitian and has rank 0 or 2.

Fact 3.7.18. Let $x, y \in \mathbb{F}^n$. Then, the following statements hold:

- i) xy^T is idempotent if and only if either $xy^T = 0$ or $x^Ty = 1$.
- ii) xy^T is Hermitian if and only if there exists $\alpha \in \mathbb{R}$ such that either $y = \alpha \bar{x}$ or $x = \alpha \bar{y}$.

Fact 3.7.19. Let $x, y \in \mathbb{F}^n$, and define $A \triangleq I - xy^T$. Then, the following statements hold:

- i) $\det A = 1 - x^Ty$.
- ii) A is nonsingular if and only if $x^Ty \neq 1$.
- iii) A is nonsingular if and only if A is elementary.
- iv) $\operatorname{rank} A = n - 1$ if and only if $x^Ty = 1$.
- v) A is Hermitian if and only if there exists $\alpha \in \mathbb{R}$ such that either $y = \alpha \bar{x}$ or $x = \alpha \bar{y}$.
- vi) A is positive semidefinite if and only if A is Hermitian and $x^Ty \leq 1$.

- vii) A is positive definite if and only if A is Hermitian and $x^T y < 1$.
- viii) A is idempotent if and only if either $xy^T = 0$ or $x^T y = 1$.
- ix) A is orthogonal if and only if either $x = 0$ or $y = \frac{1}{2}y^T y x$.
- x) A is involutory if and only if $x^T y = 2$.
- xi) A is a projector if and only if either $y = 0$ or $x = x^* x y$.
- xii) A is a reflector if and only if either $y = 0$ or $2x = x^* x y$.
- xiii) A is an elementary projector if and only if $x \neq 0$ and $y = (x^* x)^{-1} x$.
- xiv) A is an elementary reflector if and only if $x \neq 0$ and $y = 2(x^* x)^{-1} x$.

(Remark: See Fact 3.13.9.)

Fact 3.7.20. Let $x, y \in \mathbb{F}^n$ satisfy $x^T y \neq 1$. Then, $I - xy^T$ is nonsingular and

$$(I - xy^T)^{-1} = I - \frac{1}{x^T y - 1} xy^T.$$

(Remark: The inverse of an elementary matrix is an elementary matrix.)

Fact 3.7.21. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, $\det A$ is real.

Fact 3.7.22. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then,

$$(\operatorname{tr} A)^2 \leq (\operatorname{rank} A) \operatorname{tr} A^2.$$

Furthermore, equality holds if and only if there exists $\alpha \in \mathbb{R}$ such that $A^2 = \alpha A$.

(Remark: See Fact 5.11.10 and Fact 9.13.12.)

Fact 3.7.23. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is skew symmetric. Then, $\operatorname{tr} A = 0$. If, in addition, $B \in \mathbb{R}^{n \times n}$ is symmetric, then $\operatorname{tr} AB = 0$.

Fact 3.7.24. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is skew Hermitian. Then, $\operatorname{Re} \operatorname{tr} A = 0$. If, in addition, $B \in \mathbb{F}^{n \times n}$ is Hermitian, then $\operatorname{Re} \operatorname{tr} AB = 0$.

Fact 3.7.25. Let $A \in \mathbb{F}^{n \times m}$. Then, $A^* A$ is positive semidefinite. Furthermore, $A^* A$ is positive definite if and only if A is left invertible. In this case, $A^L \in \mathbb{F}^{m \times n}$ defined by

$$A^L \triangleq (A^* A)^{-1} A^*$$

is a left inverse of A . (Remark: See Fact 2.15.2, Fact 3.7.26, and Fact 3.13.6.)

Fact 3.7.26. Let $A \in \mathbb{F}^{n \times m}$. Then, AA^* is positive semidefinite. Furthermore, AA^* is positive definite if and only if A is right invertible. In this case, $A^R \in \mathbb{F}^{m \times n}$ defined by

$$A^R \triangleq A^*(AA^*)^{-1}$$

is a right inverse of A . (Remark: See Fact 2.15.2, Fact 3.13.6, and Fact 3.7.25.)

Fact 3.7.27. Let $A \in \mathbb{F}^{n \times m}$. Then, $A^* A$, AA^* , and $\begin{bmatrix} 0 & A^* \\ A & 0 \end{bmatrix}$ are Hermitian, and $\begin{bmatrix} 0 & A^* \\ -A & 0 \end{bmatrix}$ is skew Hermitian. Now, assume that $n = m$. Then, $A + A^*$, $j(A - A^*)$,

and $\frac{1}{2j}(A - A^*)$ are Hermitian, while $A - A^*$ is skew Hermitian. Finally,

$$A = \frac{1}{2}(A + A^*) + \frac{1}{2j}(A - A^*)$$

and

$$A = \frac{1}{2}(A + A^*) + j[\frac{1}{2j}(A - A^*)].$$

(Remark: The last two identities are Cartesian decompositions.)

Fact 3.7.28. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist a unique Hermitian matrix $B \in \mathbb{F}^{n \times n}$ and a unique skew-Hermitian matrix $C \in \mathbb{F}^{n \times n}$ such that $A = B + C$. Specifically, if $A = \hat{B} + j\hat{C}$, where $\hat{B}, \hat{C} \in \mathbb{R}^{n \times n}$, then $\hat{B}$ and $\hat{C}$ are given by

$$B = \frac{1}{2}(A + A^*) = \frac{1}{2}(\hat{B} + \hat{B}^T) + j\frac{1}{2}(\hat{C} - \hat{C}^T)$$

and

$$C = \frac{1}{2j}(A - A^*) = \frac{1}{2}(\hat{B} - \hat{B}^T) + j\frac{1}{2}(\hat{C} + \hat{C}^T).$$

Furthermore, A is normal if and only if $BC = CB$. (Remark: See Fact 11.13.9.)

Fact 3.7.29. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist unique Hermitian matrices $B, C \in \mathbb{C}^{n \times n}$ such that $A = B + jC$. Specifically, if $A = \hat{B} + j\hat{C}$, where $\hat{B}, \hat{C} \in \mathbb{R}^{n \times n}$, then $\hat{B}$ and $\hat{C}$ are given by

$$B = \frac{1}{2}(A + A^*) = \frac{1}{2}(\hat{B} + \hat{B}^T) + j\frac{1}{2}(\hat{C} - \hat{C}^T)$$

and

$$C = \frac{1}{2j}(A - A^*) = \frac{1}{2}(\hat{C} + \hat{C}^T) - j\frac{1}{2}(\hat{B} - \hat{B}^T).$$

Furthermore, A is normal if and only if $BC = CB$. (Remark: This result is the *Cartesian decomposition*.)

Fact 3.7.30. Let $A, B \in \mathbb{C}^{n \times n}$, assume that A is either Hermitian or skew Hermitian, and assume that B is either Hermitian or skew Hermitian. Then,

$$\text{rank } AB = \text{rank } BA.$$

(Proof: AB and $(AB)^* = BA$ have the same singular values. See Fact 5.11.19.)
(Remark: See Fact 2.10.26.)

Fact 3.7.31. Let $A, B \in \mathbb{R}^{3 \times 3}$, and assume that A and B are skew symmetric. Then,

$$\text{tr } AB^3 = \frac{1}{2}(\text{tr } AB)(\text{tr } B^2)$$

and

$$\text{tr } A^3B^3 = \frac{1}{4}(\text{tr } A^2)(\text{tr } AB)(\text{tr } B^2) + \frac{1}{3}(\text{tr } A^3)(\text{tr } B^3).$$

(Proof: See [79].)

Fact 3.7.32. Let $A \in \mathbb{F}^{n \times n}$ and $k \geq 1$. If A is (normal, Hermitian, unitary, involutory, positive semidefinite, positive definite, idempotent, nilpotent), then so is A^k . If A is (skew Hermitian, skew involutory), then so is A^{2k+1} . If A is Hermitian, then A^{2k} is positive semidefinite. If A is tripotent, then so is A^{3k} .

Fact 3.7.33. Let $a, b, c, d, e, f \in \mathbb{R}$, and define the skew-symmetric matrix $A \in \mathbb{R}^{4 \times 4}$ given by

$$A \triangleq \begin{bmatrix} 0 & a & b & c \\ -a & 0 & d & e \\ -b & -d & 0 & f \\ -c & -e & -f & 0 \end{bmatrix}.$$

Then,

$$\det A = (af - be + cd)^2.$$

(Proof: See [1184, p. 63].) (Remark: See Fact 4.8.14 and Fact 4.10.2.)

Fact 3.7.34. Let $A \in \mathbb{R}^{2n \times 2n}$, and assume that A is skew symmetric. Then, there exists a nonsingular matrix $S \in \mathbb{R}^{2n \times 2n}$ such that $S^T A S = J_{2n}$. (Proof: See [103, p. 231].)

Fact 3.7.35. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is positive definite. Then,

$$\mathcal{E} \triangleq \{x \in \mathbb{R}^n: x^T A x \leq 1\}$$

is a hyperellipsoid. Furthermore, the volume V of $\mathcal{E}$ is given by

$$V = \frac{\alpha(n)}{\sqrt{\det A}},$$

where

$$\alpha(n) \triangleq \begin{cases} \pi^{n/2}/(n/2)!, & n \text{ even,} \\ 2^n \pi^{(n-1)/2} [(n-1)/2]!/n!, & n \text{ odd.} \end{cases}$$

In particular, the area of the ellipse $\{x \in \mathbb{R}^2: x^T A x \leq 1\}$ is $\pi/\det A$. (Remark: $\alpha(n)$ is the volume of the unit n -dimensional hypersphere.) (Remark: See [801, p. 36].)

3.8 Facts on Commutators

Fact 3.8.1. Let $A, B \in \mathbb{F}^{n \times n}$. If either A and B are Hermitian or A and B are skew Hermitian, then $[A, B]$ is skew Hermitian. Furthermore, if A is Hermitian and B is skew Hermitian, or vice versa, then $[A, B]$ is Hermitian.

Fact 3.8.2. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $\text{tr } A = 0$.
- ii) There exist matrices $B, C \in \mathbb{F}^{n \times n}$ such that B is Hermitian, $\text{tr } C = 0$, and $A = [B, C]$.
- iii) There exist matrices $B, C \in \mathbb{F}^{n \times n}$ such that $A = [B, C]$.

(Proof: See [535] and Fact 5.9.18. If every diagonal entry of A is zero, then let $B \triangleq \text{diag}(1, \dots, n)$, $C_{(i,i)} \triangleq 0$, and, for $i \neq j$, $C_{(i,j)} \triangleq A_{(i,j)}/(i-j)$. See [1487, p. 110]. See also [1098, p. 172].)

Fact 3.8.3. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is Hermitian, and $\text{tr } A = 0$.
- ii) There exists a nonsingular matrix $B \in \mathbb{F}^{n \times n}$ such that $A = [B, B^*]$.
- iii) There exist a Hermitian matrix $B \in \mathbb{F}^{n \times n}$ and a skew-Hermitian matrix $C \in \mathbb{F}^{n \times n}$ such that $A = [B, C]$.
- iv) There exist a skew-Hermitian matrix $B \in \mathbb{F}^{n \times n}$ and a Hermitian matrix $C \in \mathbb{F}^{n \times n}$ such that $A = [B, C]$.

(Proof: See [535] and [1266].)

Fact 3.8.4. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is skew Hermitian, and $\text{tr } A = 0$.
- ii) There exists a nonsingular matrix $B \in \mathbb{F}^{n \times n}$ such that $A = [{}_jB, B^*]$.
- iii) If $A \in \mathbb{C}^{n \times n}$ is skew Hermitian, then there exist Hermitian matrices $B, C \in \mathbb{F}^{n \times n}$ such that $A = [B, C]$.

(Proof: See [535] or use Fact 3.8.3.)

Fact 3.8.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is skew symmetric. Then, there exist symmetric matrices $B, C \in \mathbb{F}^{n \times n}$ such that $A = [B, C]$. (Proof: Use Fact 5.15.24. See [1098, pp. 83, 89].) (Remark: “Symmetric” is correct for $\mathbb{F} = \mathbb{C}$.)

Fact 3.8.6. Let $A \in \mathbb{F}^{n \times n}$, and assume that $[A, [A, A^*]] = 0$. Then, A is normal. (Remark: See [1487, p. 32].)

Fact 3.8.7. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist $B, C \in \mathbb{F}^{n \times n}$ such that B is normal, C is Hermitian, and

$$A = B + [C, B].$$

(Remark: See [440].)

3.9 Facts on Linear Interpolation

Fact 3.9.1. Let $y \in \mathbb{F}^n$ and $x \in \mathbb{F}^m$. Then, there exists a matrix $A \in \mathbb{F}^{n \times m}$ such that $y = Ax$ if and only if either $y = 0$ or $x \neq 0$. If $y = 0$, then one such matrix is $A = 0$. If $x \neq 0$, then one such matrix is

$$A = (x^*x)^{-1}yx^*.$$

(Remark: This is a linear interpolation problem. See [773].)

Fact 3.9.2. Let $x, y \in \mathbb{F}^n$, and assume that $x \neq 0$. Then, there exists a Hermitian matrix $A \in \mathbb{F}^{n \times n}$ such that $y = Ax$ if and only if x^*y is real. One such matrix is

$$A = (x^*x)^{-1}[yx^* + xy^* - x^*yI].$$

Now, assume that x and y are real. Then,

$$\sigma_{\max}(A) = \frac{\|x\|_2}{\|y\|_2} = \min\{\sigma_{\max}(B): B \in \mathbb{R}^{n \times n} \text{ is symmetric and } y = Bx\}.$$

(Proof: The last statement is given in [1205].)

Fact 3.9.3. Let $x, y \in \mathbb{F}^n$, and assume that $x \neq 0$. Then, there exists a positive-definite matrix $A \in \mathbb{F}^{n \times n}$ such that $y = Ax$ if and only if x^*y is real and positive. One such matrix is

$$A = I + (x^*y)^{-1}yy^* - (x^*x)^{-1}xx^*.$$

(Proof: To show that A is positive definite, note that the elementary projector $I - (x^*x)^{-1}xx^*$ is positive semidefinite and $\text{rank}[I - (x^*x)^{-1}xx^*] = n - 1$. Since $(x^*y)^{-1}yy^*$ is positive semidefinite, it follows that $\mathcal{N}(A) \subseteq \mathcal{N}[I - (x^*x)^{-1}xx^*]$. Next, since $x^*y > 0$, it follows that $y^*x \neq 0$ and $y \neq 0$, and thus $x \notin \mathcal{N}(A)$. Consequently, $\mathcal{N}(A) \subset \mathcal{N}[I - (x^*x)^{-1}xx^*]$ (note proper inclusion), and thus $\text{def } A < 1$. Hence, A is nonsingular.)

Fact 3.9.4. Let $x, y \in \mathbb{F}^n$. Then, there exists a skew-Hermitian matrix $A \in \mathbb{F}^{n \times n}$ such that $y = Ax$ if and only if either $y = 0$ or $x \neq 0$ and $x^*y = 0$. If $x \neq 0$ and $x^*y = 0$, then one such matrix is

$$A = (x^*x)^{-1}(yx^* - xy^*).$$

(Proof: See [924].)

Fact 3.9.5. Let $x, y \in \mathbb{R}^n$. Then, there exists an orthogonal matrix $A \in \mathbb{R}^{n \times n}$ such that $Ax = y$ if and only if $x^T x = y^T y$. (Remark: One such matrix is given by a product of n plane rotations given by Fact 5.15.16. Another matrix is given by the product of elementary reflectors given by Fact 5.15.15. For $n = 3$, one such matrix is given by Fact 3.11.8, while another is given by the exponential of a skew-symmetric matrix given by Fact 11.11.7. See Fact 3.14.4.) (Problem: Extend this result to $\mathbb{C}^n$.) (Remark: See Fact 9.15.6.)

3.10 Facts on the Cross Product

Fact 3.10.1. Let $x, y, z, w \in \mathbb{R}^3$, and define the cross-product matrix $K(x) \in \mathbb{R}^{3 \times 3}$ by

$$K(x) \triangleq \begin{bmatrix} 0 & -x_{(3)} & x_{(2)} \\ x_{(3)} & 0 & -x_{(1)} \\ -x_{(2)} & x_{(1)} & 0 \end{bmatrix}.$$

Then, the following statements hold:

- i) $x \times x = K(x)x = 0$.
- ii) $x^T K(x) = 0$.
- iii) $K^T(x) = -K(x)$.
- iv) $K^2(x) = xx^T - (x^T x)I$.

$$v) \operatorname{tr} K^T(x)K(x) = -\operatorname{tr} K^2(x) = 2x^T x.$$

$$vi) K^3(x) = -(x^T x)K(x).$$

$$vii) [I - K(x)]^{-1} = I + (1 + x^T x)^{-1}[K(x) + K^2(x)].$$

$$viii) [I + \frac{1}{2}K(x)][I - \frac{1}{2}K(x)]^{-1} = I + \frac{4}{4+x^T x}[K(x) + \frac{1}{2}K^2(x)].$$

ix) Define

$$H(x) \triangleq \frac{1}{2}[\frac{1}{2}(1 - x^T x)I + xx^T + K(x)].$$

Then,

$$H(x)H^T(x) = \frac{1}{16}(1 + x^T x)^2 I.$$

$$x) \text{ For all } \alpha, \beta \in \mathbb{R}, K(\alpha x + \beta y) = \alpha K(x) + \beta K(y).$$

$$xi) x \times y = -(y \times x) = K(x)y = -K(y)x = K^T(y)x.$$

$$xii) \text{ If } x \times y \neq 0, \text{ then } \mathcal{N}[(x \times y)^T] = \{x \times y\}^\perp = \mathcal{R}[\begin{bmatrix} x & y \end{bmatrix}].$$

$$xiii) K(x \times y) = K[K(x)y] = [K(x), K(y)].$$

$$xiv) K(x \times y) = yx^T - xy^T = \begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} -y^T \\ x^T \end{bmatrix} = -\begin{bmatrix} x & y \end{bmatrix} J_2 \begin{bmatrix} x & y \end{bmatrix}^T.$$

$$xv) (x \times y) \times x = (x^T x I - xx^T)y.$$

$$xvi) K[(x \times y) \times x] = (x^T x)K(y) - (x^T y)K(x).$$

$$xvii) (x \times y)^T(x \times y) = \det \begin{bmatrix} x & y & x \times y \end{bmatrix}.$$

$$xviii) (x \times y)^T z = x^T(y \times z) = \det \begin{bmatrix} x & y & z \end{bmatrix}.$$

$$xix) x \times (y \times z) = (x^T z)y - (x^T y)z.$$

$$xx) (x \times y) \times z = (x^T z)y - (y^T z)x.$$

$$xxi) K[(x \times y) \times z] = (x^T z)K(y) - (y^T z)K(x).$$

$$xxii) K[x \times (y \times z)] = (x^T z)K(y) - (x^T y)K(z).$$

$$xxiii) (x \times y)^T(x \times y) = x^T x y^T y - (x^T y)^2.$$

$$xxiv) K(x)K(y) = yx^T - x^T y I_3.$$

$$xxv) K(x)K(y)K(x) = -(x^T y)K(x).$$

$$xxvi) K^2(x)K(y) + K(y)K^2(x) = -(x^T x)K(y) - (x^T y)K(x).$$

$$xxvii) K^2(x)K^2(y) - K^2(y)K^2(x) = -(x^T y)K(x \times y).$$

$$xxviii) K(x)K(z)(x^T w y - x^T y w) = K(x)K(w)x^T z y.$$

$$xxix) \sqrt{(x \times y)^T(x \times y)} = \sqrt{x^T x y^T y} \sin \theta, \text{ where } \theta \text{ is the angle between } x \text{ and } y.$$

$$xxx) (x \times y)^T(x \times y) = x^T x y^T y - (x^T y)^2.$$

$$xxxi) 2xx^T K(y) = (x \times y)x^T + x(x \times y)^T + x^T x K(y) - x^T y K(x).$$

$$xxxii) (x \times y)^T(z \times w) = x^T z y^T w - x^T w y^T z = \det \begin{bmatrix} x^T z & x^T w \\ y^T z & y^T w \end{bmatrix}.$$

$$xxxiii) (x \times y) \times (z \times w) = x^T(y \times w)z - x^T(y \times z)w = x^T(z \times w)y - y^T(z \times w)x.$$

$$xxxiv) x \times [y \times (z \times w)] = (y^T w)(x \times z) - (y^T z)(x \times w).$$

$$xxv) \quad x \times [y \times (y \times x)] = y \times [x \times (y \times x)] = (y^T x)(x \times y).$$

xxvi) Let $A \in \mathbb{R}^{3 \times 3}$. Then,

$$A^T K(Ax)A = (\det A)K(x),$$

and thus

$$A^T(Ax \times Ay) = (\det A)(x \times y).$$

xxvii) Let $A \in \mathbb{R}^{3 \times 3}$, and assume that A is orthogonal. Then,

$$K(Ax)A = (\det A)AK(x),$$

and thus

$$Ax \times Ay = (\det A)A(x \times y).$$

xxviii) Let $A \in \mathbb{R}^{3 \times 3}$, and assume that A is orthogonal and $\det A = 1$. Then,

$$K(Ax)A = AK(x),$$

and thus

$$Ax \times Ay = A(x \times y).$$

$$xxix) \quad \begin{bmatrix} x & y & z \end{bmatrix}^A = \begin{bmatrix} y \times z & z \times x & x \times y \end{bmatrix}^T.$$

$$xli) \quad \det \begin{bmatrix} K(x) & y \\ -y^T & 0 \end{bmatrix} = (x^T y)^2.$$

$$xlii) \quad \begin{bmatrix} K(x) & y \\ -y^T & 0 \end{bmatrix}^A = -x^T y \begin{bmatrix} K(y) & x \\ -x^T & 0 \end{bmatrix}.$$

xliii) If $x^T y \neq 0$, then

$$\begin{bmatrix} K(x) & y \\ -y^T & 0 \end{bmatrix}^{-1} = \frac{-1}{x^T y} \begin{bmatrix} K(y) & x \\ -x^T & 0 \end{bmatrix}.$$

xliiii) If $x \neq 0$, then $K^+(x) = (x^T x)^{-1}K(x)$.

xliv) If $x^T y = 0$ and $x^T x + y^T y \neq 0$, then

$$\begin{bmatrix} K(x) & y \\ -y^T & 0 \end{bmatrix}^+ = \frac{-1}{x^T x + y^T y} \begin{bmatrix} K(x) & y \\ -y^T & 0 \end{bmatrix}.$$

(Proof: Results vii), viii), and xrv)–xxvii) are given in [746, p. 363]. Result ix) is given in [1341]. Statement xxviii) is a consequence of a result given in [572, p. 58]. Statement xxx) is equivalent to the fact that $\sin^2 \theta + \cos^2 \theta = 1$. Using xviii),

$$e_i^T A^T (Ax \times Ay) = \det \begin{bmatrix} Ax & Ay & Ae_i \end{bmatrix} = (\det A)e_i^T (x \times y)$$

for all $i = 1, 2, 3$, which proves xxvi). Result xxxix) is given in [1319]. Results xli)–xliv) are proved in [1334]. (Proof: See [410, 474, 746, 1058, 1192, 1262, 1327].) (Remark: Cross products of complex vectors are considered in [599].) (Remark: A cross product can be defined on $\mathbb{R}^7$. See [477, pp. 297–299].) (Remark: An extension of the cross product to higher dimensions is given by the outer product in Clifford algebras. See Fact 9.7.5 and [349, 425, 555, 605, 671, 672, 870, 934].)

(Remark: See Fact 11.11.11.) (Problem: Extend these identities to complex vectors and matrices.)

Fact 3.10.2. Let $A \in \mathbb{R}^{3 \times 3}$, assume that A is orthogonal, let $B \in \mathbb{C}^{3 \times 3}$, and assume that B is symmetric. Then,

$$\sum_{i=1}^3 (Ae_i) \times (BAe_i) = 0.$$

(Proof: For $i = 1, 2, 3$, multiply by $e_i^T A^T$.)

Fact 3.10.3. Let α_1, α_2 , and α_3 be distinct positive numbers, let $A \in \mathbb{R}^{3 \times 3}$, assume that A is orthogonal, and assume that

$$\sum_{i=1}^3 \alpha_i e_i \times Ae_i = 0.$$

Then,

$$A \in \{I, \text{diag}(1, -1, -1), \text{diag}(-1, 1, -1), \text{diag}(-1, -1, 1)\}.$$

(Remark: This result characterizes equilibria for a dynamical system on $\text{SO}(3)$. See [306].)

3.11 Facts on Unitary and Shifted-Unitary Matrices

Fact 3.11.1. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$, assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are subspaces, and assume that $\dim \mathcal{S}_1 \leq \dim \mathcal{S}_2$. Then, there exists a unitary matrix $A \in \mathbb{F}^{n \times n}$ such that $A\mathcal{S}_1 \subseteq \mathcal{S}_2$.

Fact 3.11.2. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$, assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are subspaces, and assume that $\dim \mathcal{S}_1 + \dim \mathcal{S}_2 \leq n$. Then, there exists a unitary matrix $A \in \mathbb{F}^{n \times n}$ such that $A\mathcal{S}_1 \subseteq \mathcal{S}_2^\perp$. (Proof: Use Fact 3.11.1.)

Fact 3.11.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is unitary. Then, the following statements hold:

- i) $A = A^{-*}$.
- ii) $A^T = \overline{A}^{-1} = \overline{A}^*$.
- iii) $\overline{A} = A^{-T} = \overline{A}^{-*}$.
- iv) $A^* = A^{-1}$.

Fact 3.11.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then, the following statements are equivalent:

- i) A is normal.
- ii) $A^{-1}A^*$ is unitary.
- iii) $[A, A^*] = 0$.

$$iv) [A, A^{-*}] = 0.$$

$$v) [A^{-1}, A^{-*}] = 0.$$

(Proof: See [589].) (Remark: See Fact 3.7.12, Fact 5.15.4, Fact 6.3.16, and Fact 6.6.10.)

Fact 3.11.5. Let $A \in \mathbb{F}^{n \times m}$. If A is (left inner, right inner), then A is (left invertible, right invertible) and A^* is a (left inverse, right inverse) of A .

Fact 3.11.6. Let $\theta \in \mathbb{R}$, and define the orthogonal matrix

$$A(\theta) \triangleq \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}.$$

Now, let $\theta_1, \theta_2 \in \mathbb{R}$. Then,

$$A(\theta_1)A(\theta_2) = A(\theta_1 + \theta_2).$$

Consequently,

$$\cos(\theta_1 + \theta_2) = (\cos \theta_1) \cos \theta_2 - (\sin \theta_1) \sin \theta_2,$$

$$\sin(\theta_1 + \theta_2) = (\cos \theta_1) \sin \theta_2 + (\sin \theta_1) \cos \theta_2.$$

Furthermore,

$$\text{SO}(2) = \{A(\theta): \theta \in \mathbb{R}\}$$

and

$$\text{O}(2) \setminus \text{SO}(2) = \left\{ \begin{bmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{bmatrix} : \theta \in \mathbb{R} \right\}.$$

(Remark: See Proposition 3.3.6 and Fact 11.11.3.)

Fact 3.11.7. Let $A \in \text{O}(3) \setminus \text{SO}(3)$. Then, $-A \in \text{SO}(3)$.

Fact 3.11.8. Let $x, y \in \mathbb{R}^3$, assume that $x^T x = y^T y \neq 0$, let $\theta \in (0, \pi)$ denote the angle between x and y , define $z \in \mathbb{R}^3$ by

$$z \triangleq \frac{1}{\|x \times y\|_2} x \times y,$$

and define $A \in \mathbb{R}^{3 \times 3}$ by

$$A \triangleq (\cos \theta)I + (\sin \theta)K(z) + (1 - \cos \theta)zz^T.$$

Then,

$$A = I + (\sin \theta)K(z) + (1 - \cos \theta)K^2(z),$$

$y = Ax$, A is orthogonal, and $\det A = 1$. Furthermore,

$$A = (I - B)(I + B)^{-1},$$

where

$$B \triangleq -\tan\left(\frac{1}{2}\theta\right)K(z).$$

(Proof: The expression for A in terms of B is derived in [11]. The expression involving B is derived in [1008, pp. 244, 245].) (Remark: θ is given by

$$\theta = \cos^{-1} \frac{x^T y}{\|x\|_2 \|y\|_2}.$$

Furthermore,

$$\sin \theta = \frac{\|x \times y\|_2}{\|x\|_2 \|y\|_2}.$$

(Remark: A can be written as

$$\begin{aligned} A &= (\cos \theta)I + \frac{1}{\|x\|_2^2}(yx^T - xy^T) + \frac{1 - \cos \theta}{\|x \times y\|_2^2}(x \times y)(x \times y)^T \\ &= \frac{x^T y}{x^T x}I + \frac{1}{x^T x}(yx^T - xy^T) + \frac{1 - \cos \theta}{(x^T x \sin \theta)^2}(x \times y)(x \times y)^T \\ &= \frac{x^T y}{x^T x}I + \frac{1}{x^T x}(yx^T - xy^T) + \frac{\tan(\frac{1}{2}\theta)}{(x^T x)^2 \sin \theta}(x \times y)(x \times y)^T \\ &= \frac{x^T y}{x^T x}I + \frac{1}{x^T x}(yx^T - xy^T) + \frac{1}{(x^T x)^2(1 + \cos \theta)}(x \times y)(x \times y)^T \\ &= \frac{x^T y}{x^T x}I + \frac{1}{x^T x}(yx^T - xy^T) + \frac{1}{x^T x(x^T x + x^T y)}(x \times y)(x \times y)^T. \end{aligned}$$

As a check, note that

$$\begin{aligned} Ax &= (\cos \theta)x + \frac{1}{\|x\|_2^2}(x^T xy - y^T xx) + \frac{1 - \cos \theta}{\|x \times y\|_2^2}(x \times y)(x \times y)^T x \\ &= \frac{x^T y}{\|x\|_2^2}x + \frac{1}{\|x\|_2^2}(x^T xy - y^T xx) \\ &= y. \end{aligned}$$

Furthermore, B can be written as

$$B = \frac{1}{x^T x + x^T y}(xy^T - yx^T).$$

These expressions satisfy $A + B + AB = I$. (Remark: The matrix A represents a right-hand rule rotation of the nonzero vector x through the angle θ around z to yield the vector y , which has the same length as x . In the cases $x = y$ and $x = -y$, which correspond, respectively, to $\theta = 0$ and $\theta = \pi$, the pivot vector z is not unique. Letting $z \in \mathbb{R}^3$ be arbitrary in these cases yields $A = I$ and $A = -I$, respectively, and thus $y = Ax$ holds in both cases. However, $-I$ has determinant -1 .) (Remark: See Fact 11.11.6.) (Remark: This is a linear interpolation problem. See Fact 3.9.5, Fact 11.11.7, and [135, 773].) (Remark: Extensions of the Cayley transform are discussed in [1342].)

Fact 3.11.9. Let $A \in \mathbb{R}^{3 \times 3}$, and let $z \triangleq \begin{bmatrix} b \\ c \\ d \end{bmatrix}$, where $b^2 + c^2 + d^2 = 1$. Then, $A \in \text{SO}(3)$, and A rotates every vector in $\mathbb{R}^3$ by the angle π about z if and only if

$$A = \begin{bmatrix} 2b^2 - 1 & 2bc & 2bd \\ 2bc & 2c^2 - 1 & 2cd \\ 2bd & 2cd & 2d^2 - 1 \end{bmatrix}.$$

(Proof: This formula follows from the last expression for A in Fact 3.11.10 with $\theta = \pi$. See [357, p. 30].) (Remark: A is a reflector.) (Problem: Solve for b , c , and d in terms of the entries of A .)

Fact 3.11.10. Let $A \in \mathbb{R}^{3 \times 3}$. Then, $A \in \text{SO}(3)$ if and only if there exist real numbers a, b, c, d such that $a^2 + b^2 + c^2 + d^2 = 1$ and

$$A = \begin{bmatrix} a^2 + b^2 - c^2 - d^2 & 2(bc - ad) & 2(ac + bd) \\ 2(ad + bc) & a^2 - b^2 + c^2 - d^2 & 2(cd - ab) \\ 2(bd - ac) & 2(ab + cd) & a^2 - b^2 - c^2 + d^2 \end{bmatrix}.$$

In this case,

$$a = \pm \frac{1}{2} \sqrt{1 + \text{tr } A}.$$

If, in addition, $a \neq 0$, then b , c , and d are given by

$$b = \frac{A_{(3,2)} - A_{(2,3)}}{4a}, \quad c = \frac{A_{(1,3)} - A_{(3,1)}}{4a}, \quad d = \frac{A_{(2,1)} - A_{(1,2)}}{4a}.$$

Now, define $v \triangleq \begin{bmatrix} b & c & d \end{bmatrix}^T$. Then, A represents a rotation about the unit-length vector $z \triangleq (\csc \frac{\theta}{2})v$ through the angle $\theta \in [0, 2\pi]$ that satisfies

$$a = \cos \frac{\theta}{2},$$

where the direction of rotation is determined by the right-hand rule. Therefore,

$$\theta \triangleq 2 \cos^{-1} a.$$

If $a \in [0, 1]$, then

$$\theta = 2 \cos^{-1} \left(\frac{1}{2} \sqrt{1 + \text{tr } A} \right) = \cos^{-1} \left(\frac{1}{2} [(\text{tr } A) - 1] \right),$$

whereas, if $a \in [-1, 0]$, then

$$\theta = 2 \cos^{-1} \left(-\frac{1}{2} \sqrt{1 + \text{tr } A} \right) = \pi + \cos^{-1} \left(\frac{1}{2} [1 - \text{tr } A] \right).$$

In particular, $a = 1$ if and only if $\theta = 0$; $a = 0$ if and only if $\theta = \pi$; and $a = -1$ if and only if $\theta = 2\pi$. Furthermore,

$$\begin{aligned} A &= (2a^2 - 1)I_n + 2aK(v) + 2vv^T \\ &= (\cos \theta)I + (\sin \theta)K(z) + (1 - \cos \theta)zz^T \\ &= I + (\sin \theta)K(z) + (1 - \cos \theta)K^2(z). \end{aligned}$$

Furthermore,

$$A - A^T = 4aK(v) = 2(\sin \theta)K(z),$$

and thus

$$2a \sin \frac{\theta}{2} = \sin \theta.$$

If $\theta = 0$ or $\theta = 2\pi$, then $v = z = 0$, whereas, if $\theta = \pi$, then

$$K^2(z) = \frac{1}{2}(A - I).$$

Conversely, let $\theta \in \mathbb{R}$, let $z \in \mathbb{R}^3$, assume that $z^T z = 1$, and define

$$\begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} \cos \frac{\theta}{2} \\ (\sin \frac{\theta}{2})z \end{bmatrix}.$$

Then, A represents a rotation about the unit-length vector z through the angle θ , where the direction of rotation is determined by the right-hand rule. In this case, A is given by

$$A = \begin{bmatrix} z_{(1)}^2 + (z_{(2)}^2 + z_{(3)}^2) \cos \theta & z_{(1)} z_{(2)} (1 - \cos \theta) - z_{(3)} \sin \theta & z_{(1)} z_{(3)} (1 - \cos \theta) + z_{(2)} \sin \theta \\ z_{(1)} z_{(2)} (1 - \cos \theta) + z_{(3)} \sin \theta & z_{(2)}^2 + (z_{(1)}^2 + z_{(3)}^2) \cos \theta & z_{(2)} z_{(3)} (1 - \cos \theta) - z_{(1)} \sin \theta \\ z_{(1)} z_{(3)} (1 - \cos \theta) - z_{(2)} \sin \theta & z_{(2)} z_{(3)} (1 - \cos \theta) + z_{(1)} \sin \theta & z_{(3)}^2 + (z_{(1)}^2 + z_{(2)}^2) \cos \theta \end{bmatrix}.$$

(Proof: See [477, p. 162], [555, p. 22], [1185, p. 19], and use Fact 3.11.8.) (Remark: This result is due to Rodrigues.) (Remark: The numbers a, b, c, d , which are *Euler parameters*, are elements of S^3 , which is the sphere in $\mathbb{R}^4$. The elements of S^3 can be viewed as unit quaternions, thus giving S^3 a group structure. See Fact 3.21.2. Conversely, a, b, c, d can be expressed in terms of the entries of a 3×3 orthogonal matrix, which are the *direction cosines*. See [152, pp. 384–387]. See also Fact 3.22.1.) (Remark: Replacing a by $-a$ in A but keeping b, c, d unchanged yields the transpose of A .) (Remark: Note that A is unchanged when a, b, c, d are replaced by $-a, -b, -c, -d$. Conversely, given the direction cosines of a rotation matrix A , there exist exactly two distinct quadruples (a, b, c, d) of Euler parameters that parameterize A . Therefore, the Euler parameters, which parameterize the unit sphere S^3 in $\mathbb{R}^4$, provide a *double cover* of $SO(3)$. See [969, p. 304] and Fact 3.22.1.) (Remark: $Sp(1)$ is a double cover of $SO(3)$, $Sp(1) \times Sp(1)$ is a double cover of $SO(4)$, $Sp(2)$ is a double cover of $SO(5)$, and $SU(4)$ is a double cover of $SO(3)$. For each n , $SO(n)$ is double covered by the *spin group* $Spin(n)$. See [362, p. 141], [1256, p. 130], and [1436, pp. 42–47]. $Sp(2)$ is defined in Fact 3.22.4.) (Remark: Rotation matrices in $\mathbb{R}^{2 \times 2}$ are discussed in [1196].) (Remark: A history of Rodrigues's contributions is given in [27].) (Remark: See Fact 8.9.26 and Fact 11.15.10.) (Remark: Extensions to $n \times n$ matrices are considered in [538].)

Fact 3.11.11. Let $\theta_1, \theta_2 \in \mathbb{R}$, let $z_1, z_2 \in \mathbb{R}^3$, assume that $z_1^T z_1 = z_2^T z_2 = 1$, and, for $i = 1, 2$, let $A_i \in \mathbb{R}^{3 \times 3}$ be the rotation matrix that represents the rotation about the unit-length vector z_i through the angle θ_i , where the direction of rotation is determined by the right-hand rule. Then, $A_3 \triangleq A_2 A_1$ represents the rotation about the unit-length vector z_3 through the angle θ_3 , where the direction of rotation is determined by the right-hand rule, and where θ_3 and z_3 are given by

$$\cos \frac{\theta_3}{2} = (\cos \frac{\theta_2}{2}) \cos \frac{\theta_1}{2} - (\sin \frac{\theta_2}{2}) \sin \frac{\theta_1}{2} z_2^T z_1$$

and

$$\begin{aligned} z_3 &= (\csc \frac{\theta_3}{2}) [(\sin \frac{\theta_2}{2})(\cos \frac{\theta_1}{2})z_2 + (\cos \frac{\theta_2}{2})(\sin \frac{\theta_1}{2})z_1 + (\sin \frac{\theta_2}{2})(\sin \frac{\theta_1}{2})(z_2 \times z_1)] \\ &= \frac{\cot \frac{\theta_3}{2}}{1 - z_2^T z_1 (\tan \frac{\theta_2}{2}) \tan \frac{\theta_1}{2}} [(\tan \frac{\theta_2}{2})z_2 + (\tan \frac{\theta_1}{2})z_1 + (\tan \frac{\theta_2}{2})(\tan \frac{\theta_1}{2})(z_2 \times z_1)]. \end{aligned}$$

(Proof: See [555, pp. 22–24].) (Remark: These expressions are *Rodrigues's formu-*

las, which are identical to the quaternion multiplication formula given by

$$\begin{bmatrix} a_3 \\ b_3 \\ c_3 \\ d_3 \end{bmatrix} = \begin{bmatrix} \cos \frac{\theta_3}{2} \\ (\sin \frac{\theta_3}{2})z_3 \end{bmatrix} = \begin{bmatrix} a_1a_2 - z_2^T z_1 \\ a_1z_2 + a_2z_1 + z_2 \times z_1 \end{bmatrix}$$

with

$$\begin{bmatrix} a_2 \\ b_2 \\ c_2 \\ d_2 \end{bmatrix} = \begin{bmatrix} \cos \frac{\theta_2}{2} \\ (\sin \frac{\theta_2}{2})z_2 \end{bmatrix}, \quad \begin{bmatrix} a_1 \\ b_1 \\ c_1 \\ d_1 \end{bmatrix} = \begin{bmatrix} \cos \frac{\theta_1}{2} \\ (\sin \frac{\theta_1}{2})z_1 \end{bmatrix}$$

in Fact 3.22.1. See [27].)

Fact 3.11.12. Let $x, y, z \in \mathbb{R}^2$. If x is rotated according to the right-hand rule through an angle $\theta \in \mathbb{R}$ about y , then the resulting vector $\hat{x} \in \mathbb{R}^2$ is given by

$$\hat{x} = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} x + \begin{bmatrix} y_{(1)}(1 - \cos \theta) + y_{(2)} \sin \theta \\ y_{(2)}(1 - \cos \theta) + y_{(1)} \sin \theta \end{bmatrix}.$$

If x is reflected across the line passing through 0 and z and parallel to the line passing through 0 and y , then the resulting vector $\hat{x} \in \mathbb{R}^2$ is given by

$$\hat{x} = \begin{bmatrix} y_{(1)}^2 - y_{(2)}^2 & 2y_{(1)}y_{(2)} \\ 2y_{(1)}y_{(2)} & y_{(2)}^2 - y_{(1)}^2 \end{bmatrix} x + \begin{bmatrix} -z_{(1)}(y_{(1)}^2 - y_{(2)}^2 - 1) - 2z_{(2)}y_{(1)}y_{(2)} \\ -z_{(2)}(y_{(1)}^2 - y_{(2)}^2 - 1) - 2z_{(1)}y_{(1)}y_{(2)} \end{bmatrix}.$$

(Remark: These *affine planar transformations* are used in computer graphics. See [62, 498, 1095].) (Remark: See Fact 3.11.13 and Fact 3.11.31.)

Fact 3.11.13. Let $x, y \in \mathbb{R}^3$, and assume that $y^T y = 1$. If x is rotated according to the right-hand rule through an angle $\theta \in \mathbb{R}$ about the line passing through 0 and y , then the resulting vector $\hat{x} \in \mathbb{R}^3$ is given by

$$\hat{x} = x + (\sin \theta)(y \times x) + (1 - \cos \theta)[y \times (y \times x)].$$

(Proof: See [23].) (Remark: See Fact 3.11.12 and Fact 3.11.31.)

Fact 3.11.14. Let $x, y \in \mathbb{F}^n$, let $A \in \mathbb{F}^{n \times n}$, and assume that A is unitary. Then, $x^* y = 0$ if and only if $(Ax)^* Ay = 0$.

Fact 3.11.15. Let $A \in \mathbb{F}^{n \times n}$, assume that A is unitary, and let $x \in \mathbb{F}^n$ be such that $x^* x = 1$ and $Ax = -x$. Then, the following statements hold:

- i) $\det(A + I) = 0$.
- ii) $A + 2xx^*$ is unitary.
- iii) $A = (A + 2xx^*)(I_n - 2xx^*) = (I_n - 2xx^*)(A + 2xx^*)$.
- iv) $\det(A + 2xx^*) = -\det A$.

Fact 3.11.16. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is unitary. Then,

$$|\operatorname{Re tr} A| \leq n,$$

$$|\operatorname{Im} \operatorname{tr} A| \leq n,$$

and

$$|\operatorname{tr} A| \leq n.$$

(Remark: The third inequality does not follow from the first two inequalities.)

Fact 3.11.17. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is orthogonal. Then,

$$-1_{n \times n} \leq A \leq 1_{n \times n}$$

and

$$-n \leq \operatorname{tr} A \leq n.$$

Furthermore, the following statements are equivalent:

- i) $A = I$.
- ii) $\operatorname{diag}(A) = I$.
- iii) $\operatorname{tr} A = n$.

Finally, if n is odd and $\det A = 1$, then

$$2 - n \leq \operatorname{tr} A \leq n.$$

(Remark: See Fact 3.11.18.)

Fact 3.11.18. Let $A \in \mathbb{R}^{n \times n}$, assume that A is orthogonal, let $B \in \mathbb{R}^{n \times n}$, and assume that B is diagonal and positive definite. Then,

$$-B1_{n \times n} \leq BA \leq B1_{n \times n}$$

and

$$-\operatorname{tr} B \leq \operatorname{tr} BA \leq \operatorname{tr} B.$$

Furthermore, the following statements are equivalent:

- i) $BA = B$.
- ii) $\operatorname{diag}(BA) = B$.
- iii) $\operatorname{tr} BA = \operatorname{tr} B$.

(Remark: See Fact 3.11.17.)

Fact 3.11.19. Let $x \in \mathbb{C}^n$, where $n \geq 2$. Then, the following statements are equivalent:

- i) There exists a unitary matrix $A \in \mathbb{C}^{n \times n}$ such that

$$x = \begin{bmatrix} A_{(1,1)} \\ \vdots \\ A_{(n,n)} \end{bmatrix}.$$

- ii) For all $j = 1, \dots, n$, $|x_{(j)}| \leq 1$ and

$$2(1 - |x_{(j)}|) + \sum_{i=1}^n |x_{(i)}| \leq n.$$

(Proof: See [1338].) (Remark: This result is equivalent to the Schur-Horn theorem given by Fact 8.17.10.) (Remark: The inequalities in *ii*) define a polytope.)

Fact 3.11.20. Let $A \in \mathbb{C}^{n \times n}$, and assume that A is unitary. Then, $|\det A| = 1$.

Fact 3.11.21. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is orthogonal. Then, either $\det A = 1$ or $\det A = -1$.

Fact 3.11.22. Let $A, B \in \text{SO}(3)$. Then,

$$\det(A + B) \geq 0.$$

(Proof: See [1013].)

Fact 3.11.23. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is unitary. Then,

$$|\det(I + A)| \leq 2^n.$$

If, in addition, A is real, then

$$0 \leq \det(I + A) \leq 2^n.$$

Fact 3.11.24. Let $M \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, and assume that M is unitary. Then,

$$\det A = (\det M) \overline{\det D}.$$

(Proof: Let $\begin{bmatrix} \hat{A} & \hat{B} \\ \hat{C} & \hat{D} \end{bmatrix} \triangleq A^{-1}$, and take the determinant of $A \begin{bmatrix} I & \hat{B} \\ 0 & \hat{D} \end{bmatrix} = \begin{bmatrix} A & 0 \\ C & I \end{bmatrix}$. See [12] or [1188].) (Remark: See Fact 2.14.28 and Fact 2.14.7.)

Fact 3.11.25. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is block diagonal. Then, A is (unitary, shifted unitary) if and only if every diagonally located block has the same property.

Fact 3.11.26. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is unitary. Then, $\frac{1}{\sqrt{2}} \begin{bmatrix} A & -A \\ A & A \end{bmatrix}$ is unitary.

Fact 3.11.27. Let $A, B \in \mathbb{R}^{n \times n}$. Then, $A + jB$ is (Hermitian, skew Hermitian, unitary) if and only if $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$ is (symmetric, skew symmetric, orthogonal). (Remark: See Fact 2.19.7.)

Fact 3.11.28. The following statements hold:

- i*) If $A \in \mathbb{F}^{n \times n}$ is skew Hermitian, then $I + A$ is nonsingular, $B \triangleq (I - A)(I + A)^{-1}$ is unitary, and $I + B = 2(I + A)^{-1}$. If, in addition, $\text{mspec}(A) = \text{mspec}(A)$, then $\det B = 1$.
- ii*) If $B \in \mathbb{F}^{n \times n}$ is unitary and $\lambda \in \mathbb{C}$ is such that $|\lambda| = 1$ and $I + \lambda B$ is nonsingular, then $A \triangleq (I + \lambda B)^{-1}(I - \lambda B)$ is skew Hermitian and $I + A = 2(I + \lambda B)^{-1}$.
- iii*) If $A \in \mathbb{F}^{n \times n}$ is skew Hermitian, then there exists a unique unitary matrix $B \in \mathbb{F}^{n \times n}$ such that $I + B$ is nonsingular and $A = (I + B)^{-1}(I - B)$. In fact, $B \triangleq (I - A)(I + A)^{-1}$.

- iv)* If B is unitary and $\lambda \in \mathbb{C}$ is such that $|\lambda| = 1$ and $I + \lambda B$ is nonsingular, then there exists a unique skew-Hermitian matrix $A \in \mathbb{F}^{n \times n}$ such that $B = \overline{\lambda}(I - A)(I + A)^{-1}$. In fact, $A \triangleq (I + \lambda B)^{-1}(I - \lambda B)$.

(Proof: See [508, p. 184] and [711, p. 440].) (Remark: $\mathcal{C}(A) \triangleq (A - I)(A + I)^{-1} = I - 2(A + I)^{-1}$ is the *Cayley transform* of A . See Fact 3.11.8, Fact 3.11.29, Fact 3.11.30, Fact 3.11.31, Fact 3.19.12, Fact 8.9.30, and Fact 11.21.8.)

Fact 3.11.29. The following statements hold:

- i)* If $A \in \mathbb{F}^{n \times n}$ is Hermitian, then $A + jI$ is nonsingular, $B \triangleq (jI - A)(jI + A)^{-1}$ is unitary, and $I + B = 2j(jI + A)^{-1}$.
- ii)* If $B \in \mathbb{F}^{n \times n}$ is unitary and $\lambda \in \mathbb{C}$ is such that $|\lambda| = 1$ and $I + \lambda B$ is nonsingular, then $A \triangleq j(I - \lambda B)(I + \lambda B)^{-1}$ is Hermitian and $jI + A = 2j(I + \lambda B)^{-1}$.
- iii)* If $A \in \mathbb{F}^{n \times n}$ is Hermitian, then there exists a unique unitary matrix $B \in \mathbb{F}^{n \times n}$ such that $I + B$ is nonsingular and $A = j(I - B)(I + B)^{-1}$. In fact, $B = (jI - A)(jI + A)^{-1}$.
- iv)* If $B \in \mathbb{F}^{n \times n}$ is unitary and $\lambda \in \mathbb{C}$ is such that $|\lambda| = 1$ and $I + \lambda B$ is nonsingular, then there exists a unique Hermitian matrix $A \in \mathbb{F}^{n \times n}$ such that $\lambda B = (jI - A)(jI + A)^{-1}$. In fact, $A \triangleq j(I - \lambda B)(I + \lambda B)^{-1}$.

(Proof: See [508, pp. 168, 169].) (Remark: The linear fractional transformation $f(s) \triangleq (j - s)/(j + s)$ maps the upper half plane of $\mathbb{C}$ onto the unit disk in $\mathbb{C}$, and the real line onto the unit circle in $\mathbb{C}$.)

Fact 3.11.30. The following statements hold:

- i)* If $A \in \mathbb{R}^{n \times n}$ is skew symmetric, then $I + A$ is nonsingular, $B \triangleq (I - A)(I + A)^{-1}$ is orthogonal, $I + B = 2(I + A)^{-1}$, and $\det B = 1$.
- ii)* If $B \in \mathbb{R}^{n \times n}$ is orthogonal, $C \in \mathbb{R}^{n \times n}$ is diagonal with diagonally located entries ± 1 , and $I + CB$ is nonsingular, then $A \triangleq (I + CB)^{-1}(I - CB)$ is skew symmetric, $I + A = 2(I + CB)^{-1}$, and $\det CB = 1$.
- iii)* If $A \in \mathbb{R}^{n \times n}$ is skew symmetric, then there exists a unique orthogonal matrix $B \in \mathbb{R}^{n \times n}$ such that $I + B$ is nonsingular and $A = (I + B)^{-1}(I - B)$. In fact, $B \triangleq (I - A)(I + A)^{-1}$.
- iv)* If $B \in \mathbb{R}^{n \times n}$ is orthogonal and $C \in \mathbb{R}^{n \times n}$ is diagonal with diagonally located entries ± 1 , then there exists a unique skew-symmetric matrix $A \in \mathbb{R}^{n \times n}$ such that $CB = (I - A)(I + A)^{-1}$. In fact, $A = (I + CB)^{-1}(I - CB)$.

(Remark: The last statement is due to Hsu. See [1098, p. 101].) (Remark: The Cayley transform is a one-to-one and onto map from the set of skew-symmetric matrices to the set of orthogonal matrices whose spectrum does not include -1 .)

Fact 3.11.31. Let $x \in \mathbb{R}^3$, assume that $x^T x = 1$, let $\theta \in [0, 2\pi)$, assume that $\theta \neq \pi$, and define the skew-symmetric matrix $A \in \mathbb{R}^{3 \times 3}$ by

$$A \triangleq -(\tan \frac{\theta}{2})K(x) = \begin{bmatrix} 0 & x_{(3)} \tan \frac{\theta}{2} & -x_{(2)} \tan \frac{\theta}{2} \\ -x_{(3)} \tan \frac{\theta}{2} & 0 & x_{(1)} \tan \frac{\theta}{2} \\ x_{(2)} \tan \frac{\theta}{2} & -x_{(1)} \tan \frac{\theta}{2} & 0 \end{bmatrix}.$$

Then, the matrix $B \in \mathbb{R}^{3 \times 3}$ defined by

$$B \triangleq (I - A)(I + A)^{-1}$$

is an orthogonal matrix that rotates vectors about x through an angle equal to θ according to the right-hand rule. (Proof: See [1008, pp. 243, 244].) (Remark: Every 3×3 skew-symmetric matrix has a representation of the form given by A .) (Remark: See Fact 3.11.10, Fact 3.11.11, Fact 3.11.12, Fact 3.11.13, Fact 3.11.30, and Fact 11.11.7.)

Fact 3.11.32. Furthermore, if $A, B \in \mathbb{F}^{n \times n}$ are unitary, then

$$\sqrt{1 - \left| \frac{1}{n} \text{tr } AB \right|^2} \leq \sqrt{1 - \left| \frac{1}{n} \text{tr } A \right|^2} + \sqrt{1 - \left| \frac{1}{n} \text{tr } B \right|^2}.$$

(Proof: See [1391].) (Remark: See Fact 2.12.1.)

Fact 3.11.33. If $A \in \mathbb{F}^{n \times n}$ is shifted unitary, then $B \triangleq 2A - I$ is unitary. Conversely, If $B \in \mathbb{F}^{n \times n}$ is unitary, then $A \triangleq \frac{1}{2}(B + I)$ is shifted unitary. (Remark: The affine mapping $f(A) \triangleq 2A - I$ from the shifted-unitary matrices to the unitary matrices is one-to-one and onto. See Fact 3.14.1 and Fact 3.15.2.) (Remark: See Fact 3.7.14 and Fact 3.13.13.)

Fact 3.11.34. If $A \in \mathbb{F}^{n \times n}$ is shifted unitary, then A is normal. Hence, the following statements are equivalent:

- i) A is shifted unitary.
- ii) $A + A^* = 2A^*A$.
- iii) $A + A^* = 2AA^*$.

(Proof: By Fact 3.11.33 there exists a unitary matrix B such that $A = \frac{1}{2}(B + I)$. Since B is normal, it follows from Fact 3.7.14 that A is normal.)

3.12 Facts on Idempotent Matrices

Fact 3.12.1. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$ be complementary subspaces, and let $A \in \mathbb{F}^{n \times n}$ be the idempotent matrix onto $\mathcal{S}_1$ along $\mathcal{S}_2$. Then, A^* is the idempotent matrix onto $\mathcal{S}_2^\perp$ along $\mathcal{S}_1^\perp$, and $A_\perp^*$ is the idempotent matrix onto $\mathcal{S}_1^\perp$ along $\mathcal{S}_2^\perp$. (Remark: See Fact 2.9.18.)

Fact 3.12.2. Let $A \in \mathbb{F}^{n \times n}$. Then, A is idempotent if and only if there exists a positive integer k such that $A^{k+1} = A^k$.

Fact 3.12.3. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is idempotent.
- ii) $\mathcal{N}(A) = \mathcal{R}(A_{\perp})$.
- iii) $\mathcal{R}(A) = \mathcal{N}(A_{\perp})$.

In this case, the following statements hold:

- iv) A is the idempotent matrix onto $\mathcal{R}(A)$ along $\mathcal{N}(A)$.
- v) $A_{\perp}$ is the idempotent matrix onto $\mathcal{N}(A)$ along $\mathcal{R}(A)$.
- vi) A^* is the idempotent matrix onto $\mathcal{N}(A)^{\perp}$ along $\mathcal{R}(A)^{\perp}$.
- vii) $A_{\perp}^*$ is the idempotent matrix onto $\mathcal{R}(A)^{\perp}$ along $\mathcal{N}(A)^{\perp}$.

(Proof: See [654, p. 146].) (Remark: See Fact 2.10.1 and Fact 5.12.18.)

Fact 3.12.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then,

$$\mathcal{R}(I - AA^*) = \mathcal{R}(2I - A - A^*).$$

(Proof: See [1287].)

Fact 3.12.5. Let $A \in \mathbb{F}^{n \times n}$. Then, A is idempotent if and only if $-A$ is skew idempotent.

Fact 3.12.6. Let $A \in \mathbb{F}^{n \times n}$. Then, A is idempotent and $\text{rank } A = 1$ if and only if there exist vectors $x, y \in \mathbb{F}^n$ such that $y^T x = 1$ and $A = xy^T$.

Fact 3.12.7. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, A^T , $\overline{A}$, and A^* are idempotent.

Fact 3.12.8. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent and skew Hermitian. Then, $A = 0$.

Fact 3.12.9. Let $A \in \mathbb{F}^{n \times n}$. Then, A is idempotent if and only if $\text{rank } A + \text{rank}(I - A) = n$.

Fact 3.12.10. Let $A \in \mathbb{F}^{n \times m}$. If $A^L \in \mathbb{F}^{m \times n}$ is a left inverse of A , then AA^L is idempotent and $\text{rank } A^L = \text{rank } A$. Furthermore, if $A^R \in \mathbb{F}^{m \times n}$ is a right inverse of A , then $A^R A$ is idempotent and $\text{rank } A^R = \text{rank } A$.

Fact 3.12.11. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular and idempotent. Then, $A = I_n$.

Fact 3.12.12. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, so is $A_{\perp} \triangleq I - A$, and, furthermore, $AA_{\perp} = A_{\perp}A = 0$.

Fact 3.12.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then,

$$\det(I + A) = 2^{\text{tr } A}$$

and

$$(I + A)^{-1} = I - \frac{1}{2}A.$$

Fact 3.12.14. Let $A \in \mathbb{F}^{n \times n}$ and $\alpha \in \mathbb{F}$, where $\alpha \neq 0$. Then, the matrices

$$\begin{bmatrix} A & A^* \\ A^* & A \end{bmatrix}, \quad \begin{bmatrix} A & \alpha^{-1}A \\ \alpha(I - A) & I - A \end{bmatrix}, \quad \begin{bmatrix} A & \alpha^{-1}A \\ -\alpha A & -A \end{bmatrix}$$

are, respectively, normal, idempotent, and nilpotent.

Fact 3.12.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then,

$$\mathcal{R}([A, B]) = \mathcal{R}(A - B) \cap \mathcal{R}(A_{\perp} - B)$$

and

$$\mathcal{N}([A, B]) = \mathcal{N}(A - B) \cap \mathcal{N}(A_{\perp} - B).$$

(Proof: See [1424].)

Fact 3.12.16. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nilpotent. Then, there exist idempotent matrices $B, C \in \mathbb{F}^{n \times n}$ such that $A = [B, C]$. (Proof: See [439].) (Remark: A necessary and sufficient condition for a matrix to be a commutator of a pair of idempotents is given in [439].) (Remark: See Fact 9.9.9 for the case of projectors.)

Fact 3.12.17. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are idempotent, and define $A_{\perp} \triangleq I - A$ and $B_{\perp} \triangleq I - B$. Then, the following identities hold:

- i) $(A - B)^2 + (A_{\perp} - B)^2 = I$.
- ii) $[A, B] = [B, A_{\perp}] = [B_{\perp}, A] = [A_{\perp}, B_{\perp}]$.
- iii) $A - B = AB_{\perp} - A_{\perp}B$.
- iv) $AB_{\perp} + BA_{\perp} = AB_{\perp}A + A_{\perp}BA_{\perp}$.
- v) $A[A, B] = [A, B]A_{\perp}$.
- vi) $B[A, B] = [A, B]B_{\perp}$.

(Proof: See [1044].)

Fact 3.12.18. Let $A, B \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) Assume that $A^3 = -A$ and $B = I + A + A^2$. Then, $B^4 = I$, $B^{-1} = I - A + A^2$, $B^3 - B^2 + B - I = 0$, $A = \frac{1}{2}(B - B^3)$, and $I + A^2$ is idempotent.
- ii) Assume that $B^3 - B^2 + B - I = 0$ and $A = \frac{1}{2}(B - B^3)$. Then, $A^3 = -A$ and $B = I + A + A^2$.
- iii) Assume that $B^4 = I$ and $A = \frac{1}{2}(B - B^{-1})$. Then, $A^3 = -A$, and $\frac{1}{4}(I + B + B^2 + B^3)$ is idempotent.

(Remark: The geometric meaning of these results is discussed in [474, pp. 153, 212–214, 242].)

Fact 3.12.19. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{l \times n}$, and assume that A is idempotent, $\text{rank} \begin{bmatrix} C^* & B \end{bmatrix} = n$, and $CB = 0$. Then,

$$\text{rank } CAB = \text{rank } CA + \text{rank } AB - \text{rank } A.$$

(Proof: See [1307].) (Remark: See Fact 3.12.20.)

Fact 3.12.20. $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, and assume that A is idempotent. Then,

$$\begin{aligned} \text{rank } A &= \text{rank} \begin{bmatrix} A_{12} \\ A_{22} \end{bmatrix} + \text{rank} \begin{bmatrix} A_{11} & A_{12} \end{bmatrix} - \text{rank } A_{12} \\ &= \text{rank} \begin{bmatrix} A_{11} \\ A_{21} \end{bmatrix} + \text{rank} \begin{bmatrix} A_{21} & A_{22} \end{bmatrix} - \text{rank } A_{21}. \end{aligned}$$

(Proof: See [1307] and Fact 3.12.19.) (Remark: See Fact 3.13.12 and Fact 6.5.13.)

Fact 3.12.21. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$, and assume that AB is nonsingular. Then, $B(AB)^{-1}A$ is idempotent.

Fact 3.12.22. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are idempotent, and let $\alpha, \beta \in \mathbb{F}$ be nonzero and satisfy $\alpha + \beta \neq 0$. Then,

$$\begin{aligned} \text{rank}(A + B) &= \text{rank}(\alpha A + \beta B) \\ &= \text{rank } A + \text{rank}(A_{\perp}BA_{\perp}) \\ &= n - \dim[\mathcal{N}(A_{\perp}B) \cap \mathcal{N}(A)] \\ &= \text{rank} \begin{bmatrix} 0 & A & B \\ A & 0 & 0 \\ B & 0 & 2B \end{bmatrix} - \text{rank } A - \text{rank } B \\ &= \text{rank} \begin{bmatrix} A & B \\ B & 0 \end{bmatrix} - \text{rank } B = \text{rank} \begin{bmatrix} B & A \\ A & 0 \end{bmatrix} - \text{rank } A \\ &= \text{rank}(B_{\perp}AB_{\perp}) + \text{rank } B = \text{rank}(A_{\perp}BA_{\perp}) + \text{rank } A \\ &= \text{rank}(A + A_{\perp}B) = \text{rank}(A + BA_{\perp}) \\ &= \text{rank}(B + B_{\perp}A) = \text{rank}(B + AB_{\perp}) \\ &= \text{rank}(I - A_{\perp}B_{\perp}) = \text{rank}(I - B_{\perp}A_{\perp}) \\ &= \text{rank} \begin{bmatrix} AB_{\perp} & B \end{bmatrix} = \text{rank} \begin{bmatrix} BA_{\perp} & A \end{bmatrix} \\ &= \text{rank} \begin{bmatrix} B_{\perp}A \\ B \end{bmatrix} = \text{rank} \begin{bmatrix} A_{\perp}B \\ A \end{bmatrix} \\ &= \text{rank } A + \text{rank } B - n + \text{rank} \begin{bmatrix} A_{\perp} & A_{\perp}B_{\perp} \\ B_{\perp}A_{\perp} & B_{\perp} \end{bmatrix}. \end{aligned}$$

Furthermore, the following statements hold:

i) If $AB = 0$, then

$$\begin{aligned} \text{rank}(A + B) &= \text{rank}(BA_{\perp}) + \text{rank } A \\ &= \text{rank}(B_{\perp}A) + \text{rank } B. \end{aligned}$$

ii) If $BA = 0$, then

$$\begin{aligned}\text{rank}(A + B) &= \text{rank}(AB_{\perp}) + \text{rank } B \\ &= \text{rank}(A_{\perp}B) + \text{rank } A.\end{aligned}$$

iii) If $AB = BA$, then

$$\begin{aligned}\text{rank}(A + B) &= \text{rank}(AB_{\perp}) + \text{rank } B \\ &= \text{rank}(BA_{\perp}) + \text{rank } A.\end{aligned}$$

iv) $A + B$ is idempotent if and only if $AB = BA = 0$.

v) $A + B = I$ if and only if $AB = BA = 0$ and $\text{rank } [A, B] = \text{rank } A + \text{rank } B = n$.

(Remark: See Fact 6.4.33.) (Proof: See [597, 835, 836, 1306, 1309]. To prove necessity in iv) note that $AB + BA = 0$ implies $AB + ABA = ABA + BA = 0$, which implies that $AB - BA = 0$, and hence $AB = 0$. See [630, p. 250] and [654, p. 435].)

Fact 3.12.23. Let $A \in \mathbb{F}^{n \times n}$, let $r \triangleq \text{rank } A$, and let $B \in \mathbb{F}^{n \times r}$ and $C \in \mathbb{F}^{r \times n}$ satisfy $A = BC$. Then, A is idempotent if and only if $CB = I$. (Proof: See [1396, p. 16].) (Remark: $A = BC$ is a full-rank factorization.)

Fact 3.12.24. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are idempotent, and let $C \in \mathbb{F}^{n \times m}$. Then,

$$\begin{aligned}\text{rank}(AC - CB) &= \text{rank}(AC - ACB) + \text{rank}(ACB - CB) \\ &= \text{rank} \begin{bmatrix} AC \\ B \end{bmatrix} + \text{rank} \begin{bmatrix} CB & A \end{bmatrix} - \text{rank } A - \text{rank } B.\end{aligned}$$

(Proof: See [1281].)

Fact 3.12.25. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then,

$$\begin{aligned}\text{rank}(A - B) &= \text{rank} \begin{bmatrix} 0 & A & B \\ A & 0 & 0 \\ B & 0 & 0 \end{bmatrix} - \text{rank } A - \text{rank } B \\ &= \text{rank} \begin{bmatrix} A \\ B \end{bmatrix} + \text{rank} \begin{bmatrix} A & B \end{bmatrix} - \text{rank } A - \text{rank } B \\ &= n - \dim[\mathcal{N}(A) \cap \mathcal{N}(B)] - \dim[\mathcal{R}(A) \cap \mathcal{R}(B)] \\ &= \text{rank}(AB_{\perp}) + \text{rank}(A_{\perp}B) \\ &\leq \text{rank}(A + B) \\ &\leq \text{rank } A + \text{rank } B.\end{aligned}$$

Furthermore, if either $AB = 0$ or $BA = 0$, then

$$\text{rank}(A - B) = \text{rank}(A + B) = \text{rank } A + \text{rank } B.$$

(Proof: See [597, 836, 1306, 1309]. The inequality $\text{rank}(A - B) \leq \text{rank}(A + B)$ follows from Fact 2.11.13 and the block 3×3 expressions in this result and in

Fact 3.12.22. To prove the last statement in the case $AB = 0$, first note that $\text{rank } A + \text{rank } B = \text{rank}(A - B)$, which yields $\text{rank}(A - B) \leq \text{rank}(A + B) \leq \text{rank } A + \text{rank } B = \text{rank}(A - B)$. (Remark: See Fact 6.4.33.)

Fact 3.12.26. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then, the following statements are equivalent:

- i) $A + B$ is nonsingular.
- ii) There exist $\alpha, \beta \in \mathbb{F}$ such that $\alpha + \beta \neq 0$ and $\alpha A + \beta B$ is nonsingular.
- iii) For all nonzero $\alpha, \beta \in \mathbb{F}$ such that $\alpha + \beta \neq 0$, $\alpha A + \beta B$ is nonsingular.

(Proof: See [104, 833, 1309].)

Fact 3.12.27. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then, the following statements are equivalent:

- i) $A - B$ is idempotent.
- ii) $\text{rank}(A_{\perp} + B) + \text{rank}(A - B) = n$.
- iii) $ABA = B$.
- iv) $\text{rank}(A - B) = \text{rank } A - \text{rank } B$.
- v) $\mathcal{R}(B) \subseteq \mathcal{R}(A)$ and $\mathcal{R}(B^*) \subseteq \mathcal{R}(A^*)$.

(Proof: See [1308].) (Remark: This result is due to Hartwig and Styan.)

Fact 3.12.28. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then, the following statements are equivalent:

- i) $A - B$ is nonsingular.
- ii) $I - AB$ is nonsingular, and there exist $\alpha, \beta \in \mathbb{F}$ such that $\alpha + \beta \neq 0$ and $\alpha A + \beta B$ is nonsingular.
- iii) $I - AB$ is nonsingular, and $\alpha A + \beta B$ is nonsingular for all $\alpha, \beta \in \mathbb{F}$ such that $\alpha + \beta \neq 0$.
- iv) $I - AB$ and $A + A_{\perp}B$ are nonsingular.
- v) $I - AB$ and $A + B$ are nonsingular.
- vi) $\mathcal{R}(A) + \mathcal{R}(B) = \mathbb{F}^n$ and $\mathcal{R}(A^*) + \mathcal{R}(B^*) = \mathbb{F}^n$.
- vii) $\mathcal{R}(A) + \mathcal{R}(B) = \mathbb{F}^n$ and $\mathcal{N}(A) + \mathcal{N}(B) = \mathbb{F}^n$.
- viii) $\mathcal{R}(A) \cap \mathcal{R}(B) = \{0\}$ and $\mathcal{N}(A) \cap \mathcal{N}(B) = \{0\}$.
- ix) $\text{rank}\begin{bmatrix} A & B \end{bmatrix} = \text{rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank } A + \text{rank } B = n$.

(Proof: See [104, 597, 834, 836, 1306].)

Fact 3.12.29. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are idempotent. Then, the following statements hold:

- i) $\mathcal{R}(A) \cap \mathcal{R}(B) \subseteq \mathcal{R}(AB)$.

- ii) $\mathcal{N}(B) + [\mathcal{N}(A) \cap \mathcal{R}(B)] \subseteq \mathcal{N}(AB) \subseteq \mathcal{R}(I - AB) \subseteq \mathcal{N}(A) + \mathcal{N}(B)$.
- iii) If $AB = BA$, then AB is the idempotent matrix onto $\mathcal{R}(A) \cap \mathcal{R}(B)$ along $\mathcal{N}(A) + \mathcal{N}(B)$.

Furthermore, the following statements are equivalent:

- iv) $AB = BA$.
- v) $\text{rank } AB = \text{rank } BA$, and AB is the idempotent matrix onto $\mathcal{R}(A) \cap \mathcal{R}(B)$ along $\mathcal{N}(A) + \mathcal{N}(B)$.
- vi) $\text{rank } AB = \text{rank } BA$, and $A + B - AB$ is the idempotent matrix onto $\mathcal{R}(A) + \mathcal{R}(B)$ along $\mathcal{N}(A) \cap \mathcal{N}(B)$.

In addition, the following statements are equivalent:

- vii) AB is idempotent.
- viii) $\mathcal{R}(AB) \subseteq \mathcal{R}(B) + [\mathcal{N}(A) \cap \mathcal{N}(B)]$.
- ix) $\mathcal{R}(AB) = \mathcal{R}(A) \cap (\mathcal{R}(B) + [\mathcal{N}(A) \cap \mathcal{N}(B)])$.
- x) $\mathcal{N}(B) + [\mathcal{N}(A) \cap \mathcal{R}(B)] = \mathcal{R}(I - AB)$.

Finally, the following statements hold:

- xi) $A - B$ is idempotent if and only if B is the idempotent matrix onto $\mathcal{R}(A) \cap \mathcal{R}(B)$ along $\mathcal{N}(A) + \mathcal{N}(B)$.
- xii) $A + B$ is idempotent if and only if A is the idempotent matrix onto $\mathcal{R}(A) \cap \mathcal{N}(B)$ along $\mathcal{N}(A) + \mathcal{R}(B)$.

(Proof: See [536, p. 53] and [596].) (Remark: See Fact 5.12.19.)

Fact 3.12.30. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are idempotent, and assume that $AB = BA$. Then, the following statements are equivalent:

- i) $A - B$ is nonsingular.
- ii) $(A - B)^2 = I$.
- iii) $A + B = I$.

(Proof: See [597].)

Fact 3.12.31. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then,

$$\begin{aligned} \text{rank } [A, B] &= \text{rank}(A - B) + \text{rank}(A_{\perp} - B) - n \\ &= \text{rank}(A - B) + \text{rank } AB + \text{rank } BA - \text{rank } A - \text{rank } B. \end{aligned}$$

Furthermore, the following statements hold:

- i) $AB = BA$ if and only if $\mathcal{R}(AB) = \mathcal{R}(BA)$ and $\mathcal{R}[(AB)^*] = \mathcal{R}[(BA)^*]$.
- ii) $AB = BA$ if and only if

$$\text{rank}(A - B) + \text{rank}(A_{\perp} - B) = n.$$

- iii) $[A, B]$ is nonsingular if and only if $A - B$ and $A_{\perp} - B$ are nonsingular.

- iv)* $\max\{\text{rank } AB, \text{rank } BA\} \leq \text{rank}(AB + BA)$.
- v)* $AB + BA = 0$ if and only if $AB = BA = 0$.
- vi)* $AB + BA$ is nonsingular if and only if $A + B$ and $A_{\perp} - B$ are nonsingular.
- vii)* $\text{rank}(AB + BA) = \text{rank}(\alpha AB + \beta BA)$.
- viii)* $A_{\perp} - B$ is nonsingular if and only if $\text{rank } A = \text{rank } B = \text{rank } AB = \text{rank } BA$.
In this case, A and B are similar.
- ix)* $\text{rank}(A + B) + \text{rank}(AB - BA) = \text{rank}(A - B) + \text{rank}(AB + BA)$.
- x)* $\text{rank}(AB - BA) \leq \text{rank}(AB + BA)$.

(Proof: See [836].)

Fact 3.12.32. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are idempotent, and assume that $A - B$ is nonsingular. Then, $A + B$ is nonsingular. Now, define $F, G \in \mathbb{F}^{n \times n}$ by

$$F \triangleq A(A - B)^{-1} = (A - B)^{-1}(I - B)$$

and

$$G \triangleq (A - B)^{-1}A = (I - A)(A - B)^{-1}.$$

Then, F and G are idempotent. In particular, F is the idempotent matrix onto $\mathcal{R}(A)$ along $\mathcal{N}(B)$, and G^* is the idempotent matrix onto $\mathcal{R}(A^*)$ along $\mathcal{R}(B^*)$. Furthermore,

$$FB = AG = 0,$$

$$(A - B)^{-1} = F - G_{\perp},$$

$$(A - B)^{-1} = (A + B)^{-1}(A - B)(A + B)^{-1},$$

$$(A + B)^{-1} = I - G_{\perp}F - GF_{\perp},$$

$$(A + B)^{-1} = (A - B)^{-1}(A + B)(A - B)^{-1}.$$

(Proof: See [836].) (Remark: See [836] for an explicit expression for $(A + B)^{-1}$ in the case $A - B$ is nonsingular.) (Remark: See Proposition 3.5.3.)

Fact 3.12.33. If $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times (n-m)}$, assume that $[A \ B]$ is nonsingular, and define

$$P \triangleq \begin{bmatrix} A & 0 \end{bmatrix} \begin{bmatrix} A & B \end{bmatrix}^{-1}$$

and

$$Q \triangleq \begin{bmatrix} 0 & B \end{bmatrix} \begin{bmatrix} A & B \end{bmatrix}^{-1}.$$

Then, the following statements hold:

- i)* P and Q are idempotent.
- ii)* $P + Q = I_n$.
- iii)* $PQ = 0$.
- iv)* $P \begin{bmatrix} A & 0 \end{bmatrix} = \begin{bmatrix} A & 0 \end{bmatrix}$.
- v)* $Q \begin{bmatrix} 0 & B \end{bmatrix} = \begin{bmatrix} 0 & B \end{bmatrix}$.

- vi)* $\mathcal{R}(P) = \mathcal{R}(A)$ and $\mathcal{N}(P) = \mathcal{R}(B)$.
- vii)* $\mathcal{R}(Q) = \mathcal{R}(B)$ and $\mathcal{N}(Q) = \mathcal{R}(A)$.
- viii)* If $A^*B = 0$, then $P = A(A^*A)^{-1}A$ and $Q = B(B^*B)^{-1}B^*$.
- ix)* $\mathcal{R}(A)$ and $\mathcal{R}(B)$ are complementary subspaces.
- x)* P is the idempotent matrix onto $\mathcal{R}(A)$ along $\mathcal{R}(B)$.
- xi)* Q is the idempotent matrix onto $\mathcal{R}(B)$ along $\mathcal{R}(A)$.

(Proof: See [1497].) (Remark: See Fact 3.13.24, Fact 6.4.18, and Fact 6.4.19.)

3.13 Facts on Projectors

Fact 3.13.1. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i)* A is a projector.
- ii)* $A = AA^*$.
- iii)* $A = A^*A$.
- iv)* A is idempotent and normal.
- v)* A and A^*A are idempotent.
- vi)* $AA^*A = A$, and A is idempotent.
- vii)* A and $\frac{1}{2}(A + A^*)$ are idempotent.
- viii)* A is idempotent, and $AA^* + A^*A = A + A^*$.
- ix)* A is tripotent, and $A^2 = A^*$.
- x)* $AA^* = A^*AA^*$.
- xi)* A is idempotent, and $\text{rank } A + \text{rank}(I - A^*A) = n$.
- xii)* A is idempotent, and, for all $x \in \mathbb{F}^n$, $x^*Ax \geq 0$.

(Remark: See Fact 3.13.2, Fact 3.13.3, and Fact 6.3.27.) (Remark: The matrix $A = \begin{bmatrix} 1/2 & 1/2 \\ 0 & 0 \end{bmatrix}$ satisfies $\text{tr } A = \text{tr } A^*A$ but is not a projector. See Fact 3.7.13.)

Fact 3.13.2. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, the following statements are equivalent:

- i)* A is a projector.
- ii)* $\text{rank } A = \text{tr } A = \text{tr } A^2$.

(Proof: See [1184, p. 55].) (Remark: See Fact 3.13.1 and Fact 3.13.3.)

Fact 3.13.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, the following statements are equivalent:

- i)* A is a projector.
- ii)* $AA^*A = A$.

- iii) A is Hermitian.
- iv) A is normal.
- v) A is range Hermitian.

(Proof: See [1335].) (Remark: See Fact 3.13.1 and Fact 3.13.2.)

Fact 3.13.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is a projector. Then, A is positive semidefinite.

Fact 3.13.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is a projector, and let $x \in \mathbb{F}^n$. Then, $x \in \mathcal{R}(A)$ if and only if $x = Ax$.

Fact 3.13.6. Let $A \in \mathbb{F}^{n \times m}$. If $\text{rank } A = m$, then $B \triangleq A(A^*A)^{-1}A^*$ is a projector and $\text{rank } B = m$. If $\text{rank } A = n$, then $B \triangleq A^*(AA^*)^{-1}A$ is a projector and $\text{rank } B = n$. (Remark: See Fact 2.15.2, Fact 3.7.25, and Fact 3.7.26.)

Fact 3.13.7. Let $x \in \mathbb{F}^n$ be nonzero, and define the elementary projector $A \triangleq I - (x^*x)^{-1}xx^*$. Then, the following statements hold:

- i) $\text{rank } A = n - 1$.
- ii) $\mathcal{N}(A) = \text{span}\{x\}$.
- iii) $\mathcal{R}(A) = \{x\}^\perp$.
- iv) $2A - I$ is the elementary reflector $I - 2(x^*x)^{-1}xx^*$.

(Remark: If $y \in \mathbb{F}^n$, then Ay is the *projection* of y on $\{x\}^\perp$.)

Fact 3.13.8. Let $n > 1$, let $\mathcal{S} \subset \mathbb{F}^n$, and assume that $\mathcal{S}$ is a hyperplane. Then, there exists a unique elementary projector $A \in \mathbb{F}^{n \times n}$ such that $\mathcal{R}(A) = \mathcal{S}$ and $\mathcal{N}(A) = \mathcal{S}^\perp$. Furthermore, if $x \in \mathbb{F}^n$ is nonzero and $\mathcal{S} \triangleq \{x\}^\perp$, then $A = I - (x^*x)^{-1}xx^*$.

Fact 3.13.9. Let $A \in \mathbb{F}^{n \times n}$. Then, A is a projector and $\text{rank } A = n - 1$ if and only if there exists a nonzero vector $x \in \mathcal{N}(A)$ such that

$$A = I - (x^*x)^{-1}xx^*.$$

In this case, it follows that, for all $y \in \mathbb{F}^n$,

$$y^*y - y^*Ay = \frac{|y^*x|^2}{x^*x}.$$

Furthermore, for $y \in \mathbb{F}^n$, the following statements are equivalent:

- i) $y^*Ay = y^*y$.
- ii) $y^*x = 0$.
- iii) $Ay = y$.

(Remark: See Fact 3.7.19.)

Fact 3.13.10. Let $A \in \mathbb{F}^{n \times n}$, assume that A is a projector, and let $x \in \mathbb{F}^n$. Then,

$$x^*Ax \leq x^*x.$$

Furthermore, the following statements are equivalent:

- i) $x^*Ax = x^*x$.
- ii) $Ax = x$.
- iii) $x \in \mathcal{R}(A)$.

Fact 3.13.11. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, A is a projector if and only if, for all $x \in \mathbb{F}^n$, $x^*Ax \leq x^*x$. (Proof: See [1098, p. 105].)

Fact 3.13.12. $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, and assume that A is a projector. Then,

$$\text{rank } A = \text{rank } A_{11} + \text{rank } A_{22} - \text{rank } A_{12}.$$

(Proof: See [1308] and Fact 3.12.20.) (Remark: See Fact 3.12.20 and Fact 6.5.13.)

Fact 3.13.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that A satisfies two out of the three properties (Hermitian, shifted unitary, idempotent). Then, A satisfies the remaining property. Furthermore, these matrices are the projectors. (Proof: If A is idempotent and shifted unitary, then $(2A - I)^{-1} = 2A - I = (2A^* - I)^{-1}$. Hence, A is Hermitian.) (Remark: The condition $A + A^* = 2AA^*$ is considered in Fact 3.11.33.) (Remark: See Fact 3.14.2 and Fact 3.14.6.)

Fact 3.13.14. Let $A \in \mathbb{F}^{n \times n}$, let $B \in \mathbb{F}^{n \times m}$, assume that A is a projector, and assume that $\mathcal{R}(AB) = \mathcal{R}(B)$. Then, $AB = B$. (Proof: $0 = \mathcal{R}(A_{\perp}AB) = A_{\perp}\mathcal{R}(AB) = A_{\perp}\mathcal{R}(B) = \mathcal{R}(A_{\perp}B)$. Hence, $A_{\perp}B = 0$. Consequently, $B = (A + A_{\perp})B = AB$.) (Remark: See Fact 6.4.16.)

Fact 3.13.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, $\mathcal{R}(A) = \mathcal{R}(B)$ if and only if $A = B$. (Remark: See Proposition 3.5.1.)

Fact 3.13.16. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are projectors, and assume that $\text{rank } A = \text{rank } B$. Then, there exists a reflector $S \in \mathbb{F}^{n \times n}$ such that $A = SBS$. If, in addition, $A + B - I$ is nonsingular, then one such reflector is given by $S = \langle A + B - I \rangle (A + B - I)^{-1}$. (Proof: See [327].)

Fact 3.13.17. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the following statements are equivalent:

- i) $\mathcal{R}(A) \subseteq \mathcal{R}(B)$.
- ii) $A \leq B$.
- iii) $AB = A$.
- iv) $BA = A$.
- v) $B - A$ is a projector

(Proof: See [1184, pp. 24, 169].) (Remark: See Fact 9.8.3.)

Fact 3.13.18. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then,

$$\mathcal{R}(I - AB) = \mathcal{N}(A) + \mathcal{N}(B)$$

and

$$\mathcal{R}(A + A_{\perp}B) = \mathcal{R}(A) + \mathcal{R}(B).$$

(Proof: See [594, 1328].)

Fact 3.13.19. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the following statements are equivalent:

- i) $AB = 0$.
- ii) $BA = 0$.
- iii) $\mathcal{R}(A) = \mathcal{R}(B)^{\perp}$.
- iv) $A + B$ is a projector.

In this case, $\mathcal{R}(A + B) = \mathcal{R}(A) + \mathcal{R}(B)$. (Proof: See [530, pp. 42–44].) (Remark: See [537].)

Fact 3.13.20. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the following statements are equivalent:

- i) AB is a projector.
- ii) $AB = BA$.
- iii) AB is idempotent.
- iv) AB is Hermitian.
- v) AB is normal.
- vi) AB is range Hermitian.

In this case, the following statements hold:

- vii) $\mathcal{R}(AB) = \mathcal{R}(A) \cap \mathcal{R}(B)$.
- viii) AB is the projector onto $\mathcal{R}(A) \cap \mathcal{R}(B)$.
- ix) $A + A_{\perp}B$ is a projector.
- x) $A + A_{\perp}B$ is the projector onto $\mathcal{R}(A) + \mathcal{R}(B)$.

(Proof: See [530, pp. 42–44] and [1321, 1423].) (Remark: See Fact 5.12.16 and Fact 6.4.23.) (Problem: If $A + A_{\perp}B$ is a projector, then does it follow that A and B commute?)

Fact 3.13.21. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, AB is group invertible. (Proof: $\mathcal{N}(BA) \subseteq \mathcal{N}(BABA) \subseteq \mathcal{N}(ABABA) = \mathcal{N}(ABAABA) = \mathcal{N}(ABA) = \mathcal{N}(ABBA) = \mathcal{N}(BA)$.) (Remark: See [1423].)

Fact 3.13.22. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the $ln \times ln$ matrix below has rank

$$\text{rank} \begin{bmatrix} A+B & AB & & & \\ AB & A+B & \ddots & & \\ & \ddots & \ddots & \ddots & \\ & & \ddots & A+B & AB \\ & & & AB & A+B \end{bmatrix} = l \text{rank}(A+B).$$

(Proof: See [1309].)

Fact 3.13.23. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then,

$$\text{rank}(A+B) = \text{rank } A + \text{rank } B - n + \text{rank}(A_{\perp} + B_{\perp}),$$

$$\text{rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank } A + \text{rank } B - n + \text{rank} \begin{bmatrix} A_{\perp} & B_{\perp} \end{bmatrix},$$

$$\text{rank} [A, B] = 2(\text{rank} \begin{bmatrix} A & B \end{bmatrix} + \text{rank } AB - \text{rank } A - \text{rank } B).$$

(Proof: See [1306, 1309].)

Fact 3.13.24. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the following statements are equivalent:

- i) $A - B$ is nonsingular.
- ii) $\text{rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank } A + \text{rank } B = n$.
- iii) $\mathcal{R}(A)$ and $\mathcal{R}(B)$ are complementary subspaces.

Now, assume that i)–iii) hold. Then, the following statements hold:

- iv) $I - BA$ is nonsingular.
- v) $A + B - AB$ is nonsingular.
- vi) The idempotent matrix $M \in \mathbb{F}^{n \times n}$ onto $\mathcal{R}(B)$ along $\mathcal{R}(A)$ is given by

$$\begin{aligned} M &= (I - BA)^{-1}B(I - BA) \\ &= B(I - AB)^{-1}(I - BA) \\ &= (I - AB)^{-1}(I - A) \\ &= A(A + B - AB)^{-1}. \end{aligned}$$

- vii) M satisfies

$$M + M^* = (B - A)^{-1} + I,$$

that is,

$$(B - A)^{-1} = M + M^* - I = M - M_{\perp}^*.$$

(Proof: See Fact 5.12.17 and [6, 271, 537, 588, 744, 1115]. The uniqueness of M follows from Proposition 3.5.3, while vii) follows from Fact 5.12.18.) (Remark: See

Fact 3.12.33, Fact 5.12.18, Fact 6.4.18, and Fact 6.4.19.)

3.14 Facts on Reflectors

Fact 3.14.1. If $A \in \mathbb{F}^{n \times n}$ is a projector, then $B \triangleq 2A - I$ is a reflector. Conversely, if $B \in \mathbb{F}^{n \times n}$ is a reflector, then $A \triangleq \frac{1}{2}(B + I)$ is a projector. (Remark: See Fact 3.15.2.) (Remark: The affine mapping $f(A) \triangleq 2A - I$ from the projectors to the reflectors is one-to-one and onto. See Fact 3.11.33 and Fact 3.15.2.)

Fact 3.14.2. Let $A \in \mathbb{F}^{n \times n}$, and assume that A satisfies two out of the three properties (Hermitian, unitary, involutory). Then, A also satisfies the remaining property. Furthermore, these matrices are the reflectors. (Remark: See Fact 3.13.13 and Fact 3.14.6.)

Fact 3.14.3. Let $x \in \mathbb{F}^n$ be nonzero, and define the elementary reflector $A \triangleq I - 2(x^*x)^{-1}xx^*$. Then, the following statements hold:

- i) $\det A = -1$.
- ii) If $y \in \mathbb{F}^n$, then Ay is the reflection of y across $\{x\}^\perp$.
- iii) $Ax = -x$.
- iv) $\frac{1}{2}(A + I)$ is the elementary projector $I - (x^*x)^{-1}xx^*$.

Fact 3.14.4. Let $x, y \in \mathbb{F}^n$. Then, there exists a unique elementary reflector $A \in \mathbb{F}^{n \times n}$ such that $Ax = y$ if and only if x^*y is real and $x^*x = y^*y$. If, in addition, $x \neq y$, then A is given by

$$A = I - 2[(x - y)^*(x - y)]^{-1}(x - y)(x - y)^*.$$

(Remark: This result is the *reflection theorem*. See [558, pp. 16–18] and [1129, p. 357]. See Fact 3.9.5.)

Fact 3.14.5. Let $n > 1$, let $\mathcal{S} \subset \mathbb{F}^n$, and assume that $\mathcal{S}$ is a hyperplane. Then, there exists a unique elementary reflector $A \in \mathbb{F}^{n \times n}$ such that, for all $y = y_1 + y_2 \in \mathbb{F}^n$, where $y_1 \in \mathcal{S}$ and $y_2 = \mathcal{S}^\perp$, it follows that $Ay = y_1 - y_2$. Furthermore, if $\mathcal{S} = \{x\}^\perp$, then $A = I - 2(x^*x)^{-1}xx^*$.

Fact 3.14.6. Let $A \in \mathbb{F}^{n \times n}$, and assume that A satisfies two out of the three properties (skew Hermitian, unitary, skew involutory). Then, A also satisfies the remaining property. Furthermore, these matrices are the skew reflectors. (Remark: See Fact 3.13.13, Fact 3.14.2, and Fact 3.14.7.)

Fact 3.14.7. Let $A \in \mathbb{C}^{n \times n}$. Then, A is a reflector if and only if jA is a skew reflector. (Remark: The mapping $f(A) \triangleq jA$ relates Fact 3.14.2 to Fact 3.14.6.) (Problem: When A is real and n is even, determine a real transformation between the reflectors and the skew reflectors.)

Fact 3.14.8. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is a reflector.
- ii) $A = AA^* + A^* - I$.
- iii) $A = \frac{1}{2}(A + I)(A^* + I) - I$.

3.15 Facts on Involutory Matrices

Fact 3.15.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is involutory. Then, either $\det A = 1$ or $\det A = -1$.

Fact 3.15.2. If $A \in \mathbb{F}^{n \times n}$ is idempotent, then $B \triangleq 2A - I$ is involutory. Conversely, if $B \in \mathbb{F}^{n \times n}$ is involutory, then $A_1 \triangleq \frac{1}{2}(I + B)$ and $A_2 \triangleq \frac{1}{2}(I - B)$ are idempotent. (Remark: See Fact 3.14.1.) (Remark: The affine mapping $f(A) \triangleq 2A - I$ from the idempotent matrices to the involutory matrices is one-to-one and onto. See Fact 3.11.33 and Fact 3.14.1.)

Fact 3.15.3. Let $A \in \mathbb{F}^{n \times n}$. Then, A is involutory if and only if

$$(A + I)(A - I) = 0.$$

Fact 3.15.4. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are involutory. Then,

$$\mathcal{R}([A, B]) = \mathcal{R}(A - B) \cap \mathcal{R}(A + B)$$

and

$$\mathcal{N}([A, B]) = \mathcal{N}(A - B) \cap \mathcal{N}(A + B).$$

(Proof: See [1292].)

Fact 3.15.5. Let $A \in \mathbb{F}^{n \times m}$, let $B \in \mathbb{F}^{m \times n}$, and define

$$C \triangleq \begin{bmatrix} I - BA & B \\ 2A - ABA & AB - I \end{bmatrix}.$$

Then, C is involutory. (Proof: See [998, p. 113].)

Fact 3.15.6. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is skew involutory. Then, n is even.

3.16 Facts on Tripotent Matrices

Fact 3.16.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is tripotent. Then, A^2 is idempotent. (Remark: The converse is false. A counterexample is $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.)

Fact 3.16.2. Let $A \in \mathbb{F}^{n \times n}$. Then, A is nonsingular and tripotent if and only if A is involutory.

Fact 3.16.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, A is tripotent if and only if

$$\text{rank } A = \text{rank}(A + A^2) + \text{rank}(A - A^2).$$

(Proof: See [1184, p. 176].)

Fact 3.16.4. Let $A \in \mathbb{R}^{n \times n}$ be tripotent. Then,

$$\text{rank } A = \text{rank } A^2 = \text{tr } A^2.$$

Fact 3.16.5. If $A, B \in \mathbb{F}^{n \times n}$ are idempotent and $AB = 0$, then $A + BA_\perp$ is idempotent and $C \triangleq A - B$ is tripotent. Conversely, if $C \in \mathbb{F}^{n \times n}$ is tripotent, then $A \triangleq \frac{1}{2}(C^2 + C)$ and $B \triangleq \frac{1}{2}(C^2 - C)$ are idempotent and satisfy $C = A - B$ and $AB = BA = 0$. (Proof: See [987, p. 114].)

3.17 Facts on Nilpotent Matrices

Fact 3.17.1. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $\mathcal{R}(A) = \mathcal{N}(A)$.
- ii) A is similar to a block-diagonal matrix each of whose diagonal blocks is N_2 .

(Proof: To prove $i) \implies ii)$, let $S \in \mathbb{F}^{n \times n}$ transform A into its Jordan form. Then, it follows from Fact 2.10.2 that $\mathcal{R}(SAS^{-1}) = S\mathcal{R}(AS^{-1}) = S\mathcal{R}(A) = S\mathcal{N}(A) = S\mathcal{N}(AS^{-1}S) = \mathcal{N}(AS^{-1}) = \mathcal{N}(SAS^{-1})$. The only Jordan block J that satisfies $\mathcal{R}(J) = \mathcal{N}(J)$ is $J = N_2$. Using $\mathcal{R}(N_2) = \mathcal{N}(N_2)$ and reversing these steps yields the converse result.) (Remark: The fact that n is even follows from $\text{rank } A + \text{def } A = n$ and $\text{rank } A = \text{def } A$.) (Remark: See Fact 3.17.2 and Fact 3.17.3.)

Fact 3.17.2. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $\mathcal{N}(A) \subseteq \mathcal{R}(A)$.
- ii) A is similar to a block-diagonal matrix each of whose diagonal blocks is either nonsingular or N_2 .

(Remark: See Fact 3.17.1 and Fact 3.17.3.)

Fact 3.17.3. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $\mathcal{R}(A) \subseteq \mathcal{N}(A)$.
- ii) A is similar to a block-diagonal matrix each of whose diagonal blocks is either zero or N_2 .

(Remark: See Fact 3.17.1 and Fact 3.17.2.)

Fact 3.17.4. Let $n \in \mathbb{P}$ and $k \in \{0, \dots, n\}$. Then, $\text{rank } N_n^k = n - k$.

Fact 3.17.5. Let $A \in \mathbb{R}^{n \times n}$. Then, $\text{rank } A^k$ is a nonincreasing function of $k \geq 1$. Furthermore, if there exists $k \in \{1, \dots, n\}$ such that $\text{rank } A^{k+1} = \text{rank } A^k$,

then $\text{rank } A^l = \text{rank } A^k$ for all $l \geq k$. Finally, if A is nilpotent and $A^l \neq 0$, then $\text{rank } A^{k+1} < \text{rank } A^k$ for all $k = 1, \dots, l$.

Fact 3.17.6. Let $A \in \mathbb{F}^{n \times n}$. Then, A is nilpotent if and only if, for all $k = 1, \dots, n$, $\text{tr } A^k = 0$. (Proof: See [1098, p. 103] or use Fact 4.8.2 with $p = \chi_A$ and $\mu_1 = \dots = \mu_n = 0$.)

Fact 3.17.7. Let $\lambda \in \mathbb{F}$ and $n, k \in \mathbb{P}$. Then,

$$(\lambda I_n + N_n)^k = \begin{cases} \lambda^k I_n + \binom{k}{1} \lambda^{k-1} N_n + \dots + \binom{k}{k} N_n^k, & k < n-1, \\ \lambda^k I_n + \binom{k}{1} \lambda^{k-1} N_n + \dots + \binom{k}{n-1} \lambda^{k-n+1} N_n^{n-1}, & k \geq n-1, \end{cases}$$

that is, for $k \geq n-1$,

$$\begin{bmatrix} \lambda & 1 & \cdots & 0 & 0 \\ 0 & \lambda & \ddots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \ddots & \lambda & 1 \\ 0 & 0 & \cdots & 0 & \lambda \end{bmatrix}^k = \begin{bmatrix} \lambda^k & \binom{k}{1} \lambda^{k-1} & \cdots & \binom{k}{n-2} \lambda^{k-n+1} & \binom{k}{n-1} \lambda^{k-n+1} \\ 0 & \lambda^k & \ddots & \binom{k}{n-3} \lambda^{k-n+2} & \binom{k}{n-2} \lambda^{k-n+2} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \ddots & \lambda^k & \binom{k}{1} \lambda^{k-1} \\ 0 & 0 & \cdots & 0 & \lambda^k \end{bmatrix}.$$

Fact 3.17.8. Let $A \in \mathbb{R}^{n \times n}$, assume that A is nilpotent, and let $k \geq 1$ be such that $A^k = 0$. Then,

$$\det(I - A) = 1$$

and

$$(I - A)^{-1} = \sum_{i=0}^{k-1} A^i.$$

Fact 3.17.9. Let $A, B \in \mathbb{F}^{n \times n}$, assume that B is nilpotent, and assume that $AB = BA$. Then, $\det(A + B) = \det A$. (Proof: Use Fact 5.17.4.)

Fact 3.17.10. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A and B are nilpotent, and assume that $AB = BA$. Then, $A + B$ is nilpotent. (Proof: If $A^k = B^l = 0$, then $(A + B)^{k+l} = 0$.)

Fact 3.17.11. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are either both upper triangular or both lower triangular. Then,

$$[A, B]^n = 0.$$

Hence, $[A, B]$ is nilpotent. (Remark: See [499, 500].) (Remark: See Fact 5.17.6.)

Fact 3.17.12. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $[A, [A, B]] = 0$. Then, $[A, B]$ is nilpotent. (Remark: This result is due to Jacobson. See [492] or [709, p. 98].)

Fact 3.17.13. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that there exist $k \in \mathbb{P}$ and nonzero $\alpha \in \mathbb{R}$ such that $[A^k, B] = \alpha A$. Then, A is nilpotent. (Proof: For all $l \in \mathbb{N}$,

$A^{k+l}B - A^lBA^k = \alpha A^{l+1}$, and thus $\text{tr } A^{l+1} = 0$. The result now follows from Fact 3.17.6.) (Remark: See [1145].)

3.18 Facts on Hankel and Toeplitz Matrices

Fact 3.18.1. Let $A \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

- i) If A is Toeplitz, then $\hat{I}A$ and $A\hat{I}$ are Hankel.
- ii) If A is Hankel, then $\hat{I}A$ and $A\hat{I}$ are Toeplitz.
- iii) A is Toeplitz if and only if $\hat{I}A\hat{I}$ is Toeplitz.
- iv) A is Hankel if and only if $\hat{I}A\hat{I}$ is Hankel.

Fact 3.18.2. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hankel, and consider the following conditions:

- i) A is Hermitian.
- ii) A is real.
- iii) A is symmetric.

Then, $i) \implies ii) \implies iii)$.

Fact 3.18.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is a partitioned matrix, each of whose blocks is a $k \times k$ (circulant, Hankel, Toeplitz) matrix. Then, A is similar to a block-(circulant, Hankel, Toeplitz) matrix. (Proof: See [140].)

Fact 3.18.4. For all $i, j = 1, \dots, n$, define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{1}{i+j-1}.$$

Then, A is Hankel, positive definite, and

$$\det A = \frac{[1!2! \cdots (n-1)!]^4}{1!2! \cdots (2n-1)!}.$$

Furthermore, for all $i, j = 1, \dots, n$, A^{-1} has integer entries given by

$$(A^{-1})_{(i,j)} = (-1)^{i+j}(i+j-1) \binom{n+i-1}{n-j} \binom{n+j-1}{n-i} \binom{i+j-2}{i-1}^2.$$

Finally, for large n ,

$$\det A \approx 2^{-2n^2}.$$

(Remark: A is the *Hilbert matrix*, which is a Cauchy matrix. See [681, p. 513], Fact 1.10.36, Fact 3.20.14, Fact 3.20.15, and Fact 12.21.18.) (Remark: See [325].)

Fact 3.18.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Toeplitz. Then, A is reverse symmetric.

Fact 3.18.6. Let $A \in \mathbb{F}^{n \times n}$. Then, A is Toeplitz if and only if there exist $a_0, \dots, a_n \in \mathbb{F}$ and $b_1, \dots, b_n \in \mathbb{F}$ such that

$$A = \sum_{i=1}^n b_i N_n^{iT} + \sum_{i=0}^n a_i N_n^i.$$

Fact 3.18.7. Let $A \in \mathbb{F}^{n \times n}$, let $k \geq 1$, and assume that A is (lower triangular, strictly lower triangular, upper triangular, strictly upper triangular). Then, so is A^k . If, in addition, A is Toeplitz, then so is A^k . (Remark: If A is Toeplitz, then A^2 is not necessarily Toeplitz.) (Remark: See Fact 11.13.1.)

3.19 Facts on Hamiltonian and Symplectic Matrices

Fact 3.19.1. Let $A \in \mathbb{F}^{2n \times 2n}$. Then, A is Hamiltonian if and only if $JA = (JA)^T$. Furthermore, A is symplectic if and only if $A^T J A = J$.

Fact 3.19.2. Assume that $n \in \mathbb{P}$ is even, let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hamiltonian and symplectic. Then, A is skew involutory. (Remark: See Fact 3.19.3.)

Fact 3.19.3. The following statements hold:

- i) I_{2n} is orthogonal, shifted orthogonal, a projector, a reflector, and symplectic.
- ii) J_{2n} is skew symmetric, orthogonal, skew involutory, a skew reflector, symplectic, and Hamiltonian.
- iii) $\hat{I}_{2n}$ is symmetric, orthogonal, involutory, shifted orthogonal, a projector, a reflector, and Hamiltonian.

(Remark: See Fact 3.19.2 and Fact 5.9.25.)

Fact 3.19.4. Let $A \in \mathbb{F}^{2n \times 2n}$, assume that A is Hamiltonian, and let $S \in \mathbb{F}^{2n \times 2n}$ be symplectic. Then, SAS^{-1} is Hamiltonian.

Fact 3.19.5. Let $A \in \mathbb{F}^{2n \times 2n}$, and assume that A is Hamiltonian and nonsingular. Then, A^{-1} is Hamiltonian.

Fact 3.19.6. Let $\mathcal{A} \in \mathbb{F}^{2n \times 2n}$. Then, $\mathcal{A}$ is Hamiltonian if and only if there exist $A, B, C, D \in \mathbb{F}^{n \times n}$ such that B and C are symmetric and

$$\mathcal{A} = \begin{bmatrix} A & B \\ C & -A^T \end{bmatrix}.$$

(Remark: See Fact 4.9.23.)

Fact 3.19.7. Let $A \in \mathbb{F}^{2n \times 2n}$, and assume that A is Hamiltonian. Then, $\text{tr } A = 0$.

Fact 3.19.8. Let $\mathcal{A} \in \mathbb{F}^{2n \times 2n}$. Then, $\mathcal{A}$ is skew symmetric and Hamiltonian if and only if there exist a skew-symmetric matrix $A \in \mathbb{F}^{n \times n}$ and a symmetric matrix $B \in \mathbb{F}^{n \times n}$ such that

$$\mathcal{A} = \begin{bmatrix} A & B \\ -B & A \end{bmatrix}.$$

Fact 3.19.9. Let $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{F}^{2n \times 2n}$, where $A, B, C, D \in \mathbb{F}^{n \times n}$. Then, $\mathcal{A}$ is symplectic if and only if $A^T C$ and $B^T D$ are symmetric and $A^T D - C^T B = I$.

Fact 3.19.10. Let $A \in \mathbb{F}^{2n \times 2n}$, and assume that A is symplectic. Then, $\det A = 1$. (Proof: Using Fact 2.14.16 and Fact 3.19.9 it follows that $\det \mathcal{A} = \det(A^T D - C^T B) = \det I = 1$. See also [103, p. 27], [423], [624, p. 8], or [1186, p. 128].)

Fact 3.19.11. Let $A \in \mathbb{F}^{2 \times 2}$. Then, A is symplectic if and only if $\det A = 1$. Hence, $\text{SL}_{\mathbb{F}}(2) = \text{Symp}_{\mathbb{F}}(2)$.

Fact 3.19.12. The following statements hold:

- i) If $A \in \mathbb{F}^{2n \times 2n}$ is Hamiltonian and $A + I$ is nonsingular, then $B \triangleq (A - I)(A + I)^{-1}$ is symplectic, $I - B$ is nonsingular, and $(I - B)^{-1} = \frac{1}{2}(A + I)$.
- ii) If $B \in \mathbb{F}^{2n \times 2n}$ is symplectic and $I - B$ is nonsingular, then $A = (I + B)(I - B)^{-1}$ is Hamiltonian, $A + I$ is nonsingular, and $(A + I)^{-1} = \frac{1}{2}(I - B)$.
- iii) If $A \in \mathbb{F}^{2n \times 2n}$ is Hamiltonian, then there exists a unique symplectic matrix $B \in \mathbb{F}^{2n \times 2n}$ such that $I - B$ is nonsingular and $A = (I + B)(I - B)^{-1}$. In fact, $B = (A - I)(A + I)^{-1}$.
- iv) If $B \in \mathbb{F}^{2n \times 2n}$ is symplectic and $I - B$ is nonsingular, then there exists a unique Hamiltonian matrix $A \in \mathbb{F}^{2n \times 2n}$ such that $B = (A - I)(A + I)^{-1}$. In fact, $A = (I + B)(I - B)^{-1}$.

(Remark: See Fact 3.11.28, Fact 3.11.29, and Fact 3.11.30.)

Fact 3.19.13. Let $\mathcal{A} \in \mathbb{R}^{2n \times 2n}$. Then, $\mathcal{A} \in \text{osymp}_{\mathbb{R}}(2n)$ if and only if there exist $A, B \in \mathbb{R}^{n \times n}$ such that A is skew symmetric, B is symmetric, and $\mathcal{A} = \begin{bmatrix} A & B \\ -B & A \end{bmatrix}$. (Proof: See [395].) (Remark: $\text{OSymp}_{\mathbb{R}}(2n)$ is the *orthosymplectic group*.)

3.20 Facts on Miscellaneous Types of Matrices

Fact 3.20.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that there exists $i \in \{1, \dots, n\}$ such that either $\text{row}_i(A) = 0$ or $\text{col}_i(A) = 0$. Then, A is reducible.

Fact 3.20.2. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is reducible. Then, A has at least $n - 1$ entries that are equal to zero.

Fact 3.20.3. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is a permutation matrix. Then, A is irreducible if and only if there exists a permutation matrix $S \in \mathbb{R}^{n \times n}$ such that SAS^{-1} is the primary circulant. (Proof: See [1184, p. 177].) (Remark: The primary circulant is defined in Fact 5.16.7.)

Fact 3.20.4. Let $A \in \mathbb{F}^{n \times n}$. Then, A is reducible if and only if $|A|$ is reducible. Furthermore, A is irreducible if and only if $|A|$ is irreducible.

Fact 3.20.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $l \in \{0, \dots, n\}$ and $k \in \{1, \dots, n\}$. Then, the following statements are equivalent:

- i) Every submatrix B of A whose entries are entries of A lying above the l th superdiagonal of A satisfies $\text{rank } B \leq k - 1$.
- ii) Every submatrix C of A whose entries are entries of A^{-1} lying above the l th subdiagonal of A^{-1} satisfies $\text{rank } C \leq l + k - 1$.

Specifically, the following statements hold:

- iii) A is lower triangular if and only if A^{-1} is lower triangular.
- iv) A is diagonal if and only if A^{-1} is diagonal.
- v) A is lower Hessenberg if and only if every submatrix C of A^{-1} whose entries are entries of A^{-1} lying on or above the diagonal of A^{-1} satisfies $\text{rank } C \leq 1$.
- vi) A is tridiagonal if and only if every submatrix C of A^{-1} whose entries are entries of A^{-1} lying on or above the diagonal of A^{-1} satisfies $\text{rank } C \leq 1$ and every submatrix C of A^{-1} whose entries are entries of A^{-1} lying on or below the diagonal of A^{-1} satisfies $\text{rank } C \leq 1$.

(Remark: The 0th subdiagonal and the 0th superdiagonal are the diagonal.) (Proof: See [1242].) (Remark: Statement iii) corresponds to $l = 0$ and $k = 1$, iv) corresponds to $l = 0$ and $k = 1$ applied to A and A^T , v) corresponds to $l = 1$ and $k = 1$, and vi) corresponds to $l = 1$ and $k = 1$ applied to A and A^T . (Remark: See Fact 2.11.20.) (Remark: Extensions to generalized inverses are considered in [131, 1131].)

Fact 3.20.6. Let $A \in \mathbb{F}^{n \times n}$ be the tridiagonal matrix

$$A \triangleq \begin{bmatrix} a+b & ab & 0 & \cdots & 0 & 0 \\ 1 & a+b & ab & \cdots & 0 & 0 \\ 0 & 1 & a+b & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & a+b & ab \\ 0 & 0 & 0 & \cdots & 1 & a+b \end{bmatrix}.$$

Then,

$$\det A = \begin{cases} (n+1)a^n, & a = b, \\ \frac{a^{n+1} - b^{n+1}}{a - b}, & a \neq b. \end{cases}$$

(Proof: See [841, pp. 401, 621].)

Fact 3.20.7. Let $A \in \mathbb{R}^{n \times n}$ be the tridiagonal, Toeplitz matrix

$$A \triangleq \begin{bmatrix} b & c & 0 & \cdots & 0 & 0 \\ a & b & c & \cdots & 0 & 0 \\ 0 & a & b & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & b & c \\ 0 & 0 & 0 & \cdots & a & b \end{bmatrix},$$

and define

$$\alpha \triangleq \frac{1}{2}(b + \sqrt{b^2 - 4ac}), \quad \beta \triangleq \frac{1}{2}(b - \sqrt{b^2 - 4ac}).$$

Then,

$$\det A = \begin{cases} b^n, & ac = 0, \\ (n+1)(b/2)^n, & b^2 = 4ac, \\ (\alpha^{n+1} - \beta^{n+1})/(\alpha - \beta), & b^2 \neq 4ac. \end{cases}$$

(Proof: See [1490, pp. 101, 102].) (Remark: See Fact 3.20.6 and Fact 5.11.43.)

Fact 3.20.8. Let $A \in \mathbb{R}^{n \times n}$, assume that A is tridiagonal with positive diagonal entries, and assume that, for all $i = 2, \dots, n$,

$$A_{(i,i-1)}A_{(i-1,i)} < \frac{1}{4} \left(\cos \frac{\pi}{n+1} \right)^{-2} A_{(i,i)}A_{(i-1,i-1)}.$$

Then, $\det A > 0$. If, in addition, A is symmetric, then A is positive definite. (Proof: See [766].) (Remark: Related results are given in [324].) (Remark: See Fact 8.8.18.)

Fact 3.20.9. Let $A \in \mathbb{R}^{n \times n}$, assume that A is tridiagonal, assume that every entry of the superdiagonal and subdiagonal of A is nonzero, assume that every leading principal subdeterminant of A and every trailing principal subdeterminant of A is nonzero. Then, every entry of A^{-1} is nonzero. (Proof: See [700].)

Fact 3.20.10. Define $A \in \mathbb{R}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} 2 & -1 & 0 & \cdots & 0 & 0 \\ -1 & 2 & -1 & \cdots & 0 & 0 \\ 0 & -1 & 2 & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & 2 & -1 \\ 0 & 0 & 0 & \cdots & -1 & 1 \end{bmatrix}.$$

Then,

$$A^{-1} = \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & 2 & 2 & \cdots & 2 & 2 \\ 1 & 2 & 3 & \ddots & 3 & 3 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 1 & 2 & 3 & \ddots & n-1 & n-1 \\ 1 & 2 & 3 & \cdots & n-1 & n \end{bmatrix}.$$

(Proof: See [1184, p. 182], where the (n, n) entry of A is incorrect.) (Remark: See Fact 3.20.9.)

Fact 3.20.11. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and assume that $A_{(2,2)}, \dots, A_{(n-1,n-1)}$ are nonzero. Then, A^{-1} is tridiagonal if and only if, for all $i, j = 1, \dots, n$ such that $|i - j| \geq 2$, and for all k satisfying $\min\{i, j\} < k < \max\{i, j\}$, it follows that

$$A_{(i,j)} = \frac{A_{(i,k)}A_{(k,j)}}{A_{(k,k)}}.$$

(Proof: See [147].)

Fact 3.20.12. Let $A \in \mathbb{F}^{n \times m}$. Then, A is (semicontractive, contractive) if and only if A^* is.

Fact 3.20.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is dissipative. Then, A is nonsingular. (Proof: Suppose that A is singular, and let $x \in \mathcal{N}(A)$. Then, $x^*(A + A^*)x = 0$.) (Remark: If $A + A^*$ is nonsingular, then A is not necessarily nonsingular. Consider $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.)

Fact 3.20.14. Let $a_1, \dots, a_n, b_1, \dots, b_n \in \mathbb{R}$, assume that $a_i + b_j \neq 0$ for all $i, j = 1, \dots, n$, and, for all $i, j = 1, \dots, n$, define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{1}{a_i + b_j}.$$

Then,

$$\det A = \frac{\prod_{1 \leq i < j \leq n} (a_j - a_i)(b_j - b_i)}{\prod_{1 \leq i, j \leq n} (a_i + b_j)}.$$

Now, assume that $a_1, \dots, a_n$ are distinct and $b_1, \dots, b_n$ are distinct. Then, A is nonsingular and

$$(A^{-1})_{(i,j)} = \frac{\prod_{1 \leq k \leq n} (a_j + b_k)(a_k + b_i)}{(a_j + b_i) \prod_{\substack{1 \leq k \leq n \\ k \neq j}} (a_j - a_k) \prod_{\substack{1 \leq k \leq n \\ k \neq i}} (b_i - b_k)}.$$

Furthermore,

$$1_{1 \times n} A^{-1} 1_{n \times 1} = \sum_{i=1}^n (a_i + b_i).$$

(Remark: A is a *Cauchy matrix*. See [199], [681, p. 515], Fact 3.18.4, Fact 3.20.15, and Fact 12.21.18.)

Fact 3.20.15. Let $x_1, \dots, x_n$ be distinct positive numbers, let $y_1, \dots, y_n$ be distinct positive numbers, and let $A \in \mathbb{R}^{n \times n}$, where, for all $i, j = 1, \dots, n$,

$$A_{(i,j)} \triangleq \frac{1}{x_i + y_j}.$$

Then, A is nonsingular. (Proof: See [854].) (Remark: A is a Cauchy matrix. See Fact 3.18.4, Fact 3.20.14, and Fact 12.21.18.)

Fact 3.20.16. Let $A \in \mathbb{F}^{n \times m}$. Then, A is centrosymmetric if and only if $A^T = A^{\tilde{}}$. Furthermore, A is centrohermitian if and only if $A^* = A^{\tilde{}}$.

Fact 3.20.17. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. If A and B are both (centrohermitian, centrosymmetric), then so is AB . (Proof: See [685].)

Fact 3.20.18. Let $A, B \in \mathbb{F}^n$, and assume that A and B are (upper triangular, lower triangular). Then, AB is (upper triangular, lower triangular). If, in addition, either A or B is (strictly upper triangular, strictly lower triangular), then AB is (strictly upper triangular, strictly lower triangular). (Remark: See Fact 3.21.5.)

3.21 Facts on Groups

Fact 3.21.1. The following subsets of $\mathbb{R}$ are groups:

- i) $\{x \in \mathbb{R}: x \neq 0\}$.
- ii) $\{x \in \mathbb{R}: x > 0\}$.
- iii) $\{x \in \mathbb{R}: x \neq 0 \text{ and } x \text{ is rational}\}$.
- iv) $\{x \in \mathbb{R}: x > 0 \text{ and } x \text{ is rational}\}$.
- v) $\{-1, 1\}$.
- vi) $\{1\}$.

Fact 3.21.2. Let n be a nonnegative integer, and define $S^n \triangleq \{x \in \mathbb{R}^{n+1}: x^T x = 1\}$, which is the unit sphere in $\mathbb{R}^{n+1}$. Then, the following statements hold:

- i) $\text{SO}(1) = \text{SU}(1) = \{1\}$.
- ii) $S^0 = \text{O}(1) = \{-1, 1\}$.
- iii) $\{1, -1, j, -j\}$.
- iv) $\text{U}(1) = \{e^{j\theta}: \theta \in [0, 2\pi)\} \approx \text{SO}(2)$.

- v) $S^1 = \{ \begin{bmatrix} \cos \theta & \sin \theta \end{bmatrix}^T \in \mathbb{R}^2: \theta \in [0, 2\pi) \} = \{ \begin{bmatrix} \operatorname{Re} z & \operatorname{Im} z \end{bmatrix}^T: z \in U(1) \}.$
- vi) $SU(2) = \{ \begin{bmatrix} z & w \\ -\bar{w} & \bar{z} \end{bmatrix} \in \mathbb{C}^{2 \times 2}: z, w \in \mathbb{C} \text{ and } |z|^2 + |w|^2 = 1 \} \approx Sp(1).$
- vii) $S^3 = \{ \begin{bmatrix} \operatorname{Re} z & \operatorname{Im} z & \operatorname{Re} w & \operatorname{Im} w \end{bmatrix}^T \in \mathbb{R}^4: \begin{bmatrix} z & w \end{bmatrix}^T \in \mathbb{C}^2 \text{ and } |z|^2 + |w|^2 = 1 \}.$

(Proof: See [1256, p. 40].) (Remark: $Sp(1) \subset \mathbb{H}^{1 \times 1}$ is the group of unit quaternions. See Fact 3.22.1.) (Remark: A group operation can be defined on S^n if and only if $n = 0, 1$, or 3 . See [1256, p. 40].)

Fact 3.21.3. The groups $U(n)$ and $O(2n) \cap \operatorname{Symp}_{\mathbb{R}}(2n)$ are isomorphic. In particular, $U(1)$ and $O(2) \cap \operatorname{Symp}_{\mathbb{R}}(2) = SO(2)$ are isomorphic. (Proof: See [97].)

Fact 3.21.4. The following subsets of $\mathbb{F}^{n \times n}$ are Lie algebras:

- i) $\mathfrak{ut}(n) \triangleq \{A \in \mathfrak{gl}_{\mathbb{F}}(n): A \text{ is upper triangular}\}.$
- ii) $\mathfrak{sut}(n) \triangleq \{A \in \mathfrak{gl}_{\mathbb{F}}(n): A \text{ is strictly upper triangular}\}.$
- iii) $\{0_{n \times n}\}.$

Fact 3.21.5. The following subsets of $\mathbb{F}^{n \times n}$ are groups:

- i) $UT(n) \triangleq \{A \in GL_{\mathbb{F}}(n): A \text{ is upper triangular}\}.$
- ii) $UT_+(n) \triangleq \{A \in UT(n): A_{(i,i)} > 0 \text{ for all } i = 1, \dots, n\}.$
- iii) $UT_{\pm 1}(n) \triangleq \{A \in UT(n): A_{(i,i)} = \pm 1 \text{ for all } i = 1, \dots, n\}.$
- iv) $SUT(n) \triangleq \{A \in UT(n): A_{(i,i)} = 1 \text{ for all } i = 1, \dots, n\}.$
- v) $\{I_n\}.$

(Remark: The matrices in $SUT(n)$ are unipotent. See Fact 5.15.9.) (Remark: $SUT(3)$ for $\mathbb{F} = \mathbb{R}$ is the *Heisenberg group*.) (Remark: See Fact 3.20.18.)

Fact 3.21.6. Let $P \in \mathbb{R}^{n \times n}$, and assume that P is a permutation matrix. Then, there exist transposition matrices $T_1, \dots, T_k \in \mathbb{R}^{n \times n}$ such that

$$P = T_1 \cdots T_k.$$

(Remark: The permutation matrix T_i is a *transposition matrix* if it has exactly two off-diagonal entries that are nonzero.) (Remark: Every permutation of n objects can be realized as a sequence of pairwise transpositions. See [445, pp. 106, 107] or [497, p. 82].) (Example:

$$\begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

which represents a 3-cycle.) (Remark: This factorization in terms of transpositions is not unique. However, Fact 5.16.8 shows that every permutation can be written essentially uniquely as a product of disjoint cycles.)

Fact 3.21.7. The following subsets of $\mathbb{R}^{n \times n}$ are finite groups:

- i) $P(n) \triangleq \{A \in \text{GL}_{\mathbb{R}}(n): A \text{ is a permutation matrix}\}.$
- ii) $\text{SP}(n) \triangleq \{A \in P(n): \det A = 1\}.$

Furthermore, let k be a positive integer, and define $R, S \in \mathbb{R}^{2 \times 2}$ by

$$R \triangleq \begin{bmatrix} \cos \frac{2\pi}{k} & \sin \frac{2\pi}{k} \\ -\sin \frac{2\pi}{k} & \cos \frac{2\pi}{k} \end{bmatrix}, \quad S \triangleq \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = \hat{I}_2.$$

Then, $R^k = S^2 = I_2$, and the following subsets of $\mathbb{R}^{2 \times 2}$ are finite groups:

- iii) $\text{O}_k(2) \triangleq \{I, R, \dots, R^{k-1}, S, SR, \dots, SR^{k-1}\}.$
- iv) $\text{SO}_k(2) \triangleq \{I, R, \dots, R^{k-1}\}.$

Finally, the cardinality of $P(n)$, $\text{SP}(n)$, $\text{O}_k(2)$, and $\text{SO}_k(2)$ is $n!$, $\frac{1}{2}n!$, $2k$, and k , respectively. (Remark: The elements of $P(n)$ permute n -tuples arbitrarily, while the elements of $\text{SP}(n)$ permute n -tuples evenly. See Fact 5.16.8. The elements of $\text{SO}_k(2)$ perform counterclockwise rotations of planar figures by the angle $2\pi/k$ about a line perpendicular to the plane and passing through 0, while the elements of $\text{O}_k(2)$ perform the rotations of $\text{SO}_k(2)$ and reflect planar figures across the line $y = x$. See [445, pp. 41, 845].) (Remark: These groups are matrix representations of *symmetry groups*, which are groups of transformations that map a set onto itself. Specifically, $P(k)$, $\text{SP}(k)$, $\text{O}_k(2)$, and $\text{SO}_k(2)$, are matrix representations of the *permutation group* S_k , the *alternating group* A_k , the *dihedral group* D_k , and the *cyclic group* C_k , respectively, all of which can be viewed as abstract groups having matrix representations. Matrix representations of groups are discussed in [520].) (Remark: An *abstract group* is a collection of objects (not necessarily matrices) that satisfy the properties of a group as defined by Definition 3.3.3.) (Remark: Every finite subgroup of $\text{O}(2)$ is a representation of either D_k or C_k for some k . Furthermore, every finite subgroup of $\text{SO}(3)$ is a representation of either D_k or C_k for some k or A_4 , S_4 , or A_5 . The symmetry groups A_4 , S_4 , and A_5 are represented by bijective transformations of regular solids. Specifically, A_4 is represented by the *tetrahedral group*, which consists of 12 rotation matrices that map a regular tetrahedron onto itself; S_4 is represented by the *octahedral group*, which consists of 24 rotation matrices that map an octahedron or a cube onto itself; and A_5 is represented by the *icosahedral group*, which consists of 60 rotation matrices that map a regular icosahedron or a regular dodecahedron onto itself. The 12 elements of the tetrahedral group representing A_4 are given by DR^k , where $D \in \{I_3, \text{diag}(1, -1, -1), \text{diag}(-1, -1, 1), \text{diag}(-1, 1, -1)\}$, $R \triangleq \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$, and $k = 0, 1, 2$. The 24 elements of the octahedral group representing S_4 are given by the 3×3 signed permutation matrices with determinant 1, where a *signed permutation matrix* has exactly one nonzero entry, which is either 1 or -1 , in each row and column. See [75, p. 184], [346, p. 32], [571, pp. 176–193], [603, pp. 9–23], [1149, p. 69], [1187, pp. 35–43], or [1256, pp. 45–47].) (Remark: The dihedral group D_2 is also called the *Klein four group*.) (Remark: The permutation group S_k is not Abelian for all $k \geq 3$. The alternating group A_3 is Abelian, whereas A_k is not Abelian for all $k \geq 4$. A_5 is essential to the classical result of Abel and Galois that there exist fifth-order polynomials whose roots cannot be expressed in terms of radicals involving the coefficients. Two such polynomials are $p(x) = x^5 - x - 1$ and $p(x) = x^5 - 16x + 2$. See [75, p. 574] and [445, pp. 32, 625–639].)

Fact 3.21.8. The following sets of matrices are groups:

- i) $P(2) = O_1(2) = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \right\}$.
- ii) $SO_2(2) = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \right\}$.
- iii) $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \right\}$.
- iv) $SP(3) = \{I_3, P_3, P_3^2\}$, where $P_3 \triangleq \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$.
- v) $O_2(2) = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \right\}$.
- vi) $\{I_4, P_4, P_4^2, P_4^3\}$, where $P_4 \triangleq \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \end{bmatrix}$.
- vii) $P(3) = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \right\}$.
- viii) $\left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ -1 & 1 \end{bmatrix}, \begin{bmatrix} -1 & -1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} -1 & -1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix} \right\}$.
- ix) For all $k \geq 0$, $SU_k(1) \triangleq \{1, e^{2\pi j/k}, e^{4\pi j/k}, \dots, e^{2(k-1)\pi j/k}\}$.
- x) $\{I, P_k, P_k^2, \dots, P_k^{k-1}\}$.

(Remark: i), ii), and iii) are representations of the cyclic group C_2 , which is identical to the permutation group S_2 and the dihedral group D_1 ; iv) is a representation of the cyclic group C_3 , which is identical to alternating group A_3 ; v) is a representation of the dihedral group D_2 , which is also called the Klein four group, see Fact 3.21.7; vi) is a representation of the cyclic group C_4 ; vii) is a representation of the permutation group S_3 , which is identical to the dihedral group D_3 , with $A^2 = B^3 = (AB)^2 = I_3$, where $A \triangleq \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ and $B \triangleq \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}$; viii) is a representation of the dihedral group D_3 , where $\begin{bmatrix} 0 & 1 \\ -1 & -1 \end{bmatrix}^3 = I_2$; ix) is a matrix representation of the cyclic group C_k and its real representation $SO_k(2)$; x) is a matrix representation of the cyclic group C_k , where P_k is the $k \times k$ primary circulant defined in Fact 5.16.7. The groups $P(n)$ and $SP(n)$ are defined in Fact 3.21.7. Representations of groups are discussed in [316, 631, 703].)

Fact 3.21.9. The following statements hold:

- i) There exists exactly one isomorphically distinct group consisting of one element. A representation is $\{I_n\}$.
- ii) There exists exactly one isomorphically distinct group consisting of two elements, namely, the cyclic group C_2 , which is identical to the permutation group S_2 and the dihedral group D_1 . Representations of C_2 are given by $P(2)$, $O_1(2)$, $SO_2(2)$, and $SU_2(1) = \{1, -1\}$.
- iii) There exists exactly one isomorphically distinct group consisting of three elements, namely, the cyclic group C_3 , which is identical to the alternating group A_3 . Representations of C_3 are given by $SP(3)$, $SO_3(2)$, $SU_3(1)$, and $\{I_3, P_3, P_3^2\}$.
- iv) There exist exactly two isomorphically distinct groups consisting of four elements, namely, the cyclic group C_4 and the dihedral group D_2 . Representations of C_4 are given by $SO_4(2)$ and $SU_4(1) = \{1, -1, j, -j\}$. A

representation of D_2 is given by $O_2(2)$.

- v) There exists exactly one isomorphically distinct group consisting of five elements, namely, the cyclic group C_5 . Representations of C_5 are given by $SO_5(2)$, $SU_5(1)$, and $\{I_5, P_5, P_5^2, P_5^3, P_5^4\}$.
- vi) There exist exactly two isomorphically distinct groups consisting of six elements, namely, the cyclic group C_6 and the dihedral group D_3 , which is identical to S_3 . Representations of C_6 are given by $SO_6(2)$, $SU_6(1)$, and $\{I_6, P_6, P_6^2, P_6^3, P_6^4, P_6^5\}$. Representations of D_3 are given by $P(3)$ and $O_3(2)$.
- vii) There exists exactly one isomorphically distinct group consisting of seven elements, namely, the cyclic group C_7 . Representations of C_7 are given by $SO_7(2)$, $SU_7(1)$, and $\{I_7, P_7, P_7^2, P_7^3, P_7^4, P_7^5, P_7^6\}$.
- viii) There exist exactly five isomorphically distinct groups consisting of eight elements, namely, C_8 , $D_2 \times C_2$, $C_4 \times C_2$, D_4 , and the quaternion group $\{\pm 1, \pm \hat{i}, \pm \hat{j}, \pm \hat{k}\}$. Representations of C_8 are given by $SO_8(2)$, $SU_8(1)$, and $\{I_8, P_8, P_8^2, P_8^3, P_8^4, P_8^5, P_8^6, P_8^7\}$. A representation of D_4 is given by $O_8(2)$. Representations of the quaternion group are given by *ii*) of Fact 3.22.3 and *v*) of Fact 3.22.6.

(Proof: See [555, pp. 4–7].) (Remark: P_k is the $k \times k$ primary circulant defined in Fact 5.16.7.)

Fact 3.21.10. Let $S \subset \mathbb{F}^{n \times n}$, and assume that S is a group. Then, $\{A^T: A \in S\}$ and $\{\overline{A}: A \in S\}$ are groups.

Fact 3.21.11. Let $P \in \mathbb{F}^{n \times n}$, and define $S \triangleq \{A \in \mathbb{F}^{n \times n}: A^T P A = P\}$. Then, S is a group. If, in addition, P is nonsingular and skew symmetric, then, for every matrix $P \in S$, it follows that $\det P = 1$. (Proof: See [341].) (Remark: If $\mathbb{F} = \mathbb{R}$, n is even, and $P = J_n$, then $S = \text{Symp}_{\mathbb{R}}(n)$.) (Remark: Weaker conditions on P such that $\det P = 1$ for all $P \in S$ are given in [341].)

3.22 Facts on Quaternions

Fact 3.22.1. Let $\hat{i}, \hat{j}, \hat{k}$ satisfy

$$\hat{i}^2 = \hat{j}^2 = \hat{k}^2 = -1,$$

$$\hat{i}\hat{j} = \hat{k} = -\hat{j}\hat{i},$$

$$\hat{j}\hat{k} = \hat{i} = -\hat{k}\hat{j},$$

$$\hat{k}\hat{i} = \hat{j} = -\hat{i}\hat{k},$$

and define

$$\mathbb{H} \triangleq \{a + b\hat{i} + c\hat{j} + d\hat{k}: a, b, c, d \in \mathbb{R}\}.$$

Furthermore, for $a, b, c, d \in \mathbb{R}$, define $q \triangleq a + b\hat{i} + c\hat{j} + d\hat{k}$, $\overline{q} \triangleq a - b\hat{i} - c\hat{j} - d\hat{k}$, and $|q| \triangleq \sqrt{q\overline{q}} = \sqrt{a^2 + b^2 + c^2 + d^2} = |\overline{q}|$. Then,

$$qI_4 = UQ(q)U,$$

where

$$\mathcal{Q}(q) \triangleq \begin{bmatrix} a & -b & -c & -d \\ b & a & -d & c \\ c & d & a & -b \\ d & -c & b & a \end{bmatrix}$$

and

$$U \triangleq \frac{1}{2} \begin{bmatrix} 1 & \hat{i} & \hat{j} & \hat{k} \\ -\hat{i} & 1 & \hat{k} & -\hat{j} \\ -\hat{j} & -\hat{k} & 1 & \hat{i} \\ -\hat{k} & \hat{j} & -\hat{i} & 1 \end{bmatrix}$$

satisfies $U^2 = I_4$. In addition,

$$\det \mathcal{Q}(q) = (a^2 + b^2 + c^2 + d^2)^2.$$

Furthermore, if $|q| = 1$, then $\begin{bmatrix} a & -b & -c & -d \\ b & a & -d & c \\ c & d & a & -b \\ d & -c & b & a \end{bmatrix}$ is orthogonal. Next, for $i = 1, 2$, let

$a_i, b_i, c_i, d_i \in \mathbb{R}$, define $q_i \triangleq a_i + b_i \hat{i} + c_i \hat{j} + d_i \hat{k}$, and define

$$q_3 \triangleq q_2 q_1 = a_3 + b_3 \hat{i} + c_3 \hat{j} + d_3 \hat{k}.$$

Then,

$$\overline{q_3} = \overline{q_2} \overline{q_1},$$

$$|q_3| = |q_2 q_1| = |q_1 q_2| = |q_1 \overline{q_2}| = |\overline{q_1} q_2| = |\overline{q_1}| |q_2|,$$

$$\mathcal{Q}(q_3) = \mathcal{Q}(q_2) \mathcal{Q}(q_1),$$

and

$$\begin{bmatrix} a_3 \\ b_3 \\ c_3 \\ d_3 \end{bmatrix} = \mathcal{Q}(q_2) \begin{bmatrix} a_1 \\ b_1 \\ c_1 \\ d_1 \end{bmatrix}.$$

Next, for $i = 1, 2$, define $v_i \triangleq [b_i \ c_i \ d_i]^T$. Then,

$$\begin{bmatrix} a_3 \\ b_3 \\ c_3 \\ d_3 \end{bmatrix} = \begin{bmatrix} a_2 a_1 - v_2^T v_1 \\ a_1 v_2 + a_2 v_1 + v_2 \times v_1 \end{bmatrix}.$$

(Remark: q is a *quaternion*. See [477, pp. 287–294]. Note the analogy between $\hat{i}, \hat{j}, \hat{k}$ and the unit vectors in $\mathbb{R}^3$ under cross-product multiplication. See [103, p. 119].) (Remark: The group $\text{Sp}(1)$ of unit-length quaternions is isomorphic to $\text{SU}(2)$. See [362, p. 30], [1256, p. 40], and Fact 3.19.11.) (Remark: The unit-length quaternions, whose coefficients comprise the unit sphere $S^3 \subset \mathbb{R}^4$ and are called *Euler parameters*, provide a double cover of $\text{SO}(3)$ as shown by Fact 3.11.10. See [152, p. 380] and [26, 346, 850, 1195].) (Remark: An equivalent formulation of quaternion multiplication is given by Rodrigues's formulas. See Fact 3.11.11.) (Remark: Determinants of matrices with quaternion entries are discussed in [80] and [1256, p. 31].) (Remark: The *Clifford algebras* include the *quaternion algebra* $\mathbb{H}$ and the *octonion algebra* $\mathbb{O}$, which involves the *Cayley numbers*. See [477, pp.

295–300]. These ideas form the basis for *geometric algebra*. See [1217, p. 100] and [98, 346, 349, 364, 411, 425, 426, 477, 605, 607, 636, 670, 671, 672, 684, 831, 870, 934, 1098, 1185, 1250, 1256, 1279].)

Fact 3.22.2. Let $a, b, c, d \in \mathbb{R}$, and let $q \triangleq a + b\hat{i} + c\hat{j} + d\hat{k} \in \mathbb{H}$. Then,

$$q = a + b\hat{i} + (c + d\hat{i})\hat{j}.$$

(Remark: For all $q \in \mathbb{H}$, there exist $z, w \in \mathbb{C}$ such that $q = z + w\hat{j}$, where we interpret $\mathbb{C}$ as $\{a + b\hat{i} : a, b \in \mathbb{R}\}$. This observation is analogous to the fact that, for all $z \in \mathbb{C}$, there exist $a, b \in \mathbb{R}$ such that $z = a + bj$, where $j \triangleq \sqrt{-1}$. See [1256, p. 10].)

Fact 3.22.3. The following sets are groups:

- i) $\mathbb{Q} \triangleq \{\pm 1, \pm \hat{i}, \pm \hat{j}, \pm \hat{k}\}$.
- ii) $\text{GL}_{\mathbb{H}}(1) \triangleq \mathbb{H} \setminus \{0\} = \{a + b\hat{i} + c\hat{j} + d\hat{k} : a, b, c, d \in \mathbb{R} \text{ and } a^2 + b^2 + c^2 + d^2 > 0\}$.
- iii) $\text{Sp}(1) \triangleq \{a + b\hat{i} + c\hat{j} + d\hat{k} : a, b, c, d \in \mathbb{R} \text{ and } a^2 + b^2 + c^2 + d^2 = 1\}$.
- iv) $\mathbb{Q}_{\mathbb{R}} \triangleq \left\{ \pm I_4, \pm \begin{bmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{bmatrix}, \pm \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix}, \pm \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix} \right\}$.
- v) $\text{GL}_{\mathbb{H}, \mathbb{R}}(1) \triangleq \left\{ \begin{bmatrix} a & -b & -c & -d \\ b & a & -d & c \\ c & d & a & -b \\ d & -c & b & a \end{bmatrix} : a^2 + b^2 + c^2 + d^2 > 0 \right\}$.
- vi) $\text{GL}'_{\mathbb{H}, \mathbb{R}}(1) \triangleq \left\{ \begin{bmatrix} a & -b & -c & -d \\ b & a & -d & c \\ c & d & a & -b \\ d & -c & b & a \end{bmatrix} : a^2 + b^2 + c^2 + d^2 = 1 \right\}$.

Furthermore, $\mathbb{Q}$ and $\mathbb{Q}_{\mathbb{R}}$ are isomorphic, $\text{GL}_{\mathbb{H}}(1)$ and $\text{GL}_{\mathbb{H}, \mathbb{R}}(1)$ are isomorphic, $\text{Sp}(1)$ and $\text{GL}'_{\mathbb{H}, \mathbb{R}}(1)$ are isomorphic, and $\text{GL}'_{\mathbb{H}, \mathbb{R}}(1) \subset \text{SO}(4) \cap \text{Symp}_{\mathbb{R}}(4)$. (Remark: J_4 is an element of $\text{Symp}_{\mathbb{R}}(4) \cap \text{SO}(4)$ but is not contained in $\text{GL}'_{\mathbb{H}, \mathbb{R}}(1)$.) (Remark: See Fact 3.22.1.)

Fact 3.22.4. Define

$$\text{Sp}(n) \triangleq \{A \in \mathbb{H}^{n \times n} : A^*A = I\},$$

where $\mathbb{H}$ is the quaternion algebra, $A^* \triangleq \overline{A}^T$, and, for $q = a + b\hat{i} + c\hat{j} + d\hat{k} \in \mathbb{H}$, $\overline{q} \triangleq a - b\hat{i} - c\hat{j} - d\hat{k}$. Then, the groups $\text{Sp}(n)$ and $\text{U}(2n) \cap \text{Symp}_{\mathbb{C}}(2n)$ are isomorphic. In particular, $\text{Sp}(1)$ and $\text{U}(2) \cap \text{Symp}_{\mathbb{C}}(2) = \text{SU}(2)$ are isomorphic. (Proof: See [97].) (Remark: $\text{U}(n)$ and $\text{O}(2n) \cap \text{Symp}_{\mathbb{R}}(2n)$ are isomorphic.) (Remark: See Fact 3.22.3.)

Fact 3.22.5. Let n be a positive integer. Then, $\text{SO}(2n) \cap \text{Symp}_{\mathbb{R}}(2n)$ is a matrix group whose Lie algebra is $\text{so}(2n) \cap \text{symp}_{\mathbb{R}}(2n)$. Furthermore, $A \in \text{SO}(2n) \cap \text{Symp}_{\mathbb{R}}(2n)$ if and only if $A \in \text{Symp}_{\mathbb{R}}(2n)$ and $AJ_{2n} = J_{2n}A$. Finally, $A \in \text{so}(2n) \cap \text{symp}_{\mathbb{R}}(2n)$ if and only if $A \in \text{symp}_{\mathbb{R}}(2n)$ and $AJ_{2n} = J_{2n}A$. (Proof: See [194].)

Fact 3.22.6. Define $Q_0, Q_1, Q_2, Q_3 \in \mathbb{C}^{2 \times 2}$ by

$$Q_0 \triangleq I_2, \quad Q_1 \triangleq \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad Q_2 \triangleq \begin{bmatrix} -j & 0 \\ 0 & j \end{bmatrix}, \quad Q_3 \triangleq \begin{bmatrix} 0 & -j \\ -j & 0 \end{bmatrix}.$$

Then, the following statements hold:

- i) $Q_0^* = Q_0$ and $Q_i^* = -Q_i$ for all $i = 1, 2, 3$.
- ii) $Q_0^2 = Q_0$ and $Q_i^2 = -Q_0$ for all $i = 1, 2, 3$.
- iii) $Q_i Q_j = -Q_j Q_i$ for all $1 \leq i < j \leq 3$.
- iv) $Q_1 Q_2 = Q_3$, $Q_2 Q_3 = Q_1$, and $Q_3 Q_1 = Q_2$.
- v) $\{\pm Q_0, \pm Q_1, \pm Q_2, \pm Q_3\}$ is a group.

For $\beta \triangleq [\beta_0 \quad \beta_1 \quad \beta_2 \quad \beta_3]^T \in \mathbb{R}^4$ define

$$Q(\beta) \triangleq \sum_{i=0}^3 \beta_i Q_i = \begin{bmatrix} \beta_0 + \beta_1 j & -(\beta_2 + \beta_3 j) \\ \beta_2 - \beta_3 j & \beta_0 - \beta_1 j \end{bmatrix}.$$

Then,

$$Q(\beta) Q^*(\beta) = \beta^T \beta I_2$$

and

$$\det Q(\beta) = \beta^T \beta.$$

Hence, if $\beta^T \beta = 1$, then $Q(\beta)$ is unitary. Furthermore, the complex matrices Q_0, Q_1, Q_2, Q_3 , and $Q(\beta)$ have the real representations

$$\begin{aligned} Q_0 &= I_4, & Q_1 &= \begin{bmatrix} -J_2 & 0 \\ 0 & -J_2 \end{bmatrix}, \\ Q_2 &= \begin{bmatrix} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{bmatrix}, & Q_3 &= \begin{bmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{bmatrix}, \\ Q(\beta) &= \begin{bmatrix} \beta_0 & -\beta_1 & -\beta_2 & -\beta_3 \\ \beta_1 & \beta_0 & -\beta_3 & \beta_2 \\ \beta_2 & \beta_3 & \beta_0 & -\beta_1 \\ \beta_3 & -\beta_2 & \beta_1 & \beta_0 \end{bmatrix}. \end{aligned}$$

Hence,

$$Q(\beta) Q^T(\beta) = \beta^T \beta I_4$$

and

$$\det Q(\beta) = (\beta^T \beta)^2.$$

(Remark: Q_0, Q_1, Q_2, Q_3 represent the quaternions $1, \hat{i}, \hat{j}, \hat{k}$. See Fact 3.22.1. An alternative representation is given by the *Pauli spin matrices* given by $\sigma_0 = I_2, \sigma_1 = jQ_3, \sigma_2 = jQ_1, \sigma_3 = jQ_2$. See [636, pp. 143–144], [777].) (Remark: For applications of quaternions, see [26, 607, 636, 850].) (Remark: $Q(\beta)$ has the form $\begin{bmatrix} A & B \\ -B & A \end{bmatrix}$, where A and $\hat{I}B$ are rotation-dilations. See Fact 2.19.1.)

Fact 3.22.7. Let $A, B, C, D \in \mathbb{R}^{n \times m}$, define $\hat{i}, \hat{j}, \hat{k}$ as in Fact 3.22.1, and let $Q \triangleq A + \hat{i}B + \hat{j}C + \hat{k}D$. Then,

$$\text{diag}(Q, Q) = U_n^* \begin{bmatrix} A + \hat{i}B & -C - \hat{i}D \\ C - \hat{i}D & A - \hat{i}B \end{bmatrix} U_m,$$

where

$$U_n \triangleq \frac{1}{\sqrt{2}} \begin{bmatrix} I_n & -\hat{i}I_n \\ -\hat{j}I_n & \hat{k}I_n \end{bmatrix}.$$

Furthermore, $U_n U_n^* = I_{2n}$. (Proof: See [1304, 1305].) (Remark: When $n = m$, this identity uses a similarity transformation to construct a complex representation of quaternions.) (Remark: The complex conjugate U_n^* is constructed as in Fact 3.22.7.)

Fact 3.22.8. Let $A, B, C, D \in \mathbb{R}^{n \times n}$, define $\hat{i}, \hat{j}, \hat{k}$ as in Fact 3.22.1, and let $Q \triangleq A + \hat{i}B + \hat{j}C + \hat{k}D$. Then,

$$\text{diag}(Q, Q, Q, Q) = U_n \begin{bmatrix} A & -B & -C & -D \\ B & A & -D & C \\ C & D & A & -B \\ D & -C & B & A \end{bmatrix} U_m,$$

where

$$U_n \triangleq \frac{1}{2} \begin{bmatrix} I_n & \hat{i}I_n & \hat{j}I_n & \hat{k}I_n \\ -\hat{i}I_n & I_n & \hat{k}I_n & -\hat{j}I_n \\ -\hat{j}I_n & -\hat{k}I_n & I_n & \hat{i}I_n \\ -\hat{k}I_n & \hat{j}I_n & -\hat{i}I_n & I_n \end{bmatrix}.$$

Furthermore, $U_n^* = U_n$ and $U_n^2 = I_{4n}$. (Proof: See [1304, 1305]. See also [80, 257, 470, 600, 1488].) (Remark: When $n = m$, this identity uses a similarity transformation to construct a real representation of quaternions. See Fact 2.14.11.) (Remark: The complex conjugate U_n^* is constructed by replacing $\hat{i}, \hat{j}, \hat{k}$ by $-\hat{i}, -\hat{j}, -\hat{k}$, respectively, in U_n^T .)

Fact 3.22.9. Let $A \in \mathbb{C}^{2 \times 2}$. Then, A is unitary if and only if there exist $\theta \in \mathbb{R}$ and $\beta \in \mathbb{R}^4$ such that $A = e^{j\theta}Q(\beta)$, where $Q(\beta)$ is defined in Fact 3.22.6. (Proof: See [1129, p. 228].)

3.23 Notes

In the literature on generalized inverses, range-Hermitian matrices are traditionally called *EP matrices*. Elementary reflectors are traditionally called *Householder matrices* or *Householder reflections*.

An alternative term for irreducible is *indecomposable*, see [963, p. 147].

Left equivalence, right equivalence, and biequivalence are treated in [1129]. Each of the groups defined in Proposition 3.3.6 is a *Lie group*; see Definition 11.6.1. Elementary treatments of Lie algebras and Lie groups are given in [75, 77, 103, 362, 459, 473, 553, 554, 724, 1077, 1147, 1185], while an advanced treatment ap-

pears in [1366]. Some additional groups of structured matrices are given in [944]. Applications of group theory are discussed in [781].

“Almost nonnegative matrices” are called “ML-matrices” in [1184, p. 208] and “essentially nonnegative matrices” in [182, 190, 617].

The terminology “idempotent” and “projector” is not standardized in the literature. Some writers use “projector,” “oblique projector,” or “projection” [536] for idempotent, and “orthogonal projector” or “orthoprojector” for projector. Centrosymmetric and centrohermitian matrices are discussed in [883, 1410].

Matrices with set-valued entries are discussed in [551]. Matrices with entries having physical dimensions are discussed in [641, 1062].

Chapter Four

Polynomial Matrices and Rational Transfer Functions

In this chapter we consider matrices whose entries are polynomials or rational functions. The decomposition of polynomial matrices in terms of the Smith form provides the foundation for developing canonical forms in Chapter 5. In this chapter we also present some basic properties of eigenvalues and eigenvectors as well as the minimal and characteristic polynomials of a square matrix. Finally, we consider the extension of the Smith form to the Smith-McMillan form for rational transfer functions.

4.1 Polynomials

A function $p: \mathbb{C} \mapsto \mathbb{C}$ of the form

$$p(s) = \beta_k s^k + \beta_{k-1} s^{k-1} + \cdots + \beta_1 s + \beta_0, \quad (4.1.1)$$

where $k \in \mathbb{N}$ and $\beta_0, \dots, \beta_k \in \mathbb{F}$, is a *polynomial*. The set of polynomials is denoted by $\mathbb{F}[s]$. If the coefficient $\beta_k \in \mathbb{F}$ is nonzero, then the *degree* of p , denoted by $\deg p$, is k . If, in addition, $\beta_k = 1$, then p is *monic*. If $k = 0$, then p is *constant*. The degree of a nonzero constant polynomial is zero, while the degree of the zero polynomial is defined to be $-\infty$.

Let p_1 and p_2 be polynomials. Then,

$$\deg p_1 p_2 = \deg p_1 + \deg p_2. \quad (4.1.2)$$

If $p_1 = 0$ or $p_2 = 0$, then $\deg p_1 p_2 = \deg p_1 + \deg p_2 = -\infty$. If p_2 is a nonzero constant, then $\deg p_2 = 0$, and thus $\deg p_1 p_2 = \deg p_1$. Furthermore,

$$\deg(p_1 + p_2) \leq \max\{\deg p_1, \deg p_2\}. \quad (4.1.3)$$

Therefore, $\deg(p_1 + p_2) = \max\{\deg p_1, \deg p_2\}$ if and only if either *i*) $\deg p_1 \neq \deg p_2$ or *ii*) $p_1 = p_2 = 0$ or *iii*) $r \triangleq \deg p_1 = \deg p_2 \neq -\infty$ and the sum of the coefficients of s^r in p_1 and p_2 is not zero. Equivalently, $\deg(p_1 + p_2) < \max\{\deg p_1, \deg p_2\}$ if and only if $r \triangleq \deg p_1 = \deg p_2 \neq -\infty$ and the sum of the coefficients of s^r in p_1 and p_2 is zero.

Let $p \in \mathbb{F}[s]$ be a polynomial of degree $k \geq 1$. Then, it follows from the *fundamental theorem of algebra* that p has k possibly repeated complex roots $\lambda_1, \dots, \lambda_k$ and thus can be factored as

$$p(s) = \beta \prod_{i=1}^k (s - \lambda_i), \quad (4.1.4)$$

where $\beta \in \mathbb{F}$. The multiplicity of a root $\lambda \in \mathbb{C}$ of p is denoted by $\text{mult}_p(\lambda)$. If λ is not a root of p , then $\text{mult}_p(\lambda) = 0$. The multiset consisting of the roots of p including multiplicity is $\text{mroots}(p) = \{\lambda_1, \dots, \lambda_k\}_{\text{ms}}$, while the set of roots of p ignoring multiplicity is $\text{roots}(p) = \{\hat{\lambda}_1, \dots, \hat{\lambda}_l\}$, where $\sum_{i=1}^l \text{mult}_p(\hat{\lambda}_i) = k$. If $\mathbb{F} = \mathbb{R}$, then the multiplicity of a root λ_i whose imaginary part is nonzero is equal to the multiplicity of its complex conjugate $\bar{\lambda}_i$. Hence, $\text{mroots}(p)$ is *self-conjugate*, that is, $\text{mroots}(p) = \overline{\text{mroots}(p)}$.

Let $p \in \mathbb{F}[s]$. If $p(-s) = p(s)$ for all $s \in \mathbb{C}$, then p is *even*, while, if $p(-s) = -p(s)$ for all $s \in \mathbb{C}$, then p is *odd*. If p is either odd or even, then $\text{mroots}(p) = -\text{mroots}(p)$. If $p \in \mathbb{R}[s]$ and there exists a polynomial $q \in \mathbb{R}[s]$ such that $p(s) = q(s)q(-s)$ for all $s \in \mathbb{C}$, then p has a *spectral factorization*. If p has a spectral factorization, then p is even and $\deg p$ is an even integer.

Proposition 4.1.1. Let $p \in \mathbb{R}[s]$. Then, the following statements are equivalent:

- i) p has a spectral factorization.
- ii) p is even, and every imaginary root of p has even multiplicity.
- iii) p is even, and $p(j\omega) \geq 0$ for all $\omega \in \mathbb{R}$.

Proof. The equivalence of i) and ii) is immediate. To prove i) $\implies$ iii), note that, for all $\omega \in \mathbb{R}$,

$$p(j\omega) = q(j\omega)q(-j\omega) = |q(j\omega)|^2 \geq 0.$$

Conversely, to prove iii) $\implies$ i) write $p = p_1 p_2$, where every root of p_1 is imaginary and none of the roots of p_2 are imaginary. Now, let z be a root of p_2 . Then, $-z$, $\bar{z}$, and $-\bar{z}$ are also roots of p_2 with the same multiplicity as z . Hence, there exists a polynomial $p_{20} \in \mathbb{R}[s]$ such that $p_2(s) = p_{20}(s)p_{20}(-s)$ for all $s \in \mathbb{C}$.

Next, assuming that p has at least one imaginary root, write $p_1(s) = \prod_{i=1}^k (s^2 + \omega_i^2)^{m_i}$, where $0 \leq \omega_1 < \dots < \omega_k$ and $m_i \triangleq \text{mult}_p(j\omega_i)$. Let ω_{i_0} denote the smallest element of the set $\{\omega_1, \dots, \omega_k\}$ such that m_i is odd. Then, it follows that $p_1(j\omega) = \prod_{i=1}^k (\omega_i^2 - \omega^2)^{m_i} < 0$ for all $\omega \in (\omega_{i_0}, \omega_{i_0+1})$, where $\omega_{k+1} \triangleq \infty$. However, note that $p_1(j\omega) = p(j\omega)/p_2(j\omega) = p(j\omega)/|p_{20}(j\omega)|^2 \geq 0$ for all $\omega \in \mathbb{R}$, which is a contradiction. Therefore, m_i is even for all $i = 1, \dots, k$, and thus $p_1(s) = p_{10}(s)p_{10}(-s)$ for all $s \in \mathbb{C}$, where $p_{10}(s) \triangleq \prod_{i=1}^k (s^2 + \omega_i^2)^{m_i/2}$. Consequently, $p(s) = p_{10}(s)p_{20}(s)p_{10}(-s)p_{20}(-s)$ for all $s \in \mathbb{C}$. Finally, if p has no imaginary roots, then $p_1 = 1$, and $p(s) = p_{20}(s)p_{20}(-s)$ for all $s \in \mathbb{C}$. $\square$

The following division algorithm is essential to the study of polynomials.

Lemma 4.1.2. Let $p_1, p_2 \in \mathbb{F}[s]$, and assume that p_2 is not the zero polynomial. Then, there exist unique polynomials $q, r \in \mathbb{F}[s]$ such that $\deg r < \deg p_2$ and

$$p_1 = qp_2 + r. \quad (4.1.5)$$

Proof. Define $n \triangleq \deg p_1$ and $m \triangleq \deg p_2$. If $n < m$, then $q = 0$ and $r = p_1$. Hence, $\deg r = \deg p_1 = n < m = \deg p_2$.

Now, assume that $n \geq m \geq 0$, and write $p_1(s) = \beta_n s^n + \cdots + \beta_0$ and $p_2(s) = \gamma_m s^m + \cdots + \gamma_0$. If $n = 0$, then $m = 0$, $\gamma_0 \neq 0$, $q = \beta_0/\gamma_0$, and $r = 0$. Hence, $-\infty = \deg r < 0 = \deg p_2$.

If $n = 1$, then either $m = 0$ or $m = 1$. If $m = 0$, then $p_2(s) = \gamma_0 \neq 0$, and (4.1.5) is satisfied with $q(s) = p_1(s)/\gamma_0$ and $r = 0$, in which case $-\infty = \deg r < 0 = \deg p_2$. If $m = 1$, then (4.1.5) is satisfied with $q(s) = \beta_1/\gamma_1$ and $r(s) = \beta_0 - \beta_1\gamma_0/\gamma_1$. Hence, $\deg r \leq 0 < 1 = \deg p_2$.

Now, suppose that $n = 2$. Then, $\hat{p}_1(s) = p_1(s) - (\beta_2/\gamma_m)s^{2-m}p_2(s)$ has degree 1. Applying (4.1.5) with p_1 replaced by $\hat{p}_1$, it follows that there exist polynomials $q_1, r_1 \in \mathbb{F}[s]$ such that $\hat{p}_1 = q_1p_2 + r_1$ and such that $\deg r_1 < \deg p_2$. It thus follows that $p_1(s) = q_1(s)p_2(s) + r_1(s) + (\beta_2/\gamma_m)s^{2-m}p_2(s) = q(s)p_2(s) + r(s)$, where $q(s) = q_1(s) + (\beta_2/\gamma_m)s^{n-m}$ and $r = r_1$, which verifies (4.1.5). Similar arguments apply to successively larger values of n .

To prove uniqueness, suppose there exist polynomials $\hat{q}$ and $\hat{r}$ such that $\deg \hat{r} < \deg p_2$ and $p_1 = \hat{q}p_2 + \hat{r}$. Then, it follows that $(\hat{q} - q)p_2 = r - \hat{r}$. Next, note that $\deg(r - \hat{r}) < \deg p_2$. If $\hat{q} \neq q$, then $\deg p_2 \leq \deg[(\hat{q} - q)p_2]$ so that $\deg(r - \hat{r}) < \deg[(\hat{q} - q)p_2]$, which is a contradiction. Thus, $\hat{q} = q$, and, hence, $r = \hat{r}$. $\square$

In Lemma 4.1.2, q is the *quotient* of p_1 and p_2 , while r is the *remainder*. If $r = 0$, then p_2 divides p_1 , or, equivalently, p_1 is a *multiple* of p_2 . Note that, if $p_2(s) = s - \alpha$, where $\alpha \in \mathbb{F}$, then r is constant and is given by $r(s) = p_1(\alpha)$.

If a polynomial $p_3 \in \mathbb{F}[s]$ divides two polynomials $p_1, p_2 \in \mathbb{F}[s]$, then p_3 is a *common divisor* of p_1 and p_2 . Given polynomials $p_1, p_2 \in \mathbb{F}[s]$, there exists a unique monic polynomial $p_3 \in \mathbb{F}[s]$, the *greatest common divisor* of p_1 and p_2 , such that p_3 is a common divisor of p_1 and p_2 and such that every common divisor of p_1 and p_2 divides p_3 . In addition, there exist polynomials $q_1, q_2 \in \mathbb{F}[s]$ such that the greatest common divisor p_3 of p_1 and p_2 is given by $p_3 = q_1p_1 + q_2p_2$. See [1081, p. 113] for proofs of these results. Finally, p_1 and p_2 are *coprime* if their greatest common divisor is $p_3 = 1$, while a polynomial $p \in \mathbb{F}[s]$ is *irreducible* if there do not exist nonconstant polynomials $p_1, p_2 \in \mathbb{F}[s]$ such that $p = p_1p_2$. For example, if $\mathbb{F} = \mathbb{R}$, then $p(s) = s^2 + s + 1$ is irreducible.

If a polynomial $p_3 \in \mathbb{F}[s]$ is a multiple of two polynomials $p_1, p_2 \in \mathbb{F}[s]$, then p_3 is a *common multiple* of p_1 and p_2 . Given nonzero polynomials p_1 and p_2 , there exists (see [1081, p. 113]) a unique monic polynomial $p_3 \in \mathbb{F}[s]$ that is a common multiple of p_1 and p_2 and that divides every common multiple of p_1 and p_2 . The polynomial p_3 is the *least common multiple* of p_1 and p_2 .

The polynomial $p \in \mathbb{F}[s]$ given by (4.1.1) can be evaluated with a square matrix argument $A \in \mathbb{F}^{n \times n}$ by defining

$$p(A) \triangleq \beta_k A^k + \beta_{k-1} A^{k-1} + \cdots + \beta_1 A + \beta_0 I. \quad (4.1.6)$$

4.2 Polynomial Matrices

The set $\mathbb{F}^{n \times m}[s]$ of *polynomial matrices* consists of matrix functions $P: \mathbb{C} \mapsto \mathbb{C}^{n \times m}$ whose entries are elements of $\mathbb{F}[s]$. A polynomial matrix $P \in \mathbb{F}^{n \times m}[s]$ can thus be written as

$$P(s) = s^k B_k + s^{k-1} B_{k-1} + \cdots + s B_1 + B_0, \quad (4.2.1)$$

where $B_0, \dots, B_k \in \mathbb{F}^{n \times m}$. If B_k is nonzero, then the *degree* of P , denoted by $\deg P$, is k , whereas, if $P = 0$, then $\deg P = -\infty$. If $n = m$ and B_k is nonsingular, then P is *regular*, while, if $B_k = I$, then P is *monic*.

The following result, which generalizes Lemma 4.1.2, provides a division algorithm for polynomial matrices.

Lemma 4.2.1. Let $P_1, P_2 \in \mathbb{F}^{n \times n}[s]$, where P_2 is regular. Then, there exist unique polynomial matrices $Q, R, \hat{Q}, \hat{R} \in \mathbb{F}^{n \times n}[s]$ such that $\deg R < \deg P_2$, $\deg \hat{R} < \deg P_2$,

$$P_1 = QP_2 + R, \quad (4.2.2)$$

and

$$P_1 = P_2 \hat{Q} + \hat{R}. \quad (4.2.3)$$

Proof. See [559, p. 90] or [1081, pp. 134–135]. $\square$

If $R = 0$, then P_2 *right divides* P_1 , while, if $\hat{R} = 0$, then P_2 *left divides* P_1 .

Let the polynomial matrix $P \in \mathbb{F}^{n \times m}[s]$ be given by (4.2.1). Then, P can be evaluated with a square matrix argument in two different ways, either from the right or from the left. For $A \in \mathbb{C}^{m \times m}$ define

$$P_R(A) \triangleq B_k A^k + B_{k-1} A^{k-1} + \cdots + B_1 A + B_0, \quad (4.2.4)$$

while, for $A \in \mathbb{C}^{n \times n}$, define

$$P_L(A) \triangleq A^k B_k + A^{k-1} B_{k-1} + \cdots + A B_1 + B_0. \quad (4.2.5)$$

$P_R(A)$ and $P_L(A)$ are *matrix polynomials*.

If $n = m$, then $P_R(A)$ and $P_L(A)$ can be evaluated for all $A \in \mathbb{F}^{n \times n}$, although these matrices may be different.

The following result is useful.

Lemma 4.2.2. Let $Q, \hat{Q} \in \mathbb{F}^{n \times n}[s]$ and $A \in \mathbb{F}^{n \times n}$. Furthermore, define $P, \hat{P} \in \mathbb{F}^{n \times n}[s]$ by $P(s) \triangleq Q(s)(sI - A)$ and $\hat{P}(s) \triangleq (sI - A)\hat{Q}(s)$. Then, $P_R(A) = 0$ and $\hat{P}_L(A) = 0$.

Let $p \in \mathbb{F}[s]$ be given by (4.1.1), and define $P(s) \triangleq p(s)I_n = s^k \beta_k I_n + s^{k-1} \beta_{k-1} I_n + \cdots + s \beta_1 I_n + \beta_0 I_n \in \mathbb{F}^{n \times n}[s]$. For $A \in \mathbb{C}^{n \times n}$ it follows that $p(A) = P(A) = P_R(A) = P_L(A)$.

The following result specializes Lemma 4.2.1 to the case of polynomial matrix divisors of degree 1.

Corollary 4.2.3. Let $P \in \mathbb{F}^{n \times n}[s]$ and $A \in \mathbb{F}^{n \times n}$. Then, there exist unique polynomial matrices $Q, \hat{Q} \in \mathbb{F}^{n \times n}[s]$ and unique matrices $R, \hat{R} \in \mathbb{F}^{n \times n}$ such that

$$P(s) = Q(s)(sI - A) + R \quad (4.2.6)$$

and

$$P(s) = (sI - A)\hat{Q}(s) + \hat{R}. \quad (4.2.7)$$

Furthermore, $R = P_R(A)$ and $\hat{R} = P_L(A)$.

Proof. In Lemma 4.2.1 set $P_1 = P$ and $P_2(s) = sI - A$. Since $\deg P_2 = 1$, it follows that $\deg R = \deg \hat{R} = 0$, and thus R and $\hat{R}$ are constant. Finally, the last statement follows from Lemma 4.2.2. $\square$

Definition 4.2.4. Let $P \in \mathbb{F}^{n \times m}[s]$. Then, $\text{rank } P$ is defined by

$$\text{rank } P \triangleq \max_{s \in \mathbb{C}} \text{rank } P(s). \quad (4.2.8)$$

Let $P \in \mathbb{F}^{n \times n}[s]$. Then, $P(s) \in \mathbb{C}^{n \times n}$ for all $s \in \mathbb{C}$. Furthermore, $\det P$ is a polynomial in s , that is, $\det P \in \mathbb{F}[s]$.

Definition 4.2.5. Let $P \in \mathbb{F}^{n \times n}[s]$. Then, P is *nonsingular* if $\det P$ is not the zero polynomial; otherwise, P is *singular*.

Proposition 4.2.6. Let $P \in \mathbb{F}^{n \times n}[s]$, and assume that P is regular. Then, P is nonsingular.

Let $P \in \mathbb{F}^{n \times n}[s]$. If P is nonsingular, then the *inverse* P^{-1} of P can be constructed according to (2.7.22). In general, the entries of P^{-1} are rational functions of s (see Definition 4.7.1). For example, if $P(s) = \begin{bmatrix} s+2 & s+1 \\ s-2 & s-1 \end{bmatrix}$, then $P^{-1}(s) = \frac{1}{2s} \begin{bmatrix} s-1 & -s-1 \\ -s+2 & s+2 \end{bmatrix}$. In certain cases, P^{-1} is also a polynomial matrix. For example, if $P(s) = \begin{bmatrix} s & 1 \\ s^2+s-1 & s+1 \end{bmatrix}$, then $P^{-1}(s) = \begin{bmatrix} s+1 & -1 \\ -s^2-s+1 & s \end{bmatrix}$.

The following result is an extension of Proposition 2.7.7 from constant matrices to polynomial matrices.

Proposition 4.2.7. Let $P \in \mathbb{F}^{n \times m}[s]$. Then, $\text{rank } P$ is the order of the largest nonsingular polynomial matrix that is a submatrix of P .

Proof. For all $s \in \mathbb{C}$ it follows from Proposition 2.7.7 that $\text{rank } P(s)$ is the order of the largest nonsingular submatrix of $P(s)$. Now, let $s_0 \in \mathbb{C}$ be such that $\text{rank } P(s_0) = \text{rank } P$. Then, $P(s_0)$ has a nonsingular submatrix of maximal order $\text{rank } P$. Therefore, P has a nonsingular polynomial submatrix of maximal order $\text{rank } P$. $\square$

A polynomial matrix can be transformed by performing elementary row and column operations of the following types:

- i) Multiply a row or a column by a nonzero constant.
- ii) Interchange two rows or two columns.
- iii) Add a polynomial multiple of one (row, column) to another (row, column).

These operations correspond respectively to left multiplication or right multiplication by the elementary matrices

$$I_n + (\alpha - 1)E_{i,i} = \begin{bmatrix} I_{i-1} & 0 & 0 \\ 0 & \alpha & 0 \\ 0 & 0 & I_{n-i} \end{bmatrix}, \quad (4.2.9)$$

where $\alpha \in \mathbb{F}$ is nonzero,

$$I_n + E_{i,j} + E_{j,i} - E_{i,i} - E_{j,j} = \begin{bmatrix} I_{i-1} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & I_{j-i-1} & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & I_{n-j} \end{bmatrix}, \quad (4.2.10)$$

where $i \neq j$, and the *elementary polynomial matrix*

$$I_n + pE_{i,j} = \begin{bmatrix} I_{i-1} & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & p & 0 \\ 0 & 0 & I_{j-i-1} & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & I_{n-j} \end{bmatrix}, \quad (4.2.11)$$

where $i \neq j$ and $p \in \mathbb{F}[s]$. The matrices shown in (4.2.10) and (4.2.11) illustrate the case $i < j$. Applying these operations sequentially corresponds to forming products of elementary matrices and elementary polynomial matrices. Note that the elementary polynomial matrix $I + pE_{i,j}$ is nonsingular, and that $(I + pE_{i,j})^{-1} = I - pE_{i,j}$. Therefore, the inverse of an elementary polynomial matrix is an elementary polynomial matrix.

4.3 The Smith Decomposition and Similarity Invariants

Definition 4.3.1. Let $P \in \mathbb{F}^{n \times n}[s]$. Then, P is *unimodular* if P is the product of elementary matrices and elementary polynomial matrices.

The following result provides a canonical form, known as the *Smith form*, for polynomial matrices under unimodular transformation.

Theorem 4.3.2. Let $P \in \mathbb{F}^{n \times m}[s]$, and let $r \triangleq \text{rank } P$. Then, there exist unimodular matrices $S_1 \in \mathbb{F}^{n \times n}[s]$ and $S_2 \in \mathbb{F}^{m \times m}[s]$ and monic polynomials $p_1, \dots, p_r \in \mathbb{F}[s]$ such that p_i divides p_{i+1} for all $i = 1, \dots, r-1$ and such that

$$P = S_1 \begin{bmatrix} p_1 & & & 0 \\ & \ddots & & \\ & & p_r & \\ 0 & & & 0_{(n-r) \times (m-r)} \end{bmatrix} S_2. \quad (4.3.1)$$

Furthermore, for all $i = 1, \dots, r$, let Δ_i denote the monic greatest common divisor of all $i \times i$ subdeterminants of P . Then, p_i is uniquely determined by

$$\Delta_i = p_1 \cdots p_i. \quad (4.3.2)$$

Proof. The result is obtained by sequentially applying elementary row and column operations to P . For details, see [787, pp. 390–392] or [1081, pp. 125–128]. $\square$

Definition 4.3.3. The monic polynomials $p_1, \dots, p_r \in \mathbb{F}[s]$ of the Smith form (4.3.1) of $P \in \mathbb{F}^{n \times m}[s]$ are the *Smith polynomials* of P . The *Smith zeros* of P are the roots of $p_1, \dots, p_r$. Let

$$\text{Szeros}(P) \triangleq \text{roots}(p_r) \quad (4.3.3)$$

and

$$\text{mSzeros}(P) \triangleq \bigcup_{i=1}^r \text{mroots}(p_i). \quad (4.3.4)$$

Proposition 4.3.4. Let $P \in \mathbb{R}^{n \times m}[s]$, and assume there exist unimodular matrices $S_1 \in \mathbb{F}^{n \times n}[s]$ and $S_2 \in \mathbb{F}^{m \times m}[s]$ and monic polynomials $p_1, \dots, p_r \in \mathbb{F}[s]$ satisfying (4.3.1). Then, $\text{rank } P = r$.

Proposition 4.3.5. Let $P \in \mathbb{F}^{n \times m}[s]$, and let $r \triangleq \text{rank } P$. Then, r is the largest order of all nonsingular submatrices of P .

Proof. Let r_0 denote the largest order of all nonsingular submatrices of P , and let $P_0 \in \mathbb{F}^{r_0 \times r_0}[s]$ be a nonsingular submatrix of P . First, assume that $r < r_0$. Then, there exists $s_0 \in \mathbb{C}$ such that $\text{rank } P(s_0) = \text{rank } P_0(s_0) = r_0$. Thus, $r = \text{rank } P = \max_{s \in \mathbb{C}} \text{rank } P(s) \geq \text{rank } P(s_0) = r_0$, which is a contradiction. Next, assume that $r > r_0$. Then, it follows from (4.3.1) that there exists $s_0 \in \mathbb{C}$ such that $\text{rank } P(s_0) = r$. Consequently, $P(s_0)$ has a nonsingular $r \times r$ submatrix. Let $\hat{P}_0 \in \mathbb{F}^{r \times r}[s]$ denote the corresponding submatrix of P . Thus, $\hat{P}_0$ is nonsingular, which implies that P has a nonsingular submatrix whose order is greater than r_0 , which is a contradiction. Consequently, $r = r_0$. $\square$

Proposition 4.3.6. Let $P \in \mathbb{F}^{n \times m}[s]$, and let $\mathcal{S} \subset \mathbb{C}$ be a finite set. Then,

$$\text{rank } P = \max_{s \in \mathbb{C} \setminus \mathcal{S}} \text{rank } P(s). \quad (4.3.5)$$

Proposition 4.3.7. Let $P \in \mathbb{F}^{n \times n}[s]$. Then, the following statements are equivalent:

- i) P is unimodular.
- ii) $\det P$ is a nonzero constant.
- iii) The Smith form of P is the identity.
- iv) P is nonsingular, and P^{-1} is a polynomial matrix.
- v) P is nonsingular, and P^{-1} is unimodular.

Proof. To prove $i) \implies ii)$, note that every elementary matrix and every elementary polynomial matrix has a constant nonzero determinant. Since P is a product of elementary matrices and elementary polynomial matrices, its determinant is a constant.

To prove $ii) \implies iii)$, note that it follows from (4.3.1) that $\text{rank } P = n$ and $\det P = (\det S_1)(\det S_2)p_1 \cdots p_n$, where $S_1, S_2 \in \mathbb{F}^{n \times n}$ are unimodular and $p_1, \dots, p_n$ are monic polynomials. From the result $i) \implies ii)$, it follows that $\det S_1$ and $\det S_2$ are nonzero constants. Since $\det P$ is a nonzero constant, it follows that $p_1 \cdots p_n = \det P / [(\det S_1)(\det S_2)]$ is a nonzero constant. Since $p_1, \dots, p_n$ are monic polynomials, it follows that $p_1 = \cdots = p_n = 1$.

Next, to prove $iii) \implies iv)$, note that P is unimodular, and thus it follows that $\det P$ is a nonzero constant. Furthermore, since P^A is a polynomial matrix, it follows that $P^{-1} = (\det P)^{-1}P^A$ is a polynomial matrix.

To prove $iv) \implies v)$, note that $\det P^{-1}$ is a polynomial. Since $\det P$ is a polynomial and $\det P^{-1} = 1/\det P$ it follows that $\det P$ is a nonzero constant. Hence, P is unimodular, and thus $P^{-1} = (\det P)^{-1}P^A$ is unimodular.

Finally, to prove $v) \implies i)$, note that $\det P^{-1}$ is a nonzero constant, and thus $P = [\det P^{-1}]^{-1}[P^{-1}]^A$ is a polynomial matrix. Furthermore, since $\det P = 1/\det P^{-1}$, it follows that $\det P$ is a nonzero constant. Hence, P is unimodular. $\square$

Proposition 4.3.8. Let $A_1, B_1, A_2, B_2 \in \mathbb{F}^{n \times n}$, where A_2 is nonsingular, and define the polynomial matrices $P_1, P_2 \in \mathbb{F}^{n \times n}[s]$ by $P_1(s) \triangleq sA_1 + B_1$ and $P_2(s) \triangleq sA_2 + B_2$. Then, P_1 and P_2 have the same Smith polynomials if and only if there exist nonsingular matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that $P_2 = S_1P_1S_2$.

Proof. The sufficiency result is immediate. To prove necessity, note that it follows from Theorem 4.3.2 that there exist unimodular matrices $T_1, T_2 \in \mathbb{F}^{n \times n}[s]$ such that $P_2 = T_2P_1T_1$. Now, since P_2 is regular, it follows from Lemma 4.2.1 that there exist polynomial matrices $Q, \hat{Q} \in \mathbb{F}^{n \times n}[s]$ and constant matrices $R, \hat{R} \in \mathbb{F}^{n \times n}$

such that $T_1 = QP_2 + R$ and $T_2 = P_2\hat{Q} + \hat{R}$. Next, we have

$$\begin{aligned}
 P_2 &= T_2P_1T_1 \\
 &= (P_2\hat{Q} + \hat{R})P_1T_1 \\
 &= \hat{R}P_1T_1 + P_2\hat{Q}T_2^{-1}P_2 \\
 &= \hat{R}P_1(QP_2 + R) + P_2\hat{Q}T_2^{-1}P_2 \\
 &= \hat{R}P_1R + (T_2 - P_2\hat{Q})P_1QP_2 + P_2\hat{Q}T_2^{-1}P_2 \\
 &= \hat{R}P_1R + T_2P_1QP_2 + P_2(-\hat{Q}P_1Q + \hat{Q}T_2^{-1})P_2 \\
 &= \hat{R}P_1R + P_2(T_1^{-1}Q - \hat{Q}P_1Q + \hat{Q}T_2^{-1})P_2.
 \end{aligned}$$

Since P_2 is regular and has degree 1, it follows that, if $T_1^{-1}Q - \hat{Q}P_1Q + \hat{Q}T_2^{-1}$ is not zero, then $\deg P_2(T_1^{-1}Q - \hat{Q}P_1Q + \hat{Q}T_2^{-1})P_2 \geq 2$. However, since P_2 and $\hat{R}P_1R$ have degree less than 2, it follows that $T_1^{-1}Q - \hat{Q}P_1Q + \hat{Q}T_2^{-1} = 0$. Hence, $P_2 = \hat{R}P_1R$.

Next, to show that $\hat{R}$ and R are nonsingular, note that, for all $s \in \mathbb{C}$,

$$P_2(s) = \hat{R}P_1(s)R = s\hat{R}A_1R + \hat{R}B_1R,$$

which implies that $A_2 = S_1A_1S_2$, where $S_1 = \hat{R}$ and $S_2 = R$. Since A_2 is nonsingular, it follows that S_1 and S_2 are nonsingular. $\square$

Definition 4.3.9. Let $A \in \mathbb{F}^{n \times n}$. Then, the *similarity invariants* of A are the Smith polynomials of $sI - A$.

The following result provides necessary and sufficient conditions for two matrices to be similar.

Theorem 4.3.10. Let $A, B \in \mathbb{F}^{n \times n}$. Then, A and B are similar if and only if they have the same similarity invariants.

Proof. To prove necessity, assume that A and B are similar. Then, the matrices $sI - A$ and $sI - B$ have the same Smith form and thus the same similarity invariants. To prove sufficiency, it follows from Proposition 4.3.8 that there exist nonsingular matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that $sI - A = S_1(sI - B)S_2$. Thus, $S_1 = S_2^{-1}$, and, hence, $A = S_1BS_1^{-1}$. $\square$

Corollary 4.3.11. Let $A \in \mathbb{F}^{n \times n}$. Then, A and A^T are similar.

An improved form of Corollary 4.3.11 is given by Corollary 5.3.8.

4.4 Eigenvalues

Let $A \in \mathbb{F}^{n \times n}$. Then, the polynomial matrix $sI - A \in \mathbb{F}^{n \times n}[s]$ is monic and has degree 1.

Definition 4.4.1. Let $A \in \mathbb{F}^{n \times n}$. Then, the *characteristic polynomial* of A is the polynomial $\chi_A \in \mathbb{F}[s]$ given by

$$\chi_A(s) \triangleq \det(sI - A). \quad (4.4.1)$$

Since $sI - A$ is a polynomial matrix, its determinant is the product of its Smith polynomials, that is, the similarity invariants of A .

Proposition 4.4.2. Let $A \in \mathbb{F}^{n \times n}$, and let $p_1, \dots, p_n \in \mathbb{F}[s]$ denote the similarity invariants of A . Then,

$$\chi_A = \prod_{i=1}^n p_i. \quad (4.4.2)$$

Proposition 4.4.3. Let $A \in \mathbb{F}^{n \times n}$. Then, χ_A is monic and $\deg \chi_A = n$.

Let $A \in \mathbb{F}^{n \times n}$, and write the characteristic polynomial of A as

$$\chi_A(s) = s^n + \beta_{n-1}s^{n-1} + \dots + \beta_1s + \beta_0, \quad (4.4.3)$$

where $\beta_0, \dots, \beta_{n-1} \in \mathbb{F}$. The *eigenvalues* of A are the n possibly repeated roots $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ of χ_A , that is, the solutions of the *characteristic equation*

$$\chi_A(s) = 0. \quad (4.4.4)$$

It is often convenient to denote the eigenvalues of A by $\lambda_1(A), \dots, \lambda_n(A)$ or just $\lambda_1, \dots, \lambda_n$. This notation may be ambiguous, however, since it does not uniquely specify which eigenvalue is denoted by λ_i . If, however, every eigenvalue of A is real, then we employ the notational convention

$$\lambda_1 \geq \dots \geq \lambda_n, \quad (4.4.5)$$

and we define

$$\lambda_{\max}(A) \triangleq \lambda_1, \quad \lambda_{\min}(A) \triangleq \lambda_n. \quad (4.4.6)$$

Definition 4.4.4. Let $A \in \mathbb{F}^{n \times n}$. The *algebraic multiplicity* of an eigenvalue λ of A , denoted by $\text{amult}_A(\lambda)$, is the algebraic multiplicity of λ as a root of χ_A , that is,

$$\text{amult}_A(\lambda) \triangleq \text{mult}_{\chi_A}(\lambda). \quad (4.4.7)$$

The multiset consisting of the eigenvalues of A including their algebraic multiplicity, denoted by $\text{mspec}(A)$, is the *multispectrum* of A , that is,

$$\text{mspec}(A) \triangleq \text{mroots}(\chi_A). \quad (4.4.8)$$

Ignoring algebraic multiplicity, $\text{spec}(A)$ denotes the *spectrum* of A , that is,

$$\text{spec}(A) \triangleq \text{roots}(\chi_A). \quad (4.4.9)$$

Note that

$$\text{Szeros}(sI - A) = \text{spec}(A) \quad (4.4.10)$$

and

$$\text{mSzeros}(sI - A) = \text{mspec}(A). \quad (4.4.11)$$

If $\lambda \notin \text{spec}(A)$, then $\lambda \notin \text{roots}(\chi_A)$, and thus $\text{amult}_A(\lambda) = \text{mult}_{\chi_A}(\lambda) = 0$.

Let $A \in \mathbb{F}^{n \times n}$ and $\text{mroots}(\chi_A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\chi_A(s) = \prod_{i=1}^n (s - \lambda_i). \quad (4.4.12)$$

If $\mathbb{F} = \mathbb{R}$, then $\chi_A(s)$ has real coefficients, and thus the eigenvalues of A occur in complex conjugate pairs, that is, $\text{mroots}(\chi_A) = \text{mroots}(\overline{\chi_A})$. Now, let $\text{spec}(A) = \{\lambda_1, \dots, \lambda_r\}$, and, for all $i = 1, \dots, r$, let n_i denote the algebraic multiplicity of λ_i . Then,

$$\chi_A(s) = \prod_{i=1}^r (s - \lambda_i)^{n_i}. \quad (4.4.13)$$

The following result gives some basic properties of the spectrum of a matrix.

Proposition 4.4.5. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) $\chi_{A^T} = \chi_A$.
- ii) For all $s \in \mathbb{C}$, $\chi_{-A}(s) = (-1)^n \chi_A(-s)$.
- iii) $\text{mspec}(A^T) = \text{mspec}(A)$.
- iv) $\text{mspec}(\overline{A}) = \overline{\text{mspec}(A)}$.
- v) $\text{mspec}(A^*) = \overline{\text{mspec}(A)}$.
- vi) $0 \in \text{spec}(A)$ if and only if $\det A = 0$.
- vii) If $k \in \mathbb{N}$ or if A is nonsingular and $k \in \mathbb{Z}$, then

$$\text{mspec}(A^k) = \{\lambda^k: \lambda \in \text{mspec}(A)\}_{\text{ms}}. \quad (4.4.14)$$
- viii) If $\alpha \in \mathbb{F}$, then $\chi_{\alpha A + I}(s) = \chi_A(s - \alpha)$.
- ix) If $\alpha \in \mathbb{F}$, then $\text{mspec}(\alpha I + A) = \alpha + \text{mspec}(A)$.
- x) If $\alpha \in \mathbb{F}$, then $\text{mspec}(\alpha A) = \alpha \text{mspec}(A)$.
- xi) If A is Hermitian, then $\text{spec}(A) \subset \mathbb{R}$.
- xii) If A and B are similar, then $\chi_A = \chi_B$ and $\text{mspec}(A) = \text{mspec}(B)$.

Proof. To prove i), note that

$$\det(sI - A^T) = \det(sI - A)^T = \det(sI - A).$$

To prove ii), note that

$$\chi_{-A}(s) = \det(sI + A) = (-1)^n \det(-sI - A) = (-1)^n \chi_A(-s).$$

Next, *iii*) follows from *i*). Next, *iv*) follows from

$$\det(sI - \overline{A}) = \det(\overline{sI - A}) = \overline{\det(sI - A)},$$

while *v*) follows from *iii*) and *iv*).

Next, *vi*) follows from the fact that $\chi_A(0) = (-1)^n \det A$. To prove “ $\supseteq$ ” in *vii*), note that, if $\lambda \in \text{spec}(A)$ and $x \in \mathbb{C}^n$ is an eigenvector of A associated with λ (see Section 4.5), then $A^2x = A(Ax) = A(\lambda x) = \lambda Ax = \lambda^2x$. Similarly, if A is nonsingular, then $Ax = \lambda x$ implies that $A^{-1}x = \lambda^{-1}x$, and thus $A^{-2}x = \lambda^{-2}x$. Similar arguments apply to arbitrary $k \in \mathbb{Z}$. The reverse inclusion follows from the Jordan decomposition given by Theorem 5.3.3.

To prove *viii*), note that

$$\chi_{\alpha I + A}(s) = \det[sI - (\alpha I + A)] = \det[(s - \alpha)I - A] = \chi_A(s - \alpha).$$

Statement *ix*) follows immediately.

Statement *x*) is true for $\alpha = 0$. For $\alpha \neq 0$, it follows that

$$\chi_{\alpha A}(s) = \det(sI - \alpha A) = \alpha^{-1} \det[(s/\alpha)I - A] = \chi_A(s/\alpha).$$

To prove *xi*), assume that $A = A^*$, let $\lambda \in \text{spec}(A)$, and let $x \in \mathbb{C}^n$ be an eigenvector of A associated with λ . Then, $\lambda = x^*Ax/x^*x$, which is real. Finally, *xii*) is immediate. $\square$

The following result characterizes the coefficients of χ_A in terms of the eigenvalues of A .

Proposition 4.4.6. Let $A \in \mathbb{F}^{n \times n}$, let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, and, for all $i = 1, \dots, n$, let γ_i denote the sum of all $i \times i$ principal subdeterminants of A . Then, for all $i = 1, \dots, n-1$,

$$\gamma_i = \sum_{1 \leq j_1 < \dots < j_i \leq n} \lambda_{j_1} \cdots \lambda_{j_i}. \quad (4.4.15)$$

Furthermore, for all $i = 0, \dots, n-1$, the coefficient β_i of s^i in (4.4.3) is given by

$$\beta_i = (-1)^{n-i} \gamma_{n-i}. \quad (4.4.16)$$

In particular,

$$\beta_{n-1} = -\text{tr } A = -\sum_{i=1}^n \lambda_i, \quad (4.4.17)$$

$$\beta_{n-2} = \frac{1}{2}[(\text{tr } A)^2 - \text{tr } A^2] = \sum_{1 \leq j_1 < j_2 \leq n} \lambda_{j_1} \lambda_{j_2}, \quad (4.4.18)$$

$$\beta_1 = (-1)^{n-1} \text{tr } A^A = (-1)^{n-1} \sum_{1 \leq j_1 < \dots < j_{n-1} \leq n} \lambda_{j_1} \cdots \lambda_{j_{n-1}} = (-1)^{n-1} \sum_{i=1}^n \det A_{[i;i]}, \quad (4.4.19)$$

$$\beta_0 = (-1)^n \det A = (-1)^n \prod_{i=1}^n \lambda_i. \quad (4.4.20)$$

Proof. The expression for γ_i given by (4.4.15) follows from the factored form of $\chi_A(s)$ given by (4.4.12), while the expression for β_i given by (4.4.16) follows by examining the cofactor expansion (2.7.16) of $\det(sI - A)$. For details, see [998, p. 495]. Equation (4.4.17) follows from (4.4.16) and the fact that the $(n-1) \times (n-1)$ principal subdeterminants of A are the diagonal entries $A_{(i,i)}$. Using

$$\sum_{i=1}^n \lambda_i^2 = \left(\sum_{i=1}^n \lambda_i \right)^2 - 2 \sum \lambda_{j_1} \lambda_{j_2},$$

where the third summation is taken over all pairs of elements of $\text{mspec}(A)$, and (4.4.17) yields (4.4.18). Next, if A is nonsingular, then $\chi_{A^{-1}}(s) = (-s)^n (\det A^{-1}) \chi_A(1/s)$. Using (4.4.3) with s replaced by $1/s$ and (4.4.17), it follows that $\text{tr } A^{-1} = (-1)^{n-1} (\det A^{-1}) \beta_1$, and, hence, (4.4.19) is satisfied. Using continuity for the case in which A is singular yields (4.4.19) for arbitrary A . Finally, $\beta_0 = \chi_A(0) = \det(0I - A) = (-1)^n \det A$, which verifies (4.4.20). $\square$

From the definition of the adjugate of a matrix it follows that $(sI - A)^A \in \mathbb{F}^{n \times n}[s]$ is a monic polynomial matrix of degree $n-1$ of the form

$$(sI - A)^A = s^{n-1}I + s^{n-2}B_{n-2} + \cdots + sB_1 + B_0, \quad (4.4.21)$$

where $B_0, B_1, \dots, B_{n-2} \in \mathbb{F}^{n \times n}$. Since $(sI - A)^A$ is regular, it follows from Proposition 4.2.6 that $(sI - A)^A$ is a nonsingular polynomial matrix. The matrix $(sI - A)^{-1}$ is the *resolvent* of A , which is given by

$$(sI - A)^{-1} = \frac{1}{\chi_A(s)} (sI - A)^A. \quad (4.4.22)$$

Therefore,

$$(sI - A)^{-1} = \frac{s^{n-1}}{\chi_A(s)} I + \frac{s^{n-2}}{\chi_A(s)} B_{n-2} + \cdots + \frac{s}{\chi_A(s)} B_1 + \frac{1}{\chi_A(s)} B_0. \quad (4.4.23)$$

The next result is the *Cayley-Hamilton theorem*, which shows that every matrix is a “root” of its characteristic polynomial.

Theorem 4.4.7. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\chi_A(A) = 0. \quad (4.4.24)$$

Proof. Define $P, Q \in \mathbb{F}^{n \times n}[s]$ by $P(s) \triangleq \chi_A(s)I$ and $Q(s) \triangleq (sI - A)^A$. Then, (4.4.22) implies that $P(s) = Q(s)(sI - A)$. It thus follows from Lemma 4.2.2 that $P_R(A) = 0$. Furthermore, $\chi_A(A) = P(A) = P_R(A)$. Hence, $\chi_A(A) = 0$. $\square$

In the notation of (4.4.13), it follows from Theorem 4.4.7 that

$$\prod_{i=1}^r (\lambda_i I - A)^{n_i} = 0. \quad (4.4.25)$$

Lemma 4.4.8. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\frac{d}{ds} \chi_A(s) = \operatorname{tr}[(sI - A)^A] = \sum_{i=1}^n \det(sI - A_{[i;i]}). \quad (4.4.26)$$

Proof. It follows from (4.4.19) that $\frac{d}{ds} \chi_A(s)|_{s=0} = \beta_1 = (-1)^{n-1} \operatorname{tr} A^A$. Hence,

$$\begin{aligned} \frac{d}{ds} \chi_A(s) &= \frac{d}{dz} \det[(s+z)I - A] \Big|_{z=0} = \frac{d}{dz} \det[zI - (-sI + A)] \Big|_{z=0} \\ &= (-1)^{n-1} \operatorname{tr}[(-sI + A)^A] = \operatorname{tr}[(sI - A)^A]. \quad \square \end{aligned}$$

The following result, known as *Leverrier's algorithm*, provides a recursive formula for the coefficients $\beta_0, \dots, \beta_{n-1}$ of χ_A and $B_0, \dots, B_{n-2}$ of $(sI - A)^A$.

Proposition 4.4.9. Let $A \in \mathbb{F}^{n \times n}$, let χ_A be given by (4.4.3), and let $(sI - A)^A$ be given by (4.4.21). Then, $\beta_{n-1}, \dots, \beta_0$ and $B_{n-2}, \dots, B_0$ are given by

$$\beta_k = \frac{1}{k-n} \operatorname{tr} AB_k, \quad k = n-1, \dots, 0, \quad (4.4.27)$$

$$B_{k-1} = AB_k + \beta_k I, \quad k = n-1, \dots, 1, \quad (4.4.28)$$

where $B_{n-1} = I$.

Proof. Since $(sI - A)(sI - A)^A = \chi_A(s)I$, it follows that

$$\begin{aligned} s^n I + s^{n-1}(B_{n-2} - A) + s^{n-2}(B_{n-3} - AB_{n-2}) + \dots + s(B_0 - AB_1) - AB_0 \\ = (s^n + \beta_{n-1}s^{n-1} + \dots + \beta_1 s + \beta_0)I. \end{aligned}$$

Equating coefficients of powers of s yields (4.4.28) along with $-AB_0 = \beta_0 I$. Taking the trace of this last identity yields $\beta_0 = -\frac{1}{n} \operatorname{tr} AB_0$, which confirms (4.4.27) for $k = 0$. Next, using (4.4.26) and (4.4.21), it follows that

$$\frac{d}{ds} \chi_A(s) = \sum_{k=1}^n k \beta_k s^{k-1} = \sum_{k=1}^n (\operatorname{tr} B_{k-1}) s^{k-1},$$

where $B_{n-1} \triangleq I_n$ and $\beta_n \triangleq 1$. Equating powers of s , it follows that $k\beta_k = \operatorname{tr} B_{k-1}$ for all $k = 1, \dots, n$. Now, (4.4.28) implies that $k\beta_k = \operatorname{tr}(AB_k + \beta_k I)$ for all $k = 1, \dots, n-1$, which implies (4.4.27). $\square$

Proposition 4.4.10. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$, and assume that $m \leq n$. Then,

$$\chi_{AB}(s) = s^{n-m} \chi_{BA}(s). \quad (4.4.29)$$

Consequently, $\operatorname{mspec}(AB) = \operatorname{mspec}(BA) \cup \{0, \dots, 0\}_{\operatorname{ms}}$, (4.4.30)

where the multiset $\{0, \dots, 0\}_{\operatorname{ms}}$ contains $n - m$ 0's.

Proof. First note that

$$\begin{bmatrix} 0_{m \times m} & 0_{m \times n} \\ A & AB \end{bmatrix} = \begin{bmatrix} I_m & -B \\ 0_{n \times m} & I_n \end{bmatrix} \begin{bmatrix} BA & 0_{m \times n} \\ A & 0_{n \times n} \end{bmatrix} \begin{bmatrix} I_m & B \\ 0_{n \times m} & I_n \end{bmatrix},$$

which shows that $\begin{bmatrix} 0_{m \times m} & 0_{m \times n} \\ A & AB \end{bmatrix}$ and $\begin{bmatrix} BA & 0_{m \times n} \\ A & 0_{n \times n} \end{bmatrix}$ are similar. It thus follows from xi) of Proposition 4.4.5 that $s^m \chi_{AB}(s) = s^n \chi_{BA}(s)$, which implies (4.4.29). Finally, (4.4.30) follows immediately from (4.4.29). $\square$

If $n = m$, then Proposition 4.4.10 specializes to the following result.

Corollary 4.4.11. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\chi_{AB} = \chi_{BA}. \quad (4.4.31)$$

Consequently,

$$\text{mspec}(AB) = \text{mspec}(BA). \quad (4.4.32)$$

We define the *spectral abscissa* of $A \in \mathbb{F}^{n \times n}$ by

$$\text{spabs}(A) \triangleq \max\{\text{Re } \lambda: \lambda \in \text{spec}(A)\} \quad (4.4.33)$$

and the *spectral radius* of $A \in \mathbb{F}^{n \times n}$ by

$$\text{sprad}(A) \triangleq \max\{|\lambda|: \lambda \in \text{spec}(A)\}. \quad (4.4.34)$$

Let $A \in \mathbb{F}^{n \times n}$. Then, $\nu_-(A)$, $\nu_0(A)$, and $\nu_+(A)$ denote the number of eigenvalues of A counting algebraic multiplicity having, respectively, negative, zero, and positive real part. Define the *inertia* of A by

$$\text{In } A \triangleq \begin{bmatrix} \nu_-(A) \\ \nu_0(A) \\ \nu_+(A) \end{bmatrix} \quad (4.4.35)$$

and the *signature* of A by

$$\text{sig } A \triangleq \nu_+(A) - \nu_-(A). \quad (4.4.36)$$

Note that $\text{spabs}(A) < 0$ if and only if $\nu_-(A) = n$, while $\text{spabs}(A) = 0$ if and only if $\nu_+(A) = 0$.

4.5 Eigenvectors

Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \mathbb{C}$ be an eigenvalue of A . Then, $\chi_A(\lambda) = \det(\lambda I - A) = 0$, and thus $\lambda I - A \in \mathbb{C}^{n \times n}$ is singular. Furthermore, $\mathcal{N}(\lambda I - A)$ is a nontrivial subspace of $\mathbb{C}^n$, that is, $\text{def}(\lambda I - A) > 0$. If $x \in \mathcal{N}(\lambda I - A)$, that is, $Ax = \lambda x$, and $x \neq 0$, then x is an *eigenvector of A associated with λ* . By definition, all eigenvectors are nonzero. Note that, if A and λ are real, then there exists a real eigenvector associated with λ .

Definition 4.5.1. The *geometric multiplicity* of $\lambda \in \text{spec}(A)$, denoted by $\text{gmult}_A(\lambda)$, is the number of linearly independent eigenvectors associated with λ , that is,

$$\text{gmult}_A(\lambda) \triangleq \text{def}(\lambda I - A). \quad (4.5.1)$$

By convention, if $\lambda \notin \text{spec}(A)$, then $\text{gmult}_A(\lambda) \triangleq 0$.

Proposition 4.5.2. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \text{spec}(A)$. Then, the following statements hold:

- i) $\text{rank}(\lambda I - A) + \text{gmult}_A(\lambda) = n$.
- ii) $\text{def } A = \text{gmult}_A(0)$.
- iii) $\text{rank } A + \text{gmult}_A(0) = n$.

The spectral properties of normal matrices deserve special attention.

Lemma 4.5.3. Let $A \in \mathbb{F}^{n \times n}$ be normal, let $\lambda \in \text{spec}(A)$, and let $x \in \mathbb{C}^n$ be an eigenvector of A associated with λ . Then, x is an eigenvector of A^* associated with $\bar{\lambda} \in \text{spec}(A^*)$.

Proof. Since $\lambda \in \text{spec}(A)$, statement v) of Proposition 4.4.5 implies that $\bar{\lambda} \in \text{spec}(A^*)$. Next, since x and λ satisfy $Ax = \lambda x$, $x^*A^* = \bar{\lambda}x^*$, and $AA^* = A^*A$, it follows that

$$\begin{aligned} (A^*x - \bar{\lambda}x)^*(A^*x - \bar{\lambda}x) &= x^*AA^*x - \bar{\lambda}x^*Ax - \lambda x^*A^*x + \lambda\bar{\lambda}x^*x \\ &= x^*A^*Ax - \lambda\bar{\lambda}x^*x - \lambda\bar{\lambda}x^*x + \lambda\bar{\lambda}x^*x \\ &= \lambda\bar{\lambda}x^*x - \lambda\bar{\lambda}x^*x = 0. \end{aligned}$$

Hence, $A^*x = \bar{\lambda}x$. □

Proposition 4.5.4. Let $A \in \mathbb{F}^{n \times n}$. Then, eigenvectors associated with distinct eigenvalues of A are linearly independent. If, in addition, A is normal, then these eigenvectors are mutually orthogonal.

Proof. Let $\lambda_1, \lambda_2 \in \text{spec}(A)$ be distinct with associated eigenvectors $x_1, x_2 \in \mathbb{C}^n$. Suppose that x_1 and x_2 are linearly dependent, that is, $x_1 = \alpha x_2$, where $\alpha \in \mathbb{C}$ and $\alpha \neq 0$. Then, $Ax_1 = \lambda_1 x_1 = \lambda_1 \alpha x_2$, while also $Ax_1 = A\alpha x_2 = \alpha \lambda_2 x_2$. Hence, $\alpha(\lambda_1 - \lambda_2)x_2 = 0$, which contradicts $\alpha \neq 0$. Since pairwise linear independence does not imply the linear independence of larger sets, next, let $\lambda_1, \lambda_2, \lambda_3 \in \text{spec}(A)$ be distinct with associated eigenvectors $x_1, x_2, x_3 \in \mathbb{C}^n$. Suppose that x_1, x_2, x_3 are linearly dependent. In this case, there exist $a_1, a_2, a_3 \in \mathbb{C}$, not all zero, such that $a_1 x_1 + a_2 x_2 + a_3 x_3 = 0$. If $a_1 = 0$, then $a_2 x_2 + a_3 x_3 = 0$. However, $\lambda_2 \neq \lambda_3$ implies that x_2 and x_3 are linearly independent, which in turn implies that $a_2 = 0$ and $a_3 = 0$. Since a_1, a_2, a_3 are not all zero, it follows that $a_1 \neq 0$. Therefore, $x_1 = \alpha x_2 + \beta x_3$, where $\alpha \triangleq -a_2/a_1$ and $\beta \triangleq -a_3/a_1$ are not both zero. Thus, $Ax_1 = A(\alpha x_2 + \beta x_3) = \alpha Ax_2 + \beta Ax_3 = \alpha \lambda_2 x_2 + \beta \lambda_3 x_3$. However, $Ax_1 = \lambda_1 x_1 = \lambda_1(\alpha x_2 + \beta x_3) = \alpha \lambda_1 x_2 + \beta \lambda_1 x_3$. Subtracting these relations yields $0 = \alpha(\lambda_1 - \lambda_2)x_2 + \beta(\lambda_1 - \lambda_3)x_3$. Since x_2 and x_3 are linearly independent, it follows that $\alpha(\lambda_1 - \lambda_2) = 0$ and $\beta(\lambda_1 - \lambda_3) = 0$. Since α and β are not both zero, it follows that $\lambda_1 = \lambda_2$ or $\lambda_1 = \lambda_3$, which contradicts the assumption that $\lambda_1, \lambda_2, \lambda_3$ are distinct. The same arguments apply to sets of four or more eigenvectors.

Now, suppose that A is normal, and let $\lambda_1, \lambda_2 \in \text{spec}(A)$ be distinct eigenvalues with associated eigenvectors $x_1, x_2 \in \mathbb{C}^n$. Then, by Lemma 4.5.3, $Ax_1 = \lambda_1 x_1$ implies that $A^*x_1 = \bar{\lambda}_1 x_1$. Consequently, $x_1^*A = \lambda_1 x_1^*$, which implies that $x_1^*Ax_2 = \lambda_1 x_1^*x_2$. Furthermore, $x_1^*Ax_2 = \lambda_2 x_1^*x_2$. It thus follows that $0 = (\lambda_1 - \lambda_2)x_1^*x_2$.

Hence, $\lambda_1 \neq \lambda_2$ implies that $x_1^* x_2 = 0$. $\square$

If $A \in \mathbb{R}^{n \times n}$ is symmetric, then Lemma 4.5.3 is not needed and the proof of Proposition 4.5.4 is simpler. In this case, it follows from x of Proposition 4.4.5 that $\lambda_1, \lambda_2 \in \text{spec}(A)$ are real, and thus associated eigenvectors $x_1 \in \mathcal{N}(\lambda_1 I - A)$ and $x_2 \in \mathcal{N}(\lambda_2 I - A)$ can be chosen to be real. Hence, $Ax_1 = \lambda_1 x_1$ and $Ax_2 = \lambda_2 x_2$ imply that $x_2^T A x_1 = \lambda_1 x_2^T x_1$ and $x_1^T A x_2 = \lambda_2 x_1^T x_2$. Since $x_1^T A x_2 = x_2^T A^T x_1 = x_2^T A x_1$ and $x_1^T x_2 = x_2^T x_1$, it follows that $(\lambda_1 - \lambda_2)x_1^T x_2 = 0$. Since $\lambda_1 \neq \lambda_2$, it follows that $x_1^T x_2 = 0$.

4.6 The Minimal Polynomial

Theorem 4.4.7 showed that every square matrix $A \in \mathbb{F}^{n \times n}$ is a root of its characteristic polynomial. However, there may be polynomials of degree less than n having A as a root. In fact, the following result shows that there exists a unique monic polynomial that has A as a root and that divides all polynomials that have A as a root.

Theorem 4.6.1. Let $A \in \mathbb{F}^{n \times n}$. Then, there exists a unique monic polynomial $\mu_A \in \mathbb{F}[s]$ of minimal degree such that $\mu_A(A) = 0$. Furthermore, $\deg \mu_A \leq n$, and μ_A divides every polynomial $p \in \mathbb{F}[s]$ satisfying $p(A) = 0$.

Proof. Since $\chi_A(A) = 0$ and $\deg \chi_A = n$, it follows that there exists a minimal positive integer $n_0 \leq n$ such that there exists a monic polynomial $p_0 \in \mathbb{F}[s]$ satisfying $p_0(A) = 0$ and $\deg p_0 = n_0$. Let $p \in \mathbb{F}[s]$ satisfy $p(A) = 0$. Then, by Lemma 4.1.2, there exist polynomials $q, r \in \mathbb{F}[s]$ such that $p = qp_0 + r$ and $\deg r < \deg p_0$. However, $p(A) = p_0(A) = 0$ implies that $r(A) = 0$. If $r \neq 0$, then r can be normalized to obtain a monic polynomial of degree less than n_0 , which contradicts the definition n_0 . Hence, $r = 0$, which implies that p_0 divides p . This proves existence.

Now, suppose there exist two monic polynomials $p_0, \hat{p}_0 \in \mathbb{F}[s]$ of degree n_0 and such that $p_0(A) = \hat{p}_0(A) = 0$. By the previous argument, p_0 divides $\hat{p}_0$, and vice versa. Therefore, p_0 is a constant multiple of $\hat{p}_0$. Since p_0 and $\hat{p}_0$ are both monic, it follows that $p_0 = \hat{p}_0$. This proves uniqueness. Denote this polynomial by μ_A . $\square$

The monic polynomial μ_A of smallest degree having A as a root is the *minimal polynomial* of A .

The following result relates the characteristic and minimal polynomials of $A \in \mathbb{F}^{n \times n}$ to the similarity invariants of A . Note that $\text{rank}(sI - A) = n$, so that A has n similarity invariants $p_1, \dots, p_n \in \mathbb{F}[s]$. In this case, (4.3.1) becomes

$$sI - A = S_1(s) \begin{bmatrix} p_1(s) & & 0 \\ & \ddots & \\ 0 & & p_n(s) \end{bmatrix} S_2(s), \quad (4.6.1)$$

where $S_1, S_2 \in \mathbb{F}^{n \times n}[s]$ are unimodular and p_i divides p_{i+1} for all $i = 1, \dots, n-1$.

Proposition 4.6.2. Let $A \in \mathbb{F}^{n \times n}$, and let $p_1, \dots, p_n \in \mathbb{F}[s]$ be the similarity invariants of A , where p_i divides p_{i+1} for all $i = 1, \dots, n-1$. Then,

$$\chi_A = \prod_{i=1}^n p_i \quad (4.6.2)$$

and

$$\mu_A = p_n. \quad (4.6.3)$$

Proof. Using Theorem 4.3.2 and (4.6.1), it follows that

$$\chi_A(s) = \det(sI - A) = [\det S_1(s)] [\det S_2(s)] \prod_{i=1}^n p_i(s).$$

Since S_1 and S_2 are unimodular and χ_A and $p_1, \dots, p_n$ are monic, it follows that $[\det S_1(s)][\det S_2(s)] = 1$, which proves (4.6.2).

To prove (4.6.3), first note that it follows from Theorem 4.3.2 that $\chi_A = \Delta_{n-1} p_n$, where $\Delta_{n-1} \in \mathbb{F}[s]$ is the greatest common divisor of all $(n-1) \times (n-1)$ subdeterminants of $sI - A$. Since the $(n-1) \times (n-1)$ subdeterminants of $sI - A$ are the entries of $\pm(sI - A)^A$, it follows that Δ_{n-1} divides every entry of $(sI - A)^A$. Hence, there exists a polynomial matrix $P \in \mathbb{F}^{n \times n}[s]$ such that $(sI - A)^A = \Delta_{n-1}(s)P(s)$. Furthermore, since $(sI - A)^A(sI - A) = \chi_A(s)I$, it follows that $\Delta_{n-1}(s)P(s)(sI - A) = \chi_A(s)I = \Delta_{n-1}(s)p_n(s)I$, and thus $P(s)(sI - A) = p_n(s)I$. Lemma 4.2.2 now implies that $p_n(A) = 0$.

Since $p_n(A) = 0$, it follows from Theorem 4.6.1 that μ_A divides p_n . Hence, let $q \in \mathbb{F}[s]$ be the monic polynomial satisfying $p_n = q\mu_A$. Furthermore, since $\mu_A(A) = 0$, it follows from Corollary 4.2.3 that there exists a polynomial matrix $Q \in \mathbb{F}^{n \times n}[s]$ such that $\mu_A(s)I = Q(s)(sI - A)$. Thus, $P(s)(sI - A) = p_n(s)I = q(s)\mu_A(s)I = q(s)Q(s)(sI - A)$, which implies that $P = qQ$. Thus, q divides every entry of P . However, since P is obtained by dividing $(sI - A)^A$ by the greatest common divisor of all of its entries, it follows that the greatest common divisor of the entries of P is 1. Hence, $q = 1$, which implies that $p_n = \mu_A$, which proves (4.6.3). $\square$

Proposition 4.6.2 shows that μ_A divides χ_A , which is also a consequence of Theorem 4.4.7 and Theorem 4.6.1. Proposition 4.6.2 also shows that $\mu_A = \chi_A$ if and only if $p_1 = \dots = p_{n-1} = 1$, that is, if and only if $p_n = \chi_A$ is the only nonconstant similarity invariant of A . Note that, in general, it follows from (4.6.2) that $\sum_{i=1}^n \deg p_i = n$.

Finally, note that the similarity invariants of the $n \times n$ identity matrix I_n are given by $p_i(s) = s - 1$ for all $i = 1, \dots, n$. Thus, $\chi_{I_n}(s) = (s - 1)^n$ and $\mu_{I_n}(s) = s - 1$.

Proposition 4.6.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A and B are similar. Then,

$$\mu_A = \mu_B. \quad (4.6.4)$$

4.7 Rational Transfer Functions and the Smith-McMillan Decomposition

We now turn our attention to rational functions.

Definition 4.7.1. The set $\mathbb{F}(s)$ of *rational functions* consists of functions $g: \mathbb{C} \setminus \mathcal{S} \mapsto \mathbb{C}$, where $g(s) = p(s)/q(s)$, $p, q \in \mathbb{F}[s]$, $q \neq 0$, and $\mathcal{S} \triangleq \text{roots}(q)$. The rational function g is *strictly proper*, *proper*, *exactly proper*, *improper*, respectively, if $\deg p < \deg q$, $\deg p \leq \deg q$, $\deg p = \deg q$, $\deg p > \deg q$. If p and q are coprime, then the *zeros* of g are the elements of $\text{mroots}(p)$, while the *poles* of g are the elements of $\text{mroots}(q)$. The set of proper rational functions is denoted by $\mathbb{F}_{\text{prop}}(s)$. The *relative degree* of $g \in \mathbb{F}_{\text{prop}}(s)$, denoted by $\text{reldeg } g$, is $\deg q - \deg p$.

Definition 4.7.2. The set $\mathbb{F}^{l \times m}(s)$ of *rational transfer functions* consists of matrices whose entries are elements of $\mathbb{F}(s)$. The rational transfer function $G \in \mathbb{F}^{l \times m}(s)$ is *strictly proper* if every entry of G is strictly proper, *proper* if every entry of G is proper, *exactly proper* if every entry of G is proper and at least one entry of G is exactly proper, and *improper* if at least one entry of G is improper. The set of proper rational transfer functions is denoted by $\mathbb{F}_{\text{prop}}^{l \times m}(s)$.

Definition 4.7.3. Let $G \in \mathbb{F}_{\text{prop}}^{l \times m}(s)$. Then, the *relative degree* of G , denoted by $\text{reldeg } G$, is defined by

$$\text{reldeg } G \triangleq \min_{\substack{i=1, \dots, l \\ j=1, \dots, m}} \text{reldeg } G_{(i,j)}. \quad (4.7.1)$$

By writing $(sI - A)^{-1}$ as

$$(sI - A)^{-1} = \frac{1}{\chi_A(s)} (sI - A)^A, \quad (4.7.2)$$

it follows from (4.4.21) that $(sI - A)^{-1}$ is a strictly proper rational transfer function. In fact, for all $i = 1, \dots, n$,

$$\text{reldeg } [(sI - A)^{-1}]_{(i,i)} = 1, \quad (4.7.3)$$

and thus

$$\text{reldeg } (sI - A)^{-1} = 1. \quad (4.7.4)$$

The following definition is an extension of Definition 4.2.4 to rational transfer functions.

Definition 4.7.4. Let $G \in \mathbb{F}^{l \times m}(s)$, and, for all $i = 1, \dots, l$ and $j = 1, \dots, m$, let $G_{(i,j)} = p_{ij}/q_{ij}$, where $q_{ij} \neq 0$, and $p_{ij}, q_{ij} \in \mathbb{F}[s]$ are coprime. Then, the *poles* of G are the elements of the set

$$\text{poles}(G) \triangleq \bigcup_{i,j=1}^{l,m} \text{roots}(q_{ij}), \quad (4.7.5)$$

and the *blocking zeros* of G are the elements of the set

$$\text{bzeros}(G) \triangleq \bigcap_{i,j=1}^{l,m} \text{roots}(p_{ij}). \quad (4.7.6)$$

Finally, the rank of G is the nonnegative integer

$$\text{rank } G \triangleq \max_{s \in \mathbb{C} \setminus \text{poles}(G)} \text{rank } G(s). \quad (4.7.7)$$

The following result provides a canonical form, known as the *Smith-McMillan form*, for rational transfer functions under unimodular transformation.

Theorem 4.7.5. Let $G \in \mathbb{F}^{l \times m}(s)$, and let $r \triangleq \text{rank } G$. Then, there exist unimodular matrices $S_1 \in \mathbb{F}^{l \times l}[s]$ and $S_2 \in \mathbb{F}^{m \times m}[s]$ and monic polynomials $p_1, \dots, p_r, q_1, \dots, q_r \in \mathbb{F}[s]$ such that p_i and q_i are coprime for all $i = 1, \dots, r$, p_i divides p_{i+1} for all $i = 1, \dots, r-1$, q_{i+1} divides q_i for all $i = 1, \dots, r-1$, and

$$G = S_1 \begin{bmatrix} p_1/q_1 & & & \\ & \ddots & & \\ & & p_r/q_r & \\ & & & 0_{(l-r) \times (m-r)} \end{bmatrix} S_2. \quad (4.7.8)$$

Proof. Let n_{ij}/d_{ij} denote the (i, j) entry of G , where $n_{ij}, d_{ij} \in \mathbb{F}[s]$ are coprime, and let $d \in \mathbb{F}[s]$ denote the least common multiple of d_{ij} for all $i = 1, \dots, l$ and $j = 1, \dots, m$. From Theorem 4.3.2 it follows that the polynomial matrix dG has the Smith form $\text{diag}(\hat{p}_1, \dots, \hat{p}_r, 0, \dots, 0)$, where $\hat{p}_1, \dots, \hat{p}_r \in \mathbb{F}[s]$ and $\hat{p}_i$ divides $\hat{p}_{i+1}$ for all $i = 1, \dots, r-1$. Now, divide this Smith form by d and express every rational function $\hat{p}_i/d$ in coprime form p_i/q_i so that p_i divides p_{i+1} for all $i = 1, \dots, r-1$ and q_{i+1} divides q_i for all $i = 1, \dots, r-1$. $\square$

Proposition 4.7.6. Let $G \in \mathbb{F}^{l \times m}(s)$, and assume that there exist unimodular matrices $S_1 \in \mathbb{F}^{l \times l}[s]$ and $S_2 \in \mathbb{F}^{m \times m}[s]$ and monic polynomials $p_1, \dots, p_r, q_1, \dots, q_r \in \mathbb{F}[s]$ such that p_i and q_i are coprime for all $i = 1, \dots, r$ and such that (4.7.8) holds. Then, $\text{rank } G = r$.

Proposition 4.7.7. Let $G \in \mathbb{F}^{n \times m}[s]$, and let $r \triangleq \text{rank } G$. Then, r is the largest order of all nonsingular submatrices of G .

Proposition 4.7.8. Let $G \in \mathbb{F}^{n \times m}(s)$, and let $\mathcal{S} \subset \mathbb{C}$ be a finite set such that $\text{poles}(G) \subseteq \mathcal{S}$. Then,

$$\text{rank } G = \max_{s \in \mathbb{C} \setminus \mathcal{S}} \text{rank } G(s). \quad (4.7.9)$$

Let $g_1, \dots, g_r \in \mathbb{F}^n(s)$. Then, $g_1, \dots, g_r$ are *linearly independent* if $\alpha_1, \dots, \alpha_r \in \mathbb{F}[s]$ and $\sum_{n=1}^r \alpha_i g_i = 0$ imply that $\alpha_1 = \dots = \alpha_r = 0$. Equivalently, $g_1, \dots, g_r$ are *linearly independent* if $\alpha_1, \dots, \alpha_r \in \mathbb{F}(s)$ and $\sum_{n=1}^r \alpha_i g_i = 0$ imply that $\alpha_1 = \dots = \alpha_r = 0$. In other words, the coefficients α_i can be either polynomials or rational functions.

Proposition 4.7.9. Let $G \in \mathbb{F}^{l \times m}(s)$. Then, $\text{rank } G$ is equal to the number of linearly independent columns of G .

Since $G \in \mathbb{F}^{l \times m}[s] \subset \mathbb{F}^{l \times m}(s)$, Proposition 4.7.9 applies to polynomial matrices.

Definition 4.7.10. Let $G \in \mathbb{F}^{l \times m}(s)$, assume that $G \neq 0$, let $r \triangleq \text{rank } G$, and let $p_1, \dots, p_r, q_1, \dots, q_r \in \mathbb{F}[s]$ be given by Theorem 4.7.5. Then, the *McMillan degree* $\text{Mcdeg } G$ of G is defined by

$$\text{Mcdeg } G \triangleq \sum_{i=1}^r \deg q_i. \quad (4.7.10)$$

Furthermore, the *transmission zeros* of G are the elements of the set

$$\text{tzeros}(G) \triangleq \text{roots}(p_r). \quad (4.7.11)$$

Proposition 4.7.11. Let $G \in \mathbb{F}^{l \times m}(s)$, assume that $G \neq 0$, and assume that G has the Smith-McMillan form (4.7.8). Then,

$$\text{poles}(G) = \text{roots}(q_1) \quad (4.7.12)$$

and

$$\text{bzeros}(G) = \text{roots}(p_1). \quad (4.7.13)$$

Note that

$$\text{bzeros}(G) \subseteq \text{tzeros}(G). \quad (4.7.14)$$

Furthermore, we define the multisets

$$\text{mpoles}(G) \triangleq \bigcup_{i=1}^r \text{mroots}(q_i), \quad (4.7.15)$$

$$\text{mtzeros}(G) \triangleq \bigcup_{i=1}^r \text{mroots}(p_i), \quad (4.7.16)$$

$$\text{mbzeros}(G) \triangleq \text{mroots}(p_1). \quad (4.7.17)$$

Note that

$$\text{mbzeros}(G) \subseteq \text{mtzeros}(G). \quad (4.7.18)$$

If $G = 0$, then these multisets as well as the sets $\text{poles}(G)$, $\text{tzeros}(G)$, and $\text{bzeros}(G)$ are empty.

Proposition 4.7.12. Let $G \in \mathbb{F}_{\text{prop}}^{l \times m}(s)$, assume that $G \neq 0$, let $z \in \mathbb{C}$, and assume that z is not a pole of G . Then, z is a transmission zero of G if and only if $\text{rank } G(z) < \text{rank } G$. Furthermore, z is a blocking zero of G if and only if $G(z) = 0$.

The following example shows that a pole of G can also be a transmission zero of G .

Example 4.7.13. Define $G \in \mathbb{R}_{\text{prop}}^{2 \times 2}(s)$ by

$$G(s) = \begin{bmatrix} \frac{1}{(s+1)^2} & \frac{1}{(s+1)(s+2)} \\ \frac{1}{(s+1)(s+2)} & \frac{s+3}{(s+2)^2} \end{bmatrix}.$$

Then, $\text{rank } G = 2$. Furthermore,

$$G(s) = S_1(s) \begin{bmatrix} \frac{1}{(s+1)^2(s+2)^2} & 0 \\ 0 & s+2 \end{bmatrix} S_2(s),$$

where $S_1, S_2 \in \mathbb{R}^{2 \times 2}[s]$ are the unimodular matrices

$$S_1(s) = \begin{bmatrix} (s+2)(s^3 + 4s^2 + 5s + 1) & 1 \\ (s+1)(s^3 + 5s^2 + 8s + 3) & 1 \end{bmatrix}$$

and

$$S_2(s) = \begin{bmatrix} -(s+2) & (s+1)(s^2 + 3s + 1) \\ 1 & -s(s+2) \end{bmatrix}.$$

Hence, the McMillan degree of G is 4, the poles of G are -1 and -2 , the transmission zero of G is -2 , and G has no blocking zeros. Note that -2 is both a pole and a transmission zero of G . Note also that, although G is strictly proper, the Smith-McMillan form of G is improper.

Let $G \in \mathbb{F}_{\text{prop}}^{l \times m}(s)$. A factorization of G of the form

$$G(s) = N(s)D^{-1}(s), \quad (4.7.19)$$

where $N \in \mathbb{F}^{l \times m}[s]$ and $D \in \mathbb{F}^{m \times m}[s]$, is a *right polynomial fraction description* of G . We say that N and D are *right coprime* if every $R \in \mathbb{F}^{m \times m}[s]$ that right divides both N and D is unimodular. In this case, (4.7.19) is a *coprime right polynomial fraction description* of G .

Theorem 4.7.14. Let $N \in \mathbb{F}^{l \times m}[s]$ and $D \in \mathbb{F}^{m \times m}[s]$. Then, the following statements are equivalent:

- i) N and D are right coprime.
- ii) There exist $X \in \mathbb{F}^{m \times l}[s]$ and $Y \in \mathbb{F}^{m \times m}[s]$ such that

$$XN + YD = I. \quad (4.7.20)$$

- iii) For all $s \in \mathbb{C}$,

$$\text{rank} \begin{bmatrix} N(s) \\ D(s) \end{bmatrix} = m. \quad (4.7.21)$$

Proof. See [1150, p. 297]. □

Equation (4.7.20) is the *Bezout identity*.

The following result shows that all coprime right polynomial fraction descriptions of a proper rational transfer function G are related by a unimodular

transformation.

Proposition 4.7.15. Let $G \in \mathbb{F}_{\text{prop}}^{l \times m}(s)$, let $N, \hat{N} \in \mathbb{F}^{l \times m}[s]$, let $D, \hat{D} \in \mathbb{F}^{m \times m}[s]$, and assume that $G = ND^{-1} = \hat{N}\hat{D}^{-1}$. Then, there exists a unimodular matrix $R \in \mathbb{F}^{m \times m}[s]$ such that $N = \hat{N}R$ and $D = \hat{D}R$.

Proof. See [1150, p. 298]. $\square$

The following result uses the Smith-McMillan form to show that every proper rational transfer function has a coprime right polynomial fraction description.

Proposition 4.7.16. Let $G \in \mathbb{F}_{\text{prop}}^{l \times m}(s)$. Then, G has a coprime right polynomial fraction description. If, in addition, $G(s) = N(s)D^{-1}(s)$, where $N \in \mathbb{F}^{l \times m}[s]$ and $D \in \mathbb{F}^{m \times m}[s]$, is a coprime right polynomial fraction description of G , then

$$\text{Szeros}(N) = \text{tzeros}(G) \quad (4.7.22)$$

and

$$\text{Szeros}(D) = \text{poles}(G). \quad (4.7.23)$$

Proof. Note that (4.7.8) can be written as

$$\begin{aligned} G &= S_1 \begin{bmatrix} p_1/q_1 & & & 0 \\ & \ddots & & \\ & & p_r/q_r & \\ 0 & & & 0_{(l-r) \times (m-r)} \end{bmatrix} S_2 \\ &= S_1 \begin{bmatrix} p_1 & & & 0 \\ & \ddots & & \\ & & p_r & \\ 0 & & & 0_{(l-r) \times (m-r)} \end{bmatrix} \begin{bmatrix} q_1 & & & 0 \\ & \ddots & & \\ & & q_r & \\ 0 & & & I_{m-r} \end{bmatrix}^{-1} S_2 \\ &= S_1 \begin{bmatrix} p_1 & & & 0 \\ & \ddots & & \\ & & p_r & \\ 0 & & & 0_{(l-r) \times (m-r)} \end{bmatrix} \left(S_2^{-1} \begin{bmatrix} q_1 & & & 0 \\ & \ddots & & \\ & & q_r & \\ 0 & & & I_{m-r} \end{bmatrix} \right)^{-1}, \end{aligned}$$

which, by Theorem 4.7.14, is a right coprime polynomial fraction description of G . The last statement follows from Theorem 4.7.5 and Proposition 4.7.15. $\square$

4.8 Facts on Polynomials and Rational Functions

Fact 4.8.1. Let $p \in \mathbb{R}[s]$ be monic, and define $q(s) \triangleq s^n p(1/s)$, where $n \triangleq \deg p$. If $0 \notin \text{roots}(p)$, then $\deg(q) = n$ and

$$\text{mroots}(q) = \{1/\lambda: \lambda \in \text{mroots}(p)\}_{\text{ms}}.$$

If $0 \in \text{roots}(p)$ with multiplicity r , then $\deg(q) = n - r$ and

$$\text{mroots}(q) = \{1/\lambda: \lambda \neq 0 \text{ and } \lambda \in \text{mroots}(p)\}_{\text{ms}}.$$

(Remark: See Fact 11.17.4 and Fact 11.17.5.)

Fact 4.8.2. Let $p \in \mathbb{F}^n[s]$ be given by

$$p(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_1s + \beta_0,$$

let $\beta_n \triangleq 1$, let $\text{mroots}(p) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, and define $\mu_1, \dots, \mu_n$ by

$$\mu_i \triangleq \lambda_1^i + \cdots + \lambda_n^i.$$

Then, for all $k = 1, \dots, n$,

$$k\beta_{n-k} + \mu_1\beta_{n-k+1} + \mu_2\beta_{n-k+2} + \cdots + \mu_k\beta_n = 0.$$

That is,

$$\begin{bmatrix} n & \mu_1 & \mu_2 & \mu_3 & \mu_4 & \cdots & \mu_n \\ 0 & n-1 & \mu_1 & \mu_2 & \mu_3 & \cdots & \mu_{n-1} \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 & 2 & \mu_1 & \mu_2 \\ 0 & 0 & \cdots & 0 & 0 & 1 & \mu_1 \end{bmatrix} \begin{bmatrix} \beta_0 \\ \beta_1 \\ \vdots \\ \beta_{n-1} \\ \beta_n \end{bmatrix} = 0.$$

Consequently, $\beta_1, \dots, \beta_{n-1}$ are uniquely determined by $\mu_1, \dots, \mu_n$. In particular,

$$\begin{aligned} \beta_{n-1} &= -\mu_1, \\ \beta_{n-2} &= \frac{1}{2}(\mu_1^2 - \mu_2), \\ \beta_3 &= \frac{1}{6}(-\mu_1^3 + 3\mu_1\mu_2 - 2\mu_3). \end{aligned}$$

(Proof: See [709, p. 44] and [1002, p. 9].) (Remark: These equations are a consequence of Newton's identities given by Fact 1.15.11. Note that, for $i = 0, \dots, n$, it follows that $\beta_i = (-1)^{n-i}E_{n-i}$, where E_i is the i th elementary symmetric polynomial of the roots of p .)

Fact 4.8.3. Let $p, q \in \mathbb{F}[s]$ be monic. Then, p and q are coprime if and only if their least common multiple is pq .

Fact 4.8.4. Let $p, q \in \mathbb{F}[s]$, where $p(s) = a_ns^n + \cdots + a_1s + a_0$, $q(s) = b_ms^m + \cdots + b_1s + b_0$, $\deg p = n$, and $\deg q = m$. Furthermore, define the Toeplitz matrices $[p]^{(m)} \in \mathbb{F}^{m \times (n+m)}$ and $[q]^{(n)} \in \mathbb{F}^{n \times (n+m)}$ by

$$[p]^{(m)} \triangleq \begin{bmatrix} a_n & a_{n-1} & \cdots & a_1 & a_0 & 0 & 0 & \cdots & 0 \\ 0 & a_n & a_{n-1} & \cdots & a_1 & a_0 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \cdots & \ddots & \ddots & \ddots & \vdots \end{bmatrix}$$

and

$$[q]^{(n)} \triangleq \begin{bmatrix} b_m & b_{m-1} & \cdots & b_1 & b_0 & 0 & 0 & \cdots & 0 \\ 0 & b_m & b_{m-1} & \cdots & b_1 & b_0 & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \cdots & \ddots & \ddots & \ddots & \vdots \end{bmatrix}.$$

Then, p and q are coprime if and only if

$$\det \begin{bmatrix} [p]^{(m)} \\ [q]^{(n)} \end{bmatrix} \neq 0.$$

(Proof: See [481, p. 162] or [1098, pp. 187–191].) (Remark: $\begin{bmatrix} A \\ B \end{bmatrix}$ is the *Sylvester matrix*, and $\det \begin{bmatrix} A \\ B \end{bmatrix}$ is the *resultant* of p and q .) (Remark: The form $\begin{bmatrix} [p]^{(m)} \\ [q]^{(n)} \end{bmatrix}$ appears in [1098, pp. 187–191]. The result is given in [481, p. 162] in terms of $\begin{bmatrix} \hat{I}[p]^{(m)} \\ \hat{I}[q]^{(n)} \end{bmatrix} \hat{I}$ and in [1503, p. 85] in terms of $\begin{bmatrix} [p]^{(m)} \\ \hat{I}[q]^{(n)} \end{bmatrix}$. Interweaving the rows of $[p]^{(m)}$ and $[q]^{(n)}$ and taking the transpose yields a *step-down matrix* [389].)

Fact 4.8.5. Let $p_1, \dots, p_n \in \mathbb{F}[s]$, and let $d \in \mathbb{F}[s]$ be the greatest common divisor of $p_1, \dots, p_n$. Then, there exist polynomials $q_1, \dots, q_n \in \mathbb{F}[s]$ such that

$$d = \sum_{i=1}^n q_i p_i.$$

In addition, $p_1, \dots, p_n$ are coprime if and only if there exist polynomials $q_1, \dots, q_n \in \mathbb{F}[s]$ such that

$$1 = \sum_{i=1}^n q_i p_i.$$

(Proof: See [508, p. 16].) (Remark: The polynomial d is given by the *Bezout equation*.)

Fact 4.8.6. Let $p, q \in \mathbb{F}[s]$, where $p(s) = a_n s^n + \dots + a_1 s + a_0$ and $q(s) = b_n s^n + \dots + b_1 s + b_0$, and define $[p]^{(n)}, [q]^{(n)} \in \mathbb{F}^{n \times 2n}$ as in Fact 4.8.4. Furthermore, define

$$R(p, q) \triangleq \begin{bmatrix} [p]^{(n)} \\ [q]^{(n)} \end{bmatrix} = \begin{bmatrix} A_1 & A_2 \\ B_1 & B_2 \end{bmatrix},$$

where $A_1, A_2, B_1, B_2 \in \mathbb{F}^{n \times n}$, and define $\hat{p}(s) \triangleq s^n p(-s)$ and $\hat{q}(s) \triangleq s^n q(-s)$. Then,

$$\begin{bmatrix} A_1 & A_2 \\ B_1 & B_2 \end{bmatrix} = \begin{bmatrix} \hat{p}(N_n^T) & p(N_n) \\ \hat{q}(N_n^T) & q(N_n) \end{bmatrix},$$

$$A_1 B_1 = B_1 A_1,$$

$$A_2 B_2 = B_2 A_2,$$

$$A_1 B_2 + A_2 B_1 = B_1 A_2 + B_2 A_1.$$

Therefore,

$$\begin{bmatrix} I & 0 \\ -B_1 & A_1 \end{bmatrix} \begin{bmatrix} A_1 & A_2 \\ B_1 & B_2 \end{bmatrix} = \begin{bmatrix} A_1 & A_2 \\ 0 & A_1 B_2 - B_1 A_2 \end{bmatrix},$$

$$\begin{bmatrix} -B_2 & A_2 \\ 0 & I \end{bmatrix} \begin{bmatrix} A_1 & A_2 \\ B_1 & B_2 \end{bmatrix} = \begin{bmatrix} A_2 B_1 - B_2 A_1 & 0 \\ B_1 & B_2 \end{bmatrix},$$

and

$$\det R(p, q) = \det(A_1 B_2 - B_1 A_2) = \det(B_2 A_1 - A_2 B_1).$$

Now, define $B(p, q) \in \mathbb{F}^{n \times n}$ by

$$B(p, q) \triangleq (A_1 B_2 - B_1 A_2) \hat{I}.$$

Then, the following statements hold:

i) For all $s, \hat{s} \in \mathbb{C}$,

$$p(s)q(\hat{s}) - q(s)p(\hat{s}) = (s - \hat{s}) \begin{bmatrix} 1 \\ s \\ \vdots \\ s^{n-1} \end{bmatrix}^T B(p, q) \begin{bmatrix} 1 \\ \hat{s} \\ \vdots \\ \hat{s}^{n-1} \end{bmatrix}.$$

ii) $B(p, q) = (B_2 A_1 - A_2 B_1) \hat{I} = \hat{I} (A_1^T B_2^T - B_1^T A_2^T) = \hat{I} (B_1^T A_2^T - A_1^T B_2^T)$.

iii) $\begin{bmatrix} 0 & B(p, q) \\ -B(p, q) & 0 \end{bmatrix} = QR^T(p, q)QR(p, q)Q$, where $Q \triangleq \begin{bmatrix} 0 & \hat{I} \\ -\hat{I} & 0 \end{bmatrix}$.

iv) $|\det B(p, q)| = |\det R(p, q)| = |\det q[C(p)]|$.

v) $B(p, q)$ and $\hat{B}(p, q)$ are symmetric.

vi) $B(p, q)$ is a linear function of (p, q) .

vii) $B(p, q) = -B(q, p)$.

Now, assume that $\deg q \leq \deg p = n$ and p is monic. Then, the following statements hold:

viii) $\deg B(p, q)$ is equal to the degree of the greatest common divisor of p and q .

ix) p and q are coprime if and only if $B(p, q)$ is nonsingular.

x) If $B(p, q)$ is nonsingular, then $[B(p, q)]^{-1}$ is Hankel. In fact,

$$[B(p, q)]^{-1} = H(a/p),$$

where $a, b \in \mathbb{F}[s]$ satisfy the Bezout equation $aq + bp = 1$.

xi) If $q = q_1 q_2$, where $q_1, q_2 \in \mathbb{F}[s]$, then

$$B(p, q) = B(p, q_1)q_2[C(p)] = q_1[C^T(p)]B(p, q_2).$$

xii) $B(p, q) = B(p, q)C(p) = C^T(p)B(p, q)$.

xiii) $B(p, q) = B(p, 1)q[C(p)] = q[C^T(p)]B(p, 1)$, where $B(p, 1)$ is the Hankel matrix

$$B(p, 1) = \begin{bmatrix} a_1 & a_2 & \cdots & a_{n-1} & 1 \\ a_2 & a_3 & \ddots & 1 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ a_{n-1} & 1 & \ddots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \end{bmatrix}.$$

In particular, for $n = 3$ and $q(s) = s$, it follows that

$$\begin{bmatrix} -a_0 & 0 & 0 \\ 0 & a_2 & 1 \\ 0 & 1 & 0 \end{bmatrix} = \begin{bmatrix} a_1 & a_2 & 1 \\ a_2 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -a_0 & -a_1 & -a_2 \end{bmatrix}.$$

xiv) If A_2 is nonsingular, then

$$\begin{bmatrix} A_1 & A_2 \\ B_1 & B_2 \end{bmatrix} = \begin{bmatrix} 0 & I \\ A_2^{-1}\hat{I} & B_2A_2^{-1} \end{bmatrix} \begin{bmatrix} B(p, q) & 0 \\ 0 & I \end{bmatrix} \begin{bmatrix} I & 0 \\ A_1 & A_2 \end{bmatrix}.$$

xv) If p has distinct roots $\lambda_1, \dots, \lambda_n$, then

$$V^T(\lambda_1, \dots, \lambda_n)B(p, q)V(\lambda_1, \dots, \lambda_n) = \text{diag}[q(\lambda_1)p'(\lambda_1), \dots, q(\lambda_n)p'(\lambda_n)].$$

(Proof: See [481, pp. 164–167], [508, pp. 200–207], and [663]. To prove *ii)*, note that A_1, A_2, B_1, B_2 are square and Toeplitz, and thus reverse symmetric, that is, $A_1 = A_1^T$. See Fact 3.18.5.) (Remark: $B(p, q)$ is the *Bezout matrix* of p and q . See [145, 662, 722, 1356, 1444], [1098, p. 189], and Fact 5.15.24.) (Remark: *xiii)* is the *Barnett factorization*. See [138, 1356]. The definitions of $B(p, q)$ and *ii)* are the *Gohberg-Semencul formulas*. See [508, p. 206].) (Remark: It follows from continuity that the expressions for $\det R(p, q)$ are valid whether or not A_1 or B_2 is singular. See Fact 2.14.13.) (Remark: The inverse of a Hankel matrix is a Bezout matrix. See [481, p. 174].)

Fact 4.8.7. Let $p, q \in \mathbb{F}[s]$, where $p(s) = \alpha_1 s + \alpha_0$ and $q(s) = s^2 + \beta_1 s + \beta_0$. Then, p and q are coprime if and only if $\alpha_0^2 + \alpha_1^2 \beta_0 \neq \alpha_0 \alpha_1 \beta_1$. (Proof: Use Fact 4.8.6.)

Fact 4.8.8. Let $p, q \in \mathbb{F}[s]$, assume that q is monic, assume that $\deg p < \deg q = n$, and define $B(p, q)$ as in Fact 4.8.6. Furthermore, define $g \in \mathbb{F}(s)$ by

$$g(s) \triangleq \frac{p(s)}{q(s)} = \sum_{i=1}^{\infty} \frac{h_i}{s^i}.$$

Finally, define the Hankel matrix $H_{i,j}(g) \in \mathbb{R}^{i \times j}$ by

$$H_{i,j}(g) = \begin{bmatrix} h_1 & h_2 & h_{k+3} & \cdots & h_j \\ h_{k+2} & h_{k+3} & \ddots & \ddots & \vdots \\ h_{k+3} & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ h_i & \cdots & \cdots & \cdots & h_{j+i-1} \end{bmatrix}.$$

Then, the following statements are equivalent:

- i)* p and q are coprime.
- ii)* $H_{n,n}(g)$ is nonsingular.
- iii)* For all $i, j \geq n$, $\text{rank } H_{i,j}(g) = n$.
- iv)* There exist $i, j \geq n$ such that $\text{rank } H_{i,j}(g) = n$.

Furthermore, the following statements hold:

- v) If p and q are coprime, then $[H_{n,n}(g)]^{-1} = B(q, a)$, where $a, b \in \mathbb{F}[s]$ satisfy the Bezout equation $ap + bq = 1$.
- vi) $B(q, p) = B(q, 1)H_{n,n}(g)B(q, 1)$.
- vii) $B(q, p)$ and $H_{n,n}(g)$ are congruent.
- viii) $\text{In } B(q, p) = \text{In } H_{n,n}(g)$.
- ix) $\det H_{n,n}(g) = \det B(q, p)$.

(Proof: See [508, pp. 215–221].) (Remark: See Proposition 12.9.11.)

Fact 4.8.9. Let $q \in \mathbb{R}[s]$, define $g \in \mathbb{F}(s)$ by $g \triangleq q'/q$, and define $B(q, q')$ as in Fact 4.8.6. Then, the following statements hold:

- i) The number of distinct roots of q is $\text{rank } B(q, q')$.
- ii) q has n distinct roots if and only if $B(q, q')$ is nonsingular.
- iii) The number of distinct real roots of q is $\text{sig } B(q, q')$.
- iv) q has n distinct, real roots if and only if $B(q, q')$ is positive definite.
- v) The number of distinct complex roots of q is $2\nu_-[B(q, q')]$.
- vi) q has n distinct, complex roots if and only if n is even and $\nu_-[B(q, q')] = n/2$.
- vii) q has n real roots if and only if $B(q, q')$ is positive semidefinite.

(Proof: See [508, p. 252].) (Remark: $q'(s) \triangleq (d/ds)q(s)$.)

Fact 4.8.10. Let $q \in \mathbb{F}[s]$, where $q(s) = \sum_{i=0}^n b_i s^i$, and define

$$\text{coeff}(q) \triangleq \begin{bmatrix} b_n \\ \vdots \\ b_0 \end{bmatrix}.$$

Now, let $p \in \mathbb{F}[s]$, where $p(s) = \sum_{i=0}^n a_i s^i$. Then,

$$\text{coeff}(pq) = A \text{coeff}(q),$$

where $A \in \mathbb{F}^{2n \times (n+1)}$ is the Toeplitz matrix

$$A = \begin{bmatrix} a_n & 0 & 0 & \cdots & 0 \\ a_{n-1} & a_n & 0 & \cdots & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ a_0 & a_1 & \ddots & \ddots & a_n \\ 0 & a_0 & \ddots & \ddots & a_{n-1} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & a_0 & a_1 \end{bmatrix}.$$

In particular, if $n = 3$, then

$$A = \begin{bmatrix} a_2 & 0 & 0 \\ a_1 & a_2 & 0 \\ a_0 & a_1 & a_2 \\ 0 & a_0 & a_1 \end{bmatrix}.$$

Fact 4.8.11. Let $\lambda_1, \dots, \lambda_n \in \mathbb{C}$ be distinct and, for all $i = 1, \dots, n$, define

$$p_i(s) \triangleq \prod_{\substack{j=1 \\ j \neq i}}^n \frac{s - \lambda_i}{\lambda_i - \lambda_j}.$$

Then, for all $i = 1, \dots, n$,

$$p_i(\lambda_j) = \begin{cases} 1, & i = j, \\ 0, & i \neq j. \end{cases}$$

(Remark: This identity is the *Lagrange interpolation formula*.)

Fact 4.8.12. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\det(I + A) \neq 0$. Then, there exists $p \in \mathbb{F}[s]$ such that $\deg p \leq n - 1$ and $(I + A)^{-1} = p(A)$. (Remark: See Fact 4.8.12.)

Fact 4.8.13. Let $A \in \mathbb{F}^{n \times n}$, let $q \in \mathbb{F}[s]$, and assume that $q(A)$ is nonsingular. Then, there exists $p \in \mathbb{F}[s]$ such that $\deg p \leq n - 1$ and $[q(A)]^{-1} = p(A)$. (Proof: See Fact 5.14.24.)

Fact 4.8.14. Let $A \in \mathbb{R}^{n \times n}$, assume that A is skew symmetric, and let the components of $x_A \in \mathbb{R}^{n(n-1)/2}$ be the entries $A_{(i,j)}$ for all $i > j$. Then, there exists a polynomial function $p: \mathbb{R}^{n(n-1)/2} \mapsto \mathbb{R}$ such that, for all $\alpha \in \mathbb{R}$ and $x \in \mathbb{R}^{n(n-1)/2}$,

$$p(\alpha x) = \alpha^{n/2} p(x)$$

and

$$\det A = p^2(x_A).$$

In particular,

$$\det \begin{bmatrix} 0 & a \\ -a & 0 \end{bmatrix} = a^2$$

and

$$\det \begin{bmatrix} 0 & a & b & c \\ -a & 0 & d & e \\ -b & -d & 0 & f \\ -c & -e & -f & 0 \end{bmatrix} = (af - be + cd)^2.$$

(Proof: See [878, p. 224] and [1098, pp. 125–127].) (Remark: The polynomial p is the *Pfaffian*, and this result is *Pfaff's theorem*.) (Remark: An extension to the product of a pair of skew-symmetric matrices is given in [436].) (Remark: See Fact 3.7.33.)

Fact 4.8.15. Let $G \in \mathbb{F}^{n \times m}(s)$, and let $G_{(i,j)} = n_{ij}/d_{ij}$, where $n_{ij} \in \mathbb{F}[s]$ and $d_{ij} \in \mathbb{F}[s]$ are coprime for all $i = 1, \dots, n$ and $j = 1, \dots, m$. Then, q_1 given by the Smith-McMillan form is the least common multiple of $d_{11}, d_{12}, \dots, d_{nm}$.

Fact 4.8.16. Let $G \in \mathbb{F}^{n \times m}(s)$, assume that $\text{rank } G = m$, and let $\lambda \in \mathbb{C}$, where λ is not a pole of G . Then, λ is a transmission zero of G if and only if there exists a vector $u \in \mathbb{C}^m$ such that $G(\lambda)u = 0$. Furthermore, if G is square, then λ is a transmission zero of G if and only if $\det G(\lambda) = 0$.

Fact 4.8.17. Let $G \in \mathbb{F}^{n \times m}(s)$, let $\omega \in \mathbb{R}$, and assume that $j\omega$ is not a pole of G . Then,

$$\text{Im } G(-j\omega) = -\text{Im } G(j\omega).$$

4.9 Facts on the Characteristic and Minimal Polynomials

Fact 4.9.1. Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbb{R}^{2 \times 2}$. Then, the following identities hold:

- i) $\text{mspec}(A) = \left\{ \frac{1}{2} \left[a + d \pm \sqrt{(a-d)^2 + 4bc} \right] \right\}_{\text{ms}}$
 $= \left\{ \frac{1}{2} \left[\text{tr } A \pm \sqrt{(\text{tr } A)^2 - 4 \det A} \right] \right\}_{\text{ms}}.$
- ii) $\chi_A(s) = s^2 - (\text{tr } A)s + \det A$.
- iii) $\det A = \frac{1}{2}[(\text{tr } A)^2 - \text{tr } A^2]$.
- iv) $(sI - A)^A = sI + A - (\text{tr } A)I$.
- v) $A^{-1} = (\det A)^{-1}[(\text{tr } A)I - A]$.
- vi) $A^A = (\text{tr } A)I - A$.
- vii) $\text{tr } A^{-1} = \text{tr } A / \det A$.

Fact 4.9.2. Let $A \in \mathbb{R}^{3 \times 3}$. Then, the following identities hold:

- i) $\chi_A(s) = s^3 - (\text{tr } A)s^2 + (\text{tr } A^A)s - \det A$.
- ii) $\text{tr } A^A = \frac{1}{2}[(\text{tr } A)^2 - \text{tr } A^2]$.
- iii) $\det A = \frac{1}{3}\text{tr } A^3 - \frac{1}{2}(\text{tr } A)\text{tr } A^2 + \frac{1}{6}(\text{tr } A)^3$.
- iv) $(sI - A)^A = s^2I + s[A - (\text{tr } A)I] + A^2 - (\text{tr } A)A + \frac{1}{2}[(\text{tr } A)^2 - \text{tr } A^2]I$.

(Remark: See Fact 7.5.17.)

Fact 4.9.3. Let $A, B \in \mathbb{F}^{2 \times 2}$. Then,

$$AB + BA - (\text{tr } A)B - (\text{tr } B)A + [(\text{tr } A)(\text{tr } B) - \text{tr } AB]I = 0.$$

Furthermore,

$$\det(A + B) - \det A - \det B = (\text{tr } A)(\text{tr } B) - \text{tr } AB.$$

(Proof: Apply the Cayley-Hamilton theorem to $A + xB$, differentiate with respect to x , and set $x = 0$. For the second identity, evaluate the Cayley-Hamilton theorem with $A + B$. See [499, 500, 890, 1128] or [1186, p. 37].) (Remark: This identity is a *polarized Cayley-Hamilton theorem*. See [78].)

Fact 4.9.4. Let $A, B, C \in \mathbb{F}^{2 \times 2}$. Then,

$$\begin{aligned} 2ABC &= (\operatorname{tr} A)BC + (\operatorname{tr} B)AC + (\operatorname{tr} C)AB \\ &\quad - (\operatorname{tr} AC)B + [(\operatorname{tr} AB) - (\operatorname{tr} A)(\operatorname{tr} B)]C \\ &\quad + [(\operatorname{tr} BC) - (\operatorname{tr} B)(\operatorname{tr} C)]A \\ &\quad - [(\operatorname{tr} ACB) - (\operatorname{tr} AC)(\operatorname{tr} B)]I. \end{aligned}$$

(Remark: This identity is a *polarized Cayley-Hamilton theorem*. See [78].) (Remark: An analogous formula exists for the product of six 3×3 matrices. See [78].)

Fact 4.9.5. Let $A, B, C \in \mathbb{F}^{3 \times 3}$, and assume that $\operatorname{tr} A = \operatorname{tr} B = \operatorname{tr} C = 0$. Then,

$$4 \operatorname{tr}(A^2 B^2) + 2 \operatorname{tr}[(AB)^2] = \operatorname{tr}(A^2) \operatorname{tr}(B^2) + 2[\operatorname{tr}(AB)]^2$$

and

$$\begin{aligned} &6 \operatorname{tr}(A^2 B^2 AB) + 6 \operatorname{tr}(B^2 A^2 BA) + 2 \operatorname{tr}(AB) \operatorname{tr}[(AB)^2] + 2 \operatorname{tr}(A^3) \operatorname{tr}(B^3) \\ &= 2 \operatorname{tr}(AB) \operatorname{tr}(A^2 B^2) + \operatorname{tr}(A^2) \operatorname{tr}(AB) \operatorname{tr}(B^2) + 2[\operatorname{tr}(AB)]^3 + 6 \operatorname{tr}(A^2 B) \operatorname{tr}(AB^2). \end{aligned}$$

(Proof: See [81].)

Fact 4.9.6. Let $A, B, C \in \mathbb{F}^{3 \times 3}$. Then,

$$\begin{aligned} &\sum [A'B'C' - (\operatorname{tr} A')B'C' + (\operatorname{tr} A')(\operatorname{tr} B')C' - (\operatorname{tr} A'B')C'] \\ &\quad - [(\operatorname{tr} A)(\operatorname{tr} B)\operatorname{tr} C - (\operatorname{tr} A)\operatorname{tr} BC - (\operatorname{tr} B)\operatorname{tr} CA - (\operatorname{tr} C)\operatorname{tr} AB + \operatorname{tr} ABC \\ &\quad + \operatorname{tr} CBA]I = 0, \end{aligned}$$

where the sum is taken over all six permutations A', B', C' of A, B, C . (Remark: This identity is a *polarized Cayley-Hamilton theorem*. See [79, 890, 1128].)

Fact 4.9.7. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B commute, and define $f: \mathbb{C}^2 \mapsto \mathbb{C}$ by $f(r, s) \triangleq \det(rA - sB)$. Then, $f(B, A) = 0$. (Remark: This result is the *generalized Cayley-Hamilton theorem*. See [356, 682].)

Fact 4.9.8. Let $A \in \mathbb{F}^{n \times n}$, let $\chi_A(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_0$, and let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$A^A = (-1)^{n-1}(A^{n-1} + \beta_{n-1}A^{n-2} + \cdots + \beta_1 I).$$

Furthermore,

$$\operatorname{tr} A^A = (-1)^{n-1} \chi'_A(0) = (-1)^{n-1} \beta_1 = \sum_{1 \leq j_1 < \cdots < j_{n-1} \leq n} \lambda_{j_1} \cdots \lambda_{j_{n-1}} = \sum_{i=1}^n \det A_{[i; i]}.$$

(Proof: Use $A^{-1} \chi_A(A) = 0$. The second identity follows from (4.4.19) or Lemma 4.4.8.) (Remark: See Fact 4.10.7.)

Fact 4.9.9. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $\chi_A(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_0$. Then,

$$\begin{aligned}\chi_{A^{-1}}(s) &= \frac{1}{\det A}(-s)^n \chi_A(1/s) \\ &= s^n + (\beta_1/\beta_0)s^{n-1} + \cdots + (\beta_{n-1}/\beta_0)s + 1/\beta_0.\end{aligned}$$

(Remark: See Fact 5.16.2.)

Fact 4.9.10. Let $A \in \mathbb{F}^{n \times n}$, and assume that either A and $-A$ are similar or A^T and $-A$ are similar. Then,

$$\chi_A(s) = (-1)^n \chi_A(-s).$$

Furthermore, if n is even, then χ_A is even, whereas, if n is odd, then χ_A is odd. (Remark: A and A^T are similar. See Corollary 4.3.11 and Corollary 5.3.8.)

Fact 4.9.11. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $s \in \mathbb{C}$,

$$(sI - A)^A = \chi_A(s)(sI - A)^{-1} = \sum_{i=0}^{n-1} \chi_A^{[i]}(s)A^i,$$

where

$$\chi_A(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_1s + \beta_0$$

and, for all $i = 0, \dots, n-1$, the polynomial $\chi_A^{[i]}$ is defined by

$$\chi_A^{[i]}(s) \triangleq s^{n-i} + \beta_{n-1}s^{n-1-i} + \cdots + \beta_{i+1}.$$

Note that

$$\chi_A^{[n-1]}(s) = s + \beta_{n-1}, \quad \chi_A^{[n]}(s) = 1,$$

and that, for all $i = 0, \dots, n-1$ and with $\chi_A^{[0]} \triangleq \chi_A$, the polynomials $\chi_A^{[i]}$ satisfy the recursion

$$s\chi_A^{[i+1]}(s) = \chi_A^{[i]}(s) - \beta_i.$$

(Proof: See [1455, p. 31].)

Fact 4.9.12. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is skew symmetric. If n is even, then χ_A is even, whereas, if n is odd, then χ_A is odd.

Fact 4.9.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is skew Hermitian. Then, for all $s \in \mathbb{C}$,

$$\chi_A(-s) = (-1)^n \overline{\chi_A(\overline{s})}.$$

Fact 4.9.14. Let $A \in \mathbb{F}^{n \times n}$. Then, χ_A is even for the matrices $\mathcal{A} \in \mathbb{F}^{2n \times 2n}$ given by $\begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix}$, $\begin{bmatrix} A & 0 \\ 0 & -A \end{bmatrix}$, and $\begin{bmatrix} A & 0 \\ 0 & -A^* \end{bmatrix}$.

Fact 4.9.15. Let $A, B \in \mathbb{F}^{n \times n}$, and define $\mathcal{A} \triangleq \begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix}$. Then,

$$\chi_{\mathcal{A}}(s) = \chi_{AB}(s^2) = \chi_{BA}(s^2).$$

Consequently, $\chi_{\mathcal{A}}$ is even. (Proof: Use Fact 2.14.13 and Proposition 4.4.10.)

Fact 4.9.16. Let $x, y, z, w \in \mathbb{F}^n$, and define $A \triangleq xy^T$ and $B \triangleq xy^T + zw^T$. Then,

$$\chi_A(s) = s^{n-1}(s - x^T y)$$

and

$$\chi_B(s) = s^{n-2} [s^2 - (x^T y + z^T w)s + x^T y z^T w - y^T z x^T w].$$

(Remark: See Fact 5.11.13.)

Fact 4.9.17. Let $x, y \in \mathbb{F}^{n-1}$, and define $A \in \mathbb{F}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} 0 & x^T \\ y & 0 \end{bmatrix}.$$

Then,

$$\chi_A(s) = s^{n-1}(s^2 - y^T x).$$

(Proof: See [1333].)

Fact 4.9.18. Let $x, y, z, w \in \mathbb{F}^{n-1}$, and define $A \in \mathbb{F}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} 1 & x^T \\ y & zw^T \end{bmatrix}.$$

Then,

$$\chi_A(s) = s^{n-3} [s^3 - (1 + w^T z)s^2 + (w^T z - x^T y)s + w^T z x^T y - x^T z w^T y].$$

(Proof: See [409].) (Remark: Extensions are given in [1333].)

Fact 4.9.19. Let $x \in \mathbb{R}^3$, and define $\theta \triangleq \sqrt{x^T x}$. Then,

$$\chi_{K(x)}(s) = s^3 + \theta^2 s.$$

Hence,

$$\text{mspec}[K(x)] = \{0, j\theta, -j\theta\}_{\text{ms}}.$$

Now, assume that $x \neq 0$. Then, x is an eigenvector corresponding to the eigenvalue 0, that is, $K(x)x = 0$. Furthermore, if either $x_{(1)} \neq 0$ or $x_{(2)} \neq 0$, then

$$\begin{bmatrix} x_{(1)}x_{(3)} + j\theta x_{(2)} \\ x_{(2)}x_{(3)} - j\theta x_{(1)} \\ -x_{(1)}^2 - x_{(2)}^2 \end{bmatrix}$$

is an eigenvector corresponding to the eigenvalue $j\theta$. Finally, if $x_{(1)} = x_{(2)} = 0$, then $\begin{bmatrix} j \\ 1 \\ 0 \end{bmatrix}$ is an eigenvector corresponding to the eigenvalue $j\theta$. (Remark: See Fact 11.11.6.)

Fact 4.9.20. Let $a, b \in \mathbb{R}^3$, where $a = [a_1 \ a_2 \ a_3]^T$ and $b = [b_1 \ b_2 \ b_3]^T$, and define the skew-symmetric matrix $A \in \mathbb{R}^{4 \times 4}$ by

$$A \triangleq \begin{bmatrix} K(a) & b \\ -b^T & 0 \end{bmatrix}.$$

Then, the following statements hold:

$$i) \det A = (a^T b)^2.$$

$$ii) \chi_A(s) = s^4 + (a^T a + b^T b)s^2 + (a^T b)^2.$$

$$\text{iii) } A^A = -a^T b \begin{bmatrix} K(b) & a \\ -a^T & 0 \end{bmatrix}.$$

$$\text{iv) If } \det A \neq 0, \text{ then } A^{-1} = -(a^T b)^{-1} \begin{bmatrix} K(b) & a \\ -a^T & 0 \end{bmatrix}.$$

v) If $\det A = 0$, then

$$A^3 = -(a^T a + b^T b)^2 A$$

and

$$A^+ = -(a^T a + b^T b)^{-2} A.$$

(Proof: See [1334].) (Remark: See Fact 4.10.2 and Fact 11.11.17.)

Fact 4.9.21. Let $A \in \mathbb{R}^{2n \times 2n}$, and assume that A is Hamiltonian. Then, χ_A is even, and thus $\text{mspec}(A) = -\text{mspec}(A)$. (Remark: See Fact 5.9.24.)

Fact 4.9.22. Let $A, B, C \in \mathbb{R}^{n \times n}$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ C & -A^T \end{bmatrix}.$$

If B and C are symmetric, then $\mathcal{A}$ is Hamiltonian. If B and C are skew symmetric, then $\chi_{\mathcal{A}}$ is even, although $\mathcal{A}$ is not necessarily Hamiltonian. (Proof: For the second result replace J_{2n} by $\begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix}$.)

Fact 4.9.23. Let $A \in \mathbb{R}^{n \times n}$, $R \in \mathbb{R}^{n \times n}$, and $B \in \mathbb{R}^{n \times m}$, and define $\mathcal{A} \in \mathbb{R}^{2n \times 2n}$ by

$$\mathcal{A} \triangleq \begin{bmatrix} A & BB^T \\ R & -A^T \end{bmatrix}.$$

Then, for all $s \notin \text{spec}(A)$,

$$\chi_{\mathcal{A}}(s) = (-1)^n \chi_A(s) \chi_A(-s) \det[I + B^T(-sI - A^T)^{-1} R(sI - A)^{-1} B].$$

Now, assume that R is symmetric. Then, $\mathcal{A}$ is Hamiltonian, and $\chi_{\mathcal{A}}$ is even. If, in addition, R is positive semidefinite, then $(-1)^n \chi_{\mathcal{A}}$ has a spectral factorization. (Proof: Using (2.8.10) and (2.8.14), it follows that, for all $\pm s \notin \text{spec}(A)$,

$$\begin{aligned} \chi_{\mathcal{A}}(s) &= \det(sI - A) \det[sI + A^T - R(sI - A)^{-1} BB^T] \\ &= (-1)^n \chi_A(s) \chi_A(-s) \det[I - B^T(sI + A^T)^{-1} R(sI - A)^{-1} B]. \end{aligned}$$

To prove the second statement, note that, for all $\omega \in \mathbb{R}$ such that $j\omega \notin \text{spec}(A)$, it follows that

$$\chi_{\mathcal{A}}(j\omega) = (-1)^n \chi_A(j\omega) \overline{\chi_A(j\omega)} \det[I + B^T(j\omega I - A)^{-*} R(j\omega I - A)^{-1} B].$$

Thus, $(-1)^n \chi_{\mathcal{A}}(j\omega) \geq 0$. By continuity, $(-1)^n \chi_{\mathcal{A}}(j\omega) \geq 0$ for all $\omega \in \mathbb{R}$. Now, Proposition 4.1.1 implies that $(-1)^n \chi_{\mathcal{A}}$ has a spectral factorization. (Remark: Not all Hamiltonian matrices $\mathcal{A} \in \mathbb{R}^{2n \times 2n}$ have the property that $(-1)^n \chi_{\mathcal{A}}$ has a spectral factorization. Consider $\begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -1 & 0 & 0 & 0 \\ 0 & -3 & 0 & 0 \end{bmatrix}$, whose spectrum is $\{j, -j, \sqrt{3}j, -\sqrt{3}j\}$.)

(Remark: This result is closely related to Proposition 12.17.8.) (Remark: See Fact 3.19.6.)

Fact 4.9.24. Let $A \in \mathbb{F}^{n \times n}$. Then, $\mu_A = \chi_A$ if and only if there exists a unique monic polynomial $p \in \mathbb{F}[s]$ of degree n and such that $p(A) = 0$. (Proof: To prove necessity, note that if $\hat{p} \neq p$ is monic, of degree n , and satisfies $\hat{p}(A) = 0$, then $p - \hat{p}$ is nonzero, has degree less than n , and satisfies $(p - \hat{p})(A) = 0$. Conversely, if $\mu_A \neq \chi_A$, then $\mu_A + \chi_A$ is monic, has degree n , and satisfies $(\mu_A + \chi_A)(A) = 0$.)

4.10 Facts on the Spectrum

Fact 4.10.1. Let $A \in \mathbb{F}^{3 \times 3}$, assume that A is symmetric, let $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ denote the eigenvalues of A , where $\lambda_1 \geq \lambda_2 \geq \lambda_3$, and define

$$p = \frac{1}{6} \operatorname{tr} [A - \frac{1}{3}(\operatorname{tr} A)I]^2$$

and

$$q = \frac{1}{2} \det [A - \frac{1}{3}(\operatorname{tr} A)I].$$

Then, the following statements hold:

- i) $0 \leq |q| \leq p^{3/2}$.
- ii) $p = 0$ if and only if $\lambda_1 = \lambda_2 = \lambda_3 = \frac{1}{3} \operatorname{tr} A$.
- iii) $p > 0$ if and only if

$$\begin{aligned}\lambda_1 &= \frac{1}{3} \operatorname{tr} A + 2\sqrt{p} \cos \phi, \\ \lambda_2 &= \frac{1}{3} \operatorname{tr} A + \sqrt{3p} \sin \phi - \sqrt{p} \cos \phi, \\ \lambda_3 &= \frac{1}{3} \operatorname{tr} A - \sqrt{3p} \sin \phi - \sqrt{p} \cos \phi,\end{aligned}$$

where $\phi \in [0, \pi/3]$ is given by

$$\phi = \frac{1}{3} \cos^{-1} \frac{q}{p^{3/2}}.$$

- iv) $\phi = 0$ if and only if $q = p^{3/2} > 0$. In this case,

$$\begin{aligned}\lambda_1 &= \frac{1}{3} \operatorname{tr} A + 2\sqrt{p}, \\ \lambda_2 &= \lambda_3 = \frac{1}{3} \operatorname{tr} A - \sqrt{p}.\end{aligned}$$

- v) $\phi = \pi/6$ if and only if $p > 0$ and $q = 0$. In this case, $\sin \phi = 1/2$, $\cos \phi = \sqrt{3}/2$, and

$$\begin{aligned}\lambda_1 &= \frac{1}{3} \operatorname{tr} A + \sqrt{3p}, \\ \lambda_2 &= \frac{1}{3} \operatorname{tr} A, \\ \lambda_3 &= \frac{1}{3} \operatorname{tr} A - \sqrt{3p}.\end{aligned}$$

- vi) $\phi = \pi/3$ if and only if $q = -p^{3/2} < 0$. In this case, $\sin \phi = \sqrt{3}/2$, $\cos \phi = 1/2$, and

$$\begin{aligned}\lambda_1 &= \lambda_2 = \frac{1}{3} \operatorname{tr} A + \sqrt{p}, \\ \lambda_3 &= \frac{1}{3} \operatorname{tr} A - 2\sqrt{p}.\end{aligned}$$

(Proof: See [1203].) (Remark: This result is based on *Cardano's trigonometric solution* for the roots of a cubic polynomial. See [234, 1203].) (Remark: The inequality $q^2 \leq p^3$ follows from Fact 1.10.13.)

Fact 4.10.2. Let $a, b, c, d, \omega \in \mathbb{R}$, and define the skew-symmetric matrix $A \in \mathbb{R}^{4 \times 4}$ given by

$$A \triangleq \begin{bmatrix} 0 & \omega & a & b \\ -\omega & 0 & c & d \\ -a & -c & 0 & \omega \\ -b & -d & -\omega & 0 \end{bmatrix}.$$

Then,

$$\chi_A(s) = s^4 + (2\omega^2 + a^2 + b^2 + c^2 + d^2)s^2 + [\omega^2 - (ad - bc)]^2$$

and

$$\det A = [\omega^2 - (ad - bc)]^2.$$

Hence, A is singular if and only if $bc \leq ad$ and $\omega = \sqrt{ad - bc}$. Furthermore, A has a repeated eigenvalue if and only if either *i*) A is singular or *ii*) $a = -d$ and $b = c$. In case *i*), A has the repeated eigenvalue 0, while, in case *ii*), A has the repeated eigenvalues $j\sqrt{\omega^2 + a^2 + b^2}$ and $-j\sqrt{\omega^2 + a^2 + b^2}$. Finally, cases *i*) and *ii*) cannot occur simultaneously. (Remark: See Fact 4.9.20, Fact 3.7.33, Fact 11.11.15, and Fact 11.11.17.)

Fact 4.10.3. Define $A, B \in \mathbb{R}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} 1 & -2 & & & \\ & 1 & -2 & & \\ & & 1 & \ddots & \\ & & & \ddots & -2 \\ & & & & 1 \end{bmatrix}$$

and

$$B \triangleq \begin{bmatrix} 1 & -2 & & & \\ & 1 & -2 & & \\ & & 1 & \ddots & \\ & & & \ddots & -2 \\ \alpha & & & & 1 \end{bmatrix},$$

where $\alpha \triangleq -1/2^{n-1}$. Then,

$$\text{spec}(A) = \{1\}$$

and

$$\det B = 0.$$

Fact 4.10.4. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$|\text{spabs}(A)| \leq \text{sprad}(A).$$

Fact 4.10.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and assume that $\text{sprad}(I - A) < 1$. Then,

$$A^{-1} = \sum_{k=0}^{\infty} (I - A)^k.$$

Fact 4.10.6. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. If $\operatorname{tr} A^k = \operatorname{tr} B^k$ for all $k \in \{1, \dots, \max\{m, n\}\}$, then A and B have the same nonzero eigenvalues with the same algebraic multiplicity. Now, assume that $n = m$. Then, $\operatorname{tr} A^k = \operatorname{tr} B^k$ for all $k \in \{1, \dots, n\}$ if and only if $\operatorname{mspec}(A) = \operatorname{mspec}(B)$. (Proof: Use *Newton's identities*. See Fact 4.8.2.) (Remark: This result yields Proposition 4.4.10 since $\operatorname{tr}(AB)^k = \operatorname{tr}(BA)^k$ for all $k \geq 1$ and for all nonsquare matrices A and B .) (Remark: Setting $B = 0_{n \times n}$ yields necessity in Fact 2.12.14.)

Fact 4.10.7. Let $A \in \mathbb{F}^{n \times n}$, and let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\operatorname{mspec}(A^A) = \begin{cases} \left\{ \frac{\det A}{\lambda_1}, \dots, \frac{\det A}{\lambda_n} \right\}_{\text{ms}}, & \operatorname{rank} A = n, \\ \left\{ \sum_{i=1}^n \det A_{[i;i]}, 0, \dots, 0 \right\}_{\text{ms}}, & \operatorname{rank} A = n-1, \\ \{0\}, & \operatorname{rank} A \leq n-2. \end{cases}$$

(Remark: If $\operatorname{rank} A = n-1$ and $\lambda_n = 0$, then it follows from (4.4.19) that

$$\sum_{i=1}^n \det A_{[i;i]} = \lambda_1 \cdots \lambda_{n-1}.)$$

(Remark: See Fact 2.16.8, Fact 4.9.8, and Fact 5.11.36.)

Fact 4.10.8. Let $A \in \mathbb{F}^{n \times n}$, and let $p \in \mathbb{F}[s]$. Then, μ_A divides p if and only if $\operatorname{spec}(A) \subseteq \operatorname{roots}(p)$ and, for all $\lambda \in \operatorname{spec}(A)$, $\operatorname{ind}_A(\lambda) \leq \operatorname{mult}_p(\lambda)$.

Fact 4.10.9. Let $A \in \mathbb{F}^{n \times n}$, let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, and let $p \in \mathbb{F}[s]$. Then, the following statements hold:

- i) $\operatorname{mspec}[p(A)] = \{p(\lambda_1), \dots, p(\lambda_n)\}_{\text{ms}}$.
- ii) $\operatorname{roots}(p) \cap \operatorname{spec}(A) = \emptyset$ if and only if $p(A)$ is nonsingular.
- iii) μ_A divides p if and only if $p(A) = 0$.

Fact 4.10.10. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and let $p \in \mathbb{F}[s]$. Then,

$$p\left(\begin{bmatrix} A & B \\ 0 & C \end{bmatrix}\right) = \begin{bmatrix} p(A) & \hat{B} \\ 0 & p(C) \end{bmatrix},$$

where $\hat{B} \in \mathbb{F}^{n \times m}$.

Fact 4.10.11. Let $A_1 \in \mathbb{F}^{n \times n}$, $A_{12} \in \mathbb{F}^{n \times m}$, and $A_2 \in \mathbb{F}^{m \times m}$, and define $A \in \mathbb{F}^{(n+m) \times (n+m)}$ by

$$A \triangleq \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix}.$$

Then,

$$\chi_A = \chi_{A_1} \chi_{A_2}.$$

Furthermore,

$$\chi_{A_1}(A) = \begin{bmatrix} 0 & B_1 \\ 0 & \chi_{A_1}(A_2) \end{bmatrix}$$

and

$$\chi_{A_2}(A) = \begin{bmatrix} \chi_{A_2}(A_1) & B_2 \\ 0 & 0 \end{bmatrix},$$

where $B_1, B_2 \in \mathbb{F}^{n \times m}$. Therefore,

$$\mathcal{R}[\chi_{A_2}(A)] \subseteq \mathcal{R}\left(\begin{bmatrix} I_n \\ 0 \end{bmatrix}\right) \subseteq \mathcal{N}[\chi_{A_1}(A)]$$

and

$$\chi_{A_2}(A_1)B_1 + B_2\chi_{A_1}(A_2) = 0.$$

Hence,

$$\chi_A(A) = \chi_{A_1}(A)\chi_{A_2}(A) = \chi_{A_2}(A)\chi_{A_1}(A) = 0.$$

Fact 4.10.12. Let $A_1 \in \mathbb{F}^{n \times n}$, $A_{12} \in \mathbb{F}^{n \times m}$, and $A_2 \in \mathbb{F}^{m \times m}$, assume that $\text{spec}(A_1)$ and $\text{spec}(A_2)$ are disjoint, and define $A \in \mathbb{F}^{(n+m) \times (n+m)}$ by

$$A \triangleq \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix}.$$

Furthermore, let $\mu_1, \mu_2 \in \mathbb{F}[s]$ be such that

$$\begin{aligned} \mu_A &= \mu_1\mu_2, \\ \text{roots}(\mu_1) &= \text{spec}(A_1), \\ \text{roots}(\mu_2) &= \text{spec}(A_2). \end{aligned}$$

Then,

$$\mu_1(A) = \begin{bmatrix} 0 & B_1 \\ 0 & \mu_1(A_2) \end{bmatrix}$$

and

$$\mu_2(A) = \begin{bmatrix} \mu_2(A_1) & B_2 \\ 0 & 0 \end{bmatrix},$$

where $B_1, B_2 \in \mathbb{F}^{n \times m}$. Therefore,

$$\mathcal{R}[\mu_2(A)] \subseteq \mathcal{R}\left(\begin{bmatrix} I_n \\ 0 \end{bmatrix}\right) \subseteq \mathcal{N}[\mu_1(A)]$$

and

$$\mu_2(A_1)B_1 + B_2\mu_1(A_2) = 0.$$

Hence,

$$\mu_A(A) = \mu_1(A)\mu_2(A) = \mu_2(A)\mu_1(A) = 0.$$

Fact 4.10.13. Let $A_1, A_2, A_3, A_4, B_1, B_2 \in \mathbb{F}^{n \times n}$, and define $A \in \mathbb{F}^{4n \times 4n}$ by

$$A \triangleq \begin{bmatrix} A_1 & B_1 & 0 & 0 \\ 0 & A_2 & 0 & 0 \\ 0 & 0 & A_3 & 0 \\ 0 & 0 & B_2 & A_4 \end{bmatrix}.$$

Then,

$$\text{mspec}(A) = \bigcup_{i=1}^4 \text{mspec}(A_i).$$

Fact 4.10.14. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$, and assume that $m < n$. Then,
 $\text{mspec}(I_n + AB) = \text{mspec}(I_m + BA) \cup \{1, \dots, 1\}_{\text{ms}}.$

Fact 4.10.15. Let $a, b \in \mathbb{F}$, and define the symmetric, Toeplitz matrix $A \in \mathbb{F}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} a & b & b & \cdots & b \\ b & a & b & \cdots & b \\ b & b & a & \cdots & b \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ b & b & b & \cdots & a \end{bmatrix}.$$

Then,

$$\text{mspec}(A) = \{a + (n-1)b, a-b, \dots, a-b\}_{\text{ms}},$$

$$A1_n = [a + (n-1)b]1_n,$$

and

$$A^2 + a_1A + a_0I = 0,$$

where $a_1 \triangleq -2a + (2-n)b$ and $a_0 \triangleq a^2 + (n-2)ab + (1-n)b^2$. Finally,

$$\text{mspec}(aI_n + b1_{n \times n}) = \{a + nb, a, \dots, a\}_{\text{ms}}.$$

(Remark: See Fact 2.13.13 and Fact 8.9.34.) (Remark: For the remaining eigenvectors of A , see [1184, pp. 149, 317].)

Fact 4.10.16. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{spec}(A) \subset \bigcup_{i=1}^n \left\{ s \in \mathbb{C}: |s - A_{(i,i)}| \leq \sum_{\substack{j=1 \\ j \neq i}}^n |A_{(i,j)}| \right\}.$$

(Remark: This result is the *Gershgorin circle theorem*. See [268, 1370] for a proof and related results.) (Remark: This result yields Corollary 9.4.5 for $\|\cdot\|_{\text{col}}$ and $\|\cdot\|_{\text{row}}$.)

Fact 4.10.17. Let $A \in \mathbb{F}^{n \times n}$, and assume that, for all $i = 1, \dots, n$,

$$\sum_{\substack{j=1 \\ j \neq i}}^n |A_{(i,j)}| < |A_{(i,i)}|.$$

Then, A is nonsingular. (Proof: Apply the Gershgorin circle theorem.) (Remark: This result is the *diagonal dominance theorem*, and A is *diagonally dominant*. See [1174] for a history of this result.) (Remark: For related results, see Fact 4.10.19 and [456, 1020, 1107].)

Fact 4.10.18. Let $A \in \mathbb{F}^{n \times n}$, assume that, for all $i = 1, \dots, n$, $A_{(i,i)} \neq 0$, and assume that

$$\alpha_i \triangleq \frac{\sum_{j=1, j \neq i}^n |A_{(i,j)}|}{|A_{(i,i)}|} < 1.$$

Then,

$$|A_{(1,1)}| \prod_{i=2}^n (|A_{(i,i)}| - l_i + L_i) \leq |\det A|,$$

where

$$l_i \triangleq \sum_{j=1}^{i-1} \alpha_j |A_{(i,j)}|, \quad L_i \triangleq \left| \frac{A_{(i,1)}}{A_{(1,1)}} \right| \sum_{j=i+1}^n |A_{(i,j)}|.$$

(Proof: See [256].) (Remark: Note that, for all $i = 1, \dots, n$, $l_i = \sum_{j=1}^{i-1} \alpha_j |A_{(i,j)}| \leq \sum_{j=1, j \neq i}^n \alpha_j |A_{(i,j)}| \leq \sum_{j=1, j \neq i}^n |A_{(i,j)}| = \alpha_i |A_{(i,i)}| < |A_{(i,i)}|$. Hence, the lower bound for $|\det A|$ is positive.)

Fact 4.10.19. Let $A \in \mathbb{F}^{n \times n}$, and, for all $i = 1, \dots, n$, define

$$r_i \triangleq \sum_{\substack{j=1 \\ j \neq i}}^n |A_{(i,j)}|, \quad c_i \triangleq \sum_{\substack{j=1 \\ j \neq i}}^n |A_{(j,i)}|.$$

Furthermore, assume that at least one of the following conditions is satisfied:

- i) For all distinct $i, j = 1, \dots, n$, $r_i c_j < |A_{(i,i)} A_{(j,j)}|$.
- ii) A is irreducible, for all $i = 1, \dots, n$ it follows that $r_i \leq |A_{(i,i)}|$, and there exists $i \in \{1, \dots, n\}$ such that $r_i < |A_{(i,i)}|$.
- iii) There exist positive integers $k_1, \dots, k_n$ such that $\sum_{i=1}^n (1 + k_i)^{-1} \leq 1$ and such that, for all $i = 1, \dots, n$, $k_i \max_{j=1, \dots, n, j \neq i} |A_{(i,j)}| < |A_{(i,i)}|$.
- iv) There exists $\alpha \in [0, 1]$ such that, for all $i = 1, \dots, n$, $r_i^\alpha c_i^{1-\alpha} < |A_{(i,i)}|$.

Then, A is nonsingular. (Proof: See [101].) (Remark: All three conditions yield stronger results than Fact 4.10.17.)

Fact 4.10.20. Let $A \in \mathbb{R}^{n \times n}$, assume that A is symmetric, and, for $i = 1, \dots, n$, define

$$\alpha_i \triangleq \sum_{\substack{j=1 \\ j \neq i}}^n |A_{(i,j)}|.$$

Then,

$$\text{spec}(A) \subset \bigcup_{i=1}^n [A_{(i,i)} - \alpha_i, A_{(i,i)} + \alpha_i].$$

Furthermore, for $i = 1, \dots, n$, define

$$\beta_i \triangleq \max\{0, \max_{\substack{j=1, n \\ j \neq i}} A_{(i,j)}\}$$

and

$$\gamma_i \triangleq \min\{0, \min_{\substack{j=1, n \\ j \neq i}} A_{(i,j)}\}.$$

Then,

$$\text{spec}(A) \subset \bigcup_{i=1}^n \left[\sum_{j=1}^n A_{(i,j)} - n\beta_i, \sum_{j=1}^n A_{(i,j)} - n\gamma_i \right].$$

(Proof: The first statement is the specialization of the Gershgorin circle theorem to real, symmetric matrices. See Fact 4.10.16. The second result is given in [137].)

Fact 4.10.21. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{spec}(A) \subset \bigcup_{\substack{i,j=1 \\ i \neq j}}^n \left\{ s \in \mathbb{C}: |s - A_{(i,i)}| |s - A_{(j,j)}| \leq \sum_{\substack{k=1 \\ k \neq i}}^n |A_{(i,k)}| \sum_{\substack{k=1 \\ k \neq j}}^n |A_{(j,k)}| \right\}.$$

(Remark: The inclusion region is the *ovals of Cassini*. The result is due to Brauer. See [709, p. 380].)

Fact 4.10.22. Let $A \in \mathbb{F}^{n \times n}$, and let λ_n denote the eigenvalue of A of smallest absolute value. Then,

$$|\lambda_n| \leq \max_{i=1, \dots, n} |\text{tr } A^i|^{1/i}.$$

Furthermore,

$$\text{sprad}(A) \leq \max_{i=1, \dots, 2n-1} |\text{tr } A^i|^{1/i}$$

and

$$\text{sprad}(A) \leq \frac{5}{n} \max_{i=1, \dots, n} |\text{tr } A^i|^{1/i}.$$

(Remark: These results are *Turan's inequalities*. See [1010, p. 657].)

Fact 4.10.23. Let $A \in \mathbb{F}^{n \times n}$, and, for $j = 1, \dots, n$, define $b_j \triangleq \sum_{i=1}^n |A_{(i,j)}|$. Then,

$$\sum_{j=1}^n |A_{(j,j)}| / b_j \leq \text{rank } A.$$

(Proof: See [1098, p. 67].) (Remark: Interpret $0/0$ as 0 .) (Remark: See Fact 4.10.17.)

Fact 4.10.24. Let $A_1, \dots, A_r \in \mathbb{F}^{n \times n}$, assume that $A_1, \dots, A_r$ are normal, and let $A \in \text{co} \{A_1, \dots, A_r\}$. Then,

$$\text{spec}(A) \subseteq \text{co} \bigcup_{i=1, \dots, r} \text{spec}(A_i).$$

(Proof: See [1399].) (Remark: See Fact 8.14.7.)

Fact 4.10.25. Let $A, B \in \mathbb{R}^{n \times n}$. Then,

$$\text{mspec} \left(\begin{bmatrix} A & B \\ B & A \end{bmatrix} \right) = \text{mspec}(A + B) \cup \text{mspec}(A - B).$$

(Proof: See [1184, p. 93].) (Remark: See Fact 2.14.26.)

Fact 4.10.26. Let $A, B \in \mathbb{R}^{n \times n}$. Then,

$$\text{mspec} \left(\begin{bmatrix} A & B \\ -B & A \end{bmatrix} \right) = \text{mspec}(A + jB) \cup \text{mspec}(A - jB).$$

Now, assume that A is symmetric and B is skew symmetric. Then, $\begin{bmatrix} A & B \\ B^T & A \end{bmatrix}$ is symmetric, $A + jB$ is Hermitian, and

$$\text{mspec}\left(\begin{bmatrix} A & B \\ B^T & A \end{bmatrix}\right) = \text{mspec}(A + jB) \cup \text{mspec}(A - jB).$$

(Remark: See Fact 2.19.3 and Fact 8.15.6.)

Fact 4.10.27. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{n \times m}$, assume that A and B are Hermitian, and define $\mathcal{A}_0 \triangleq \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$ and $\mathcal{A} \triangleq \begin{bmatrix} A & C \\ C^* & B \end{bmatrix}$. Furthermore, define

$$\eta \triangleq \min_{\substack{i=1,\dots,n \\ j=1,\dots,m}} |\lambda_i(A) - \lambda_j(B)|.$$

Then, for all $i = 1, \dots, n + m$,

$$|\lambda_i(\mathcal{A}) - \lambda_i(\mathcal{A}_0)| \leq \frac{2\sigma_{\max}^2(C)}{\eta + \sqrt{\eta^2 + 4\sigma_{\max}^2(C)}}.$$

(Proof: See [200, pp. 142–146] or [893].)

Fact 4.10.28. Let $A \in \mathbb{R}^{n \times n}$, let $b, c \in \mathbb{R}^n$, define $p \in \mathbb{R}[s]$ by $p(s) \triangleq c^T(sI - A)^A b$, assume that p and $\det(sI - A)$ are coprime, define $A_\alpha \triangleq A + \alpha bc^T$ for all $\alpha \in [0, \infty)$, and let $\lambda: [0, \infty) \rightarrow \mathbb{C}$ be a continuous function such that $\lambda(\alpha) \in \text{spec}(A_\alpha)$ for all $\alpha \in [0, \infty)$. Then, either $\lim_{\alpha \rightarrow \infty} |\lambda(\alpha)| = \infty$ or $\lim_{\alpha \rightarrow \infty} \lambda(\alpha) \in \text{roots}(p)$. (Remark: This result is a consequence of *root locus* analysis from classical control theory, which determines asymptotic pole locations under high-gain feedback.)

Fact 4.10.29. Let $A \in \mathbb{F}^{n \times n}$, where $n \geq 2$, and assume that there exist $\alpha \in [0, \infty)$ and $B \in \mathbb{F}^{n \times n}$ such that $A = \alpha I - B$ and $\text{sprad}(B) \leq \alpha$. Then,

$$\text{spec}(A) \subset \{0\} \cup \text{ORHP}.$$

If, in addition, $\text{sprad}(B) < \alpha$, then

$$\text{spec}(A) \subset \text{ORHP},$$

and thus A is nonsingular. (Proof: Let $\lambda \in \text{spec}(A)$. Then, there exists $\mu \in \text{spec}(B)$ such that $\lambda = \alpha - \mu$. Hence, $\text{Re } \lambda = \alpha - \text{Re } \mu$. Since $\text{Re } \mu \leq |\text{Re } \mu| \leq |\mu| \leq \text{sprad}(B)$, it follows that $\text{Re } \lambda \geq \alpha - |\text{Re } \mu| \geq \alpha - |\mu| \geq \alpha - \text{sprad}(B) \geq 0$. Hence, $\text{Re } \lambda \geq 0$. Now, suppose that $\text{Re } \lambda = 0$. Then, since $\alpha - \lambda = \mu \in \text{spec}(B)$, it follows that $\alpha^2 + |\lambda|^2 \leq [\text{sprad}(B)]^2 \leq \alpha^2$. Hence, $\lambda = 0$. By a similar argument, if $\text{sprad}(B) < \alpha$, then $\text{Re } \lambda > 0$.) (Remark: Converses of these statements hold when B is nonnegative. See Fact 4.11.6.)

4.11 Facts on Graphs and Nonnegative Matrices

Fact 4.11.1. Let $\mathcal{G} = (\{x_1, \dots, x_n\}, \mathcal{R})$ be a graph without self-loops, assume that $\mathcal{G}$ is antisymmetric, let $A \in \mathbb{R}^{n \times n}$ denote the adjacency matrix of $\mathcal{G}$, let $L_{\text{in}} \in \mathbb{R}^{n \times n}$ and $L_{\text{out}} \in \mathbb{R}^{n \times n}$ denote the inbound and outbound Laplacians of $\mathcal{G}$, respectively, and let A_{sym} , D_{sym} , and L_{sym} denote the adjacency, degree, and

Laplacian matrices, respectively, of $\text{sym}(\mathcal{G})$. Then,

$$D_{\text{sym}} = D_{\text{in}} + D_{\text{out}},$$

$$A_{\text{sym}} = A + A^T,$$

and

$$L_{\text{sym}} = L_{\text{in}} + L_{\text{out}}^T = L_{\text{in}}^T + L_{\text{out}} = D_{\text{sym}} - A_{\text{sym}}.$$

Fact 4.11.2. Let $\mathcal{G} = (\{x_1, \dots, x_n\}, \mathcal{R})$ be a graph, and let $A \in \mathbb{R}^{n \times n}$ be the adjacency matrix of $\mathcal{G}$. Then, the following statements are equivalent:

- i) $\mathcal{G}$ is connected.
- ii) $\mathcal{G}$ has no directed cuts.
- iii) A is irreducible.

Furthermore, the following statements are equivalent:

- iv) $\mathcal{G}$ is not connected.
- v) $\mathcal{G}$ has a directed cut.
- vi) A is reducible.

Finally, suppose that A is reducible and there exist $k \geq 1$ and a permutation matrix $S \in \mathbb{R}^{n \times n}$ such that $SAS^T = \begin{bmatrix} B & C \\ 0_{k \times (n-k)} & D \end{bmatrix}$, where $B \in \mathbb{F}^{(n-k) \times (n-k)}$, $C \in \mathbb{F}^{(n-k) \times k}$, and $D \in \mathbb{F}^{k \times k}$. Then, $(\{x_{i_1}, \dots, x_{i_{n-k}}\}, \{x_{i_{n-k+1}}, \dots, x_{i_n}\})$ is a directed cut, where $\begin{bmatrix} i_1 & \dots & i_n \end{bmatrix}^T = S \begin{bmatrix} 1 & \dots & n \end{bmatrix}^T$. (Proof: See [709, p. 362].)

Fact 4.11.3. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph, where $\mathcal{X} = \{x_1, \dots, x_n\}$, and let A be the adjacency matrix of $\mathcal{G}$. Then, the following statements hold:

- i) The number of distinct walks from x_i to x_j of length $k \geq 1$ is $(A^k)_{(j,i)}$.
- ii) Let k be an integer such that $1 \leq k \leq n-1$. Then, for distinct $x_i, x_j \in \mathcal{X}$, the number of distinct walks from x_i to x_j whose length is less than or equal to k is $[(I + A)^k]_{(j,i)}$.

Fact 4.11.4. Let $A \in \mathbb{F}^{n \times n}$, and consider $\mathcal{G}(A) = (\mathcal{X}, \mathcal{R})$, where $\mathcal{X} = \{x_1, \dots, x_n\}$. Then, the following statements are equivalent:

- i) $\mathcal{G}(A)$ is connected.
- ii) There exists $k \geq 1$ such that $(I + |A|)^{k-1}$ is positive.
- iii) $(I + |A|)^{n-1}$ is positive.

(Proof: See [709, pp. 358, 359].)

Fact 4.11.5. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, and assume that A is nonnegative. Then, the following statements hold:

- i) $\text{sprad}(A)$ is an eigenvalue of A .
- ii) There exists a nonzero nonnegative vector $x \in \mathbb{R}^n$ such that $Ax =$

$\text{sprad}(A)x$.

Furthermore, the following statements are equivalent:

- iii) A is irreducible.
- iv) $(I + A)^{n-1}$ is positive.
- v) $\mathcal{G}(A)$ is connected.
- vi) A has exactly one nonnegative eigenvector whose components sum to 1, and this eigenvector is positive.

If A is irreducible, then the following statements hold:

- vii) $\text{sprad}(A) > 0$.
- viii) $\text{sprad}(A)$ is a simple eigenvalue of A .
- ix) There exists a positive vector $x \in \mathbb{R}^n$ such that $Ax = \text{sprad}(A)x$.
- x) A has exactly one positive eigenvector whose components sum to 1.
- xi) Assume that $\{\lambda_1, \dots, \lambda_k\}_{\text{ms}} = \{\lambda \in \text{mspec}(A) : |\lambda| = \text{sprad}(A)\}_{\text{ms}}$. Then, $\lambda_1, \dots, \lambda_k$ are distinct, and

$$\{\lambda_1, \dots, \lambda_k\} = \{e^{2\pi j i/k} \text{sprad}(A) : i = 1, \dots, k\}.$$

Furthermore,

$$\text{mspec}(A) = e^{2\pi j/k} \text{mspec}(A).$$

- xii) If at least one diagonal entry of A is positive, then $\text{sprad}(A)$ is the only eigenvalue of A whose absolute value is $\text{sprad}(A)$.
- xiii) If A has at least m positive diagonal entries, then A^{2n-m-1} is positive.

In addition, the following statements are equivalent:

- xiv) There exists $k \geq 1$ such that A^k is positive.
- xv) A is irreducible and $|\lambda| < \text{sprad}(A)$ for all $\lambda \in \text{spec}(A) \setminus \{\text{sprad}(A)\}$.
- xvi) A^{n^2-2n+2} is positive.
- xvii) $\mathcal{G}(A)$ is aperiodic.

A is *primitive* if *xiv)*–*xviii)* are satisfied. (Example: $\begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$ is irreducible but not primitive.) If A is primitive, then the following statements hold:

- xviii) For all $k \in \mathbb{P}$, A^k is primitive.
- xix) If $k \in \mathbb{P}$ and A^k is positive, then, for all $l \geq k$, A^l is positive.
- xx) There exists a positive integer $k \leq (n-1)n^n$ such that A^k is positive.
- xxi) If $x, y \in \mathbb{R}^n$ are positive and satisfy $Ax = \text{sprad}(A)x$ and $A^T y = \text{sprad}(A)y$, then

$$\lim_{k \rightarrow \infty} ([\text{sprad}(A)]^{-1}A)^k = \frac{1}{x^T y} xy^T.$$

- xxii) If $x_0 \in \mathbb{R}^n$ is nonzero and nonnegative and $x, y \in \mathbb{R}^n$ are positive and

satisfy $Ax = \text{sprad}(A)x$ and $A^T y = \text{sprad}(A)y$, then

$$\lim_{k \rightarrow \infty} \frac{A^k x_0 - [\text{sprad}(A)]^k y^T x_0 x}{\|A^k x_0\|_2} = 0.$$

xxiii) $\text{sprad}(A) = \lim_{k \rightarrow \infty} (\text{tr } A^k)^{1/k}$.

(Remark: For an arbitrary nonzero and nonnegative initial condition, the state $x_k = A^k x_0$ of the difference equation $x_{k+1} = Ax_k$ approaches a distribution given by the eigenvector associated with the positive eigenvalue of maximum absolute value. In demography, this eigenvector is interpreted as the *stable age distribution*. See [805, pp. 47, 63].) (Proof: See [16, pp. 45–49], [133, p. 17], [181, pp. 26–28, 32, 55], [481], and [709, pp. 507–518]. For *xxiii)*, see [1193] and [1369, p. 49].) (Remark: This result is the *Perron-Frobenius theorem*.) (Remark: See Fact 11.18.20.) (Remark: Statement *xvi)* is due to Wielandt. See [1098, p. 157].) (Remark: Statement *xvii)* is given in [1148, p. 9–3].) (Remark: See Fact 6.6.20.) (Example: Let x and y be positive numbers such that $x + y < 1$, and define

$$A \triangleq \begin{bmatrix} x & y & 1 - x - y \\ 1 - x - y & x & y \\ y & 1 - x - y & x \end{bmatrix}.$$

Then, $A1_{3 \times 1} = A^T 1_{3 \times 1} = 1_{3 \times 1}$, and thus $\lim_{k \rightarrow \infty} A^k = \frac{1}{3} 1_{3 \times 3}$. See [238, p. 213].)

Fact 4.11.6. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, and assume that A is a Z-matrix. Then, the following statements are equivalent:

- i)* There exist $\alpha \in (0, \infty)$ and $B \in \mathbb{R}^{n \times n}$ such that $A = \alpha I - B$, B is nonnegative, and $\text{sprad}(B) \leq \alpha$.
- ii)* $\text{spec}(A) \subset \text{ORHP} \cup \{0\}$.
- iii)* $\text{spec}(A) \subset \text{CRHP}$.
- iv)* If $\lambda \in \text{spec}(A)$ is real, then $\lambda \geq 0$.
- v)* Every principal subdeterminant of A is nonnegative.
- vi)* For every diagonal, positive-definite matrix $D \in \mathbb{R}^{n \times n}$, it follows that $A + D$ is nonsingular.

(Remark: A is an *M-matrix* if A is a Z-matrix and *i)–v)* hold. Example: $A = \begin{bmatrix} 0 & -1 \\ 0 & 0 \end{bmatrix} = I - \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$). In addition, the following statements are equivalent:

- vii)* There exist $\alpha \in (0, \infty)$ and $B \in \mathbb{R}^{n \times n}$ such that $A = \alpha I - B$, B is nonnegative, and $\text{sprad}(B) < \alpha$.
- viii)* $\text{spec}(A) \subset \text{ORHP}$.

(Proof: The result *i)* $\implies$ *ii)* follows from Fact 4.10.29, while *ii)* $\implies$ *iii)* is immediate. To prove *iii)* $\implies$ *i)*, let $\alpha \in (0, \infty)$ be sufficiently large that $B \triangleq \alpha I - A$ is nonnegative. Hence, for every $\mu \in \text{spec}(B)$, it follows that $\lambda \triangleq \alpha - \mu \in \text{spec}(A)$. Since $\text{Re } \lambda \geq 0$, it follows that every $\mu \in \text{spec}(B)$ satisfies $\text{Re } \mu \leq \alpha$. Since B is nonnegative, it follows from *i)* of Fact 4.11.5 that $\text{sprad}(B)$ is an eigenvalue of B . Hence, setting $\mu = \text{sprad}(B)$ implies that $\text{sprad}(B) \leq \alpha$. Conditions *iv)* and *v)* are proved in [182, pp. 149, 150]. Finally, the argument used to prove that *i)* $\implies$ *ii)*

shows in addition that $vii) \implies viii)$.) (Remark: A is a *nonsingular M-matrix* if $vii)$ and $viii)$ hold. See Fact 11.19.5.) (Remark: See Fact 11.19.3.)

Fact 4.11.7. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$. If A is a Z-matrix, then every principal submatrix of A is also a Z-matrix. Furthermore, if A is an M-matrix, then every principal submatrix of A is also an M-matrix. (Proof: See [711, p. 114].)

Fact 4.11.8. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, and assume that A is a nonsingular M-matrix, B is a Z-matrix, and $A \leq B$. Then, the following statements hold:

- i) $\text{tr}(A^{-1}A^T) \leq n$.
- ii) $\text{tr}(A^{-1}A^T) = n$ if and only if A is symmetric.
- iii) B is a nonsingular M-matrix.
- iv) $0 \leq B^{-1} \leq A^{-1}$.
- v) $0 < \det A \leq \det B$.

(Proof: See [711, pp. 117, 370].)

Fact 4.11.9. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, assume that A is a Z-matrix, and define

$$\tau(A) \triangleq \min\{\text{Re } \lambda : \lambda \in \text{spec}(A)\}.$$

Then, the following statements hold:

- i) $\tau(A) \in \text{spec}(A)$.
- ii) $\min_{i=1, \dots, n} \sum_{j=1}^n A_{(i,j)} \leq \tau(A)$.

Now, assume that A is an M-matrix. Then, the following statements hold:

- iii) If A is nonsingular, then $\tau(A) = 1/\text{sprad}(A^{-1})$.
- iv) $[\tau(A)]^n \leq \det A$.
- v) If $B \in \mathbb{R}^{n \times n}$, B is an M-matrix, and $B \leq A$, then $\tau(B) \leq \tau(A)$.

(Proof: See [711, pp. 128–131].) (Remark: $\tau(A)$ is the *minimum eigenvalue* of A .) (Remark: See Fact 7.6.15.)

Fact 4.11.10. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, and assume that A is an M-matrix. Then, the following statements hold:

- i) There exists a nonzero nonnegative vector $x \in \mathbb{R}^n$ such that Ax is nonnegative.
- ii) If A is irreducible, then there exists a positive vector $x \in \mathbb{R}^n$ such that Ax is nonnegative.

Now, assume that A is singular. Then, the following statements hold:

- iii) $\text{rank } A = n - 1$.
- iv) There exists a positive vector $x \in \mathbb{R}^n$ such that $Ax = 0$.

- v) A is group invertible.
- vi) Every principal submatrix of A of order less than n and greater than 1 is a nonsingular M-matrix.
- vii) If $x \in \mathbb{R}^n$ and Ax is nonnegative, then $Ax = 0$.

(Proof: To prove the first statement, it follows from Fact 4.11.6 that there exist $\alpha \in (0, \infty)$ and $B \in \mathbb{R}^{n \times n}$ such that $A = \alpha I - B$, B is nonnegative, and $\text{sprad}(B) \leq \alpha$. Consequently, it follows from ii) of Fact 4.11.5 that there exists a nonzero nonnegative vector $x \in \mathbb{R}^n$ such that $Bx = \text{sprad}(B)x$. Therefore, $Ax = [\alpha - \text{sprad}(B)]x$ is nonnegative. Statements iii)–vii) are given in [182, p. 156].)

Fact 4.11.11. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a symmetric graph, where $\mathcal{X} = \{x_1, \dots, x_n\}$, and let $L_{\text{in}} \in \mathbb{R}^{n \times n}$ denote the Laplacian of $\mathcal{G}$. Then, the following statements hold:

- i) $\text{spec}(L) \subset \{0\} \cup \text{ORHP}$.
- ii) $0 \in \text{spec}(L)$, and an associated eigenvector is $1_{n \times 1}$.
- iii) 0 is a semisimple eigenvalue of L .
- iv) 0 is a simple eigenvalue of L if and only if $\mathcal{G}$ has a spanning subgraph that is a tree.
- v) L is positive semidefinite.
- vi) $0 \in \text{spec}(L) \subset \{0\} \cup [0, \infty)$.
- vii) If $\mathcal{G}$ is connected, then 0 is a simple eigenvalue of L .
- viii) $\mathcal{G}$ is connected if and only if $\lambda_{n-1}(L)$ is positive.

(Proof: For the last statement, see [993, p. 147].) (Remark: See Fact 11.19.7.) (Problem: Extend these results to graphs that are not symmetric.)

Fact 4.11.12. Let $A \triangleq \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$. Then, $\chi_A(s) = s^2 - s - 1$ and $\text{spec}(A) = \{\alpha, \beta\}$, where $\alpha \triangleq \frac{1}{2}(1 + \sqrt{5}) \approx 1.61803$ and $\beta \triangleq \frac{1}{2}(1 - \sqrt{5}) \approx -0.61803$ satisfy

$$\alpha - 1 = 1/\alpha, \quad \beta - 1 = 1/\beta.$$

Furthermore, $\begin{bmatrix} \alpha \\ 1 \end{bmatrix}$ is an eigenvector of A associated with α . Now, for $k \geq 0$, consider the difference equation

$$x_{k+1} = Ax_k.$$

Then, for all $k \geq 0$,

$$x_k = A^k x_0$$

and

$$x_{k+2(1)} = x_{k+1(1)} + x_{k(1)}.$$

Furthermore, if x_0 is positive, then

$$\lim_{k \rightarrow \infty} \frac{x_{k(1)}}{x_{k(2)}} = \alpha.$$

In particular, if $x_0 \triangleq \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, then, for all $k \geq 0$,

$$x_k = \begin{bmatrix} F_{k+2} \\ F_{k+1} \end{bmatrix},$$

where $F_1 \triangleq F_2 \triangleq 1$ and, for all $k \geq 1$, F_k is given by

$$F_k = \frac{1}{\sqrt{5}}(\alpha^k - \beta^k)$$

and satisfies

$$F_{k+2} = F_{k+1} + F_k.$$

Furthermore,

$$\frac{1}{1-x-x^2} = F_1x + F_2x^2 + \dots$$

and

$$A^k = \begin{bmatrix} F_{k+1} & F_k \\ F_k & F_{k-1} \end{bmatrix}.$$

On the other hand, if $x_0 \triangleq \begin{bmatrix} 3 \\ 1 \end{bmatrix}$, then, for all $k \geq 0$,

$$x_k = \begin{bmatrix} L_{k+2} \\ L_{k+1} \end{bmatrix},$$

where $L_1 \triangleq 1$, $L_2 \triangleq 3$, and, for all $k \geq 1$, L_k is given by

$$L_k = \alpha^k + \beta^k$$

and satisfies

$$L_{k+2} = L_{k+1} + L_k.$$

Moreover,

$$\lim_{k \rightarrow \infty} \frac{F_{k+1}}{F_k} = \frac{L_{k+1}}{L_k} = \alpha.$$

In addition,

$$\alpha = \sqrt{1 + \sqrt{1 + \sqrt{1 + \sqrt{1 + \dots}}}}$$

Finally, for all $k \geq 1$,

$$F_{k+1} = \det \begin{bmatrix} 1 & j & 0 & \cdots & 0 & 0 \\ j & 1 & j & \cdots & 0 & 0 \\ 0 & j & 1 & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & \ddots & 1 & j \\ 0 & 0 & 0 & \cdots & j & 1 \end{bmatrix} = \det \begin{bmatrix} 1 & 1 & 0 & \cdots & 0 & 0 \\ -1 & 1 & 1 & \cdots & 0 & 0 \\ 0 & -1 & 1 & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & 0 & 0 & \ddots & 1 & 1 \\ 0 & 0 & 0 & \cdots & -1 & 1 \end{bmatrix},$$

where both matrices are of size $k \times k$. (Proof: Use the last statement of Fact 4.11.5.) (Remark: F_k is the k th *Fibonacci number*, L_k is the k th *Lucas number*, and α is the *golden ratio*. See [841, pp. 6–8, 239–241, 362, 363] and Fact 12.23.4. The expressions for F_k and L_k involving powers of α and β are *Binet's formulas*. See [177, p. 125]. The iterated square root identity is given in [477, p. 24]. The determinant identities are given in [279] and [1119, p. 515].) (Remark: $1/(1-x-x^2)$ is a *generating function* for the Fibonacci numbers. See [1407].)

Fact 4.11.13. Consider the nonnegative companion matrix $A \in \mathbb{R}^{n \times n}$ defined by

$$A \triangleq \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \ddots & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 1/n & 1/n & 1/n & \cdots & 1/n & 1/n \end{bmatrix}.$$

Then, A is irreducible, 1 is a simple eigenvalue of A with associated eigenvector $1_{n \times 1}$, and $|\lambda| < 1$ for all $\lambda \in \text{spec}(A) \setminus \{1\}$. Furthermore, if $x \in \mathbb{R}^n$, then

$$\lim_{k \rightarrow \infty} A^k x = \left[\frac{2}{n(n+1)} \sum_{i=1}^n i x_{(i-1)} \right] 1_{n \times 1}.$$

(Proof: See [629, pp. 82, 83, 263–266].) (Remark: The result follows from Fact 4.11.5.)

Fact 4.11.14. Let $A \in \mathbb{R}^{n \times m}$ and $b \in \mathbb{R}^m$. Then, the following statements are equivalent:

- i) If $x \in \mathbb{R}^m$ and $Ax \geq 0$, then $b^T x \geq 0$.
- ii) There exists a vector $y \in \mathbb{R}^n$ such that $y \geq 0$ and $A^T y = b$.

Equivalently, exactly one of the following two statements is satisfied:

- iii) There exists a vector $x \in \mathbb{R}^m$ such that $Ax \geq 0$ and $b^T x < 0$.
- iv) There exists a vector $y \in \mathbb{R}^n$ such that $y \geq 0$ and $A^T y = b$.

(Proof: See [157, p. 47] or [239, p. 24].) (Remark: This result is the *Farkas theorem*.)

Fact 4.11.15. Let $A \in \mathbb{R}^{n \times m}$. Then, the following statements are equivalent:

- i) There exists a vector $x \in \mathbb{R}^m$ such that $Ax >> 0$.
- ii) If $y \in \mathbb{R}^n$ is nonzero and $y \geq 0$, then $A^T y \neq 0$.

Equivalently, exactly one of the following two statements is satisfied:

- iii) There exists a vector $x \in \mathbb{R}^m$ such that $Ax >> 0$.
- iv) There exists a nonzero vector $y \in \mathbb{R}^n$ such that $y \geq 0$ and $A^T y = 0$.

(Proof: See [157, p. 47] or [239, p. 23].) (Remark: This result is *Gordan's theorem*.)

Fact 4.11.16. Let $A \in \mathbb{C}^{n \times n}$, and define $|A| \in \mathbb{R}^{n \times n}$ by $|A|_{(i,j)} \triangleq |A_{(i,j)}|$ for all $i, j = 1, \dots, n$. Then,

$$\text{sprad}(A) \leq \text{sprad}(|A|).$$

(Proof: See [998, p. 619].)

Fact 4.11.17. Let $A \in \mathbb{R}^{n \times n}$, assume that A is nonnegative, and let $\alpha \in [0, 1]$. Then,

$$\text{sprad}(A) \leq \text{sprad}[\alpha A + (1 - \alpha)A^T].$$

(Proof: See [130].)

Fact 4.11.18. Let $A, B \in \mathbb{R}^{n \times n}$, where $0 \leq A \leq B$. Then,

$$\text{sprad}(A) \leq \text{sprad}(B).$$

In particular, $B_0 \in \mathbb{R}^{m \times m}$ is a principal submatrix of B , then

$$\text{sprad}(B_0) \leq \text{sprad}(B).$$

If, in addition, $A \neq B$ and $A + B$ is irreducible, then

$$\text{sprad}(A) < \text{sprad}(B).$$

Hence, if $\text{sprad}(A) = \text{sprad}(B)$ and $A + B$ is irreducible, then $A = B$. (Proof: See [170, p. 27]. See also [447, pp. 500, 501].)

Fact 4.11.19. Let $A, B \in \mathbb{R}^{n \times n}$, assume that B is diagonal, assume that A and $A + B$ are nonnegative, and let $\alpha \in [0, 1]$. Then,

$$\text{sprad}[\alpha A + (1 - \alpha)B] \leq \alpha \text{sprad}(A) + (1 - \alpha) \text{sprad}(A + B).$$

(Proof: See [1148, p. 9-5].)

Fact 4.11.20. Let $A \in \mathbb{R}^{n \times n}$, assume that $A \gg 0$, and let $\lambda \in \text{spec}(A) \setminus \{\text{sprad}(A)\}$. Then,

$$|\lambda| \leq \frac{A_{\max} - A_{\min}}{A_{\max} + A_{\min}} \text{sprad}(A),$$

where

$$A_{\max} \triangleq \max \{A_{(i,j)} : i, j = 1, \dots, n\}$$

and

$$A_{\min} \triangleq \min \{A_{(i,j)} : i, j = 1, \dots, n\}.$$

(Remark: This result is *Hopf's theorem*.) (Remark: The equality case is discussed in [688].)

Fact 4.11.21. Let $A \in \mathbb{R}^{n \times n}$, assume that A is nonnegative and irreducible, and let $x, y \in \mathbb{R}^n$, where $x > 0$ and $y > 0$ satisfy $Ax = \text{sprad}(A)x$ and $A^T y = \text{sprad}(A)y$. Then,

$$\lim_{l \rightarrow \infty} \frac{1}{l} \sum_{k=1}^l \left[\frac{1}{\text{sprad}(A)} A \right]^k = xy^T.$$

If, in addition, A is primitive, then

$$\lim_{k \rightarrow \infty} \left[\frac{1}{\text{sprad}(A)} A \right]^k = xy^T.$$

(Proof: See [447, p. 503] and [709, p. 516].)

Fact 4.11.22. Let $A \in \mathbb{R}^{n \times n}$, assume that A is nonnegative, and let k and m be positive integers. Then,

$$[\operatorname{tr} A^k]^m \leq n^{m-1} \operatorname{tr} A^{km}.$$

(Proof: See [860].) (Remark: This result is the *JLL inequality*.)

4.12 Notes

Much of the development in this chapter is based on [1081]. Additional discussions of the Smith and Smith-McMillan forms are given in [787] and [1498]. The proofs of Lemma 4.4.8 and Leverrier's algorithm Proposition 4.4.9 are based on [1129, pp. 432, 433], where it is called the *Souriau-Frame algorithm*. Alternative proofs of Leverrier's algorithm are given in [143, 720]. The proof of Theorem 4.6.1 is based on [709]. Polynomial-based approaches to linear algebra are given in [276, 508], while polynomial matrices and rational transfer functions are studied in [559, 1368].

The term *normal rank* is often used to refer to what we call the rank of a rational transfer function.

Chapter Five

Matrix Decompositions

In this chapter we present several matrix decompositions, namely, the Smith, multicompanion, elementary multicompanion, hypercompanion, Jordan, Schur, and singular value decompositions.

5.1 Smith Form

Our first decomposition involves rectangular matrices subject to a biequivalence transformation. This result is the specialization of the Smith decomposition given by Theorem 4.3.2 to constant matrices.

Theorem 5.1.1. Let $A \in \mathbb{F}^{n \times m}$ and $r \triangleq \text{rank } A$. Then, there exist nonsingular matrices $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$ such that

$$A = S_1 \begin{bmatrix} I_r & 0_{r \times (m-r)} \\ 0_{(n-r) \times r} & 0_{(n-r) \times (m-r)} \end{bmatrix} S_2. \quad (5.1.1)$$

Corollary 5.1.2. Let $A, B \in \mathbb{F}^{n \times m}$. Then, A and B are biequivalent if and only if A and B have the same Smith form.

Proposition 5.1.3. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

- i) A and B are left equivalent if and only if $\mathcal{N}(A) = \mathcal{N}(B)$.
- ii) A and B are right equivalent if and only if $\mathcal{R}(A) = \mathcal{R}(B)$.
- iii) A and B are biequivalent if and only if $\text{rank } A = \text{rank } B$.

Proof. The proof of necessity is immediate in i)–iii). Sufficiency in iii) follows from Corollary 5.1.2. For sufficiency in i) and ii), see [1129, pp. 179–181]. $\square$

5.2 Multicompanion Form

For the monic polynomial $p(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_1s + \beta_0 \in \mathbb{F}[s]$ of degree $n \geq 1$, the *companion matrix* $C(p) \in \mathbb{F}^{n \times n}$ associated with p is defined to

be

$$C(p) \triangleq \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \ddots & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ -\beta_0 & -\beta_1 & -\beta_2 & \cdots & -\beta_{n-2} & -\beta_{n-1} \end{bmatrix}. \quad (5.2.1)$$

If $n = 1$, then $p(s) = s + \beta_0$ and $C(p) = -\beta_0$. Furthermore, if $n = 0$ and $p = 1$, then we define $C(p) \triangleq 0_{0 \times 0}$. Note that, if $n \geq 1$, then $\text{tr } C(p) = -\beta_{n-1}$ and $\det C(p) = (-1)^n \beta_0 = (-1)^n p(0)$.

It is easy to see that the characteristic polynomial of the companion matrix $C(p)$ is p . For example, let $n = 3$ so that

$$C(p) = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -\beta_0 & -\beta_1 & -\beta_2 \end{bmatrix}, \quad (5.2.2)$$

and thus

$$sI - C(p) = \begin{bmatrix} s & -1 & 0 \\ 0 & s & -1 \\ \beta_0 & \beta_1 & s + \beta_2 \end{bmatrix}. \quad (5.2.3)$$

Adding s times the second column and s^2 times the third column to the first column leaves the determinant of $sI - C(p)$ unchanged and yields

$$\begin{bmatrix} 0 & -1 & 0 \\ 0 & s & -1 \\ p(s) & \beta_1 & s + \beta_2 \end{bmatrix}. \quad (5.2.4)$$

Hence, $\chi_{C(p)} = p$. If $n = 0$ and $p = 1$, then we define $\chi_{C(p)} \triangleq \chi_{0_0 \times 0} = 1$. The following result shows that companion matrices have the same characteristic and minimal polynomials.

Proposition 5.2.1. Let $p \in \mathbb{F}[s]$ be a monic polynomial having degree n . Then, there exist unimodular matrices $S_1, S_2 \in \mathbb{F}^{n \times n}[s]$ such that

$$sI - C(p) = S_1(s) \begin{bmatrix} I_{n-1} & 0_{(n-1) \times 1} \\ 0_{1 \times (n-1)} & p(s) \end{bmatrix} S_2(s). \quad (5.2.5)$$

Furthermore,

$$\chi_{C(p)} = \mu_{C(p)} = p. \quad (5.2.6)$$

Proof. Since $\chi_{C(p)} = p$, it follows that $\text{rank}[sI - C(p)] = n$. Next, since $\det([sI - C(p)]_{[n,1]}) = (-1)^{n-1}$, it follows that $\Delta_{n-1} = 1$, where Δ_{n-1} is the greatest common divisor (which is monic by definition) of all $(n-1) \times (n-1)$ subdeterminants of $sI - C(p)$. Furthermore, since Δ_{i-1} divides Δ_i for all $i = 2, \dots, n-1$, it follows that $\Delta_1 = \cdots = \Delta_{n-2} = 1$. Consequently, $p_1 = \cdots = p_{n-1} = 1$. Since, by

Proposition 4.6.2, $\chi_{C(p)} = \prod_{i=1}^n p_i = p_n$ and $\mu_{C(p)} = p_n$, it follows that $\chi_{C(p)} = \mu_{C(p)} = p$. $\square$

Next, we consider block-diagonal matrices all of whose diagonally located blocks are companion matrices.

Lemma 5.2.2. Let $p_1, \dots, p_n \in \mathbb{F}[s]$ be monic polynomials such that p_i divides p_{i+1} for all $i = 1, \dots, n-1$ and $n = \sum_{i=1}^n \deg p_i$. Furthermore, define $C \triangleq \text{diag}[C(p_1), \dots, C(p_n)] \in \mathbb{F}^{n \times n}$. Then, there exist unimodular matrices $S_1, S_2 \in \mathbb{F}^{n \times n}[s]$ such that

$$sI - C = S_1(s) \begin{bmatrix} p_1(s) & & 0 \\ & \ddots & \\ 0 & & p_n(s) \end{bmatrix} S_2(s). \quad (5.2.7)$$

Proof. Letting $k_i = \deg p_i$, Proposition 5.2.1 implies that the Smith form of $sI_{k_i} - C(p_i)$ is $0_{0 \times 0}$ if $k_i = 0$ and $\text{diag}(I_{k_i-1}, p_i)$ if $k_i \geq 1$. Note that $p_1 = \dots = p_{n_0} = 1$, where $n_0 \triangleq \sum_{i=1}^n \max\{0, k_i - 1\}$. By combining these Smith forms and rearranging diagonal entries, it follows that there exist unimodular matrices $S_1, S_2 \in \mathbb{F}^{n \times n}[s]$ such that

$$\begin{aligned} sI - C &= \begin{bmatrix} sI_{k_1} - C(p_1) & & \\ & \ddots & \\ & & sI_{k_n} - C(p_n) \end{bmatrix} \\ &= S_1(s) \begin{bmatrix} p_1(s) & & 0 \\ & \ddots & \\ 0 & & p_n(s) \end{bmatrix} S_2(s). \end{aligned}$$

Since p_i divides p_{i+1} for all $i = 1, \dots, n-1$, it follows that this diagonal matrix is the Smith form of $sI - C$. $\square$

The following result uses Lemma 5.2.2 to construct a canonical form, known as the *multicompanion form*, for square matrices under a similarity transformation.

Theorem 5.2.3. Let $A \in \mathbb{F}^{n \times n}$, and let $p_1, \dots, p_n \in \mathbb{F}[s]$ denote the similarity invariants of A , where p_i divides p_{i+1} for all $i = 1, \dots, n-1$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A = S \begin{bmatrix} C(p_1) & & 0 \\ & \ddots & \\ 0 & & C(p_n) \end{bmatrix} S^{-1}. \quad (5.2.8)$$

Proof. Lemma 5.2.2 implies that the $n \times n$ matrix $sI - C$, where $C \triangleq \text{diag}[C(p_1), \dots, C(p_n)]$, has the Smith form $\text{diag}(p_1, \dots, p_n)$. Now, since $sI - A$ has the same similarity invariants as C , it follows from Theorem 4.3.10 that A and C are similar. $\square$

Corollary 5.2.4. Let $A \in \mathbb{F}^{n \times n}$. Then, $\mu_A = \chi_A$ if and only if A is similar to $C(\chi_A)$.

Proof. Suppose that $\mu_A = \chi_A$. Then, it follows from Proposition 4.6.2 that $p_i = 1$ for all $i = 1, \dots, n-1$ and $p_n = \chi_A$ is the only nonconstant similarity invariant of A . Thus, $C(p_i) = 0_{0 \times 0}$ for all $i = 1, \dots, n-1$, and it follows from Theorem 5.2.3 that A is similar to $C(\chi_A)$. The converse follows from (5.2.6), xi) of Proposition 4.4.5, and Proposition 4.6.3. $\square$

Corollary 5.2.5. Let $A \in \mathbb{F}^{n \times n}$ be a companion matrix. Then, $A = C(\chi_A)$ and $\mu_A = \chi_A$.

Note that, if $A = I_n$, then the similarity invariants of A are $p_i(s) = s - 1$ for all $i = 1, \dots, n$. Thus, $C(p_i) = 1$ for all $i = 1, \dots, n$, as expected.

Corollary 5.2.6. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A and B are similar.
- ii) A and B have the same similarity invariants.
- iii) A and B have the same multicompanion form.

The multicompanion form given by Theorem 5.2.3 provides a canonical form for A in terms of a block-diagonal matrix of companion matrices. As shown below, however, the multicompanion form is only one such decomposition. The goal of the remainder of this section is to obtain an additional canonical form by applying a similarity transformation to the multicompanion form.

To begin, note that, if A_i is similar to B_i for all $i = 1, \dots, r$, then $\text{diag}(A_1, \dots, A_r)$ is similar to $\text{diag}(B_1, \dots, B_r)$. Therefore, it follows from Corollary 5.2.6 that, if $sI - A_i$ and $sI - B_i$ have the same Smith form for all $i = 1, \dots, r$, then $sI - \text{diag}(A_1, \dots, A_r)$ and $sI - \text{diag}(B_1, \dots, B_r)$ have the same Smith form. The following lemma is needed.

Lemma 5.2.7. Let $A = \text{diag}(A_1, A_2)$, where $A_i \in \mathbb{F}^{n_i \times n_i}$ for $i = 1, 2$. Then, μ_A is the least common multiple of μ_{A_1} and μ_{A_2} . In particular, if μ_{A_1} and μ_{A_2} are coprime, then $\mu_A = \mu_{A_1}\mu_{A_2}$.

Proof. Since $0 = \mu_A(A) = \text{diag}[\mu_A(A_1), \mu_A(A_2)]$, it follows that $\mu_A(A_1) = 0$ and $\mu_A(A_2) = 0$. Therefore, Theorem 4.6.1 implies that μ_{A_1} and μ_{A_2} both divide μ_A . Consequently, the least common multiple q of μ_{A_1} and μ_{A_2} also divides μ_A . Since $q(A_1) = 0$ and $q(A_2) = 0$, it follows that $q(A) = 0$. Therefore, μ_A divides q . Hence, $q = \mu_A$. If, in addition, μ_{A_1} and μ_{A_2} are coprime, then $\mu_A = \mu_{A_1}\mu_{A_2}$. $\square$

Proposition 5.2.8. Let $p \in \mathbb{F}[s]$ be a monic polynomial of positive degree n , and let $p = p_1 \cdots p_r$, where $p_1, \dots, p_r \in \mathbb{F}[s]$ are monic and pairwise coprime polynomials. Then, the matrices $C(p)$ and $\text{diag}[C(p_1), \dots, C(p_r)]$ are similar.

Proof. Let $\hat{p}_2 = p_2 \cdots p_r$ and $\hat{C} \triangleq \text{diag}[C(p_1), C(\hat{p}_2)]$. Since p_1 and $\hat{p}_2$ are coprime, it follows from Lemma 5.2.7 that $\mu_{\hat{C}} = \mu_{C(p_1)}\mu_{C(\hat{p}_2)}$. Furthermore, $\chi_{\hat{C}} = \chi_{C(p_1)}\chi_{C(\hat{p}_2)} = \mu_{\hat{C}}$. Hence, Corollary 5.2.4 implies that $\hat{C}$ is similar to $C(\chi_{\hat{C}})$. However, $\chi_{\hat{C}} = p_1 \cdots p_r = p$, so that $\hat{C}$ is similar to $C(p)$. If $r > 2$, then the same argument can be used to decompose $C(\hat{p}_2)$ to show that $C(p)$ is similar to $\text{diag}[C(p_1), \dots, C(p_r)]$. $\square$

Proposition 5.2.8 can be used to decompose every companion block of a multicompanion form into smaller companion matrices. This procedure can be carried out for every companion block whose characteristic polynomial has coprime factors. For example, suppose that $A \in \mathbb{R}^{10 \times 10}$ has the similarity invariants $p_i(s) = 1$ for all $i = 1, \dots, 7$, $p_8(s) = (s+1)^2$, $p_9(s) = (s+1)^2(s+2)$, and $p_{10}(s) = (s+1)^2(s+2)(s^2+3)$, so that, by Theorem 5.2.3, the multicompanion form of A is $\text{diag}[C(p_8), C(p_9), C(p_{10})]$, where $C(p_8) \in \mathbb{R}^{2 \times 2}$, $C(p_9) \in \mathbb{R}^{3 \times 3}$, and $C(p_{10}) \in \mathbb{R}^{5 \times 5}$. According to Proposition 5.2.8, the companion matrices $C(p_9)$ and $C(p_{10})$ can be further decomposed. For example, $C(p_9)$ is similar to $\text{diag}[C(p_{9,1}), C(p_{9,2})]$, where $p_{9,1}(s) = (s+1)^2$ and $p_{9,2}(s) = s+2$ are coprime. Furthermore, $C(p_{10})$ is similar to four different diagonal matrices, three of which have two companion blocks while the fourth has three companion blocks. Since $p_8(s) = (s+1)^2$ does not have nonconstant coprime factors, however, it follows that the companion matrix $C(p_8)$ cannot be decomposed into smaller companion matrices.

The largest number of companion blocks achievable by similarity transformation is obtained by factoring every similarity invariant into *elementary divisors*, which are powers of irreducible polynomials that are nonconstant, monic, and pairwise coprime. In the above example, this factorization is given by $p_9(s) = p_{9,1}(s)p_{9,2}(s)$, where $p_{9,1}(s) = (s+1)^2$ and $p_{9,2}(s) = s+2$, and by $p_{10} = p_{10,1}p_{10,2}p_{10,3}$, where $p_{10,1}(s) = (s+1)^2$, $p_{10,2}(s) = s+2$, and $p_{10,3}(s) = s^2+3$. The elementary divisors of A are thus $(s+1)^2$, $(s+1)^2$, $s+2$, $(s+1)^2$, $s+2$, and s^2+3 , which yields six companion blocks. Viewing $A \in \mathbb{C}^{n \times n}$ we can further factor $p_{10,3}(s) = (s+j\sqrt{3})(s-j\sqrt{3})$, which yields a total of seven companion blocks. From Proposition 5.2.8 and Theorem 5.2.3 we obtain the *elementary multicompanion form*, which provides another canonical form for A .

Theorem 5.2.9. Let $A \in \mathbb{F}^{n \times n}$, and let $q_1^{l_1}, \dots, q_h^{l_h} \in \mathbb{F}[s]$ be the elementary divisors of A , where $l_1, \dots, l_h \in \mathbb{P}$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A = S \begin{bmatrix} C(q_1^{l_1}) & & 0 \\ & \ddots & \\ 0 & & C(q_h^{l_h}) \end{bmatrix} S^{-1}. \quad (5.2.9)$$

5.3 Hypercompanion Form and Jordan Form

In this section we present an alternative form of the companion blocks of the elementary multicompanion form (5.2.9). To do this we define the *hypercompanion*

matrix $\mathcal{H}_l(q)$ associated with the elementary divisor $q^l \in \mathbb{F}[s]$, where $l \in \mathbb{P}$, as follows. For $q(s) = s - \lambda \in \mathbb{C}[s]$, define the $l \times l$ Toeplitz hypercompanion matrix

$$\mathcal{H}_l(q) \triangleq \lambda I_l + N_l = \begin{bmatrix} \lambda & 1 & 0 & & & \\ 0 & \lambda & 1 & & & 0 \\ & & \ddots & \ddots & & \\ & & & \ddots & 1 & 0 \\ & 0 & & & \lambda & 1 \\ & & & & 0 & \lambda \end{bmatrix}, \quad (5.3.1)$$

while, for $q(s) = s^2 - \beta_1 s - \beta_0 \in \mathbb{R}[s]$, define the $2l \times 2l$ real, tridiagonal hypercompanion matrix

$$\mathcal{H}_l(q) \triangleq \begin{bmatrix} 0 & 1 & & & & \\ \beta_0 & \beta_1 & 1 & & & 0 \\ & 0 & 0 & 1 & & \\ & & \beta_0 & \beta_1 & 1 & \\ & & & \ddots & \ddots & \ddots \\ 0 & & & & \ddots & 0 & 1 \\ & & & & & \beta_0 & \beta_1 \end{bmatrix}. \quad (5.3.2)$$

The following result shows that the hypercompanion matrix $\mathcal{H}_l(q)$ is similar to the companion matrix $C(q^l)$ associated with the elementary divisor q^l of $\mathcal{H}_l(q)$.

Lemma 5.3.1. Let $l \in \mathbb{P}$, and let $q(s) = s - \lambda \in \mathbb{C}[s]$ or $q(s) = s^2 - \beta_1 s - \beta_0 \in \mathbb{R}[s]$. Then, q^l is the only elementary divisor of $\mathcal{H}_l(q)$, and $\mathcal{H}_l(q)$ is similar to $C(q^l)$.

Proof. Let k denote the order of $\mathcal{H}_l(q)$. Then, $\chi_{\mathcal{H}_l(q)} = q^l$ and $\det([sI - \mathcal{H}_l(q)]_{[k;1]}) = (-1)^{k-1}$. Hence, as in the proof of Proposition 5.2.1, it follows that $\chi_{\mathcal{H}_l(q)} = \mu_{\mathcal{H}_l(q)}$. Corollary 5.2.4 now implies that $\mathcal{H}_l(q)$ is similar to $C(q^l)$. $\square$

Proposition 5.2.8 and Lemma 5.3.1 yield the following canonical form, which is known as the *hypercompanion form*.

Theorem 5.3.2. Let $A \in \mathbb{F}^{n \times n}$, and let $q_1^{l_1}, \dots, q_h^{l_h} \in \mathbb{F}[s]$ be the elementary divisors of A , where $l_1, \dots, l_h \in \mathbb{P}$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A = S \begin{bmatrix} \mathcal{H}_{l_1}(q_1) & & 0 \\ & \ddots & \\ 0 & & \mathcal{H}_{l_h}(q_h) \end{bmatrix} S^{-1}. \quad (5.3.3)$$

Next, consider Theorem 5.3.2 with $\mathbb{F} = \mathbb{C}$. In this case, every elementary divisor $q_i^{l_i}$ is of the form $(s - \lambda_i)^{l_i}$, where $\lambda_i \in \mathbb{C}$. Furthermore, $S \in \mathbb{C}^{n \times n}$, and the hypercompanion form (5.3.3) is a block-diagonal matrix whose diagonally located blocks are of the form (5.3.1). The hypercompanion form (5.3.3) with every diagonally located block of the form (5.3.1) is the *Jordan form*, as given by the following

result.

Theorem 5.3.3. Let $A \in \mathbb{C}^{n \times n}$, and let $q_1^{l_1}, \dots, q_h^{l_h} \in \mathbb{C}[s]$ be the elementary divisors of A , where $l_1, \dots, l_h \in \mathbb{P}$ and each of the polynomials $q_1, \dots, q_h \in \mathbb{C}[s]$ has degree 1. Then, there exists a nonsingular matrix $S \in \mathbb{C}^{n \times n}$ such that

$$A = S \begin{bmatrix} \mathcal{H}_{l_1}(q_1) & & 0 \\ & \ddots & \\ 0 & & \mathcal{H}_{l_h}(q_h) \end{bmatrix} S^{-1}. \quad (5.3.4)$$

Corollary 5.3.4. Let $p \in \mathbb{F}[s]$, let $\lambda_1, \dots, \lambda_r$ denote the distinct roots of p , and, for $i = 1, \dots, r$, let $l_i \triangleq m_p(\lambda_i)$ and $p_i(s) \triangleq s - \lambda_i$. Then, $C(p)$ is similar to $\text{diag}[\mathcal{H}_{l_1}(p_1), \dots, \mathcal{H}_{l_r}(p_r)]$.

To illustrate the structure of the Jordan form, let $l_i = 3$ and $q_i(s) = s - \lambda_i$, where $\lambda_i \in \mathbb{C}$. Then, $\mathcal{H}_{l_i}(q_i)$ is the 3×3 matrix

$$\mathcal{H}_{l_i}(q_i) = \lambda_i I_3 + N_3 = \begin{bmatrix} \lambda_i & 1 & 0 \\ 0 & \lambda_i & 1 \\ 0 & 0 & \lambda_i \end{bmatrix} \quad (5.3.5)$$

so that $\text{mspec}[\mathcal{H}_{l_i}(q_i)] = \{\lambda_i, \lambda_i, \lambda_i\}_{\text{ms}}$. If $\mathcal{H}_{l_i}(q_i)$ is the only diagonally located block of the Jordan form associated with the eigenvalue λ_i , then the algebraic multiplicity of λ_i is equal to 3, while its geometric multiplicity is equal to 1.

Now, consider Theorem 5.3.2 with $\mathbb{F} = \mathbb{R}$. In this case, every elementary divisor $q_i^{l_i}$ is either of the form $(s - \lambda_i)^{l_i}$ or of the form $(s^2 - \beta_{1i}s - \beta_{0i})^{l_i}$, where $\beta_{0i}, \beta_{1i} \in \mathbb{R}$. Furthermore, $S \in \mathbb{R}^{n \times n}$, and the hypercompanion form (5.3.3) is a block-diagonal matrix whose diagonally located blocks are real matrices of the form (5.3.1) or (5.3.2). In this case, (5.3.3) is the *real hypercompanion form*.

Applying an additional real similarity transformation to each diagonally located block of the real hypercompanion form yields the *real Jordan form*. To do this, define the *real Jordan matrix* $\mathcal{J}_l(q)$ for $l \in \mathbb{P}$ as follows. For $q(s) = s - \lambda \in \mathbb{F}[s]$ define $\mathcal{J}_l(q) \triangleq \mathcal{H}_l(q)$, while, if $q(s) = s^2 - \beta_1 s - \beta_0 \in \mathbb{F}[s]$ is irreducible with a nonreal root $\lambda = \nu + j\omega$, then define the $2l \times 2l$ upper Hessenberg matrix

$$\mathcal{J}_l(q) \triangleq \begin{bmatrix} \nu & \omega & 1 & 0 & & & \\ -\omega & \nu & 0 & 1 & \ddots & & 0 \\ & & \nu & \omega & 1 & \ddots & \\ & & -\omega & \nu & 0 & \ddots & \ddots \\ & & & & \ddots & \ddots & 1 & 0 \\ & & & & & \ddots & 0 & 1 \\ & & & & & & & \nu & \omega \\ & & & & & & & -\omega & \nu \end{bmatrix}. \quad (5.3.6)$$

Theorem 5.3.5. Let $A \in \mathbb{R}^{n \times n}$, and let $q_1^{l_1}, \dots, q_h^{l_h} \in \mathbb{R}[s]$, where $l_1, \dots, l_h \in \mathbb{P}$ are the elementary divisors of A . Then, there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that

$$A = S \begin{bmatrix} \mathcal{J}_{l_1}(q_1) & & 0 \\ & \ddots & \\ 0 & & \mathcal{J}_{l_h}(q_h) \end{bmatrix} S^{-1}. \quad (5.3.7)$$

Proof. For the irreducible quadratic $q(s) = s^2 - \beta_1 s - \beta_0 \in \mathbb{R}[s]$ we show that $\mathcal{J}_l(q)$ and $\mathcal{H}_l(q)$ are similar. Writing $q(s) = (s - \lambda)(s - \bar{\lambda})$, it follows from Theorem 5.3.3 that $\mathcal{H}_l(q) \in \mathbb{R}^{2l \times 2l}$ is similar to $\text{diag}(\lambda I_l + N_l, \bar{\lambda} I_l + N_l)$. Next, by using a permutation similarity transformation, it follows that $\mathcal{H}_l(q)$ is similar to

$$\begin{bmatrix} \lambda & 0 & 1 & 0 & & & & & \\ 0 & \bar{\lambda} & 0 & 1 & 0 & & & & 0 \\ & 0 & \lambda & 0 & 1 & 0 & & & \\ & & 0 & \bar{\lambda} & 0 & 1 & & & \\ & & & \ddots & \ddots & \ddots & & & \\ & & & & \ddots & \ddots & 1 & 0 & \\ & & & & & \ddots & 0 & 1 & \\ & 0 & & & & & \lambda & 0 & \\ & & & & & & 0 & \bar{\lambda} \end{bmatrix},$$

Finally, applying the similarity transformation $S \triangleq \text{diag}(\hat{S}, \dots, \hat{S})$ to the above matrix, where $\hat{S} \triangleq \begin{bmatrix} -j & -j \\ 1 & -1 \end{bmatrix}$ and $\hat{S}^{-1} = \frac{1}{2} \begin{bmatrix} j & 1 \\ j & -1 \end{bmatrix}$, yields $\mathcal{J}_l(q)$. $\square$

Example 5.3.6. Let $A, B \in \mathbb{R}^{4 \times 4}$ and $C \in \mathbb{C}^{4 \times 4}$ be given by

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -16 & 0 & -8 & 0 \end{bmatrix},$$

$$B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ -4 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -4 & 0 \end{bmatrix},$$

and

$$C = \begin{bmatrix} 2j & 1 & 0 & 0 \\ 0 & 2j & 0 & 0 \\ 0 & 0 & -2j & 1 \\ 0 & 0 & 0 & -2j \end{bmatrix}.$$

Then, A is in companion form, B is in real hypercompanion form, and C is in Jordan form. Furthermore, A , B , and C are similar.

Example 5.3.7. Let $A, B \in \mathbb{R}^{6 \times 6}$ and $C \in \mathbb{C}^{6 \times 6}$ be given by

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ -27 & 54 & -63 & 44 & -21 & 6 \end{bmatrix},$$

$$B = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ -3 & 2 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & -3 & 2 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -3 & 2 \end{bmatrix},$$

and

$$C = \begin{bmatrix} 1+j\sqrt{2} & 1 & 0 & 0 & 0 & 0 \\ 0 & 1+j\sqrt{2} & 1 & 0 & 0 & 0 \\ 0 & 0 & 1+j\sqrt{2} & 0 & 0 & 0 \\ 0 & 0 & 0 & 1-j\sqrt{2} & 1 & 0 \\ 0 & 0 & 0 & 0 & 1-j\sqrt{2} & 1 \\ 0 & 0 & 0 & 0 & 0 & 1-j\sqrt{2} \end{bmatrix}.$$

Then, A is in companion form, B is in real hypercompanion form, and C is in Jordan form. Furthermore, A , B , and C are similar.

The next result shows that every matrix is similar to its transpose by means of a symmetric similarity transformation. This result, which improves Corollary 4.3.11, is due to Frobenius.

Corollary 5.3.8. Let $A \in \mathbb{F}^{n \times n}$. Then, there exists a symmetric, nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A = SA^T S^{-1}$.

Proof. It follows from Theorem 5.3.3 that there exists a nonsingular matrix $\hat{S} \in \mathbb{C}^{n \times n}$ such that $A = \hat{S}B\hat{S}^{-1}$, where $B = \text{diag}(B_1, \dots, B_r)$ is the Jordan form of A , and $B_i \in \mathbb{C}^{n_i \times n_i}$ for all $i = 1, \dots, r$. Now, define the symmetric nonsingular matrix $S \triangleq \hat{S}\tilde{I}\hat{S}^T$, where $\tilde{I} \triangleq \text{diag}(\hat{I}_{n_1}, \dots, \hat{I}_{n_r})$ is symmetric and involutory. Furthermore, note that $\hat{I}_{n_i}B_i\hat{I}_{n_i} = B_i^T$ for all $i = 1, \dots, r$ so that $\tilde{I}B\tilde{I} = B^T$, and thus $\tilde{I}B^T\tilde{I} = B$. Hence, it follows that

$$\begin{aligned} SA^T S^{-1} &= S\hat{S}^{-T}B^T\hat{S}^T S^{-1} = \hat{S}\tilde{I}\hat{S}^T\hat{S}^{-T}B^T\hat{S}^T\hat{S}^{-T}\tilde{I}\hat{S}^{-1} \\ &= \hat{S}\tilde{I}B^T\tilde{I}\hat{S}^{-1} = \hat{S}B\hat{S}^{-1} = A. \end{aligned}$$

If A is real, then a similar argument based on the real Jordan form shows that S can be chosen to be real. $\square$

An extension of Corollary 5.3.8 to the case in which A is normal is given by Fact 5.9.9.

Corollary 5.3.9. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist symmetric matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that S_2 is nonsingular and $A = S_1 S_2$.

Proof. From Corollary 5.3.8 it follows that there exists a symmetric, nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A = S A^T S^{-1}$. Now, let $S_1 \triangleq S A^T$ and $S_2 \triangleq S^{-1}$. Note that S_2 is symmetric and nonsingular. Furthermore, $S_1^T = A S = S A^T = S_1$, which shows that S_1 is symmetric. $\square$

Note that Corollary 5.3.8 follows from Corollary 5.3.9. If $A = S_1 S_2$, where S_1, S_2 are symmetric and S_2 is nonsingular, then $A = S_2^{-1} S_2 S_1 S_2 = S_2^{-1} A^T S_2$.

5.4 Schur Decomposition

The *Schur decomposition* uses a unitary similarity transformation to transform an arbitrary square matrix into an upper triangular matrix.

Theorem 5.4.1. Let $A \in \mathbb{C}^{n \times n}$. Then, there exist a unitary matrix $S \in \mathbb{C}^{n \times n}$ and an upper triangular matrix $B \in \mathbb{C}^{n \times n}$ such that

$$A = S B S^*. \quad (5.4.1)$$

Proof. Let $\lambda_1 \in \mathbb{C}$ be an eigenvalue of A with associated eigenvector $x \in \mathbb{C}^n$ chosen such that $x^* x = 1$. Furthermore, let $S_1 \triangleq \begin{bmatrix} x & \hat{S}_1 \end{bmatrix} \in \mathbb{C}^{n \times n}$ be unitary, where $\hat{S}_1 \in \mathbb{C}^{n \times (n-1)}$ satisfies $\hat{S}_1^* S_1 = I_{n-1}$ and $x^* \hat{S}_1 = 0_{1 \times (n-1)}$. Then, $S_1 e_1 = x$, and

$$\text{col}_1(S_1^{-1} A S_1) = S_1^{-1} A x = \lambda_1 S_1^{-1} x = \lambda_1 e_1.$$

Consequently,

$$A = S_1 \begin{bmatrix} \lambda_1 & C_1 \\ 0_{(n-1) \times 1} & A_1 \end{bmatrix} S_1^{-1},$$

where $C_1 \in \mathbb{C}^{1 \times (n-1)}$ and $A_1 \in \mathbb{C}^{(n-1) \times (n-1)}$. Next, let $S_{20} \in \mathbb{C}^{(n-1) \times (n-1)}$ be a unitary matrix such that

$$A_1 = S_{20} \begin{bmatrix} \lambda_2 & C_2 \\ 0_{(n-2) \times 1} & A_2 \end{bmatrix} S_{20}^{-1},$$

where $C_2 \in \mathbb{C}^{1 \times (n-2)}$ and $A_2 \in \mathbb{C}^{(n-2) \times (n-2)}$. Hence,

$$A = S_1 S_2 \begin{bmatrix} \lambda_1 & C_{11} & C_{12} \\ 0 & \lambda_2 & C_2 \\ 0 & 0 & A_2 \end{bmatrix} S_2^{-1} S_1,$$

where $C_1 = \begin{bmatrix} C_{11} & C_{12} \end{bmatrix}$, $C_{11} \in \mathbb{C}$, and $S_2 \triangleq \begin{bmatrix} 1 & 0 \\ 0 & S_{20} \end{bmatrix}$ is unitary. Proceeding in a similar manner yields (5.4.1) with $S \triangleq S_1 S_2 \cdots S_{n-1}$, where $S_1, \dots, S_{n-1} \in \mathbb{C}^{n \times n}$ are unitary. $\square$

It can be seen that the diagonal entries of B are the eigenvalues of A .

The *real Schur decomposition* uses a real orthogonal similarity transformation to transform a real matrix into an upper Hessenberg matrix with real 1×1 and 2×2 diagonally located blocks.

Corollary 5.4.2. Let $A \in \mathbb{R}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_r\}_{\text{ms}} \cup \{\nu_1 + j\omega_1, \nu_1 - j\omega_1, \dots, \nu_l + j\omega_l, \nu_l - j\omega_l\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_r \in \mathbb{R}$ and, for all $i = 1, \dots, l$, $\nu_i, \omega_i \in \mathbb{R}$ and $\omega_i \neq 0$. Then, there exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that

$$A = SBS^T, \quad (5.4.2)$$

where B is upper block triangular and the diagonally located blocks $B_1, \dots, B_r \in \mathbb{R}$ and $\hat{B}_1, \dots, \hat{B}_l \in \mathbb{R}^{2 \times 2}$ of B satisfy $B_i \triangleq [\lambda_i]$ for all $i = 1, \dots, r$ and $\text{spec}(\hat{B}_i) = \{\nu_i + j\omega_i, \nu_i - j\omega_i\}$ for all $i = 1, \dots, l$.

Proof. The proof is analogous to the proof of Theorem 5.3.5. See also [709, p. 82]. $\square$

Corollary 5.4.3. Let $A \in \mathbb{R}^{n \times n}$, and assume that the spectrum of A is real. Then, there exist an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ and an upper triangular matrix $B \in \mathbb{R}^{n \times n}$ such that

$$A = SBS^T. \quad (5.4.3)$$

The Schur decomposition reveals the structure of range-Hermitian matrices and thus, as a special case, normal matrices.

Corollary 5.4.4. Let $A \in \mathbb{F}^{n \times n}$, and define $r \triangleq \text{rank } A$. Then, A is range Hermitian if and only if there exist a unitary matrix $S \in \mathbb{F}^{n \times n}$ and a nonsingular matrix $B \in \mathbb{F}^{r \times r}$ such that

$$A = S \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} S^*. \quad (5.4.4)$$

In addition, A is normal if and only if there exist a unitary matrix $S \in \mathbb{C}^{n \times n}$ and a diagonal matrix $B \in \mathbb{C}^{r \times r}$ such that (5.4.4) is satisfied.

Proof. Suppose that A is range Hermitian, and let $A = S\hat{B}S^*$, where $\hat{B}$ is upper triangular and $S \in \mathbb{F}^{n \times n}$ is unitary. Assume that A is singular, and choose S such that $\hat{B}_{(j,j)} = \hat{B}_{(j+1,j+1)} = \dots = \hat{B}_{(n,n)} = 0$ and such that all other diagonal entries of $\hat{B}$ are nonzero. Thus, $\text{row}_n(\hat{B}) = 0$, which implies that $e_n \notin \mathcal{R}(\hat{B})$. Since A is range Hermitian, it follows that $\mathcal{R}(\hat{B}) = \mathcal{R}(\hat{B}^*)$ so that $e_n \notin \mathcal{R}(\hat{B}^*)$. Thus, $\text{col}_n(\hat{B}) = \text{row}_n(\hat{B}^*) = 0$. If, in addition, $\hat{B}_{(n-1,n-1)} = 0$, then $\text{col}_{n-1}(\hat{B}) = 0$. Repeating this argument shows that $\hat{B}$ has the form $\begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix}$, where $B \in \mathbb{F}^{r \times r}$ is nonsingular.

Now, suppose that A is normal, and let $A = S\hat{B}S^*$, where $\hat{B} \in \mathbb{C}^{n \times n}$ is upper triangular and $S \in \mathbb{C}^{n \times n}$ is unitary. Since A is normal, it follows that $AA^* = A^*A$, which implies that $\hat{B}\hat{B}^* = \hat{B}^*\hat{B}$. Since $\hat{B}$ is upper triangular, it follows that $(\hat{B}^*\hat{B})_{(1,1)} = \hat{B}_{(1,1)}\hat{B}_{(1,1)}$, whereas $(\hat{B}\hat{B}^*)_{(1,1)} = \text{row}_1(\hat{B})[\text{row}_1(\hat{B})]^* = \sum_{i=1}^n \hat{B}_{(1,i)}\hat{B}_{(1,i)}$. Since $(\hat{B}^*\hat{B})_{(1,1)} = (\hat{B}\hat{B}^*)_{(1,1)}$, it follows that $\hat{B}_{(1,i)} = 0$ for all $i = 2, \dots, n$. Continuing in a similar fashion row by row, it follows that $\hat{B}$ is

diagonal. $\square$

Corollary 5.4.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and define $r \triangleq \text{rank } A$. Then, there exist a unitary matrix $S \in \mathbb{F}^{n \times n}$ and a diagonal matrix $B \in \mathbb{R}^{r \times r}$ such that (5.4.4) is satisfied. In addition, A is positive semidefinite if and only if the diagonal entries of B are positive, and A is positive definite if and only if A is positive semidefinite and $r = n$.

Proof. Corollary 5.4.4 and x, xi of Proposition 4.4.5 imply that there exist a unitary matrix $S \in \mathbb{F}^{n \times n}$ and a diagonal matrix $B \in \mathbb{R}^{r \times r}$ such that (5.4.4) is satisfied. If A is positive semidefinite, then $x^*Ax \geq 0$ for all $x \in \mathbb{F}^n$. Choosing $x = Se_i$, it follows that $B_{(i,i)} = e_i^T S^* A S e_i \geq 0$ for all $i = 1, \dots, r$. If A is positive definite, then $r = n$ and $B_{(i,i)} > 0$ for all $i = 1, \dots, n$. $\square$

Proposition 5.4.6. Let $A \in \mathbb{F}^{n \times n}$ be Hermitian. Then, there exists a non-singular matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A = S \begin{bmatrix} -I_{\nu_-(A)} & 0 & 0 \\ 0 & 0_{\nu_0(A) \times \nu_0(A)} & 0 \\ 0 & 0 & I_{\nu_+(A)} \end{bmatrix} S^*. \quad (5.4.5)$$

Furthermore,

$$\text{rank } A = \nu_+(A) + \nu_-(A) \quad (5.4.6)$$

and

$$\text{def } A = \nu_0(A). \quad (5.4.7)$$

Proof. Since A is Hermitian, it follows from Corollary 5.4.5 that there exist a unitary matrix $\hat{S} \in \mathbb{F}^{n \times n}$ and a diagonal matrix $B \in \mathbb{R}^{n \times n}$ such that $A = \hat{S} B \hat{S}^*$. Choose S to order the diagonal entries of B such that $B = \text{diag}(B_1, 0, -B_2)$, where the diagonal matrices B_1, B_2 are both positive definite. Now, define $\hat{B} \triangleq \text{diag}(B_1, I, B_2)$. Then, $B = \hat{B}^{1/2} D \hat{B}^{1/2}$, where $D \triangleq \text{diag}(I_{\nu_-(A)}, 0_{\nu_0(A) \times \nu_0(A)}, -I_{\nu_+(A)})$. Hence, $A = \hat{S} \hat{B}^{1/2} D \hat{B}^{1/2} \hat{S}^*$. $\square$

The following result is *Sylvester's law of inertia*.

Corollary 5.4.7. Let $A, B \in \mathbb{F}^{n \times n}$ be Hermitian. Then, A and B are congruent if and only if $\text{In } A = \text{In } B$.

Proposition 4.5.4 shows that two or more eigenvectors associated with distinct eigenvalues of a normal matrix are mutually orthogonal. Thus, a normal matrix has at least as many mutually orthogonal eigenvectors as it has distinct eigenvalues. The next result, which is an immediate consequence of Corollary 5.4.4, shows that every $n \times n$ normal matrix actually has n mutually orthogonal eigenvectors. In fact, the converse is also true.

Corollary 5.4.8. Let $A \in \mathbb{C}^{n \times n}$. Then, A is normal if and only if A has n mutually orthogonal eigenvectors.

The following result concerns the *real normal form*.

Corollary 5.4.9. Let $A \in \mathbb{R}^{n \times n}$ be range symmetric. Then, there exist an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ and a nonsingular matrix $B \in \mathbb{R}^{r \times r}$, where $r \triangleq \text{rank } A$, such that

$$A = S \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} S^T. \quad (5.4.8)$$

In addition, assume that A is normal, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_r\}_{\text{ms}} \cup \{\nu_1 + j\omega_1, \nu_1 - j\omega_1, \dots, \nu_l + j\omega_l, \nu_l - j\omega_l\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_r \in \mathbb{R}$ and, for all $i = 1, \dots, l$, $\nu_i, \omega_i \in \mathbb{R}$ and $\omega_i \neq 0$. Then, there exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that

$$A = SBS^T, \quad (5.4.9)$$

where $B \triangleq \text{diag}(B_1, \dots, B_r, \hat{B}_1, \dots, \hat{B}_l)$, $B_i \triangleq [\lambda_i]$ for all $i = 1, \dots, r$, and $\hat{B}_i \triangleq \begin{bmatrix} \nu_i & \omega_i \\ -\omega_i & \nu_i \end{bmatrix}$ for all $i = 1, \dots, l$.

5.5 Eigenstructure Properties

Definition 5.5.1. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \mathbb{C}$. Then, the *index of λ with respect to A* , denoted by $\text{ind}_A(\lambda)$, is the smallest nonnegative integer k such that

$$\mathcal{R}[(\lambda I - A)^k] = \mathcal{R}[(\lambda I - A)^{k+1}]. \quad (5.5.1)$$

That is,

$$\text{ind}_A(\lambda) = \text{ind}(\lambda I - A). \quad (5.5.2)$$

Note that $\lambda \notin \text{spec}(A)$ if and only if $\text{ind}_A(\lambda) = 0$. Hence, $0 \notin \text{spec}(A)$ if and only if $\text{ind } A = \text{ind}_A(0) = 0$.

Proposition 5.5.2. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \mathbb{C}$. Then, $\text{ind}_A(\lambda)$ is the smallest nonnegative integer k such that

$$\text{rank}[(\lambda I - A)^k] = \text{rank}[(\lambda I - A)^{k+1}]. \quad (5.5.3)$$

Furthermore, $\text{ind } A$ is the smallest nonnegative integer k such that

$$\text{rank}(A^k) = \text{rank}(A^{k+1}). \quad (5.5.4)$$

Proof. Corollary 2.4.2 implies that $\mathcal{R}[(\lambda I - A)^k] \subseteq \mathcal{R}[(\lambda I - A)^{k+1}]$. Now, Lemma 2.3.4 implies that $\mathcal{R}[(\lambda I - A)^k] = \mathcal{R}[(\lambda I - A)^{k+1}]$ if and only if $\text{rank}[(\lambda I - A)^k] = \text{rank}[(\lambda I - A)^{k+1}]$. $\square$

Proposition 5.5.3. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \text{spec}(A)$. Then, the following statements hold:

- i) The order of the largest Jordan block of A associated with λ is $\text{ind}_A(\lambda)$.
- ii) The number of Jordan blocks of A associated with λ is $\text{gmult}_A(\lambda)$.
- iii) The number of linearly independent eigenvectors of A associated with λ is $\text{gmult}_A(\lambda)$.
- iv) $\text{ind}_A(\lambda) \leq \text{amult}_A(\lambda)$.

- v) $\text{gmult}_A(\lambda) \leq \text{amult}_A(\lambda)$.
- vi) $\text{ind}_A(\lambda) + \text{gmult}_A(\lambda) \leq \text{amult}_A(\lambda) + 1$.
- vii) $\text{ind}_A(\lambda) + \text{gmult}_A(\lambda) = \text{amult}_A(\lambda) + 1$ if and only if every block except possibly one block associated with λ is of order 1.

Definition 5.5.4. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \text{spec}(A)$. Then, the following terminology is defined:

- i) λ is *simple* if $\text{amult}_A(\lambda) = 1$.
- ii) A is *simple* if every eigenvalue of A is simple.
- iii) λ is *cyclic* (or *nonderogatory*) if $\text{gmult}_A(\lambda) = 1$.
- iv) A is *cyclic* (or *nonderogatory*) if every eigenvalue of A is cyclic.
- v) λ is *derogatory* if $\text{gmult}_A(\lambda) > 1$.
- vi) A is *derogatory* if A has at least one derogatory eigenvalue.
- vii) λ is *semisimple* if $\text{gmult}_A(\lambda) = \text{amult}_A(\lambda)$.
- viii) A is *semisimple* if every eigenvalue of A is semisimple.
- ix) λ is *defective* if $\text{gmult}_A(\lambda) < \text{amult}_A(\lambda)$.
- x) A is *defective* if A has at least one defective eigenvalue.
- xi) A is *diagonalizable over $\mathbb{C}$* if A is semisimple.
- xii) $A \in \mathbb{R}^{n \times n}$ is *diagonalizable over $\mathbb{R}$* if A is semisimple and every eigenvalue of A is real.

Proposition 5.5.5. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \text{spec}(A)$. Then, λ is simple if and only if λ is cyclic and semisimple.

Proposition 5.5.6. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \text{spec}(A)$. Then,

$$\text{def}[(\lambda I - A)^{\text{ind}_A(\lambda)}] = \text{amult}_A(\lambda). \quad (5.5.5)$$

Theorem 5.3.3 yields the following result, which shows that the subspaces $\mathcal{N}[(\lambda I - A)^k]$, where $\lambda \in \text{spec}(A)$ and $k = \text{ind}_A(\lambda)$, provide a decomposition of $\mathbb{F}^n$.

Proposition 5.5.7. Let $A \in \mathbb{F}^{n \times n}$, let $\text{spec}(A) = \{\lambda_1, \dots, \lambda_r\}$, and, for all $i = 1, \dots, r$, let $k_i \triangleq \text{ind}_A(\lambda_i)$. Then, the following statements hold:

- i) $\mathcal{N}[(\lambda_i I - A)^{k_i}] \cap \mathcal{N}[(\lambda_j I - A)^{k_j}] = \{0\}$ for all $i, j = 1, \dots, r$ such that $i \neq j$.
- ii) $\sum_{i=1}^r \mathcal{N}[(\lambda_i I - A)^{k_i}] = \mathbb{F}^n$.

Proposition 5.5.8. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \text{spec}(A)$. Then, the following statements are equivalent:

- i) λ is semisimple.

- ii)* $\text{def}(\lambda I - A) = \text{def}[(\lambda I - A)^2]$.
- iii)* $\mathcal{N}(\lambda I - A) = \mathcal{N}[(\lambda I - A)^2]$.
- iv)* $\text{ind}_A(\lambda) = 1$.

Proof. To prove that *i)* implies *ii)*, suppose that λ is semisimple so that $\text{gmult}_A(\lambda) = \text{amult}_A(\lambda)$, and thus $\text{def}(\lambda I - A) = \text{amult}_A(\lambda)$. Then, it follows from Proposition 5.5.6 that $\text{def}[(\lambda I - A)^k] = \text{amult}_A(\lambda)$, where $k \triangleq \text{ind}_A(\lambda)$. Therefore, it follows from Corollary 2.5.7 that $\text{amult}_A(\lambda) = \text{def}(\lambda I - A) \leq \text{def}[(\lambda I - A)^2] \leq \text{def}[(\lambda I - A)^k] = \text{amult}_A(\lambda)$, which implies that $\text{def}(\lambda I - A) = \text{def}[(\lambda I - A)^2]$.

To prove that *ii)* implies *iii)*, note that it follows from Corollary 2.5.7 that $\mathcal{N}(\lambda I - A) \subseteq \mathcal{N}[(\lambda I - A)^2]$. Since, by *ii)*, these subspaces have equal dimension, it follows from Lemma 2.3.4 that these subspaces are equal. Conversely, *iii)* implies *ii)*.

Finally, *iv)* is equivalent to the fact that every Jordan block of A associated with λ has order 1, which is equivalent to the fact that the geometric multiplicity of λ is equal to the algebraic multiplicity of λ , that is, that λ is semisimple. $\square$

Corollary 5.5.9. Let $A \in \mathbb{F}^{n \times n}$. Then, A is group invertible if and only if $\text{ind } A \leq 1$.

Proposition 5.5.10. Assume that $A, B \in \mathbb{F}^{n \times n}$ are similar. Then, the following statements hold:

- i)* $\text{mspec}(A) = \text{mspec}(B)$.
- ii)* For all $\lambda \in \text{spec}(A)$, $\text{gmult}_A(\lambda) = \text{gmult}_B(\lambda)$.

Proposition 5.5.11. Let $A \in \mathbb{F}^{n \times n}$. Then, A is semisimple if and only if A is similar to a normal matrix.

The following result is an extension of Corollary 5.3.9.

Proposition 5.5.12. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i)* A is semisimple, and $\text{spec}(A) \subset \mathbb{R}$.
- ii)* There exists a positive-definite matrix $S \in \mathbb{F}^{n \times n}$ such that $A = SA^*S^{-1}$.
- iii)* There exist a Hermitian matrix $S_1 \in \mathbb{F}^{n \times n}$ and a positive-definite matrix $S_2 \in \mathbb{F}^{n \times n}$ such that $A = S_1S_2$.

Proof. To prove that *i)* implies *ii)*, let $\hat{S} \in \mathbb{F}^{n \times n}$ be a nonsingular matrix such that $A = \hat{S}B\hat{S}^{-1}$, where $B \in \mathbb{R}^{n \times n}$ is diagonal. Then, $B = \hat{S}^{-1}A\hat{S} = \hat{S}^*A^*\hat{S}^{-*}$. Hence, $A = \hat{S}B\hat{S}^{-1} = \hat{S}(\hat{S}^*A^*\hat{S}^{-*})\hat{S}^{-1} = (\hat{S}\hat{S}^*)A^*(\hat{S}\hat{S}^*)^{-1} = SA^*S^{-1}$, where $S \triangleq \hat{S}\hat{S}^*$ is positive definite. To show that *ii)* implies *iii)*, note that $A = SA^*S^{-1} = S_1S_2$, where $S_1 \triangleq SA^*$ and $S_2 = S^{-1}$. Since $S_1^* = (SA^*)^* = AS^* = AS = SA^* = S_1$, it follows that S_1 is Hermitian. Furthermore, since S is positive definite, it follows

that S^{-1} , and hence S_2 , is also positive definite. Finally, to prove that *iii*) implies *i*), note that $A = S_1 S_2 = S_2^{-1/2} (S_2^{1/2} S_1 S_2^{1/2}) S_2^{1/2}$. Since $S_2^{1/2} S_1 S_2^{1/2}$ is Hermitian, it follows from Corollary 5.4.5 that $S_2^{1/2} S_1 S_2^{1/2}$ is unitarily similar to a real diagonal matrix. Consequently, A is semisimple and $\text{spec}(A) \subset \mathbb{R}$. $\square$

If a matrix is block triangular, then the following result shows that its eigenvalues and their algebraic multiplicity are determined by the diagonally located blocks. If, in addition, the matrix is block diagonal, then the geometric multiplicities of its eigenvalues are determined by the diagonally located blocks.

Proposition 5.5.13. Let $A \in \mathbb{F}^{n \times n}$, assume that A is partitioned as $A = \begin{bmatrix} A_{11} & \cdots & A_{1k} \\ \vdots & \ddots & \vdots \\ A_{k1} & \cdots & A_{kk} \end{bmatrix}$, where, for all $i, j = 1, \dots, k$, $A_{ij} \in \mathbb{F}^{n_i \times n_j}$, and let $\lambda \in \text{spec}(A)$. Then, the following statements hold:

i) If A_{ii} is the only nonzero block in the i th column of blocks, then

$$\text{amult}_{A_{ii}}(\lambda) \leq \text{amult}_A(\lambda). \quad (5.5.6)$$

ii) If A is upper block triangular or lower block triangular, then

$$\text{amult}_A(\lambda) = \sum_{i=1}^r \text{amult}_{A_{ii}}(\lambda) \quad (5.5.7)$$

and

$$\text{mspec}(A) = \bigcup_{i=1}^k \text{mspec}(A_{ii}). \quad (5.5.8)$$

Proposition 5.5.14. Let $A \in \mathbb{F}^{n \times n}$, assume that A is partitioned as $A = \begin{bmatrix} A_{11} & \cdots & A_{1k} \\ \vdots & \ddots & \vdots \\ A_{k1} & \cdots & A_{kk} \end{bmatrix}$, where, for all $i, j = 1, \dots, k$, $A_{ij} \in \mathbb{F}^{n_i \times n_j}$, and let $\lambda \in \text{spec}(A)$. Then, the following statements hold:

i) If A_{ii} is the only nonzero block in the i th column of blocks, then

$$\text{gmult}_{A_{ii}}(\lambda) \leq \text{gmult}_A(\lambda). \quad (5.5.9)$$

ii) If A is upper block triangular, then

$$\text{gmult}_{A_{11}}(\lambda) \leq \text{gmult}_A(\lambda). \quad (5.5.10)$$

iii) If A is lower block triangular, then

$$\text{gmult}_{A_{kk}}(\lambda) \leq \text{gmult}_A(\lambda). \quad (5.5.11)$$

iv) If A is block diagonal, then

$$\text{gmult}_A(\lambda) = \sum_{i=1}^r \text{gmult}_{A_{ii}}(\lambda). \quad (5.5.12)$$

Proposition 5.5.15. Let $A \in \mathbb{F}^{n \times n}$, let $\text{spec}(A) = \{\lambda_1, \dots, \lambda_r\}$, and let $k_i \triangleq \text{ind}_A(\lambda_i)$ for all $i = 1, \dots, r$. Then,

$$\mu_A(s) = \prod_{i=1}^r (s - \lambda_i)^{k_i} \quad (5.5.13)$$

and

$$\deg \mu_A = \sum_{i=1}^r k_i. \quad (5.5.14)$$

Furthermore, the following statements are equivalent:

- i) $\mu_A = \chi_A$.
- ii) A is cyclic.
- iii) For all $\lambda \in \text{spec}(A)$, the Jordan form of A contains exactly one block associated with λ .
- iv) A is similar to $C(\chi_A)$.

Proof. Let $A = SBS^{-1}$, where $B = \text{diag}(B_1, \dots, B_{n_h})$ denotes the Jordan form of A given by (5.3.4). Let $\lambda_i \in \text{spec}(A)$, and let B_j be a Jordan block associated with λ_i . Then, the order of B_j is less than or equal to k_i . Consequently, $(B_j - \lambda_i I)^{k_i} = 0$.

Next, let $p(s)$ denote the right-hand side of (5.5.13). Thus,

$$\begin{aligned} p(A) &= \prod_{i=1}^r (A - \lambda_i I)^{k_i} = S \left[\prod_{i=1}^r (B - \lambda_i I)^{k_i} \right] S^{-1} \\ &= S \text{diag} \left(\prod_{i=1}^r (B_1 - \lambda_i I)^{k_i}, \dots, \prod_{i=1}^r (B_{n_h} - \lambda_i I)^{k_i} \right) S^{-1} = 0. \end{aligned}$$

Therefore, it follows from Theorem 4.6.1 that μ_A divides p . Furthermore, note that, if k_i is replaced by $\hat{k}_i < k_i$, then $p(A) \neq 0$. Hence, p is the minimal polynomial of A . The equivalence of i) and ii) is now immediate, while the equivalence of ii) and iii) follows from Theorem 5.3.5. The equivalence of i) and iv) is given by Corollary 5.2.4. $\square$

Example 5.5.16. The standard nilpotent matrix N_n is in companion form, and thus is cyclic. In fact, N_n consists of a single Jordan block, and $\chi_{N_n}(s) = \mu_{N_n}(s) = s^n$.

Example 5.5.17. The matrix $\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$ is normal but is neither symmetric nor skew symmetric, while the matrix $\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$ is normal but is neither symmetric nor semisimple with real eigenvalues.

Example 5.5.18. The matrices $\begin{bmatrix} 1 & 0 \\ 2 & -1 \end{bmatrix}$ and $\begin{bmatrix} 1 & 1 \\ 0 & 2 \end{bmatrix}$ are diagonalizable over $\mathbb{R}$ but not normal, while the matrix $\begin{bmatrix} -1 & 1 \\ -2 & 1 \end{bmatrix}$ is diagonalizable but is neither normal nor diagonalizable over $\mathbb{R}$.

Example 5.5.19. The product of the Hermitian matrices $\begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$ and $\begin{bmatrix} 2 & 1 \\ 1 & -2 \end{bmatrix}$ has no real eigenvalues.

Example 5.5.20. The matrices $\begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$ and $\begin{bmatrix} 0 & 1 \\ -2 & 3 \end{bmatrix}$ are similar, whereas $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ and $\begin{bmatrix} 0 & 1 \\ -1 & 2 \end{bmatrix}$ have the same spectrum but are not similar.

Proposition 5.5.21. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) A is singular if and only if $0 \in \text{spec}(A)$.
- ii) A is group invertible if and only if either A is nonsingular or $0 \in \text{spec}(A)$ is semisimple.
- iii) A is Hermitian if and only if A is normal and $\text{spec}(A) \subset \mathbb{R}$.
- iv) A is skew Hermitian if and only if A is normal and $\text{spec}(A) \subset j\mathbb{R}$.
- v) A is positive semidefinite if and only if A is normal and $\text{spec}(A) \subset [0, \infty)$.
- vi) A is positive definite if and only if A is normal and $\text{spec}(A) \subset (0, \infty)$.
- vii) A is unitary if and only if A is normal and $\text{spec}(A) \subset \{\lambda \in \mathbb{C}: |\lambda| = 1\}$.
- viii) A is shifted unitary if and only if A is normal and

$$\text{spec}(A) \subset \{\lambda \in \mathbb{C}: |\lambda - \tfrac{1}{2}| = \tfrac{1}{2}\}.$$
- ix) A is involutory if and only if A is semisimple and $\text{spec}(A) \subseteq \{-1, 1\}$.
- x) A is skew involutory if and only if A is semisimple and $\text{spec}(A) \subseteq \{-j, j\}$.
- xi) A is idempotent if and only if A is semisimple and $\text{spec}(A) \subseteq \{0, 1\}$.
- xii) A is skew idempotent if and only if A is semisimple and $\text{spec}(A) \subseteq \{0, -1\}$.
- xiii) A is tripotent if and only if A is semisimple and $\text{spec}(A) \subseteq \{-1, 0, 1\}$.
- xiv) A is nilpotent if and only if $\text{spec}(A) = \{0\}$.
- xv) A is unipotent if and only if $\text{spec}(A) = \{1\}$.
- xvi) A is a projector if and only if A is normal and $\text{spec}(A) \subseteq \{0, 1\}$.
- xvii) A is a reflector if and only if A is normal and $\text{spec}(A) \subseteq \{-1, 1\}$.
- xviii) A is a skew reflector if and only if A is normal and $\text{spec}(A) \subseteq \{-j, j\}$.
- xi) A is an elementary projector if and only if A is normal and $\text{mspec}(A) = \{0, 1, \dots, 1\}_{\text{ms}}$.
- xx) A is an elementary reflector if and only if A is normal and $\text{mspec}(A) = \{-1, 1, \dots, 1\}_{\text{ms}}$.

If, furthermore, $A \in \mathbb{F}^{2n \times 2n}$, then the following statements hold:

- xxi) If A is Hamiltonian, then $\text{mspec}(A) = \text{mspec}(-A)$.
- xxii) If A is symplectic, then $\text{mspec}(A) = \text{mspec}(A^{-1})$.

The following result is a consequence of Proposition 5.5.12 and Proposition 5.5.21.

Corollary 5.5.22. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is either involutory, idempotent, skew idempotent, tripotent, a projector, or a reflector. Then, the following statements hold:

- i) There exists a positive-definite matrix $S \in \mathbb{F}^{n \times n}$ such that $A = SA^*S^{-1}$.
- ii) There exist a Hermitian matrix $S_1 \in \mathbb{F}^{n \times n}$ and a positive-definite matrix $S_2 \in \mathbb{F}^{n \times n}$ such that $A = S_1S_2$.

5.6 Singular Value Decomposition

The third matrix decomposition that we consider is the *singular value decomposition*. Unlike the Jordan and Schur decompositions, the singular value decomposition applies to matrices that are not necessarily square. Let $A \in \mathbb{F}^{n \times m}$, where $A \neq 0$, and consider the positive-semidefinite matrices $AA^* \in \mathbb{F}^{n \times n}$ and $A^*A \in \mathbb{F}^{m \times m}$. It follows from Proposition 4.4.10 that AA^* and A^*A have the same nonzero eigenvalues with the same algebraic multiplicities. Since AA^* and A^*A are positive semidefinite, it follows that they have the same *positive* eigenvalues with the same algebraic multiplicities. Furthermore, since AA^* is Hermitian, it follows that the number of positive eigenvalues of AA^* (or A^*A) counting algebraic multiplicity is equal to the rank of AA^* (or A^*A). Since $\text{rank } A = \text{rank } AA^* = \text{rank } A^*A$, it thus follows that AA^* and A^*A both have r positive eigenvalues, where $r \triangleq \text{rank } A$.

Definition 5.6.1. Let $A \in \mathbb{F}^{n \times m}$. Then, the *singular values* of A are the $\min\{n, m\}$ nonnegative numbers $\sigma_1(A), \dots, \sigma_{\min\{n, m\}}(A)$, where, for all $i = 1, \dots, \min\{n, m\}$,

$$\sigma_i(A) \triangleq \lambda_i^{1/2}(AA^*) = \lambda_i^{1/2}(A^*A). \quad (5.6.1)$$

Hence,

$$\sigma_1(A) \geq \dots \geq \sigma_{\min\{n, m\}}(A) \geq 0. \quad (5.6.2)$$

Let $A \in \mathbb{F}^{n \times m}$, and define $r \triangleq \text{rank } A$. If $1 \leq r < \min\{n, m\}$, then

$$\sigma_1(A) \geq \dots \geq \sigma_r(A) > \sigma_{r+1}(A) = \dots = \sigma_{\min\{n, m\}}(A) = 0, \quad (5.6.3)$$

whereas, if $r = \min\{m, n\}$, then

$$\sigma_1(A) \geq \dots \geq \sigma_r(A) = \sigma_{\min\{n, m\}}(A) > 0. \quad (5.6.4)$$

For convenience, define

$$\sigma_{\max}(A) \triangleq \sigma_1(A) \quad (5.6.5)$$

and, if $n = m$,

$$\sigma_{\min}(A) \triangleq \sigma_n(A). \quad (5.6.6)$$

If $n \neq m$, then $\sigma_{\min}(A)$ is not defined. By convention, we define

$$\sigma_{\max}(0_{n \times m}) = \sigma_{\min}(0_{n \times n}) = 0, \quad (5.6.7)$$

and, for all $i = 1, \dots, \min\{n, m\}$,

$$\sigma_i(A) = \sigma_i(A^*) = \sigma_i(\overline{A}) = \sigma_i(A^T). \quad (5.6.8)$$

Now, suppose that $n = m$. If A is Hermitian, then, for all $i = 1, \dots, n$,

$$\sigma_i(A) = |\lambda_i(A)|, \quad (5.6.9)$$

while, if A is positive semidefinite, then, for all $i = 1, \dots, n$,

$$\sigma_i(A) = \lambda_i(A). \quad (5.6.10)$$

Proposition 5.6.2. Let $A \in \mathbb{F}^{n \times m}$. If $n \leq m$, then the following statements are equivalent:

- i) $\text{rank } A = n$.
- ii) $\sigma_n(A) > 0$.

If $m \leq n$, then the following statements are equivalent:

- iii) $\text{rank } A = m$.
- iv) $\sigma_m(A) > 0$.

If $n = m$, then the following statements are equivalent:

- v) A is nonsingular.
- vi) $\sigma_{\min}(A) > 0$.

Proposition 5.6.3. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) Assume that A and B are normal. Then, A and B are unitarily similar if and only if $\text{mspec}(A) = \text{mspec}(B)$.
- ii) Assume that A and B are projectors. Then, A and B are unitarily similar if and only if $\text{rank } A = \text{rank } B$.
- iii) Assume that A and B are (projectors, reflectors). Then, A and B are unitarily similar if and only if $\text{tr } A = \text{tr } B$.
- iv) Assume that A and B are semisimple. Then, A and B are similar if and only if $\text{mspec}(A) = \text{mspec}(B)$.
- v) Assume that A and B are (involutory, skew involutory, idempotent). Then, A and B are similar if and only if $\text{tr } A = \text{tr } B$.
- vi) Assume that A and B are idempotent. Then, A and B are similar if and only if $\text{rank } A = \text{rank } B$.
- vii) Assume that A and B are tripotent. Then, A and B are similar if and only if $\text{rank } A = \text{rank } B$ and $\text{tr } A = \text{tr } B$.

We now state the singular value decomposition.

Theorem 5.6.4. Let $A \in \mathbb{F}^{n \times m}$, assume that A is nonzero, let $r \triangleq \text{rank } A$, and define $B \triangleq \text{diag}[\sigma_1(A), \dots, \sigma_r(A)]$. Then, there exist unitary matrices $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$ such that

$$A = S_1 \begin{bmatrix} B & 0_{r \times (m-r)} \\ 0_{(n-r) \times r} & 0_{(n-r) \times (m-r)} \end{bmatrix} S_2. \quad (5.6.11)$$

Furthermore, each column of S_1 is an eigenvector of AA^* , while each column of S_2^* is an eigenvector of A^*A .

Proof. For convenience, assume that $r < \min\{n, m\}$, since otherwise the zero matrices become empty matrices. By Corollary 5.4.5 there exists a unitary matrix $U \in \mathbb{F}^{n \times n}$ such that

$$AA^* = U \begin{bmatrix} B^2 & 0 \\ 0 & 0 \end{bmatrix} U^*.$$

Partition $U = \begin{bmatrix} U_1 & U_2 \end{bmatrix}$, where $U_1 \in \mathbb{F}^{n \times r}$ and $U_2 \in \mathbb{F}^{n \times (n-r)}$. Since $U^*U = I_n$, it follows that $U_1^*U_1 = I_r$ and $U_1^*U = \begin{bmatrix} I_r & 0_{r \times (n-r)} \end{bmatrix}$. Now, define $V_1 \triangleq A^*U_1B^{-1} \in \mathbb{F}^{m \times r}$, and note that

$$V_1^*V_1 = B^{-1}U_1^*AA^*U_1B^{-1} = B^{-1}U_1^*U \begin{bmatrix} B^2 & 0 \\ 0 & 0 \end{bmatrix} U^*U_1B^{-1} = I_r.$$

Next, note that, since $U_2^*U = \begin{bmatrix} 0_{(n-r) \times r} & I_{n-r} \end{bmatrix}$, it follows that

$$U_2^*AA^* = \begin{bmatrix} 0 & I \end{bmatrix} \begin{bmatrix} B^2 & 0 \\ 0 & 0 \end{bmatrix} U^* = 0.$$

However, since $\mathcal{R}(A) = \mathcal{R}(AA^*)$, it follows that $U_2^*A = 0$. Finally, let $V_2 \in \mathbb{F}^{m \times (m-r)}$ be such that $V \triangleq \begin{bmatrix} V_1 & V_2 \end{bmatrix} \in \mathbb{F}^{m \times m}$ is unitary. Hence, we have

$$\begin{aligned} U \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} V^* &= \begin{bmatrix} U_1 & U_2 \end{bmatrix} \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} V_1^* \\ V_2^* \end{bmatrix} = U_1BV_1^* = U_1BB^{-1}U_1^*A \\ &= U_1U_1^*A = (U_1U_1^* + U_2U_2^*)A = UU^*A = A, \end{aligned}$$

which yields (5.6.11) with $S_1 = U$ and $S_2 = V^*$. $\square$

An immediate corollary of the singular value decomposition is the *polar decomposition*.

Corollary 5.6.5. Let $A \in \mathbb{F}^{n \times n}$. Then, there exists a positive-semidefinite matrix $M \in \mathbb{F}^{n \times n}$ and a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A = MS. \quad (5.6.12)$$

Proof. It follows from the singular value decomposition that there exist unitary matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ and a diagonal positive-definite matrix $B \in \mathbb{F}^{r \times r}$, where $r \triangleq \text{rank } A$, such that $A = S_1 \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} S_2$. Hence,

$$A = S_1 \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} S_1^* S_1 S_2 = MS,$$

where $M \triangleq S_1 \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} S_1^*$ is positive semidefinite and $S \triangleq S_1 S_2$ is unitary. $\square$

Proposition 5.6.6. Let $A \in \mathbb{F}^{n \times m}$, let $r \triangleq \text{rank } A$, and define the Hermitian matrix $\mathcal{A} \triangleq \begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$. Then, $\text{In } \mathcal{A} = \begin{bmatrix} r & 0 & r \end{bmatrix}^T$, and the $2r$ nonzero eigenvalues of $\mathcal{A}$ are the r positive singular values of A and their negatives.

Proof. Since $\chi_{\mathcal{A}}(s) = \det(s^2I - A^*A)$, it follows that

$$\text{mspec}(\mathcal{A}) \setminus \{0, \dots, 0\}_{\text{ms}} = \{\sigma_1(A), -\sigma_1(A), \dots, \sigma_r(A), -\sigma_r(A)\}_{\text{ms}}. \quad \square$$

5.7 Pencils and the Kronecker Canonical Form

Let $A, B \in \mathbb{F}^{n \times m}$, and define the polynomial matrix $P_{A,B} \in \mathbb{F}^{n \times m}[s]$, called a *pencil*, by

$$P_{A,B}(s) \triangleq sB - A.$$

The pencil $P_{A,B}$ is *regular* if $\text{rank } P_{A,B} = \min\{n, m\}$ (see Definition 4.2.4). Otherwise, $P_{A,B}$ is *singular*.

Let $A, B \in \mathbb{F}^{n \times m}$. Since $P_{A,B} \in \mathbb{F}^{n \times m}$ we define the *generalized spectrum* of $P_{A,B}$ by

$$\text{spec}(A, B) \triangleq \text{Szeros}(P_{A,B}) \quad (5.7.1)$$

and the *generalized multispectrum* of $P_{A,B}$ by

$$\text{mspec}(A, B) \triangleq \text{mSzeros}(P_{A,B}). \quad (5.7.2)$$

Furthermore, the elements of $\text{spec}(A, B)$ are the *generalized eigenvalues* of $P_{A,B}$.

The structure of a pencil is illuminated by the following result known as the *Kronecker canonical form*.

Theorem 5.7.1. Let $A, B \in \mathbb{C}^{n \times m}$. Then, there exist nonsingular matrices $S_1 \in \mathbb{C}^{n \times n}$ and $S_2 \in \mathbb{C}^{m \times m}$ such that, for all $s \in \mathbb{C}$,

$$P_{A,B}(s) = S_1 \text{diag}(sI_{r_1} - A_1, sB_2 - I_{r_2}, [sI_{k_1} - N_{k_1} - e_{k_1}], \dots, [sI_{k_p} - N_{k_p} - e_{k_p}], \\ [sI_{l_1} - N_{l_1} - e_{l_1}]^T, \dots, [sI_{l_q} - N_{l_q} - e_{l_q}]^T, 0_{t \times u}) S_2, \quad (5.7.3)$$

where $A_1 \in \mathbb{C}^{r_1 \times r_1}$ is in Jordan form, $B_2 \in \mathbb{R}^{r_2 \times r_2}$ is nilpotent and in Jordan form, $k_1, \dots, k_p, l_1, \dots, l_q$ are positive integers, and $[sI_l - N_l - e_l] \in \mathbb{C}^{l \times (l+1)}$. Furthermore,

$$\text{rank } P_{A,B} = r_1 + r_2 + \sum_{i=1}^p k_i + \sum_{i=1}^q l_i. \quad (5.7.4)$$

Proof. See [65, Chapter 2], [541, Chapter XII], [787, pp. 395–398], [866], [872, pp. 128, 129], and [1230, Chapter VI]. $\square$

In Theorem 5.7.1, note that

$$n = r_1 + r_2 + \sum_{i=1}^p k_i + \sum_{i=1}^q l_i + q + t \quad (5.7.5)$$

and

$$m = r_1 + r_2 + \sum_{i=1}^p k_i + \sum_{i=1}^q l_i + p + u. \quad (5.7.6)$$

Proposition 5.7.2. Let $A, B \in \mathbb{C}^{n \times m}$, and consider the notation of Theorem 5.7.1. Then, $P_{A,B}$ is regular if and only if $t = u = 0$ and either $p = 0$ or $q = 0$.

Let $A, B \in \mathbb{F}^{n \times m}$, and let $\lambda \in \mathbb{C}$. Then,

$$\text{rank } P_{A,B}(\lambda) = \text{rank}(\lambda I - A_1) + r_2 + \sum_{i=1}^p k_i + \sum_{i=1}^q l_i. \quad (5.7.7)$$

Note that λ is a generalized eigenvalue of $P_{A,B}$ if and only if $\text{rank } P_{A,B}(\lambda) < \text{rank } P_{A,B}$. Consequently, λ is a generalized eigenvalue of $P_{A,B}$ if and only if λ is an eigenvalue of A_1 , that is,

$$\text{spec}(A, B) = \text{spec}(A_1) \quad (5.7.8)$$

and

$$\text{mspec}(A, B) = \text{mspec}(A_1). \quad (5.7.9)$$

The *generalized algebraic multiplicity* $\text{amult}_{A,B}(\lambda)$ of $\lambda \in \text{spec}(A, B)$ is defined by

$$\text{amult}_{A,B}(\lambda) \triangleq \text{amult}_{A_1}(\lambda). \quad (5.7.10)$$

It can be seen that, for $\lambda \in \text{spec}(A, B)$,

$$\text{gmult}_{A_1}(\lambda) \triangleq \text{rank } P_{A,B} - \text{rank } P_{A,B}(\lambda).$$

The *generalized geometric multiplicity* $\text{gmult}_{A,B}(\lambda)$ of $\lambda \in \text{spec}(A, B)$ is defined by

$$\text{gmult}_{A,B}(\lambda) \triangleq \text{gmult}_{A_1}(\lambda). \quad (5.7.11)$$

Now, assume that $A, B \in \mathbb{F}^{n \times n}$, that is, A and B are square, which, from (5.7.5) and (5.7.6), is equivalent to $q + t = p + u$. Then, the *characteristic polynomial* $\chi_{A,B} \in \mathbb{F}[s]$ of (A, B) is defined by

$$\chi_{A,B}(s) \triangleq \det P_{A,B}(s) = \det(sB - A).$$

Proposition 5.7.3. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) $P_{A,B}$ is singular if and only if $\chi_{A,B} = 0$.
- ii) $P_{A,B}$ is singular if and only if $\deg \chi_{A,B} = -\infty$.
- iii) $P_{A,B}$ is regular if and only if $\chi_{A,B}$ is not the zero polynomial.
- iv) $P_{A,B}$ is regular if and only if $0 \leq \deg \chi_{A,B} \leq n$.
- v) If $P_{A,B}$ is regular, then $\text{mult}_{\chi_{A,B}}(0) = n - \deg \chi_{B,A}$.
- vi) $\deg \chi_{A,B} = n$ if and only if B is nonsingular.
- vii) If B is nonsingular, then $\chi_{A,B} = \chi_{B^{-1}A}$, $\text{spec}(A, B) = \text{spec}(B^{-1}A)$, and $\text{mspec}(A, B) = \text{mspec}(B^{-1}A)$.
- viii) $\text{roots}(\chi_{A,B}) = \text{spec}(A, B)$.
- ix) $\text{mroots}(\chi_{A,B}) = \text{mspec}(A, B)$.

- x)* If A or B is nonsingular, then $P_{A,B}$ is regular.
- xi)* If all of the generalized eigenvalues of (A, B) are real, then $P_{A,B}$ is regular.
- xii)* If $P_{A,B}$ is regular, then $\mathcal{N}(A) \cap \mathcal{N}(B) = \{0\}$.
- xiii)* If $P_{A,B}$ is regular, then there exist nonsingular matrices $S_1, S_2 \in \mathbb{C}^{n \times n}$ such that, for all $s \in \mathbb{C}$,

$$P_{A,B}(s) = S_1 \left(s \begin{bmatrix} I_r & 0 \\ 0 & B_2 \end{bmatrix} - \begin{bmatrix} A_1 & 0 \\ 0 & I_{n-r} \end{bmatrix} \right) S_2,$$

where $r \triangleq \deg \chi_{A,B}$, $A_1 \in \mathbb{C}^{r \times r}$ is in Jordan form, and $B_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ is nilpotent and in Jordan form. Furthermore,

$$\begin{aligned} \chi_{A,B} &= \chi_{A_1}, \\ \text{roots}(\chi_{A,B}) &= \text{spec}(A_1), \end{aligned}$$

and

$$\text{mroots}(\chi_{A,B}) = \text{mspec}(A_1).$$

Proof. See [872, p. 128] and [1230, Chapter VI]. □

Statement *xiii)* is the *Weierstrass canonical form* for a square, regular pencil.

Proposition 5.7.4. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and assume that B is Hermitian. Then, the following statements hold:

- i)* $P_{A,B}$ is regular.
- ii)* There exists $\alpha \in \mathbb{F}$ such that $A + \alpha B$ is nonsingular.
- iii)* $\mathcal{N}(A) \cap \mathcal{N}(B) = \{0\}$.
- iv)* $\mathcal{N}(\begin{bmatrix} A \\ B \end{bmatrix}) = \{0\}$.
- v)* There exists nonzero $\alpha \in \mathbb{F}$ such that $\mathcal{N}(A) \cap \mathcal{N}(B + \alpha A) = \{0\}$.
- vi)* For all nonzero $\alpha \in \mathbb{F}$, $\mathcal{N}(A) \cap \mathcal{N}(B + \alpha A) = \{0\}$.
- vii)* All generalized eigenvalues of (A, B) are real.

If, in addition, B is positive semidefinite, then the following statement is equivalent to *i)–vii)*:

- viii)* There exists $\beta > 0$ such that $\beta B < A$.

Proof. The results *i)* $\implies$ *ii)* and *ii)* $\implies$ *iii)* are immediate. Next, Fact 2.10.10 and Fact 2.11.3 imply that *iii)*, *iv)*, *v)*, and *vi)* are equivalent. Next, to prove *iii)* $\implies$ *vii)*, let $\lambda \in \mathbb{C}$ be a generalized eigenvalue of (A, B) . Since $\lambda = 0$ is real, suppose $\lambda \neq 0$. Since $\det(\lambda B - A) = 0$, let nonzero $\theta \in \mathbb{C}^n$ satisfy $(\lambda B - A)\theta = 0$, and thus it follows that $\theta^* A \theta = \lambda \theta^* B \theta$. Furthermore, note that $\theta^* A \theta$ and $\theta^* B \theta$ are real. Now, suppose $\theta \in \mathcal{N}(A)$. Then, it follows from $(\lambda B - A)\theta = 0$ that $\theta \in \mathcal{N}(B)$, which contradicts $\mathcal{N}(A) \cap \mathcal{N}(B) = \{0\}$. Hence, $\theta \notin \mathcal{N}(A)$, and thus $\theta^* A \theta > 0$ and, consequently, $\theta^* B \theta \neq 0$. Hence, it follows that $\lambda = \theta^* A \theta / \theta^* B \theta$, and thus λ is real. Hence, all generalized eigenvalues of (A, B) are real.

Next, to prove $vii) \implies i)$, let $\lambda \in \mathbb{C} \setminus \mathbb{R}$ so that λ is not a generalized eigenvalue of (A, B) . Consequently, $\chi_{A,B}(s)$ is not the zero polynomial, and thus (A, B) is regular.

Next, to prove $i)-vii) \implies viii)$, let $\theta \in \mathbb{R}^n$ be nonzero, and note that $\mathcal{N}(A) \cap \mathcal{N}(B) = \{0\}$ implies that either $A\theta \neq 0$ or $B\theta \neq 0$. Hence, either $\theta^T A \theta > 0$ or $\theta^T B \theta > 0$. Thus, $\theta^T (A + B) \theta > 0$, which implies $A + B > 0$ and hence $-B < A$.

Finally, to prove $viii) \implies i)-vii)$, let $\beta \in \mathbb{R}$ be such that $\beta B < A$, so that $\beta \theta^T B \theta < \theta^T A \theta$ for all nonzero $\theta \in \mathbb{R}^n$. Next, suppose $\hat{\theta} \in \mathcal{N}(A) \cap \mathcal{N}(B)$ is nonzero. Hence, $A\hat{\theta} = 0$ and $B\hat{\theta} = 0$. Consequently, $\hat{\theta}^T B \hat{\theta} = 0$ and $\hat{\theta}^T A \hat{\theta} = 0$, which contradicts $\beta \hat{\theta}^T B \hat{\theta} < \hat{\theta}^T A \hat{\theta}$. Thus, $\mathcal{N}(A) \cap \mathcal{N}(B) = \{0\}$. $\square$

5.8 Facts on the Inertia

Fact 5.8.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then,

$$\text{rank } A = \text{sig } A = \text{tr } A$$

and

$$\text{In } A = \begin{bmatrix} 0 & & \\ & n - \text{tr } A & \\ & & \text{tr } A \end{bmatrix}.$$

Fact 5.8.2. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is involutory. Then,

$$\text{rank } A = n,$$

$$\text{sig } A = \text{tr } A,$$

and

$$\text{In } A = \begin{bmatrix} \frac{1}{2}(n - \text{tr } A) & & \\ & 0 & \\ & & \frac{1}{2}(n + \text{tr } A) \end{bmatrix}.$$

Fact 5.8.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is tripotent. Then,

$$\text{rank } A = \text{tr } A^2,$$

$$\text{sig } A = \text{tr } A,$$

and

$$\text{In } A = \begin{bmatrix} \frac{1}{2}(\text{tr } A^2 - \text{tr } A) & & \\ & n - \text{tr } A^2 & \\ & & \frac{1}{2}(\text{tr } A^2 + \text{tr } A) \end{bmatrix}.$$

Fact 5.8.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is either skew Hermitian, skew involutory, or nilpotent. Then,

$$\text{sig } A = \nu_-(A) = \nu_+(A) = 0$$

and

$$\text{In } A = \begin{bmatrix} 0 \\ n \\ 0 \end{bmatrix}.$$

Fact 5.8.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is group invertible, and assume that $\text{spec}(A) \cap j\mathbb{R} \subseteq \{0\}$. Then,

$$\text{rank } A = \nu_-(A) + \nu_+(A)$$

and

$$\text{def } A = \nu_0(A) = \text{amult}_A(0).$$

Fact 5.8.6. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then,

$$\text{rank } A = \nu_-(A) + \nu_+(A)$$

and

$$\text{In } A = \begin{bmatrix} \nu_-(A) \\ \nu_0(A) \\ \nu_+(A) \end{bmatrix} = \begin{bmatrix} \frac{1}{2}(\text{rank } A - \text{sig } A) \\ n - \text{rank } A \\ \frac{1}{2}(\text{rank } A + \text{sig } A) \end{bmatrix}.$$

Fact 5.8.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, $\text{In } A = \text{In } B$ if and only if $\text{rank } A = \text{rank } B$ and $\text{sig } A = \text{sig } B$.

Fact 5.8.8. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and let A_0 be a principal submatrix of A . Then,

$$\nu_-(A_0) \leq \nu_-(A)$$

and

$$\nu_+(A_0) \leq \nu_+(A).$$

(Proof: See [770].)

Fact 5.8.9. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then,

$$\text{rank } A = \text{sig } A = \nu_+(A)$$

and

$$\text{In } A = \begin{bmatrix} 0 \\ \text{def } A \\ \text{rank } A \end{bmatrix}.$$

Fact 5.8.10. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then,

$$\text{In } A = \begin{bmatrix} 0 \\ \text{def } A \\ \text{rank } A \end{bmatrix}.$$

If, in addition, A is positive definite, then

$$\text{In } A = \begin{bmatrix} 0 \\ 0 \\ n \end{bmatrix}.$$

Fact 5.8.11. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is an elementary projector.
- ii) A is a projector, and $\text{tr } A = n - 1$.
- iii) A is a projector, and $\text{In } A = \begin{bmatrix} 0 & \\ & 1 \\ & & n-1 \end{bmatrix}$.

Furthermore, the following statements are equivalent:

- iv) A is an elementary reflector.
- v) A is a reflector, and $\text{tr } A = n - 2$.
- vi) A is a reflector, and $\text{In } A = \begin{bmatrix} 1 & \\ & 0 \\ & & n-1 \end{bmatrix}$.

(Proof: See Proposition 5.5.21.)

Fact 5.8.12. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $A + A^*$ is positive definite.
- ii) For all Hermitian matrices $B \in \mathbb{F}^{n \times n}$, $\text{In } B = \text{In } AB$.

(Proof: See [280].)

Fact 5.8.13. Let $A, B \in \mathbb{F}^{n \times n}$, assume that AB and B are Hermitian, and assume that $\text{spec}(A) \cap [0, \infty) = \emptyset$. Then,

$$\text{In}(-AB) = \text{In } B.$$

(Proof: See [280].)

Fact 5.8.14. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian and nonsingular, and assume that $\text{spec}(AB) \cap [0, \infty) = \emptyset$. Then,

$$\nu_+(A) + \nu_+(B) = n.$$

(Proof: Use Fact 5.8.13. See [280].) (Remark: Weaker versions of this result are given in [761, 1036].)

Fact 5.8.15. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and let $S \in \mathbb{F}^{m \times n}$. Then,

$$\nu_-(SAS^*) \leq \nu_-(A)$$

and

$$\nu_+(SAS^*) \leq \nu_+(A).$$

Furthermore, consider the following conditions:

- i) $\text{rank } S = n$.
- ii) $\text{rank } SAS^* = \text{rank } A$.
- iii) $\nu_-(SAS^*) = \nu_-(A)$ and $\nu_+(SAS^*) = \nu_+(A)$.

Then, i) $\implies$ ii) $\iff$ iii). (Proof: See [447, pp. 430, 431] and [508, p. 194].)

Fact 5.8.16. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and let $S \in \mathbb{F}^{m \times n}$. Then,

$$\nu_-(SAS^*) + \nu_+(SAS^*) = \text{rank } SAS^* \leq \min\{\text{rank } A, \text{rank } S\},$$

$$\nu_-(A) + \text{rank } S - n \leq \nu_-(SAS^*) \leq \nu_-(A),$$

$$\nu_+(A) + \text{rank } S - n \leq \nu_+(SAS^*) \leq \nu_+(A).$$

(Proof: See [1060].)

Fact 5.8.17. Let $A, S \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and assume that S is nonsingular. Then, there exist $\alpha_1, \dots, \alpha_n \in [\lambda_{\min}(SS^*), \lambda_{\max}(SS^*)]$ such that, for all $i = 1, \dots, n$,

$$\lambda_i(SAS^*) = \alpha_i \lambda_i(A).$$

(Proof: See [1439].) (Remark: This result, which is due to Ostrowski, is a quantitative version of Sylvester's law of inertia given by Corollary 5.4.7.)

Fact 5.8.18. Let $A, S \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and assume that S is nonsingular. Then, the following statements are equivalent:

- i) $\text{In}(SAS^*) = \text{In } A$.
- ii) $\text{rank}(SAS^*) = \text{rank } A$.
- iii) $\mathcal{R}(A) \cap \mathcal{N}(A) = \{0\}$.

(Proof: See [109].)

Fact 5.8.19. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and assume that A is positive definite and C is negative definite. Then,

$$\text{In} \begin{bmatrix} A & B & 0 \\ B^* & C & 0 \\ 0 & 0 & 0_{l \times l} \end{bmatrix} = \begin{bmatrix} n \\ m \\ l \end{bmatrix}.$$

(Proof: The result follows from Fact 5.8.6. See [770].)

Fact 5.8.20. Let $A \in \mathbb{R}^{n \times m}$. Then,

$$\begin{aligned} \text{In} \begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix} &= \text{In} \begin{bmatrix} AA^* & 0 \\ 0 & -A^*A \end{bmatrix} \\ &= \text{In} \begin{bmatrix} AA^+ & 0 \\ 0 & -A^+A \end{bmatrix} \\ &= \begin{bmatrix} \text{rank } A \\ n + m - 2\text{rank } A \\ \text{rank } A \end{bmatrix}. \end{aligned}$$

(Proof: See [447, pp. 432, 434].)

Fact 5.8.21. Let $A \in \mathbb{C}^{n \times n}$, assume that A is Hermitian, and let $B \in \mathbb{C}^{n \times m}$. Then,

$$\text{In} \begin{bmatrix} A & B \\ B^* & 0 \end{bmatrix} \geq \begin{bmatrix} \text{rank } B \\ n - \text{rank } B \\ \text{rank } B \end{bmatrix}.$$

Furthermore, if $\mathcal{R}(A) \subseteq \mathcal{R}(B)$, then

$$\text{In} \begin{bmatrix} A & B \\ B^* & 0 \end{bmatrix} = \begin{bmatrix} \text{rank } B \\ n + m - 2\text{rank } B \\ \text{rank } B \end{bmatrix}.$$

Finally, if $\text{rank } B = n$, Then,

$$\text{In} \begin{bmatrix} A & B \\ B^* & 0 \end{bmatrix} = \begin{bmatrix} n \\ m - n \\ n \end{bmatrix}.$$

(Proof: See [447, pp. 433, 434] or [945].) (Remark: Extensions are given in [945].)
(Remark: See Fact 8.15.27.)

Fact 5.8.22. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ and a skew-Hermitian matrix $B \in \mathbb{F}^{n \times n}$ such that

$$A = S \left(\begin{bmatrix} I_{\nu_-(A+A^*)} & 0 & 0 \\ 0 & 0_{\nu_0(A+A^*) \times \nu_0(A+A^*)} & 0 \\ 0 & 0 & -I_{\nu_+(A+A^*)} \end{bmatrix} + B \right) S^*.$$

(Proof: Write $A = \frac{1}{2}(A+A^*) + \frac{1}{2}(A-A^*)$, and apply Proposition 5.4.6 to $\frac{1}{2}(A+A^*)$.)

5.9 Facts on Matrix Transformations for One Matrix

Fact 5.9.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{spec}(A) = \{1\}$. Then, A^k is similar to A for all $k \geq 1$.

Fact 5.9.2. Let $A \in \mathbb{F}^{n \times n}$, and assume there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $S^{-1}AS$ is upper triangular. Then, for all $r = 1, \dots, n$, $\mathcal{R}(S \begin{bmatrix} I_r \\ 0 \end{bmatrix})$ is an invariant subspace of A . (Remark: Analogous results hold for lower triangular matrices and block-triangular matrices.)

Fact 5.9.3. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist unique matrices $B, C \in \mathbb{F}^{n \times n}$ such that the following properties are satisfied:

- i) B is diagonalizable over $\mathbb{F}$.
- ii) C is nilpotent.
- iii) $A = B + C$.
- iv) $BC = CB$.

Furthermore, $\text{mspec}(A) = \text{mspec}(B)$. (Proof: See [691, p. 112] or [727, p. 74]. Existence follows from the real Jordan form. The last statement follows from Fact 5.17.4.) (Remark: This result is the *S-N decomposition* or the *Jordan-Chevalley*

decomposition.)

Fact 5.9.4. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is similar to a skew-Hermitian matrix.
- ii) A is semisimple, and $\text{spec}(A) \subset j\mathbb{R}$.

(Remark: See Fact 11.18.12.)

Fact 5.9.5. Let $A \in \mathbb{F}^{n \times n}$, and let $r \triangleq \text{rank } A$. Then, A is group invertible if and only if there exist a nonsingular matrix $B \in \mathbb{F}^{r \times r}$ and a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A = S \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} S^{-1}.$$

Fact 5.9.6. Let $A \in \mathbb{F}^{n \times n}$, and let $r \triangleq \text{rank } A$. Then, A is range Hermitian if and only if there exist a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ and a nonsingular matrix $B \in \mathbb{F}^{r \times r}$ such that

$$A = S \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} S^*.$$

(Remark: S need not be unitary for sufficiency. See Corollary 5.4.4.) (Proof: Use the QR decomposition Fact 5.15.8 to let $S \triangleq \hat{S}R$, where $\hat{S}$ is unitary and R is upper triangular. See [1277].)

Fact 5.9.7. Let $A \in \mathbb{F}^{n \times n}$. Then, there exists an involutory matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A^T = SAS^T.$$

(Remark: Note A^T rather than A^* .) (Proof: See [420] and [577].)

Fact 5.9.8. Let $A \in \mathbb{F}^{n \times n}$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A = SA^*S^{-1}$ if and only if there exist Hermitian matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that $A = S_1S_2$. (Proof: See [1490, pp. 215, 216].) (Remark: See Proposition 5.5.12.)

Fact 5.9.9. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is normal. Then, there exists a symmetric, nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A^T = SAS^{-1}$$

and such that $S^{-1} = \overline{S}$. (Proof: For $\mathbb{F} = \mathbb{C}$, let $A = UBU^*$, where U is unitary and B is diagonal. Then, $A^T = SA\overline{S} = SAS^{-1}$, where $S \triangleq \overline{U}U^{-1}$. For $\mathbb{F} = \mathbb{R}$, use the real normal form and let $S \triangleq U\tilde{I}U^T$, where U is orthogonal and $\tilde{I} \triangleq \text{diag}(\tilde{I}, \dots, \tilde{I})$.) (Remark: See Corollary 5.3.8.)

Fact 5.9.10. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is normal. Then, there exists a reflector $S \in \mathbb{R}^{n \times n}$ such that

$$A^T = SAS^{-1}.$$

Consequently, A and A^T are orthogonally similar. Finally, if A is skew symmetric, then A and $-A$ are orthogonally similar. (Proof: Specialize Fact 5.9.9 to the case

$\mathbb{F} = \mathbb{R}$.)

Fact 5.9.11. Let $A \in \mathbb{F}^{n \times n}$. Then, there exists a reverse-symmetric, nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A^T = SAS^{-1}$. (Proof: The result follows from Corollary 5.3.8. See [882].)

Fact 5.9.12. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist reverse-symmetric matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that S_2 is nonsingular and $A = S_1 S_2$. (Proof: The result follows from Corollary 5.3.9. See [882].)

Fact 5.9.13. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is not of the form aI , where $a \in \mathbb{R}$. Then, A is similar to a matrix with diagonal entries $0, \dots, 0, \operatorname{tr} A$. (Proof: See [1098, p. 77].) (Remark: This result is due to Gibson.)

Fact 5.9.14. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is not zero. Then, A is similar to a matrix whose diagonal entries are all nonzero. (Proof: See [1098, p. 79].) (Remark: This result is due to Marcus and Purves.)

Fact 5.9.15. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is symmetric. Then, there exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that $-1 \notin \operatorname{spec}(S)$ and SAS^T is diagonal. (Proof: See [1098, p. 101].) (Remark: This result is due to Hsu.)

Fact 5.9.16. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is symmetric. Then, there exist a diagonal matrix $B \in \mathbb{R}^{n \times n}$ and a skew-symmetric matrix $C \in \mathbb{R}^{n \times n}$ such that

$$A = [2(I + C)^{-1} - I]B[2(I + C)^{-1} - I]^T.$$

(Proof: Use Fact 5.9.15. See [1098, p. 101].)

Fact 5.9.17. Let $A \in \mathbb{F}^{n \times n}$. Then, there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that S^*AS has equal diagonal entries. (Proof: See [488] or [1098, p. 78], or use Fact 5.9.18.) (Remark: The diagonal entries are equal to $(\operatorname{tr} A)/n$.) (Remark: This result is due to Parker. See [535].)

Fact 5.9.18. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $\operatorname{tr} A = 0$.
- ii) There exist matrices $B, C \in \mathbb{F}^{n \times n}$ such that $A = [B, C]$.
- iii) A is unitarily similar to a matrix whose diagonal entries are zero.

(Proof: See [13, 535, 799, 814] or [626, p. 146].) (Remark: This result is *Shoda's theorem*.) (Remark: See Fact 5.9.19.)

Fact 5.9.19. Let $R \in \mathbb{F}^{n \times n}$, and assume that R is Hermitian. Then, the following statements are equivalent:

- i) $\operatorname{tr} R < 0$.
- ii) R is unitarily similar to a matrix all of whose diagonal entries are negative.
- iii) There exists an asymptotically stable matrix $A \in \mathbb{F}^{n \times n}$ such that $R = A + A^*$.

(Proof: See [120].) (Remark: See Fact 5.9.18.)

Fact 5.9.20. Let $A \in \mathbb{F}^{n \times n}$. Then, AA^* and A^*A are unitarily similar.

Fact 5.9.21. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, A and A^* are unitarily similar. (Proof: The result follows from Fact 5.9.27 and the fact that $\begin{bmatrix} 1 & a \\ 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 1 & 0 \\ a & 0 \end{bmatrix}$ are unitarily similar. See [419].)

Fact 5.9.22. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is symmetric. Then, there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A = SBS^T,$$

where

$$B \triangleq \text{diag}[\sigma_1(A), \dots, \sigma_n(A)].$$

(Proof: See [709, p. 207].) (Remark: A is symmetric, complex, and T-congruent to B .)

Fact 5.9.23. Let $A \in \mathbb{F}^{n \times n}$. Then, $\begin{bmatrix} A & 0 \\ 0 & -A \end{bmatrix}$ and $\begin{bmatrix} 0 & A \\ A & 0 \end{bmatrix}$ are unitarily similar. (Proof: Use the unitary transformation $\frac{1}{\sqrt{2}} \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix}$.)

Fact 5.9.24. Let $n \in \mathbb{P}$. Then,

$$\hat{I}_n = \begin{cases} S \begin{bmatrix} -I_{n/2} & 0 \\ 0 & -I_{n/2} \end{bmatrix} S^T, & n \text{ even,} \\ S \begin{bmatrix} -I_{n/2} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & I_{n/2} \end{bmatrix} S^T, & n \text{ odd,} \end{cases}$$

where

$$S \triangleq \begin{cases} \frac{1}{\sqrt{2}} \begin{bmatrix} I_{n/2} & -\hat{I}_{n/2} \\ \hat{I}_{n/2} & I_{n/2} \end{bmatrix}, & n \text{ even,} \\ \frac{1}{\sqrt{2}} \begin{bmatrix} I_{n/2} & 0 & -\hat{I}_{n/2} \\ 0 & \sqrt{2} & 0 \\ \hat{I}_{n/2} & 0 & I_{n/2} \end{bmatrix}, & n \text{ odd.} \end{cases}$$

Therefore,

$$\text{mspec}(\hat{I}_n) = \begin{cases} \{-1, 1, \dots, -1, 1\}_{\text{ms}}, & n \text{ even,} \\ \{1, -1, 1, \dots, -1, 1\}_{\text{ms}}, & n \text{ odd.} \end{cases}$$

(Remark: For even n , Fact 3.19.3 shows that $\hat{I}_n$ is Hamiltonian, and thus, by Fact 4.9.21, $\text{mspec}(\hat{I}_n) = -\text{mspec}(I_n)$.) (Remark: See [1410].)

Fact 5.9.25. Let $n \in \mathbb{P}$. Then,

$$J_{2n} = S \begin{bmatrix} jI_n & 0 \\ 0 & -jI_n \end{bmatrix} S^*,$$

where

$$S \triangleq \frac{1}{\sqrt{2}} \begin{bmatrix} I & -I \\ jI & -jI \end{bmatrix}.$$

Hence,

$$\text{mspec}(J_{2n}) = \{j, -j, \dots, j, -j\}_{\text{ms}}$$

and

$$\det J_{2n} = 1.$$

(Proof: See Fact 2.19.3.) (Remark: Fact 3.19.3 shows that J_{2n} is Hamiltonian, and thus, by Fact 4.9.21, $\text{mspec}(J_{2n}) = -\text{mspec}(J_{2n})$.)

Fact 5.9.26. Let $A \in \mathbb{F}^{n \times n}$, assume that A is idempotent, and let $r \triangleq \text{rank } A$. Then, there exists a matrix $B \in \mathbb{F}^{r \times (n-r)}$ and a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that

$$A = S \begin{bmatrix} I_r & B \\ 0 & 0_{(n-r) \times (n-r)} \end{bmatrix} S^*.$$

(Proof: See [536, p. 46].)

Fact 5.9.27. Let $A \in \mathbb{F}^{n \times n}$, assume that A is idempotent, and let $r \triangleq \text{rank } A$. Then, there exist a unitary matrix $S \in \mathbb{F}^{n \times n}$ and positive numbers $a_1, \dots, a_k$ such that

$$A = S \text{diag} \left(\begin{bmatrix} 1 & a_1 \\ 0 & 0 \end{bmatrix}, \dots, \begin{bmatrix} 1 & a_k \\ 0 & 0 \end{bmatrix}, I_{r-k}, 0_{(n-r-k) \times (n-r-k)} \right) S^*.$$

(Proof: See [419].) (Remark: This result provides a canonical form for idempotent matrices under unitary similarity. See also [537].) (Remark: See Fact 5.9.21.)

Fact 5.9.28. Let $A \in \mathbb{F}^{n \times m}$, assume that A is nonzero, let $r \triangleq \text{rank } A$, define $B \triangleq \text{diag}[\sigma_1(A), \dots, \sigma_r(A)]$, and let $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$ be unitary matrices such that

$$A = S_1 \begin{bmatrix} B & 0_{r \times (m-r)} \\ 0_{(n-r) \times r} & 0_{(n-r) \times (m-r)} \end{bmatrix} S_2.$$

Then, there exist $K \in \mathbb{F}^{r \times r}$ and $L \in \mathbb{F}^{r \times (m-r)}$ such that

$$KK^* + LL^* = I_r$$

and

$$A = S_1 \begin{bmatrix} BK & BL \\ 0_{(n-r) \times r} & 0_{(n-r) \times (m-r)} \end{bmatrix} S_1^*.$$

(Proof: See [115, 651].) (Remark: See Fact 6.3.15 and Fact 6.6.15.)

Fact 5.9.29. Let $A \in \mathbb{F}^{n \times n}$, assume that A is unitary, and partition A as

$$A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix},$$

where $A_{11} \in \mathbb{F}^{m \times k}$, $A_{12} \in \mathbb{F}^{m \times q}$, $A_{21} \in \mathbb{F}^{p \times k}$, $A_{22} \in \mathbb{F}^{p \times q}$, and $m + p = k + q = n$. Then, there exist unitary matrices $U, V \in \mathbb{F}^{n \times n}$ and $l, r \geq 0$ such that

$$A = U \begin{bmatrix} I_r & 0 & 0 & 0 & 0 & 0 \\ 0 & \Gamma & 0 & 0 & \Sigma & 0 \\ 0 & 0 & 0 & 0 & 0 & I_{m-r-l} \\ 0 & 0 & 0 & I_{q-m+r} & 0 & 0 \\ 0 & \Sigma & 0 & 0 & -\Gamma & 0 \\ 0 & 0 & I_{k-r-l} & 0 & 0 & 0 \end{bmatrix} V,$$

where $\Gamma, \Sigma \in \mathbb{R}^{l \times l}$ are diagonal and satisfy

$$0 < \Gamma_{(l,l)} \leq \cdots \leq \Gamma_{(1,1)} < 1, \quad (5.9.1)$$

$$0 < \Sigma_{(1,1)} \leq \cdots \leq \Sigma_{(l,l)} < 1, \quad (5.9.2)$$

and

$$\Gamma^2 + \Sigma^2 = I_m.$$

(Proof: See [536, p. 12] and [1230, p. 37].) (Remark: This result is the *CS decomposition*. See [1059, 1061]. The entries $\Sigma_{(i,i)}$ and $\Gamma_{(i,i)}$ can be interpreted as sines and cosines, respectively, of the principal angles between a pair of subspaces $\mathcal{S}_1 = \mathcal{R}(X_1)$ and $\mathcal{S}_2 = \mathcal{R}(Y_1)$ such that $[X_1 \ X_2]$ and $[Y_1 \ Y_2]$ are unitary and $A = [X_1 \ X_2]^* [Y_1 \ Y_2]$; see [536, pp. 25–29], [1230, pp. 40–43], and Fact 2.9.19. Principal angles can also be defined recursively; see [536, p. 25] and [537].)

Fact 5.9.30. Let $A \in \mathbb{F}^{n \times n}$, and let $r \triangleq \text{rank } A$. Then, there exist $S_1 \in \mathbb{F}^{n \times r}$, $B \in \mathbb{R}^{r \times r}$, and $S_2 \in \mathbb{F}^{n \times r}$, such that S_1 is left inner, S_2 is right inner, B is upper triangular, $I \circ B = \alpha I$, where $\alpha \triangleq \prod_{i=1}^r \sigma_i(A)$, and

$$A = S_1 B S_2.$$

(Proof: See [757].) (Remark: Note that B is real.) (Remark: This result is the *geometric mean decomposition*.)

Fact 5.9.31. Let $A \in \mathbb{C}^{n \times n}$. Then, there exists a matrix $B \in \mathbb{R}^{n \times n}$ such that $A\bar{A}$ and B^2 are similar. (Proof: See [415].)

5.10 Facts on Matrix Transformations for Two or More Matrices

Fact 5.10.1. Let $q(s) \triangleq s^2 - \beta_1 s - \beta_0 \in \mathbb{R}[s]$ be irreducible, and let $\lambda = \nu + j\omega$ denote a root of q so that $\beta_1 = 2\nu$ and $\beta_0 = -(\nu^2 + \omega^2)$. Then,

$$\mathcal{H}_1(q) = \begin{bmatrix} 0 & 1 \\ \beta_0 & \beta_1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ \nu & \omega \end{bmatrix} \begin{bmatrix} \nu & \omega \\ -\omega & \nu \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -\nu/\omega & 1/\omega \end{bmatrix} = S \mathcal{J}_1(q) S^{-1}.$$

The transformation matrix $S = \begin{bmatrix} 1 & 0 \\ \nu & \omega \end{bmatrix}$ is not unique; an alternative choice is $S = \begin{bmatrix} \omega & \nu \\ 0 & \nu^2 + \omega^2 \end{bmatrix}$. Similarly,

$$\mathcal{H}_2(q) = \begin{bmatrix} 0 & 1 & 0 & 0 \\ \beta_0 & \beta_1 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & \beta_0 & \beta_1 \end{bmatrix} = S \begin{bmatrix} \nu & \omega & 1 & 0 \\ -\omega & \nu & 0 & 1 \\ 0 & 0 & \nu & \omega \\ 0 & 0 & -\omega & \nu \end{bmatrix} S^{-1} = S \mathcal{J}_2(q) S^{-1},$$

where

$$S \triangleq \begin{bmatrix} \omega & \nu & \omega & \nu \\ 0 & \nu^2 + \omega^2 & \omega & \nu^2 + \omega^2 + \nu \\ 0 & 0 & -2\omega\nu & 2\omega^2 \\ 0 & 0 & -2\omega(\nu^2 + \omega^2) & 0 \end{bmatrix}.$$

Fact 5.10.2. Let $q(s) \triangleq s^2 - 2\nu s + \nu^2 + \omega^2 \in \mathbb{R}[s]$ with roots $\lambda = \nu + j\omega$ and $\bar{\lambda} = \nu - j\omega$. Then,

$$\mathcal{H}_1(q) = \begin{bmatrix} \nu & \omega \\ -\omega & \nu \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 1 \\ j & -j \end{bmatrix} \begin{bmatrix} \lambda & 0 \\ 0 & \bar{\lambda} \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -j \\ 1 & j \end{bmatrix}$$

and

$$\mathcal{H}_2(q) = \begin{bmatrix} \nu & \omega & 1 & 0 \\ -\omega & \nu & 0 & 1 \\ 0 & 0 & \nu & \omega \\ 0 & 0 & -\omega & \nu \end{bmatrix} = S \begin{bmatrix} \lambda & 1 & 0 & 0 \\ 0 & \lambda & 0 & 0 \\ 0 & 0 & \bar{\lambda} & 1 \\ 0 & 0 & 0 & \bar{\lambda} \end{bmatrix} S^{-1},$$

where

$$S \triangleq \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 0 & 1 & 0 \\ j & 0 & -j & 0 \\ 0 & 1 & 0 & 1 \\ 0 & j & 0 & -j \end{bmatrix}, \quad S^{-1} = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & -j & 0 & 0 \\ 0 & 0 & 1 & -j \\ 1 & j & 0 & 0 \\ 0 & 0 & 1 & j \end{bmatrix}.$$

Fact 5.10.3. Left equivalence, right equivalence, biequivalence, unitary left equivalence, unitary right equivalence, and unitary biequivalence are equivalence relations on $\mathbb{F}^{n \times m}$. Similarity, congruence, and unitary similarity are equivalence relations on $\mathbb{F}^{n \times n}$.

Fact 5.10.4. Let $A, B \in \mathbb{F}^{n \times m}$. Then, A and B are in the same equivalence class of $\mathbb{F}^{n \times m}$ induced by biequivalent transformations if and only if A and B are biequivalent to $\begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$. Now, let $n = m$. Then, A and B are in the same equivalence class of $\mathbb{F}^{n \times n}$ induced by similarity transformations if and only if A and B have the same Jordan form.

Fact 5.10.5. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are similar. Then, A is semisimple if and only if B is.

Fact 5.10.6. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is normal. Then, A is unitarily similar to its Jordan form.

Fact 5.10.7. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are normal, and assume that A and B are similar. Then, A and B are unitarily similar. (Proof: Since A and B are similar, it follows that $\text{mspec}(A) = \text{mspec}(B)$. Since A and B are

normal, it follows that they are unitarily similar to the same diagonal matrix. See Fact 5.10.6. See [627, p. 104].) (Remark: See [541, p. 8] for related results.)

Fact 5.10.8. Let $A, B \in \mathbb{F}^{n \times n}$, and let $r \triangleq 2n^2$. Then, the following statements are equivalent:

- i) A and B are unitarily similar.
- ii) For all $k_1, \dots, k_r, l_1, \dots, l_r \in \mathbb{N}$ such that $\sum_{i,j=1}^r (k_i + l_j) \leq r$, it follows that

$$\operatorname{tr} A^{k_1} A^{l_1*} \dots A^{k_r} A^{l_r*} = \operatorname{tr} B^{k_1} B^{l_1*} \dots B^{k_r} B^{l_r*}.$$

(Proof: See [1076].) (Remark: See [790, pp. 71, 72] and [220, 1190].) (Remark: The number of distinct tuples of positive integers whose sum is a positive integer k is 2^{k-1} . The number of expressions in ii) is thus $\sum_{k=1}^{2n^2} 2^{k-1} = 4^{n^2} - 1$. Because of properties of the trace function, the number of distinct expressions is less than this number. Furthermore, in special cases, the number of expressions that need to be checked is significantly less than the number of distinct expressions. In the case $n = 2$, it suffices to check three equalities, specifically, $\operatorname{tr} A = \operatorname{tr} B$, $\operatorname{tr} A^2 = \operatorname{tr} B^2$, and $\operatorname{tr} A^*A = \operatorname{tr} B^*B$. In the case $n = 3$, it suffices to check 7 equalities. See [220, 1190].)

Fact 5.10.9. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are idempotent, assume that $\operatorname{sprad}(A - B) < 1$, and define

$$S \triangleq (AB + A_{\perp}B_{\perp})[I - (A - B)^2]^{-1/2}.$$

Then, the following statements hold:

- i) S is nonsingular.
- ii) If $A = B$, then $S = I$.
- iii) $S^{-1} = (BA + B_{\perp}A_{\perp})[I - (B - A)^2]^{-1/2}$.
- iv) A and B are similar. In fact, $A = SBS^{-1}$.
- v) If A and B are projectors, then S is unitary and A and B are unitarily similar.

(Proof: See [690, p. 412].) (Remark: $[I - (A - B)^2]^{-1/2}$ is defined by ix) of Fact 10.11.24.)

Fact 5.10.10. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then, the following statements are equivalent:

- i) A and B are unitarily similar.
- ii) $\operatorname{tr} A = \operatorname{tr} B$ and, for all $i = 1, \dots, \lfloor n/2 \rfloor$, $\operatorname{tr} (AA^*)^i = \operatorname{tr} (BB^*)^i$.
- iii) $\chi_{AA^*} = \chi_{BB^*}$.

(Proof: The result follows from Fact 5.9.27. See [419].)

Fact 5.10.11. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that either A or B is nonsingular. Then, AB and BA are similar. (Proof: If A is nonsingular, then $AB = A(BA)A^{-1}$.)

Fact 5.10.12. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, AB and BA are unitarily similar. (Remark: This result is due to Dixmier. See [1114].)

Fact 5.10.13. Let $A \in \mathbb{F}^{n \times n}$. Then, A is idempotent if and only if there exists an orthogonal matrix $B \in \mathbb{F}^{n \times n}$ such that A and B are similar.

Fact 5.10.14. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are idempotent, and assume that $A + B - I$ is nonsingular. Then, A and B are similar. In particular,

$$A = (A + B - I)^{-1}B(A + B - I).$$

Fact 5.10.15. Let $A_1, \dots, A_r \in \mathbb{F}^{n \times n}$, and assume that $A_i A_j = A_j A_i$ for all $i, j = 1, \dots, r$. Then,

$$\dim \operatorname{span} \left\{ \prod_{i=1}^r A_i^{n_i} : 0 \leq n_i \leq n-1, i = 1, \dots, r \right\} \leq \frac{1}{4}n^2 + 1.$$

(Remark: This result gives a bound on the dimension of a commutative subalgebra.)
(Remark: This result is due to Schur. See [859].)

Fact 5.10.16. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $AB = BA$. Then,

$$\dim \operatorname{span} \{ A^i B^j : 0 \leq i \leq n-1, 0 \leq j \leq n-1 \} \leq n.$$

(Remark: This result gives a bound on the dimension of a commutative subalgebra generated by two matrices.) (Remark: This result is due to Gerstenhaber. See [150, 859].)

Fact 5.10.17. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are normal, nonsingular, and congruent. Then, $\operatorname{In} A = \operatorname{In} B$. (Remark: This result is due to Ando.)

Fact 5.10.18. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

- i) The matrices A and B are unitarily left equivalent if and only if $A^*A = B^*B$.
- ii) The matrices A and B are unitarily right equivalent if and only if $AA^* = BB^*$.
- iii) The matrices A and B are unitarily biequivalent if and only if A and B have the same singular values with the same multiplicity.

(Proof: See [715] and [1129, pp. 372, 373].) (Remark: In [715] A and B need not be the same size.) (Remark: The singular value decomposition provides a canonical form under unitary biequivalence in analogy with the Smith form under biequivalence.) (Remark: Note that $AA^* = BB^*$ implies that $\mathcal{R}(A) = \mathcal{R}(B)$, which implies right equivalence, which is an alternative proof of the immediate fact that unitary right equivalence implies right equivalence.)

Fact 5.10.19. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) $A^*A = B^*B$ if and only if there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that $A = SB$.

- ii) $A^*A \leq B^*B$ if and only if there exists a matrix $S \in \mathbb{F}^{n \times n}$ such that $A = SB$ and $S^*S \leq I$.
- iii) $A^*B + B^*A = 0$ if and only if there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that $(I - S)A = (I + S)B$.
- iv) $A^*B + B^*A \geq 0$ if and only if there exists a matrix $S \in \mathbb{F}^{n \times n}$ such that $(I - S)A = (I + S)B$ and $S^*S \leq I$.

(Proof: See [709, p. 406] and [1117].) (Remark: Statements iii) and iv) follow from i) and ii) by replacing A and B with $A - B$ and $A + B$, respectively.)

Fact 5.10.20. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{n \times m}$. Then, there exist matrices $X, Y \in \mathbb{F}^{n \times m}$ satisfying

$$AX + YB + C = 0$$

if and only if

$$\text{rank} \begin{bmatrix} A & 0 \\ 0 & -B \end{bmatrix} = \text{rank} \begin{bmatrix} A & C \\ 0 & -B \end{bmatrix}.$$

(Proof: See [1098, pp. 194, 195] and [1403].) (Remark: $AX + YB + C = 0$ is a generalization of Sylvester's equation. See Fact 5.10.21.) (Remark: This result is due to Roth.) (Remark: An explicit expression for all solutions is given by Fact 6.5.7, which applies to the case in which A and B are not necessarily square and thus X and Y are not necessarily the same size.)

Fact 5.10.21. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{n \times m}$. Then, there exists a matrix $X \in \mathbb{F}^{n \times m}$ satisfying

$$AX + XB + C = 0$$

if and only if the matrices

$$\begin{bmatrix} A & 0 \\ 0 & -B \end{bmatrix}, \quad \begin{bmatrix} A & C \\ 0 & -B \end{bmatrix}$$

are similar. In this case,

$$\begin{bmatrix} A & C \\ 0 & -B \end{bmatrix} = \begin{bmatrix} I & X \\ 0 & I \end{bmatrix} \begin{bmatrix} A & 0 \\ 0 & -B \end{bmatrix} \begin{bmatrix} I & -X \\ 0 & I \end{bmatrix}.$$

(Proof: See [1403]. For sufficiency, see [867, pp. 422–424] or [1098, pp. 194, 195].) (Remark: $AX + XB + C = 0$ is *Sylvester's equation*. See Proposition 7.2.4, Corollary 7.2.5, and Proposition 11.9.3.) (Remark: This result is due to Roth. See [217].)

Fact 5.10.22. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then, the matrices

$$\begin{bmatrix} A + B & A \\ 0 & -A - B \end{bmatrix}, \quad \begin{bmatrix} A + B & 0 \\ 0 & -A - B \end{bmatrix}$$

are similar. In fact,

$$\begin{bmatrix} A + B & A \\ 0 & -A - B \end{bmatrix} = \begin{bmatrix} I & X \\ 0 & I \end{bmatrix} \begin{bmatrix} A + B & 0 \\ 0 & -A - B \end{bmatrix} \begin{bmatrix} I & -X \\ 0 & I \end{bmatrix},$$

where $X \triangleq \frac{1}{4}(I + A - B)$. (Remark: This result is due to Tian.) (Remark: See Fact 5.10.21.)

Fact 5.10.23. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{n \times m}$, and assume that A and B are nilpotent. Then, the matrices

$$\begin{bmatrix} A & C \\ 0 & B \end{bmatrix}, \quad \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$$

are similar if and only if

$$\text{rank} \begin{bmatrix} A & C \\ 0 & B \end{bmatrix} = \text{rank } A + \text{rank } B$$

and

$$AC + CB = 0.$$

(Proof: See [1294].)

5.11 Facts on Eigenvalues and Singular Values for One Matrix

Fact 5.11.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is singular. If A is either simple or cyclic, then $\text{rank } A = n - 1$.

Fact 5.11.2. Let $A \in \mathbb{R}^{n \times n}$, and assume that $A \in \text{SO}(n)$. Then, $\text{amult}_A(-1)$ is even. Now, assume that $n = 3$. Then, the following statements hold:

- i) $\text{amult}_A(1)$ is either 1 or 3.
- ii) $\text{tr } A \geq -1$.
- iii) $\text{tr } A = -1$ if and only if $\text{mspec}(A) = \{1, -1, -1\}_{\text{ms}}$.

Fact 5.11.3. Let $A \in \mathbb{F}^{n \times n}$, let $\alpha \in \mathbb{F}$, and assume that $A^2 = \alpha A$. Then, $\text{spec}(A) \subseteq \{0, \alpha\}$.

Fact 5.11.4. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and let $\alpha \in \mathbb{R}$. Then, $A^2 = \alpha A$ if and only if $\text{spec}(A) \subseteq \{0, \alpha\}$. (Remark: See Fact 3.7.22.)

Fact 5.11.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then,

$$\text{spabs}(A) = \lambda_{\max}(A)$$

and

$$\text{sprad}(A) = \sigma_{\max}(A) = \max\{|\lambda_{\min}(A)|, \lambda_{\max}(A)\}.$$

If, in addition, A is positive semidefinite, then

$$\text{sprad}(A) = \sigma_{\max}(A) = \text{spabs}(A) = \lambda_{\max}(A).$$

(Remark: See Fact 5.12.2.)

Fact 5.11.6. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is skew Hermitian. Then, the eigenvalues of A are imaginary. (Proof: Let $\lambda \in \text{spec}(A)$. Since $0 \leq AA^* = -A^2$, it follows that $-\lambda^2 \geq 0$, and thus $\lambda^2 \leq 0$.)

Fact 5.11.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then, the following statements are equivalent:

- i) $\text{mspec}(A) = \text{mspec}(B)$.
- ii) $\text{rank } A = \text{rank } B$.
- iii) $\text{tr } A = \text{tr } B$.

Fact 5.11.8. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is idempotent.
 - ii) $\text{rank}(I - A) \leq \text{tr}(I - A)$, A is group invertible, and every eigenvalue of A is nonnegative.
 - iii) A and $I - A$ are group invertible, and every eigenvalue of A is nonnegative.
- (Proof: See [649].)

Fact 5.11.9. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_k, 0, \dots, 0\}_{\text{ms}}$. Then,

$$|\text{tr } A|^2 \leq \left(\sum_{i=1}^k |\lambda_i| \right)^2 \leq k \sum_{i=1}^k |\lambda_i|^2.$$

(Proof: Use Fact 1.15.3.)

Fact 5.11.10. Let $A \in \mathbb{F}^{n \times n}$, and assume that A has exactly k nonzero eigenvalues. Then,

$$\left. \begin{array}{l} |\text{tr } A|^2 \\ k |\text{tr } A^2| \leq k \text{tr } (A^2 A^2)^{1/2} \end{array} \right\} \leq k \text{tr } A^* A \leq (\text{rank } A) \text{tr } A^* A.$$

Furthermore, the upper left-hand inequality is an equality if and only if A is normal and all of the nonzero eigenvalues of A have the same absolute value, while the right-hand inequality is an equality if and only if A is group invertible. If, in addition, all of the eigenvalues of A are real, then

$$(\text{tr } A)^2 \leq k \text{tr } A^2 \leq k \text{tr } A^* A \leq (\text{rank } A) \text{tr } A^* A.$$

(Proof: The upper left-hand inequality in the first string is given in [1448]. The lower left-hand inequality in the first string is given by Fact 9.11.3. When all of the eigenvalues of A are real, the inequality $(\text{tr } A)^2 \leq k \text{tr } A^2$ follows from Fact 5.11.9.) (Remark: The inequality $|\text{tr } A|^2 \leq k |\text{tr } A^2|$ does not necessarily hold. Consider $\text{mspec}(A) = \{1, 1, j, -j\}_{\text{ms}}$.) (Remark: See Fact 3.7.22, Fact 8.17.7, Fact 9.13.17, and Fact 9.13.18.)

Fact 5.11.11. Let $A \in \mathbb{R}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\sum_{i=1}^n (\text{Re } \lambda_i)(\text{Im } \lambda_i) = 0$$

and

$$\text{tr } A^2 = \sum_{i=1}^n (\text{Re } \lambda_i)^2 - \sum_{i=1}^n (\text{Im } \lambda_i)^2.$$

Fact 5.11.12. Let $n \geq 2$, let $a_1, \dots, a_n > 0$, and define the symmetric matrix $A \in \mathbb{R}^{n \times n}$ by $A_{(i,j)} \triangleq a_i + a_j$ for all $i, j = 1, \dots, n$. Then,

$$\text{rank } A \leq 2$$

and

$$\text{mspec}(A) = \{\lambda, \mu, 0, \dots, 0\}_{\text{ms}},$$

where

$$\lambda \triangleq \sum_{i=1}^n a_i + \sqrt{n \sum_{i=1}^n a_i^2}, \quad \mu \triangleq \sum_{i=1}^n a_i - \sqrt{n \sum_{i=1}^n a_i^2}.$$

Furthermore, the following statements hold:

$$i) \quad \lambda > 0.$$

$$ii) \quad \mu \leq 0.$$

Furthermore, the following statements are equivalent:

$$iii) \quad \mu < 0.$$

$$iv) \quad \text{At least two of the numbers } a_1, \dots, a_n > 0 \text{ are distinct.}$$

$$v) \quad \text{rank } A = 2.$$

In this case,

$$\lambda_{\min}(A) = \mu < 0 < \text{tr } A = 2 \sum_{i=1}^n a_i < \lambda_{\max}(A) = \lambda.$$

(Proof: $A = a1_{1 \times n} + 1_{n \times 1}a^T$, where $a \triangleq [a_1 \ \dots \ a_n]^T$. Then, it follows from Fact 2.11.12 that $\text{rank } A \leq \text{rank}(a1_{1 \times n}) + \text{rank}(1_{n \times 1}a^T) = 2$. Furthermore, $\text{mspec}(A)$ follows from Fact 5.11.13, while Fact 1.15.14 implies that $\mu \leq 0$.) (Remark: See Fact 8.8.7.)

Fact 5.11.13. Let $x, y \in \mathbb{R}^n$. Then,

$$\text{mspec}(xy^T + yx^T) = \left\{ x^T y + \sqrt{x^T xy^T y}, x^T y - \sqrt{x^T xy^T y}, 0, \dots, 0 \right\}_{\text{ms}},$$

$$\text{sprad}(xy^T + yx^T) = \begin{cases} x^T y + \sqrt{x^T xy^T y}, & x^T y \geq 0, \\ \left| x^T y - \sqrt{x^T xy^T y} \right|, & x^T y \leq 0, \end{cases}$$

and

$$\text{spabs}(xy^T + yx^T) = x^T y + \sqrt{x^T xy^T y}.$$

If, in addition, x and y are nonzero, then $v_1, v_2 \in \mathbb{R}^n$ defined by

$$v_1 \triangleq \frac{1}{\|x\|}x + \frac{1}{\|y\|}y, \quad v_2 \triangleq \frac{1}{\|x\|}x - \frac{1}{\|y\|}y$$

are eigenvectors of $xy^T + yx^T$ corresponding to $x^T y + \sqrt{x^T xy^T y}$ and $x^T y - \sqrt{x^T xy^T y}$, respectively. (Proof: See [374, p. 539].) (Example: The spectrum of $\begin{bmatrix} 0_{n \times n} & 1_{n \times 1} \\ 1_{1 \times n} & 0 \end{bmatrix}$ is $\{-\sqrt{n}, 0, \dots, 0, \sqrt{n}\}_{\text{ms}}$.) (Problem: Extend this result to $\mathbb{C}$ and $xy^T + zw^T$. See Fact 4.9.16.)

Fact 5.11.14. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\text{mspec}[(I + A)^2] = \{(1 + \lambda_1)^2, \dots, (1 + \lambda_n)^2\}_{\text{ms}}.$$

If A is nonsingular, then

$$\text{mspec}(A^{-1}) = \{\lambda_1^{-1}, \dots, \lambda_n^{-1}\}_{\text{ms}}.$$

Finally, if $I + A$ is nonsingular, then

$$\text{mspec}[(I + A)^{-1}] = \{(1 + \lambda_1)^{-1}, \dots, (1 + \lambda_n)^{-1}\}_{\text{ms}}$$

and

$$\text{mspec}[A(I + A)^{-1}] = \{\lambda_1(1 + \lambda_1)^{-1}, \dots, \lambda_n(1 + \lambda_n)^{-1}\}_{\text{ms}}.$$

(Proof: Use Fact 5.11.15.)

Fact 5.11.15. Let $p, q \in \mathbb{F}[s]$, assume that p and q are coprime, define $g \triangleq p/q \in \mathbb{F}(s)$, let $A \in \mathbb{F}^{n \times n}$, let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, assume that $\text{roots}(q) \cap \text{spec}(A) = \emptyset$, and define $g(A) \triangleq p(A)[q(A)]^{-1}$. Then,

$$\text{mspec}[g(A)] = \{g(\lambda_1), \dots, g(\lambda_n)\}_{\text{ms}}.$$

(Proof: Statement *ii*) of Fact 4.10.9 implies that $q(A)$ is nonsingular.)

Fact 5.11.16. Let $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$. Then,

$$\sigma_{\max}(xy^*) = \sqrt{x^*xy^*y}.$$

If, in addition, $m = n$, then

$$\text{mspec}(xy^*) = \{x^*y, 0, \dots, 0\}_{\text{ms}},$$

$$\text{mspec}(I + xy^*) = \{1 + x^*y, 1, \dots, 1\}_{\text{ms}},$$

$$\text{sprad}(xy^*) = |x^*y|,$$

$$\text{spabs}(xy^*) = \max\{0, \text{Re } x^*y\}.$$

(Remark: See Fact 9.7.26.)

Fact 5.11.17. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{rank } A = 1$. Then,

$$\sigma_{\max}(A) = (\text{tr } AA^*)^{1/2}.$$

Fact 5.11.18. Let $x, y \in \mathbb{F}^n$, and assume that $x^*y \neq 0$. Then,

$$\sigma_{\max}[(x^*y)^{-1}xy^*] \geq 1.$$

Fact 5.11.19. Let $A \in \mathbb{F}^{n \times m}$, and let $\alpha \in \mathbb{F}$. Then, for all $i = 1, \dots, \min\{n, m\}$,

$$\sigma_i(\alpha A) = |\alpha| \sigma_i(A).$$

Fact 5.11.20. Let $A \in \mathbb{F}^{n \times m}$. Then, for all $i = 1, \dots, \text{rank } A$, it follows that

$$\sigma_i(A) = \sigma_i(A^*).$$

Fact 5.11.21. Let $A \in \mathbb{F}^{n \times n}$, and let $\lambda \in \text{spec}(A)$. Then, the following inequalities hold:

- i) $\sigma_{\min}(A) \leq |\lambda| \leq \sigma_{\max}(A)$.
- ii) $\lambda_{\min}\left[\frac{1}{2}(A + A^*)\right] \leq \text{Re } \lambda \leq \lambda_{\max}\left[\frac{1}{2}(A + A^*)\right]$.
- iii) $\lambda_{\min}\left[\frac{1}{2j}(A - A^*)\right] \leq \text{Im } \lambda \leq \lambda_{\max}\left[\frac{1}{2j}(A - A^*)\right]$.

(Remark: i) is *Browne's theorem*, ii) is *Bendixson's theorem*, and iii) is *Hirsch's theorem*. See [311, p. 17] and [963, pp. 140–144].) (Remark: See Fact 5.11.22, Fact 5.12.3, and Fact 9.11.8.)

Fact 5.11.22. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k [\sigma_{n-i+1}^2(A) - |\lambda_i|^2] \leq 2 \sum_{i=1}^k \left(\sigma_i^2\left[\frac{1}{2j}(A - A^*)\right] - |\text{Im } \lambda_i|^2 \right)$$

and

$$2 \sum_{i=1}^k \left(\sigma_{n-i+1}^2\left[\frac{1}{2j}(A - A^*)\right] - |\text{Im } \lambda_i|^2 \right) \leq \sum_{i=1}^k [\sigma_i^2(A) - |\lambda_i|^2].$$

Furthermore,

$$\sum_{i=1}^n [\sigma_i^2(A) - |\lambda_i|^2] = 2 \sum_{i=1}^n \left(\sigma_i^2\left[\frac{1}{2j}(A - A^*)\right] - |\text{Im } \lambda_i|^2 \right).$$

Finally, for all $i = 1, \dots, n$,

$$\sigma_n(A) \leq |\text{Re } \lambda_i| \leq \sigma_1(A)$$

and

$$\sigma_n\left[\frac{1}{2j}(A - A^*)\right] \leq |\text{Im } \lambda_i| \leq \sigma_1\left[\frac{1}{2j}(A - A^*)\right].$$

(Proof: See [552].) (Remark: See Fact 9.11.7.)

Fact 5.11.23. Let $A \in \mathbb{F}^{n \times n}$, let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, and let r denote the number of Jordan blocks in the Jordan decomposition of A . Then, for all $k = 1, \dots, r$,

$$\sum_{i=1}^k \sigma_{n-i+1}^2(A) \leq \sum_{i=1}^k |\lambda_i|^2 \leq \sum_{i=1}^k \sigma_i^2(A)$$

and

$$\sum_{i=1}^k \sigma_{n-i+1}^2\left[\frac{1}{2j}(A - A^*)\right] \leq \sum_{i=1}^k |\text{Im } \lambda_i|^2 \leq \sum_{i=1}^k \sigma_i^2\left[\frac{1}{2j}(A - A^*)\right].$$

(Proof: See [552].)

Fact 5.11.24. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1(A), \dots, \lambda_n(A)\}_{\text{ms}}$, where $\lambda_1(A), \dots, \lambda_n(A)$ are ordered such that $\text{Re } \lambda_1(A) \geq \dots \geq \text{Re } \lambda_n(A)$. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \text{Re } \lambda_i(A) \leq \sum_{i=1}^k \lambda_i\left[\frac{1}{2}(A + A^*)\right]$$

and

$$\sum_{i=1}^n \operatorname{Re} \lambda_i(A) = \operatorname{Re} \operatorname{tr} A = \operatorname{Re} \operatorname{tr} \frac{1}{2}(A + A^*) = \sum_{i=1}^n \lambda_i \left[\frac{1}{2}(A + A^*) \right].$$

In particular,

$$\lambda_{\min} \left[\frac{1}{2}(A + A^*) \right] \leq \operatorname{Re} \lambda_n(A) \leq \operatorname{spabs}(A) \leq \lambda_{\max} \left[\frac{1}{2}(A + A^*) \right].$$

Furthermore, the last right-hand inequality is an equality if and only if A is normal. (Proof: See [197, p. 74]. Also, see *xii*) and *xiv*) of Fact 11.15.7.) (Remark: $\operatorname{spabs}(A) = \operatorname{Re} \lambda_1(A)$.) (Remark: This result is due to Fan.)

Fact 5.11.25. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $i = 1, \dots, n$,

$$-\sigma_i(A) \leq \lambda_i \left[\frac{1}{2}(A + A^*) \right] \leq \sigma_i(A).$$

In particular,

$$-\sigma_{\min}(A) \leq \lambda_{\min} \left[\frac{1}{2}(A + A^*) \right] \leq \sigma_{\min}(A)$$

and

$$-\sigma_{\max}(A) \leq \lambda_{\max} \left[\frac{1}{2}(A + A^*) \right] \leq \sigma_{\max}(A).$$

(Proof: See [690, p. 447], [711, p. 151], or [971, p. 240].) (Remark: This result generalizes $\operatorname{Re} z \leq |z|$ for $z \in \mathbb{C}$.) (Remark: See Fact 8.17.4 and Fact 5.11.27.)

Fact 5.11.26. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} -\sigma_{\max}(A) &\leq -\sigma_{\min}(A) \\ &\leq \lambda_{\min} \left[\frac{1}{2}(A + A^*) \right] \\ &\leq \operatorname{spabs}(A) \\ &\leq \left\{ \begin{array}{l} |\operatorname{spabs}(A)| \leq \operatorname{sprad}(A) \\ \frac{1}{2} \lambda_{\max}(A + A^*) \end{array} \right\} \\ &\leq \sigma_{\max}(A). \end{aligned}$$

(Proof: Combine Fact 5.11.24 and Fact 5.11.25.)

Fact 5.11.27. Let $A \in \mathbb{F}^{n \times n}$, and let $\{\mu_1, \dots, \mu_n\}_{\text{ms}} = \{\frac{1}{2}|\lambda_1(A + A^*)|, \dots, \frac{1}{2}|\lambda_n(A + A^*)|\}_{\text{ms}}$, where $\mu_1 \geq \dots \geq \mu_n \geq 0$. Then, $\begin{bmatrix} \sigma_1(A) & \dots & \sigma_n(A) \end{bmatrix}$ weakly majorizes $\begin{bmatrix} \mu_1 & \dots & \mu_n \end{bmatrix}$. (Proof: See [971, p. 240].) (Remark: See Fact 5.11.25.)

Fact 5.11.28. Let $A \in \mathbb{F}^{n \times n}$, and let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_n$ are ordered such that $|\lambda_1| \geq \dots \geq |\lambda_n|$. Then, for all $k = 1, \dots, n$,

$$\prod_{i=1}^k |\lambda_i| \leq \prod_{i=1}^k \sigma_i(A)$$

with equality for $k = n$, that is,

$$|\det A| = \prod_{i=1}^n |\lambda_i| = \prod_{i=1}^n \sigma_i(A).$$

Hence, for all $k = 1, \dots, n$,

$$\prod_{i=k}^n \sigma_i(A) \leq \prod_{i=k}^n |\lambda_i|.$$

(Proof: See [197, p. 43], [690, p. 445], [711, p. 171], or [1485, p. 19].) (Remark: This result is due to Weyl.) (Remark: See Fact 8.18.21 and Fact 9.13.19.)

Fact 5.11.29. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_n$ are ordered such that $|\lambda_1| \geq \dots \geq |\lambda_n|$. Then,

$$\begin{aligned} \sigma_{\min}(A) &\leq \sigma_{\max}^{1/n}(A) \sigma_{\min}^{(n-1)/n}(A) \leq |\lambda_n| \leq |\lambda_1| \\ &\leq \sigma_{\min}^{1/n}(A) \sigma_{\max}^{(n-1)/n}(A) \leq \sigma_{\max}(A) \end{aligned}$$

and

$$\begin{aligned} \sigma_{\min}^n(A) &\leq \sigma_{\max}(A) \sigma_{\min}^{n-1}(A) \leq |\det A| \\ &\leq \sigma_{\min}(A) \sigma_{\max}^{n-1}(A) \leq \sigma_{\max}^n(A). \end{aligned}$$

(Proof: Use Fact 5.11.28. See [690, p. 445].) (Remark: See Fact 11.20.12.) (Remark: See Fact 8.13.1.)

Fact 5.11.30. Let $\beta_0, \dots, \beta_{n-1} \in \mathbb{F}$, define $A \in \mathbb{F}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \ddots & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ -\beta_0 & -\beta_1 & -\beta_2 & \cdots & -\beta_{n-2} & -\beta_{n-1} \end{bmatrix},$$

and define $\alpha \triangleq 1 + \sum_{i=0}^{n-1} |\beta_i|^2$. Then,

$$\begin{aligned} \sigma_1(A) &= \sqrt{\frac{1}{2} \left(\alpha + \sqrt{\alpha^2 - 4|\beta_0|^2} \right)}, \\ \sigma_2(A) &= \cdots = \sigma_{n-1}(A) = 1, \\ \sigma_n(A) &= \sqrt{\frac{1}{2} \left(\alpha - \sqrt{\alpha^2 - 4|\beta_0|^2} \right)}. \end{aligned}$$

In particular,

$$\sigma_1(N_n) = \cdots = \sigma_{n-1}(N_n) = 1$$

and

$$\sigma_{\min}(N_n) = 0.$$

(Proof: See [681, p. 523] or [802, 817].) (Remark: See Fact 6.3.28 and Fact 11.20.12.)

Fact 5.11.31. Let $\beta \in \mathbb{C}$. Then,

$$\sigma_{\max}\left(\begin{bmatrix} 1 & 2\beta \\ 0 & 1 \end{bmatrix}\right) = |\beta| + \sqrt{1 + |\beta|^2}$$

and

$$\sigma_{\min}\left(\begin{bmatrix} 1 & 2\beta \\ 0 & 1 \end{bmatrix}\right) = \sqrt{1 + |\beta|^2} - |\beta|.$$

(Proof: See [897].) (Remark: Inequalities involving the singular values of block-triangular matrices are given in [897].)

Fact 5.11.32. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\sigma_{\max}\left(\begin{bmatrix} I & 2A \\ 0 & I \end{bmatrix}\right) = \sigma_{\max}(A) + \sqrt{1 + \sigma_{\max}^2(A)}.$$

(Proof: See [681, p. 116].)

Fact 5.11.33. For $i = 1, \dots, l$, let $A_i \in \mathbb{F}^{n_i \times m_i}$. Then,

$$\sigma_{\max}[\text{diag}(A_1, \dots, A_l)] = \max\{\sigma_{\max}(A_1), \dots, \sigma_{\max}(A_l)\}.$$

Fact 5.11.34. Let $A \in \mathbb{F}^{n \times m}$, and let $r \triangleq \text{rank } A$. Then, for all $i = 1, \dots, r$,

$$\lambda_i(AA^*) = \lambda_i(A^*A) = \sigma_i(AA^*) = \sigma_i(A^*A) = \sigma_i^2(A).$$

In particular,

$$\sigma_{\max}(AA^*) = \sigma_{\max}^2(A),$$

and, if $n = m$, then

$$\sigma_{\min}(AA^*) = \sigma_{\min}^2(A).$$

Furthermore, for all $i = 1, \dots, r$,

$$\sigma_i(AA^*A) = \sigma_i^3(A).$$

Fact 5.11.35. Let $A \in \mathbb{F}^{n \times n}$. Then, $\sigma_{\max}(A) \leq 1$ if and only if $A^*A \leq I$.

Fact 5.11.36. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $i = 1, \dots, n$,

$$\sigma_i(A^A) = \prod_{\substack{j=1 \\ j \neq n+1-i}}^n \sigma_j(A).$$

(Proof: See Fact 4.10.7 and [1098, p. 149].)

Fact 5.11.37. Let $A \in \mathbb{F}^{n \times n}$. Then, $\sigma_1(A) = \sigma_n(A)$ if and only if there exist $\lambda \in \mathbb{F}$ and a unitary matrix $B \in \mathbb{F}^{n \times n}$ such that $A = \lambda B$. (Proof: See [1098, pp. 149, 165].)

Fact 5.11.38. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, the following statements hold:

- i) If σ is a singular value of A , then either $\sigma = 0$ or $\sigma \geq 1$.
- ii) If $A \neq 0$, then $\sigma_{\max}(A) \geq 1$.

iii) $\sigma_{\max}(A) = 1$ if and only if A is a projector.

iv) If $1 \leq \text{rank } A \leq n - 1$, then

$$\sigma_{\max}(A) = \sigma_{\max}(A_{\perp}).$$

v) If $A \neq 0$, then

$$\sigma_{\max}(A) = \sigma_{\max}(A + A^* - I) = \sigma_{\max}(A + A^*) - 1$$

and

$$\sigma_{\max}(I - 2A) = \sigma_{\max}(A) + [\sigma_{\max}^2(A) - 1]^{1/2}.$$

(Proof: See [537, 723, 744]. Statement *iv*) is given in [536, p. 61] and follows from Fact 5.11.39.) (Problem: Use Fact 5.9.26 to prove *iv*.)

Fact 5.11.39. Let $A \in \mathbb{F}^{n \times n}$, assume that A is idempotent, and assume that $1 \leq \text{rank } A \leq n - 1$. Then,

$$\sigma_{\max}(A) = \sigma_{\max}(A + A^* - I) = \frac{1}{\sin \theta},$$

where $\theta \in (0, \pi/2]$ is defined by

$$\cos \theta = \max\{|x^*y| : (x, y) \in \mathcal{R}(A) \times \mathcal{N}(A) \text{ and } x^*x = y^*y = 1\}.$$

(Proof: See [537, 744].) (Remark: θ is the minimal principal angle. See Fact 2.9.19 and Fact 5.12.17.) (Remark: Note that $\mathcal{N}(A) = \mathcal{R}(A_{\perp})$. See Fact 3.12.3.) (Remark: This result is due to Ljance.) (Remark: This result yields statement *iii*) of Fact 5.11.38.) (Remark: See Fact 10.9.18.)

Fact 5.11.40. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, be the tridiagonal matrix

$$A \triangleq \begin{bmatrix} b_1 & c_1 & 0 & \cdots & 0 & 0 \\ a_1 & b_2 & c_2 & \cdots & 0 & 0 \\ 0 & a_2 & b_3 & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & b_{n-1} & c_{n-1} \\ 0 & 0 & 0 & \cdots & a_{n-1} & b_n \end{bmatrix},$$

and assume that, for all $i = 1, \dots, n - 1$, $a_i c_i > 0$. Then, A is simple, and every eigenvalue of A is real. Hence, $\text{rank } A \geq n - 1$. (Proof: SAS^{-1} is symmetric, where $S \triangleq \text{diag}(d_1, \dots, d_n)$, $d_1 \triangleq 1$, and $d_{i+1} \triangleq (c_i/a_i)^{1/2} d_i$ for all $i = 1, \dots, n - 1$. For a proof of the fact that A is simple, see [481, p. 198].) (Remark: See Fact 5.11.41.)

Fact 5.11.41. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, be the tridiagonal matrix

$$A \triangleq \begin{bmatrix} b_1 & c_1 & 0 & \cdots & 0 & 0 \\ a_1 & b_2 & c_2 & \cdots & 0 & 0 \\ 0 & a_2 & b_3 & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & b_{n-1} & c_{n-1} \\ 0 & 0 & 0 & \cdots & a_{n-1} & b_n \end{bmatrix},$$

and assume that, for all $i = 1, \dots, n-1$, $a_i c_i \neq 0$. Then, A is reducible. Furthermore, let k_+ and k_- denote, respectively, the number of positive and negative numbers in the sequence

$$1, a_1 c_1, a_1 a_2 c_1 c_2, \dots, a_1 a_2 \cdots a_{n-1} c_1 c_2 \cdots c_{n-1}.$$

Then, A has at least $|k_+ - k_-|$ distinct real eigenvalues, of which at least $\max\{0, n - 3 \min\{k_+, k_-\}\}$ are simple. (Proof: See [1376].) (Remark: Note that $k_+ + k_- = n$ and $|k_+ - k_-| = n - 2 \min\{k_+, k_-\}$.) (Remark: This result yields Fact 5.11.40 as a special case.)

Fact 5.11.42. Let $A \in \mathbb{R}^{n \times n}$ be the tridiagonal matrix

$$A \triangleq \begin{bmatrix} 0 & 1 & 0 & & & & \\ n-1 & 0 & 2 & & & & 0 \\ 0 & n-2 & 0 & \ddots & & & \\ & \ddots & \ddots & \ddots & \ddots & & \\ & & \ddots & \ddots & 0 & n-2 & 0 \\ & 0 & & \ddots & 2 & 0 & n-1 \\ & & & & 0 & 1 & 0 \end{bmatrix}.$$

Then,

$$\chi_A(s) = \prod_{i=1}^n [s - (n + 1 - 2i)].$$

Hence,

$$\text{spec}(A) = \begin{cases} \{n-1, -(n-1), \dots, 1, -1\}, & n \text{ even,} \\ \{n-1, -(n-1), \dots, 2, -2, 0\}, & n \text{ odd.} \end{cases}$$

(Proof: See [1260].)

Fact 5.11.43. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 1$, be the tridiagonal, Toeplitz matrix

$$A \triangleq \begin{bmatrix} b & c & 0 & \cdots & 0 & 0 \\ a & b & c & \cdots & 0 & 0 \\ 0 & a & b & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & b & c \\ 0 & 0 & 0 & \cdots & a & b \end{bmatrix},$$

and assume that $ac > 0$. Then,

$$\text{spec}(A) = \left\{ b + 2\sqrt{ac} \cos \frac{i\pi}{n+1} : i = 1, \dots, n \right\}.$$

(Remark: See [681, p. 522].) (Remark: See Fact 3.20.7.)

Fact 5.11.44. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 1$, be the tridiagonal, Toeplitz matrix

$$A \triangleq \begin{bmatrix} 0 & 1/2 & 0 & \cdots & 0 & 0 \\ 1/2 & 0 & 1/2 & \cdots & 0 & 0 \\ 0 & 1/2 & 0 & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & 0 & 1/2 \\ 0 & 0 & 0 & \cdots & 1/2 & 0 \end{bmatrix}.$$

Then,

$$\text{spec}(A) = \left\{ \cos \frac{i\pi}{n+1} : i = 1, \dots, n \right\},$$

and, for $i = 1, \dots, n$, associated mutually orthogonal eigenvectors satisfying $\|v_i\|_2 = 1$ are, respectively,

$$v_i = \sqrt{\frac{2}{n+1}} \begin{bmatrix} \sin \frac{i\pi}{n+1} \\ \sin \frac{2i\pi}{n+1} \\ \vdots \\ \sin \frac{ni\pi}{n+1} \end{bmatrix}.$$

(Remark: See [822].)

Fact 5.11.45. Let $A \in \mathbb{F}^{n \times n}$, and assume that A has real eigenvalues. Then,

$$\begin{aligned} \frac{1}{n} \text{tr } A - \sqrt{\frac{n-1}{n} [\text{tr } A^2 - \frac{1}{n} (\text{tr } A)^2]} &\leq \lambda_{\min}(A) \\ &\leq \frac{1}{n} \text{tr } A - \sqrt{\frac{1}{n^2-n} [\text{tr } A^2 - \frac{1}{n} (\text{tr } A)^2]} \\ &\leq \frac{1}{n} \text{tr } A + \sqrt{\frac{1}{n^2-n} [\text{tr } A^2 - \frac{1}{n} (\text{tr } A)^2]} \\ &\leq \lambda_{\max}(A) \\ &\leq \frac{1}{n} \text{tr } A + \sqrt{\frac{n-1}{n} [\text{tr } A^2 - \frac{1}{n} (\text{tr } A)^2]}. \end{aligned}$$

Furthermore, for all $i = 1, \dots, n$,

$$|\lambda_i(A) - \frac{1}{n} \operatorname{tr} A| \leq \sqrt{\frac{n-1}{n} [\operatorname{tr} A^2 - \frac{1}{n} (\operatorname{tr} A)^2]}.$$

Finally, if $n = 2$, then

$$\frac{1}{n} \operatorname{tr} A - \sqrt{\frac{1}{n} \operatorname{tr} A^2 - \frac{1}{n^2} (\operatorname{tr} A)^2} = \lambda_{\min}(A) \leq \lambda_{\max}(A) = \frac{1}{n} \operatorname{tr} A + \sqrt{\frac{1}{n} \operatorname{tr} A^2 - \frac{1}{n^2} (\operatorname{tr} A)^2}.$$

(Proof: See [1448, 1449].) (Remark: These inequalities are related to Fact 1.15.12.)

Fact 5.11.46. Let $A \in \mathbb{F}^{n \times n}$, and let $\mu(A) \triangleq \min\{|\lambda| : \lambda \in \operatorname{spec}(A)\}$. Then,

$$\frac{1}{n} |\operatorname{tr} A| - \sqrt{\frac{n-1}{n} (\operatorname{tr} A A^* - \frac{1}{n} |\operatorname{tr} A|^2)} \leq \mu(A) \leq \sqrt{\frac{1}{n} \operatorname{tr} A A^*}$$

and

$$\frac{1}{n} |\operatorname{tr} A| \leq \operatorname{sprad}(A) \leq \frac{1}{n} |\operatorname{tr} A| + \sqrt{\frac{n-1}{n} (\operatorname{tr} A A^* - \frac{1}{n} |\operatorname{tr} A|^2)}.$$

(Proof: See Theorem 3.1 of [1448].)

Fact 5.11.47. Let $A \in \mathbb{F}^{n \times n}$, where $n \geq 2$, be the bidiagonal matrix

$$A \triangleq \begin{bmatrix} a_1 & b_1 & 0 & \cdots & 0 & 0 \\ 0 & a_2 & b_2 & \cdots & 0 & 0 \\ 0 & 0 & a_3 & \ddots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & a_{n-1} & b_{n-1} \\ 0 & 0 & 0 & \cdots & 0 & a_n \end{bmatrix},$$

and assume that $a_1, \dots, a_n, b_1, \dots, b_{n-1}$ are nonzero. Then, the following statements hold:

- i) The singular values of A are distinct.
- ii) If $B \in \mathbb{F}^{n \times n}$ is bidiagonal and $|B| = |A|$, then A and B have the same singular values.
- iii) If $B \in \mathbb{F}^{n \times n}$ is bidiagonal, $|A| \leq |B|$, and $|A| \neq |B|$, then $\sigma_{\max}(A) < \sigma_{\max}(B)$.
- iv) If $B \in \mathbb{F}^{n \times n}$ is bidiagonal, $|I \circ A| \leq |I \circ B|$, and $|I \circ A| \neq |I \circ B|$, then $\sigma_{\min}(A) < \sigma_{\min}(B)$.
- v) If $B \in \mathbb{F}^{n \times n}$ is bidiagonal, $|I_{\sup} \circ A| \leq |I_{\sup} \circ B|$, and $|I_{\sup} \circ A| \neq |I_{\sup} \circ B|$, where $I_{\sup}$ denotes the matrix all of whose entries on the superdiagonal are 1 and are 0 otherwise, then $\sigma_{\min}(B) < \sigma_{\min}(A)$.

(Proof: See [981, p. 17-5].)

5.12 Facts on Eigenvalues and Singular Values for Two or More Matrices

Fact 5.12.1. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{n \times m}$, let $r \triangleq \text{rank } B$, and define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & 0 \end{bmatrix}$. Then, $\nu_-(\mathcal{A}) \geq r$, $\nu_0(\mathcal{A}) \geq 0$, and $\nu_+(\mathcal{A}) \geq r$. If, in addition, $n = m$ and B is nonsingular, then $\text{In } \mathcal{A} = \begin{bmatrix} n & 0 & n \end{bmatrix}^T$. (Proof: See [717].) (Remark: See Proposition 5.6.6.)

Fact 5.12.2. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\text{sprad}(A + B) \leq \sigma_{\max}(A + B) \leq \sigma_{\max}(A) + \sigma_{\max}(B).$$

If, in addition, A and B are Hermitian, then

$$\text{sprad}(A + B) = \sigma_{\max}(A + B) \leq \sigma_{\max}(A) + \sigma_{\max}(B) = \text{sprad}(A) + \text{sprad}(B)$$

and

$$\lambda_{\min}(A) + \lambda_{\min}(B) \leq \lambda_{\min}(A + B) \leq \lambda_{\max}(A + B) \leq \lambda_{\max}(A) + \lambda_{\max}(B).$$

(Proof: Use Lemma 8.4.3 for the last string of inequalities.) (Remark: See Fact 5.11.5.)

Fact 5.12.3. Let $A, B \in \mathbb{F}^{n \times n}$, and let λ be an eigenvalue of $A + B$. Then,

$$\frac{1}{2}\lambda_{\min}(A^* + A) + \frac{1}{2}\lambda_{\min}(B^* + B) \leq \text{Re } \lambda \leq \frac{1}{2}\lambda_{\max}(A^* + A) + \frac{1}{2}\lambda_{\max}(B^* + B).$$

(Proof: See [311, p. 18].) (Remark: See Fact 5.11.21.)

Fact 5.12.4. Let $A, B \in \mathbb{F}^{n \times n}$ be normal, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}$ and $\text{mspec}(B) = \{\mu_1, \dots, \mu_n\}$. Then,

$$\min \text{Re} \sum_{i=1}^n \lambda_i \mu_{\sigma(i)} \leq \text{Re } \text{tr } AB \leq \max \text{Re} \sum_{i=1}^n \lambda_i \mu_{\sigma(i)},$$

where “max” and “min” are taken over all permutations σ of the eigenvalues of B . Now, assume that A and B are Hermitian. Then, $\text{tr } AB$ is real, and

$$\sum_{i=1}^n \lambda_i(A) \lambda_{n-i+1}(B) \leq \text{tr } AB \leq \sum_{i=1}^n \lambda_i(A) \lambda_i(B).$$

Furthermore, the last inequality is an identity if and only if there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that $A = S \text{diag}[\lambda_1(A), \dots, \lambda_n(A)] S^*$ and $B = S \text{diag}[\lambda_1(B), \dots, \lambda_n(B)] S^*$. (Proof: See [957]. For the second string of inequalities, use Fact 1.16.4. For the last statement, see [239, p. 10] or [891].) (Remark: The upper bound for $\text{tr } AB$ is due to Fan.) (Remark: See Fact 5.12.5, Fact 5.12.8, Proposition 8.4.13, Fact 8.12.28, and Fact 8.18.18.)

Fact 5.12.5. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that B is Hermitian. Then,

$$\sum_{i=1}^n \lambda_i \left[\frac{1}{2}(A + A^*) \right] \lambda_{n-i+1}(B) \leq \text{Re } \text{tr } AB \leq \sum_{i=1}^n \lambda_i \left[\frac{1}{2}(A + A^*) \right] \lambda_i(B).$$

(Proof: Apply the second string of inequalities in Fact 5.12.4.) (Remark: For A, B real, these inequalities are given in [837]. The complex case is given in [871].) (See

Proposition 8.4.13 for the case in which B is positive semidefinite.)

Fact 5.12.6. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$, and let $r \triangleq \min\{\text{rank } A, \text{rank } B\}$. Then,

$$|\text{tr } AB| \leq \sum_{i=1}^r \sigma_i(A) \sigma_i(B).$$

(Proof: See [971, pp. 514, 515] or [1098, p. 148].) (Remark: Applying Fact 5.12.4 to $\begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix}$ and $\begin{bmatrix} 0 & B^* \\ B & 0 \end{bmatrix}$ and using Proposition 5.6.6 yields the weaker result

$$|\text{Re tr } AB| \leq \sum_{i=1}^r \sigma_i(A) \sigma_i(B).$$

See [239, p. 14].) (Remark: This result is due to Mirsky.) (Remark: See Fact 5.12.7.) (Remark: A generalization is given by Fact 9.14.3.)

Fact 5.12.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that B is positive semidefinite. Then,

$$|\text{tr } AB| \leq \sigma_{\max}(A) \text{tr } B.$$

(Proof: Apply Fact 5.12.6.) (Remark: A generalization is given by Fact 9.14.4.)

Fact 5.12.8. Let $A, B \in \mathbb{R}^{n \times n}$, assume that B is symmetric, and define $C \triangleq \frac{1}{2}(A + A^T)$. Then,

$$\begin{aligned} \lambda_{\min}(C) \text{tr } B - \lambda_{\min}(B) [n \lambda_{\min}(C) - \text{tr } A] \\ \leq \text{tr } AB \leq \lambda_{\max}(C) \text{tr } B - \lambda_{\max}(B) [n \lambda_{\max}(C) - \text{tr } A]. \end{aligned}$$

(Proof: See [468].) (Remark: See Fact 5.12.4, Proposition 8.4.13, and Fact 8.12.28. Extensions are given in [1071].)

Fact 5.12.9. Let $A, B, Q, S_1, S_2 \in \mathbb{R}^{n \times n}$, assume that A and B are symmetric, assume that Q, S_1 , and S_2 are orthogonal, assume that $S_1^T A S_1$ and $S_2^T B S_2$ are diagonal with the diagonal entries arranged in nonincreasing order, and define the orthogonal matrices $Q_1, Q_2 \in \mathbb{R}^{n \times n}$ by $Q_1 \triangleq S_1 \text{revdiag}(\pm 1, \dots, \pm 1) S_1^T$ and $Q_2 \triangleq S_2 \text{diag}(\pm 1, \dots, \pm 1) S_2^T$. Then,

$$\text{tr } A Q_1 B Q_1^T \leq \text{tr } A Q B Q^T \leq \text{tr } A Q_2 B Q_2^T.$$

(Proof: See [156, 891].) (Remark: See Fact 5.12.8.)

Fact 5.12.10. Let $A_1, \dots, A_k, B_1, \dots, B_k \in \mathbb{F}^{n \times n}$, and assume that $A_1, \dots, A_k$ are unitary. Then,

$$|\text{tr } A_1 B_1 \cdots A_k B_k| \leq \sum_{i=1}^n \sigma_i(B_1) \cdots \sigma_i(B_k).$$

(Proof: See [971, p. 516].) (Remark: This result is due to Fan.) (Remark: See Fact 5.12.9.)

Fact 5.12.11. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that $AB = BA$. Then,

$$\text{sprad}(AB) \leq \text{sprad}(A) \text{sprad}(B)$$

and

$$\text{sprad}(A + B) \leq \text{sprad}(A) + \text{sprad}(B).$$

(Proof: Use Fact 5.17.4.) (Remark: If $AB \neq BA$, then both of these inequalities may be violated. Consider $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$.)

Fact 5.12.12. Let $A, B \in \mathbb{C}^{n \times n}$, assume that A and B are normal, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$ and $\text{mspec}(B) = \{\mu_1, \dots, \mu_n\}_{\text{ms}}$. Then,

$$|\det(A + B)| \leq \min \left\{ \prod_{i=1}^n \max_{j=1, \dots, n} |\lambda_i + \mu_j|, \prod_{j=1}^n \max_{i=1, \dots, n} |\lambda_i + \mu_j| \right\}.$$

(Proof: See [1110].) (Remark: Equality is discussed in [161].) (Remark: See Fact 9.14.18.)

Fact 5.12.13. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times m}$. Then,

$$\det(ABB^*A^*) \leq \left[\prod_{i=1}^m \sigma_i(B) \right] \det(AA^*).$$

(Proof: See [447, p. 218].)

Fact 5.12.14. Let $A, B, C \in \mathbb{F}^{n \times n}$, assume that $\text{spec}(A) \cap \text{spec}(B) = \emptyset$, and assume that $[A + B, C] = 0$ and $[AB, C] = 0$. Then, $[A, C] = [B, C] = 0$. (Proof: The result follows from Corollary 7.2.5.) (Remark: This result is due to Embry. See [217].)

Fact 5.12.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then,

$$\text{spec}(AB) \subset [0, 1]$$

and

$$\text{spec}(A - B) \subset [-1, 1].$$

(Proof: See [38], [536, p. 53], or [1098, p. 147].) (Remark: The first result is due to Afriat.)

Fact 5.12.16. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the following statements are equivalent:

- i) AB is a projector.
- ii) $\text{spec}(A + B) \subset \{0\} \cup [1, \infty)$.
- iii) $\text{spec}(A - B) \subset \{-1, 0, 1\}$.

(Proof: See [537, 598].) (Remark: See Fact 3.13.20 and Fact 6.4.23.)

Fact 5.12.17. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are nonzero projectors, and define the minimal principal angle $\theta \in [0, \pi/2]$ by

$$\cos \theta = \max\{|x^*y| : (x, y) \in \mathcal{R}(A) \times \mathcal{R}(B) \text{ and } x^*x = y^*y = 1\}.$$

Then, the following statements hold:

- i)* $\sigma_{\max}(AB) = \sigma_{\max}(BA) = \cos \theta$.
- ii)* $\sigma_{\max}(A + B) = 1 + \sigma_{\max}(AB) = 1 + \cos \theta$.
- iii)* $1 \leq \sigma_{\max}(AB) + \sigma_{\max}(A - B)$.
- iv)* If $\sigma_{\max}(A - B) < 1$, then $\text{rank } A = \text{rank } B$.
- v)* $\theta > 0$ if and only if $\mathcal{R}(A) \cap \mathcal{R}(B) = \{0\}$.

Furthermore, the following statements are equivalent:

- vi)* $A - B$ is nonsingular.
- vii)* $\mathcal{R}(A)$ and $\mathcal{R}(B)$ are complementary subspaces.
- viii)* $\sigma_{\max}(A + B - I) < 1$.

Now, assume that $A - B$ is nonsingular. Then, the following statements hold:

- ix)* $\sigma_{\max}(AB) < 1$.
- x)* $\sigma_{\max}[(A - B)^{-1}] = \frac{1}{\sqrt{1 - \sigma_{\max}^2(AB)}} = 1/\sin \theta$.
- xi)* $\sigma_{\min}(A - B) = \sin \theta$.
- xii)* $\sigma_{\min}^2(A - B) + \sigma_{\max}^2(AB) = 1$.
- xiii)* $I - AB$ is nonsingular.
- xiv)* If $\text{rank } A = \text{rank } B$, then $\sigma_{\max}(A - B) = \sin \theta$.

(Proof: Statement *i)* is given in [744]. Statement *ii)* is given in [537]. Statement *iii)* follows from the first inequality in Fact 8.18.11. For *iv)*, see [447, p. 195] or [560, p. 389]. Statement *v)* is given in [560, p. 393]. Fact 3.13.24 shows that *vi)* and *vii)* are equivalent. Statement *viii)* is given in [272]; see also [536, p. 236]. Statement *xiv)* follows from [1230, pp. 92, 93].) (Remark: Additional conditions for the nonsingularity of $A - B$ are given in Fact 3.13.24.) (Remark: See Fact 2.9.19 and Fact 5.11.39.) (Remark: See Fact 5.12.18.)

Fact 5.12.18. Let $A \in \mathbb{F}^{n \times n}$, assume that A is idempotent, and let $P, Q \in \mathbb{F}^{n \times n}$, where P is the projector onto $\mathcal{R}(A)$ and Q is the projector onto $\mathcal{N}(A)$. Then, the following statements hold:

- i)* $P - Q$ is nonsingular.
- ii)* $(P - Q)^{-1} = A + A^* - I = A - A_{\perp}^*$.
- iii)* $\sigma_{\max}(A) = \frac{1}{\sqrt{1 - \sigma_{\max}^2(PQ)}} = \sigma_{\max}[(P - Q)^{-1}] = \sigma_{\max}(A + A^* - I)$.
- iv)* $\sigma_{\max}(A) = 1/\sin \theta$, where θ is the minimal principal angle $\theta \in [0, \pi/2]$ defined by

$$\cos \theta = \max\{|x^*y| : (x, y) \in \mathcal{R}(P) \times \mathcal{R}(Q) \text{ and } x^*x = y^*y = 1\}.$$

- v)* $\sigma_{\min}^2(P - Q) = 1 - \sigma_{\max}^2(PQ)$.
- vi)* $\sigma_{\max}(PQ) = \sigma_{\max}(QP) = \sigma_{\max}(P + Q - I) < 1$.

(Proof: See [1115] and Fact 5.12.17. The nonsingularity of $P - Q$ follows from Fact

3.13.24. Statement *ii*) is given by Fact 3.13.24 and Fact 6.3.25. The first identity in *iii*) is given in [272]. See also [537].) (Remark: $A_{\perp}^*$ is the idempotent matrix onto $\mathcal{R}(A)^{\perp}$ along $\mathcal{N}(A)^{\perp}$. See Fact 3.12.3.) (Remark: $P = AA^+$ and $Q = I - A^+A$.)

Fact 5.12.19. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are idempotent. Then, $A - B$ is idempotent if and only if $A - B$ is group invertible and every eigenvalue of $A - B$ is nonnegative. (Proof: See [649].) (Remark: This result is due to Makelainen and Styán.) (Remark: See Fact 3.12.29.) (Remark: Conditions for a matrix to be expressible as a difference of idempotents are given in [649].)

Fact 5.12.20. Let $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, and $C \in \mathbb{R}^{m \times m}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^T & C \end{bmatrix} \in \mathbb{R}^{(n+m) \times (n+m)}$, and assume that $\mathcal{A}$ is symmetric. Then,

$$\lambda_{\min}(\mathcal{A}) + \lambda_{\max}(\mathcal{A}) \leq \lambda_{\max}(A) + \lambda_{\max}(C).$$

(Proof: See [223, p. 56].)

Fact 5.12.21. Let $M \in \mathbb{R}^{r \times r}$, assume that M is positive definite, let $C, K \in \mathbb{R}^{r \times r}$, assume that C and K are positive semidefinite, and consider the equation

$$M\ddot{q} + C\dot{q} + Kq = 0.$$

Then, $x(t) \triangleq \begin{bmatrix} q(t) \\ \dot{q}(t) \end{bmatrix}$ satisfies $\dot{x}(t) = Ax(t)$, where A is the $2r \times 2r$ matrix

$$A \triangleq \begin{bmatrix} 0 & I \\ -M^{-1}K & -M^{-1}C \end{bmatrix}.$$

Furthermore, the following statements hold:

i) A , K , and M satisfy

$$\det A = \frac{\det K}{\det M}.$$

ii) A and K satisfy

$$\text{rank } A = r + \text{rank } K.$$

iii) A is nonsingular if and only if K is positive definite. In this case,

$$A^{-1} = \begin{bmatrix} -K^{-1}C & -K^{-1}M \\ I & 0 \end{bmatrix}.$$

iv) Let $\lambda \in \mathbb{C}$. Then, $\lambda \in \text{spec}(A)$ if and only if $\det(\lambda^2 M + \lambda C + K) = 0$.

v) If $\lambda \in \text{spec}(A)$, $\text{Re } \lambda = 0$, and $\text{Im } \lambda \neq 0$, then λ is semisimple.

vi) $\text{mspec}(A) \subset \text{CLHP}$.

vii) If $C = 0$, then $\text{spec}(A) \subset j\mathbb{R}$.

viii) If C and K are positive definite, then $\text{spec}(A) \subset \text{OLHP}$.

ix) $\hat{x}(t) \triangleq \begin{bmatrix} \frac{1}{\sqrt{2}} K^{1/2} q(t) \\ \frac{1}{\sqrt{2}} M^{1/2} \dot{q}(t) \end{bmatrix}$ satisfies $\dot{x}(t) = \hat{A}x(t)$, where

$$\hat{A} \triangleq \begin{bmatrix} 0 & K^{1/2} M^{-1/2} \\ -M^{-1/2} K^{1/2} & -M^{-1/2} C M^{-1/2} \end{bmatrix}.$$

If, in addition, $C = 0$, then $\hat{A}$ is skew symmetric.

$x) \hat{x}(t) \triangleq \begin{bmatrix} M^{1/2}q(t) \\ M^{1/2}\dot{q}(t) \end{bmatrix}$ satisfies $\dot{x}(t) = \hat{A}x(t)$, where

$$\hat{A} \triangleq \begin{bmatrix} 0 & I \\ -M^{-1/2}KM^{-1/2} & -M^{-1/2}CM^{-1/2} \end{bmatrix}.$$

If, in addition, $C = 0$, then $\hat{A}$ is Hamiltonian.

(Remark: M, C , and K are mass, damping, and stiffness matrices, respectively. See [186].) (Remark: See Fact 11.18.38.) (Problem: Prove v .)

Fact 5.12.22. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that A and B are positive semidefinite. Then, every eigenvalue λ of $\begin{bmatrix} 0 & B \\ -A & 0 \end{bmatrix}$ satisfies $\operatorname{Re} \lambda = 0$. (Proof: Square this matrix.) (Problem: What happens if A and B have different dimensions?) In addition, let $C \in \mathbb{R}^{n \times n}$, and assume that C is (positive semidefinite, positive definite). Then, every eigenvalue of $\begin{bmatrix} 0 & A \\ -B & -C \end{bmatrix}$ satisfies ($\operatorname{Re} \lambda \leq 0$, $\operatorname{Re} \lambda < 0$). (Problem: Consider also $\begin{bmatrix} -C & A \\ -B & -C \end{bmatrix}$ and $\begin{bmatrix} -C & A \\ -A & -C \end{bmatrix}$.)

5.13 Facts on Matrix Pencils

Fact 5.13.1. Let $A, B \in \mathbb{F}^{n \times n}$, assume that $P_{A,B}$ is a regular pencil, let $\mathcal{S} \subseteq \mathbb{F}^n$, assume that $\mathcal{S}$ is a subspace, let $k \triangleq \dim \mathcal{S}$, let $S \in \mathbb{F}^{n \times k}$, and assume that $\mathcal{R}(S) = \mathcal{S}$. Then, the following statements are equivalent:

$i) \dim(AS + BS) = \dim \mathcal{S}$.

$ii) \text{ There exists a matrix } M \in \mathbb{F}^{k \times k} \text{ such that } AS = BSM.$

(Proof: See [872, p. 144].) (Remark: $\mathcal{S}$ is a *deflating subspace* of $P_{A,B}$. This result generalizes Fact 2.9.25.)

5.14 Facts on Matrix Eigenstructure

Fact 5.14.1. Let $A \in \mathbb{F}^{n \times n}$. Then, $\operatorname{rank} A = 1$ if and only if $\operatorname{gmult}_A(0) = n - 1$. In this case, $\operatorname{mspec}(A) = \{\operatorname{tr} A, 0, \dots, 0\}_{\operatorname{ms}}$. (Proof: Use Proposition 5.5.3.) (Remark: See Fact 2.10.19.)

Fact 5.14.2. Let $A \in \mathbb{F}^{n \times n}$, let $\lambda \in \operatorname{spec}(A)$, assume that λ is cyclic, let $i \in \{1, \dots, n\}$ be such that $\operatorname{rank}(A - \lambda I)_{(\{i\}^{\sim}, \{1, \dots, n\})} = n - 1$, and define $x \in \mathbb{C}^n$ by

$$x \triangleq \begin{bmatrix} \det(A - \lambda I)_{[i,1]} \\ -\det(A - \lambda I)_{[i,2]} \\ \vdots \\ (-1)^{n+1} \det(A - \lambda I)_{[i,n]} \end{bmatrix}.$$

Then, x is an eigenvector of A associated with λ . (Proof: See [1339].)

Fact 5.14.3. Let $n \geq 2$, $x, y \in \mathbb{F}^n$, define $A \triangleq xy^T$, and assume that $\text{rank } A = 1$, that is, A is nonzero. Then, the following statements are equivalent:

- i) A is semisimple.
- ii) $y^T x \neq 0$.
- iii) $\text{tr } A \neq 0$.
- iv) A is group invertible.
- v) $\text{ind } A = 1$.
- vi) $\text{amult}_A(0) = n - 1$.

Furthermore, the following statements are equivalent:

- vii) A is defective.
- viii) $y^T x = 0$.
- ix) $\text{tr } A = 0$.
- x) A is not group invertible.
- xi) $\text{ind } A = 2$.
- xii) A is nilpotent.
- xiii) $\text{amult}_A(0) = n$.
- xiv) $\text{spec}(A) = \{0\}$.

(Remark: See Fact 2.10.19.)

Fact 5.14.4. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is group invertible.
- ii) $\mathcal{R}(A) = \mathcal{R}(A^2)$.
- iii) $\text{ind } A \leq 1$.
- iv) $\text{rank } A = \sum_{i=1}^r \text{amult}_A(\lambda_i)$, where $\lambda_1, \dots, \lambda_r$ are the nonzero eigenvalues of A .

Fact 5.14.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is diagonalizable over $\mathbb{F}$. Then, A^T , $\bar{A}$, A^* , and A^A are diagonalizable. If, in addition, A is nonsingular, then A^{-1} is diagonalizable. (Proof: See Fact 2.16.10 and Fact 3.7.10.)

Fact 5.14.6. Let $A \in \mathbb{F}^{n \times n}$, assume that A is diagonalizable over $\mathbb{F}$ with eigenvalues $\lambda_1, \dots, \lambda_n$, and let $B \triangleq \text{diag}(\lambda_1, \dots, \lambda_n)$. If, $x_1, \dots, x_n \in \mathbb{F}^n$ are linearly independent eigenvectors of A associated with $\lambda_1, \dots, \lambda_n$, respectively, then $A = SBS^{-1}$, where $S \triangleq \begin{bmatrix} x_1 & \cdots & x_n \end{bmatrix}$. Conversely, if $S \in \mathbb{F}^{n \times n}$ is nonsingular and $A = SBS^{-1}$, then, for all $i = 1, \dots, n$, $\text{col}_i(S)$ is an associated eigenvector.

Fact 5.14.7. Let $A \in \mathbb{F}^{n \times n}$, let $S \in \mathbb{F}^{n \times n}$, assume that S is nonsingular, let $\lambda \in \mathbb{C}$, and assume that $\text{row}_1(S^{-1}AS) = \lambda e_1^T$. Then, $\lambda \in \text{spec}(A)$, and $\text{col}_1(S)$ is an associated eigenvector.

Fact 5.14.8. Let $A \in \mathbb{C}^{n \times n}$. Then, there exist $v_1, \dots, v_n \in \mathbb{C}^n$ such that the following statements hold:

- i) $v_1, \dots, v_n \in \mathbb{C}^n$ are linearly independent.
- ii) For each $k \times k$ Jordan block of A associated with $\lambda \in \text{spec}(A)$, there exist $v_{i_1}, \dots, v_{i_k}$ such that

$$\begin{aligned} Av_{i_1} &= \lambda v_{i_1}, \\ Av_{i_2} &= \lambda v_{i_2} + v_{i_1}, \\ &\vdots \\ Av_{i_k} &= \lambda v_{i_k} + v_{i_{k-1}}. \end{aligned}$$

- iii) Let λ and $v_{i_1}, \dots, v_{i_k}$ be given by ii). Then,

$$\text{span}\{v_{i_1}, \dots, v_{i_k}\} = \mathcal{N}[(\lambda I - A)^k].$$

(Remark: $v_1, \dots, v_n$ are *generalized eigenvectors* of A .) (Remark: $(v_{i_1}, \dots, v_{i_k})$ is a *Jordan chain* of A associated with λ . See [867, pp. 229–231].) (Remark: See Fact 11.13.7.)

Fact 5.14.9. Let $A \in \mathbb{F}^{n \times n}$. Then, A is cyclic if and only if there exists a vector $b \in \mathbb{F}^n$ such that $\begin{bmatrix} b & Ab & \cdots & A^{n-1}b \end{bmatrix}$ is nonsingular. (Proof: See Fact 12.20.13.) (Remark: (A, b) is controllable. See Corollary 12.6.3.)

Fact 5.14.10. Let $A \in \mathbb{F}^{n \times n}$, and define the positive integer m by

$$m \triangleq \max_{\lambda \in \text{spec}(A)} \text{gmult}_A(\lambda).$$

Then, m is the smallest integer such that there exists $B \in \mathbb{F}^{n \times m}$ such that $\text{rank} \begin{bmatrix} B & AB & \cdots & A^{n-1}B \end{bmatrix} = n$. (Proof: See Fact 12.20.13.) (Remark: (A, B) is controllable. See Corollary 12.6.3.)

Fact 5.14.11. Let $A \in \mathbb{R}^{n \times n}$. Then, A is cyclic and semisimple if and only if A is simple.

Fact 5.14.12. Let $A = \text{revdiag}(a_1, \dots, a_n) \in \mathbb{R}^{n \times n}$. Then, A is semisimple if and only if, for all $i = 1, \dots, n$, a_i and a_{n+1-i} are either both zero or both nonzero. (Proof: See [626, p. 116], [804], or [1098, pp. 68, 86].)

Fact 5.14.13. Let $A \in \mathbb{F}^{n \times n}$. Then, A has at least m real eigenvalues and m associated linearly independent eigenvectors if and only if there exists a positive-semidefinite matrix $S \in \mathbb{F}^{n \times n}$ such that $\text{rank } S = m$ and $AS = SA^*$. (Proof: See [1098, pp. 68, 86].) (Remark: See Proposition 5.5.12.) (Remark: This result is due to Drazin and Haynsworth.)

Fact 5.14.14. Let $A \in \mathbb{F}^{n \times n}$, assume that A is normal, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then, there exist vectors $x_1, \dots, x_n \in \mathbb{C}^n$ such that $x_i^* x_j = \delta_{ij}$ for all $i, j = 1, \dots, n$ and

$$A = \sum_{i=1}^n \lambda_i x_i x_i^*.$$

(Remark: This result is a restatement of Corollary 5.4.4.)

Fact 5.14.15. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, where $|\lambda_1| \geq \dots \geq |\lambda_n|$. Then, the following statements are equivalent:

- i) A is normal.
- ii) For all $i = 1, \dots, n$, $|\lambda_i| = \sigma_i(A)$.
- iii) $\sum_{i=1}^n |\lambda_i|^2 = \sum_{i=1}^n \sigma_i^2(A)$.
- iv) There exists $p \in \mathbb{F}[s]$ such that $A = p(A^*)$.
- v) Every eigenvector of A is also an eigenvector of A^* .
- vi) $AA^* - A^*A$ is either positive semidefinite or negative semidefinite.
- vii) For all $x \in \mathbb{F}^n$, $x^*A^*Ax = x^*AA^*x$.
- viii) For all $x, y \in \mathbb{F}^n$, $x^*A^*Ay = x^*AA^*y$.

In this case,

$$\text{sprad}(A) = \sigma_{\max}(A).$$

(Proof: See [589] or [1098, p. 146].) (Remark: See Fact 9.11.2 and Fact 9.8.13.)

Fact 5.14.16. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is (simple, cyclic, derogatory, semisimple, defective, diagonalizable over $\mathbb{F}$).
- ii) There exists $\alpha \in \mathbb{F}$ such that $A + \alpha I$ is (simple, cyclic, derogatory, semisimple, defective, diagonalizable over $\mathbb{F}$).
- iii) For all $\alpha \in \mathbb{F}$, $A + \alpha I$ is (simple, cyclic, derogatory, semisimple, defective, diagonalizable over $\mathbb{F}$).

Fact 5.14.17. Let $x, y \in \mathbb{F}^n$, assume that $x^T y \neq 1$, and define the elementary matrix $A \triangleq I - xy^T$. Then, A is semisimple if and only if either $xy^T = 0$ or $x^T y \neq 0$. (Remark: Use Fact 5.14.3 and Fact 5.14.16.)

Fact 5.14.18. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nilpotent. Then, A is nonzero if and only if A is defective.

Fact 5.14.19. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is either involutory or skew involutory. Then, A is semisimple.

Fact 5.14.20. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is involutory. Then, A is diagonalizable over $\mathbb{R}$.

Fact 5.14.21. Let $A \in \mathbb{F}^{n \times n}$, assume that A is semisimple, and assume that $A^3 = A^2$. Then, A is idempotent.

Fact 5.14.22. Let $A \in \mathbb{F}^{n \times n}$. Then, A is cyclic if and only if every matrix $B \in \mathbb{F}^{n \times n}$ satisfying $AB = BA$ is a polynomial in A . (Proof: See [711, p. 275].) (Remark: See Fact 2.18.9, Fact 5.14.23, Fact 5.14.24, and Fact 7.5.2.)

Fact 5.14.23. Let $A \in \mathbb{F}^{n \times n}$, assume that A is simple, let $B \in \mathbb{F}^{n \times n}$, and assume that $AB = BA$. Then, B is a polynomial in A whose degree is not greater than $n - 1$. (Proof: See [1490, p. 59].) (Remark: See Fact 5.14.22.)

Fact 5.14.24. Let $A, B \in \mathbb{F}^{n \times n}$. Then, B is a polynomial in A if and only if B commutes with every matrix that commutes with A . (Proof: See [711, p. 276].) (Remark: See Fact 4.8.13.) (Remark: See Fact 2.18.9, Fact 5.14.22, Fact 5.14.23, and Fact 7.5.2.)

Fact 5.14.25. Let $A, B \in \mathbb{C}^{n \times n}$, assume that $AB = BA$, let $x \in \mathbb{C}^n$ be an eigenvector of A with associated eigenvalue $\lambda \in \mathbb{C}$, and assume that $Bx \neq 0$. Then, Bx is an eigenvector of A with associated eigenvalue $\lambda \in \mathbb{C}$. (Proof: $A(Bx) = BAx = B(\lambda x) = \lambda(Bx)$.)

Fact 5.14.26. Let $A \in \mathbb{C}^{n \times n}$, and let $x \in \mathbb{C}^n$ be an eigenvector of A with associated eigenvalue λ . If A is nonsingular, then x is an eigenvector of A^A with associated eigenvalue $(\det A)/\lambda$. If $\text{rank } A = n - 1$, then x is an eigenvector of A^A with associated eigenvalue $\text{tr } A^A$ or 0. Finally, if $\text{rank } A \leq n - 2$, then x is an eigenvector of A^A with associated eigenvalue 0. (Proof: Use Fact 5.14.25 and the fact that $A^A A = A A^A$. See [354].) (Remark: See Fact 2.16.8 or Fact 6.3.6.)

Fact 5.14.27. Let $A, B \in \mathbb{C}^{n \times n}$. Then, the following statements are equivalent:

- i) $\cap_{k,l=1}^{n-1} \mathcal{N}([A^k, B^l]) \neq \{0\}$.
- ii) $\sum_{k,l=1}^{n-1} [A^k, B^l]^* [A^k, B^l]$ is singular.
- iii) A and B have a common eigenvector.

(Proof: See [547].) (Remark: This result is due to Shemesh.) (Remark: See Fact 5.17.1.)

Fact 5.14.28. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that $AB = BA$. Then, there exists a nonzero vector $x \in \mathbb{C}^n$ that is an eigenvector of both A and B . (Proof: See [709, p. 51].)

Fact 5.14.29. Let $A, B \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) Assume that A and B are Hermitian. Then, AB is Hermitian if and only if $AB = BA$.
- ii) A is normal if and only if, for all $C \in \mathbb{F}^{n \times n}$, $AC = CA$ implies that $A^*C = CA^*$.
- iii) Assume that B is Hermitian and $AB = BA$. Then, $A^*B = BA^*$.
- iv) Assume that A and B are normal and $AB = BA$. Then, AB is normal.
- v) Assume that A , B , and AB are normal. Then, BA is normal.
- vi) Assume that A and B are normal and either A or B has the property that distinct eigenvalues have unequal absolute values. Then, AB is normal if and only if $AB = BA$.

(Proof: See [358, 1428], [630, p. 157], and [1098, p. 102].)

Fact 5.14.30. Let $A, B, C \in \mathbb{F}^{n \times n}$, and assume that A and B are normal and $AC = CB$. Then, $A^*C = CB^*$. (Proof: Consider $\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}$ and $\begin{bmatrix} 0 & C \\ 0 & 0 \end{bmatrix}$ in *ii*) of Fact 5.14.29. See [627, p. 104] or [630, p. 321].) (Remark: This result is the *Putnam-Fuglede theorem*.)

Fact 5.14.31. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is dissipative and B is range Hermitian. Then,

$$\text{ind } B = \text{ind } AB.$$

(Proof: See [189].)

Fact 5.14.32. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$. Then,

$$\max\{\text{ind } A, \text{ind } C\} \leq \text{ind} \begin{bmatrix} A & B \\ 0 & C \end{bmatrix} \leq \text{ind } A + \text{ind } C.$$

If C is nonsingular, then

$$\text{ind} \begin{bmatrix} A & B \\ 0 & C \end{bmatrix} = \text{ind } A,$$

whereas, if A is nonsingular, then

$$\text{ind} \begin{bmatrix} A & B \\ 0 & C \end{bmatrix} = \text{ind } C.$$

(Proof: See [265, 999].) (Remark: See Fact 6.6.13.) (Remark: The eigenstructure of a partitioned Hamiltonian matrix is considered in Fact 12.23.1.)

Fact 5.14.33. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that A and B are skew symmetric. Then, there exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that

$$A = S \begin{bmatrix} 0_{(n-l) \times (n-l)} & A_{12} \\ -A_{12}^T & A_{22} \end{bmatrix} S^T$$

and

$$B = S \begin{bmatrix} B_{11} & B_{12} \\ -B_{12}^T & 0_{l \times l} \end{bmatrix} S^T,$$

where $l \triangleq \lfloor n/2 \rfloor$. Consequently,

$$\text{mspec}(AB) = \text{mspec}(-A_{12}B_{12}^T) \cup \text{mspec}(-A_{12}^TB_{12}),$$

and thus every nonzero eigenvalue of AB has even algebraic multiplicity. (Proof: See [30].)

Fact 5.14.34. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that A and B are skew symmetric. If n is even, then there exists a monic polynomial p of degree $n/2$ such that $\chi_{AB}(s) = p^2(s)$ and $p(AB) = 0$. If n is odd, then there exists a monic polynomial $p(s)$ of degree $(n-1)/2$ such that $\chi_{AB}(s) = sp^2(s)$ and $ABp(AB) = 0$. Consequently, if n is (even, odd), then χ_{AB} is (even, odd) and (every, every nonzero) eigenvalue of AB has even algebraic multiplicity and geometric multiplicity of at least 2. (Proof: See [418, 578].)

Fact 5.14.35. Let $q(t)$ denote the displacement of a mass $m > 0$ connected to a spring $k \geq 0$ and dashpot $c \geq 0$ and subject to a force $f(t)$. Then, $q(t)$ satisfies

$$m\ddot{q}(t) + c\dot{q}(t) + kq(t) = f(t)$$

or

$$\ddot{q}(t) + \frac{c}{m}\dot{q}(t) + \frac{k}{m}q(t) = \frac{1}{m}f(t).$$

Now, define the *natural frequency* $\omega_n \triangleq \sqrt{k/m}$ and, if $k > 0$, the *damping ratio* $\zeta \triangleq c/2\sqrt{km}$ to obtain

$$\ddot{q}(t) + 2\zeta\omega_n\dot{q}(t) + \omega_n^2q(t) = \frac{1}{m}f(t).$$

If $k = 0$, then set $\omega_n = 0$ and $\zeta\omega_n = c/2m$. Next, define $x_1(t) \triangleq q(t)$ and $x_2(t) \triangleq \dot{q}(t)$ so that this equation can be written as

$$\begin{bmatrix} \dot{x}_1(t) \\ \dot{x}_2(t) \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -\omega_n^2 & -2\zeta\omega_n \end{bmatrix} \begin{bmatrix} x_1(t) \\ x_2(t) \end{bmatrix} + \begin{bmatrix} 0 \\ 1/m \end{bmatrix} f(t).$$

The eigenvalues of the companion matrix $A_c \triangleq \begin{bmatrix} 0 & 1 \\ -\omega_n^2 & -2\zeta\omega_n \end{bmatrix}$ are given by

$$\text{mspec}(A_c) = \begin{cases} \{-\zeta\omega_n - j\omega_d, -\zeta\omega_n + j\omega_d\}_{\text{ms}}, & 0 \leq \zeta \leq 1, \\ \{(-\zeta - \sqrt{\zeta^2 - 1})\omega_n, (-\zeta + \sqrt{\zeta^2 - 1})\omega_n\}, & \zeta > 1, \end{cases}$$

where $\omega_d \triangleq \omega_n\sqrt{1 - \zeta^2}$ is the *damped natural frequency*. The matrix A_c has repeated eigenvalues in exactly two cases, namely,

$$\text{mspec}(A_c) = \begin{cases} \{0, 0\}_{\text{ms}}, & \omega_n = 0, \\ \{-\omega_n, -\omega_n\}_{\text{ms}}, & \zeta = 1. \end{cases}$$

In both of these cases the matrix A_c is defective. In the case $\omega_n = 0$, the matrix A_c is also in Jordan form, while, in the case $\zeta = 1$, it follows that $A_c = S A_J S^{-1}$, where $S \triangleq \begin{bmatrix} -1 & 0 \\ \omega_n & -1 \end{bmatrix}$ and A_J is the Jordan form matrix $A_J \triangleq \begin{bmatrix} -\omega_n & 1 \\ 0 & -\omega_n \end{bmatrix}$. If A_c is not defective, that is, if $\omega_n \neq 0$ and $\zeta \neq 1$, then the Jordan form A_J of A_c is given by

$$A_J \triangleq \begin{cases} \begin{bmatrix} -\zeta\omega_n + j\omega_d & 0 \\ 0 & -\zeta\omega_n - j\omega_d \end{bmatrix}, & 0 \leq \zeta < 1, \omega_n \neq 0, \\ \begin{bmatrix} (-\zeta - \sqrt{\zeta^2 - 1})\omega_n & 0 \\ 0 & (-\zeta + \sqrt{\zeta^2 - 1})\omega_n \end{bmatrix}, & \zeta > 1, \omega_n \neq 0. \end{cases}$$

In the case $0 \leq \zeta < 1$ and $\omega_n \neq 0$, define the real normal form

$$A_n \triangleq \begin{bmatrix} -\zeta\omega_n & \omega_d \\ -\omega_d & -\zeta\omega_n \end{bmatrix}.$$

The matrices A_c , A_J , and A_n are related by the similarity transformations

$$A_c = S_1 A_J S_1^{-1} = S_2 A_n S_2^{-1}, \quad A_J = S_3 A_n S_3^{-1},$$

where

$$\begin{aligned} S_1 &\triangleq \begin{bmatrix} 1 & 1 \\ -\zeta\omega_n + j\omega_d & -\zeta\omega_n - j\omega_d \end{bmatrix}, & S_1^{-1} &= \frac{j}{2\omega_d} \begin{bmatrix} -\zeta\omega_n - j\omega_d & -1 \\ \zeta\omega_n - j\omega_d & 1 \end{bmatrix}, \\ S_2 &\triangleq \frac{1}{\omega_d} \begin{bmatrix} 1 & 0 \\ -\zeta\omega_n & \omega_d \end{bmatrix}, & S_2^{-1} &= \begin{bmatrix} \omega_d & 0 \\ \zeta\omega_n & 1 \end{bmatrix}, \\ S_3 &\triangleq \frac{1}{2\omega_d} \begin{bmatrix} 1 & -j \\ 1 & j \end{bmatrix}, & S_3^{-1} &= \omega_d \begin{bmatrix} 1 & 1 \\ j & -j \end{bmatrix}. \end{aligned}$$

In the case $\zeta > 1$ and $\omega_n \neq 0$, the matrices A_c and A_J are related by

$$A_c = S_4 A_J S_4^{-1},$$

where

$$S_4 \triangleq \begin{bmatrix} 1 & 1 \\ -\zeta\omega_n + j\omega_d & -\zeta\omega_n - j\omega_d \end{bmatrix}, \quad S_4^{-1} = \frac{j}{2\omega_d} \begin{bmatrix} -\zeta\omega_n - j\omega_d & -1 \\ \zeta\omega_n - j\omega_d & 1 \end{bmatrix}.$$

Finally, define the energy-coordinates matrix

$$A_e \triangleq \begin{bmatrix} 0 & \omega_n \\ -\omega_n & -2\zeta\omega_n \end{bmatrix}.$$

Then, $A_e = S_5 A_c S_5^{-1}$, where

$$S_5 \triangleq \sqrt{\frac{m}{2}} \begin{bmatrix} 1 & 0 \\ 0 & 1/\omega_n \end{bmatrix}.$$

(Remark: m and k are not necessarily integers here.)

5.15 Facts on Matrix Factorizations

Fact 5.15.1. Let $A \in \mathbb{F}^{n \times n}$. Then, A is normal if and only if there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that $A^* = AS$. (Proof: See [1098, pp. 102, 113].)

Fact 5.15.2. Let $A \in \mathbb{C}^{n \times n}$. Then, there exists a nonsingular matrix $S \in \mathbb{C}^{n \times n}$ such that SAS^{-1} is symmetric. (Proof: See [709, p. 209].) (Remark: The symmetric matrix is a *complex symmetric Jordan form*.) (Remark: See Corollary 5.3.8.) (Remark: The coefficient of the last matrix in [709, p. 209] should be $j/2$.)

Fact 5.15.3. Let $A \in \mathbb{C}^{n \times n}$, and assume that A^2 is normal. Then, the following statements hold:

- i) There exists a unitary matrix $S \in \mathbb{C}^{n \times n}$ such that SAS^{-1} is symmetric.
- ii) There exists a symmetric unitary matrix $S \in \mathbb{C}^{n \times n}$ such that $A^T = SAS^{-1}$.

(Proof: See [1375].)

Fact 5.15.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then, A^{-1} and A^* are similar if and only if there exists a nonsingular matrix $B \in \mathbb{F}^{n \times n}$ such that $A = B^{-1}B^*$. Furthermore, A is unitary if and only if there exists a normal,

nonsingular matrix $B \in \mathbb{F}^{n \times n}$ such that $A = B^{-1}B^*$. (Proof: See [398]. Sufficiency in the second statement follows from Fact 3.11.4.)

Fact 5.15.5. Let $A \in \mathbb{F}^{m \times m}$ and $B \in \mathbb{F}^{n \times n}$. Then, there exist matrices $C \in \mathbb{F}^{m \times n}$ and $D \in \mathbb{F}^{n \times m}$ such that $A = CD$ and $B = DC$ if and only if the following statements hold:

- i) The Jordan blocks associated with nonzero eigenvalues are identical in A and B .
- ii) Let $n_1 \geq n_2 \geq \dots \geq n_r$ denote the orders of the Jordan blocks of A associated with $0 \in \text{spec}(A)$, and let $m_1 \geq m_2 \geq \dots \geq m_r$ denote the orders of the Jordan blocks of B associated with $0 \in \text{spec}(B)$, where $n_i = 0$ or $m_i = 0$ as needed. Then, $|n_i - m_i| \leq 1$ for all $i = 1, \dots, r$.

(Proof: See [771].) (Remark: See Fact 5.15.6.)

Fact 5.15.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are nonsingular. Then, A and B are similar if and only if there exist nonsingular matrices $C, D \in \mathbb{F}^{n \times n}$ such that $A = CD$ and $B = DC$. (Proof: Sufficiency follows from Fact 5.10.11. Necessity is a special case of Fact 5.15.5.)

Fact 5.15.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are nonsingular. Then, $\det A = \det B$ if and only if there exist nonsingular matrices $C, D, E \in \mathbb{R}^{n \times n}$ such that $A = CDE$ and $B = EDC$. (Remark: This result is due to Shoda and Taussky-Todd. See [258].)

Fact 5.15.8. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist matrices $B, C \in \mathbb{F}^{n \times n}$ such that B is unitary, C is upper triangular, and $A = BC$. If, in addition, A is nonsingular, then there exist unique matrices $B, C \in \mathbb{F}^{n \times n}$ such that B is unitary, C is upper triangular with positive diagonal entries, and $A = BC$. (Proof: See [709, p. 112] or [1129, p. 362].) (Remark: This result is the *QR decomposition*. The orthogonal matrix B is constructed as a product of elementary reflectors.)

Fact 5.15.9. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = m$. Then, there exist a unique matrix $B \in \mathbb{F}^{n \times m}$ and a matrix $C \in \mathbb{F}^{m \times m}$ such that $B^*B = I_m$, C is upper triangular with positive diagonal entries, and $A = BC$. (Proof: See [709, p. 15] or [1129, p. 206].) (Remark: $C \in \text{UT}_+(n)$. See Fact 3.21.5.) (Remark: This factorization is a consequence of *Gram-Schmidt orthonormalization*.)

Fact 5.15.10. Let $A \in \mathbb{F}^{n \times n}$, let $r \triangleq \text{rank } A$, and assume that the first r leading principal subdeterminants of A are nonzero. Then, there exist matrices $B, C \in \mathbb{F}^{n \times n}$ such that B is lower triangular, C is upper triangular, and $A = BC$. Either B or C can be chosen to be nonsingular. Furthermore, both B and C are nonsingular if and only if A is nonsingular. (Proof: See [709, p. 160].) (Remark: This result is the *LU decomposition*.) (Remark: All LU factorizations of a singular matrix are characterized in [424].)

Fact 5.15.11. Let $\theta \in (-\pi, \pi)$. Then,

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} = \begin{bmatrix} 1 & -\tan(\theta/2) \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ \sin \theta & 1 \end{bmatrix} \begin{bmatrix} 1 & -\tan(\theta/2) \\ 0 & 1 \end{bmatrix}.$$

(Remark: This result is a *ULU factorization* involving three *shear* factors. The matrix $-I_2$ requires four factors. In general, all factors may be different. See [1240, 1311].)

Fact 5.15.12. Let $A \in \mathbb{F}^{n \times n}$. Then, A is nonsingular if and only if A is the product of elementary matrices. (Problem: How many factors are needed?)

Fact 5.15.13. Let $A \in \mathbb{F}^{n \times n}$, assume that A is a projector, and let $r \triangleq \text{rank } A$. Then, there exist nonzero vectors $x_1, \dots, x_{n-r} \in \mathbb{F}^n$ such that $x_i^* x_j = 0$ for all $i \neq j$ and such that

$$A = \prod_{i=1}^{n-r} [I - (x_i^* x_i)^{-1} x_i x_i^*].$$

(Remark: Every projector is the product of mutually orthogonal elementary projectors.) (Proof: A is unitarily similar to $\text{diag}(1, \dots, 1, 0, \dots, 0)$, which can be written as the product of elementary projectors.)

Fact 5.15.14. Let $A \in \mathbb{F}^{n \times n}$. Then, A is a reflector if and only if there exist $m \leq n$ nonzero vectors $x_1, \dots, x_m \in \mathbb{F}^n$ such that $x_i^* x_j = 0$ for all $i \neq j$ and such that

$$A = \prod_{i=1}^m [I - 2(x_i^* x_i)^{-1} x_i x_i^*].$$

In this case, m is the algebraic multiplicity of $-1 \in \text{spec}(A)$. (Remark: Every reflector is the product of mutually orthogonal elementary reflectors.) (Proof: A is unitarily similar to $\text{diag}(\pm 1, \dots, \pm 1)$, which can be written as the product of elementary reflectors.)

Fact 5.15.15. Let $A \in \mathbb{R}^{n \times n}$. Then, A is orthogonal if and only if there exist $m \in \mathbb{P}$ and nonzero vectors $x_1, \dots, x_m \in \mathbb{R}^n$ such that $\det A = (-1)^m$ and

$$A = \prod_{i=1}^m [I - 2(x_i^T x_i)^{-1} x_i x_i^T].$$

(Remark: Every orthogonal matrix is the product of elementary reflectors. This factorization is a result of Cartan and Dieudonné. See [103, p. 24] and [1168, 1354]. The minimal number of factors is unsettled. See Fact 3.14.4 and Fact 3.9.5. The complex case is open.)

Fact 5.15.16. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$. Then, A is orthogonal and $\det A = 1$ if and only if there exist $m \in \mathbb{P}$ such that $1 \leq m \leq n(n-1)/2$, $\theta_1, \dots, \theta_m \in \mathbb{R}$, and $j_1, \dots, j_m, k_1, \dots, k_m \in \{1, \dots, n\}$ such that

$$A = \prod_{i=1}^m P(\theta_i, j_i, k_i),$$

where

$$P(\theta, j, k) \triangleq I_n + [(\cos \theta) - 1](E_{j,j} + E_{k,k}) + (\sin \theta)(E_{j,k} - E_{k,j}).$$

(Proof: See [471].) (Remark: $P(\theta, j, k)$ is a *plane* or *Givens rotation*. See Fact 3.9.5.) (Remark: Suppose that $\det A = -1$, and let $B \in \mathbb{R}^{n \times n}$ be an elementary reflector. Then, $AB \in \text{SO}(n)$. Therefore, the factorization given above holds with an additional elementary reflector.) (Problem: Generalize this result to $\mathbb{C}^{n \times n}$.) (Remark: See [887].)

Fact 5.15.17. Let $A \in \mathbb{F}^{n \times n}$. Then, $A^{2*}A = A^*A^2$ if and only if there exist a projector $B \in \mathbb{F}^{n \times n}$ and a Hermitian matrix $C \in \mathbb{F}^{n \times n}$ such that $A = BC$. (Proof: See [1114].)

Fact 5.15.18. Let $A \in \mathbb{R}^{n \times n}$. Then, $|\det A| = 1$ if and only if A is the product of $n + 2$ or fewer involutory matrices that have exactly one negative eigenvalue. In addition, the following statements hold:

- i) If $n = 2$, then 3 or fewer factors are needed.
- ii) If $A \neq \alpha I$ for all $\alpha \in \mathbb{R}$ and $\det A = (-1)^n$, then n or fewer factors are needed.
- iii) If $\det A = (-1)^{n+1}$, then $n + 1$ or fewer factors are needed.

(Proof: See [298, 1112].) (Remark: The minimal number of factors for a unitary matrix A is given in [417].)

Fact 5.15.19. Let $A \in \mathbb{C}^{n \times n}$, and define $r_0 \triangleq n$ and $r_k \triangleq \text{rank } A^k$ for all $k = 1, 2, \dots$. Then, there exists a matrix $B \in \mathbb{C}^{n \times n}$ such that $A = B^2$ if and only if the sequence $(r_k - r_{k+1})_{k=0}^{\infty}$ does not contain two elements that are the same odd integer and, if $r_0 - r_1$ is odd, then $r_0 + r_2 \geq 1 + 2r_1$. Now, assume that $A \in \mathbb{R}^{n \times n}$. Then, there exists $B \in \mathbb{R}^{n \times n}$ such that $A = B^2$ if and only if the above condition holds and, for every negative eigenvalue λ of A and for every positive integer k , the Jordan form of A has an even number of $k \times k$ blocks associated with λ . (Proof: See [711, p. 472].) (Remark: See Fact 11.18.36.) (Remark: For all $l \geq 2$, $A \triangleq N_l$ does not have a square root.) (Remark: Uniqueness is discussed in [769]. Square roots of A that are functions of A are defined in [678].) (Remark: The principal square root is considered in Theorem 10.6.1.) (Remark: m th roots are considered in [329, 683, 1101, 1263].)

Fact 5.15.20. Let $A \in \mathbb{C}^{n \times n}$, and assume that A is group invertible. Then, there exists $B \in \mathbb{C}^{n \times n}$ such that $A = B^2$.

Fact 5.15.21. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular and has no negative eigenvalues. Furthermore, define $(P_k)_{k=0}^{\infty} \subset \mathbb{F}^{n \times n}$ and $(Q_k)_{k=0}^{\infty} \subset \mathbb{F}^{n \times n}$ by

$$P_0 \triangleq A, \quad Q_0 \triangleq I,$$

and, for all $k \geq 1$,

$$P_{k+1} \triangleq \frac{1}{2}(P_k + Q_k^{-1}),$$

$$Q_{k+1} \triangleq \frac{1}{2}(Q_k + P_k^{-1}).$$

Then,

$$B \triangleq \lim_{k \rightarrow \infty} P_k$$

exists, satisfies $B^2 = A$, and is the unique square root of A satisfying $\text{spec}(B) \subset \text{ORHP}$. Furthermore,

$$\lim_{k \rightarrow \infty} Q_k = A^{-1}.$$

(Proof: See [397, 677].) (Remark: All indicated inverses exist.) (Remark: This sequence is related to Newton's iteration for the matrix sign function. See Fact 10.10.2.) (Remark: See Fact 8.9.32.)

Fact 5.15.22. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $r \triangleq \text{rank } A$. Then, there exists $B \in \mathbb{F}^{n \times r}$ such that $A = BB^*$.

Fact 5.15.23. Let $A \in \mathbb{F}^{n \times n}$, and let $k \geq 1$. Then, there exists a unique matrix $B \in \mathbb{F}^{n \times n}$ such that

$$A = B(B^*B)^k.$$

(Proof: See [1091].)

Fact 5.15.24. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist symmetric matrices $B, C \in \mathbb{F}^{n \times n}$, one of which is nonsingular, such that $A = BC$. (Proof: See [1098, p. 82].) (Remark: Note that

$$\begin{bmatrix} \beta_1 & \beta_2 & 1 \\ \beta_2 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -\beta_0 & -\beta_1 & -\beta_2 \end{bmatrix} = \begin{bmatrix} -\beta_0 & 0 & 0 \\ 0 & \beta_2 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

and use Theorem 5.2.3.) (Remark: This result is due to Frobenius. The identity is a *Bezout matrix factorization*; see Fact 4.8.6. See [240, 241, 628].) (Remark: B and C are symmetric for $\mathbb{F} = \mathbb{C}$.)

Fact 5.15.25. Let $A \in \mathbb{C}^{n \times n}$. Then, $\det A$ is real if and only if A is the product of four Hermitian matrices. Furthermore, four is the smallest number of factors in general. (Proof: See [1459].)

Fact 5.15.26. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) A is the product of two positive-semidefinite matrices if and only if A is similar to a positive-semidefinite matrix.
- ii) If A is nilpotent, then A is the product of three positive-semidefinite matrices.
- iii) If A is singular, then A is the product of four positive-semidefinite matrices.
- iv) $\det A > 0$ and $A \neq \alpha I$ for all $\alpha \leq 0$ if and only if A is the product of four positive-definite matrices.

v) $\det A > 0$ if and only if A is the product of five positive-definite matrices.

(Proof: [117, 628, 1458, 1459].) (Remark: See [1459] for factorizations of complex matrices and operators.) (Example for v):

$$\begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 2 & 0 \\ 0 & 1/2 \end{bmatrix} \begin{bmatrix} 5 & 7 \\ 7 & 10 \end{bmatrix} \begin{bmatrix} 13/2 & -5 \\ -5 & 4 \end{bmatrix} \begin{bmatrix} 8 & 5 \\ 5 & 13/4 \end{bmatrix} \begin{bmatrix} 25/8 & -11/2 \\ -11/2 & 10 \end{bmatrix}.$$

Fact 5.15.27. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) $A = BC$, where $B \in \mathbf{S}^n$ and $C \in \mathbf{N}^n$, if and only if A^2 is diagonalizable over $\mathbb{R}$ and $\text{spec}(A) \subset [0, \infty)$.
- ii) $A = BC$, where $B \in \mathbf{S}^n$ and $C \in \mathbf{P}^n$, if and only if A is diagonalizable over $\mathbb{R}$.
- iii) $A = BC$, where $B, C \in \mathbf{N}^n$, if and only if $A = DE$, where $D \in \mathbf{N}^n$ and $E \in \mathbf{P}^n$.
- iv) $A = BC$, where $B \in \mathbf{N}^n$ and $C \in \mathbf{P}^n$, if and only if A is diagonalizable over $\mathbb{R}$ and $\text{spec}(A) \subset [0, \infty)$.
- v) $A = BC$, where $B, C \in \mathbf{P}^n$, if and only if A is diagonalizable over $\mathbb{R}$ and $\text{spec}(A) \subset [0, \infty)$.

(Proof: See [706, 1453, 1458].)

Fact 5.15.28. Let $A \in \mathbb{F}^{n \times n}$. Then, A is singular or the identity if and only if A is the product of n or fewer idempotent matrices in $\mathbb{F}^{n \times n}$, each of whose rank is equal to $\text{rank } A$. Furthermore, $\text{rank}(A - I) \leq k \text{def } A$, where $k \geq 1$, if and only if A is the product of k idempotent matrices. (Examples:

$$\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1/2 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & 1/2 \\ 0 & 1 \end{bmatrix}$$

and

$$\begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}.)$$

(Proof: See [71, 125, 378, 460].)

Fact 5.15.29. Let $A \in \mathbb{R}^{n \times n}$, assume that A is singular, and assume that A is not a 2×2 nilpotent matrix. Then, there exist nilpotent matrices $B, C \in \mathbb{R}^{n \times n}$ such that $A = BC$ and $\text{rank } A = \text{rank } B = \text{rank } C$. (Proof: See [1215, 1457]. See also [1248].)

Fact 5.15.30. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, there exist $B, C \in \mathbb{F}^{n \times n}$ such that B is positive definite, C is positive semidefinite, and $A = BC$. (Proof: See [1324].)

Fact 5.15.31. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is nonsingular. Then, A is similar to A^{-1} if and only if A is the product of two involutory matrices. If, in addition, A is orthogonal, then A is the product of two reflectors. (Proof: See [123, 414, 1451, 1452] or [1098, p. 108].) (Problem: Construct these reflectors for $A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$.)

Fact 5.15.32. Let $A \in \mathbb{R}^{n \times n}$. Then, $|\det A| = 1$ if and only if A is the product of four or fewer involutory matrices. (Proof: [124, 611, 1214].)

Fact 5.15.33. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$. Then, A is the product of two commutators. (Proof: See [1459].)

Fact 5.15.34. Let $A \in \mathbb{R}^{n \times n}$, and assume that $\det A = 1$. Then, there exist nonsingular matrices $B, C \in \mathbb{R}^{n \times n}$ such that $A = BCB^{-1}C^{-1}$. (Proof: See [1191].) (Remark: The product is a *multiplicative commutator*. This result is due to Shoda.)

Fact 5.15.35. Let $A \in \mathbb{R}^{n \times n}$, assume that A is orthogonal, and assume that $\det A = 1$. Then, there exist reflectors $B, C \in \mathbb{R}^{n \times n}$ such that $A = BCB^{-1}C^{-1}$. (Proof: See [1268].)

Fact 5.15.36. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then, there exists an involutory matrix $B \in \mathbb{F}^{n \times n}$ and a symmetric matrix $C \in \mathbb{F}^{n \times n}$ such that $A = BC$. (Proof: See [577].)

Fact 5.15.37. Let $A \in \mathbb{F}^{n \times n}$, and assume that n is even. Then, the following statements are equivalent:

- i) A is the product of two skew-symmetric matrices.
- ii) Every elementary divisor of A has even algebraic multiplicity.
- iii) There exists a matrix $B \in \mathbb{F}^{n/2 \times n/2}$ such that A is similar to $\begin{bmatrix} B & 0 \\ 0 & B \end{bmatrix}$.

(Remark: In i) the factors are skew symmetric even when A is complex.) (Proof: See [578, 1459].)

Fact 5.15.38. Let $A \in \mathbb{C}^{n \times n}$, and assume that $n \geq 4$ and n is even. Then, A is the product of five skew-symmetric matrices in $\mathbb{C}^{n \times n}$. (Proof: See [857, 858].)

Fact 5.15.39. Let $A \in \mathbb{F}^{n \times n}$. Then, there exist a symmetric matrix $B \in \mathbb{F}^{n \times n}$ and a skew-symmetric matrix $C \in \mathbb{F}^{n \times n}$ such that $A = BC$ if and only if A is similar to $-A$. (Proof: See [1135].)

Fact 5.15.40. Let $A \in \mathbb{F}^{n \times m}$, and let $r \triangleq \text{rank } A$. Then, there exist matrices $B \in \mathbb{F}^{n \times r}$ and $C \in \mathbb{R}^{r \times m}$ such that $A = BC$ and $\text{rank } B = \text{rank } C = r$.

Fact 5.15.41. Let $A \in \mathbb{F}^{n \times n}$. Then, A is diagonalizable over $\mathbb{F}$ with (nonnegative, positive) eigenvalues if and only if there exist (positive-semidefinite, positive-definite) matrices $B, C \in \mathbb{F}^{n \times n}$ such that $A = BC$. (Proof: To prove sufficiency, use Theorem 8.3.5 and note that

$$A = S^{-1}(SBS^*)(S^{-*}CS^{-1})S.)$$

5.16 Facts on Companion, Vandermonde, and Circulant Matrices

Fact 5.16.1. Let $p \in \mathbb{F}[s]$, where $p(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_0$, and define $C_b(p), C_r(p), C_t(p), C_l(p) \in \mathbb{F}^{n \times n}$ by

$$C_b(p) \triangleq \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \ddots & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ -\beta_0 & -\beta_1 & -\beta_2 & \cdots & -\beta_{n-2} & -\beta_{n-1} \end{bmatrix},$$

$$C_r(p) \triangleq \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & -\beta_0 \\ 1 & 0 & 0 & \cdots & 0 & -\beta_1 \\ 0 & 1 & 0 & \cdots & 0 & -\beta_2 \\ \vdots & \ddots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \ddots & 0 & -\beta_{n-2} \\ 0 & 0 & 0 & \cdots & 1 & -\beta_{n-1} \end{bmatrix},$$

$$C_t(p) \triangleq \begin{bmatrix} -\beta_{n-1} & -\beta_{n-2} & \cdots & -\beta_2 & -\beta_1 & -\beta_0 \\ 1 & 0 & \cdots & 0 & 0 & 0 \\ \vdots & \ddots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \ddots & 0 & 0 & 0 \\ 0 & 0 & \ddots & 1 & 0 & 0 \\ 0 & 0 & \cdots & 0 & 1 & 0 \end{bmatrix},$$

$$C_l(p) \triangleq \begin{bmatrix} -\beta_{n-1} & 1 & \cdots & 0 & 0 & 0 \\ -\beta_{n-2} & 0 & \ddots & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ -\beta_2 & 0 & \cdots & 0 & 1 & 0 \\ -\beta_1 & 0 & \cdots & 0 & 0 & 1 \\ -\beta_0 & 0 & \cdots & 0 & 0 & 0 \end{bmatrix}.$$

Then,

$$C_r(p) = C_b^T(p), \quad C_l(p) = C_t^T(p),$$

$$C_t(p) = \hat{I}C_b(p)\hat{I}, \quad C_l(p) = \hat{I}C_r(p)\hat{I},$$

$$C_l(p) = C_b^{\hat{T}}(p), \quad C_t(p) = C_r^{\hat{T}}(p),$$

and

$$\chi_{C_b(p)} = \chi_{C_r(p)} = \chi_{C_t(p)} = \chi_{C_l(p)} = p.$$

Furthermore,

$$C_r(p) = SC_b(p)S^{-1}$$

and

$$C_l(p) = \hat{S}C_t(p)\hat{S}^{-1},$$

where $S, \hat{S} \in \mathbb{F}^{n \times n}$ are the Hankel matrices

$$S \triangleq \begin{bmatrix} \beta_1 & \beta_2 & \cdots & \beta_{n-1} & 1 \\ \beta_2 & \beta_3 & \ddots & 1 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \beta_{n-1} & 1 & \ddots & 0 & 0 \\ 1 & 0 & \cdots & 0 & 0 \end{bmatrix}$$

and

$$\hat{S} \triangleq \hat{I}S\hat{I} = \begin{bmatrix} 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \ddots & 1 & \beta_{n-1} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 1 & \ddots & \beta_3 & \beta_2 \\ 1 & \beta_{n-1} & \cdots & \beta_2 & \beta_1 \end{bmatrix}.$$

(Remark: $(C_b(p), C_r(p), C_t(p), C_l(p))$ are the (*bottom, right, top, left*) companion matrices. Note that $C_b(p) = C(p)$. See [144, p. 282] and [787, p. 659].) (Remark: $S = B(p, 1)$, where $B(p, 1)$ is a Bezout matrix. See Fact 4.8.6.)

Fact 5.16.2. Let $p \in \mathbb{F}[s]$, where $p(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_0$, assume that $\beta_0 \neq 0$, and let

$$C_b(p) \triangleq \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \ddots & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ -\beta_0 & -\beta_1 & -\beta_2 & \cdots & -\beta_{n-2} & -\beta_{n-1} \end{bmatrix}.$$

Then,

$$C_b^{-1}(p) = C_t(\hat{p}) = \begin{bmatrix} -\beta_1/\beta_0 & \cdots & -\beta_{n-2}/\beta_0 & -\beta_{n-1}/\beta_0 & -1/\beta_0 \\ 1 & \cdots & 0 & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & \cdots & 1 & 0 & 0 \\ 0 & \cdots & 0 & 1 & 0 \end{bmatrix},$$

where $\hat{p}(s) \triangleq \beta_0^{-1}s^n p(1/s)$. (Remark: See Fact 4.9.9.)

Fact 5.16.3. Let $\lambda_1, \dots, \lambda_n \in \mathbb{F}$, and define the *Vandermonde matrix* $V(\lambda_1, \dots, \lambda_n) \in \mathbb{F}^{n \times n}$ by

$$V(\lambda_1, \dots, \lambda_n) \triangleq \begin{bmatrix} 1 & 1 & \cdots & 1 \\ \lambda_1 & \lambda_2 & \cdots & \lambda_n \\ \lambda_1^2 & \lambda_2^2 & \cdots & \lambda_n^2 \\ \lambda_1^3 & \lambda_2^3 & \cdots & \lambda_n^3 \\ \vdots & \vdots & \ddots & \vdots \\ \lambda_1^{n-1} & \lambda_2^{n-1} & \cdots & \lambda_n^{n-1} \end{bmatrix}.$$

Then,

$$\det V(\lambda_1, \dots, \lambda_n) = \prod_{1 \leq i < j \leq n} (\lambda_i - \lambda_j).$$

Thus, $V(\lambda_1, \dots, \lambda_n)$ is nonsingular if and only if $\lambda_1, \dots, \lambda_n$ are distinct. (Remark: This result yields Proposition 4.5.4. Let $x_1, \dots, x_k$ be eigenvectors of $V(\lambda_1, \dots, \lambda_n)$ associated with distinct eigenvalues $\lambda_1, \dots, \lambda_k$ of $V(\lambda_1, \dots, \lambda_n)$. Suppose that $\alpha_1 x_1 + \cdots + \alpha_k x_k = 0$ so that $V^i(\lambda_1, \dots, \lambda_n)(\alpha_1 x_1 + \cdots + \alpha_k x_k) = \alpha_1 \lambda_1^i x_1 + \cdots + \alpha_k \lambda_k^i x_k = 0$ for all $i = 0, 1, \dots, k-1$. Let $X \triangleq [x_1 \cdots x_k] \in \mathbb{F}^{n \times k}$ and $D \triangleq \text{diag}(\alpha_1, \dots, \alpha_k)$. Then, $XD V^T(\lambda_1, \dots, \lambda_k) = 0$, which implies that $XD = 0$. Hence, $\alpha_i x_i = 0$ for all $i = 1, \dots, k$, and thus $\alpha_1 = \cdots = \alpha_k = 0$.) (Remark: Connections between the Vandermonde matrix and the Pascal matrix, *Stirling matrix*, *Bernoulli matrix*, *Bernstein matrix*, and companion matrices are discussed in [5]. See also Fact 11.11.4.)

Fact 5.16.4. Let $p \in \mathbb{F}[s]$, where $p(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_1 s + \beta_0$, and assume that p has distinct roots $\lambda_1, \dots, \lambda_n \in \mathbb{C}$. Then,

$$C(p) = V(\lambda_1, \dots, \lambda_n) \text{diag}(\lambda_1, \dots, \lambda_n) V^{-1}(\lambda_1, \dots, \lambda_n).$$

Consequently, for all $i = 1, \dots, n$, λ_i is an eigenvalue of $C(p)$ with associated eigenvector $\text{col}_i(V)$. Finally,

$$(VV^T)^{-1} C V V^T = C^T.$$

(Proof: See [139].) (Remark: Case in which $C(p)$ has repeated eigenvalues is considered in [139].)

Fact 5.16.5. Let $A \in \mathbb{F}^{n \times n}$. Then, A is cyclic if and only if A is similar to a companion matrix. (Proof: The result follows from Corollary 5.3.4. Alternatively,

let $\text{spec}(A) = \{\lambda_1, \dots, \lambda_r\}$ and $A = SBS^{-1}$, where $S \in \mathbb{C}^{n \times n}$ is nonsingular and $B = \text{diag}(B_1, \dots, B_r)$ is the Jordan form of A , where, for all $i = 1, \dots, r$, $B_i \in \mathbb{C}^{n_i \times n_i}$ and $\lambda_i, \dots, \lambda_i$ are the diagonal entries of B_i . Now, define $R \in \mathbb{C}^{n \times n}$ by $R \triangleq \begin{bmatrix} R_1 & \cdots & R_r \end{bmatrix} \in \mathbb{C}^{n \times n}$, where, for all $i = 1, \dots, r$, $R_i \in \mathbb{C}^{n \times n_i}$ is the matrix

$$R_i \triangleq \begin{bmatrix} 1 & 0 & \cdots & 0 \\ \lambda_i & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ \lambda_i^{n-2} & \binom{n-2}{1} \lambda_i^{n-3} & \cdots & \binom{n-2}{n_i-1} \lambda_i^{n-n_i-1} \\ \lambda_i^{n-1} & \binom{n-1}{1} \lambda_i^{n-2} & \cdots & \binom{n-1}{n_i-1} \lambda_i^{n-n_i} \end{bmatrix}.$$

Then, since $\lambda_1, \dots, \lambda_r$ are distinct, it follows that R is nonsingular. Furthermore, $C = RBR^{-1}$ is in companion form, and thus $A = SR^{-1}CRS$. If $n_i = 1$ for all $i = 1, \dots, r$, then R is a Vandermonde matrix. See Fact 5.16.3 and Fact 5.16.4.)

Fact 5.16.6. Let $\lambda_1, \dots, \lambda_n \in \mathbb{F}$ and, for $i = 1, \dots, n$, define

$$p_i(s) \triangleq \prod_{\substack{j=1 \\ j \neq i}}^n (s - \lambda_j).$$

Furthermore, define $A \in \mathbb{F}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} p_1(0) & \frac{1}{1!} p_1'(0) & \cdots & \frac{1}{(n-1)!} p_1^{(n-1)}(0) \\ \vdots & \ddots & \ddots & \vdots \\ p_n(0) & \frac{1}{1!} p_n'(0) & \cdots & \frac{1}{(n-1)!} p_n^{(n-1)}(0) \end{bmatrix}.$$

Then,

$$\text{diag}[p_1(\lambda_1), \dots, p_n(\lambda_n)] = AV(\lambda_1, \dots, \lambda_n).$$

(Proof: See [481, p. 159].)

Fact 5.16.7. Let $a_0, \dots, a_{n-1} \in \mathbb{F}$, and define $\text{circ}(a_0, \dots, a_{n-1}) \in \mathbb{F}^{n \times n}$ by

$$\text{circ}(a_0, \dots, a_{n-1}) \triangleq \begin{bmatrix} a_0 & a_1 & a_2 & \cdots & a_{n-2} & a_{n-1} \\ a_{n-1} & a_0 & a_1 & \cdots & a_{n-3} & a_{n-2} \\ a_{n-2} & a_{n-1} & a_0 & \ddots & a_{n-4} & a_{n-3} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ a_2 & a_3 & a_4 & \ddots & a_0 & a_1 \\ a_1 & a_2 & a_3 & \cdots & a_{n-1} & a_0 \end{bmatrix}.$$

A matrix of this form is *circulant*. Furthermore, for $n \geq 2$, define the $n \times n$ *primary circulant*

$$P_n \triangleq \text{circ}(0, 1, 0, \dots, 0) \triangleq \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ 0 & 0 & 1 & \ddots & 0 & 0 \\ 0 & 0 & 0 & \ddots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \ddots & 0 & 1 \\ 1 & 0 & 0 & \cdots & 0 & 0 \end{bmatrix}.$$

Define $P_1 \triangleq 1$. Finally, define $p(s) \triangleq a_{n-1}s^{n-1} + \cdots + a_1s + a_0 \in \mathbb{F}[s]$, and let $\theta \triangleq e^{2\pi j/n}$. Then, the following statements hold:

- i) $p(P_n) = \text{circ}(a_0, \dots, a_{n-1})$.
- ii) $P_n = C(q)$, where $q \in \mathbb{F}[s]$ is defined by $q(s) \triangleq s^n - 1$.
- iii) $\text{spec}(P_n) = \{1, \theta, \theta^2, \dots, \theta^{n-1}\}$.
- iv) $\det P_n = (-1)^{n-1}$.
- v) $\text{mspec}[\text{circ}(a_0, \dots, a_{n-1})] = \{p(1), p(\theta), p(\theta^2), \dots, p(\theta^{n-1})\}_{\text{ms}}$.
- vi) If $A, B \in \mathbb{F}^{n \times n}$ are circulant, then $AB = BA$ and AB is circulant.
- vii) If A is circulant, then $\overline{A}$, A^T , and A^* are circulant.
- viii) If A is circulant and $k \geq 0$, then A^k is circulant.
- ix) If A is nonsingular and circulant, then A^{-1} is circulant.
- x) $A \in \mathbb{F}^{n \times n}$ is circulant if and only if $A = P_n A P_n^T$.
- xi) P_n is an orthogonal matrix, and $P_n^n = I_n$.
- xii) If $A \in \mathbb{F}^{n \times n}$ is circulant, then A is reverse symmetric, Toeplitz, and normal.
- xiii) If $A \in \mathbb{F}^{n \times n}$ is circulant and nonzero, then A is irreducible.
- xiv) $A \in \mathbb{F}^{n \times n}$ is normal if and only if A is unitarily similar to a circulant matrix.

Next, define the *Fourier matrix* $S \in \mathbb{C}^{n \times n}$ by

$$S \triangleq n^{-1/2} V(1, \theta, \dots, \theta^{n-1}) = \frac{1}{\sqrt{n}} \begin{bmatrix} 1 & 1 & 1 & \cdots & 1 \\ 1 & \theta & \theta^2 & \cdots & \theta^{n-1} \\ 1 & \theta^2 & \theta^4 & \cdots & \theta^{n-2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & \theta^{n-1} & \theta^{n-2} & \cdots & \theta \end{bmatrix}.$$

Then, the following statements hold:

- xv) S is symmetric and unitary, but not Hermitian.
- xvi) $S^4 = I_n$.
- xvii) $\text{spec}(S) \subseteq \{1, -1, j, -j\}$.

xviii) $\operatorname{Re} S$ and $\operatorname{Im} S$ are symmetric, commute, and satisfy

$$(\operatorname{Re} S)^2 + (\operatorname{Im} S)^2 = I_n.$$

$$xix) S^{-1}P_n S = \operatorname{diag}(1, \theta, \dots, \theta^{n-1}).$$

$$xx) S^{-1}\operatorname{circ}(a_0, \dots, a_{n-1})S = \operatorname{diag}[p(1), p(\theta), \dots, p(\theta^{n-1})].$$

(Proof: See [16, pp. 81–98], [377, p. 81], and [1490, pp. 106–110].) (Remark: Circulant matrices play a role in digital signal processing, specifically, in the efficient implementation of the *fast Fourier transform*. See [997, pp. 356–380], [1142], and [1361, pp. 206, 207].) (Remark: S is a *Fourier matrix* and a Vandermonde matrix.) (Remark: If a real Toeplitz matrix is normal, then it must be either symmetric, skew symmetric, circulant, or skew circulant. See [72, 472]. A unified treatment of the solutions of quadratic, cubic, and quartic equations using circulant matrices is given in [788].) (Remark: The set $\{I, P_k, P_k^2, \dots, P_k^{k-1}\}$ is a group. See Fact 3.21.8 and Fact 3.21.9.) (Remark: Circulant matrices are generalized by *cycle matrices*, which correspond to visual geometric symmetries. See [548].)

Fact 5.16.8. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is a permutation matrix. Then, there exists a permutation matrix $S \in \mathbb{R}^{n \times n}$ such that

$$A = S \operatorname{diag}(P_{n_1}, \dots, P_{n_r}) S^{-1},$$

and, for all $i = 1, \dots, r$, $P_{n_i} \in \mathbb{R}^{n_i \times n_i}$ is a primary circulant (see Fact 5.16.7.) Furthermore, the primary circulants $P_{n_1}, \dots, P_{n_r}$ are unique up to a relabeling. Consequently,

$$\operatorname{mspec}(A) = \bigcup_{i=1}^r \{1, \theta_i, \dots, \theta_i^{n_i-1}\}_{\operatorname{ms}},$$

where $\theta_i \triangleq e^{2\pi j/n_i}$. Hence,

$$\det A = (-1)^{n-r}.$$

Finally, the smallest positive integer m such that $A^m = I$ is given by the least common multiple of $n_1, \dots, n_r$. (Proof: See [377, p. 29]. The last statement follows from [445, pp. 32, 33].) (Remark: This result provides a canonical form for permutation matrices under unitary similarity with a permutation matrix.) (Remark: It follows that A can be written as the product

$$A = S \begin{bmatrix} P_{n_1} & 0 \\ 0 & I \end{bmatrix} \cdots \begin{bmatrix} I & 0 & 0 \\ 0 & P_{n_i} & 0 \\ 0 & 0 & I \end{bmatrix} \cdots \begin{bmatrix} I & 0 \\ 0 & P_{n_r} \end{bmatrix} S^{-1},$$

where the factors represent disjoint cycles. The factorization reveals the *cycle decomposition* for an element of the permutation group S_n on a set having n elements, where S_n is represented by the group of $n \times n$ permutation matrices. See [445, pp. 29–32], [1149, p. 18] and Fact 3.21.7.) (Remark: The number of possible canonical forms is given by p_n , where p_n is the number of integral partitions of n . For example, $p_1 = 1$, $p_2 = 2$, $p_3 = 3$, $p_4 = 5$, and $p_5 = 7$. For all n , p_n is given by the expansion

$$1 + \sum_{n=1}^{\infty} p_n x^n = \frac{1}{(1-x)(1-x^2)(1-x^3)\cdots}.$$

See [1503, pp. 210, 211].)

5.17 Facts on Simultaneous Transformations

Fact 5.17.1. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^{-1} and SBS^{-1} are upper triangular. Then, A and B have a common eigenvector with corresponding eigenvalues $(SAS^{-1})_{(1,1)}$ and $(SBS^{-1})_{(1,1)}$. (Proof: See [547].) (Remark: See Fact 5.14.27.)

Fact 5.17.2. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that $P_{A,B}$ is regular. Then, there exist unitary matrices $S_1, S_2 \in \mathbb{C}^{n \times n}$ such that S_1AS_2 and S_1BS_2 are upper triangular. (Proof: See [1230, p. 276].)

Fact 5.17.3. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that $P_{A,B}$ is regular. Then, there exist orthogonal matrices $S_1, S_2 \in \mathbb{R}^{n \times n}$ such that S_1AS_2 is upper triangular and S_1BS_2 is upper Hessenberg with 2×2 diagonally located blocks. (Proof: See [1230, p. 290].) (Remark: This result is due to Moler and Stewart.)

Fact 5.17.4. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$, and assume that $AB = BA$ for all $A, B \in \mathcal{S}$. Then, there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that, for all $A \in \mathcal{S}$, SAS^* is upper triangular. (Proof: See [709, p. 81] and [1113].) (Remark: See Fact 5.17.9.)

Fact 5.17.5. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that either

$$[A, [A, B]] = [B, [A, B]] = 0$$

or

$$\text{rank } [A, B] \leq 1.$$

Then, there exists a nonsingular matrix $S \in \mathbb{C}^{n \times n}$ such that SAS^{-1} and SBS^{-1} are upper triangular. (Proof: The first result is due to McCoy, and the second result is due to Laffey. See [547, 1113].)

Fact 5.17.6. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that A and B are idempotent. Then, there exists a unitary matrix $S \in \mathbb{C}^{n \times n}$ such that SAS^* and SBS^* are upper triangular if and only if $[A, B]$ is nilpotent. (Proof: See [1251].) (Remark: Necessity follows from Fact 3.17.11.) (Remark: See Fact 5.17.4.)

Fact 5.17.7. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$, and assume that every matrix $A \in \mathcal{S}$ is normal. Then, $AB = BA$ for all $A, B \in \mathcal{S}$ if and only if there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that, for all $A \in \mathcal{S}$, SAS^* is diagonal. (Remark: See Fact 8.16.1 and [709, pp. 103, 172].)

Fact 5.17.8. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$, and assume that every matrix $A \in \mathcal{S}$ is diagonalizable over $\mathbb{F}$. Then, $AB = BA$ for all $A, B \in \mathcal{S}$ if and only if there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that, for all $A \in \mathcal{S}$, SAS^{-1} is diagonal. (Proof: See [709, p. 52].)

Fact 5.17.9. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $\{x \in \mathbb{F}^n : x^*Ax = x^*Bx = 0\} = \{0\}$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and SBS^* are upper triangular. (Proof: See [1098, p. 96].) (Remark: A and B need not be Hermitian.) (Remark: See Fact 5.17.4 and Fact 8.16.6.) (Remark: Simultaneous triangularization by means of a unitary biequivalence transformation

is given in Proposition 5.7.3.)

5.18 Facts on the Polar Decomposition

Fact 5.18.1. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$(AA^*)^{1/2}A = A(A^*A)^{1/2}.$$

(Remark: See Fact 5.18.4.) (Remark: The positive-semidefinite square root is defined in (8.5.3).)

Fact 5.18.2. Let $A \in \mathbb{F}^{n \times m}$, where $n \leq m$. Then, there exist $M \in \mathbb{F}^{n \times n}$ and $S \in \mathbb{F}^{n \times m}$ such that M is positive semidefinite, S satisfies $SS^* = I_n$, and $A = MS$. Furthermore, M is given uniquely by $M = (AA^*)^{1/2}$. If, in addition, $\text{rank } A = n$, then S is given uniquely by

$$S = (AA^*)^{-1/2}A = \frac{2}{\pi}A^* \int_0^\infty (t^2I + AA^*)^{-1} dt.$$

(Proof: See [683, Chapter 8].)

Fact 5.18.3. Let $A \in \mathbb{F}^{n \times m}$, where $m \leq n$. Then, there exist $M \in \mathbb{F}^{m \times m}$ and $S \in \mathbb{F}^{n \times m}$ such that M is positive semidefinite, S satisfies $S^*S = I_m$, and $A = SM$. Furthermore, M is given uniquely by $M = (A^*A)^{1/2}$. If, in addition, $\text{rank } A = m$, then M is positive definite and S is given uniquely by

$$S = A(A^*A)^{-1/2} = \frac{2}{\pi}A \int_0^\infty (t^2I + A^*A)^{-1} dt.$$

(Proof: See [683, Chapter 8].)

Fact 5.18.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then, there exist unique matrices $M, S \in \mathbb{F}^{n \times n}$ such that $A = MS$, M is positive definite, and S is unitary. In particular, $M = (AA^*)^{1/2}$ and $S = (AA^*)^{-1/2}A$. (Remark: See Fact 5.18.1.)

Fact 5.18.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then, there exist unique matrices $M, S \in \mathbb{F}^{n \times n}$ such that $A = SM$, M is positive definite, and S is unitary. In particular, $M = (A^*A)^{1/2}$ and $S = (AA^*)^{-1/2}A$.

Fact 5.18.6. Let $M_1, M_2 \in \mathbb{F}^{n \times n}$, assume that M_1, M_2 are positive definite, let $S_1, S_2 \in \mathbb{F}^{n \times n}$, assume that S_1, S_2 are unitary, and assume that $M_1S_1 = S_2M_2$. Then, $S_1 = S_2$. (Proof: Let $A = M_1S_1 = S_2M_2$. Then, $S_1 = (S_2M_2^2S_2^*)^{-1/2}S_2M_2 = S_2$.)

Fact 5.18.7. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is singular. Then, there exist a matrix $S \in \mathbb{F}^{n \times n}$ and unique matrices $M_1, M_2 \in \mathbb{F}^{n \times n}$ such that $A = M_1S = SM_2$. In particular, $M_1 = (AA^*)^{1/2}$ and $M_2 = (A^*A)^{1/2}$. (Remark: S is not uniquely determined.)

Fact 5.18.8. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $M, S \in \mathbb{F}^{n \times n}$ be such that $A = MS$, M is positive semidefinite, and S is unitary. Then, A is normal if and only if $MS = SM$. (Proof: See [709, p. 414].)

Fact 5.18.9. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are unitary, and assume that $A + B$ is nonsingular. Then, the unitary factor in the polar decomposition of $A + B$ is $A(A^*B)^{1/2}$. (Proof: See [1013] or [683, p. 216].) (Remark: The principal square root of A^*B exists since $A + B$ is nonsingular.)

5.19 Facts on Additive Decompositions

Fact 5.19.1. Let $A \in \mathbb{C}^{n \times n}$. Then, there exist unitary matrices $B, C \in \mathbb{C}^{n \times n}$ such that

$$A = \frac{1}{2}\sigma_{\max}(A)(B + C).$$

(Proof: See [899, 1484].)

Fact 5.19.2. Let $A \in \mathbb{R}^{n \times n}$. Then, there exist orthogonal matrices $B, C, D, E \in \mathbb{R}^{n \times n}$ such that

$$A = \frac{1}{2}\sigma_{\max}(A)(B + C + D - E).$$

(Proof: See [899]. See also [1484].) (Remark: $A/\sigma_{\max}(A)$ is expressed as an affine combination of B, C, D, E since the sum of the coefficients is 1.)

Fact 5.19.3. Let $A \in \mathbb{R}^{n \times n}$, assume that $\sigma_{\max}(A) \leq 1$, and define $r \triangleq \text{rank}(I - A^*A)$. Then, A is a convex combination of not more than $h(r)$ orthogonal matrices, where

$$h(r) \triangleq \begin{cases} 1 + r, & r \leq 4, \\ 3 + \log_2 r, & r > 4. \end{cases}$$

(Proof: See [899].)

Fact 5.19.4. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) A is positive semidefinite, $\text{tr } A$ is an integer, and $\text{rank } A \leq \text{tr } A$.
- ii) There exist projectors $B_1, \dots, B_l \in \mathbb{F}^{n \times n}$, where $l = \text{tr } A$, such that $A = \sum_{i=1}^l B_i$.

(Proof: See [489, 1460].) (Remark: The minimal number of projectors needed in general is $\text{tr } A$.) (Remark: See Fact 5.19.7.)

Fact 5.19.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, $0 \leq A \leq I$, and $\text{tr } A$ is a rational number. Then, A is the average of a finite set of projectors in $\mathbb{F}^{n \times n}$. (Proof: See [327].) (Remark: The required number of projectors can be arbitrarily large.)

Fact 5.19.6. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and assume that $0 \leq A \leq I$. Then, A is a convex combination of $\lfloor \log_2 n \rfloor + 2$ projectors in $\mathbb{F}^{n \times n}$. (Proof: See [327].)

Fact 5.19.7. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) $\text{tr } A$ is an integer, and $\text{rank } A \leq \text{tr } A$.
- ii) There exist idempotent matrices $B_1, \dots, B_m \in \mathbb{F}^{n \times n}$ such that $A = \sum_{i=1}^m B_i$.
- iii) There exist a positive integer m and idempotent matrices $B_1, \dots, B_m \in \mathbb{F}^{n \times n}$ such that, for all $i = 1, \dots, m$, $\text{rank } B_i = 1$ and $\mathcal{R}(B_i) \subseteq A$, and such that $A = \sum_{i=1}^m B_i$.
- iv) There exist idempotent matrices $B_1, \dots, B_l \in \mathbb{F}^{n \times n}$, where $l \triangleq \text{tr } A$, such that $A = \sum_{i=1}^l B_i$.

(Proof: See [650, 1216, 1460].) (Remark: The minimal number of idempotent matrices is discussed in [1397].) (Remark: See Fact 5.19.8.)

Fact 5.19.8. Let $A \in \mathbb{F}^{n \times n}$, and assume that $2\text{rank } A - 2 \leq \text{tr } A \leq 2n$. Then, there exist idempotent matrices $B, C, D, E \in \mathbb{F}^{n \times n}$ such that $A = B + C + D + E$. (Proof: See [874].) (Remark: See Fact 5.19.10.)

Fact 5.19.9. Let $A \in \mathbb{F}^{n \times n}$. If $n = 2$ or $n = 3$, then there exist $b, c \in \mathbb{F}$ and idempotent matrices $B, C \in \mathbb{F}^{n \times n}$ such that $A = bB + cC$. Furthermore, if $n \geq 4$, then there exist $b, c, d \in \mathbb{F}$ and idempotent matrices $B, C, D \in \mathbb{F}^{n \times n}$ such that $A = bB + cC + dD$. (Proof: See [1111].)

Fact 5.19.10. Let $A \in \mathbb{C}^{n \times n}$, and assume that A is Hermitian. If $n = 2$ or $n = 3$, then there exist $b, c \in \mathbb{C}$ and projectors $B, C \in \mathbb{C}^{n \times n}$ such that $A = bB + cC$. Furthermore, if $4 \leq n \leq 7$, then there exist $b, c, d \in \mathbb{F}$ and projectors $B, C, D \in \mathbb{F}^{n \times n}$ such that $A = bB + cC + dD$. If $n \geq 8$, then there exist $b, c, d, e \in \mathbb{C}$ and projectors $B, C, D, E \in \mathbb{C}^{n \times n}$ such that $A = bB + cC + dD + eE$. (Proof: See [1029].) (Remark: See Fact 5.19.8.)

5.20 Notes

The multicompanion form and the elementary multicompanion form are known as *rational canonical forms* [445, pp. 472–488], while the multicompanion form is traditionally called the *Frobenius canonical form* [146]. The derivation of the Jordan form by means of the elementary multicompanion form and the hypercompanion form follows [1081]. Corollary 5.3.8, Corollary 5.3.9, and Proposition 5.5.12 are given in [240, 241, 1257, 1258, 1261]. Corollary 5.3.9 is due to Frobenius. Canonical forms for congruence transformations are given in [884, 1275].

It is sometimes useful to define block-companion form matrices in which the scalars are replaced by matrix blocks [559, 560, 562]. The companion form provides only one of many connections between matrices and polynomials. Additional connections are given by the *Leslie*, *Schwarz*, and *Routh* forms [139]. Given a polynomial expressed in terms of an arbitrary polynomial basis, the corresponding matrix is in *confederate form*, which specializes to the *comrade form* when the polynomials are orthogonal, which in turn specializes to the *colleague form* when

Chebyshev polynomials are used. The companion, confederate, comrade, and colleague forms are called *congenial* matrices. See [139, 141, 144] and Fact 11.18.25 and Fact 11.18.27 for the Schwarz and Routh forms. The companion matrix is sometimes called a *Frobenius matrix* or the *Frobenius canonical form*, see [5].

Matrix pencils are discussed in [85, 163, 224, 842, 1340, 1352]. Computational algorithms for the Kronecker canonical form are given in [917, 1358]. Applications to linear system theory are discussed in [311, pp. 52–55] and [791].

Application of the polar decomposition to the elastic deformation of solids is discussed in [1072, pp. 140–142].

Chapter Six

Generalized Inverses

Generalized inverses provide a useful extension of the matrix inverse to singular matrices and to rectangular matrices that are neither left nor right invertible.

6.1 Moore-Penrose Generalized Inverse

Let $A \in \mathbb{F}^{n \times m}$. If A is nonzero, then, by the singular value decomposition Theorem 5.6.4, there exist orthogonal matrices $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$ such that

$$A = S_1 \begin{bmatrix} B & 0_{r \times (m-r)} \\ 0_{(n-r) \times r} & 0_{(n-r) \times (m-r)} \end{bmatrix} S_2, \quad (6.1.1)$$

where $B \triangleq \text{diag}[\sigma_1(A), \dots, \sigma_r(A)]$, $r \triangleq \text{rank } A$, and $\sigma_1(A) \geq \sigma_2(A) \geq \dots \geq \sigma_r(A) > 0$ are the positive singular values of A . In (6.1.1), some of the bordering zero matrices may be empty. Then, the (*Moore-Penrose*) *generalized inverse* A^+ of A is the $m \times n$ matrix

$$A^+ \triangleq S_2^* \begin{bmatrix} B^{-1} & 0_{r \times (n-r)} \\ 0_{(m-r) \times r} & 0_{(m-r) \times (n-r)} \end{bmatrix} S_1^*. \quad (6.1.2)$$

If $A = 0_{n \times m}$, then $A^+ \triangleq 0_{m \times n}$, while, if $m = n$ and $\det A \neq 0$, then $A^+ = A^{-1}$. In general, it is helpful to remember that A^+ and A^* are the same size. It is easy to verify that A^+ satisfies

$$AA^+A = A, \quad (6.1.3)$$

$$A^+AA^+ = A^+, \quad (6.1.4)$$

$$(AA^+)^* = AA^+, \quad (6.1.5)$$

$$(A^+A)^* = A^+A. \quad (6.1.6)$$

Hence, for each $A \in \mathbb{F}^{n \times m}$ there exists a matrix $X \in \mathbb{F}^{m \times n}$ satisfying the four conditions

$$AXA = A, \quad (6.1.7)$$

$$XAX = X, \quad (6.1.8)$$

$$(AX)^* = AX, \quad (6.1.9)$$

$$(XA)^* = XA. \quad (6.1.10)$$

We now show that X is uniquely defined by (6.1.7)–(6.1.10).

Theorem 6.1.1. Let $A \in \mathbb{F}^{n \times m}$. Then, $X = A^+$ is the unique matrix $X \in \mathbb{F}^{m \times n}$ satisfying (6.1.7)–(6.1.10).

Proof. Suppose there exists a matrix $X \in \mathbb{F}^{m \times n}$ satisfying (6.1.7)–(6.1.10). Then,

$$\begin{aligned} X &= XAX = X(AX)^* = XX^*A^* = XX^*(AA^+A)^* = XX^*A^*A^+A^* \\ &= X(AX)^*(AA^+)^* = XAXAA^+ = XAA^+ = (XA)^*A^+ = A^*X^*A^+ \\ &= (AA^+A)^*X^*A^+ = A^*A^+A^*X^*A^+ = (A^+A)^*(XA)^*A^+ \\ &= A^+AXAA^+ = A^+AA^+ = A^+. \end{aligned} \quad \square$$

Given $A \in \mathbb{F}^{n \times m}$, $X \in \mathbb{F}^{m \times n}$ is a *(1)-inverse* of A if (6.1.7) holds, a *(1,2)-inverse* of A if (6.1.7) and (6.1.8) hold, and so forth.

Proposition 6.1.2. Let $A \in \mathbb{F}^{n \times m}$, and assume that A is right invertible. Then, $X \in \mathbb{F}^{m \times n}$ is a right inverse of A if and only if X is a (1)-inverse of A . Furthermore, every right inverse (or, equivalently, every (1)-inverse) of A is also a (2,3)-inverse of A .

Proof. Suppose that $AX = I_n$, that is, $X \in \mathbb{F}^{m \times n}$ is a right inverse of A . Then, $AXA = A$, which implies that X is a (1)-inverse of A . Conversely, let X be a (1)-inverse of A , that is, $AXA = A$. Then, letting $\hat{X} \in \mathbb{F}^{m \times n}$ denote a right inverse of A , it follows that $AX = AXA\hat{X} = A\hat{X} = I_n$. Hence, X is a right inverse of A . Finally, if X is a right inverse of A , then it is also a (2,3)-inverse of A . $\square$

Proposition 6.1.3. Let $A \in \mathbb{F}^{n \times m}$, and assume that A is left invertible. Then, $X \in \mathbb{F}^{m \times n}$ is a left inverse of A if and only if X is a (1)-inverse of A . Furthermore, every left inverse (or, equivalently, every (1)-inverse) of A is also a (2,4)-inverse of A .

It can now be seen that A^+ is a particular (right, left) inverse when A is (right, left) invertible.

Corollary 6.1.4. Let $A \in \mathbb{F}^{n \times m}$. If A is right invertible, then A^+ is a right inverse of A . Furthermore, if A is left invertible, then A^+ is a left inverse of A .

The following result provides an explicit expression for A^+ when A is either right invertible or left invertible. It is helpful to note that A is (right, left) invertible if and only if (AA^*, A^*A) is positive definite.

Proposition 6.1.5. Let $A \in \mathbb{F}^{n \times m}$. If A is right invertible, then

$$A^+ = A^*(AA^*)^{-1} \quad (6.1.11)$$

and A^+ is a right inverse of A . If A is left invertible, then

$$A^+ = (A^*A)^{-1}A^* \quad (6.1.12)$$

and A^+ is a left inverse of A .

Proof. It suffices to verify (6.1.7)–(6.1.10) with $X = A^+$. $\square$

Proposition 6.1.6. Let $A \in \mathbb{F}^{n \times m}$. Then, the following statements hold:

- i) $A = 0$ if and only if $A^+ = 0$.
- ii) $(A^+)^+ = A$.
- iii) $\overline{A^+} = \overline{A}^+$.
- iv) $(A^T)^+ = (A^+)^T = A^{+T}$.
- v) $(A^*)^+ = (A^+)^* \triangleq A^{+*}$.
- vi) $\mathcal{R}(A) = \mathcal{R}(AA^*) = \mathcal{R}(AA^+) = \mathcal{R}(A^{+*}) = \mathcal{N}(I - AA^+) = \mathcal{N}(A^*)^\perp$.
- vii) $\mathcal{R}(A^*) = \mathcal{R}(A^*A) = \mathcal{R}(A^+A) = \mathcal{R}(A^+) = \mathcal{N}(I - A^+A) = \mathcal{N}(A)^\perp$.
- viii) $\mathcal{N}(A) = \mathcal{N}(A^+A) = \mathcal{N}(A^*A) = \mathcal{N}(A^{+*}) = \mathcal{R}(I - A^+A) = \mathcal{R}(A^*)^\perp$.
- ix) $\mathcal{N}(A^*) = \mathcal{N}(AA^+) = \mathcal{N}(AA^*) = \mathcal{N}(A^+) = \mathcal{R}(I - AA^+) = \mathcal{R}(A)^\perp$.
- x) AA^+ and A^+A are positive semidefinite.
- xi) $\text{spec}(AA^+) \subseteq \{0, 1\}$ and $\text{spec}(A^+A) \subseteq \{0, 1\}$.
- xii) AA^+ is the projector onto $\mathcal{R}(A)$.
- xiii) A^+A is the projector onto $\mathcal{R}(A^*)$.
- xiv) $I_m - A^+A$ is the projector onto $\mathcal{N}(A)$.
- xv) $I_n - AA^+$ is the projector onto $\mathcal{N}(A^*)$.
- xvi) $x \in \mathcal{R}(A)$ if and only if $x = AA^+x$.
- xvii) $\text{rank } A = \text{rank } A^+ = \text{rank } AA^+ = \text{rank } A^+A = \text{tr } AA^+ = \text{tr } A^+A$.
- xviii) $\text{rank}(I_m - A^+A) = m - \text{rank } A$.
- xix) $\text{rank}(I_n - AA^+) = n - \text{rank } A$.
- xx) $(A^*A)^+ = A^+A^{+*}$.
- xxi) $(AA^*)^+ = A^{+*}A^+$.
- xxii) $AA^+ = A(A^*A)^+A^*$.
- xxiii) $A^+A = A^*(AA^*)^+A$.
- xxiv) $A = AA^*A^{+*} = A^{+*}A^+A$.
- xxv) $A^* = A^*AA^+ = A^+AA^*$.
- xxvi) $A^+ = A^*(AA^*)^+ = (A^*A)^+A^* = A^*(A^*AA^*)^+A^*$.
- xxvii) $A^{+*} = (AA^*)^+A = A(A^*A)^+$.
- xxviii) $A = A(A^*A)^+A^*A = AA^*A(A^*A)^+$.
- xxix) $A = AA^*(AA^*)^+A = (AA^*)^+AA^*A$.
- xxx) If $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$ are unitary, then $(S_1AS_2)^+ = S_2^*A^+S_1^*$.

- xxxi) A is (range Hermitian, normal, Hermitian, positive semidefinite, positive definite) if and only if A^+ is.
- xxxi) If A is a projector, then $A^+ = A$.
- xxxi) $A^+ = A$ if and only if A is tripotent and A^2 is Hermitian.

Proof. The last equality in xxi) is given in [1502]. $\square$

Theorem 2.6.4 showed that the equation $Ax = b$, where $A \in \mathbb{F}^{n \times m}$ and $b \in \mathbb{F}^n$, has a solution $x \in \mathbb{F}^m$ if and only if $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix}$. In particular, $Ax = b$ has a unique solution $x \in \mathbb{F}^m$ if and only if $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix} = m$, while $Ax = b$ has infinitely many solutions if and only if $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix} < m$. We are now prepared to characterize these solutions.

Proposition 6.1.7. Let $A \in \mathbb{F}^{n \times m}$ and $b \in \mathbb{F}^n$. Then, the following statements are equivalent:

- i) There exists a vector $x \in \mathbb{F}^m$ satisfying $Ax = b$.
- ii) $\text{rank } A = \text{rank } \begin{bmatrix} A & b \end{bmatrix}$.
- iii) $b \in \mathcal{R}(A)$.
- iv) $AA^+b = b$.

Now, assume that i)–iv) are satisfied. Then, the following statements hold:

- v) $x \in \mathbb{F}^m$ satisfies $Ax = b$ if and only if

$$x = A^+b + (I - A^+A)x. \quad (6.1.13)$$

- vi) For all $y \in \mathbb{F}^m$, $x \in \mathbb{F}^m$ given by

$$x = A^+b + (I - A^+A)y \quad (6.1.14)$$

satisfies $Ax = b$.

- vii) Let $x \in \mathbb{F}^m$ be given by (6.1.14), where $y \in \mathbb{F}^m$. Then, $y = 0$ minimizes x^*x .

- viii) Assume that $\text{rank } A = m$. Then, there exists a unique vector $x \in \mathbb{F}^m$ satisfying $Ax = b$ given by $x = A^+b$. If, in addition, $A^L \in \mathbb{F}^{m \times m}$ is a left inverse of A , then $A^Lb = A^+b$.

- ix) Assume that $\text{rank } A = n$, and let $A^R \in \mathbb{F}^{m \times n}$ be a right inverse of A . Then, $x = A^Rb$ satisfies $Ax = b$.

Proof. The equivalence of i)–iii) is immediate. To prove the equivalence of iv), note that, if there exists a vector $x \in \mathbb{F}^m$ satisfying $Ax = b$, then $b = Ax = AA^+Ax = AA^+b$. Conversely, if $b = AA^+b$, then $x = A^+b$ satisfies $Ax = b$.

Now, suppose that i)–iv) hold. To prove v), let $x \in \mathbb{F}^m$ satisfy $Ax = b$ so that $A^+Ax = A^+b$. Hence, $x = x + A^+b - A^+Ax = A^+b + (I - A^+A)x$. To prove vi), let $y \in \mathbb{F}^m$, and let $x \in \mathbb{F}^m$ be given by (6.1.14). Then, $Ax = AA^+b = b$. To prove vii), let $y \in \mathbb{F}^m$, and let $x \in \mathbb{F}^m$ be given by (6.1.14). Then, $x^*x = b^*A^+A^+b + y^*(I - A^+A)y$. Therefore, x^*x is minimized by $y = 0$. See also Fact 9.15.1.

To prove *viii*), suppose that $\text{rank } A = m$. Then, A is left invertible, and it follows from Corollary 6.1.4 that A^+ is a left inverse of A . Hence, it follows from (6.1.13) that $x = A^+b$ is the unique solution of $Ax = b$. In addition, $x = A^Lb$. To prove *ix*), let $x = A^Rb$, and note that $AA^Rb = b$. $\square$

Definition 6.1.8. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times l}$, and define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{F}^{(n+k) \times (m+l)}$. Then, the *Schur complement* $A|\mathcal{A}$ of A with respect to $\mathcal{A}$ is defined by

$$A|\mathcal{A} \triangleq D - CA^+B. \quad (6.1.15)$$

Likewise, the *Schur complement* $D|\mathcal{A}$ of D with respect to $\mathcal{A}$ is defined by

$$D|\mathcal{A} \triangleq A - BD^+C. \quad (6.1.16)$$

6.2 Drazin Generalized Inverse

We now introduce a different type of generalized inverse, which applies only to square matrices yet is more useful in certain applications. Let $A \in \mathbb{F}^{n \times n}$. Then, A has a decomposition

$$A = S \begin{bmatrix} J_1 & 0 \\ 0 & J_2 \end{bmatrix} S^{-1}, \quad (6.2.1)$$

where $S \in \mathbb{F}^{n \times n}$ is nonsingular, $J_1 \in \mathbb{F}^{m \times m}$ is nonsingular, and $J_2 \in \mathbb{F}^{(n-m) \times (n-m)}$ is nilpotent. Then, the *Drazin generalized inverse* A^D of A is the matrix

$$A^D \triangleq S \begin{bmatrix} J_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} S^{-1}. \quad (6.2.2)$$

Let $A \in \mathbb{F}^{n \times n}$. Then, it follows from Definition 5.5.1 that $\text{ind } A = \text{ind}_A(0)$. Furthermore, A is nonsingular if and only if $\text{ind } A = 0$, whereas $\text{ind } A = 1$ if and only if A is singular and the zero eigenvalue of A is semisimple. In particular, $\text{ind } 0_{n \times n} = 1$. Note that $\text{ind } A$ is the order of the largest Jordan block of A associated with the zero eigenvalue of A .

It can be seen that A^D satisfies

$$A^D A A^D = A^D, \quad (6.2.3)$$

$$A A^D = A^D A, \quad (6.2.4)$$

$$A^{k+1} A^D = A^k, \quad (6.2.5)$$

where $k = \text{ind } A$. Hence, for all $A \in \mathbb{F}^{n \times n}$ such that $\text{ind } A = k$ there exists a matrix $X \in \mathbb{F}^{n \times n}$ satisfying the three conditions

$$XAX = X, \quad (6.2.6)$$

$$AX = XA, \quad (6.2.7)$$

$$A^{k+1}X = A^k. \quad (6.2.8)$$

We now show that X is uniquely defined by (6.2.6)–(6.2.8).

Theorem 6.2.1. Let $A \in \mathbb{F}^{n \times n}$, and let $k \triangleq \text{ind } A$. Then, $X = A^D$ is the unique matrix $X \in \mathbb{F}^{n \times n}$ satisfying (6.2.6)–(6.2.8).

Proof. Let $X \in \mathbb{F}^{n \times n}$ satisfy (6.2.6)–(6.2.8). If $k = 0$, then it follows from (6.2.8) that $X = A^{-1}$. Hence, let $A = S \begin{bmatrix} J_1 & 0 \\ 0 & J_2 \end{bmatrix} S^{-1}$, where $k = \text{ind } A \geq 1$, $S \in \mathbb{F}^{n \times n}$ is nonsingular, $J_1 \in \mathbb{F}^{m \times m}$ is nonsingular, and $J_2 \in \mathbb{F}^{(n-m) \times (n-m)}$ is nilpotent. Now, let $\hat{X} \triangleq S^{-1}XS = \begin{bmatrix} \hat{X}_1 & \hat{X}_{12} \\ \hat{X}_{21} & \hat{X}_2 \end{bmatrix}$ be partitioned conformably with $S^{-1}AS = \begin{bmatrix} J_1 & 0 \\ 0 & J_2 \end{bmatrix}$. Since, by (6.2.7), $\hat{A}\hat{X} = \hat{X}\hat{A}$, it follows that $J_1\hat{X}_1 = \hat{X}_1J_1$, $J_1\hat{X}_{12} = \hat{X}_{12}J_2$, $J_2\hat{X}_{21} = \hat{X}_{21}J_1$, and $J_2\hat{X}_2 = \hat{X}_2J_2$. Since $J_2^k = 0$, it follows that $J_1\hat{X}_{12}J_2^{k-1} = 0$, and thus $\hat{X}_{12}J_2^{k-1} = 0$. By repeating this argument, it follows that $J_1\hat{X}_{12}J_2 = 0$, and thus $\hat{X}_{12}J_2 = 0$, which implies that $J_1\hat{X}_{12} = 0$, and thus $\hat{X}_{12} = 0$. Similarly, $\hat{X}_{21} = 0$, so that $\hat{X} = \begin{bmatrix} \hat{X}_1 & 0 \\ 0 & \hat{X}_2 \end{bmatrix}$. Now, (6.2.8) implies that $J_1^{k+1}\hat{X}_1 = J_1^k$, and hence $\hat{X}_1 = J_1^{-1}$. Next, (6.2.6) implies that $\hat{X}_2J_2\hat{X}_2 = \hat{X}_2$, which, together with $J_2\hat{X}_2 = \hat{X}_2J_2$, yields $\hat{X}_2^2J_2 = \hat{X}_2$. Consequently, $0 = \hat{X}_2^2J_2^k = \hat{X}_2J_2^{k-1}$, and thus, by repeating this argument, $\hat{X}_2 = 0$. Therefore, $A^D = S \begin{bmatrix} J_1^{-1} & 0 \\ 0 & 0 \end{bmatrix} S^{-1} = S \begin{bmatrix} \hat{X}_1 & 0 \\ 0 & 0 \end{bmatrix} S^{-1} = S\hat{X}S^{-1} = X$. $\square$

Proposition 6.2.2. Let $A \in \mathbb{F}^{n \times n}$, and define $k \triangleq \text{ind } A$. Then, the following statements hold:

- i) $\overline{A}^D = \overline{A^D}$.
- ii) $A^{DT} \triangleq A^{TD} \triangleq (A^T)^D = (A^D)^T$.
- iii) $A^{D*} \triangleq A^{*D} \triangleq (A^*)^D = (A^D)^*$.
- iv) If $r \in \mathbb{P}$, then $A^{Dr} \triangleq A^{rD} \triangleq (A^D)^r = (A^r)^D$.
- v) $\mathcal{R}(A^k) = \mathcal{R}(A^D) = \mathcal{R}(AA^D) = \mathcal{N}(I - AA^D)$.
- vi) $\mathcal{N}(A^k) = \mathcal{N}(A^D) = \mathcal{N}(AA^D) = \mathcal{R}(I - AA^D)$.
- vii) $\text{rank } A^k = \text{rank } A^D = \text{rank } AA^D = \text{def}(I - AA^D)$.
- viii) $\text{def } A^k = \text{def } A^D = \text{def } AA^D = \text{rank}(I - AA^D)$.
- ix) AA^D is the idempotent matrix onto $\mathcal{R}(A^D)$ along $\mathcal{N}(A^D)$.
- x) $A^D = 0$ if and only if A is nilpotent.
- xi) A^D is group invertible.
- xii) $\text{ind } A^D = 0$ if and only if A is nonsingular.
- xiii) $\text{ind } A^D = 1$ if and only if A is singular.
- xiv) $(A^D)^D = (A^D)^\# = A^2A^D$.
- xv) $(A^D)^D = A$ if and only if A is group invertible.
- xvi) If A is idempotent, then $k = 1$ and $A^D = A$.
- xvii) $A = A^D$ if and only if A is tripotent.

Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{ind } A \leq 1$ so that, by Corollary 5.5.9, A is group invertible. In this case, the Drazin generalized inverse A^D is denoted by $A^\#$, which is the *group generalized inverse* of A . Therefore, $A^\#$ satisfies

$$A^\# A A^\# = A^\#, \quad (6.2.9)$$

$$A A^\# = A^\# A, \quad (6.2.10)$$

$$A A^\# A = A, \quad (6.2.11)$$

while $A^\#$ is the unique matrix $X \in \mathbb{F}^{n \times n}$ satisfying

$$X A X = X, \quad (6.2.12)$$

$$A X = X A, \quad (6.2.13)$$

$$A X A = A. \quad (6.2.14)$$

Proposition 6.2.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is group invertible. Then, the following statements hold:

- i) $\overline{A^\#} = \overline{A}^\#$.
- ii) $A^{\#T} \triangleq A^{T\#} \triangleq (A^T)^\# = (A^\#)^T$.
- iii) $A^{\#*} \triangleq A^{*\#} \triangleq (A^*)^\# = (A^\#)^*$.
- iv) If $r \in \mathbb{P}$, then $A^{\#r} \triangleq A^{r\#} \triangleq (A^\#)^r = (A^r)^\#$.
- v) $\mathcal{R}(A) = \mathcal{R}(A A^\#) = \mathcal{N}(I - A A^\#) = \mathcal{R}(A A^+) = \mathcal{N}(I - A A^+)$.
- vi) $\mathcal{N}(A) = \mathcal{N}(A A^\#) = \mathcal{R}(I - A A^\#) = \mathcal{N}(A^+ A) = \mathcal{R}(I - A^+ A)$.
- vii) $\text{rank } A = \text{rank } A^\# = \text{rank } A A^\# = \text{rank } A^\# A$.
- viii) $\text{def } A = \text{def } A^\# = \text{def } A A^\# = \text{def } A^\# A$.
- ix) $A A^\#$ is the idempotent matrix onto $\mathcal{R}(A)$ along $\mathcal{N}(A)$.
- x) $A^\# = 0$ if and only if $A = 0$.
- xi) $A^\#$ is group invertible.
- xii) $(A^\#)^\# = A$.
- xiii) If A is idempotent, then $A^\# = A$.
- xiv) $A = A^\#$ if and only if A is tripotent.

An alternative expression for the idempotent matrix onto $\mathcal{R}(A)$ along $\mathcal{N}(A)$ is given by Proposition 3.5.9.

6.3 Facts on the Moore-Penrose Generalized Inverse for One Matrix

Fact 6.3.1. Let $A \in \mathbb{F}^{n \times m}$, $x \in \mathbb{F}^m$, $b \in \mathbb{F}^n$, and $y \in \mathbb{F}^m$, assume that A is right invertible, and assume that

$$x = A^+ b + (I - A^+ A) y,$$

which satisfies $Ax = b$. Then, there exists a right inverse $A^R \in \mathbb{F}^{m \times n}$ of A such that $x = A^R b$. Furthermore, if $S \in \mathbb{F}^{m \times n}$ is such that $z^T S b \neq 0$, where $z \triangleq (I - A^+ A)y$, then one such right inverse is given by

$$A^R = A^+ + \frac{1}{z^T S b} z z^T S.$$

Fact 6.3.2. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = 1$. Then,

$$A^+ = (\text{tr } A A^*)^{-1} A^*.$$

Consequently, if $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^n$ are nonzero, then

$$(xy^*)^+ = (x^* x y^* y)^{-1} y x^* = \frac{1}{\|x\|_2^2 \|y\|_2^2} y x^*.$$

In particular,

$$1_{n \times m}^+ = \frac{1}{nm} 1_{m \times n}.$$

Fact 6.3.3. Let $x \in \mathbb{F}^n$, and assume that x is nonzero. Then, the projector $A \in \mathbb{F}^{n \times n}$ onto $\text{span}\{x\}$ is given by

$$A = (x^* x)^{-1} x x^*.$$

Fact 6.3.4. Let $x, y \in \mathbb{F}^n$, assume that x, y are nonzero, and assume that $x^* y = 0$. Then, the projector $A \in \mathbb{F}^{n \times n}$ onto $\text{span}\{x, y\}$ is given by

$$A = (x^* x)^{-1} x x^* + (y^* y)^{-1} y y^*.$$

Fact 6.3.5. Let $x, y \in \mathbb{F}^n$, and assume that x, y are linearly independent. Then, the projector $A \in \mathbb{F}^{n \times n}$ onto $\text{span}\{x, y\}$ is given by

$$A = (x^* x y^* y - |x^* y|^2)^{-1} (y^* y x x^* - y^* x y x^* - x^* y x y^* + x^* x y y^*).$$

Furthermore, define $z \triangleq [I - (x^* x)^{-1} x x^*]y$. Then,

$$A = (x^* x)^{-1} x x^* + (z^* z)^{-1} z z^*.$$

(Remark: For $\mathbb{F} = \mathbb{R}$, this result is given in [1206, p. 178].)

Fact 6.3.6. Let $A \in \mathbb{F}^{n \times m}$, assume that $\text{rank } A = n - 1$, let $x \in \mathcal{N}(A)$ be nonzero, let $y \in \mathcal{N}(A^*)$ be nonzero, let $\alpha = 1$ if $\text{spec}(A) = \{0\}$ and the product of the nonzero eigenvalues of A otherwise, and define $k \triangleq \text{amult}_A(0)$. Then,

$$A^A = \frac{(-1)^{k+1} \alpha}{y^* (A^{k-1})^+ x} x y^*.$$

In particular,

$$N_n^A = (-1)^{n+1} E_{1,n}.$$

If, in addition, $k = 1$, then

$$A^A = \frac{\alpha}{y^* x} x y^*.$$

(Proof: See [948, p. 41] and Fact 3.17.4.) (Remark: This result provides an expression for ii) of Fact 2.16.8.) (Remark: If A is range Hermitian, then $\mathcal{N}(A) = \mathcal{N}(A^*)$ and $y^* x \neq 0$, and thus Fact 5.14.3 implies that A^A is semisimple.) (Remark: See Fact 5.14.26.)

Fact 6.3.7. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = n - 1$. Then,

$$A^+ = \frac{1}{\det[AA^* + (AA^*)^A]} A^* [AA^* + (AA^*)^A]^A.$$

(Proof: See [345].) (Remark: Extensions to matrices of arbitrary rank are given in [345].)

Fact 6.3.8. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{k \times n}$, and $C \in \mathbb{F}^{m \times l}$, and assume that B is left inner and C is right inner. Then,

$$(BAC)^+ = C^* A^+ B^*.$$

(Proof: See [654, p. 506].)

Fact 6.3.9. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} \text{rank } [A, A^+] &= 2\text{rank} \begin{bmatrix} A & A^* \end{bmatrix} - 2\text{rank } A \\ &= \text{rank}(A - A^2 A^+) \\ &= \text{rank}(A - A^+ A^2). \end{aligned}$$

Furthermore, the following statements are equivalent:

- i) A is range Hermitian.
- ii) $[A, A^+] = 0$.
- iii) $\text{rank} \begin{bmatrix} A & A^* \end{bmatrix} = \text{rank } A$.
- iv) $A = A^2 A^+$.
- v) $A = A^+ A^2$.

(Proof: See [1306].) (Remark: See Fact 3.6.3, Fact 6.3.10, and Fact 6.3.11.)

Fact 6.3.10. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is range Hermitian.
- ii) $\mathcal{R}(A) = \mathcal{R}(A^+)$.
- iii) $A^+ A = A A^+$.
- iv) $(I - A^+ A)_\perp = A A^+$.
- v) $A = A^2 A^+$.
- vi) $A = A^+ A^2$.
- vii) $A A^+ = A^2 (A^+)^2$.
- viii) $(A A^+)^2 = A^2 (A^+)^2$.
- ix) $(A^+ A)^2 = (A^+)^2 A^2$.
- x) $\text{ind } A \leq 1$, and $(A^+)^2 = (A^2)^+$.
- xi) $\text{ind } A \leq 1$, and $A A^+ A^* A = A^* A^2 A^+$.
- xii) $A^2 A^+ + A^* A^+ A = 2A$.

$$xiii) A^2A^+ + (A^2A^+)^* = A + A^*.$$

$$xiv) \mathcal{R}(A - A^+) = \mathcal{R}(A - A^3).$$

$$xv) \mathcal{R}(A + A^+) = \mathcal{R}(A + A^3).$$

(Proof: See [323, 1281, 1296, 1331] and Fact 6.6.8.) (Remark: See Fact 3.6.3, Fact 6.3.9, and Fact 6.3.11.)

Fact 6.3.11. Let $A \in \mathbb{F}^{n \times n}$, let $r \triangleq \text{rank } A$, let $B \in \mathbb{F}^{n \times r}$ and $C \in \mathbb{F}^{r \times n}$, and assume that $A = BC$ and $\text{rank } B = \text{rank } C = r$. Then, the following statements are equivalent:

$$i) A \text{ is range Hermitian.}$$

$$ii) BB^+ = C^+C.$$

$$iii) \mathcal{N}(B^*) = \mathcal{N}(C).$$

$$iv) B = C^+CB \text{ and } C = CBB^+.$$

$$v) B^+ = B^+C^+C \text{ and } C = CBB^+.$$

$$vi) B = C^+CB \text{ and } C^+ = BB^+C^+.$$

$$vii) B^+ = B^+C^+C \text{ and } C^+ = BB^+C^+.$$

(Proof: See [438].) (Remark: See Fact 3.6.3, Fact 6.3.9, and Fact 6.3.10.)

Fact 6.3.12. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

$$i) A + A^+ = 2AA^+.$$

$$ii) A + A^+ = 2A^+A.$$

$$iii) A + A^+ = AA^+ + A^+A.$$

$$iv) A \text{ is range Hermitian, and } A^2 + AA^+ = 2A.$$

$$v) A \text{ is range Hermitian, and } (I - A)^2A = 0.$$

(Proof: See [1323, 1330].)

Fact 6.3.13. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

$$i) A^+A^* = A^*A^+.$$

$$ii) AA^+A^*A = AA^*A^+A.$$

$$iii) AA^*A^2 = A^2A^*A.$$

If these conditions hold, then A is *star-dagger*. If A is star-dagger, then $A^2(A^+)^2$ and $(A^+)^2A^2$ are positive semidefinite. (Proof: See [651, 1281].) (Remark: See Fact 6.3.16.)

Fact 6.3.14. Let $A \in \mathbb{F}^{n \times m}$, let $B, C \in \mathbb{F}^{m \times n}$, assume that B is a $(1, 3)$ inverse of A , and assume that C is a $(1, 4)$ inverse of A . Then,

$$A^+ = CAB.$$

(Proof: See [174, p. 48].) (Remark: This result is due to Urquhart.)

Fact 6.3.15. Let $A \in \mathbb{F}^{n \times m}$, assume that A is nonzero, let $r \triangleq \text{rank } A$, define $B \triangleq \text{diag}[\sigma_1(A), \dots, \sigma_r(A)]$, and let $S \in \mathbb{F}^{n \times n}$, $K \in \mathbb{F}^{r \times r}$, and $L \in \mathbb{F}^{r \times (m-r)}$ be such that S is unitary,

$$KK^* + LL^* = I_r,$$

and

$$A = S \begin{bmatrix} BK & BL \\ 0_{(n-r) \times r} & 0_{(n-r) \times (m-r)} \end{bmatrix} S^*.$$

Then,

$$A^+ = S \begin{bmatrix} K^*B^{-1} & 0_{r \times (n-r)} \\ L^*B^{-1} & 0_{(m-r) \times (n-r)} \end{bmatrix} S^*.$$

(Proof: See [115, 651].) (Remark: See Fact 5.9.28 and Fact 6.6.15.)

Fact 6.3.16. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is normal.
- ii) $AA^*A^+ = A^+AA^*$.
- iii) A is range Hermitian, and $A^+A^* = A^*A^+$.
- iv) $A(AA^*A)^+ = (AA^*A)^+A$.
- v) $AA^+A^*A^2A^+ = AA^*$.
- vi) $A(A^* + A^+) = (A^* + A^+)A$.
- vii) $A^*A(AA^*)^+A^*A = AA^*$.
- viii) $2AA^*(AA^* + A^*A)^+AA^* = AA^*$.
- ix) There exists a matrix $X \in \mathbb{F}^{n \times n}$ such that $AA^*X = A^*A$ and $A^*AX = AA^*$.
- x) There exists a matrix $X \in \mathbb{F}^{n \times n}$ such that $AX = A^*$ and $A^{+*}X = A^+$.

(Proof: See [323].) (Remark: See Fact 3.7.12, Fact 3.11.4, Fact 5.15.4, Fact 6.3.13, and Fact 6.6.10.)

Fact 6.3.17. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is Hermitian.
- ii) $AA^+ = A^*A^+$.
- iii) $A^2A^+ = A^*$.
- iv) $AA^*A^+ = A$.

(Proof: See [115].)

Fact 6.3.18. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = m$. Then,

$$(AA^*)^+ = A(A^*A)^{-2}A^*.$$

(Remark: See Fact 6.4.7.)

Fact 6.3.19. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$A^+ = \lim_{\alpha \downarrow 0} A^*(AA^* + \alpha I)^{-1} = \lim_{\alpha \downarrow 0} (A^*A + \alpha I)^{-1}A^*.$$

Fact 6.3.20. Let $A \in \mathbb{F}^{n \times m}$, let $\chi_{AA^*}(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_1s + \beta_0$, and let $n - k$ denote the smallest integer in $\{0, \dots, n-1\}$ such that $\beta_k \neq 0$. Then,

$$A^+ = -\beta_{n-k}^{-1}A^*[(AA^*)^{k-1} + \beta_{n-1}(AA^*)^{k-2} + \cdots + \beta_{n-k+1}I].$$

(Proof: See [394].)

Fact 6.3.21. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then,

$$\text{In } A = \text{In } A^+ = \text{In } A^D.$$

If, in addition, A is nonsingular, then

$$\text{In } A = A^{-1}.$$

Fact 6.3.22. Let $A \in \mathbb{F}^{n \times n}$, and consider the following statements:

- i) A is idempotent.
- ii) $\text{rank } A = \text{tr } A$.
- iii) $\text{rank } A \leq \text{tr } A^2A^+A^*$.

Then, i) $\implies$ ii) $\implies$ iii). Furthermore, the following statements are equivalent:

- iv) A is idempotent.
- v) $\text{rank } A = \text{tr } A = \text{tr } A^2A^+A^*$.
- vi) There exist projectors $B, C \in \mathbb{F}^{n \times n}$ such that $A^+ = BC$.
- vii) $A^*A^+ = A^+$.
- viii) $A^+A^* = A^+$.

(Proof: See [807] and [1184, p. 166].)

Fact 6.3.23. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then,

$$A^*A^+A = A^+A$$

and

$$AA^+A^* = AA^+.$$

(Proof: Note that A^*A^+A is a projector, and $\mathcal{R}(A^*A^+A) = \mathcal{R}(A^*) = \mathcal{R}(A^+A)$. Alternatively, use Fact 6.3.22.)

Fact 6.3.24. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then,

$$A^+A + (I - A)(I - A)^+ = I$$

and

$$AA^+ + (I - A)^+(I - A) = I.$$

(Proof: $\mathcal{N}(A) = \mathcal{R}(I - A^+A) = \mathcal{R}(I - A) = \mathcal{R}[(I - A)(I - A^+)]$.) (Remark: The first identity states that the projector onto the null space of A is the same as

the projector onto the range of $I - A$, while the second identity states that the projector onto the range of A is the same as the projector onto the null space of $I - A$.) (Remark: See Fact 3.13.24 and Fact 5.12.18.)

Fact 6.3.25. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, $A + A^* - I$ is nonsingular, and

$$(A + A^* - I)^{-1} = AA^+ + A^+A - I.$$

(Proof: Use Fact 6.3.23.) (Remark: See Fact 3.13.24, Fact 5.12.18, or [998, p. 457] for a geometric interpretation of this identity.)

Fact 6.3.26. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, $2A(A + A^*)^+A^*$ is the projector onto $\mathcal{R}(A) \cap \mathcal{R}(A^*)$. (Proof: See [1320].)

Fact 6.3.27. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A^+ is idempotent.
- ii) $AA^*A = A^2$.

If A is range Hermitian, then the following statements are equivalent:

- iii) A^+ is idempotent.
- iv) $AA^* = A^*A = A$.

The following statements are equivalent:

- v) A^+ is a projector.
- vi) A is a projector.
- vii) A is idempotent, and A and A^+ are similar.
- viii) A is idempotent, and $A = A^+$.
- ix) A is idempotent, and $AA^+ = AA^*$.
- x) $A^+ = A$, and $A^2 = A^*$.
- xi) A and A^+ are idempotent.
- xii) $A = AA^+$.

(Proof: See [1184, pp. 167, 168] and [1281, 1326, 1423].) (Remark: See Fact 3.13.1.)

Fact 6.3.28. Let $A \in \mathbb{F}^{n \times m}$, and let $r \triangleq \text{rank } A$. Then, the following statements are equivalent:

- i) AA^* is a projector.
- ii) A^*A is a projector.
- iii) $AA^*A = A$.
- iv) $A^*AA^* = A^*$.
- v) $A^+ = A^*$.
- vi) $\sigma_1(A) = \sigma_r(A) = 1$.

In particular, $N_n^+ = N_n^T$. (Proof: See [174, pp. 219–220].) (Remark: A is a *partial isometry*, which preserves lengths and distances with respect to the Euclidean norm on $\mathcal{R}(A^*)$. See [174, p. 219].) (Remark: See Fact 5.11.30.)

Fact 6.3.29. Let $A \in \mathbb{F}^{n \times m}$, assume that A is nonzero, and let $r \triangleq \text{rank } A$. Then, for all $i = 1, \dots, r$, the singular values of A^+ are given by

$$\sigma_i(A^+) = \sigma_{r+1-i}^{-1}(A).$$

In particular,

$$\sigma_r(A) = 1/\sigma_{\max}(A^+).$$

If, in addition, $A \in \mathbb{F}^{n \times n}$ and A is nonsingular, then

$$\sigma_{\min}(A) = 1/\sigma_{\max}(A^{-1}).$$

Fact 6.3.30. Let $A \in \mathbb{F}^{n \times m}$. Then, $X = A^+$ is the unique matrix satisfying

$$\text{rank} \begin{bmatrix} A & AA^+ \\ A^+A & X \end{bmatrix} = \text{rank } A.$$

(Remark: See Fact 2.17.10 and Fact 6.6.2.) (Proof: See [483].)

Fact 6.3.31. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is centrohermitian. Then, A^+ is centrohermitian. (Proof: See [883].)

Fact 6.3.32. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $A^2 = AA^*A$.
- ii) A is the product of two projectors.
- iii) $A = A(A^+)^2A$.

(Remark: This result is due to Crimmins. See [1114].)

Fact 6.3.33. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$A^+ = 4(I + A^+A)^+A^+(I + AA^+)^+.$$

(Proof: Use Fact 6.4.36 with $B = A$.)

Fact 6.3.34. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is unitary. Then,

$$\lim_{k \rightarrow \infty} \frac{1}{k} \sum_{i=0}^{k-1} A^i = I - (A - I)(A - I)^+.$$

(Remark: $I - (A - I)(A - I)^+$ is the projector onto $\{x: Ax = x\} = \mathcal{N}(A - I)$.)

(Remark: This result is the *ergodic theorem*.) (Proof: Use Fact 11.21.11 and Fact 11.21.13, and note that $(A - I)^* = (A - I)^+$. See [626, p. 185].)

Fact 6.3.35. Let $A \in \mathbb{F}^{n \times m}$, and define $\{B_i\}_{i=1}^\infty$ by

$$B_{i+1} \triangleq 2B_i - B_iAB_i,$$

where $B_0 \triangleq \alpha A^*$ and $\alpha \in (0, 2/\sigma_{\max}^2(A))$. Then,

$$\lim_{i \rightarrow \infty} B_i = A^+.$$

(Proof: See [144, p. 259] or [283, p. 250]. This result is due to Ben-Israel.) (Remark: This sequence is a Newton-Raphson algorithm.) (Remark: B_0 satisfies $\text{sprad}(I - B_0 A) < 1$.) (Remark: For the case in which A is square and nonsingular, see Fact 2.16.29.) (Problem: Does convergence hold for all $B_0 \in \mathbb{F}^{n \times n}$ satisfying $\text{sprad}(I - B_0 A) < 1$?)

Fact 6.3.36. Let $A \in \mathbb{F}^{n \times m}$, let $(A_i)_{i=1}^{\infty} \subset \mathbb{F}^{n \times m}$, and assume that $\lim_{i \rightarrow \infty} A_i = A$. Then, $\lim_{i \rightarrow \infty} A_i^+ = A^+$ if and only if there exists a positive integer k such that, for all $i > k$, $\text{rank } A_i = \text{rank } A$. (Proof: See [283, pp. 218, 219].)

6.4 Facts on the Moore-Penrose Generalized Inverse for Two or More Matrices

Fact 6.4.1. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then, the following statements are equivalent:

- i) $B = A^+$.
- ii) $A^*AB = A^*$ and $B^*BA = B^*$.
- iii) $BAA^* = A^*$ and $ABB^* = B^*$.

(Remark: See [654, pp. 503, 513].)

Fact 6.4.2. Let $A \in \mathbb{F}^{n \times n}$, and let $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$ be nonzero. Furthermore, define

$$\begin{aligned} d &\triangleq A^+x, & e &\triangleq A^{+*}y, & f &\triangleq (I - AA^+)x, & g &\triangleq (I - A^+A)y, \\ \delta &\triangleq d^*d, & \eta &\triangleq e^*e, & \phi &\triangleq f^*f, & \psi &\triangleq g^*g, \\ \lambda &\triangleq 1 + y^*A^+x, & \mu &\triangleq |\lambda|^2 + \delta\psi, & \nu &\triangleq |\lambda|^2 + \eta\phi. \end{aligned}$$

Then,

$$\text{rank}(A + xy^*) = \text{rank } A - 1$$

if and only if

$$x \in \mathcal{R}(A), \quad y \in \mathcal{R}(A^*), \quad \lambda = 0.$$

In this case,

$$(A + xy^*)^+ = A^+ - \delta^{-1}dd^*A^+ - \eta^{-1}A^+ee^* + (\delta\eta)^{-1}d^*A^+ede^*.$$

Furthermore,

$$\text{rank}(A + xy^*) = \text{rank } A$$

if and only if

$$\begin{cases} x \in \mathcal{R}(A), & y \in \mathcal{R}(A^*), & \lambda \neq 0, \\ x \in \mathcal{R}(A), & y \notin \mathcal{R}(A^*), \\ x \notin \mathcal{R}(A), & y \in \mathcal{R}(A^*). \end{cases}$$

In this case, respectively,

$$\begin{cases} (A + xy^*)^+ = A^+ - \lambda^{-1}de^*, \\ (A + xy^*)^+ = A^+ - \mu^{-1}(\psi dd^*A^+ + \delta ge^*) + \mu^{-1}(\lambda gd^*A^+ - \bar{\lambda}de^*), \\ (A + xy^*)^+ = A^+ - \nu^{-1}(\phi A^+ ee^* + \eta df^*) + \nu^{-1}(\lambda A^+ ef^* - \bar{\lambda}de^*). \end{cases}$$

Finally,

$$\text{rank}(A + xy^*) = \text{rank } A + 1$$

if and only if

$$x \notin \mathcal{R}(A), \quad y \notin \mathcal{R}(A^*).$$

In this case,

$$(A + xy^*)^+ = A^+ - \phi^{-1}df^* - \psi^{-1}ge^* + \lambda(\phi\psi)^{-1}gf^*.$$

(Proof: See [108]. To prove sufficiency in the first alternative of the third statement, let $\hat{x}, \hat{y} \in \mathbb{F}^n$ be such that $x = A\hat{x}$ and $y = A^*\hat{y}$. Then, $A + xy^* = A(I + \hat{x}\hat{y}^*)$. Since $\alpha \neq 0$ it follows that $-1 \neq y^*A^+x = \hat{y}^*AA^+A\hat{x} = \hat{y}^*A\hat{x} = y^*\hat{x}$. It now follows that $I + \hat{x}\hat{y}^*$ is an elementary matrix and thus, by Fact 3.7.19, is nonsingular.) (Remark: An equivalent version of the first statement is given in [330] and [721, p. 33]. A detailed treatment of the generalized inverse of an outer-product perturbation is given in [1396, pp. 152–157].) (Remark: See Fact 2.10.25.)

Fact 6.4.3. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let $b \in \mathbb{F}^n$, and define $S \triangleq I - A^+A$. Then,

$$(A + bb^*)^+ = \begin{cases} [I - (b^*(A^+)^2b)^{-1}A^+bb^*A^+]A^+[I - (b^*(A^+)^2b)^{-1}A^+bb^*A^+], & 1 + b^*A^+b = 0, \\ A^+ - (1 + b^*A^+b)^{-1}A^+bb^*A^+, & 1 + b^*A^+b \neq 0, \\ [I - (b^*Sb)^{-1}Sbb^*]A^+[I - (b^*Sb)^{-1}bb^*S] + (b^*Sb)^{-2}Sbb^*S, & b^*Sb \neq 0. \end{cases}$$

(Proof: See [1006].)

Fact 6.4.4. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, let $C \in \mathbb{F}^{m \times m}$, assume that C is positive definite, and let $B \in \mathbb{F}^{n \times m}$. Then,

$$(A + BCB^*)^+ = A^+ - A^+B(C^{-1} + B^*A^+B)^{-1}B^*A^+$$

if and only if

$$AA^+B = B.$$

(Proof: See [1049].) (Remark: $AA^+B = B$ is equivalent to $\mathcal{R}(B) \subseteq \mathcal{R}(A)$.) (Remark: Extensions of the matrix inversion lemma are considered in [384, 487, 1006, 1126] and [654, pp. 426–428, 447, 448].)

Fact 6.4.5. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, $AB = 0$ if and only if $B^+A^+ = 0$.

Fact 6.4.6. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then, $A^+B = 0$ if and only if $A^*B = 0$.

Fact 6.4.7. Let $A \in \mathbb{F}^{n \times m}$, assume that $\text{rank } A = m$, let $B \in \mathbb{F}^{n \times n}$, and assume that B is positive definite. Then,

$$(ABA^*)^+ = A(A^*A)^{-1}B^{-1}(A^*A)^{-1}A^*.$$

(Proof: Use Fact 6.3.18.)

Fact 6.4.8. Let $A \in \mathbb{F}^{n \times m}$, let $S \in \mathbb{F}^{m \times m}$, assume that S is nonsingular, and define $B \triangleq AS$. Then,

$$BB^+ = AA^+.$$

(Proof: See [1184, p. 144].)

Fact 6.4.9. Let $A \in \mathbb{F}^{n \times r}$ and $B \in \mathbb{F}^{r \times m}$, and assume that $\text{rank } A = \text{rank } B = r$. Then,

$$(AB)^+ = B^+A^+ = B^*(BB^*)^{-1}(A^*A)^{-1}A^*.$$

(Remark: AB is a full-rank factorization.)

Fact 6.4.10. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$(AB)^+ = (A^+AB)^+(ABB^+)^+.$$

If, in addition, $\mathcal{R}(B) = \mathcal{R}(A^*)$, then $A^+AB = B$, $ABB^+ = A$, and

$$(AB)^+ = B^+A^+.$$

(Proof: See [1177, pp. 192] or [1301].) (Remark: This result is due to Cline and Greville.)

Fact 6.4.11. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, and define $B_1 \triangleq A^+AB$ and $A_1 \triangleq AB_1B_1^+$. Then,

$$AB = A_1B_1$$

and

$$(AB)^+ = B_1^+A_1^+.$$

(Proof: See [1177, pp. 191, 192].)

Fact 6.4.12. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, the following statements are equivalent:

- i) $(AB)^+ = B^+A^+$.
- ii) $\mathcal{R}(A^*AB) \subseteq \mathcal{R}(B)$ and $\mathcal{R}(BB^*A^*) \subseteq \mathcal{R}(A^*)$.
- iii) $(AB)(AB)^+ = (AB)B^+A^+$ and $(AB)^+(AB) = B^+A^+AB$.
- iv) $A^*AB = BB^+A^*AB$ and $ABB^* = ABB^*A^+A$.
- v) $AB(AB)^+A = ABB^+$ and $A^+AB = B(AB)^+AB$.
- vi) A^*ABB^+ and A^+ABB^* are Hermitian.
- vii) $(ABB^+)^+ = BB^+A^+$ and $(A^+AB)^+ = B^+A^+A$.
- viii) $B^+(ABB^+)^+ = B^+A^+$ and $(A^+AB)^+A = B^+A^+$.
- ix) $A^*ABB^* = BB^+A^*ABB^*A^+A$.

(Proof: See [15, p. 53] and [587, 1291].) (Remark: The equivalence of *i*) and *ii*) is due to Greville.) (Remark: Conditions under which B^+A^+ is a (1)-inverse of AB are given in [1291].) (Remark: See [1416].)

Fact 6.4.13. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, $AB = 0$ if and only if $B^+A^+ = 0$. Furthermore, $A^+B = 0$ if and only if $A^*B = 0$. (Proof: The first statement follows from $ix) \implies i)$ of Fact 6.4.12. The second statement follows from Proposition 6.1.6.)

Fact 6.4.14. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, the following statements are equivalent:

$$i) (AB)^+ = B^+A^+ - B^+[(I - BB^+)(I - A^+A)]^+A^+.$$

$$ii) \mathcal{R}(AA^*AB) = \mathcal{R}(AB) \text{ and } \mathcal{R}[(ABB^*B)^*] = \mathcal{R}[(AB)^*].$$

(Proof: See [1289].)

Fact 6.4.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then,

$$\mathcal{R}([A, B]) = \mathcal{R}[(A - B)^+ - (A - B)].$$

Consequently, $(A - B)^+ = (A - B)$ if and only if $AB = BA$. (Proof: See [1288].)

Fact 6.4.16. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the following statements hold:

$$i) (AB)^+ = B(AB)^+.$$

$$ii) (AB)^+ = (AB)^+A.$$

$$iii) (AB)^+ = B(AB)^+A.$$

$$iv) (AB)^+ = BA - B(B_\perp A_\perp)^+A.$$

$$v) (AB)^+, B(AB)^+, (AB)^+A, B(AB)^+A, \text{ and } BA - B(B_\perp A_\perp)^+A \text{ are idempotent.}$$

$$vi) AB = A(AB)^+B.$$

$$vii) (AB)^2 = AB + AB(B_\perp A_\perp)^+AB.$$

(Proof: To prove *i*) note that $\mathcal{R}[(AB)^+] = \mathcal{R}[(AB)^*] = \mathcal{R}(BA)$, and thus $\mathcal{R}[B(AB)^+] = \mathcal{R}[B(AB)^*] = \mathcal{R}(BA)$. Hence, $\mathcal{R}[(AB)^+] = \mathcal{R}[B(AB)^+]$. It now follows from Fact 3.13.14 that $(AB)^+ = B(AB)^+$. Statement *iv*) follows from Fact 6.4.14. Statements *v*) and *vi*) follow from *iii*). Statement *vii*) follows from *iv*) and *vi*).) (Remark: The converse of the first result in *v*) is given by Fact 6.4.17.) (Remark: See Fact 6.3.27, Fact 6.4.10, and Fact 6.4.21. See [1289, 1423].)

Fact 6.4.17. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is idempotent. Then, there exist projectors $B, C \in \mathbb{F}^{n \times n}$ such that $A = (BC)^+$. (Proof: See [322, 537].) (Remark: The converse of this result is given by *v*) of Fact 6.4.16.) (Remark: This result is due to Penrose.)

Fact 6.4.18. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $\mathcal{R}(A)$ and $\mathcal{R}(B)$ are complementary subspaces. Furthermore, define $P \triangleq AA^+$ and $Q \triangleq BB^+$. Then, the matrix $(Q_{\perp}P)^+$ is the idempotent matrix onto $\mathcal{R}(B)$ along $\mathcal{R}(A)$. (Proof: See [588].) (Remark: See Fact 3.12.33, Fact 3.13.24, and Fact 6.4.19.)

Fact 6.4.19. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are projectors, and assume that $\mathcal{R}(A)$ and $\mathcal{R}(B)$ are complementary subspaces. Then, $(A_{\perp}B)^+$ is the idempotent matrix onto $\mathcal{R}(B)$ along $\mathcal{R}(A)$. (Proof: See Fact 6.4.18, [593], or [744].) (Remark: It follows from Fact 6.4.16 that $(A_{\perp}B)^+$ is idempotent.) (Remark: See Fact 3.12.33, Fact 3.13.24, and Fact 6.4.18.)

Fact 6.4.20. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are projectors, and assume that $A - B$ is nonsingular. Then, $I - BA$ is nonsingular, and

$$(A_{\perp}B)^+ = (I - BA)^{-1}B(I - BA).$$

(Proof: Combine Fact 3.13.24 and Fact 6.4.19.)

Fact 6.4.21. Let $k \geq 1$, let $A_1, \dots, A_k \in \mathbb{F}^{n \times n}$, assume that $A_1, \dots, A_k$ are projectors, and define $B_1, \dots, B_{k-1} \in \mathbb{F}^{n \times n}$ by

$$B_i = (A_1 \cdots A_{k-i+1})^+ A_1 \cdots A_{k-i}, \quad i = 1, \dots, k-2,$$

and

$$B_{k-1} = A_2 \cdots A_k (A_1 \cdots A_k)^+.$$

Then, $B_1, \dots, B_{k-1}$ are idempotent, and

$$(A_1 \cdots A_k)^+ = B_1 \cdots B_{k-1}.$$

(Proof: See [1298].) (Remark: When $k = 2$, the result that B_1 is idempotent is given by *vi*) of Fact 6.4.16.)

Fact 6.4.22. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times n}$, and assume that A is idempotent. Then,

$$A^*(BA)^+ = (BA)^+.$$

(Proof: See [654, p. 514].)

Fact 6.4.23. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the following statements are equivalent:

- i) AB is a projector.
- ii) $[(AB)^+]^2 = [(AB)^2]^+.$

(Proof: See [1321].) (Remark: See Fact 3.13.20 and Fact 5.12.16.)

Fact 6.4.24. Let $A \in \mathbb{F}^{n \times m}$. Then, $B \in \mathbb{F}^{m \times m}$ satisfies $BAB = B$ if and only if there exist projectors $C \in \mathbb{F}^{n \times n}$ and $D \in \mathbb{F}^{m \times m}$ such that $B = (CAD)^+$. (Proof: See [588].)

Fact 6.4.25. Let $A \in \mathbb{F}^{n \times n}$. Then, A is idempotent if and only if there exist projectors $B, C \in \mathbb{F}^{n \times n}$ such that $A = (BC)^+$. (Proof: Let $A = I$ in Fact 6.4.24.) (Remark: See [594].)

Fact 6.4.26. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is range Hermitian. Then, $AB = BA$ if and only if $A^+B = BA^+$. (Proof: See [1280].)

Fact 6.4.27. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are range Hermitian. Then, the following statements are equivalent:

- i) $AB = BA$.
- ii) $A^+B = BA^+$.
- iii) $AB^+ = B^+A$.
- iv) $A^+B^+ = B^+A^+$.

(Proof: See [1280].)

Fact 6.4.28. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are range Hermitian, and assume that $(AB)^+ = A^+B^+$. Then, AB is range Hermitian. (Proof: See [648].) (Remark: See Fact 8.20.21.)

Fact 6.4.29. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are range Hermitian. Then, the following statements are equivalent:

- i) AB is range Hermitian.
- ii) $AB(I - A^+A) = 0$ and $(I - B^+B)AB = 0$.
- iii) $\mathcal{N}(A) \subseteq \mathcal{N}(AB)$ and $\mathcal{R}(AB) \subseteq \mathcal{R}(B)$.
- iv) $\mathcal{N}(AB) = \mathcal{N}(A) + \mathcal{N}(B)$ and $\mathcal{R}(AB) = \mathcal{R}(A) \cap \mathcal{R}(B)$.

(Proof: See [648, 832].)

Fact 6.4.30. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, and assume that $\text{rank } B = m$. Then,

$$AB(AB)^+ = AA^+.$$

Fact 6.4.31. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times n}$, and $C \in \mathbb{F}^{m \times n}$, and assume that $BAA^* = A^*$ and $A^*AC = A^*$. Then,

$$A^+ = BAC.$$

(Proof: See [15, p. 36].) (Remark: This result is due to Decell.)

Fact 6.4.32. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $A + B$ is nonsingular. Then, the following statements are equivalent:

- i) $\text{rank } A + \text{rank } B = n$.
- ii) $A(A + B)^{-1}B = 0$.
- iii) $B(A + B)^{-1}A = 0$.
- iv) $A(A + B)^{-1}A = A$.
- v) $B(A + B)^{-1}B = B$.
- vi) $A(A + B)^{-1}B + B(A + B)^{-1}A = 0$.

$$vii) \quad A(A+B)^{-1}A + B(A+B)^{-1}B = A + B.$$

$$viii) \quad (A+B)^{-1} = [(I - BB^+)A(I - B^+B)]^+ + [(I - AA^+)B(I - A^+A)]^+.$$

(Proof: See [1302].) (Remark: See Fact 2.11.4 and Fact 8.20.23.)

Fact 6.4.33. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$, and assume that A and B are projectors. Then, the following statements hold:

$$i) \quad A(A-B)^+B = B(A-B)^+A = 0.$$

$$ii) \quad A - B = A(A-B)^+A - B(B-A)^+B.$$

$$iii) \quad (A-B)^+ = (A-AB)^+ + (AB-B)^+.$$

$$iv) \quad (A-B)^+ = (A-BA)^+ + (BA-B)^+.$$

$$v) \quad (A-B)^+ = A - B + B(A-BA)^+ - (B-BA)^+A.$$

$$vi) \quad (A-B)^+ = A - B + (A-AB)^+B - A(B-AB)^+.$$

$$vii) \quad (I - A - B)^+ = (A_{\perp}B_{\perp})^+ - (AB)^+.$$

$$viii) \quad (I - A - B)^+ = (B_{\perp}A_{\perp})^+ - (BA)^+.$$

Furthermore, the following statements are equivalent:

$$ix) \quad AB = BA.$$

$$x) \quad (A-B)^+ = A - B.$$

$$xi) \quad B(A-BA)^+ = (B-BA)^+A.$$

$$xii) \quad (A-B)^3 = A - B.$$

$$xiii) \quad A - B \text{ is tripotent.}$$

(Proof: See [322].) (Remark: See Fact 3.12.22.)

Fact 6.4.34. Let $A, B \in \mathbb{F}^{n \times m}$, and assume that $A^*B = 0$ and $BA^* = 0$. Then,

$$(A+B)^+ = A^+ + B^+.$$

(Proof: Use Fact 2.10.29 and Fact 6.4.35. See [339] and [654, p. 513].) (Remark: This result is due to Penrose.)

Fact 6.4.35. Let $A, B \in \mathbb{F}^{n \times m}$, and assume that $\text{rank}(A+B) = \text{rank } A + \text{rank } B$. Then,

$$(A+B)^+ = (I - C^+B)A^+(I - BC^+) + C^+,$$

where $C \triangleq (I - AA^+)B(I - A^+A)$. (Proof: See [339].)

Fact 6.4.36. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$(A+B)^+ = (I + A^+B)^+(A^+ + A^+BA^+)(I + BA^+)^+$$

if and only if $AA^+B = B = BA^+A$. Furthermore, if $n = m$ and A is nonsingular, then

$$(A+B)^+ = (I + A^{-1}B)^+(A^{-1} + A^{-1}BA^{-1})(I + BA^{-1})^+.$$

(Proof: See [339].) (Remark: If A and $A+B$ are nonsingular, then the last state-

ment yields $(A + B)^{-1} = (A + B)^{-1}(A + B)(A + B)^{-1}$ for which the assumption that A is nonsingular is superfluous.)

Fact 6.4.37. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\begin{aligned} A^+ - B^+ &= B^+(B - A)A^+ + (I - B^+B)(A^* - B^*)A^{+*}A^+ + B^+B^{+*}(A^* - B^*)(I - AA^+) \\ &= A^+(B - A)B^+ + (I - A^+A)(A^* - B^*)B^{+*}B^+ + A^+A^{+*}(A^* - B^*)(I - BB^+). \end{aligned}$$

Furthermore, if B is left invertible, then

$$A^+ - B^+ = B^+(B - A)A^+ + B^+B^{+*}(A^* - B^*)(I - AA^+),$$

while, if B is right invertible, then

$$A^+ - B^+ = A^+(B - A)B^+ + (I - A^+A)(A^* - B^*)B^{+*}B^+.$$

(Proof: See [283, p. 224].)

Fact 6.4.38. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{l \times k}$, and $C \in \mathbb{F}^{n \times k}$. Then, there exists a matrix $X \in \mathbb{F}^{m \times l}$ satisfying $AXB = C$ if and only if $AA^+CB^+B = C$. Furthermore, X satisfies $AXB = C$ if and only if there exists a matrix $Y \in \mathbb{F}^{m \times l}$ such that

$$X = A^+CB^+ + Y - A^+AYBB^+.$$

Finally, if $Y = 0$, then $\text{tr } X^*X$ is minimized. (Proof: Use Proposition 6.1.7. See [948, p. 37] and, for Hermitian solutions, see [808].)

Fact 6.4.39. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = m$. Then, $A^L \in \mathbb{F}^{m \times n}$ is a left inverse of A if and only if there exists a matrix $B \in \mathbb{F}^{m \times n}$ such that

$$A^L = A^+ + B(I - AA^+).$$

(Proof: Use Fact 6.4.3 with $A = C = I_m$.)

Fact 6.4.40. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = n$. Then, $A^R \in \mathbb{F}^{m \times n}$ is a right inverse of A if and only if there exists a matrix $B \in \mathbb{F}^{m \times n}$ such that

$$A^R = A^+ + (I - A^+A)B.$$

(Proof: Use Fact 6.4.38 with $B = C = I_n$.)

Fact 6.4.41. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then,

$$\text{glb}\{A, B\} = \lim_{k \rightarrow \infty} A(BA)^k = 2A(A + B)^+B.$$

Furthermore, $2A(A + B)^+B$ is the projector onto $\mathcal{R}(A) \cap \mathcal{R}(B)$. (Proof: See [39] and [627, pp. 64, 65, 121, 122].) (Remark: See Fact 6.4.42 and Fact 8.20.18.)

Fact 6.4.42. Let $A \in \mathbb{R}^{n \times m}$ and $B \in \mathbb{R}^{n \times l}$. Then,

$$\mathcal{R}(A) \cap \mathcal{R}(B) = \mathcal{R}[AA^+(AA^+ + BB^+)^+BB^+].$$

(Remark: See Theorem 2.3.1 and Fact 8.20.18.)

Fact 6.4.43. Let $A \in \mathbb{R}^{n \times m}$ and $B \in \mathbb{R}^{n \times l}$. Then, $\mathcal{R}(A) \subseteq \mathcal{R}(B)$ if and only if $BB^+A = A$. (Proof: See [15, p. 35].)

Fact 6.4.44. Let $A \in \mathbb{R}^{n \times m}$ and $B \in \mathbb{R}^{n \times l}$. Then,

$$\begin{aligned} \dim[\mathcal{R}(A) \cap \mathcal{R}(B)] &= \text{rank } AA^+(AA^+ + BB^+)^+BB^+ \\ &= \text{rank } A + \text{rank } B - \text{rank} \begin{bmatrix} A & B \end{bmatrix}. \end{aligned}$$

(Proof: Use Fact 2.11.1, Fact 2.11.12, and Fact 6.4.42.) (Remark: See Fact 2.11.8.)

Fact 6.4.45. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then,

$$\text{lub}\{A, B\} = (A + B)(A + B)^+.$$

Furthermore, $\text{lub}\{A, B\}$ is the projector onto $\mathcal{R}(A) + \mathcal{R}(B) = \text{span}[\mathcal{R}(A) \cup \mathcal{R}(B)]$. (Proof: Use Fact 2.9.13 and Fact 8.7.3.) (Remark: See Fact 8.7.2.)

Fact 6.4.46. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then,

$$\text{lub}\{A, B\} = I - \lim_{k \rightarrow \infty} A_{\perp}(B_{\perp}A_{\perp})^k = I - 2A_{\perp}(A_{\perp} + B_{\perp})^+B_{\perp}.$$

Furthermore, $I - 2A_{\perp}(A_{\perp} + B_{\perp})^+B_{\perp}$ is the projector onto

$$\begin{aligned} [\mathcal{R}(A_{\perp}) \cap \mathcal{R}(B_{\perp})]^{\perp} &= [\mathcal{N}(A) \cap \mathcal{N}(B)]^{\perp} \\ &= [\mathcal{N}(A)]^{\perp} + [\mathcal{N}(B)]^{\perp} \\ &= \mathcal{R}(A) + \mathcal{R}(B) \\ &= \text{span}[\mathcal{R}(A) \cup \mathcal{R}(B)]. \end{aligned}$$

Consequently,

$$I - 2A_{\perp}(A_{\perp} + B_{\perp})^+B_{\perp} = (A + B)(A + B)^+.$$

(Proof: See [39] and [627, pp. 64, 65, 121, 122].) (Remark: See Fact 6.4.42 and Fact 8.20.18.)

Fact 6.4.47. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$A \overset{*}{\leq} B$$

if and only if

$$A^+A = A^+B$$

and

$$AA^+ = BA^+.$$

(Proof: See [652].) (Remark: See Fact 2.10.35.)

6.5 Facts on the Moore-Penrose Generalized Inverse for Partitioned Matrices

Fact 6.5.1. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$(A + B)^+ = \frac{1}{2} \begin{bmatrix} I_n & I_n \end{bmatrix} \begin{bmatrix} A & B \\ B & A \end{bmatrix}^+ \begin{bmatrix} I_m \\ I_m \end{bmatrix}.$$

(Proof: See [1278, 1282, 1302].) (Remark: See Fact 2.17.5 and Fact 2.19.7.)

Fact 6.5.2. Let $A_1, \dots, A_k \in \mathbb{F}^{n \times m}$. Then,

$$(A_1 + \dots + A_k)^+ = \frac{1}{k} \begin{bmatrix} I_n & \cdots & I_n \end{bmatrix} \begin{bmatrix} A_1 & A_2 & \cdots & A_k \\ A_k & A_1 & \cdots & A_{k-1} \\ \vdots & \vdots & \ddots & \vdots \\ A_2 & A_3 & \cdots & A_1 \end{bmatrix}^+ \begin{bmatrix} I_m \\ \vdots \\ I_m \end{bmatrix}.$$

(Proof: See [1282].) (Remark: The partitioned matrix is *block circulant*. See Fact 6.6.1 and Fact 2.17.6.)

Fact 6.5.3. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

- i) $\mathcal{R}(\begin{bmatrix} A \\ A^*A \end{bmatrix}) = \mathcal{R}(\begin{bmatrix} B \\ B^*B \end{bmatrix})$.
- ii) $\mathcal{R}(\begin{bmatrix} A \\ A^+A \end{bmatrix}) = \mathcal{R}(\begin{bmatrix} B \\ B^+B \end{bmatrix})$.
- iii) $A = B$.

(Remark: This result is due to Tian.)

Fact 6.5.4. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times l}$. Then,

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} I & 0 \\ CA^+ & I \end{bmatrix} \begin{bmatrix} A & B - AA^+B \\ C - CA^+A & D - CA^+B \end{bmatrix} \begin{bmatrix} I & A^+B \\ 0 & I \end{bmatrix}.$$

(Proof: See [1290].)

Fact 6.5.5. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$, and assume that $B = AA^+B$. Then,

$$\text{In } \mathcal{A} = \text{In } A + \text{In}(A|A).$$

(Remark: This result is the *Haynsworth inertia additivity formula*. See [1103].)

(Remark: If $\mathcal{A}$ is positive semidefinite, then $B = AA^+B$. See Proposition 8.2.4.)

Fact 6.5.6. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times l}$. Then,

$$\begin{aligned} \text{rank} \begin{bmatrix} A & B \end{bmatrix} &= \text{rank } A + \text{rank}(B - AA^+B) \\ &= \text{rank } B + \text{rank}(A - BB^+A) \\ &= \text{rank } A + \text{rank } B - \dim[\mathcal{R}(A) \cap \mathcal{R}(B)], \end{aligned}$$

$$\begin{aligned} \text{rank} \begin{bmatrix} A \\ C \end{bmatrix} &= \text{rank } A + \text{rank}(C - CA^+A) \\ &= \text{rank } C + \text{rank}(A - AC^+C) \\ &= \text{rank } A + \text{rank } C - \dim[\mathcal{R}(A^*) \cap \mathcal{R}(C^*)], \end{aligned}$$

$$\text{rank} \begin{bmatrix} A & B \\ C & 0 \end{bmatrix} = \text{rank } B + \text{rank } C + \text{rank}[(I_n - BB^+)A(I_m - C^+C)],$$

and

$$\begin{aligned} \operatorname{rank} \begin{bmatrix} A & B \\ C & D \end{bmatrix} &= \operatorname{rank} A + \operatorname{rank} X + \operatorname{rank} Y \\ &\quad + \operatorname{rank} [(I_k - YY^+)(D - CA^+B)(I_l - X^+X)], \end{aligned}$$

where $X \triangleq B - AA^+B$ and $Y \triangleq C - CA^+A$. Consequently,

$$\operatorname{rank} A + \operatorname{rank}(D - CA^+B) \leq \operatorname{rank} \begin{bmatrix} A & B \\ C & D \end{bmatrix},$$

and, if $AA^+B = B$ and $CA^+A = C$, then

$$\operatorname{rank} A + \operatorname{rank}(D - CA^+B) = \operatorname{rank} \begin{bmatrix} A & B \\ C & D \end{bmatrix}.$$

Finally, if $n = m$ and A is nonsingular, then

$$n + \operatorname{rank}(D - CA^{-1}B) = \operatorname{rank} \begin{bmatrix} A & B \\ C & D \end{bmatrix}.$$

(Proof: See [290, 968], Fact 2.11.8, and Fact 2.11.11.) (Remark: With certain restrictions the generalized inverses can be replaced by (1)-inverses.) (Remark: See Proposition 2.8.3 and Proposition 8.2.3.)

Fact 6.5.7. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{k \times l}$, and $C \in \mathbb{F}^{n \times l}$. Then,

$$\min_{X \in \mathbb{F}^{m \times l}, Y \in \mathbb{F}^{n \times k}} \operatorname{rank}(AX + YB + C) = \operatorname{rank} \begin{bmatrix} A & C \\ 0 & -B \end{bmatrix} - \operatorname{rank} A - \operatorname{rank} B.$$

Furthermore, X, Y is a minimizing solution if and only if there exist $U \in \mathbb{F}^{m \times k}$, $U_1 \in \mathbb{F}^{m \times l}$, and $U_2 \in \mathbb{F}^{n \times k}$, such that

$$X = -A^+C + UB + (I_m - A^+A)U_1,$$

$$Y = (AA^+ - I)CB^+ - AU + U_2(I_k - BB^+).$$

Finally, all such matrices $X \in \mathbb{F}^{m \times l}$ and $Y \in \mathbb{F}^{n \times k}$ satisfy

$$AX + YB + C = 0$$

if and only if

$$\operatorname{rank} \begin{bmatrix} A & C \\ 0 & -B \end{bmatrix} = \operatorname{rank} A + \operatorname{rank} B.$$

(Proof: See [1285, 1303].) (Remark: See Fact 5.10.20. Note that A and B are square in Fact 5.10.20.)

Fact 6.5.8. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and assume that $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$ is a projector. Then,

$$\operatorname{rank}(D - B^*A^+B) = \operatorname{rank} C - \operatorname{rank} B^*A^+B.$$

(Proof: See [1295].) (Remark: See [107].)

Fact 6.5.9. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then, the following statements are equivalent:

$$i) \operatorname{rank} \begin{bmatrix} A & B \end{bmatrix} = \operatorname{rank} A + \operatorname{rank} B.$$

- ii) $\mathcal{R}(A) \cap \mathcal{R}(B) = \emptyset$.
- iii) $\text{rank}(AA^* + BB^*) = \text{rank } A + \text{rank } B$.
- iv) $A^*(AA^* + BB^*)^+A$ is idempotent.
- v) $A^*(AA^* + BB^*)^+A = A^+A$.
- vi) $A^*(AA^* + BB^*)^+B = 0$.

(Proof: See [948, pp. 56, 57].) (Remark: See Fact 2.11.8.)

Fact 6.5.10. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$, and define the projectors $P \triangleq AA^+$ and $Q \triangleq BB^+$. Then, the following statements are equivalent:

- i) $\text{rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank } A + \text{rank } B = n$.
- ii) $P - Q$ is nonsingular.

In this case,

$$\begin{aligned} (P - Q)^{-1} &= (P - PQ)^+ + (PQ - Q)^+ \\ &= (P - QP)^+ + (QP - Q)^+ \\ &= P - Q + Q(P - QP)^+ - (Q - QP)^+P. \end{aligned}$$

(Proof: See [322].)

Fact 6.5.11. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{l \times n}$, $D \in \mathbb{F}^{l \times l}$, and assume that D is nonsingular. Then,

$$\text{rank } A = \text{rank}(A - BD^{-1}C) + \text{rank } BD^{-1}C$$

if and only if there exist matrices $X \in \mathbb{F}^{m \times l}$ and $Y \in \mathbb{F}^{l \times n}$ such that $B = AX$, $C = YA$, and $D = YAX$. (Proof: See [330].)

Fact 6.5.12. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times l}$. Then,

$$\text{rank } A + \text{rank}(D - CA^+B) = \text{rank} \begin{bmatrix} A^*AA^* & A^*B \\ CA^* & D \end{bmatrix}.$$

(Proof: See [1286].)

Fact 6.5.13. Let $A_{11} \in \mathbb{F}^{n \times m}$, $A_{12} \in \mathbb{F}^{n \times l}$, $A_{21} \in \mathbb{F}^{k \times m}$, and $A_{22} \in \mathbb{F}^{k \times l}$, and define $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \in \mathbb{F}^{(n+k) \times (m+l)}$ and $B \triangleq AA^+ = \begin{bmatrix} B_{11} & B_{12} \\ B_{12}^T & B_{22} \end{bmatrix}$, where $B_{11} \in \mathbb{F}^{n \times m}$, $B_{12} \in \mathbb{F}^{n \times l}$, $B_{21} \in \mathbb{F}^{k \times m}$, and $B_{22} \in \mathbb{F}^{k \times l}$. Then,

$$\text{rank } B_{12} = \text{rank} \begin{bmatrix} A_{11} & A_{12} \end{bmatrix} + \text{rank} \begin{bmatrix} A_{21} & A_{22} \end{bmatrix} - \text{rank } A.$$

(Proof: See [1308].) (Remark: See Fact 3.12.20 and Fact 3.13.12.)

Fact 6.5.14. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} \operatorname{rank} \begin{bmatrix} 0 & A \\ B & I \end{bmatrix} &= \operatorname{rank} A + \operatorname{rank} \begin{bmatrix} B & I - A^+A \end{bmatrix} \\ &= \operatorname{rank} \begin{bmatrix} A \\ I - BB^+ \end{bmatrix} + \operatorname{rank} B \\ &= \operatorname{rank} A + \operatorname{rank} B + \operatorname{rank} [(I - BB^+)(I - A^+A)] \\ &= n + \operatorname{rank} AB. \end{aligned}$$

Hence, the following statements hold:

- i) $\operatorname{rank} AB = \operatorname{rank} A + \operatorname{rank} B - n$ if and only if $(I - BB^+)(I - A^+A) = 0$.
- ii) $\operatorname{rank} AB = \operatorname{rank} A$ if and only if $\begin{bmatrix} B & I - A^+A \end{bmatrix}$ is right invertible.
- iii) $\operatorname{rank} AB = \operatorname{rank} B$ if and only if $\begin{bmatrix} A \\ I - BB^+ \end{bmatrix}$ is left invertible.

(Proof: See [968].) (Remark: The generalized inverses can be replaced by arbitrary (1)-inverses.)

Fact 6.5.15. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $C \in \mathbb{F}^{l \times k}$. Then,

$$\begin{aligned} \operatorname{rank} \begin{bmatrix} 0 & AB \\ BC & B \end{bmatrix} &= \operatorname{rank} B + \operatorname{rank} ABC \\ &= \operatorname{rank} AB + \operatorname{rank} BC \\ &\quad + \operatorname{rank} [(I - BC)(BC)^+]B[(I - (AB)^+(AB))]. \end{aligned}$$

Furthermore, the following statements are equivalent:

- i) $\operatorname{rank} \begin{bmatrix} 0 & AB \\ BC & B \end{bmatrix} = \operatorname{rank} AB + \operatorname{rank} BC$.
- ii) $\operatorname{rank} ABC = \operatorname{rank} AB + \operatorname{rank} BC - \operatorname{rank} B$.
- iii) There exist matrices $X \in \mathbb{F}^{k \times l}$ and $Y \in \mathbb{F}^{m \times n}$ such that

$$BCX + YAB = B.$$

(Proof: See [968, 1308] and Fact 5.10.20.) (Remark: This result is related to the Frobenius inequality. See Fact 2.11.14.)

Fact 6.5.16. Let $x, y \in \mathbb{R}^3$, and assume that x and y are linearly independent. Then,

$$\begin{bmatrix} x & y \end{bmatrix}^+ = \begin{bmatrix} x^+(I_3 - y\phi^T) \\ \phi^T \end{bmatrix},$$

where $x^+ = (x^T x)^{-1} x^T$, $\alpha \triangleq y^T(I - xx^+)y$, and $\phi \triangleq \alpha^{-1}(I - xx^+)y$. Now, let $x, y, z \in \mathbb{R}^3$, and assume that x and y are linearly independent. Then,

$$\begin{bmatrix} x & y & z \end{bmatrix}^+ = \begin{bmatrix} (I_2 - \beta w w^T) \begin{bmatrix} x & y \end{bmatrix}^+ \\ \beta w^T \begin{bmatrix} x & y \end{bmatrix}^+ \end{bmatrix},$$

where $w \triangleq [x \ y]^+ z$ and $\beta \triangleq 1/(1 + w^T w)$. (Proof: See [1319].)

Fact 6.5.17. Let $A \in \mathbb{F}^{n \times m}$ and $b \in \mathbb{F}^n$. Then,

$$\begin{bmatrix} A & b \end{bmatrix}^+ = \begin{bmatrix} A^+(I_n - b\phi^*) \\ \phi^* \end{bmatrix}$$

and

$$\begin{bmatrix} b & A \end{bmatrix}^+ = \begin{bmatrix} \phi^* \\ A^+(I_n - b\phi^*) \end{bmatrix},$$

where

$$\phi \triangleq \begin{cases} (b - AA^+b)^{+*}, & b \neq AA^+b, \\ \gamma^{-1}(AA^*)^+b, & b = AA^+b. \end{cases}$$

and $\gamma \triangleq 1 + b^*(AA^*)^+b$. (Proof: See [15, p. 44], [481, p. 270], or [1186, p. 148].) (Remark: This result is due to Greville.)

Fact 6.5.18. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\begin{bmatrix} A & B \end{bmatrix}^+ = \begin{bmatrix} A^+ - A^+B(C^+ + D) \\ C^+ + D \end{bmatrix},$$

where

$$C \triangleq (I - AA^+)B$$

and

$$D \triangleq (I - C^+C)[I + (I - C^+C)B^*(AA^*)^+B(I - C^+C)]^{-1}B^*(AA^*)^+(I - BC^+).$$

Furthermore,

$$\begin{bmatrix} A & B \end{bmatrix}^+ = \begin{cases} \begin{bmatrix} A^*(AA^* + BB^*)^{-1} \\ B^*(AA^* + BB^*)^{-1} \end{bmatrix}, & \text{rank} \begin{bmatrix} A & B \end{bmatrix} = n, \\ \begin{bmatrix} A^*A & A^*B \\ B^*A & B^*B \end{bmatrix}^{-1} \begin{bmatrix} A^* \\ B^* \end{bmatrix}, & \text{rank} \begin{bmatrix} A & B \end{bmatrix} = m + l, \\ \begin{bmatrix} A^*(AA^*)^{-1}(I - BE) \\ E \end{bmatrix}, & \text{rank } A = n, \end{cases}$$

where

$$E \triangleq [I + B^*(AA^*)^{-1}B]^{-1}B^*(AA^*)^{-1}.$$

(Proof: See [338] or [947, p. 14].) (Remark: If $\begin{bmatrix} A & B \end{bmatrix}$ is square and nonsingular and $A^*B = 0$, then the second expression yields Fact 2.17.8.)

Fact 6.5.19. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\text{rank} \left(\begin{bmatrix} A & B \end{bmatrix}^+ - \begin{bmatrix} A^+ \\ B^+ \end{bmatrix} \right) = \text{rank} \begin{bmatrix} AA^*B & BB^*A \end{bmatrix}.$$

Hence, if $A^*B = 0$, then

$$\begin{bmatrix} A & B \end{bmatrix}^+ = \begin{bmatrix} A^+ \\ B^+ \end{bmatrix}.$$

(Proof: See [1289].)

Fact 6.5.20. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then, the following statements are equivalent:

$$i) \begin{bmatrix} A & B \end{bmatrix} \begin{bmatrix} A & B \end{bmatrix}^+ = \frac{1}{2}(AA^+ + BB^+).$$

$$ii) \mathcal{R}(A) = \mathcal{R}(B).$$

Furthermore, the following statements are equivalent:

$$iii) \begin{bmatrix} A & B \end{bmatrix}^+ = \frac{1}{2} \begin{bmatrix} A^+ \\ B^+ \end{bmatrix}.$$

$$iv) AA^* = BB^*.$$

(Proof: See [1300].)

Fact 6.5.21. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{k \times l}$. Then,

$$\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}^+ = \begin{bmatrix} A^+ & 0 \\ 0 & B^+ \end{bmatrix}.$$

Fact 6.5.22. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\begin{bmatrix} I_n & A \\ 0_{m \times n} & 0_{m \times m} \end{bmatrix}^+ = \begin{bmatrix} (I_n + AA^*)^{-1} & 0_{n \times m} \\ A^*(I_n + AA^*)^{-1} & 0_{m \times m} \end{bmatrix}.$$

(Proof: See [17, 1326].)

Fact 6.5.23. Let $A \in \mathbb{F}^{n \times n}$, let $B \in \mathbb{F}^{n \times m}$, and assume that $BB^* = I$. Then,

$$\begin{bmatrix} A & B \\ B^* & 0 \end{bmatrix}^+ = \begin{bmatrix} 0 & B \\ B^* & -B^*AB \end{bmatrix}.$$

(Proof: See [447, p. 237].)

Fact 6.5.24. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $B \in \mathbb{F}^{n \times m}$. Then,

$$\begin{bmatrix} A & B \\ B^* & 0 \end{bmatrix}^+ = \begin{bmatrix} C^+ - C^+BD^+B^*C^+ & C^+BD^+ \\ (C^+BD^+)^* & DD^+ - D^+ \end{bmatrix},$$

where

$$C \triangleq A + BB^*, \quad D \triangleq B^*C^+B.$$

(Proof: See [948, p. 58].) (Remark: Representations for the generalized inverse of a partitioned matrix are given in [174, Chapter 5] and [105, 112, 134, 172, 277, 283, 296, 595, 643, 645, 736, 904, 996, 997, 999, 1000, 1001, 1046, 1120, 1137, 1278, 1310, 1418].) (Problem: Show that the generalized inverses in this result and in Fact 6.5.23 are identical when A is positive semidefinite and $BB^* = I$.)

Fact 6.5.25. Let $A \in \mathbb{F}^{n \times n}$, $x, y \in \mathbb{F}^n$, and $a \in \mathbb{F}$, and assume that $x \in \mathcal{R}(A)$. Then,

$$\begin{bmatrix} A & x \\ y^T & a \end{bmatrix} = \begin{bmatrix} I & 0 \\ y^T & 1 \end{bmatrix} \begin{bmatrix} A & 0 \\ y^T - y^T A & a - y^T A^+ x \end{bmatrix} \begin{bmatrix} I & A^+ x \\ 0 & 1 \end{bmatrix}.$$

(Remark: See Fact 2.16.2 and Fact 2.14.9, and note that $x = AA^+x$.) (Problem: Obtain a factorization for the case $x \notin \mathcal{R}(A)$.)

Fact 6.5.26. Let $A \in \mathbb{F}^{n \times m}$, assume that A is partitioned as

$$A = \begin{bmatrix} A_1 \\ \vdots \\ A_k \end{bmatrix},$$

and define

$$B \triangleq \begin{bmatrix} A_1^+ & \cdots & A_k^+ \end{bmatrix}.$$

Then, the following statements hold:

- i) $\det AB = 0$ if and only if $\text{rank } A < n$.
- ii) $0 < \det AB \leq 1$ if and only if $\text{rank } A = n$.
- iii) If $\text{rank } A = n$, then

$$\det AB = \frac{\det AA^*}{\prod_{i=1}^k \det A_i A_i^*},$$

and thus

$$\det AA^* \leq \prod_{i=1}^k \det A_i A_i^*.$$

- iv) $\det AB = 1$ if and only if $AB = I$.
- v) AB is group invertible.
- vi) Every eigenvalue of AB is nonnegative.
- vii) $\text{rank } A = \text{rank } B = \text{rank } AB = \text{rank } BA$.

Now, assume that $\text{rank } A = \sum_{i=1}^k \text{rank } A_i$, and let β denote the product of the positive eigenvalues of AB . Then, the following statements hold:

- viii) $0 < \beta \leq 1$.
- ix) $\beta = 1$ if and only if $B = A^+$.

(Proof: See [875, 1247].) (Remark: Result iii) yields Hadamard's inequality as given by Fact 8.13.34 in the case that A is square and each A_i has a single row.)

Fact 6.5.27. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\begin{aligned} \det \begin{bmatrix} A^*A & B^*A \\ B^*A & B^*B \end{bmatrix} &= \det(A^*A) \det[B^*(I - AA^+)B] \\ &= \det(B^*B) \det[A^*(I - BB^+)A]. \end{aligned}$$

(Remark: See Fact 2.14.25.)

Fact 6.5.28. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, and $D \in \mathbb{F}^{m \times m}$, assume that either $\text{rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank } A$ or $\text{rank} \begin{bmatrix} A \\ C \end{bmatrix} = \text{rank } A$, and let $A^- \in \mathbb{F}^{n \times n}$ be a (1)-inverse of A . Then,

$$\det \begin{bmatrix} A & B \\ C & D \end{bmatrix} = (\det A) \det(D - CA^-B).$$

(Proof: See [144, p. 266].)

Fact 6.5.29. Let $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, $B \in \mathbb{F}^{(n+m) \times l}$, $C \in \mathbb{F}^{l \times (n+m)}$, $D \in \mathbb{F}^{l \times l}$, and $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix}$, and assume that A and A_{11} are nonsingular. Then,

$$A|\mathcal{A} = (A_{11}|A)|(A_{11}|\mathcal{A}).$$

(Proof: See [1098, pp. 18, 19].) (Remark: This result is the *Crabtree-Haynsworth quotient formula*. See [717].) (Remark: Extensions are given in [1495].) (Problem: Extend this result to the case in which either A or A_{11} is singular.)

Fact 6.5.30. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

- i) $A \overset{\text{rs}}{\leq} B$.
- ii) $AA^+B = BA^+A = BA^+B = B$.
- iii) $\text{rank } A = \text{rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank} \begin{bmatrix} A \\ B \end{bmatrix}$ and $BA^+B = B$.

(Proof: See [1184, p. 45].) (Remark: See Fact 8.20.7.)

6.6 Facts on the Drazin and Group Generalized Inverses

Fact 6.6.1. Let $A_1, \dots, A_k \in \mathbb{F}^{n \times m}$. Then,

$$(A_1 + \dots + A_k)^D = \frac{1}{k} \begin{bmatrix} I_n & \dots & I_n \end{bmatrix} \begin{bmatrix} A_1 & A_2 & \dots & A_k \\ A_k & A_1 & \dots & A_{k-1} \\ \vdots & \vdots & \ddots & \vdots \\ A_2 & A_3 & \dots & A_1 \end{bmatrix}^D \begin{bmatrix} I_m \\ \vdots \\ I_m \end{bmatrix}.$$

(Proof: See [1282].) (Remark: See Fact 6.5.2.)

Fact 6.6.2. Let $A \in \mathbb{F}^{n \times n}$. Then, $X = A^D$ is the unique matrix satisfying

$$\text{rank} \begin{bmatrix} A & AA^D \\ A^D A & X \end{bmatrix} = \text{rank } A.$$

(Remark: See Fact 2.17.10 and Fact 6.3.30.) (Proof: See [1417, 1496].)

Fact 6.6.3. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $AB = 0$. Then,

$$(AB)^D = A(BA)^{2D}B.$$

(Remark: This result is *Cline's formula*.)

Fact 6.6.4. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $AB = BA$. Then,

$$(AB)^D = B^D A^D,$$

$$A^D B = B A^D,$$

$$AB^D = B^D A.$$

Fact 6.6.5. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $AB = BA = 0$. Then,

$$(A + B)^D = A^D + B^D.$$

(Proof: See [653].) (Remark: This result is due to Drazin.)

Fact 6.6.6. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{ind } A = \text{rank } A = 1$. Then,

$$A^\# = (\text{tr } A^2)^{-1} A.$$

Consequently, if $x, y \in \mathbb{F}^n$ satisfy $x^* y \neq 0$, then

$$(xy^*)^\# = (x^* y)^{-2} xy^*.$$

In particular,

$$1_{n \times n}^\# = n^{-2} 1_{n \times n}.$$

Fact 6.6.7. Let $A \in \mathbb{F}^{n \times n}$, and let $k \triangleq \text{ind } A$. Then,

$$A^D = A^k (A^{2k+1})^+ A^k.$$

If, in particular, $\text{ind } A \leq 1$, then

$$A^\# = A(A^3)^+ A.$$

(Proof: See [174, pp. 165, 174].)

Fact 6.6.8. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is range Hermitian.
- ii) $A^+ = A^D$.
- iii) $\text{ind } A \leq 1$, and $A^+ = A^\#$.
- iv) $\text{ind } A \leq 1$, and $A^* A^\# A + A A^\# A^* = 2A^*$.
- v) $\text{ind } A \leq 1$, and $A^+ A^\# A + A A^\# A^+ = 2A^+$.

(Proof: See [323].) (Remark: See Fact 6.3.10.)

Fact 6.6.9. Let $A \in \mathbb{F}^{n \times n}$, assume that A is group invertible, and let $S, B \in \mathbb{F}^{n \times n}$, where S is nonsingular, B is a Jordan canonical form of A , and $A = SBS^{-1}$. Then,

$$A^\# = SB^\# S^{-1} = SB^+ S^{-1}.$$

(Proof: Since B is range Hermitian, it follows from Fact 6.6.8 that $B^\# = B^+$. See [174, p. 158].)

Fact 6.6.10. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is normal.

ii) $\text{ind } A \leq 1$, and $A^\#A^* = A^*A^\#$.

(Proof: See [323].) (Remark: See Fact 3.7.12, Fact 3.11.4, Fact 5.15.4, and Fact 6.3.16.)

Fact 6.6.11. Let $A \in \mathbb{F}^{n \times n}$, and let $k \geq 1$. Then, the following statements are equivalent:

i) $k \geq \text{ind } A$.

ii) $\lim_{\alpha \rightarrow 0} \alpha^k (A + \alpha I)^{-1}$ exists.

iii) $\lim_{\alpha \rightarrow 0} (A^{k+1} + \alpha I)^{-1} A^k$ exists.

In this case,

$$A^D = \lim_{\alpha \rightarrow 0} (A^{k+1} + \alpha I)^{-1} A^k$$

and

$$\lim_{\alpha \rightarrow 0} \alpha^k (A + \alpha I)^{-1} = \begin{cases} (-1)^{k-1} (I - AA^D) A^{k-1}, & k = \text{ind } A > 0, \\ A^{-1}, & k = \text{ind } A = 0, \\ 0, & k > \text{ind } A. \end{cases}$$

(Proof: See [999].)

Fact 6.6.12. Let $A \in \mathbb{F}^{n \times n}$, let $r \triangleq \text{rank } A$, let $B \in \mathbb{R}^{n \times r}$ and $C \in \mathbb{R}^{r \times n}$, and assume that $A = BC$. Then, A is group invertible if and only if BA is nonsingular. In this case,

$$A^\# = B(CB)^{-2}C.$$

(Proof: See [174, p. 157].) (Remark: This result is due to Cline.)

Fact 6.6.13. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$. If A and C are singular, then $\text{ind} \begin{bmatrix} A & B \\ 0 & C \end{bmatrix} = 1$ if and only if $\text{ind } A = \text{ind } C = 1$, and $(I - AA^D)B(I - CC^D) = 0$. (Proof: See [999].) (Remark: See Fact 5.14.32.)

Fact 6.6.14. Let $A \in \mathbb{F}^{n \times n}$. Then, A is group invertible if and only if $\lim_{\alpha \rightarrow 0} (A + \alpha I)^{-1} A$ exists. In this case,

$$\lim_{\alpha \rightarrow 0} (A + \alpha I)^{-1} A = AA^\#.$$

(Proof: See [283, p. 138].)

Fact 6.6.15. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonzero and group invertible, let $r \triangleq \text{rank } A$, define $B \triangleq \text{diag}[\sigma_1(A), \dots, \sigma_r(A)]$, and let $S \in \mathbb{F}^{n \times n}$, $K \in \mathbb{F}^{r \times r}$, and $L \in \mathbb{F}^{r \times (n-r)}$ be such that S is unitary,

$$KK^* + LL^* = I_r,$$

and

$$A = S \begin{bmatrix} BK & BL \\ 0_{(n-r) \times r} & 0_{(n-r) \times (n-r)} \end{bmatrix} S^*.$$

Then,

$$A^\# = S \begin{bmatrix} K^{-1}B^{-1} & K^{-1}B^{-1}K^{-1}L \\ 0_{(n-r) \times r} & 0_{(n-r) \times (n-r)} \end{bmatrix} S^*.$$

(Proof: See [115, 651].) (Remark: See Fact 5.9.28 and Fact 6.3.15.)

Fact 6.6.16. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is range Hermitian.
- ii) A is group invertible and $AA^+A^+ = A^\#$.
- iii) A is group invertible and $AA^\#A^+ = A^\#$.
- iv) A is group invertible and $A^*AA^\# = A^*$.
- v) A is group invertible and $A^+AA^\# = A^+$.
- vi) A is group invertible and $A^\#A^+A = A^+$.
- vii) A is group invertible and $AA^\# = A^+A$.
- viii) A is group invertible and $A^*A^+ = A^*A^\#$.
- ix) A is group invertible and $A^+A^* = A^\#A^*$.
- x) A is group invertible and $A^+A^+ = A^+A^\#$.
- xi) A is group invertible and $A^+A^+ = A^\#A^+$.
- xii) A is group invertible and $A^+A^+ = A^\#A^\#$.
- xiii) A is group invertible and $A^+A^\# = A^\#A^\#$.
- xiv) A is group invertible and $A^\#A^+ = A^\#A^\#$.
- xv) A is group invertible and $A^+A^\# = A^\#A^+$.
- xvi) A is group invertible and $AA^+A^* = A^*AA^+$.
- xvii) A is group invertible and $AA^+A^\# = A^+A^\#A$.
- xviii) A is group invertible and $AA^+A^\# = A^\#AA^+$.
- ixx) A is group invertible and $AA^\#A^* = A^*AA^\#$.
- xx) A is group invertible and $AA^\#A^+ = A^+AA^\#$.
- xxi) A is group invertible and $AA^\#A^+ = A^\#A^+A$.
- xxii) A is group invertible and $A^*A^+A = A^+AA^*$.
- xxiii) A is group invertible and $A^+AA^\# = A^\#A^+A$.
- xxiv) A is group invertible and $A^+A^+A^\# = A^+A^\#A^+$.
- xxv) A is group invertible and $A^+A^+A^\# = A^\#A^+A^+$.
- xxvi) A is group invertible and $A^+A^\#A^+ = A^\#A^+A^+$.
- xxvii) A is group invertible and $A^+A^\#A^\# = A^\#A^+A^\#$.
- xxviii) A is group invertible and $A^+A^\#A^\# = A^\#A^\#A^+$.

xxix) A is group invertible and $A^\#A^\#A^+ = A^\#A^+A^\#$.

(Proof: See [115].)

Fact 6.6.17. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i)* A is normal.
- ii)* A is group invertible and $A^*A^+ = A^\#A^*$.
- iii)* A is group invertible and $A^*A^\# = A^+A^*$.
- iv)* A is group invertible and $A^*A^\# = A^\#A^*$.
- v)* A is group invertible and $AA^*A^\# = A^*A^\#A$.
- vi)* A is group invertible and $AA^*A^\# = A^\#AA^*$.
- vii)* A is group invertible and $AA^\#A^* = A^\#A^*A$.
- viii)* A is group invertible and $A^*AA^\# = A^\#A^*A$.
- ix)* A is group invertible and $A^{*2}A^\# = A^*A^\#A^*$.
- x)* A is group invertible and $A^*A^+A^\# = A^\#A^*A^+$.
- xi)* A is group invertible and $A^*A^\#A^* = A^\#A^{*2}$.
- xii)* A is group invertible and $A^*A^\#A^+ = A^+A^*A^\#$.
- xiii)* A is group invertible and $A^*A^\#A^\# = A^\#A^*A^\#$.
- xiv)* A is group invertible and $A^+A^*A^\# = A^\#A^+A^*$.
- xv)* A is group invertible and $A^+A^\#A^* = A^\#A^+A^*$.
- xvi)* A is group invertible and $A^\#A^*A^\# = A^\#A^\#A^*$.

(Proof: See [115].)

Fact 6.6.18. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i)* A is Hermitian.
- ii)* A is group invertible and $AA^\# = A^*A^+$.
- iii)* A is group invertible and $AA^\# = A^*A^\#$.
- iv)* A is group invertible and $AA^\# = A^+A^*$.
- v)* A is group invertible and $A^+A = A^\#A^*$.
- vi)* A is group invertible and $A^*AA^\# = A$.
- vii)* A is group invertible and $A^{*2}A^\# = A^*$.
- viii)* A is group invertible and $A^*A^+A^+ = A^\#$.
- ix)* A is group invertible and $A^*A^+A^\# = A^+$.
- x)* A is group invertible and $A^*A^+A^\# = A^\#$.
- xi)* A is group invertible and $A^*A^\#A^\# = A^\#$.

xi) A is group invertible and $A^\# A^* A^\# = A^+$.

(Proof: See [115].)

Fact 6.6.19. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are group invertible, and consider the following conditions:

i) $ABA = B$.

ii) $BAB = A$.

iii) $A^2 = B^2$.

Then, if two of the above conditions are satisfied, then the third condition is satisfied. Furthermore, if *i)*–*iii)* are satisfied, then the following statements hold:

iv) A and B are group invertible.

v) $A^\# = A^3$ and $B^\# = B^3$.

vi) $A^5 = A$ and $B^5 = B$.

vii) $A^4 = B^4 = (AB)^4$.

viii) If A and B are nonsingular, then $A^4 = B^4 = (AB)^4 = I$.

(Proof: See [469].)

Fact 6.6.20. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, assume that A is positive, define $B \triangleq \text{sprad}(A)I - A$, let $x, y \in \mathbb{R}^n$ be positive, and assume that $Ax = \text{sprad}(A)x$ and $A^T y = \text{sprad}(A)y$. Then, the following statements hold:

i) $B + \frac{1}{x^T y} xy^T$ is nonsingular.

ii) $B^\# = (B + \frac{1}{x^T y} xy^T)^{-1} (I - \frac{1}{x^T y} xy^T)$.

iii) $I - BB^\# = \frac{1}{x^T y} xy^T$.

iv) $B^\# = \lim_{k \rightarrow \infty} \left[\sum_{i=0}^{k-1} \frac{1}{[\text{sprad}(A)]^i} A^i - \frac{k}{x^T y} xy^T \right]$.

(Proof: See [1148, p. 9-4].) (Remark: See Fact 4.11.5.)

6.7 Notes

A brief history of the generalized inverse is given in [173] and [174, p. 4]. The proof of the uniqueness of A^+ is given in [948, p. 32]. Additional books on generalized inverses include [174, 245, 1118, 1396]. The terminology “range Hermitian” is used in [174]; the terminology “EP” is more common. Generalized inverses are widely used in least squares methods; see [237, 283, 876]. Applications to singular differential equations are considered in [282]. Applications to Markov chains are discussed in [737].

Chapter Seven

Kronecker and Schur Algebra

In this chapter we introduce Kronecker matrix algebra, which is useful for solving linear matrix equations.

7.1 Kronecker Product

For $A \in \mathbb{F}^{n \times m}$ define the *vec* operator as

$$\text{vec } A \triangleq \begin{bmatrix} \text{col}_1(A) \\ \vdots \\ \text{col}_m(A) \end{bmatrix} \in \mathbb{F}^{nm}, \quad (7.1.1)$$

which is the column vector of size $nm \times 1$ obtained by stacking the columns of A . We recover A from $\text{vec } A$ by writing

$$A = \text{vec}^{-1}(\text{vec } A). \quad (7.1.2)$$

Proposition 7.1.1. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then,

$$\text{tr } AB = (\text{vec } A^T)^T \text{vec } B = (\text{vec } B^T)^T \text{vec } A. \quad (7.1.3)$$

Proof. Note that

$$\begin{aligned} \text{tr } AB &= \sum_{i=1}^n \text{row}_i(A) \text{col}_i(B) \\ &= \sum_{i=1}^n [\text{col}_i(A^T)]^T \text{col}_i(B) \\ &= \begin{bmatrix} \text{col}_1^T(A^T) & \cdots & \text{col}_n^T(A^T) \end{bmatrix} \begin{bmatrix} \text{col}_1(B) \\ \vdots \\ \text{col}_n(B) \end{bmatrix} \\ &= (\text{vec } A^T)^T \text{vec } B. \end{aligned}$$

□

Next, we introduce the Kronecker product.

Definition 7.1.2. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times k}$. Then, the *Kronecker product* $A \otimes B \in \mathbb{F}^{nl \times mk}$ of A is the partitioned matrix

$$A \otimes B \triangleq \begin{bmatrix} A_{(1,1)}B & A_{(1,2)}B & \cdots & A_{(1,m)}B \\ \vdots & \vdots & \ddots & \vdots \\ A_{(n,1)}B & A_{(n,2)}B & \cdots & A_{(n,m)}B \end{bmatrix}. \quad (7.1.4)$$

Unlike matrix multiplication, the Kronecker product $A \otimes B$ does not entail a restriction on either the size of A or the size of B .

The following results are immediate consequences of the definition of the Kronecker product.

Proposition 7.1.3. Let $\alpha \in \mathbb{F}$, $A \in \mathbb{F}^{n \times m}$, and $B \in \mathbb{F}^{l \times k}$. Then,

$$A \otimes (\alpha B) = (\alpha A) \otimes B = \alpha(A \otimes B), \quad (7.1.5)$$

$$\overline{A \otimes B} = \overline{A} \otimes \overline{B}, \quad (7.1.6)$$

$$(A \otimes B)^T = A^T \otimes B^T, \quad (7.1.7)$$

$$(A \otimes B)^* = A^* \otimes B^*. \quad (7.1.8)$$

Proposition 7.1.4. Let $A, B \in \mathbb{F}^{n \times m}$ and $C \in \mathbb{F}^{l \times k}$. Then,

$$(A + B) \otimes C = A \otimes C + B \otimes C \quad (7.1.9)$$

and

$$C \otimes (A + B) = C \otimes A + C \otimes B. \quad (7.1.10)$$

Proposition 7.1.5. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{l \times k}$, and $C \in \mathbb{F}^{p \times q}$. Then,

$$A \otimes (B \otimes C) = (A \otimes B) \otimes C. \quad (7.1.11)$$

Hence, we write $A \otimes B \otimes C$ for $A \otimes (B \otimes C)$ and $(A \otimes B) \otimes C$.

The next result illustrates a useful form of compatibility between matrix multiplication and the Kronecker product.

Proposition 7.1.6. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{l \times k}$, $C \in \mathbb{F}^{m \times q}$, and $D \in \mathbb{F}^{k \times p}$. Then,

$$(A \otimes B)(C \otimes D) = AC \otimes BD. \quad (7.1.12)$$

Proof. Note that the ij block of $(A \otimes B)(C \otimes D)$ is given by

$$\begin{aligned} [(A \otimes B)(C \otimes D)]_{ij} &= \begin{bmatrix} A_{(i,1)}B & \cdots & A_{(i,m)}B \end{bmatrix} \begin{bmatrix} C_{(1,j)}D \\ \vdots \\ C_{(m,j)}D \end{bmatrix} \\ &= \sum_{k=1}^m A_{(i,k)}C_{(k,j)}BD = (AC)_{(i,j)}BD \\ &= (AC \otimes BD)_{ij}. \end{aligned} \quad \square$$

Next, we consider the inverse of a Kronecker product.

Proposition 7.1.7. Assume that $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$ are nonsingular. Then,

$$(A \otimes B)^{-1} = A^{-1} \otimes B^{-1}. \quad (7.1.13)$$

Proof. Note that

$$(A \otimes B)(A^{-1} \otimes B^{-1}) = AA^{-1} \otimes BB^{-1} = I_n \otimes I_m = I_{nm}. \quad \square$$

Proposition 7.1.8. Let $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$. Then,

$$xy^T = x \otimes y^T = y^T \otimes x \quad (7.1.14)$$

and

$$\text{vec } xy^T = y \otimes x. \quad (7.1.15)$$

The following result concerns the vec of the product of three matrices.

Proposition 7.1.9. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $C \in \mathbb{F}^{l \times k}$. Then,

$$\text{vec}(ABC) = (C^T \otimes A)\text{vec } B. \quad (7.1.16)$$

Proof. Using (7.1.12) and (7.1.15), it follows that

$$\begin{aligned} \text{vec } ABC &= \text{vec} \sum_{i=1}^l A \text{col}_i(B) e_i^T C = \sum_{i=1}^l \text{vec} \left[A \text{col}_i(B) (C^T e_i)^T \right] \\ &= \sum_{i=1}^l [C^T e_i] \otimes [A \text{col}_i(B)] = (C^T \otimes A) \sum_{i=1}^l e_i \otimes \text{col}_i(B) \\ &= (C^T \otimes A) \sum_{i=1}^l \text{vec} [\text{col}_i(B) e_i^T] = (C^T \otimes A) \text{vec } B. \end{aligned} \quad \square$$

The following result concerns the eigenvalues and eigenvectors of the Kronecker product of two matrices.

Proposition 7.1.10. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$\text{mspec}(A \otimes B) = \{\lambda\mu : \lambda \in \text{mspec}(A), \mu \in \text{mspec}(B)\}_{\text{ms}}. \quad (7.1.17)$$

If, in addition, $x \in \mathbb{C}^n$ is an eigenvector of A associated with $\lambda \in \text{spec}(A)$ and $y \in \mathbb{C}^n$ is an eigenvector of B associated with $\mu \in \text{spec}(B)$, then $x \otimes y$ is an eigenvector of $A \otimes B$ associated with $\lambda\mu$.

Proof. Using (7.1.12), we have

$$(A \otimes B)(x \otimes y) = (Ax) \otimes (By) = (\lambda x) \otimes (\mu y) = \lambda\mu(x \otimes y). \quad \square$$

Proposition 7.1.10 shows that $\text{mspec}(A \otimes B) = \text{mspec}(B \otimes A)$. Consequently, it follows that $\det(A \otimes B) = \det(B \otimes A)$ and $\text{tr}(A \otimes B) = \text{tr}(B \otimes A)$. The following results are generalizations of these identities.

Proposition 7.1.11. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$\det(A \otimes B) = \det(B \otimes A) = (\det A)^m (\det B)^n. \quad (7.1.18)$$

Proof. Let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$ and $\text{mspec}(B) = \{\mu_1, \dots, \mu_m\}_{\text{ms}}$. Then, Proposition 7.1.10 implies that

$$\begin{aligned} \det(A \otimes B) &= \prod_{i,j=1}^{n,m} \lambda_i \mu_j = \left(\lambda_1^m \prod_{j=1}^m \mu_j \right) \cdots \left(\lambda_n^m \prod_{j=1}^m \mu_j \right) \\ &= (\lambda_1 \cdots \lambda_n)^m (\mu_1 \cdots \mu_m)^n = (\det A)^m (\det B)^n. \end{aligned} \quad \square$$

Proposition 7.1.12. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$\text{tr}(A \otimes B) = \text{tr}(B \otimes A) = (\text{tr } A)(\text{tr } B). \quad (7.1.19)$$

Proof. Note that

$$\begin{aligned} \text{tr}(A \otimes B) &= \text{tr}(A_{(1,1)}B) + \cdots + \text{tr}(A_{(n,n)}B) \\ &= [A_{(1,1)} + \cdots + A_{(n,n)}] \text{tr } B \\ &= (\text{tr } A)(\text{tr } B). \end{aligned} \quad \square$$

Next, define the *Kronecker permutation matrix* $P_{n,m} \in \mathbb{F}^{nm \times nm}$ by

$$P_{n,m} \triangleq \sum_{i,j=1}^{n,m} E_{i,j,n \times m} \otimes E_{j,i,m \times n}. \quad (7.1.20)$$

Proposition 7.1.13. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\text{vec } A^T = P_{n,m} \text{vec } A. \quad (7.1.21)$$

7.2 Kronecker Sum and Linear Matrix Equations

Next, we define the Kronecker sum of two square matrices.

Definition 7.2.1. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then, the *Kronecker sum* $A \oplus B \in \mathbb{F}^{nm \times nm}$ of A and B is

$$A \oplus B \triangleq A \otimes I_m + I_n \otimes B. \quad (7.2.1)$$

Proposition 7.2.2. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{l \times l}$. Then,

$$A \oplus (B \oplus C) = (A \oplus B) \oplus C. \quad (7.2.2)$$

Hence, we write $A \oplus B \oplus C$ for $A \oplus (B \oplus C)$ and $(A \oplus B) \oplus C$.

Proposition 7.1.10 shows that, if $\lambda \in \text{spec}(A)$ and $\mu \in \text{spec}(B)$, then $\lambda\mu \in \text{spec}(A \otimes B)$. Next, we present an analogous result involving Kronecker sums.

Proposition 7.2.3. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$\text{mspec}(A \oplus B) = \{\lambda + \mu: \lambda \in \text{mspec}(A), \mu \in \text{mspec}(B)\}_{\text{ms}}. \quad (7.2.3)$$

Now, let $x \in \mathbb{C}^n$ be an eigenvector of A associated with $\lambda \in \text{spec}(A)$, and let $y \in \mathbb{C}^m$ be an eigenvector of B associated with $\mu \in \text{spec}(B)$. Then, $x \otimes y$ is an eigenvector of $A \oplus B$ associated with $\lambda + \mu$.

Proof. Note that

$$\begin{aligned} (A \oplus B)(x \otimes y) &= (A \otimes I_m)(x \otimes y) + (I_n \otimes B)(x \otimes y) \\ &= (Ax \otimes y) + (x \otimes By) = (\lambda x \otimes y) + (x \otimes \mu y) \\ &= \lambda(x \otimes y) + \mu(x \otimes y) = (\lambda + \mu)(x \otimes y). \end{aligned} \quad \square$$

The next result concerns the existence and uniqueness of solutions to *Sylvester's equation*. See Fact 5.10.21 and Proposition 11.9.3.

Proposition 7.2.4. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{n \times m}$. Then, $X \in \mathbb{F}^{n \times m}$ satisfies

$$AX + XB + C = 0 \quad (7.2.4)$$

if and only if X satisfies

$$(B^T \oplus A) \text{vec } X + \text{vec } C = 0. \quad (7.2.5)$$

Consequently, $B^T \oplus A$ is nonsingular if and only if there exists a unique matrix $X \in \mathbb{F}^{n \times m}$ satisfying (7.2.4). In this case, X is given by

$$X = -\text{vec}^{-1} \left[(B^T \oplus A)^{-1} \text{vec } C \right]. \quad (7.2.6)$$

Furthermore, $B^T \oplus A$ is singular and $\text{rank } B^T \oplus A = \text{rank} \begin{bmatrix} B^T \oplus A & \text{vec } C \end{bmatrix}$ if and only if there exist infinitely many matrices $X \in \mathbb{F}^{n \times m}$ satisfying (7.5.8). In this case, the set of solutions of (7.2.4) is given by $X + \mathcal{N}(B^T \oplus A)$.

Proof. Note that (7.2.4) is equivalent to

$$\begin{aligned} 0 &= \text{vec}(AXI + IXB) + \text{vec } C = (I \otimes A)\text{vec } X + (B^T \otimes I)\text{vec } X + \text{vec } C \\ &= (B^T \otimes I + I \otimes A)\text{vec } X + \text{vec } C = (B^T \oplus A)\text{vec } X + \text{vec } C, \end{aligned}$$

which yields (7.2.5). The remaining results follow from Corollary 2.6.7. $\square$

For the following corollary, note Fact 5.10.21.

Corollary 7.2.5. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{n \times m}$, and assume that $\text{spec}(A)$ and $\text{spec}(-B)$ are disjoint. Then, there exists a unique matrix $X \in \mathbb{F}^{n \times m}$ satisfying (7.2.4). Furthermore, the matrices $\begin{bmatrix} A & 0 \\ 0 & -B \end{bmatrix}$ and $\begin{bmatrix} A & C \\ 0 & -B \end{bmatrix}$ are similar and satisfy

$$\begin{bmatrix} A & C \\ 0 & -B \end{bmatrix} = \begin{bmatrix} I & X \\ 0 & I \end{bmatrix} \begin{bmatrix} A & 0 \\ 0 & -B \end{bmatrix} \begin{bmatrix} I & -X \\ 0 & I \end{bmatrix}. \quad (7.2.7)$$

7.3 Schur Product

An alternative form of vector and matrix multiplication is given by the *Schur product*. If $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times m}$, then $A \circ B \in \mathbb{F}^{n \times m}$ is defined by

$$(A \circ B)_{(i,j)} \triangleq A_{(i,j)} B_{(i,j)}, \quad (7.3.1)$$

that is, $A \circ B$ is formed by means of entry-by-entry multiplication. For matrices $A, B, C \in \mathbb{F}^{n \times m}$, the commutative, associative, and distributive identities

$$A \circ B = B \circ A, \quad (7.3.2)$$

$$A \circ (B \circ C) = (A \circ B) \circ C, \quad (7.3.3)$$

$$A \circ (B + C) = A \circ B + A \circ C \quad (7.3.4)$$

hold. For a real scalar $\alpha \geq 0$ and $A \in \mathbb{F}^{n \times m}$, the *Schur power* $A^{\circ\alpha}$ is defined by

$$(A^{\circ\alpha})_{(i,j)} \triangleq (A_{(i,j)})^\alpha. \quad (7.3.5)$$

Thus, $A^{\circ 2} = A \circ A$. Note that $A^{\circ 0} = 1_{n \times m}$. Furthermore, $\alpha < 0$ is allowed if A has no zero entries. In particular, $A^{\circ -1}$ is the matrix whose entries are the reciprocals of the entries of A . For all $A \in \mathbb{F}^{n \times m}$,

$$A \circ 1_{n \times m} = 1_{n \times m} \circ A = A. \quad (7.3.6)$$

Finally, if A is square, then $I \circ A$ is the diagonal part of A .

The following result shows that $A \circ B$ is a submatrix of $A \otimes B$.

Proposition 7.3.1. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$A \circ B = (A \otimes B)_{(\{1, n+2, 2n+3, \dots, n^2\}, \{1, m+2, 2m+3, \dots, m^2\})}. \quad (7.3.7)$$

If, in addition, $n = m$, then

$$A \circ B = (A \otimes B)_{(\{1, n+2, 2n+3, \dots, n^2\})}, \quad (7.3.8)$$

and thus $A \circ B$ is a principal submatrix of $A \otimes B$.

Proof. See [711, p. 304] or [962]. □

7.4 Facts on the Kronecker Product

Fact 7.4.1. Let $x, y \in \mathbb{F}^n$. Then,

$$x \otimes y = (x \otimes I_n)y = (I_n \otimes y)x.$$

Fact 7.4.2. Let $x, y, w, z \in \mathbb{F}^n$. Then,

$$x^T w y^T z = (x^T \otimes y^T)(w \otimes z) = (x \otimes y)^T(w \otimes z).$$

Fact 7.4.3. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$, and assume that A and B are (diagonal, upper triangular, lower triangular). Then, so is $A \otimes B$.

Fact 7.4.4. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $l \in \mathbb{P}$. Then,

$$(A \otimes B)^l = A^l \otimes B^l.$$

Fact 7.4.5. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\text{vec } A = (I_m \otimes A) \text{vec } I_m = (A^T \otimes I_n) \text{vec } I_n.$$

Fact 7.4.6. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\text{vec } AB = (I_l \otimes A) \text{vec } B = (B^T \otimes A) \text{vec } I_m = \sum_{i=1}^m \text{col}_i(B^T) \otimes \text{col}_i(A).$$

Fact 7.4.7. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $C \in \mathbb{F}^{l \times n}$. Then,

$$\text{tr } ABC = (\text{vec } A)^T (B \otimes I) \text{vec } C^T.$$

Fact 7.4.8. Let $A, B, C \in \mathbb{F}^{n \times n}$, and assume that C is symmetric. Then,

$$(\text{vec } C)^T (A \otimes B) \text{vec } C = (\text{vec } C)^T (B \otimes A) \text{vec } C.$$

Fact 7.4.9. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, $C \in \mathbb{F}^{l \times k}$, and $D \in \mathbb{F}^{k \times n}$. Then,

$$\text{tr } ABCD = (\text{vec } A)^T (B \otimes D^T) \text{vec } C^T.$$

Fact 7.4.10. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $k \geq 1$. Then,

$$(AB)^{\otimes k} = A^{\otimes k} B^{\otimes k},$$

where $A^{\otimes k} \triangleq A \otimes A \otimes \cdots \otimes A$, with A appearing k times.

Fact 7.4.11. Let $A, C \in \mathbb{F}^{n \times m}$ and $B, D \in \mathbb{F}^{l \times k}$, assume that A is (left equivalent, right equivalent, biequivalent) to C , and assume that B is (left equivalent, right equivalent, biequivalent) to D . Then, $A \otimes B$ is (left equivalent, right equivalent, biequivalent) to $C \otimes D$.

Fact 7.4.12. Let $A, B, C, D \in \mathbb{F}^{n \times n}$, assume that A is (similar, congruent, unitarily similar) to C , and assume that B is (similar, congruent, unitarily similar) to D . Then, $A \otimes B$ is (similar, congruent, unitarily similar) to $C \otimes D$.

Fact 7.4.13. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$, and let $\gamma \in \text{spec}(A \otimes B)$. Then,

$$\begin{aligned} \sum \text{gmult}_A(\lambda) \text{gmult}_B(\mu) &\leq \text{gmult}_{A \otimes B}(\gamma) \\ &\leq \text{amult}_{A \otimes B}(\gamma) \\ &= \sum \text{amult}_A(\lambda) \text{amult}_B(\mu), \end{aligned}$$

where both sums are taken over all $\lambda \in \text{spec}(A)$ and $\mu \in \text{spec}(B)$ such that $\lambda\mu = \gamma$.

Fact 7.4.14. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{sprad}(A \otimes A) = [\text{sprad}(A)]^2.$$

Fact 7.4.15. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$, and let $\gamma \in \text{spec}(A \otimes B)$. Then, $\text{ind}_{A \otimes B}(\gamma) = 1$ if and only if $\text{ind}_A(\lambda) = 1$ and $\text{ind}_B(\mu) = 1$ for all $\lambda \in \text{spec}(A)$ and $\mu \in \text{spec}(B)$ such that $\lambda\mu = \gamma$.

Fact 7.4.16. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{n \times n}$, and assume that A and B are (group invertible, range Hermitian, range symmetric, Hermitian, symmetric, normal, positive semidefinite, positive definite, unitary, orthogonal, projectors, reflectors, involutory, idempotent, tripotent, nilpotent, semisimple). Then, so is $A \otimes B$. (Remark: See Fact 7.4.31.)

Fact 7.4.17. Let $A_1, \dots, A_l \in \mathbb{F}^{n \times n}$, and assume that $A_1, \dots, A_l$ are skew Hermitian. If l is (even, odd), then $A_1 \otimes \dots \otimes A_l$ is (Hermitian, skew Hermitian).

Fact 7.4.18. Let $A_{i,j} \in \mathbb{F}^{n_i \times n_j}$ for all $i = 1, \dots, k$ and $j = 1, \dots, l$. Then,

$$\begin{bmatrix} A_{11} & A_{22} & \cdots \\ A_{21} & A_{22} & \ddots \\ \vdots & \ddots & \ddots \end{bmatrix} \otimes B = \begin{bmatrix} A_{11} \otimes B & A_{22} \otimes B & \cdots \\ A_{21} \otimes B & A_{22} \otimes B & \ddots \\ \vdots & \ddots & \ddots \end{bmatrix}.$$

Fact 7.4.19. Let $x \in \mathbb{F}^k$, and let $A_i \in \mathbb{F}^{n \times n_i}$ for all $i = 1, \dots, l$. Then,

$$x \otimes \begin{bmatrix} A_1 & \cdots & A_l \end{bmatrix} = \begin{bmatrix} x \otimes A_1 & \cdots & x \otimes A_l \end{bmatrix}.$$

Fact 7.4.20. Let $x \in \mathbb{F}^m$, let $A \in \mathbb{F}^{n \times m}$, and let $B \in \mathbb{F}^{m \times l}$. Then,

$$(A \otimes x)B = (A \otimes x)(B \otimes 1) = (AB) \otimes x.$$

Fact 7.4.21. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then, the eigenvalues of $\sum_{i,j=1,l}^{k,l} \gamma_{ij} A^i \otimes B^j$ are of the form $\sum_{i,j=1,l}^{k,l} \gamma_{ij} \lambda^i \mu^j$, where $\lambda \in \text{spec}(A)$ and $\mu \in \text{spec}(B)$ and an associated eigenvector is given by $x \otimes y$, where $x \in \mathbb{F}^n$ is an eigenvector of A associated with $\lambda \in \text{spec}(A)$ and $y \in \mathbb{F}^m$ is an eigenvector of B associated with $\mu \in \text{spec}(B)$. (Remark: This result is due to Stephanos.) (Proof: Let $Ax = \lambda x$ and $By = \mu y$. Then, $\gamma_{ij} (A^i \otimes B^j)(x \otimes y) = \gamma_{ij} \lambda^i \mu^j (x \otimes y)$. See [519], [867, p. 411], or [942, p. 83].)

Fact 7.4.22. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times k}$. Then,

$$\mathcal{R}(A \otimes B) = \mathcal{R}(A \otimes I_{l \times l}) \cap \mathcal{R}(I_{n \times n} \otimes B).$$

(Proof: See [1293].)

Fact 7.4.23. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times k}$. Then,

$$\text{rank}(A \otimes B) = (\text{rank } A)(\text{rank } B) = \text{rank}(B \otimes A).$$

Consequently, $A \otimes B = 0$ if and only if either $A = 0$ or $B = 0$. (Proof: Use the singular value decomposition of $A \otimes B$.) (Remark: See Fact 8.21.16.)

Fact 7.4.24. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{l \times k}$, $C \in \mathbb{F}^{n \times p}$, $D \in \mathbb{F}^{l \times q}$. Then,

$$\begin{aligned} & \text{rank} \begin{bmatrix} A \otimes B & C \otimes D \end{bmatrix} \\ & \leq \begin{cases} (\text{rank } A)\text{rank} \begin{bmatrix} B & D \end{bmatrix} + (\text{rank } D)\text{rank} \begin{bmatrix} A & C \end{bmatrix} - (\text{rank } A)\text{rank } D \\ (\text{rank } B)\text{rank} \begin{bmatrix} A & C \end{bmatrix} + (\text{rank } C)\text{rank} \begin{bmatrix} B & D \end{bmatrix} - (\text{rank } B)\text{rank } C. \end{cases} \end{aligned}$$

(Proof: See [1297].)

Fact 7.4.25. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$\text{rank}(I - A \otimes B) \leq nm - [n - \text{rank}(I - A)][m - \text{rank}(I - B)].$$

(Proof: See [333].)

Fact 7.4.26. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$\text{ind } A \otimes B = \max\{\text{ind } A, \text{ind } B\}.$$

Fact 7.4.27. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times k}$, and assume that $nl = mk$ and $n \neq m$. Then, $A \otimes B$ is singular. (Proof: See [711, p. 250].)

Fact 7.4.28. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then,

$$|n - m| \min\{n, m\} \leq \text{amult}_{A \otimes B}(0).$$

(Proof: See [711, p. 249].)

Fact 7.4.29. The Kronecker permutation matrix $P_{n,m} \in \mathbb{R}^{nm \times nm}$ has the following properties:

- i) $P_{n,m}$ is a permutation matrix.
- ii) $P_{n,m}^T = P_{n,m}^{-1} = P_{m,n}$.
- iii) $P_{n,m}$ is orthogonal.
- iv) $P_{n,m}P_{m,n} = I_{nm}$.
- v) $P_{n,n}$ is orthogonal, symmetric, and involutory.
- vi) $P_{n,n}$ is a reflector.
- vii) $\text{sig } P_{n,n} = \text{tr } P_{n,n} = n$.

viii) The inertia of $P_{n,n}$ is given by

$$\text{In } P_{n,n} = \begin{bmatrix} \frac{1}{2}(n^2 - n) & & \\ & 0 & \\ & & \frac{1}{2}(n^2 + n) \end{bmatrix}.$$

ix) $P_{1,m} = I_m$ and $P_{n,1} = I_n$.

x) If $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$, then

$$P_{n,m}(y \otimes x) = x \otimes y.$$

xi) If $A \in \mathbb{F}^{n \times m}$ and $b \in \mathbb{F}^k$, then

$$P_{k,n}(A \otimes b) = b \otimes A$$

and

$$P_{n,k}(b \otimes A) = A \otimes b.$$

xii) If $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times k}$, then

$$P_{l,n}(A \otimes B)P_{m,k} = B \otimes A$$

and

$$\text{vec}(A \otimes B) = (I_m \otimes P_{k,n} \otimes I_l)[(\text{vec } A) \otimes (\text{vec } B)].$$

xiii) If $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{l \times l}$, then

$$P_{l,n}(A \otimes B)P_{n,l} = P_{l,n}(A \otimes B)P_{l,n}^{-1} = B \otimes A.$$

Hence, $A \otimes B$ and $B \otimes A$ are similar.

xiv) If $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$, then

$$\text{tr } AB = \text{tr}[P_{m,n}(A \otimes B)].$$

Fact 7.4.30. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times k}$. Then,

$$(A \otimes B)^+ = A^+ \otimes B^+.$$

Fact 7.4.31. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$(A \otimes B)^D = A^D \otimes B^D.$$

Now, assume that A and B are group invertible. Then, $A \otimes B$ is group invertible, and

$$(A \otimes B)^\# = A^\# \otimes B^\#.$$

(Remark: See Fact 7.4.16.)

Fact 7.4.32. For all $i = 1, \dots, p$, let $A_i \in \mathbb{F}^{n_i \times n_i}$. Then,

$$\begin{aligned} \text{mspec}(A_1 \otimes \cdots \otimes A_p) \\ = \{\lambda_1 \cdots \lambda_p : \lambda_i \in \text{mspec}(A_i) \text{ for all } i = 1, \dots, p\}_{\text{ms}}. \end{aligned}$$

If, in addition, for all $i = 1, \dots, p$, $x_i \in \mathbb{C}^{n_i}$ is an eigenvector of A_i associated with $\lambda_i \in \text{spec}(A_i)$, then $x_1 \otimes \cdots \otimes x_p$ is an eigenvector of $A_1 \otimes \cdots \otimes A_p$ associated with $\lambda_1 \cdots \lambda_p$.

7.5 Facts on the Kronecker Sum

Fact 7.5.1. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$(A \oplus A)^2 = A^2 \oplus A^2 + 2A \otimes A.$$

Fact 7.5.2. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$n \leq \dim(A^T \oplus -A) = \dim \{X \in \mathbb{F}^{n \times n} : AX = XA\}$$

and

$$\text{rank}(A^T \oplus -A) = \dim \{[A, X] : X \in \mathbb{F}^{n \times n}\} \leq n^2 - n.$$

(Proof: See Fact 2.18.9.) (Remark: $\text{rank}(A^T \oplus -A)$ is the dimension of the commutant or centralizer of A . See Fact 2.18.9.) (Problem: Express $\text{rank}(A^T \oplus -A)$ in terms of the eigenstructure of A .) (Remark: See Fact 5.14.22 and Fact 5.14.24.)

Fact 7.5.3. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nilpotent, and assume that $A^T \oplus -A = 0$. Then, $A = 0$. (Proof: Note that $A^T \otimes A^k = I \otimes A^{k+1}$, and use Fact 7.4.23.)

Fact 7.5.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that, for all $X \in \mathbb{F}^{n \times n}$, $AX = XA$. Then, there exists $\alpha \in \mathbb{F}$ such that $A = \alpha I$. (Proof: It follows from Proposition 7.2.3 that all of the eigenvalues of A are equal. Hence, there exists $\alpha \in \mathbb{F}$ such that $A = \alpha I + B$, where B is nilpotent. Now, Fact 7.5.3 implies that $B = 0$.)

Fact 7.5.5. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$, and let $\gamma \in \text{spec}(A \oplus B)$. Then,

$$\begin{aligned} \sum \text{gmult}_A(\lambda) \text{gmult}_B(\mu) &\leq \text{gmult}_{A \oplus B}(\gamma) \\ &\leq \text{amult}_{A \oplus B}(\gamma) \\ &= \sum \text{amult}_A(\lambda) \text{amult}_B(\mu), \end{aligned}$$

where both sums are taken over all $\lambda \in \text{spec}(A)$ and $\mu \in \text{spec}(B)$ such that $\lambda + \mu = \gamma$.

Fact 7.5.6. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{spabs}(A \oplus A) = 2 \text{spabs}(A).$$

Fact 7.5.7. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$, and let $\gamma \in \text{spec}(A \oplus B)$. Then, $\text{ind}_{A \oplus B}(\gamma) = 1$ if and only if $\text{ind}_A(\lambda) = 1$ and $\text{ind}_B(\mu) = 1$ for all $\lambda \in \text{spec}(A)$ and $\mu \in \text{spec}(B)$ such that $\lambda + \mu = \gamma$.

Fact 7.5.8. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$, and assume that A and B are (group invertible, range Hermitian, Hermitian, symmetric, skew Hermitian, skew symmetric, normal, positive semidefinite, positive definite, semidissipative, dissipative, nilpotent, semisimple). Then, so is $A \oplus B$.

Fact 7.5.9. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$P_{m,n}(A \oplus B)P_{n,m} = P_{m,n}(A \oplus B)P_{m,n}^{-1} = B \oplus A.$$

Hence, $A \oplus B$ and $B \oplus A$ are similar, and thus

$$\text{rank}(A \oplus B) = \text{rank}(B \oplus A).$$

(Proof: Use *xiii*) of Fact 7.4.29.)

Fact 7.5.10. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$\begin{aligned} n \text{rank } B + m \text{rank } A - 2(\text{rank } A)(\text{rank } B) \\ \leq \text{rank}(A \oplus B) \\ \leq \begin{cases} nm - [n - \text{rank}(I + A)][m - \text{rank}(I - B)] \\ nm - [n - \text{rank}(I - A)][m - \text{rank}(I + B)]. \end{cases} \end{aligned}$$

If, in addition, $-A$ and B are idempotent, then

$$\text{rank}(A \oplus B) = n \text{rank } B + m \text{rank } A - 2(\text{rank } A)(\text{rank } B).$$

Equivalently,

$$\text{rank}(A \oplus B) = (\text{rank } (-A)_\perp) \text{rank } B + (\text{rank } B_\perp) \text{rank } A.$$

(Proof: See [333].) (Remark: Equality may not hold for the upper bounds when $-A$ and B are idempotent.)

Fact 7.5.11. Let $A \in \mathbb{F}^{n \times n}$, let $B \in \mathbb{F}^{m \times m}$, assume that A is positive definite, and define $p(s) \triangleq \det(I - sA)$, and let $\text{mroots}(p) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\det(A \oplus B) = (\det A)^m \prod_{i=1}^n \det(\lambda_i B + I).$$

(Proof: Specialize Fact 7.5.12.)

Fact 7.5.12. Let $A, C \in \mathbb{F}^{n \times n}$, let $B, D \in \mathbb{F}^{m \times m}$, assume that A is positive definite, assume that C is positive semidefinite, define $p(s) \triangleq \det(C - sA)$, and let $\text{mroots}(p) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\det(A \otimes B + C \otimes D) = (\det A)^m \prod_{i=1}^n \det(\lambda_i D + B).$$

(Proof: See [1002, pp. 40, 41].) (Remark: The Kronecker product definition in [1002] follows the convention of [942], where “ $A \otimes B$ ” denotes $B \otimes A$.)

Fact 7.5.13. Let $A, D \in \mathbb{F}^{n \times n}$, let $C, B \in \mathbb{F}^{m \times m}$, assume that $\text{rank } C = 1$, and assume that A is nonsingular. Then,

$$\det(A \otimes B + C \otimes D) = (\det A)^m (\det B)^{n-1} \det[B + (\text{tr } CA^{-1})D].$$

(Proof: See [1002, p. 41].)

Fact 7.5.14. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then, $\text{spec}(A)$ and $\text{spec}(-B)$ are disjoint if and only if, for all $C \in \mathbb{F}^{n \times m}$, the matrices $\begin{bmatrix} A & 0 \\ 0 & -B \end{bmatrix}$ and $\begin{bmatrix} A & C \\ 0 & -B \end{bmatrix}$ are similar. (Proof: Sufficiency follows from Fact 5.10.21, while necessity follows from Corollary 2.6.6 and Proposition 7.2.3.)

Fact 7.5.15. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{n \times m}$, and assume that $\det(B^T \oplus A) \neq 0$. Then, $X \in \mathbb{F}^{n \times m}$ satisfies

$$A^2X + 2AXB + XB^2 + C = 0$$

if and only if

$$X = -\text{vec}^{-1}\left[(B^T \oplus A)^{-2} \text{vec } C\right].$$

Fact 7.5.16. For all $i = 1, \dots, p$, let $A_i \in \mathbb{F}^{n_i \times n_i}$. Then,

$$\begin{aligned} \text{mspec}(A_1 \oplus \dots \oplus A_p) \\ = \{\lambda_1 + \dots + \lambda_p : \lambda_i \in \text{mspec}(A_i) \text{ for all } i = 1, \dots, p\}_{\text{ms}}. \end{aligned}$$

If, in addition, for all $i = 1, \dots, p$, $x_i \in \mathbb{C}^{n_i}$ is an eigenvector of A_i associated with $\lambda_i \in \text{spec}(A_i)$, then $x_1 \oplus \dots \oplus x_p$ is an eigenvector of $A_1 \oplus \dots \oplus A_p$ associated with $\lambda_1 + \dots + \lambda_p$.

Fact 7.5.17. Let $A \in \mathbb{F}^{n \times m}$, and let $k \in \mathbb{P}$ satisfy $1 \leq k \leq \min\{n, m\}$. Furthermore, define the k th *compound* $A^{(k)}$ to be the $\binom{n}{k} \times \binom{m}{k}$ matrix whose entries are $k \times k$ subdeterminants of A , ordered lexicographically. (Example: For $n = k = 3$, subsets of the rows and columns of A are chosen in the order $\{1, 1, 1\}, \{1, 1, 2\}, \{1, 1, 3\}, \{1, 2, 1\}, \{1, 2, 2\}, \dots$) Specifically, $(A^{(k)})_{(i,j)}$ is the $k \times k$ subdeterminant of A corresponding to the i th selection of k rows of A and the j th selection of k columns of A . Then, the following statements hold:

- i) $A^{(1)} = A$.
- ii) $(\alpha A)^{(k)} = \alpha^k A^{(k)}$.
- iii) $(A^T)^{(k)} = (A^{(k)})^T$.
- iv) $\overline{A}^{(k)} = \overline{A^{(k)}}$.
- v) $(A^*)^{(k)} = (A^{(k)})^*$.
- vi) If $B \in \mathbb{F}^{m \times l}$ and $1 \leq k \leq \min\{n, m, l\}$, then $(AB)^{(k)} = A^{(k)}B^{(k)}$.
- vii) If $B \in \mathbb{F}^{m \times n}$, then $\det AB = A^{(k)}B^{(k)}$.

Now, assume that $m = n$, let $1 \leq k \leq n$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then, the following statements hold:

- viii) If A is (diagonal, lower triangular, upper triangular, Hermitian, positive semidefinite, positive definite, unitary), then so is $A^{(k)}$.
- ix) Assume that A is skew Hermitian. If k is odd, then $A^{(k)}$ is skew Hermitian. If k is even, then $A^{(k)}$ is Hermitian.
- x) Assume that A is diagonal, upper triangular, or lower triangular, and let $1 \leq i_1 < \dots < i_k \leq n$. Then, the $(i_1 + \dots + i_k, i_1 + \dots + i_k)$ entry of $A^{(k)}$ is $A_{(i_1, i_1)} \cdots A_{(i_k, i_k)}$. In particular, $I_n^{(k)} = I_{\binom{n}{k}}$.
- xi) $\det A^{(k)} = (\det A)^{\binom{n-1}{k-1}}$.
- xii) $A^{(n)} = \det A$.

xiii) $SA^{(n-1)T}S = A^A$, where $S \triangleq \text{diag}(1, -1, 1, \dots)$.

xiv) $\det A^{(n-1)} = \det A^A = (\det A)^{n-1}$.

xv) $\text{tr } A^{(n-1)} = \text{tr } A^A$.

xvi) If A is nonsingular, then $(A^{(k)})^{-1} = (A^{-1})^{(k)}$.

xvii) $\text{mspec}(A^{(k)}) = \{\lambda_{i_1} \cdots \lambda_{i_k} : 1 \leq i_1 < \cdots < i_k \leq n\}_{\text{ms}}$. In particular,

$$\text{mspec}(A^{(2)}) = \{\lambda_i \lambda_j : i, j = 1, \dots, n, i < j\}_{\text{ms}}.$$

xviii) $\text{tr } A^{(k)} = \sum_{1 \leq i_1 < \cdots < i_k \leq n} \lambda_{i_1} \cdots \lambda_{i_k}$.

xix) If A has exactly k nonzero eigenvalues, then $A^{(k)}$ has exactly one nonzero eigenvalue.

xx) If $k < n$ and A has exactly k nonzero eigenvalues, then $\text{spec}(A^{(k+1)}) = \{0\}$, and thus $A^{(k+1)}$ is nilpotent.

xxi) If $B \in \mathbb{F}^{n \times n}$, then $\det(A + B) = \begin{bmatrix} A & I \end{bmatrix}^{(n)} \begin{bmatrix} I \\ B \end{bmatrix}^{(n)}$.

xxii) The characteristic polynomial of A is given by

$$\chi_A(s) = s^n + \sum_{i=1}^{n-1} (-1)^{n+i} [\text{tr } A^{(n-i)}] s^i + (-1)^n \det A.$$

xxiii) $\det(I + A) = 1 + \det A + \sum_{i=1}^{n-1} \text{tr } A^{(n-i)}$.

Now, for $i = 0, \dots, k$, define $A^{(k,i)}$ by

$$(A + sI)^{(k)} = s^k A^{(k,0)} + s^{k-1} A^{(k,1)} + \cdots + s A^{(k,k-1)} + A^{(k,k)}.$$

Then, the following statements hold:

xxiv) $A^{(k,0)} = I$.

xxv) $A^{(k,k)} = A^{(k)}$.

xxvi) If $B \in \mathbb{F}^{n \times n}$ and $\alpha, \beta \in \mathbb{F}$, then

$$(\alpha A + \beta B)^{(k,1)} = \alpha A^{(k,1)} + \beta B^{(k,1)}.$$

xxvii) $\text{mspec}(A^{(k,1)}) = \{\lambda_{i_1} + \cdots + \lambda_{i_k} : 1 \leq i_1 < \cdots < i_k \leq n\}_{\text{ms}}$.

xxviii) $\text{tr } A^{(k,1)} = \binom{n-1}{k-1} \text{tr } A$.

xxix) $\text{mspec}(A^{(2,1)}) = \{\lambda_i + \lambda_j : i, j = 1, \dots, n, i < j\}_{\text{ms}}$.

xxx) $\text{mspec}\left[\left(A^{(2,1)}\right)^2 - 4A^{(2)}\right] = \{(\lambda_i - \lambda_j)^2 : i, j = 1, \dots, n, i < j\}_{\text{ms}}$.

(Proof: See [481, pp. 142–155], [709, p. 11], [958, pp. 116–130], [971, pp. 502–506], [1098, p. 124], and [1099].) (Remark: Statement *vi*) is the *Binet-Cauchy theorem*. See [971, p. 503]. The special case given by statement *vii*) is also given by Fact 2.13.4. Another special case is given by statement *xxi*). Statement *xi*) is the *Sylvester-Franke theorem*. See [958, p. 130].) (Remark: $A^{(k,1)}$ is the k th additive compound of A .) (Remark: $(A^{(2,1)})^2 - 4A^{(2)}$ is the *discriminant* of A ,

which is singular if and only if A has a repeated eigenvalue.) (Remark: Additional expressions for the determinant of a sum of matrices are given in [1099].) (Remark: The compound operation is related to the *bialternate product* since $\text{mspec}(2A \cdot I) = \text{mspec}(A^{(2,1)})$ and $\text{mspec}(A \cdot A) = \text{mspec}(A^{(2)})$. See [519, 576], [782, pp. 313–320], and [942, pp. 84, 85].) (Remark: Induced norms of compound matrices are considered in [451].) (Remark: See Fact 11.17.12.) (Remark: Fact 4.9.2 and Fact 8.13.42.) (Problem: Express $A \cdot B$ in terms of compounds.)

7.6 Facts on the Schur Product

Fact 7.6.1. Let $x, y, z \in \mathbb{F}^n$. Then,

$$x^T(y \circ z) = z^T(x \circ y) = y^T(x \circ z).$$

Fact 7.6.2. Let $w, y \in \mathbb{F}^n$ and $x, z \in \mathbb{F}^m$. Then,

$$(wx^T) \circ (yz^T) = (w \circ y)(x \circ z)^T.$$

Fact 7.6.3. Let $A \in \mathbb{F}^{n \times n}$ and $d \in \mathbb{F}^n$. Then,

$$\text{diag}(d)A = A \circ d1_{1 \times n}.$$

Fact 7.6.4. Let $A, B \in \mathbb{F}^{n \times m}$, $D_1 \in \mathbb{F}^{n \times n}$, and $D_2 \in \mathbb{F}^{m \times m}$, and assume that D_1 and D_2 are diagonal. Then,

$$(D_1A) \circ (BD_2) = D_1(A \circ B)D_2.$$

Fact 7.6.5. Let $A_1, \dots, A_k \in \mathbb{F}^{n \times n}$. Then,

$$\mathcal{R}[(A_1A_1^*) \circ \dots \circ (A_kA_k^*)] = \text{span} \{(A_1x_1) \circ \dots \circ (A_kx_k) : x_1, \dots, x_k \in \mathbb{F}^n\}.$$

Furthermore, if $A_1, \dots, A_k$ are positive semidefinite, then

$$\begin{aligned} \mathcal{R}(A_1 \circ \dots \circ A_k) &= \text{span} \{(A_1x_1) \circ \dots \circ (A_kx_k) : x_1, \dots, x_k \in \mathbb{F}^n\} \\ &= \text{span} \{(A_1x) \circ \dots \circ (A_kx) : x \in \mathbb{F}^n\}. \end{aligned}$$

(Proof: See [1109].)

Fact 7.6.6. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\text{rank}(A \circ B) \leq \text{rank}(A \otimes B) = (\text{rank } A)(\text{rank } B).$$

(Proof: Use Proposition 7.3.1.) (Remark: See Fact 8.21.16.)

Fact 7.6.7. Let $x, a \in \mathbb{F}^n$, $y, b \in \mathbb{F}^m$, and $A \in \mathbb{F}^{n \times m}$. Then,

$$x^T(A \circ ab^T)y = (a \circ x)^T A(b \circ y).$$

Fact 7.6.8. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\text{tr}[(A \circ B)(A \circ B)^T] = \text{tr}[(A \circ A)(B \circ B)^T].$$

Fact 7.6.9. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times n}$, $a \in \mathbb{F}^m$, and $b \in \mathbb{F}^n$. Then,

$$\operatorname{tr}[A(B \circ ab^T)] = a^T(A^T \circ B)b.$$

In particular,

$$\operatorname{tr} AB = 1_m^T(A^T \circ B)1_n.$$

Fact 7.6.10. Let $A, B \in \mathbb{F}^{n \times m}$ and $C \in \mathbb{F}^{m \times n}$. Then,

$$I \circ [A(B^T \circ C)] = I \circ [(A \circ B)C] = I \circ [(A \circ C^T)B^T].$$

Hence,

$$\operatorname{tr}[A(B^T \circ C)] = \operatorname{tr}[(A \circ B)C] = \operatorname{tr}[(A \circ C^T)B^T].$$

Fact 7.6.11. Let $x \in \mathbb{R}^m$ and $A \in \mathbb{R}^{n \times m}$, and define $x^A \in \mathbb{R}^n$ by

$$x^A \triangleq \begin{bmatrix} \prod_{i=1}^m x_{(i)}^{A_{(1,i)}} \\ \vdots \\ \prod_{i=1}^m x_{(i)}^{A_{(n,i)}} \end{bmatrix},$$

where every component of x^A is assumed to exist. Then, the following statements hold:

- i) If $a \in \mathbb{R}$, then $a^x = \begin{bmatrix} a^{x_{(1)}} \\ \vdots \\ a^{x_{(m)}} \end{bmatrix}$.
- ii) $x^{-A} = (x^A)^{\circ -1}$.
- iii) If $y \in \mathbb{R}^m$, then $(x \circ y)^A = x^A \circ y^A$.
- iv) If $B \in \mathbb{R}^{n \times m}$, then $x^{A+B} = x^A \circ x^B$.
- v) If $B \in \mathbb{R}^{l \times n}$, then $(x^A)^B = x^{BA}$.
- vi) If $a \in \mathbb{R}$, then $(a^x)^A = a^{Ax}$.
- vii) If $A^L \in \mathbb{R}^{m \times n}$ is a left inverse of A and $y = x^A$, then $x = y^{A^L}$.
- viii) If $A \in \mathbb{R}^{n \times n}$ is nonsingular and $y = x^A$, then $x = y^{A^{-1}}$.
- ix) Define $f(x) \triangleq x^A$. Then, $f'(x) = \operatorname{diag}(x^A)A\operatorname{diag}(x^{\circ -1})$.
- x) Let $x_1, \dots, x_n \in \mathbb{R}^n$, let $a \in \mathbb{R}^n$, and assume that $0 < x_1 < \dots < x_n$ and $a_{(1)} < \dots < a_{(n)}$. Then,

$$\det \begin{bmatrix} x_1^a & \dots & x_n^a \end{bmatrix} > 0.$$

(Remark: These operations arise in modeling chemical reaction kinetics. See [892].)
(Proof: Result x) is given in [1130].)

Fact 7.6.12. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is nonsingular. Then,

$$(A \circ A^{-T})1_{n \times 1} = 1_{n \times 1}$$

and

$$1_{1 \times n}(A \circ A^{-T}) = 1_{1 \times n}.$$

(Proof: See [772].)

Fact 7.6.13. Let $A \in \mathbb{R}^{n \times n}$, and assume that $A \geq 0$. Then,

$$\text{sprad}\left[(A \circ A^T)^{\circ 1/2}\right] \leq \text{sprad}(A) \leq \text{sprad}\left[\frac{1}{2}(A + A^T)\right].$$

(Proof: See [1180].)

Fact 7.6.14. Let $A_1, \dots, A_r \in \mathbb{R}^{n \times n}$ and $\alpha_1, \dots, \alpha_r \in \mathbb{R}$, and assume that $A_i \geq 0$ for all $i = 1, \dots, r$, $\alpha_i > 0$ for all $i = 1, \dots, r$, and $\sum_{i=1}^r \alpha_i \geq 1$. Then,

$$\text{sprad}(A_1^{\circ \alpha_1} \circ \dots \circ A_r^{\circ \alpha_r}) \leq \prod_{i=1}^r [\text{sprad}(A_i)]^{\alpha_i}.$$

In particular, let $A \in \mathbb{R}^{n \times n}$, and assume that $A \geq 0$. Then, for all $\alpha \geq 1$,

$$\text{sprad}(A^{\circ \alpha}) \leq [\text{sprad}(A)]^\alpha,$$

whereas, for all $\alpha \leq 1$,

$$[\text{sprad}(A)]^\alpha \leq \text{sprad}(A^{\circ \alpha}).$$

Furthermore,

$$\text{sprad}\left(A^{\circ 1/2} \circ A^{T \circ 1/2}\right) \leq \text{sprad}(A)$$

and

$$[\text{sprad}(A \circ A)]^{1/2} \leq \text{sprad}(A) = [\text{sprad}(A \otimes A)]^{1/2}.$$

If, in addition, $B \in \mathbb{R}^{n \times n}$ is such that $B \geq 0$, then

$$\text{sprad}(A \circ B) \leq [\text{sprad}(A \circ A) \text{sprad}(B \circ B)]^{1/2} \leq \text{sprad}(A) \text{sprad}(B),$$

$$\begin{aligned} \text{sprad}(A \circ B) &\leq \text{sprad}(A) \text{sprad}(B) \\ &\quad + \max_{i=1, \dots, n} [2A_{(i,i)}B_{(i,i)} - \text{sprad}(A)B_{(i,i)} - \text{sprad}(B)A_{(i,i)}] \\ &\leq \text{sprad}(A) \text{sprad}(B), \end{aligned}$$

and

$$\text{sprad}\left(A^{\circ 1/2} \circ B^{\circ 1/2}\right) \leq \sqrt{\text{sprad}(A) \text{sprad}(B)}.$$

If, in addition, $A \gg 0$ and $B \gg 0$, then

$$\text{sprad}(A \circ B) < \text{sprad}(A) \text{sprad}(B).$$

(Proof: See [453, 467, 792]. The identity $\text{sprad}(A) = [\text{sprad}(A \otimes A)]^{1/2}$ follows from Fact 7.4.14.) (Remark: The inequality $\text{sprad}(A \circ A) \leq \text{sprad}(A \otimes A)$ follows from Fact 4.11.18 and Proposition 7.3.1.) (Remark: Some extensions are given in [731].)

Fact 7.6.15. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that A and B are nonsingular M-matrices. Then, the following statements hold:

- i) $A \circ B^{-1}$ is a nonsingular M-matrix.
- ii) If $n = 2$, then $\tau(A \circ A^{-1}) = 1$.
- iii) If $n \geq 3$, then $\frac{1}{n} < \tau(A \circ A^{-1}) \leq 1$.
- iv) $\tau(A) \min_{i=1, \dots, n} (B^{-1})_{(i,i)} \leq \tau(A \circ B^{-1})$.

$$v) \quad [\tau(A)\tau(B)]^n \leq |\det(A \circ B)|.$$

$$vi) \quad |(A \circ B)^{-1}| \leq A^{-1} \circ B^{-1}.$$

(Proof: See [711, pp. 359, 370, 375, 380].) (Remark: The minimum eigenvalue $\tau(A)$ is defined in Fact 4.11.9.) (Remark: Some extensions are given in [731].)

Fact 7.6.16. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\text{sprad}(A \circ B) \leq \sqrt{\text{sprad}(A \circ \overline{A}) \text{sprad}(B \circ \overline{B})}.$$

Consequently,

$$\left. \begin{array}{l} \text{sprad}(A \circ A) \\ \text{sprad}(A \circ A^T) \\ \text{sprad}(A \circ A^*) \end{array} \right\} \leq \text{sprad}(A \circ \overline{A}).$$

(Proof: See [1193].) (Remark: See Fact 9.14.34.)

Fact 7.6.17. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A and B are nonnegative, and let $\alpha \in [0, 1]$. Then,

$$\text{sprad}(A^{\circ\alpha} \circ B^{\circ(1-\alpha)}) \leq \text{sprad}^\alpha(A) \text{sprad}^{1-\alpha}(B).$$

In particular,

$$\text{sprad}(A^{\circ 1/2} \circ B^{\circ 1/2}) \leq \sqrt{\text{sprad}(A) \text{sprad}(B)}.$$

Finally,

$$\text{sprad}(A^{\circ 1/2} \circ A^{\circ 1/2T}) \leq \text{sprad}(A^{\circ\alpha} \circ A^{\circ(1-\alpha)T}) \leq \text{sprad}(A).$$

(Proof: See [1193].) (Remark: See Fact 9.14.35.)

7.7 Notes

A history of the Kronecker product is given in [665]. Kronecker matrix algebra is discussed in [259, 579, 667, 948, 994, 1219, 1379]. Applications are discussed in [1121, 1122, 1362].

The fact that the Schur product is a principal submatrix of the Kronecker product is noted in [962]. A variation of Kronecker matrix algebra for symmetric matrices can be developed in terms of the half-vectorization operator “vech” and the associated elimination and duplication matrices [667, 947, 1344].

Generalizations of the Schur and Kronecker products, known as the block-Kronecker, strong Kronecker, Khatri-Rao, and Tracy-Singh products, are discussed in [385, 714, 739, 840, 923, 925, 926, 928] and [1119, pp. 216, 217]. A related operation is the *bialternate product*, which is a variation of the compound operation discussed in Fact 7.5.17. See [519, 576], [782, pp. 313–320], and [942, pp. 84, 85]. The Schur product is also called the Hadamard product.

The Kronecker product is associated with tensor analysis and multilinear algebra [421, 545, 585, 958, 959, 994].

Chapter Eight

Positive-Semidefinite Matrices

In this chapter we focus on positive-semidefinite and positive-definite matrices. These matrices arise in a variety of applications, such as covariance analysis in signal processing and controllability analysis in linear system theory, and they have many special properties.

8.1 Positive-Semidefinite and Positive-Definite Orderings

Let $A \in \mathbb{F}^{n \times n}$ be a Hermitian matrix. As shown in Corollary 5.4.5, A is unitarily similar to a real diagonal matrix whose diagonal entries are the eigenvalues of A . We denote these eigenvalues by $\lambda_1, \dots, \lambda_n$ or, for clarity, by $\lambda_1(A), \dots, \lambda_n(A)$. As in Chapter 4, we employ the convention

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n, \quad (8.1.1)$$

and, for convenience, we define

$$\lambda_{\max}(A) \triangleq \lambda_1, \quad \lambda_{\min}(A) \triangleq \lambda_n. \quad (8.1.2)$$

Then, A is positive semidefinite if and only if $\lambda_{\min}(A) \geq 0$, while A is positive definite if and only if $\lambda_{\min}(A) > 0$.

For convenience, let $\mathbf{H}^n$, $\mathbf{N}^n$, and $\mathbf{P}^n$ denote, respectively, the Hermitian, positive-semidefinite, and positive-definite matrices in $\mathbb{F}^{n \times n}$. Hence, $\mathbf{P}^n \subset \mathbf{N}^n \subset \mathbf{H}^n$. If $A \in \mathbf{N}^n$, then we write $A \geq 0$, while, if $A \in \mathbf{P}^n$, then we write $A > 0$. If $A, B \in \mathbf{H}^n$, then $A - B \in \mathbf{N}^n$ is possible even if neither A nor B is positive semidefinite. In this case, we write $A \geq B$ or $B \leq A$. Similarly, $A - B \in \mathbf{P}^n$ is denoted by $A > B$ or $B < A$. This notation is consistent with the case $n = 1$, where $\mathbf{H}^1 = \mathbb{R}$, $\mathbf{N}^1 = [0, \infty)$, and $\mathbf{P}^1 = (0, \infty)$.

Since $0 \in \mathbf{N}^n$, it follows that $\mathbf{N}^n$ is a pointed cone. Furthermore, if $A, -A \in \mathbf{N}^n$, then $x^*Ax = 0$ for all $x \in \mathbb{F}^n$, which implies that $A = 0$. Hence, $\mathbf{N}^n$ is a one-sided cone. Finally, $\mathbf{N}^n$ and $\mathbf{P}^n$ are convex cones since, if $A, B \in \mathbf{N}^n$, then $\alpha A + \beta B \in \mathbf{N}^n$ for all $\alpha, \beta > 0$, and likewise for $\mathbf{P}^n$. The following result shows that the relation “ $\leq$ ” is a partial ordering on $\mathbf{H}^n$.

Proposition 8.1.1. The relation “ $\leq$ ” is reflexive, antisymmetric, and transitive on $\mathbf{H}^n$, that is, if $A, B, C \in \mathbf{H}^n$, then the following statements hold:

- i) $A \leq A$.
- ii) If $A \leq B$ and $B \leq A$, then $A = B$.
- iii) If $A \leq B$ and $B \leq C$, then $A \leq C$.

Proof. Since $\mathbf{N}^n$ is a pointed, one-sided, convex cone, it follows from Proposition 2.3.6 that the relation “ $\leq$ ” is reflexive, antisymmetric, and transitive. $\square$

Additional properties of “ $\leq$ ” and “ $<$ ” are given by the following result.

Proposition 8.1.2. Let $A, B, C, D \in \mathbf{H}^n$. Then, the following statements hold:

- i) If $A \geq 0$, then $\alpha A \geq 0$ for all $\alpha \geq 0$, and $\alpha A \leq 0$ for all $\alpha \leq 0$.
- ii) If $A > 0$, then $\alpha A > 0$ for all $\alpha > 0$, and $\alpha A < 0$ for all $\alpha < 0$.
- iii) $\alpha A + \beta B \in \mathbf{H}^n$ for all $\alpha, \beta \in \mathbb{R}$.
- iv) If $A \geq 0$ and $B \geq 0$, then $\alpha A + \beta B \geq 0$ for all $\alpha, \beta \geq 0$.
- v) If $A \geq 0$ and $B > 0$, then $A + B > 0$.
- vi) $A^2 \geq 0$.
- vii) $A^2 > 0$ if and only if $\det A \neq 0$.
- viii) If $A \leq B$ and $B < C$, then $A < C$.
- ix) If $A < B$ and $B \leq C$, then $A < C$.
- x) If $A \leq B$ and $C \leq D$, then $A + C \leq B + D$.
- xi) If $A \leq B$ and $C < D$, then $A + C < B + D$.

Furthermore, let $S \in \mathbb{F}^{m \times n}$. Then, the following statements hold:

- xii) If $A \leq B$, then $SAS^* \leq SBS^*$.
- xiii) If $A < B$ and $\text{rank } S = m$, then $SAS^* < SBS^*$.
- xiv) If $SAS^* \leq SBS^*$ and $\text{rank } S = n$, then $A \leq B$.
- xv) If $SAS^* < SBS^*$ and $\text{rank } S = n$, then $m = n$ and $A < B$.
- xvi) If $A \leq B$, then $SAS^* < SBS^*$ if and only if $\text{rank } S = m$ and $\mathcal{R}(S) \cap \mathcal{N}(B - A) = \{0\}$.

Proof. Results i)–xi) are immediate. To prove xii), note that $A < B$ implies that $(B - A)^{1/2}$ is positive definite. Thus, $\text{rank } S(A - B)^{1/2} = m$, which implies that $S(A - B)S^*$ is positive definite. To prove xiii), note that, since $\text{rank } S = n$, it follows that S has a left inverse $S^L \in \mathbb{F}^{n \times m}$. Thus, xi) implies that $A = S^L SAS^* S^{L*} \leq S^L SBS^* S^{L*} = B$. To prove xv), note that, since $S(B - A)S^*$ is positive definite, it follows that $\text{rank } S = m$. Hence, $m = n$ and S is nonsingular. Thus, xii) implies that $A = S^{-1} SAS^* S^{-*} < S^{-1} SBS^* S^{-*} = B$. Statement xvi) is proved in [285]. $\square$

The following result is an immediate consequence of Corollary 5.4.7.

Corollary 8.1.3. Let $A, B \in \mathbf{H}^n$, and assume that A and B are congruent. Then, A is positive semidefinite if and only if B is positive semidefinite. Furthermore, A is positive definite if and only if B is positive definite.

8.2 Submatrices

We first consider some identities involving a partitioned positive-semidefinite matrix.

Lemma 8.2.1. Let $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbf{N}^{n+m}$. Then,

$$A_{12} = A_{11}A_{11}^+A_{12}, \quad (8.2.1)$$

$$A_{12} = A_{12}A_{22}A_{22}^+. \quad (8.2.2)$$

Proof. Since $A \geq 0$, it follows from Corollary 5.4.5 that $A = BB^*$, where $B = \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} \in \mathbb{F}^{(n+m) \times r}$ and $r \triangleq \text{rank } A$. Thus, $A_{11} = B_1B_1^*$, $A_{12} = B_1B_2^*$, and $A_{22} = B_2B_2^*$. Since A_{11} is Hermitian, it follows from *xxvii*) of Proposition 6.1.6 that A_{11}^+ is also Hermitian. Next, defining $S \triangleq B_1 - B_1B_1^*(B_1B_1^*)^+B_1$, it follows that $SS^* = 0$, and thus $\text{tr } SS^* = 0$. Hence, Lemma 2.2.3 implies that $S = 0$, and thus $B_1 = B_1B_1^*(B_1B_1^*)^+B_1$. Consequently, $B_1B_2^* = B_1B_1^*(B_1B_1^*)^+B_1B_2^*$, that is, $A_{12} = A_{11}A_{11}^+A_{12}$. The second result is analogous. $\square$

Corollary 8.2.2. Let $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbf{N}^{n+m}$. Then, the following statements hold:

- i) $\mathcal{R}(A_{12}) \subseteq \mathcal{R}(A_{11})$.
- ii) $\mathcal{R}(A_{12}^*) \subseteq \mathcal{R}(A_{22})$.
- iii) $\text{rank} \begin{bmatrix} A_{11} & A_{12} \end{bmatrix} = \text{rank } A_{11}$.
- iv) $\text{rank} \begin{bmatrix} A_{12}^* & A_{22} \end{bmatrix} = \text{rank } A_{22}$.

Proof. Results *i*) and *ii*) follow from (8.2.1) and (8.2.2), while *iii*) and *iv*) are consequences of *i*) and *ii*). $\square$

Next, if (8.2.1) holds, then the partitioned Hermitian matrix $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$ can be factored as

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} = \begin{bmatrix} I & 0 \\ A_{12}^*A_{11}^+ & I \end{bmatrix} \begin{bmatrix} A_{11} & 0 \\ 0 & A_{11}|A \end{bmatrix} \begin{bmatrix} I & A_{11}^+A_{12} \\ 0 & I \end{bmatrix}, \quad (8.2.3)$$

while, if (8.2.2) holds, then

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} = \begin{bmatrix} I & A_{12}A_{22}^+ \\ 0 & I \end{bmatrix} \begin{bmatrix} A_{22}|A & 0 \\ 0 & A_{22} \end{bmatrix} \begin{bmatrix} I & 0 \\ A_{22}^+A_{12}^* & I \end{bmatrix}, \quad (8.2.4)$$

where

$$A_{11}|A = A_{22} - A_{12}^*A_{11}^+A_{12} \quad (8.2.5)$$

and

$$A_{22}|A = A_{11} - A_{12}A_{22}^+A_{12}^*. \quad (8.2.6)$$

Hence, it follows from Lemma 8.2.1 that, if A is positive semidefinite, then (8.2.3) and (8.2.4) are valid, and, furthermore, the Schur complements (see Definition 6.1.8) $A_{11}|A$ and $A_{22}|A$ are both positive semidefinite. Consequently, we have the following results.

Proposition 8.2.3. Let $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbf{N}^{n+m}$. Then,

$$\text{rank } A = \text{rank } A_{11} + \text{rank } A_{11}|A \quad (8.2.7)$$

$$= \text{rank } A_{22}|A + \text{rank } A_{22} \quad (8.2.8)$$

$$\leq \text{rank } A_{11} + \text{rank } A_{22}. \quad (8.2.9)$$

Furthermore,

$$\det A = (\det A_{11}) \det(A_{11}|A) \quad (8.2.10)$$

and

$$\det A = (\det A_{22}) \det(A_{22}|A). \quad (8.2.11)$$

Proposition 8.2.4. Let $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbf{H}^{n+m}$. Then, the following statements are equivalent:

- i) $A \geq 0$.
- ii) $A_{11} \geq 0$, $A_{12} = A_{11}A_{11}^+A_{12}$, and $A_{12}^*A_{11}^+A_{12} \leq A_{22}$.
- iii) $A_{22} \geq 0$, $A_{12} = A_{12}A_{22}^+A_{12}$, and $A_{12}A_{22}^+A_{12}^* \leq A_{11}$.

The following statements are also equivalent:

- iv) $A > 0$.
- v) $A_{11} > 0$ and $A_{12}^*A_{11}^{-1}A_{12} < A_{22}$.
- vi) $A_{22} > 0$ and $A_{12}A_{22}^{-1}A_{12}^* < A_{11}$.

The following result follows from (2.8.16) and (2.8.17) or from (8.2.3) and (8.2.4).

Proposition 8.2.5. Let $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbf{P}^{n+m}$. Then,

$$A^{-1} = \begin{bmatrix} A_{11}^{-1} + A_{11}^{-1}A_{12}(A_{11}|A)^{-1}A_{12}^*A_{11}^{-1} & -A_{11}^{-1}A_{12}(A_{11}|A)^{-1} \\ -(A_{11}|A)^{-1}A_{12}^*A_{11}^{-1} & (A_{11}|A)^{-1} \end{bmatrix} \quad (8.2.12)$$

and

$$A^{-1} = \begin{bmatrix} (A_{22}|A)^{-1} & -(A_{22}|A)^{-1}A_{12}A_{22}^{-1} \\ -A_{22}^{-1}A_{12}^*(A_{22}|A)^{-1} & A_{22}^{-1}A_{12}^*(A_{22}|A)^{-1}A_{12}A_{22}^{-1} + A_{22}^{-1} \end{bmatrix}, \quad (8.2.13)$$

where

$$A_{11}|A = A_{22} - A_{12}^*A_{11}^{-1}A_{12} \quad (8.2.14)$$

and

$$A_{22}|A = A_{11} - A_{12}A_{22}^{-1}A_{12}^*. \quad (8.2.15)$$

Now, let $A^{-1} = \begin{bmatrix} B_{11} & B_{12} \\ B_{12}^* & B_{22} \end{bmatrix}$. Then,

$$B_{11}|A^{-1} = A_{22}^{-1} \quad (8.2.16)$$

and

$$B_{22}|A^{-1} = A_{11}^{-1}. \quad (8.2.17)$$

Lemma 8.2.6. Let $A \in \mathbb{F}^{n \times n}$, $b \in \mathbb{F}^n$, and $a \in \mathbb{R}$, and define $\mathcal{A} \triangleq \begin{bmatrix} A & b \\ b^* & a \end{bmatrix}$. Then, the following statements are equivalent:

- i) $\mathcal{A}$ is positive semidefinite.
- ii) A is positive semidefinite, $b = AA^+b$, and $b^*A^+b \leq a$.
- iii) Either A is positive semidefinite, $a = 0$, and $b = 0$, or $a > 0$ and $bb^* \leq aA$.

Furthermore, the following statements are equivalent:

- i) $\mathcal{A}$ is positive definite.
- ii) A is positive definite, and $b^*A^{-1}b < a$.
- iii) $a > 0$ and $bb^* < aA$.

In this case,

$$\det \mathcal{A} = (\det A)(a - b^*A^{-1}b). \quad (8.2.18)$$

For the following result note that a matrix is a principal submatrix of itself, while the determinant of a matrix is also a principal subdeterminant of the matrix.

Proposition 8.2.7. Let $A \in \mathbf{H}^n$. Then, the following statements are equivalent:

- i) A is positive semidefinite.
- ii) Every principal submatrix of A is positive semidefinite.
- iii) Every principal subdeterminant of A is nonnegative.
- iv) For all $i = 1, \dots, n$, the sum of all $i \times i$ principal subdeterminants of A is nonnegative.
- v) $\beta_0, \dots, \beta_{n-1} \geq 0$, where $\chi_A(s) = s^n + \beta_{n-1}s^{n-1} + \dots + \beta_1s + \beta_0$.

Proof. To prove i) $\implies$ ii), let $\hat{A} \in \mathbb{F}^{m \times m}$ be the principal submatrix of A obtained from A by retaining rows and columns $i_1, \dots, i_m$. Then, $\hat{A} = S^TAS$, where $S \triangleq \begin{bmatrix} e_{i_1} & \dots & e_{i_m} \end{bmatrix} \in \mathbb{R}^{n \times m}$. Now, let $\hat{x} \in \mathbb{F}^m$. Since A is positive semidefinite, it follows that $\hat{x}^*\hat{A}\hat{x} = \hat{x}^*S^TAS\hat{x} \geq 0$, and thus $\hat{A}$ is positive semidefinite.

Next, the implications ii) $\implies$ iii) $\implies$ iv) are immediate. To prove iv) $\implies$ i), note that it follows from Proposition 4.4.6 that

$$\chi_A(s) = \sum_{i=0}^n \beta_i s^i = \sum_{i=0}^n (-1)^{n-i} \gamma_{n-i} s^i = (-1)^n \sum_{i=0}^n \gamma_{n-i} (-s)^i, \quad (8.2.19)$$

where, for all $i = 1, \dots, n$, γ_i is the sum of all $i \times i$ principal subdeterminants of A , and $\beta_n = \gamma_0 = 1$. By assumption, $\gamma_i \geq 0$ for all $i = 1, \dots, n$. Now, suppose there

exists $\lambda \in \text{spec}(A)$ such that $\lambda < 0$. Then, $0 = (-1)^n \chi_A(\lambda) = \sum_{i=0}^n \gamma_{n-i}(-\lambda)^i > 0$, which is a contradiction. The equivalence of *iv*) and *v*) follows from Proposition 4.4.6. $\square$

Proposition 8.2.8. Let $A \in \mathbf{H}^n$. Then, the following statements are equivalent:

- i*) A is positive definite.
- ii*) Every principal submatrix of A is positive definite.
- iii*) Every principal subdeterminant of A is positive.
- iv*) Every leading principal submatrix of A is positive definite.
- v*) Every leading principal subdeterminant of A is positive.

Proof. To prove *i*) $\implies$ *ii*), let $\hat{A} \in \mathbb{F}^{m \times m}$ and S be as in the proof of Proposition 8.2.7, and let $\hat{x}$ be nonzero so that $S\hat{x}$ is nonzero. Since A is positive definite, it follows that $\hat{x}^* \hat{A} \hat{x} = \hat{x}^* S^T A S \hat{x} > 0$, and hence $\hat{A}$ is positive definite.

Next, the implications *i*) $\implies$ *ii*) $\implies$ *iii*) $\implies$ *v*) and *ii*) $\implies$ *iv*) $\implies$ *v*) are immediate. To prove *v*) $\implies$ *i*), suppose that the leading principal submatrix $A_i \in \mathbb{F}^{i \times i}$ has positive determinant for all $i = 1, \dots, n$. The result is true for $n = 1$. For $n \geq 2$, we show that, if A_i is positive definite, then so is A_{i+1} . Writing $A_{i+1} = \begin{bmatrix} A_i & b_i \\ b_i^* & a_i \end{bmatrix}$, it follows from Lemma 8.2.6 that $\det A_{i+1} = (\det A_i)(a_i - b_i^* A_i^{-1} b_i) > 0$, and hence $a_i - b_i^* A_i^{-1} b_i = \det A_{i+1} / \det A_i > 0$. Lemma 8.2.6 now implies that A_{i+1} is positive definite. Using this argument for all $i = 2, \dots, n$ implies that A is positive definite. $\square$

The example $A = \begin{bmatrix} 0 & 0 \\ 0 & -1 \end{bmatrix}$ shows that every principal subdeterminant of A , rather than just the leading principal subdeterminants of A , must be checked to determine whether A is positive semidefinite. A less obvious example is $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 0 \end{bmatrix}$, whose eigenvalues are $0, 1 + \sqrt{3}$, and $1 - \sqrt{3}$. In this case, the principal subdeterminant $\det A_{[1;1]} = \det \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix} < 0$.

Note that condition *iii*) of Proposition 8.2.8 includes $\det A > 0$ since the determinant of A is also a subdeterminant of A . The matrix $A = \begin{bmatrix} 3/2 & -1 & 1 \\ -1 & 2 & 1 \\ 1 & 1 & 2 \end{bmatrix}$ has the property that every 1×1 and 2×2 subdeterminant is positive but is not positive definite. This example shows that the result *iii*) $\implies$ *ii*) of Proposition 8.2.8 is false if the requirement that the determinant of A be positive is omitted.

8.3 Simultaneous Diagonalization

This section considers the simultaneous diagonalization of a pair of matrices $A, B \in \mathbf{H}^n$. There are two types of simultaneous diagonalization. *Cogredient diagonalization* involves a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and SBS^* are both diagonal, whereas *contragredient diagonalization* involves finding a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and $S^{-*}BS^{-1}$ are both diagonal. Both types

of simultaneous transformation involve only congruence transformations. We begin by assuming that one of the matrices is positive definite, in which case the results are quite simple to prove. Our first result involves cogredient diagonalization.

Theorem 8.3.1. Let $A, B \in \mathbf{H}^n$, and assume that A is positive definite. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $SAS^* = I$ and SBS^* is diagonal.

Proof. Setting $S_1 = A^{-1/2}$, it follows that $S_1AS_1^* = I$. Now, since $S_1BS_1^*$ is Hermitian, it follows from Corollary 5.4.5 that there exists a unitary matrix $S_2 \in \mathbb{F}^{n \times n}$ such that $SBS^* = S_2S_1BS_1^*S_2^*$ is diagonal, where $S = S_2S_1$. Finally, $SAS^* = S_2S_1AS_1^*S_2^* = S_2IS_2^* = I$. $\square$

An analogous result holds for contragredient diagonalization.

Theorem 8.3.2. Let $A, B \in \mathbf{H}^n$, and assume that A is positive definite. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $SAS^* = I$ and $S^{-*}BS^{-1}$ is diagonal.

Proof. Setting $S_1 = A^{-1/2}$, it follows that $S_1AS_1^* = I$. Since $S_1^{-*}BS_1^{-1}$ is Hermitian, it follows that there exists a unitary matrix $S_2 \in \mathbb{F}^{n \times n}$ such that $S^{-*}BS^{-1} = S_2^{-*}S_1^{-*}BS_1^{-1}S_2^{-1} = S_2(S_1^{-*}BS_1^{-1})S_2^*$ is diagonal, where $S = S_2S_1$. Finally, $SAS^* = S_2S_1AS_1^*S_2^* = S_2IS_2^* = I$. $\square$

Corollary 8.3.3. Let $A, B \in \mathbf{P}^n$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and $S^{-*}BS^{-1}$ are equal and diagonal.

Proof. By Theorem 8.3.2 there exists a nonsingular matrix $S_1 \in \mathbb{F}^{n \times n}$ such that $S_1AS_1^* = I$ and $B_1 = S_1^{-*}BS_1^{-1}$ is diagonal. Defining $S \triangleq B_1^{1/4}S_1$ yields $SAS^* = S^{-*}BS^{-1} = B_1^{1/2}$. $\square$

The transformation S of Corollary 8.3.3 is a *balancing transformation*.

Next, we weaken the requirement in Theorem 8.3.1 and Theorem 8.3.2 that A be positive definite by assuming only that A is positive semidefinite. In this case, however, we assume that B is also positive semidefinite.

Theorem 8.3.4. Let $A, B \in \mathbf{N}^n$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $SAS^* = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$ and SBS^* is diagonal.

Proof. Let the nonsingular matrix $S_1 \in \mathbb{F}^{n \times n}$ be such that $S_1AS_1^* = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$, and similarly partition $S_1BS_1^* = \begin{bmatrix} B_{11} & B_{12} \\ B_{12}^* & B_{22} \end{bmatrix}$, which is positive semidefinite. Letting $S_2 \triangleq \begin{bmatrix} I & -B_{12}B_{22}^+ \\ 0 & I \end{bmatrix}$, it follows from Lemma 8.2.1 that

$$S_2S_1BS_1^*S_2^* = \begin{bmatrix} B_{11} - B_{12}B_{22}^+B_{12}^* & 0 \\ 0 & B_{22} \end{bmatrix}.$$

Next, let U_1 and U_2 be unitary matrices such that $U_1(B_{11} - B_{12}B_{22}^+B_{12}^*)U_1^*$ and

$U_2 B_{22} U_2^*$ are diagonal. Then, defining $S_3 \triangleq \begin{bmatrix} U_1 & 0 \\ 0 & U_2 \end{bmatrix}$ and $S \triangleq S_3 S_2 S_1$, it follows that $SAS^* = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$ and $SB S^* = S_3 S_2 S_1 B S_1^* S_2^* S_3^*$ is diagonal. $\square$

Theorem 8.3.5. Let $A, B \in \mathbb{N}^n$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $SAS^* = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$ and $S^{-*}BS^{-1}$ is diagonal.

Proof. Let $S_1 \in \mathbb{F}^{n \times n}$ be a nonsingular matrix such that $S_1 A S_1^* = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$, and similarly partition $S_1^{-*} B S_1^{-1} = \begin{bmatrix} B_{11} & B_{12} \\ B_{12}^* & B_{22} \end{bmatrix}$, which is positive semidefinite. Letting $S_2 \triangleq \begin{bmatrix} I & B_{11}^+ B_{12} \\ 0 & I \end{bmatrix}$, it follows that

$$S_2^{-*} S_1^{-*} B S_1^{-1} S_2^{-1} = \begin{bmatrix} B_{11} & 0 \\ 0 & B_{22} - B_{12}^* B_{11}^+ B_{12} \end{bmatrix}.$$

Now, let U_1 and U_2 be unitary matrices such that $U_1 B_{11} U_1^*$ and $U_2 (B_{22} - B_{12}^* B_{11}^+ B_{12}) U_2^*$ are diagonal. Then, defining $S_3 \triangleq \begin{bmatrix} U_1 & 0 \\ 0 & U_2 \end{bmatrix}$ and $S \triangleq S_3 S_2 S_1$, it follows that $SAS^* = \begin{bmatrix} I & 0 \\ 0 & 0 \end{bmatrix}$ and $S^{-*} B S^{-1} = S_3^{-*} S_2^{-*} S_1^{-*} B S_1^{-1} S_2^{-1} S_3^{-1}$ is diagonal. $\square$

Corollary 8.3.6. Let $A, B \in \mathbb{N}^n$. Then, AB is semisimple, and every eigenvalue of AB is nonnegative. If, in addition, A and B are positive definite, then every eigenvalue of AB is positive.

Proof. It follows from Theorem 8.3.5 that there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that $A_1 = SAS^*$ and $B_1 = S^{-*} B S^{-1}$ are diagonal with nonnegative diagonal entries. Hence, $AB = S^{-1} A_1 B_1 S$ is semisimple and has nonnegative eigenvalues. $\square$

A more direct approach to showing that AB has nonnegative eigenvalues is to use Corollary 4.4.11 and note that $\lambda_i(AB) = \lambda_i(B^{1/2} A B^{1/2}) \geq 0$.

Corollary 8.3.7. Let $A, B \in \mathbb{N}^n$, and assume that $\text{rank } A = \text{rank } B = \text{rank } AB$. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $SAS^* = S^{-*} B S^{-1}$ and such that SAS^* is diagonal.

Proof. By Theorem 8.3.5 there exists a nonsingular matrix $S_1 \in \mathbb{F}^{n \times n}$ such that $S_1 A S_1^* = \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}$, where $r \triangleq \text{rank } A$, and such that $B_1 = S_1^{-*} B S_1^{-1}$ is diagonal. Hence, $AB = S_1^{-1} \begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix} B_1 S_1$. Since $\text{rank } A = \text{rank } B = \text{rank } AB = r$, it follows that $B_1 = \begin{bmatrix} \hat{B}_1 & 0 \\ 0 & 0 \end{bmatrix}$, where $\hat{B}_1 \in \mathbb{F}^{r \times r}$ is positive diagonal. Hence, $S_1^{-*} B S_1^{-1} = \begin{bmatrix} \hat{B}_1 & 0 \\ 0 & 0 \end{bmatrix}$. Now, define $S_2 \triangleq \begin{bmatrix} \hat{B}_1^{1/4} & 0 \\ 0 & I \end{bmatrix}$ and $S \triangleq S_2 S_1$. Then, $SAS^* = S_2 S_1 A S_1^* S_2^* = \begin{bmatrix} \hat{B}_1^{1/2} & 0 \\ 0 & 0 \end{bmatrix} = S_2^{-*} S_1^{-*} B S_1^{-1} S_2^{-1} = S^{-*} B S^{-1}$. $\square$

8.4 Eigenvalue Inequalities

Next, we turn our attention to inequalities involving eigenvalues. We begin with a series of lemmas.

Lemma 8.4.1. Let $A \in \mathbf{H}^n$, and let $\beta \in \mathbb{R}$. Then, the following statements hold:

- i) $\beta I \leq A$ if and only if $\beta \leq \lambda_{\min}(A)$.
- ii) $\beta I < A$ if and only if $\beta < \lambda_{\min}(A)$.
- iii) $A \leq \beta I$ if and only if $\lambda_{\max}(A) \leq \beta$.
- iv) $A < \beta I$ if and only if $\lambda_{\max}(A) < \beta$.

Proof. To prove i), assume that $\beta I \leq A$, and let $S \in \mathbb{F}^{n \times n}$ be a unitary matrix such that $B = SAS^*$ is diagonal. Then, $\beta I \leq B$, which yields $\beta \leq \lambda_{\min}(B) = \lambda_{\min}(A)$. Conversely, let $S \in \mathbb{F}^{n \times n}$ be a unitary matrix such that $B = SAS^*$ is diagonal. Since the diagonal entries of B are the eigenvalues of A , it follows that $\lambda_{\min}(A)I \leq B$, which implies that $\beta I \leq \lambda_{\min}(A)I \leq S^*BS = A$. Results ii), iii), and iv) are proved in a similar manner. $\square$

Corollary 8.4.2. Let $A \in \mathbf{H}^n$. Then,

$$\lambda_{\min}(A)I \leq A \leq \lambda_{\max}(A)I. \quad (8.4.1)$$

Proof. The result follows from i) and iii) of Lemma 8.4.1 with $\beta = \lambda_{\min}(A)$ and $\beta = \lambda_{\max}(A)$, respectively. $\square$

The following result concerns the maximum and minimum values of the *Rayleigh quotient*.

Lemma 8.4.3. Let $A \in \mathbf{H}^n$. Then,

$$\lambda_{\min}(A) = \min_{x \in \mathbb{F}^n \setminus \{0\}} \frac{x^*Ax}{x^*x} \quad (8.4.2)$$

and

$$\lambda_{\max}(A) = \max_{x \in \mathbb{F}^n \setminus \{0\}} \frac{x^*Ax}{x^*x}. \quad (8.4.3)$$

Proof. It follows from (8.4.1) that $\lambda_{\min}(A) \leq x^*Ax/x^*x$ for all nonzero $x \in \mathbb{F}^n$. Letting $x \in \mathbb{F}^n$ be an eigenvector of A associated with $\lambda_{\min}(A)$, it follows that this lower bound is attained. This proves (8.4.2). An analogous argument yields (8.4.3). $\square$

The following result is the *Cauchy interlacing theorem*.

Lemma 8.4.4. Let $A \in \mathbf{H}^n$, and let A_0 be an $(n-1) \times (n-1)$ principal submatrix of A . Then, for all $i = 1, \dots, n-1$,

$$\lambda_{i+1}(A) \leq \lambda_i(A_0) \leq \lambda_i(A). \quad (8.4.4)$$

Proof. Note that (8.4.4) is the chain of inequalities

$$\lambda_n(A) \leq \lambda_{n-1}(A_0) \leq \lambda_{n-1}(A) \leq \dots \leq \lambda_2(A) \leq \lambda_1(A_0) \leq \lambda_1(A).$$

Suppose that this chain of inequalities does not hold. In particular, first suppose that the rightmost inequality that is not true is $\lambda_j(A_0) \leq \lambda_j(A)$, so that $\lambda_j(A) <$

$\lambda_j(A_0)$. Choose δ such that $\lambda_j(A) < \delta < \lambda_j(A_0)$ and such that δ is not an eigenvalue of A_0 . If $j = 1$, then $A - \delta I$ is negative definite, while, if $j \geq 2$, then $\lambda_j(A) < \delta < \lambda_j(A_0) \leq \lambda_{j-1}(A_0) \leq \lambda_{j-1}(A)$, so that $A - \delta I$ has $j - 1$ positive eigenvalues. Thus, $\nu_+(A - \delta I) = j - 1$. Furthermore, since $\delta < \lambda_i(A_0)$, it follows that $\nu_+(A_0 - \delta I) \geq j$.

Now, assume for convenience that the rows and columns of A are ordered so that A_0 is the $(n-1) \times (n-1)$ leading principal submatrix of A . Thus, $A = \begin{bmatrix} A_0 & \beta \\ \beta^* & \gamma \end{bmatrix}$, where $\beta \in \mathbb{F}^{n-1}$ and $\gamma \in \mathbb{F}$. Next, note the identity

$$\begin{aligned} A - \delta I &= \begin{bmatrix} I & 0 \\ \beta^*(A_0 - \delta I)^{-1} & 1 \end{bmatrix} \begin{bmatrix} A_0 - \delta I & 0 \\ 0 & \gamma - \delta - \beta^*(A_0 - \delta I)^{-1}\beta \end{bmatrix} \begin{bmatrix} I & (A_0 - \delta I)^{-1}\beta \\ 0 & 1 \end{bmatrix}, \end{aligned}$$

where $A_0 - \delta I$ is nonsingular since δ is chosen to not be an eigenvalue of A_0 . Since the right-hand side of this identity involves a congruence transformation, and since $\nu_+(A_0 - \delta I) \geq j$, it follows from Corollary 5.4.7 that $\nu_+(A - \delta I) \geq j$. However, this inequality contradicts the fact that $\nu_+(A - \delta I) = j - 1$.

Finally, suppose that the rightmost inequality in (8.4.4) that is not true is $\lambda_{j+1}(A) \leq \lambda_j(A_0)$, so that $\lambda_j(A_0) < \lambda_{j+1}(A)$. Choose δ such that $\lambda_j(A_0) < \delta < \lambda_{j+1}(A)$ and such that δ is not an eigenvalue of A_0 . Then, it follows that $\nu_+(A - \delta I) \geq j + 1$ and $\nu_+(A_0 - \delta I) = j - 1$. Using the congruence transformation as in the previous case, it follows that $\nu_+(A - \delta I) \leq j$, which contradicts the fact that $\nu_+(A - \delta I) \geq j + 1$. $\square$

The following result is the *inclusion principle*.

Theorem 8.4.5. Let $A \in \mathbf{H}^n$, and let $A_0 \in \mathbf{H}^k$ be a $k \times k$ principal submatrix of A . Then, for all $i = 1, \dots, k$,

$$\lambda_{i+n-k}(A) \leq \lambda_i(A_0) \leq \lambda_i(A). \quad (8.4.5)$$

Proof. For $k = n - 1$, the result is given by Lemma 8.4.4. Hence, let $k = n - 2$, and let A_1 denote an $(n - 1) \times (n - 1)$ principal submatrix of A such that the $(n - 2) \times (n - 2)$ principal submatrix A_0 of A is also a principal submatrix of A_1 . Therefore, Lemma 8.4.4 implies that $\lambda_n(A) \leq \lambda_{n-1}(A_1) \leq \dots \leq \lambda_2(A_1) \leq \lambda_2(A) \leq \lambda_1(A_1) \leq \lambda_1(A)$ and $\lambda_{n-1}(A_1) \leq \lambda_{n-2}(A_0) \leq \dots \leq \lambda_2(A_0) \leq \lambda_2(A_1) \leq \lambda_1(A_0) \leq \lambda_1(A_1)$. Combining these inequalities yields $\lambda_{i+2}(A) \leq \lambda_i(A_0) \leq \lambda_i(A)$ for all $i = 1, \dots, n - 2$, while proceeding in a similar manner with $k < n - 2$ yields (8.4.5). $\square$

Corollary 8.4.6. Let $A \in \mathbf{H}^n$, and let $A_0 \in \mathbf{H}^k$ be a $k \times k$ principal submatrix of A . Then,

$$\lambda_{\min}(A) \leq \lambda_{\min}(A_0) \leq \lambda_{\max}(A_0) \leq \lambda_{\max}(A) \quad (8.4.6)$$

and

$$\lambda_{\min}(A_0) \leq \lambda_k(A). \quad (8.4.7)$$

The following result compares the maximum and minimum eigenvalues with the maximum and minimum diagonal entries.

Corollary 8.4.7. Let $A \in \mathbf{H}^n$. Then,

$$\lambda_{\min}(A) \leq d_{\min}(A) \leq d_{\max}(A) \leq \lambda_{\max}(A). \quad (8.4.8)$$

Lemma 8.4.8. Let $A, B \in \mathbf{H}^n$, and assume that $A \leq B$ and $\text{mspec}(A) = \text{mspec}(B)$. Then, $A = B$.

Proof. Let $\alpha \geq 0$ be such that $0 < \hat{A} \leq \hat{B}$, where $\hat{A} \triangleq A + \alpha I$ and $\hat{B} \triangleq B + \alpha I$. Note that $\text{mspec}(\hat{A}) = \text{mspec}(\hat{B})$, and thus $\det \hat{A} = \det \hat{B}$. Next, it follows that $I \leq \hat{A}^{-1/2} \hat{B} \hat{A}^{-1/2}$. Hence, it follows from *i*) of Lemma 8.4.1 that $\lambda_{\min}(\hat{A}^{-1/2} \hat{B} \hat{A}^{-1/2}) \geq 1$. Furthermore, $\det(\hat{A}^{-1/2} \hat{B} \hat{A}^{-1/2}) = \det \hat{B} / \det \hat{A} = 1$, which implies that $\lambda_i(\hat{A}^{-1/2} \hat{B} \hat{A}^{-1/2}) = 1$ for all $i = 1, \dots, n$. Hence, $\hat{A}^{-1/2} \hat{B} \hat{A}^{-1/2} = I$, and thus $\hat{A} = \hat{B}$. Hence, $A = B$. $\square$

The following result is the *monotonicity theorem* or *Weyl's inequality*.

Theorem 8.4.9. Let $A, B \in \mathbf{H}^n$, and assume that $A \leq B$. Then, for all $i = 1, \dots, n$,

$$\lambda_i(A) \leq \lambda_i(B). \quad (8.4.9)$$

If $A \neq B$, then there exists $i \in \{1, \dots, n\}$ such that

$$\lambda_i(A) < \lambda_i(B). \quad (8.4.10)$$

If $A < B$, then (8.4.10) holds for all $i = 1, \dots, n$.

Proof. Since $A \leq B$, it follows from Corollary 8.4.2 that $\lambda_{\min}(A)I \leq A \leq B \leq \lambda_{\max}(B)I$. Hence, it follows from *iii*) and *i*) of Lemma 8.4.1 that $\lambda_{\min}(A) \leq \lambda_{\min}(B)$ and $\lambda_{\max}(A) \leq \lambda_{\max}(B)$. Next, let $S \in \mathbb{F}^{n \times n}$ be a unitary matrix such that $SAS^* = \text{diag}[\lambda_1(A), \dots, \lambda_n(A)]$. Furthermore, for $2 \leq i \leq n-1$, let $A_0 = \text{diag}[\lambda_1(A), \dots, \lambda_i(A)]$, and let B_0 denote the $i \times i$ leading principal submatrices of SAS^* and SBS^* , respectively. Since $A \leq B$, it follows that $A_0 \leq B_0$, which implies that $\lambda_{\min}(A_0) \leq \lambda_{\min}(B_0)$. It now follows from (8.4.7) that

$$\lambda_i(A) = \lambda_{\min}(A_0) \leq \lambda_{\min}(B_0) \leq \lambda_i(SBS^*) = \lambda_i(B),$$

which proves (8.4.9). If $A \neq B$, then it follows from Lemma 8.4.8 that $\text{mspec}(A) \neq \text{mspec}(B)$ and thus there exists $i \in \{1, \dots, n\}$ such that (8.4.10) holds. If $A < B$, then $\lambda_{\min}(A_0) < \lambda_{\min}(B_0)$, which implies that (8.4.10) holds for all $i = 1, \dots, n$. $\square$

Corollary 8.4.10. Let $A, B \in \mathbf{H}^n$. Then, the following statements hold:

- i*) If $A \leq B$, then $\text{tr } A \leq \text{tr } B$.
- ii*) If $A \leq B$ and $\text{tr } A = \text{tr } B$, then $A = B$.
- iii*) If $A < B$, then $\text{tr } A < \text{tr } B$.
- iv*) If $0 \leq A \leq B$, then $0 \leq \det A \leq \det B$.
- v*) If $0 \leq A < B$, then $0 \leq \det A < \det B$.

vi) If $0 < A \leq B$ and $\det A = \det B$, then $A = B$.

Proof. Statements *i)*, *iii)*, *iv)*, and *v)* follow from Theorem 8.4.9. To prove *ii)*, note that, since $A \leq B$ and $\operatorname{tr} A = \operatorname{tr} B$, it follows from Theorem 8.4.9 that $\operatorname{mspec}(A) = \operatorname{mspec}(B)$. Now, Lemma 8.4.8 implies that $A = B$. A similar argument yields *vi)*. $\square$

The following result, which is a generalization of Theorem 8.4.9, is due to Weyl.

Theorem 8.4.11. Let $A, B \in \mathbf{H}^n$. If $i + j \geq n + 1$, then

$$\lambda_i(A) + \lambda_j(B) \leq \lambda_{i+j-n}(A + B). \quad (8.4.11)$$

If $i + j \leq n + 1$, then

$$\lambda_{i+j-1}(A + B) \leq \lambda_i(A) + \lambda_j(B). \quad (8.4.12)$$

In particular, for all $i = 1, \dots, n$,

$$\lambda_i(A) + \lambda_{\min}(B) \leq \lambda_i(A + B) \leq \lambda_i(A) + \lambda_{\max}(B), \quad (8.4.13)$$

$$\lambda_{\min}(A) + \lambda_{\min}(B) \leq \lambda_{\min}(A + B) \leq \lambda_{\min}(A) + \lambda_{\max}(B), \quad (8.4.14)$$

$$\lambda_{\max}(A) + \lambda_{\min}(B) \leq \lambda_{\max}(A + B) \leq \lambda_{\max}(A) + \lambda_{\max}(B). \quad (8.4.15)$$

Furthermore,

$$\nu_-(A + B) \leq \nu_-(A) + \nu_-(B) \quad (8.4.16)$$

and

$$\nu_+(A + B) \leq \nu_+(A) + \nu_+(B). \quad (8.4.17)$$

Proof. See [709, p. 182]. The last two inequalities are noted in [393]. $\square$

Lemma 8.4.12. Let $A, B, C \in \mathbf{H}^n$. If $A \leq B$ and C is positive semidefinite, then

$$\operatorname{tr} AC \leq \operatorname{tr} BC. \quad (8.4.18)$$

If $A < B$ and C is positive definite, then

$$\operatorname{tr} AC < \operatorname{tr} BC. \quad (8.4.19)$$

Proof. Since $C^{1/2}AC^{1/2} \leq C^{1/2}BC^{1/2}$, it follows from *i)* of Corollary 8.4.10 that

$$\operatorname{tr} AC = \operatorname{tr} C^{1/2}AC^{1/2} \leq \operatorname{tr} C^{1/2}BC^{1/2} = \operatorname{tr} BC.$$

Result (8.4.19) follows from *ii)* of Corollary 8.4.10 in a similar fashion. $\square$

Proposition 8.4.13. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that B is positive semidefinite. Then,

$$\frac{1}{2}\lambda_{\min}(A + A^*) \operatorname{tr} B \leq \operatorname{Re} \operatorname{tr} AB \leq \frac{1}{2}\lambda_{\max}(A + A^*) \operatorname{tr} B. \quad (8.4.20)$$

If, in addition, A is Hermitian, then

$$\lambda_{\min}(A) \operatorname{tr} B \leq \operatorname{tr} AB \leq \lambda_{\max}(A) \operatorname{tr} B. \quad (8.4.21)$$

Proof. It follows from Corollary 8.4.2 that $\frac{1}{2}\lambda_{\min}(A+A^*)I \leq \frac{1}{2}(A+A^*)$, while Lemma 8.4.12 implies that $\frac{1}{2}\lambda_{\min}(A+A^*)\operatorname{tr} B = \operatorname{tr} \frac{1}{2}\lambda_{\min}(A+A^*)IB \leq \operatorname{tr} \frac{1}{2}(A+A^*)B = \operatorname{Re} \operatorname{tr} AB$, which proves the left-hand inequality of (8.4.20). Similarly, the right-hand inequality holds. $\square$

For results relating to Proposition 8.4.13, see Fact 5.12.4, Fact 5.12.5, Fact 5.12.8, and Fact 8.18.18.

Proposition 8.4.14. Let $A, B \in \mathbf{P}^n$, and assume that $\det B = 1$. Then,

$$(\det A)^{1/n} \leq \frac{1}{n} \operatorname{tr} AB. \quad (8.4.22)$$

Furthermore, equality holds if and only if $B = (\det A)^{1/n}A^{-1}$.

Proof. Using the arithmetic-mean–geometric-mean inequality given by Fact 1.15.14, it follows that

$$\begin{aligned} (\det A)^{1/n} &= \left(\det B^{1/2}AB^{1/2} \right)^{1/n} = \left[\prod_{i=1}^n \lambda_i \left(B^{1/2}AB^{1/2} \right) \right]^{1/n} \\ &\leq \frac{1}{n} \sum_{i=1}^n \lambda_i \left(B^{1/2}AB^{1/2} \right) = \frac{1}{n} \operatorname{tr} AB. \end{aligned}$$

Equality holds if and only if there exists $\beta > 0$ such that $B^{1/2}AB^{1/2} = \beta I$. In this case, $\beta = (\det A)^{1/n}$ and $B = (\det A)^{1/n}A^{-1}$. $\square$

The following corollary of Proposition 8.4.14 is *Minkowski's determinant theorem*.

Corollary 8.4.15. Let $A, B \in \mathbf{N}^n$, and let $p \in [1, n]$. Then,

$$\det A + \det B \leq \left[(\det A)^{1/p} + (\det B)^{1/p} \right]^p \quad (8.4.23)$$

$$\leq \left[(\det A)^{1/n} + (\det B)^{1/n} \right]^n \quad (8.4.24)$$

$$\leq \det(A+B). \quad (8.4.25)$$

Furthermore, the following statements hold:

- i) If $A = 0$ or $B = 0$ or $\det(A+B) = 0$, then (8.4.23)–(8.4.25) are identities.
- ii) If there exists $\alpha \geq 0$ such that $B = \alpha A$, then (8.4.25) is an identity.
- iii) If $A+B$ is positive definite and (8.4.25) holds as an identity, then there exists $\alpha \geq 0$ such that either $B = \alpha A$ or $A = \alpha B$.
- iv) If $n \geq 2$, $p > 1$, A is positive definite, and (8.4.23) holds as an identity, then $\det B = 0$.
- v) If $n \geq 2$, $p < n$, A is positive definite, and (8.4.24) holds as an identity, then $\det B = 0$.
- vi) If $n \geq 2$, A is positive definite, and $\det A + \det B = \det(A+B)$, then $B = 0$.

Proof. Inequalities (8.4.23) and (8.4.24) are consequences of the power-sum inequality Fact 1.15.34. Now, assume that $A+B$ is positive definite, since otherwise (8.4.23)–(8.4.25) are identities. To prove (8.4.25), Proposition 8.4.14 implies that

$$\begin{aligned} (\det A)^{1/n} + (\det B)^{1/n} &\leq \frac{1}{n} \operatorname{tr} \left[A[\det(A+B)]^{1/n} (A+B)^{-1} \right] \\ &\quad + \frac{1}{n} \operatorname{tr} \left[B[\det(A+B)]^{1/n} (A+B)^{-1} \right] \\ &= [\det(A+B)]^{1/n}. \end{aligned}$$

Statements *i*) and *ii*) are immediate. To prove *iii*), suppose that $A+B$ is positive definite and that (8.4.25) holds as an identity. Then, either A or B is positive definite. Hence, suppose that A is positive definite. Multiplying the identity $(\det A)^{1/n} + (\det B)^{1/n} = [\det(A+B)]^{1/n}$ by $(\det A)^{-1/n}$ yields

$$1 + \left(\det A^{-1/2} B A^{-1/2} \right)^{1/n} = \left[\det \left(I + A^{-1/2} B A^{-1/2} \right) \right]^{1/n}.$$

Letting $\lambda_1, \dots, \lambda_n$ denote the eigenvalues of $A^{-1/2} B A^{-1/2}$, it follows that $1 + (\lambda_1 \cdots \lambda_n)^{1/n} = [(1 + \lambda_1) \cdots (1 + \lambda_n)]^{1/n}$. It now follows from Fact 1.15.33 that $\lambda_1 = \cdots = \lambda_n$.

To prove *iv*), note that, since $1/p < 1$, $\det A > 0$, and identity holds in (8.4.23), it follows from Fact 1.15.34 that $\det B = 0$.

To prove *v*), note that, since $1/n < 1/p$, $\det A > 0$, and identity holds in (8.4.24), it follows from Fact 1.15.34 that $\det B = 0$.

To prove *vi*), note that (8.4.23) and (8.4.24) hold as identities for all $p \in [1, n]$. Therefore, $\det B = 0$. Consequently, $\det A = \det(A+B)$. Since $0 < A \leq A+B$, it follows from *vi*) of Corollary 8.4.10 that $B = 0$. $\square$

8.5 Exponential, Square Root, and Logarithm of Hermitian Matrices

Let $A = SBS^* \in \mathbb{F}^{n \times n}$ be Hermitian, where $S \in \mathbb{F}^{n \times n}$ is unitary, $B \in \mathbb{R}^{n \times n}$ is diagonal, $\operatorname{spec}(A) \subset \mathcal{D}$, and $\mathcal{D} \subseteq \mathbb{R}$. Furthermore, let $f: \mathcal{D} \mapsto \mathbb{R}$. Then, we define $f(A) \in \mathbb{H}^n$ by

$$f(A) \triangleq S f(B) S^*, \quad (8.5.1)$$

where $[f(B)]_{(i,i)} \triangleq f[B_{(i,i)}]$. Hence, with an obvious extension of notation, $f: \{X \in \mathbb{H}^n: \operatorname{spec}(X) \subset \mathcal{D}\} \mapsto \mathbb{H}^n$. If $f: \mathcal{D} \mapsto \mathbb{R}$ is one-to-one, then its inverse $f^{-1}: \{X \in \mathbb{H}^n: \operatorname{spec}(X) \subset f(\mathcal{D})\} \mapsto \mathbb{H}^n$ exists.

Let $A = SBS^* \in \mathbb{F}^{n \times n}$ be Hermitian, where $S \in \mathbb{F}^{n \times n}$ is unitary and $B \in \mathbb{R}^{n \times n}$ is diagonal. Then, the *matrix exponential* is defined by

$$e^A \triangleq S e^B S^* \in \mathbb{H}^n, \quad (8.5.2)$$

where, for all $i = 1, \dots, n$, $(e^B)_{(i,i)} \triangleq e^{B_{(i,i)}}$.

Let $A = SBS^* \in \mathbb{F}^{n \times n}$ be positive semidefinite, where $S \in \mathbb{F}^{n \times n}$ is unitary and $B \in \mathbb{R}^{n \times n}$ is diagonal with nonnegative entries. Then, for all $r \geq 0$ (not necessarily an integer), $A^r = SB^rS^*$ is positive semidefinite, where, for all $i = 1, \dots, n$, $(B^r)_{(i,i)} = [B_{(i,i)}]^r$. Note that $A^0 \triangleq I$. In particular, the positive-semidefinite matrix

$$A^{1/2} = SB^{1/2}S^* \quad (8.5.3)$$

is a square root of A since

$$A^{1/2}A^{1/2} = SB^{1/2}S^*SB^{1/2}S^* = SBS^* = A. \quad (8.5.4)$$

The uniqueness of the *positive-semidefinite square root* of A given by (8.5.3) follows from Theorem 10.6.1; see also [711, p. 410] or [877]. Uniqueness can also be shown directly; see [447, pp. 265, 266] or [709, p. 405]. Hence, if $C \in \mathbb{F}^{n \times m}$, then C^*C is positive semidefinite, and we define

$$\langle C \rangle \triangleq (C^*C)^{1/2}. \quad (8.5.5)$$

If A is positive definite, then A^r is positive definite for all $r \in \mathbb{R}$, and, if $r \neq 0$, then $(A^r)^{1/r} = A$.

Now, assume that A is positive definite. Then, the *matrix logarithm* is defined by

$$\log A \triangleq S(\log B)S^* \in \mathbf{H}^n, \quad (8.5.6)$$

where, for all $i = 1, \dots, n$, $(\log B)_{(i,i)} \triangleq \log[B_{(i,i)}]$.

In chapters 10 and 11, the matrix exponential, square root, and logarithm are extended to matrices that are not necessarily Hermitian.

8.6 Matrix Inequalities

Lemma 8.6.1. Let $A, B \in \mathbb{F}^n$, assume that A and B are Hermitian, and assume that $0 \leq A \leq B$. Then, $\mathcal{R}(A) \subseteq \mathcal{R}(B)$.

Proof. Let $x \in \mathcal{N}(B)$. Then, $x^*Bx = 0$, and thus $x^*Ax = 0$, which implies that $Ax = 0$. Hence, $\mathcal{N}(B) \subseteq \mathcal{N}(A)$, and thus $\mathcal{N}(A)^\perp \subseteq \mathcal{N}(B)^\perp$. Since A and B are Hermitian, it follows from Theorem 2.4.3 that $\mathcal{R}(A) = \mathcal{N}(A)^\perp$ and $\mathcal{R}(B) = \mathcal{N}(B)^\perp$. Hence, $\mathcal{R}(A) \subseteq \mathcal{R}(B)$. $\square$

The following result is the *Douglas-Fillmore-Williams lemma* [427, 490].

Theorem 8.6.2. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then, the following statements are equivalent:

- i) There exists a matrix $C \in \mathbb{F}^{l \times m}$ such that $A = BC$.
- ii) There exists $\alpha > 0$ such that $AA^* \leq \alpha BB^*$.
- iii) $\mathcal{R}(A) \subseteq \mathcal{R}(B)$.

Proof. First we prove that $i)$ implies $ii)$. Since $A = BC$, it follows that $AA^* = BCC^*B^*$. Since $CC^* \leq \lambda_{\max}(CC^*)I$, it follows that $AA^* \leq \alpha BB^*$, where $\alpha \triangleq \lambda_{\max}(CC^*)$. To prove that $ii)$ implies $iii)$, first note that Lemma 8.6.1 implies that $\mathcal{R}(AA^*) \subseteq \mathcal{R}(\alpha BB^*) = \mathcal{R}(BB^*)$. Since, by Theorem 2.4.3, $\mathcal{R}(AA^*) = \mathcal{R}(A)$ and $\mathcal{R}(BB^*) = \mathcal{R}(B)$, it follows that $\mathcal{R}(A) \subseteq \mathcal{R}(B)$. Finally, to prove that $iii)$ implies $i)$, use Theorem 5.6.4 to write $B = S_1 \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix} S_2$, where $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{l \times l}$ are unitary and $D \in \mathbb{R}^{r \times r}$ is diagonal with positive diagonal entries, where $r \triangleq \text{rank } B$. Since $\mathcal{R}(S_1^*A) \subseteq \mathcal{R}(S_1^*B)$ and $S_1^*B = \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix} S_2$, it follows that $S_1^*A = \begin{bmatrix} A_1 \\ 0 \end{bmatrix}$, where $A_1 \in \mathbb{F}^{r \times m}$. Consequently,

$$A = S_1 \begin{bmatrix} A_1 \\ 0 \end{bmatrix} = S_1 \begin{bmatrix} D & 0 \\ 0 & 0 \end{bmatrix} S_2 S_2^* \begin{bmatrix} D^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A_1 \\ 0 \end{bmatrix} = BC,$$

where $C \triangleq S_2^* \begin{bmatrix} D^{-1} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A_1 \\ 0 \end{bmatrix} \in \mathbb{F}^{l \times m}$. $\square$

Proposition 8.6.3. Let $(A_i)_{i=1}^\infty \subset \mathbf{N}^n$ satisfy $0 \leq A_i \leq A_j$ for all $i \leq j$, and assume there exists $B \in \mathbf{N}^n$ satisfying $A_i \leq B$ for all $i \in \mathbb{P}$. Then, $A \triangleq \lim_{i \rightarrow \infty} A_i$ exists and satisfies $0 \leq A \leq B$.

Proof. Let $k \in \{1, \dots, n\}$. Then, the sequence $(A_{i(k,k)})_{i=1}^\infty$ is nondecreasing and bounded from above. Hence, $A_{(k,k)} \triangleq \lim_{i \rightarrow \infty} A_{i(k,k)}$ exists. Now, let $k, l \in \{1, \dots, n\}$, where $k \neq l$. Since $A_i \leq A_j$ for all $i < j$, it follows that $(e_k + e_l)^* A_i (e_k + e_l) \leq (e_k + e_l)^* A_j (e_k + e_l)$, which implies that $A_{i(k,k)} - A_{j(k,l)} \leq \frac{1}{2} [A_{j(k,k)} - A_{i(k,k)} + A_{j(l,l)} - A_{i(l,l)}]$. Alternatively, replacing $e_k + e_l$ by $e_k - e_l$ yields $A_{j(k,l)} - A_{i(k,l)} \leq \frac{1}{2} [A_{j(k,k)} - A_{i(k,k)} + A_{j(l,l)} - A_{i(l,l)}]$. Thus, $A_{i(k,l)} - A_{j(k,l)} \rightarrow 0$ as $i, j \rightarrow \infty$, which implies that $A_{(k,l)} \triangleq \lim_{i \rightarrow \infty} A_{i(k,l)}$ exists. Hence, $A \triangleq \lim_{i \rightarrow \infty} A_i$ exists. Since $A_i \leq B$ for all $i = 1, 2, \dots$, it follows that $A \leq B$. $\square$

Proposition 8.6.4. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and let $p > 0$. Then,

$$A^{-1}(A - I) \leq \log A \leq p^{-1}(A^p - I) \quad (8.6.1)$$

and

$$\log A = \lim_{p \downarrow 0} p^{-1}(A^p - I). \quad (8.6.2)$$

Proof. The result follows from Fact 1.9.26. $\square$

Lemma 8.6.5. Let $A \in \mathbf{P}^n$. If $A \leq I$, then $I \leq A^{-1}$. Furthermore, if $A < I$, then $I < A^{-1}$.

Proof. Since $A \leq I$, it follows from $xi)$ of Proposition 8.1.2 that $I = A^{-1/2} A A^{-1/2} \leq A^{-1/2} I A^{-1/2} = A^{-1}$. Similarly, $A < I$ implies that $I = A^{-1/2} A A^{-1/2} < A^{-1/2} I A^{-1/2} = A^{-1}$. $\square$

Proposition 8.6.6. Let $A, B \in \mathbf{H}^n$, and assume that either A and B are positive definite or A and B are negative definite. If $A \leq B$, then $B^{-1} \leq A^{-1}$. If, in addition, $A < B$, then $B^{-1} < A^{-1}$.

Proof. Suppose that A and B are positive definite. Since $A \leq B$, it follows that $B^{-1/2}AB^{-1/2} \leq I$. Now, Lemma 8.6.5 implies that $I \leq B^{1/2}A^{-1}B^{1/2}$, which implies that $B^{-1} \leq A^{-1}$. If A and B are negative definite, then $A \leq B$ is equivalent to $-B \leq -A$. The case $A < B$ is proved in a similar manner. $\square$

The following result is the *Furuta inequality*.

Proposition 8.6.7. Let $A, B \in \mathbf{N}^n$, and assume that $0 \leq A \leq B$. Furthermore, let $p, q, r \in \mathbb{R}$ satisfy $p \geq 0$, $q \geq 1$, $r \geq 0$, and $p + 2r \leq (1 + 2r)q$. Then,

$$A^{(p+2r)/q} \leq (A^r B^p A^r)^{1/q} \quad (8.6.3)$$

and

$$(B^r A^p B^r)^{1/q} \leq B^{(p+2r)/q}. \quad (8.6.4)$$

Proof. See [522] or [530, pp. 129, 130]. $\square$

Corollary 8.6.8. Let $A, B \in \mathbf{N}^n$, and assume that $0 \leq A \leq B$. Then,

$$A^2 \leq (AB^2A)^{1/2} \quad (8.6.5)$$

and

$$(BA^2B)^{1/2} \leq B^2. \quad (8.6.6)$$

Proof. In Proposition 8.6.7 set $r = 1$, $p = 2$, and $q = 2$. $\square$

Corollary 8.6.9. Let $A, B, C \in \mathbf{N}^n$, and assume that $0 \leq A \leq C \leq B$. Then,

$$(CA^2C)^{1/2} \leq C^2 \leq (CB^2C)^{1/2}. \quad (8.6.7)$$

Proof. The result follows from Corollary 8.6.8. See also [1395]. $\square$

The following result provides representations for A^r , where $r \in (0, 1)$.

Proposition 8.6.10. Let $A \in \mathbf{P}^n$ and $r \in (0, 1)$. Then,

$$A^r = \left(\cos \frac{r\pi}{2} \right) I + \frac{\sin r\pi}{\pi} \int_0^\infty \left[\frac{x^{r+1}}{1+x^2} I - (A + xI)^{-1} x^r \right] dx \quad (8.6.8)$$

and

$$A^r = \frac{\sin r\pi}{\pi} \int_0^\infty (A + xI)^{-1} A x^{r-1} dx. \quad (8.6.9)$$

Proof. Let $t \geq 0$. As shown in [193], [197, p. 143],

$$\int_0^\infty \left[\frac{x^{r+1}}{1+x^2} - \frac{x^r}{t+x} \right] dx = \frac{\pi}{\sin r\pi} \left(t^r - \cos \frac{r\pi}{2} \right).$$

Solving for t^r and replacing t by A yields (8.6.8). Likewise, replacing t by A in *xxvii*) of Fact 1.19.1 yields (8.6.9). $\square$

The following result is the *Löwner-Heinz inequality*.

Corollary 8.6.11. Let $A, B \in \mathbf{N}^n$, assume that $0 \leq A \leq B$, and let $r \in [0, 1]$. Then, $A^r \leq B^r$. If, in addition, $A < B$ and $r \in (0, 1]$, then $A^r < B^r$.

Proof. Let $0 < A \leq B$, and let $r \in (0, 1)$. In Proposition 8.6.7, replace p, q, r with $r, 1, 0$. The first result now follows from (8.6.3). Alternatively, it follows from (8.6.8) of Proposition 8.6.10 that

$$B^r - A^r = \frac{\sin r\pi}{\pi} \int_0^\infty [(A + xI)^{-1} - (B + xI)^{-1}] x^r dx.$$

Since $A \leq B$, it follows from Proposition 8.6.6 that, for all $x \geq 0$, $(B + xI)^{-1} \leq (A + xI)^{-1}$. Hence, $A^r \leq B^r$. By continuity, the result holds for $A, B \in \mathbf{N}^n$ and $r \in [0, 1]$. In the case $A < B$, it follows from Proposition 8.6.6 that, for all $x \geq 0$, $(B + xI)^{-1} < (A + xI)^{-1}$, so that $A^r < B^r$.

Alternatively, it follows from (8.6.9) of Proposition 8.6.10 that

$$B^r - A^r = \frac{\sin r\pi}{\pi} \int_0^\infty [(A + xI)^{-1}A - (B + xI)^{-1}B] x^{r-1} dx.$$

Since $A \leq B$, it follows that, for all $x \geq 0$, $(B + xI)^{-1}B \leq (A + xI)^{-1}A$. Hence, $A^r \leq B^r$. Alternative proofs are given in [530, p. 127] and [1485, p. 2].

For the case $r = 1/2$, let $\lambda \in \mathbb{R}$ be an eigenvalue of $B^{1/2} - A^{1/2}$, and let $x \in \mathbb{F}^n$ be an associated eigenvector. Then,

$$\begin{aligned} \lambda x^* (B^{1/2} + A^{1/2}) x &= x^* (B^{1/2} + A^{1/2}) (B^{1/2} - A^{1/2}) x \\ &= x^* (B - B^{1/2}A^{1/2} + A^{1/2}B^{1/2} - A) x \\ &= x^* (B - A) x \geq 0. \end{aligned}$$

Since $B^{1/2} + A^{1/2}$ is positive semidefinite, it follows that either $\lambda \geq 0$ or $x^* (B^{1/2} + A^{1/2}) x = 0$. In the latter case, $B^{1/2}x = A^{1/2}x = 0$, which implies that $\lambda = 0$. $\square$

The Löwner-Heinz inequality does not extend to $r > 1$. In fact, $A \triangleq \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}$ and $B \triangleq \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ satisfy $A \geq B \geq 0$, whereas, for all $r > 1$, $A^r \not\geq B^r$. For details, see [530, pp. 127, 128].

Many of the results given so far involve functions that are nondecreasing or increasing on suitable sets of matrices.

Definition 8.6.12. Let $\mathcal{D} \subseteq \mathbf{H}^n$, and let $\phi: \mathcal{D} \mapsto \mathbf{H}^m$. Then, the following terminology is defined:

- i) ϕ is *nondecreasing* if, for all $A, B \in \mathcal{D}$ such that $A \leq B$, it follows that $\phi(A) \leq \phi(B)$.

- ii) ϕ is *increasing* if ϕ is nondecreasing and, for all $A, B \in \mathcal{D}$ such that $A < B$, it follows that $\phi(A) < \phi(B)$.
- iii) ϕ is *strongly increasing* if ϕ is nondecreasing and, for all $A, B \in \mathcal{D}$ such that $A \leq B$ and $A \neq B$, it follows that $\phi(A) < \phi(B)$.
- iv) ϕ is (*nonincreasing, decreasing, strongly decreasing*) if $-\phi$ is (nondecreasing, increasing, strongly increasing).

Proposition 8.6.13. The following functions are nondecreasing:

- i) $\phi: \mathbf{H}^n \mapsto \mathbf{H}^m$ defined by $\phi(A) \triangleq BAB^*$, where $B \in \mathbb{F}^{m \times n}$.
- ii) $\phi: \mathbf{H}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq \text{tr } AB$, where $B \in \mathbf{N}^n$.
- iii) $\phi: \mathbf{N}^{n+m} \mapsto \mathbf{N}^n$ defined by $\phi(A) \triangleq A_{22}|A$, where $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$.
- iv) $\phi: \mathbf{N}^n \times \mathbf{N}^m \mapsto \mathbf{N}^{nm}$ defined by $\phi(A, B) \triangleq A^{r_1} \otimes B^{r_2}$, where $r_1, r_2 \in [0, 1]$ satisfy $r_1 + r_2 \leq 1$.
- v) $\phi: \mathbf{N}^n \times \mathbf{N}^n \mapsto \mathbf{N}^n$ defined by $\phi(A, B) \triangleq A^{r_1} \circ B^{r_2}$, where $r_1, r_2 \in [0, 1]$ satisfy $r_1 + r_2 \leq 1$.

The following functions are increasing:

- vi) $\phi: \mathbf{H}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq \lambda_i(A)$, where $i \in \{1, \dots, n\}$.
- vii) $\phi: \mathbf{N}^n \mapsto \mathbf{N}^n$ defined by $\phi(A) \triangleq A^r$, where $r \in [0, 1]$.
- viii) $\phi: \mathbf{N}^n \mapsto \mathbf{N}^n$ defined by $\phi(A) \triangleq A^{1/2}$.
- ix) $\phi: \mathbf{P}^n \mapsto -\mathbf{P}^n$ defined by $\phi(A) \triangleq -A^{-r}$, where $r \in [0, 1]$.
- x) $\phi: \mathbf{P}^n \mapsto -\mathbf{P}^n$ defined by $\phi(A) \triangleq -A^{-1}$.
- xi) $\phi: \mathbf{P}^n \mapsto -\mathbf{P}^n$ defined by $\phi(A) \triangleq -A^{-1/2}$.
- xii) $\phi: -\mathbf{P}^n \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq (-A)^{-r}$, where $r \in [0, 1]$.
- xiii) $\phi: -\mathbf{P}^n \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq -A^{-1}$.
- xiv) $\phi: -\mathbf{P}^n \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq -A^{-1/2}$.
- xv) $\phi: \mathbf{H}^n \mapsto \mathbf{H}^m$ defined by $\phi(A) \triangleq BAB^*$, where $B \in \mathbb{F}^{m \times n}$ and $\text{rank } B = m$.
- xvi) $\phi: \mathbf{P}^{n+m} \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq A_{22}|A$, where $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$.
- xvii) $\phi: \mathbf{P}^{n+m} \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq -(A_{22}|A)^{-1}$, where $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$.
- xviii) $\phi: \mathbf{P}^n \mapsto \mathbf{H}^n$ defined by $\phi(A) \triangleq \log A$.

The following functions are strongly increasing:

- xix) $\phi: \mathbf{H}^n \mapsto [0, \infty)$ defined by $\phi(A) \triangleq \text{tr } BAB^*$, where $B \in \mathbb{F}^{m \times n}$ and $\text{rank } B = m$.
- xx) $\phi: \mathbf{H}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq \text{tr } AB$, where $B \in \mathbf{P}^n$.

xxi) $\phi: \mathbf{N}^n \mapsto [0, \infty)$ defined by $\phi(A) \triangleq \operatorname{tr} A^r$, where $r > 0$.

xxii) $\phi: \mathbf{N}^n \mapsto [0, \infty)$ defined by $\phi(A) \triangleq \det A$.

Proof. For the proof of *iii)*, see [896]. To prove *xviii)*, let $A, B \in \mathbf{P}^n$, and assume that $A \leq B$. Then, for all $r \in [0, 1]$, it follows from *vii)* that $r^{-1}(A^r - I) \leq r^{-1}(B^r - I)$. Letting $r \downarrow 0$ and using Proposition 8.6.4 yields $\log A \leq \log B$, which proves that $\log$ is nondecreasing. See [530, p. 139] and Fact 8.19.2. To prove that $\log$ is increasing, assume that $A < B$, and let $\varepsilon > 0$ be such that $A + \varepsilon I < B$. Then, it follows that $\log A < \log(A + \varepsilon I) \leq \log B$. $\square$

Finally, we consider convex functions defined with respect to matrix inequalities. The following definition generalizes Definition 1.2.3 in the case $n = m = p = 1$.

Definition 8.6.14. Let $\mathcal{D} \subseteq \mathbb{F}^{n \times m}$ be a convex set, and let $\phi: \mathcal{D} \mapsto \mathbf{H}^p$. Then, the following terminology is defined:

i) ϕ is *convex* if, for all $\alpha \in [0, 1]$ and $A_1, A_2 \in \mathcal{D}$,

$$\phi[\alpha A_1 + (1 - \alpha)A_2] \leq \alpha\phi(A_1) + (1 - \alpha)\phi(A_2). \quad (8.6.10)$$

ii) ϕ is *concave* if $-\phi$ is convex.

iii) ϕ is *strictly convex* if, for all $\alpha \in (0, 1)$ and distinct $A_1, A_2 \in \mathcal{D}$,

$$\phi[\alpha A_1 + (1 - \alpha)A_2] < \alpha\phi(A_1) + (1 - \alpha)\phi(A_2). \quad (8.6.11)$$

iv) ϕ is *strictly concave* if $-\phi$ is strictly convex.

Theorem 8.6.15. Let $\mathcal{S} \subseteq \mathbb{R}$, let $\phi: \mathcal{S}_1 \mapsto \mathcal{S}_2$, and assume that ϕ is continuous. Then, the following statements hold:

i) Assume that $\mathcal{S}_1 = \mathcal{S}_2 = (0, \infty)$ and $\phi: \mathbf{P}^n \mapsto \mathbf{P}^n$ is increasing. Then, $\psi: \mathbf{P}^n \mapsto \mathbf{P}^n$ defined by $\psi(x) = 1/\phi(x)$ is convex.

ii) Assume that $\mathcal{S}_1 = \mathcal{S}_2 = [0, \infty)$. Then, $\phi: \mathbf{N}^n \mapsto \mathbf{N}^n$ is increasing if and only if $\phi: \mathbf{N}^n \mapsto \mathbf{N}^n$ is concave.

iii) Assume that $\mathcal{S}_1 = [0, \infty)$ and $\mathcal{S}_2 = \mathbb{R}$. Then, $\phi: \mathbf{N}^n \mapsto \mathbf{H}^n$ is convex and $\phi(0) \leq 0$ if and only if $\psi: \mathbf{P}^n \mapsto \mathbf{H}^n$ defined by $\psi(x) = \phi(x)/x$ is increasing.

Proof. See [197, pp. 120–122]. $\square$

Lemma 8.6.16. Let $\mathcal{D} \subseteq \mathbb{F}^{n \times m}$ and $\mathcal{S} \subseteq \mathbf{H}^p$ be convex sets, and let $\phi_1: \mathcal{D} \mapsto \mathcal{S}$ and $\phi_2: \mathcal{S} \mapsto \mathbf{H}^q$. Then, the following statements hold:

i) If ϕ_1 is convex and ϕ_2 is nondecreasing and convex, then $\phi_2 \bullet \phi_1: \mathcal{D} \mapsto \mathbf{H}^q$ is convex.

ii) If ϕ_1 is concave and ϕ_2 is nonincreasing and convex, then $\phi_2 \bullet \phi_1: \mathcal{D} \mapsto \mathbf{H}^q$ is convex.

iii) If $\mathcal{S}$ is symmetric, $\phi_2(-A) = -\phi_2(A)$ for all $A \in \mathcal{S}$, ϕ_1 is concave, and ϕ_2 is nonincreasing and concave, then $\phi_2 \bullet \phi_1: \mathcal{D} \mapsto \mathbf{H}^q$ is convex.

iv) If $\mathcal{S}$ is symmetric, $\phi_2(-A) = -\phi_2(A)$ for all $A \in \mathcal{S}$, ϕ_1 is convex, and ϕ_2 is

nondecreasing and concave, then $\phi_2 \bullet \phi_1: \mathcal{D} \mapsto \mathbf{H}^q$ is convex.

Proof. To prove *i)* and *ii)*, let $\alpha \in [0, 1]$ and $A_1, A_2 \in \mathcal{D}$. In both cases it follows that

$$\begin{aligned} \phi_2(\phi_1[\alpha A_1 + (1 - \alpha)A_2]) &\leq \phi_2[\alpha \phi_1(A_1) + (1 - \alpha)\phi_1(A_2)] \\ &\leq \alpha \phi_2[\phi_1(A_1)] + (1 - \alpha)\phi_2[\phi_1(A_2)]. \end{aligned}$$

Statements *iii)* and *iv)* follow from *i)* and *ii)*, respectively. $\square$

Proposition 8.6.17. The following functions are convex:

- i)* $\phi: \mathbf{N}^n \mapsto \mathbf{N}^n$ defined by $\phi(A) \triangleq A^r$, where $r \in [1, 2]$.
- ii)* $\phi: \mathbf{N}^n \mapsto \mathbf{N}^n$ defined by $\phi(A) \triangleq A^2$.
- iii)* $\phi: \mathbf{P}^n \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq A^{-r}$, where $r \in [0, 1]$.
- iv)* $\phi: \mathbf{P}^n \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq A^{-1}$.
- v)* $\phi: \mathbf{P}^n \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq A^{-1/2}$.
- vi)* $\phi: \mathbf{N}^n \mapsto -\mathbf{N}^n$ defined by $\phi(A) \triangleq -A^r$, where $r \in [0, 1]$.
- vii)* $\phi: \mathbf{N}^n \mapsto -\mathbf{N}^n$ defined by $\phi(A) \triangleq -A^{1/2}$.
- viii)* $\phi: \mathbf{N}^n \mapsto \mathbf{H}^m$ defined by $\phi(A) \triangleq \gamma BAB^*$, where $\gamma \in \mathbb{R}$ and $B \in \mathbb{F}^{m \times n}$.
- ix)* $\phi: \mathbf{N}^n \mapsto \mathbf{N}^m$ defined by $\phi(A) \triangleq BA^r B^*$, where $B \in \mathbb{F}^{m \times n}$ and $r \in [1, 2]$.
- x)* $\phi: \mathbf{P}^n \mapsto \mathbf{N}^m$ defined by $\phi(A) \triangleq BA^{-r} B^*$, where $B \in \mathbb{F}^{m \times n}$ and $r \in [0, 1]$.
- xi)* $\phi: \mathbf{N}^n \mapsto -\mathbf{N}^m$ defined by $\phi(A) \triangleq -BA^r B^*$, where $B \in \mathbb{F}^{m \times n}$ and $r \in [0, 1]$.
- xii)* $\phi: \mathbf{P}^n \mapsto -\mathbf{P}^m$ defined by $\phi(A) \triangleq -(BA^{-r} B^*)^{-p}$, where $B \in \mathbb{F}^{m \times n}$ has rank m and $r, p \in [0, 1]$.
- xiii)* $\phi: \mathbb{F}^{n \times m} \mapsto \mathbf{N}^n$ defined by $\phi(A) \triangleq ABA^*$, where $B \in \mathbf{N}^m$.
- xiv)* $\phi: \mathbf{P}^n \times \mathbb{F}^{m \times n} \mapsto \mathbf{N}^m$ defined by $\phi(A, B) \triangleq BA^{-1} B^*$.
- xv)* $\phi: \mathbf{P}^n \times \mathbb{F}^{m \times n} \mapsto \mathbf{N}^m$ defined by $\phi(A) \triangleq (A^{-1} + A^{-*})^{-1}$.
- xvi)* $\phi: \mathbf{N}^n \times \mathbf{N}^n \mapsto \mathbf{N}^n$ defined by $\phi(A, B) \triangleq -A(A + B)^+ B$.
- xvii)* $\phi: \mathbf{N}^{n+m} \mapsto \mathbf{N}^n$ defined by $\phi(A) \triangleq -A_{22}|A$, where $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$.
- xviii)* $\phi: \mathbf{P}^{n+m} \mapsto \mathbf{P}^n$ defined by $\phi(A) \triangleq (A_{22}|A)^{-1}$, where $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$.
- xix)* $\phi: \mathbf{H}^n \mapsto [0, \infty)$ defined by $\phi(A) \triangleq \text{tr } A^k$, where k is a nonnegative even integer.
- xx)* $\phi: \mathbf{P}^n \mapsto (0, \infty)$ defined by $\phi(A) \triangleq \text{tr } A^{-r}$, where $r > 0$.
- xxi)* $\phi: \mathbf{P}^n \mapsto (-\infty, 0)$ defined by $\phi(A) \triangleq -(\text{tr } A^{-r})^{-p}$, where $r, p \in [0, 1]$.

- xxii)* $\phi: \mathbf{N}^n \times \mathbf{N}^n \mapsto (-\infty, 0]$ defined by $\phi(A, B) \triangleq -\operatorname{tr}(A^r + B^r)^{1/r}$, where $r \in [0, 1]$.
- xxiii)* $\phi: \mathbf{N}^n \times \mathbf{N}^n \mapsto [0, \infty)$ defined by $\phi(A, B) \triangleq \operatorname{tr}(A^2 + B^2)^{1/2}$.
- xxiv)* $\phi: \mathbf{N}^n \times \mathbf{N}^m \mapsto \mathbb{R}$ defined by $\phi(A, B) \triangleq -\operatorname{tr} A^r X B^p X^*$, where $X \in \mathbb{F}^{n \times m}$, $r, p \geq 0$, and $r + p \leq 1$.
- xxv)* $\phi: \mathbf{N}^n \mapsto (-\infty, 0)$ defined by $\phi(A) \triangleq -\operatorname{tr} A^r X A^p X^*$, where $X \in \mathbb{F}^{n \times n}$, $r, p \geq 0$, and $r + p \leq 1$.
- xxvi)* $\phi: \mathbf{P}^n \times \mathbf{P}^m \times \mathbb{F}^{m \times n} \mapsto \mathbb{R}$ defined by $\phi(A, B, X) \triangleq (\operatorname{tr} A^{-p} X B^{-r} X^*)^q$, where $r, p \geq 0$, $r + p \leq 1$, and $q \geq (2 - r - p)^{-1}$.
- xxvii)* $\phi: \mathbf{P}^n \times \mathbb{F}^{n \times n} \mapsto [0, \infty)$ defined by $\phi(A, X) \triangleq \operatorname{tr} A^{-p} X A^{-r} X^*$, where $r, p \geq 0$ and $r + p \leq 1$.
- xxviii)* $\phi: \mathbf{P}^n \times \mathbb{F}^{n \times n} \mapsto [0, \infty)$ defined by $\phi(A) \triangleq \operatorname{tr} A^{-p} X A^{-r} X^*$, where $r, p \in [0, 1]$ and $X \in \mathbb{F}^{n \times n}$.
- xxix)* $\phi: \mathbf{P}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq -\operatorname{tr}([A^r, X][A^{1-r}, X])$, where $r \in (0, 1)$ and $X \in \mathbf{H}^n$.
- xxx)* $\phi: \mathbf{P}^n \mapsto \mathbf{H}^n$ defined by $\phi(A) \triangleq -\log A$.
- xxxi)* $\phi: \mathbf{P}^n \mapsto \mathbf{H}^m$ defined by $\phi(A) \triangleq A \log A$.
- xxxii)* $\phi: \mathbf{N}^n \setminus \{0\} \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq -\log \operatorname{tr} A^r$, where $r \in [0, 1]$.
- xxxiii)* $\phi: \mathbf{P}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq \log \operatorname{tr} A^{-1}$.
- xxxiv)* $\phi: \mathbf{P}^n \times \mathbf{P}^n \mapsto (0, \infty)$ defined by $\phi(A, B) \triangleq \operatorname{tr}[A(\log A - \log B)]$.
- xxxv)* $\phi: \mathbf{P}^n \times \mathbf{P}^n \mapsto [0, \infty)$ defined by $\phi(A, B) \triangleq -e^{[1/(2n)]\operatorname{tr}(\log A + \log B)}$.
- xxxvi)* $\phi: \mathbf{N}^n \mapsto (-\infty, 0]$ defined by $\phi(A) \triangleq -(\det A)^{1/n}$.
- xxxvii)* $\phi: \mathbf{P}^n \mapsto (0, \infty)$ defined by $\phi(A) \triangleq \log \det B A^{-1} B^*$, where $B \in \mathbb{F}^{m \times n}$ and $\operatorname{rank} B = m$.
- xxxviii)* $\phi: \mathbf{P}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq -\log \det A$.
- xxxix)* $\phi: \mathbf{P}^n \mapsto (0, \infty)$ defined by $\phi(A) \triangleq \det A^{-1}$.
- xl)* $\phi: \mathbf{P}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq \log(\det A_k / \det A)$, where $k \in \{1, \dots, n-1\}$ and A_k is the leading $k \times k$ principal submatrix of A .
- xli)* $\phi: \mathbf{P}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq -\det A / \det A_{[n;n]}$.
- xlii)* $\phi: \mathbf{N}^n \times \mathbf{N}^m \mapsto -\mathbf{N}^{nm}$ defined by $\phi(A, B) \triangleq -A^{r_1} \otimes B^{r_2}$, where $r_1, r_2 \in [0, 1]$ satisfy $r_1 + r_2 \leq 1$.
- xliii)* $\phi: \mathbf{P}^n \times \mathbf{N}^m \mapsto \mathbf{N}^{nm}$ defined by $\phi(A, B) \triangleq A^{-r} \otimes B^{1+r}$, where $r \in [0, 1]$.
- xliv)* $\phi: \mathbf{N}^n \times \mathbf{N}^n \mapsto -\mathbf{N}^n$ defined by $\phi(A, B) \triangleq -A^{r_1} \circ B^{r_2}$, where $r_1, r_2 \in [0, 1]$ satisfy $r_1 + r_2 \leq 1$.
- xl v)* $\phi: \mathbf{H}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq \sum_{i=1}^k \lambda_i(A)$, where $k \in \{1, \dots, n\}$.

$xlvi)$ $\phi: \mathbf{H}^n \mapsto \mathbb{R}$ defined by $\phi(A) \triangleq -\sum_{i=k}^n \lambda_i(A)$, where $k \in \{1, \dots, n\}$.

Proof. Statements $i)$ and $iii)$ are proved in [43] and [197, p. 123].

Let $\alpha \in [0, 1]$ for the remainder of the proof.

To prove $ii)$ directly, let $A_1, A_2 \in \mathbf{H}^n$. Since

$$\alpha(1 - \alpha) = (\alpha - \alpha^2)^{1/2} [(1 - \alpha) - (1 - \alpha)^2]^{1/2},$$

it follows that

$$\begin{aligned} 0 &\leq [(\alpha - \alpha^2)^{1/2} A_1 - [(1 - \alpha) - (1 - \alpha)^2]^{1/2} A_2]^2 \\ &= (\alpha - \alpha^2) A_1^2 + [(1 - \alpha) - (1 - \alpha)^2] A_2^2 - \alpha(1 - \alpha)(A_1 A_2 + A_2 A_1). \end{aligned}$$

Hence,

$$[\alpha A_1 + (1 - \alpha) A_2]^2 \leq \alpha A_1^2 + (1 - \alpha) A_2^2,$$

which shows that $\phi(A) = A^2$ is convex.

To prove $iv)$ directly, let $A_1, A_2 \in \mathbf{P}^n$. Then, $\begin{bmatrix} A_1^{-1} & I \\ I & A_1 \end{bmatrix}$ and $\begin{bmatrix} A_2^{-1} & I \\ I & A_2 \end{bmatrix}$ are positive semidefinite, and thus

$$\begin{aligned} \alpha \begin{bmatrix} A_1^{-1} & I \\ I & A_1 \end{bmatrix} + (1 - \alpha) \begin{bmatrix} A_2^{-1} & I \\ I & A_2 \end{bmatrix} \\ = \begin{bmatrix} \alpha A_1^{-1} + (1 - \alpha) A_2^{-1} & I \\ I & \alpha A_1 + (1 - \alpha) A_2 \end{bmatrix} \end{aligned}$$

is positive semidefinite. It now follows from Proposition 8.2.4 that $[\alpha A_1 + (1 - \alpha) A_2]^{-1} \leq \alpha A_1^{-1} + (1 - \alpha) A_2^{-1}$, which shows that $\phi(A) = A^{-1}$ is convex.

To prove $v)$ directly, note that $\phi(A) = A^{-1/2} = \phi_2[\phi_1(A)]$, where $\phi_1(A) \triangleq A^{1/2}$ and $\phi_2(B) \triangleq B^{-1}$. It follows from $vii)$ that ϕ_1 is concave, while it follows from $iv)$ that ϕ_2 is convex. Furthermore, $x)$ of Proposition 8.6.13 implies that ϕ_2 is nonincreasing. It thus follows from $ii)$ of Lemma 8.6.16 that $\phi(A) = A^{-1/2}$ is convex.

To prove $vi)$, let $A \in \mathbf{P}^n$ and note that $\phi(A) = -A^r = \phi_2[\phi_1(A)]$, where $\phi_1(A) \triangleq A^{-r}$ and $\phi_2(B) \triangleq -B^{-1}$. It follows from $iii)$ that ϕ_1 is convex, while it follows from $iv)$ that ϕ_2 is concave. Furthermore, $x)$ of Proposition 8.6.13 implies that ϕ_2 is nondecreasing. It thus follows from $iv)$ of Lemma 8.6.16 that $\phi(A) = A^r$ is convex on $\mathbf{P}^n$. Continuity implies that $\phi(A) = A^r$ is convex on $\mathbf{N}^n$.

To prove $vii)$ directly, let $A_1, A_2 \in \mathbf{N}^n$. Then,

$$0 \leq \alpha(1 - \alpha) (A_1^{1/2} - A_2^{1/2})^2,$$

which is equivalent to

$$[\alpha A_1^{1/2} + (1 - \alpha) A_2^{1/2}]^2 \leq \alpha A_1 + (1 - \alpha) A_2.$$

Using *viii)* of Proposition 8.6.13 yields

$$\alpha A_1^{1/2} + (1 - \alpha) A_2^{1/2} \leq [\alpha A_1 + (1 - \alpha) A_2]^{1/2}.$$

Finally, multiplying by -1 shows that $\phi(A) = -A^{1/2}$ is convex.

The proof of *viii)* is immediate. Statements *ix)*, *x)*, and *xi)* follow from *i)*, *iii)*, and *vi)*, respectively.

To prove *xii)*, note that $\phi(A) = -(BA^{-r}B^*)^{-p} = \phi_2[\phi_1(A)]$, where $\phi_1(A) = -BA^{-r}B^*$ and $\phi_2(C) = C^{-p}$. Statement *x)* implies that ϕ_1 is concave, while *iii)* implies that ϕ_2 is convex. Furthermore, *ix)* of Proposition 8.6.13 implies that ϕ_2 is nonincreasing. It thus follows from *ii)* of Lemma 8.6.16 that $\phi(A) = -(BA^{-r}B^*)^{-p}$ is convex.

To prove *xiii)*, let $A_1, A_2 \in \mathbb{F}^{n \times m}$, and let $B \in \mathbf{N}^m$. Then,

$$\begin{aligned} 0 &\leq \alpha(1 - \alpha)(A_1 - A_2)B(A_1 - A_2)^* \\ &= \alpha A_1 B A_1^* + (1 - \alpha) A_2 B A_2^* - [\alpha A_1 + (1 - \alpha) A_2] B [\alpha A_1 + (1 - \alpha) A_2]^*. \end{aligned}$$

Thus,

$$[\alpha A_1 + (1 - \alpha) A_2] B [\alpha A_1 + (1 - \alpha) A_2]^* \leq \alpha A_1 B A_1^* + (1 - \alpha) A_2 B A_2^*,$$

which shows that $\phi(A) = ABA^*$ is convex.

To prove *xiv)*, let $A_1, A_2 \in \mathbf{P}^n$ and $B_1, B_2 \in \mathbb{F}^{m \times n}$. Then, it follows from Proposition 8.2.4 that $\begin{bmatrix} B_1 A_1^{-1} B_1^* & B_1 \\ B_1^* & A_1 \end{bmatrix}$ and $\begin{bmatrix} B_2 A_2^{-1} B_2^* & B_2 \\ B_2^* & A_2 \end{bmatrix}$ are positive semidefinite, and thus

$$\begin{aligned} &\alpha \begin{bmatrix} B_1 A_1^{-1} B_1^* & B_1 \\ B_1^* & A_1 \end{bmatrix} + (1 - \alpha) \begin{bmatrix} B_2 A_2^{-1} B_2^* & B_2 \\ B_2^* & A_2 \end{bmatrix} \\ &= \begin{bmatrix} \alpha B_1 A_1^{-1} B_1^* + (1 - \alpha) B_2 A_2^{-1} B_2^* & \alpha B_1 + (1 - \alpha) B_2 \\ \alpha B_1^* + (1 - \alpha) B_2^* & \alpha A_1 + (1 - \alpha) A_2 \end{bmatrix} \end{aligned}$$

is positive semidefinite. It thus follows from Proposition 8.2.4 that

$$\begin{aligned} &[\alpha B_1 + (1 - \alpha) B_2][\alpha A_1 + (1 - \alpha) A_2]^{-1}[\alpha B_1 + (1 - \alpha) B_2]^* \\ &\leq \alpha B_1 A_1^{-1} B_1^* + (1 - \alpha) B_2 A_2^{-1} B_2^*, \end{aligned}$$

which shows that $\phi(A, B) = BA^{-1}B^*$ is convex.

Result *xv)* is given in [978].

Result *xvi)* follows from Fact 8.20.18.

To prove *xvii)*, let $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbf{P}^{n+m}$ and $B \triangleq \begin{bmatrix} B_{11} & B_{12} \\ B_{12} & B_{22} \end{bmatrix} \in \mathbf{P}^{n+m}$. Then, it follows from *xiv)* with A_1, B_1, A_2, B_2 replaced by $A_{22}, A_{12}, B_{22}, B_{12}$, respectively,

that

$$\begin{aligned} & [\alpha A_{12} + (1 - \alpha)B_{12}][\alpha A_{22} + (1 - \alpha)B_{22}]^{-1}[\alpha A_{12} + (1 - \alpha)B_{12}]^* \\ & \leq \alpha A_{12}A_{22}^{-1}A_{12}^* + (1 - \alpha)B_{12}B_{22}^{-1}B_{12}^*. \end{aligned}$$

Hence,

$$\begin{aligned} & -[\alpha A_{22} + (1 - \alpha)B_{22}][\alpha A + (1 - \alpha)B] \\ & = [\alpha A_{12} + (1 - \alpha)B_{12}][\alpha A_{22} + (1 - \alpha)B_{22}]^{-1}[\alpha A_{12} + (1 - \alpha)B_{12}]^* \\ & \quad - [\alpha A_{11} + (1 - \alpha)B_{11}] \\ & \leq \alpha(A_{12}A_{22}^{-1}A_{12}^* - A_{11}) + (1 - \alpha)(B_{12}B_{22}^{-1}B_{12}^* - B_{11}) \\ & = \alpha(-A_{22}|A) + (1 - \alpha)(-B_{22}|B), \end{aligned}$$

which shows that $\phi(A) \triangleq -A_{22}|A$ is convex. By continuity, the result holds for $A \in \mathbf{N}^{n+m}$.

To prove *xviii*), note that $\phi(A) = (A_{22}|A)^{-1} = \phi_2[\phi_1(A)]$, where $\phi_1(A) = A_{22}|A$ and $\phi_2(B) = B^{-1}$. It follows from *xv*) that ϕ_1 is concave, while it follows from *iv*) that ϕ_2 is convex. Furthermore, *x*) of Proposition 8.6.13 implies that ϕ_2 is nonincreasing. It thus follows from Lemma 8.6.16 that $\phi(A) \triangleq (A_{22}|A)^{-1}$ is convex.

Result *xix*) is given in [239, p. 106].

Result *xx*) is given in by Theorem 9 of [905].

To prove *xxi*), note that $\phi(A) = -(\text{tr } A^{-r})^{-p} = \phi_2[\phi_1(A)]$, where $\phi_1(A) = \text{tr } A^{-r}$ and $\phi_2(B) = -B^{-p}$. Statement *iii*) implies that ϕ_1 is convex and that ϕ_2 is concave. Furthermore, *ix*) of Proposition 8.6.13 implies that ϕ_2 is nondecreasing. It thus follows from *iv*) of Lemma 8.6.16 that $\phi(A) = -(\text{tr } A^{-r})^{-p}$ is convex.

Results *xxii*) and *xxiii*) are proved in [286].

Results *xxiv*)–*xxviii*) are given by Corollary 1.1, Theorem 1, Corollary 2.1, Theorem 2, and Theorem 8, respectively, of [286]. A proof of *xxiv*) in the case $p = 1 - r$ is given in [197, p. 273].

Result *xxix*) is proved in [197, p. 274] and [286].

Result *xxx*) is given in [201, p. 113].

Result *xxxi*) is given in [197, p. 123], [201, p. 113], and [529].

To prove *xxxi*), note that $\phi(A) = -\log \text{tr } A^r = \phi_2[\phi_1(A)]$, where $\phi_1(A) = \text{tr } A^r$ and $\phi_2(x) = -\log x$. Statement *vi*) implies that ϕ_1 is concave. Furthermore, ϕ_2 is convex and nonincreasing. It thus follows from *ii*) of Lemma 8.6.16 that $\phi(A) = -\log \text{tr } A^r$ is convex.

Result *xxxii*) is given in [1024].

Result *xxxiv*) is given in [197, p. 275].

Result *xxv*) is given in [54].

To prove *xxvi*), let $A_1, A_2 \in \mathbf{N}^n$. From Corollary 8.4.15 it follows that $(\det A_1)^{1/n} + (\det A_2)^{1/n} \leq [\det(A_1 + A_2)]^{1/n}$. Replacing A_1 and A_2 by αA_1 and $(1 - \alpha)A_2$, respectively, and multiplying by -1 shows that $\phi(A) = -(\det A)^{1/n}$ is convex.

Result *xxvii*) is proved in [1024].

Result *xxviii*) is a special case of result *xxvii*). This result is due to Fan. See [352] or [353, p. 679]. To prove *xxviii*), note that $\phi(A) = -n \log[(\det A)^{1/n}] = \phi_2[\phi_1(A)]$, where $\phi_1(A) = (\det A)^{1/n}$ and $\phi_2(x) = -n \log x$. It follows from *xix*) that ϕ_1 is concave. Since ϕ_2 is nonincreasing and convex, it follows from *ii*) of Lemma 8.6.16 that $\phi(A) = -\log \det A$ is convex.

To prove *xxix*), note that $\phi(A) = \det A^{-1} = \phi_2[\phi_1(A)]$, where $\phi_1(A) = \log \det A^{-1}$ and $\phi_2(x) = e^x$. It follows from *xx*) that ϕ_1 is convex. Since ϕ_2 is nondecreasing and convex, it follows from *i*) of Lemma 8.6.16 that $\phi(A) = \det A^{-1}$ is convex.

Results *xl*) and *xli*) are given in [352] and [353, pp. 684, 685].

Next, *xlii*) is given in [197, p. 273], [201, p. 114], and [1485, p. 9]. Statement *xliii*) is given in [201, p. 114]. Statement *xliv*) is given in [1485, p. 9].

Finally, *xlvi*) is given in [971, p. 478]. Statement *xlvi*) follows immediately from *xlvi*). $\square$

The following result is a corollary of *xv*) of Proposition 8.6.17 for the case $\alpha = 1/2$. Versions of this result appear in [290, 658, 896, 922] and [1098, p. 152].

Corollary 8.6.18. Let $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbb{F}^{n+m}$ and $B \triangleq \begin{bmatrix} B_{11} & B_{12} \\ B_{12}^* & B_{22} \end{bmatrix} \in \mathbb{F}^{n+m}$, and assume that A and B are positive semidefinite. Then,

$$A_{11}|A + B_{11}|B \leq (A_{11} + B_{11})|(A + B).$$

The following corollary of *xlvi*) and *xlvi*) of Proposition 8.6.17 gives a strong majorization condition for the eigenvalues of a pair of Hermitian matrices.

Corollary 8.6.19. Let $A, B \in \mathbf{H}^n$. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \lambda_i(A) + \sum_{i=1}^k \lambda_{n-k+i}(B) \leq \sum_{i=1}^k \lambda_i(A + B) \leq \sum_{i=1}^k [\lambda_i(A) + \lambda_i(B)] \quad (8.6.12)$$

with equality for $k = n$. Furthermore, for all $k = 1, \dots, n$,

$$\sum_{i=k}^n [\lambda_i(A) + \lambda_i(B)] \leq \sum_{i=k}^n \lambda_i(A+B) \quad (8.6.13)$$

with equality for $k = 1$.

Proof. The lower bound in (8.6.12) is given in [1177, p. 116]. See also [197, p. 69], [320], [711, p. 201], or [971, p. 478]. $\square$

Equality in Corollary 8.6.19 is discussed in [320].

8.7 Facts on Range and Rank

Fact 8.7.1. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, there exists $\alpha > 0$ such that $A \leq \alpha B$ if and only if $\mathcal{R}(A) \subseteq \mathcal{R}(B)$. In this case, $\text{rank } A \leq \text{rank } B$. (Proof: Use Theorem 8.6.2 and Corollary 8.6.11.)

Fact 8.7.2. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\mathcal{R}(A) + \mathcal{R}(B) = \mathcal{R}[(AA^* + BB^*)^{1/2}].$$

(Proof: The result follows from Fact 2.11.1 and Theorem 2.4.3.) (Remark: See [40].)

Fact 8.7.3. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite and B is either positive semidefinite or skew Hermitian. Then, the following identities hold:

$$i) \quad \mathcal{R}(A+B) = \mathcal{R}(A) + \mathcal{R}(B).$$

$$ii) \quad \mathcal{N}(A+B) = \mathcal{N}(A) \cap \mathcal{N}(B).$$

(Proof: Use $[(\mathcal{N}(A) \cap \mathcal{N}(B))^\perp]^\perp = \mathcal{R}(A) + \mathcal{R}(B)$.)

Fact 8.7.4. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, $(A+B)(A+B)^+$ is the projector onto $\mathcal{R}(A) + \mathcal{R}(B) = \text{span}[\mathcal{R}(A) \cup \mathcal{R}(B)]$. (Proof: Use Fact 2.9.13 and Fact 8.7.3.) (Remark: See Fact 6.4.45.)

Fact 8.7.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that $A + A^* \geq 0$. Then, the following identities hold:

$$i) \quad \mathcal{N}(A) = \mathcal{N}(A + A^*) \cap \mathcal{N}(A - A^*).$$

$$ii) \quad \mathcal{R}(A) = \mathcal{R}(A + A^*) + \mathcal{R}(A - A^*).$$

$$iii) \quad \text{rank } A = \text{rank} \begin{bmatrix} A + A^* & A - A^* \end{bmatrix}.$$

Fact 8.7.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\text{rank}(A+B) = \text{rank} \begin{bmatrix} A & B \\ A & B \end{bmatrix} = \text{rank} \begin{bmatrix} A \\ B \end{bmatrix}$$

and

$$\operatorname{rank} \begin{bmatrix} A & B \\ 0 & A \end{bmatrix} = \operatorname{rank} A + \operatorname{rank}(A + B).$$

(Proof: Using Fact 8.7.3,

$$\begin{aligned} \mathcal{R} \left(\begin{bmatrix} A & B \end{bmatrix} \right) &= \mathcal{R} \left(\begin{bmatrix} A & B \end{bmatrix} \begin{bmatrix} A \\ B \end{bmatrix} \right) = \mathcal{R}(A^2 + B^2) = \mathcal{R}(A^2) + \mathcal{R}(B^2) \\ &= \mathcal{R}(A) + \mathcal{R}(B) = \mathcal{R}(A + B). \end{aligned}$$

Alternatively, it follows from Fact 6.5.6 that

$$\begin{aligned} \operatorname{rank} \begin{bmatrix} A & B \end{bmatrix} &= \operatorname{rank} \begin{bmatrix} A + B & B \end{bmatrix} \\ &= \operatorname{rank}(A + B) + \operatorname{rank}[B - (A + B)(A + B)^+ B]. \end{aligned}$$

Next, note that

$$\begin{aligned} \operatorname{rank}[B - (A + B)(A + B)^+ B] &= \operatorname{rank} \left(B^{1/2} [I - (A + B)(A + B)^+] B^{1/2} \right) \\ &\leq \operatorname{rank} \left(B^{1/2} [I - BB^+] B^{1/2} \right) = 0. \end{aligned}$$

For the second result use Theorem 8.3.4 to simultaneously diagonalize A and B .)

Fact 8.7.7. Let $A \in \mathbb{F}^{n \times n}$, and let $\mathcal{S} \subseteq \{1, \dots, n\}$. If A is either positive semidefinite or an irreducible, singular M-matrix, then the following statements hold:

i) If $\alpha \subset \{1, \dots, n\}$, then

$$\operatorname{rank} A \leq \operatorname{rank} A_{(\alpha)} + \operatorname{rank} A_{(\alpha^c)}.$$

ii) If $\alpha, \beta \subseteq \{1, \dots, n\}$, then

$$\operatorname{rank} A_{(\alpha \cup \beta)} \leq \operatorname{rank} A_{(\alpha)} + \operatorname{rank} A_{(\beta)} - \operatorname{rank} A_{(\alpha \cap \beta)}.$$

iii) If $1 \leq k \leq n - 1$, then

$$k \sum_{\{\alpha: \operatorname{card}(\alpha)=k+1\}} \det A_{(\alpha)} \leq (n - k) \sum_{\{\alpha: \operatorname{card}(\alpha)=k\}} \det A_{(\alpha)}.$$

If, in addition, A is either positive definite, a nonsingular M-matrix, or totally positive, then all three inclusions hold as identities. (Proof: See [938].) (Remark: See Fact 8.13.36.) (Remark: Totally positive means that every subdeterminant of A is positive. See Fact 11.18.23.)

8.8 Facts on Structured Positive-Semidefinite Matrices

Fact 8.8.1. Let $\phi: \mathbb{R} \mapsto \mathbb{C}$, and assume that, for all $x_1, \dots, x_n \in \mathbb{R}$, the matrix $A \in \mathbb{C}^{n \times n}$, where $A_{(i,j)} \triangleq \phi(x_i - x_j)$, is positive semidefinite. (The function ϕ is *positive semidefinite*.) Then, the following statements hold:

i) For all $x_1, x_2 \in \mathbb{R}$, it follows that

$$|\phi(x_1) - \phi(x_2)|^2 \leq 2\phi(0)\operatorname{Re}[\phi(0) - \phi(x_1 - x_2)].$$

- ii) The function $\psi: \mathbb{R} \mapsto \mathbb{C}$, where, for all $x \in \mathbb{R}$, $\psi(x) \triangleq \overline{\phi(x)}$, is positive semidefinite.
- iii) For all $\alpha \in \mathbb{R}$, the function $\psi: \mathbb{R} \mapsto \mathbb{C}$, where, for all $x \in \mathbb{R}$, $\psi(x) \triangleq \phi(\alpha x)$, is positive semidefinite.
- iv) The function $\psi: \mathbb{R} \mapsto \mathbb{C}$, where, for all $x \in \mathbb{R}$, $\psi(x) \triangleq |\phi(x)|$, is positive semidefinite.
- v) The function $\psi: \mathbb{R} \mapsto \mathbb{C}$, where, for all $x \in \mathbb{R}$, $\psi(x) \triangleq \operatorname{Re} \phi(x)$, is positive semidefinite.
- vi) If $\phi_1: \mathbb{R} \mapsto \mathbb{C}$ and $\phi_2: \mathbb{R} \mapsto \mathbb{C}$ are positive semidefinite, then $\phi_3: \mathbb{R} \mapsto \mathbb{C}$, where, for all $x \in \mathbb{R}$, $\phi_3(x) \triangleq \phi_1(x)\phi_2(x)$, is positive semidefinite.
- vii) If $\phi_1: \mathbb{R} \mapsto \mathbb{C}$ and $\phi_2: \mathbb{R} \mapsto \mathbb{C}$ are positive semidefinite and α_1, α_2 are positive numbers, then $\phi_3: \mathbb{R} \mapsto \mathbb{C}$, where, for all $x \in \mathbb{R}$, $\phi_3(x) \triangleq \alpha_1\phi_1(x) + \alpha_2\phi_2(x)$, is positive semidefinite.
- viii) Let $\psi: \mathbb{R} \mapsto \mathbb{C}$, for all $x, y \in \mathbb{R}$, define $K: \mathbb{R} \times \mathbb{R} \mapsto \mathbb{C}$ by $K(x, y) \triangleq \phi(x - y)$, and assume that K is bounded and continuous. Then, ψ is positive semidefinite if and only if, for every continuous integrable function $f: \mathbb{R} \mapsto \mathbb{C}$, it follows that

$$\int_{\mathbb{R}^2} K(x, y) f(x) \overline{f(y)} dx dy \geq 0.$$

(Proof: See [201, pp. 141–144].) (Remark: The function K is a *kernel function* associated with a reproducing kernel space. See [546] for extensions to vector arguments. For applications, see [1175] and Fact 8.8.2.)

Fact 8.8.2. Let $a_1, \dots, a_n \in \mathbb{R}$, and define $A \in \mathbb{C}^{n \times n}$ by either of the following expressions:

- i) $A_{(i,j)} \triangleq \frac{1}{1+j(a_i-a_j)}.$
- ii) $A_{(i,j)} \triangleq \frac{1}{1-j(a_i-a_j)}.$
- iii) $A_{(i,j)} \triangleq \frac{1}{1+(a_i-a_j)^2}.$
- iv) $A_{(i,j)} \triangleq \frac{1}{1+|a_i-a_j|}.$
- v) $A_{(i,j)} \triangleq e^{j(a_i-a_j)}.$
- vi) $A_{(i,j)} \triangleq \cos(a_i - a_j).$
- vii) $A_{(i,j)} \triangleq \frac{\sin[(a_i-a_j)]}{a_i-a_j}.$
- viii) $A_{(i,j)} \triangleq \frac{a_i-a_j}{\sinh[(a_i-a_j)]}.$
- ix) $A_{(i,j)} \triangleq \frac{\sinh p(a_i-a_j)}{\sinh(a_i-a_j)},$ where $p \in (0, 1).$
- x) $A_{(i,j)} \triangleq \frac{\tanh[(a_i-a_j)]}{a_i-a_j}.$

$$xi) \quad A_{(i,j)} \triangleq \frac{\sinh[(a_i - a_j)]}{(a_i - a_j)[\cosh(a_i - a_j) + p]}, \text{ where } p \in (-1, 1].$$

$$xii) \quad A_{(i,j)} \triangleq \frac{1}{\cosh(a_i - a_j) + p}, \text{ where } p \in (-1, 1].$$

$$xiii) \quad A_{(i,j)} \triangleq \frac{\cosh p(a_i - a_j)}{\cosh(a_i - a_j)}, \text{ where } p \in [-1, 1].$$

$$xiv) \quad A_{(i,j)} \triangleq e^{-(a_i - a_j)^2}.$$

$$xv) \quad A_{(i,j)} \triangleq e^{-|a_i - a_j|^p}, \text{ where } p \in [0, 2].$$

$$xvi) \quad A_{(i,j)} \triangleq \frac{1}{1 + |a_i - a_j|}.$$

$$xvii) \quad A_{(i,j)} \triangleq \frac{1 + p(a_i - a_j)^2}{1 + q(a_i - a_j)^2}, \text{ where } 0 \leq p \leq q.$$

$$xviii) \quad A_{(i,j)} \triangleq \text{tr } e^{B + J(a_i - a_j)C}, \text{ where } B, C \in \mathbb{C}^{n \times n} \text{ are Hermitian and commute.}$$

Then, A is positive semidefinite. Finally, if, α is a nonnegative number and A is defined by either $ix)$, $x)$, $xii)$, $xiii)$, $xvi)$, or $xvii)$, then $A^{\circ\alpha}$ is positive semidefinite. (Proof: See [201, pp. 141–144, 153, 177, 188], [216], [422, p. 90], and [709, pp. 400, 401, 456, 457, 462, 463].) (Remark: In each case, A is associated with a positive-semidefinite function. See Fact 8.8.1.) (Remark: $xv)$ is related to the Bessis-Moussa-Villani conjecture. See Fact 8.12.30 and Fact 8.12.31.) (Problem: In each case, determine rank A and determine when A is positive definite.)

Fact 8.8.3. Let $a_1, \dots, a_n$ be positive numbers, and define $A \in \mathbb{R}^{n \times n}$ by either of the following expressions:

$$i) \quad A_{(i,j)} \triangleq \min\{a_i, a_j\}.$$

$$ii) \quad A_{(i,j)} \triangleq \frac{1}{\max\{a_i, a_j\}}.$$

$$iii) \quad A_{(i,j)} \triangleq \frac{a_i}{a_j}, \text{ where } a_1 \leq \dots \leq a_n.$$

$$iv) \quad A_{(i,j)} \triangleq \frac{a_i^p - a_j^p}{a_i - a_j}, \text{ where } p \in [0, 1].$$

$$v) \quad A_{(i,j)} \triangleq \frac{a_i^p + a_j^p}{a_i + a_j}, \text{ where } p \in [-1, 1].$$

$$vi) \quad A_{(i,j)} \triangleq \frac{\log a_i - \log a_j}{a_i - a_j}.$$

Then, A is positive semidefinite. If, in addition, α is a positive number, then $A^{\circ\alpha}$ is positive semidefinite. (Proof: See [199], [201, p. 153, 178, 189], and [422, p. 90].) (Remark: The matrix A in $iii)$ is the Schur product of the matrices defined in $i)$ and $ii)$.)

Fact 8.8.4. Let $a_1 < \dots < a_n$ be positive numbers, and define $A \in \mathbb{R}^{n \times n}$ by $A_{(i,j)} \triangleq \min\{a_i, a_j\}$. Then, A is positive definite,

$$\det A = \prod_{i=1}^n (a_i - a_{i-1}),$$

and, for all $x \in \mathbb{R}^n$,

$$x^T A^{-1} x = \sum_{i=1}^n \frac{[x_{(i)} - x_{(i-1)}]^2}{a_i - a_{i-1}},$$

where $a_0 \triangleq 0$ and $x_0 \triangleq 0$. (Remark: The matrix A is a covariance matrix arising in the theory of Brownian motion. See [673, p. 132] and [1454, p. 50].)

Fact 8.8.5. Define $A \in \mathbb{R}^{n \times n}$ by either of the following expressions:

- i) $A_{(i,j)} \triangleq \binom{i+j}{i}$.
- ii) $A_{(i,j)} \triangleq (i+j)!$.
- iii) $A_{(i,j)} \triangleq \min\{i, j\}$.
- iv) $A_{(i,j)} \triangleq \gcd\{i, j\}$.
- v) $A_{(i,j)} \triangleq \frac{i}{j}$.

Then, A is positive semidefinite. If, in addition, α is a nonnegative number, then $A^{\circ \alpha}$ is positive semidefinite. (Remark: Fact 8.21.2 guarantees the weaker result that $A^{\circ \alpha}$ is positive semidefinite for all $\alpha \in [0, n-2]$.) (Remark: i) is the *Pascal matrix*. See [5, 199, 448]. The fact that A is positive semidefinite follows from the identity

$$\binom{i+j}{i} = \sum_{k=0}^{\min\{i,j\}} \binom{i}{k} \binom{j}{k}.$$

(Remark: The matrix defined in v), which is a special case of iii) of Fact 8.8.3, is the *Lehmer matrix*.) (Remark: The determinant of A defined in iv) can be expressed in terms of the *Euler totient function*. See [66, 253].)

Fact 8.8.6. Let $a_1, \dots, a_n \geq 0$ and $p \in \mathbb{R}$, assume that either $a_1, \dots, a_n$ are positive or p is positive, and, for all $i, j = 1, \dots, n$, define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq (a_i a_j)^p.$$

Then, A is positive semidefinite. (Proof: Let $a \triangleq [a_1 \ \cdots \ a_n]^T$ and $A \triangleq a^{\circ p} a^{\circ p T}$.)

Fact 8.8.7. Let $a_1, \dots, a_n > 0$, let $\alpha > 0$, and, for all $i, j = 1, \dots, n$, define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{1}{(a_i + a_j)^\alpha}.$$

Then, A is positive semidefinite. (Proof: See [199], [201, pp. 24, 25], or [1092].) (Remark: See Fact 5.11.12.) (Remark: For $\alpha = 1$, A is a Cauchy matrix. See Fact 3.20.14.)

Fact 8.8.8. Let $a_1, \dots, a_n > 0$, let $r \in [-1, 1]$, and, for all $i, j = 1, \dots, n$, define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{a_i^r + a_j^r}{a_i + a_j}.$$

Then, A is positive semidefinite. (Proof: See [1485, p. 74].)

Fact 8.8.9. Let $a_1, \dots, a_n > 0$, let $q > 0$, let $p \in [-q, q]$, and, for all $i, j = 1, \dots, n$, define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{a_i^p + a_j^p}{a_i^q + a_j^q}.$$

Then, A is positive semidefinite. (Proof: Let $r = p/q$ and $b_i = a_i^q$. Then, $A_{(i,j)} = (b_i^r + b_j^r)/(b_i + b_j)$. Now, use Fact 8.8.8. See [979] for the case $q \geq p \geq 0$.) (Remark: The case $q = 1$ and $p = 0$ yields a Cauchy matrix. In the case $n = 2$, $A \geq 0$ yields Fact 1.10.33.) (Problem: When is A positive definite?)

Fact 8.8.10. Let $a_1, \dots, a_n > 0$, let $p \in (-2, 2]$, and define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{1}{a_i^2 + pa_i a_j + a_j^2}.$$

Then, A is positive semidefinite. (Proof: See [204].)

Fact 8.8.11. Let $a_1, \dots, a_n > 0$, let $p \in (-1, \infty)$, and define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{1}{a_i^3 + p(a_i^2 a_j + a_i a_j^2) + a_j^3}.$$

Then, A is positive semidefinite. (Proof: See [204].)

Fact 8.8.12. Let $a_1, \dots, a_n > 0$, $p \in [-1, 1]$, $q \in (-2, 2]$, and, for all $i, j = 1, \dots, n$, define $A \in \mathbb{R}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{a_i^p + a_j^p}{a_i^2 + qa_i a_j + a_j^2}.$$

Then, A is positive semidefinite. (Proof: See [1482] or [1485, p. 76].)

Fact 8.8.13. Let $A \in \mathbb{R}^{n \times n}$, assume that A is positive semidefinite, assume that $A_{(i,i)} > 0$ for all $i = 1, \dots, n$, and define $B \in \mathbb{R}^{n \times n}$ by

$$B_{(i,j)} \triangleq \frac{A_{(i,j)}}{\mu_\alpha(A_{(i,i)}, A_{(j,j)})},$$

where, for positive scalars α, x, y ,

$$\mu_\alpha(x, y) \triangleq \left[\frac{1}{2}(x^\alpha + y^\alpha) \right]^{1/\alpha}.$$

Then, B is positive semidefinite. If, in addition, A is positive definite, then B is positive definite. In particular, letting $\alpha \downarrow 0$, $\alpha = 1$, and $\alpha \rightarrow \infty$, respectively, the matrices $C, D, E \in \mathbb{R}^{n \times n}$ defined by

$$\begin{aligned} C_{(i,j)} &\triangleq \frac{A_{(i,j)}}{\sqrt{A_{(i,i)} A_{(j,j)}}}, \\ D_{(i,j)} &\triangleq \frac{2A_{(i,j)}}{A_{(i,i)} + A_{(j,j)}}, \\ E_{(i,j)} &\triangleq \frac{A_{(i,j)}}{\max\{A_{(i,i)}, A_{(j,j)}\}} \end{aligned}$$

are positive semidefinite. Finally, if A is positive definite, then C , D , and E are positive definite. (Proof: See [1151].) (Remark: The assumption that all of the diagonal entries of A are positive can be weakened. See [1151].) (Remark: See Fact 1.10.34.) (Problem: Extend this result to Hermitian matrices.)

Fact 8.8.14. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian, $A_{(i,i)} > 0$ for all $i = 1, \dots, n$, and, for all $i, j = 1, \dots, n$,

$$|A_{(i,j)}| < \frac{1}{n-1} \sqrt{A_{(i,i)} A_{(j,j)}}.$$

Then, A is positive definite. (Proof: Note that

$$x^* A x = \sum_{i=1}^{n-1} \sum_{j=i+1}^n \begin{bmatrix} x_{(i)} \\ x_{(j)} \end{bmatrix}^* \begin{bmatrix} \frac{1}{n-1} A_{(i,i)} & A_{(i,j)} \\ \overline{A_{(i,j)}} & \frac{1}{n-1} A_{(j,j)} \end{bmatrix} \begin{bmatrix} x_{(i)} \\ x_{(j)} \end{bmatrix}.)$$

(Remark: This result is due to Roup.)

Fact 8.8.15. Let $\alpha, \beta, \gamma \in [0, \pi]$, and define $A \in \mathbb{R}^{3 \times 3}$ by

$$A = \begin{bmatrix} 1 & \cos \alpha & \cos \gamma \\ \cos \alpha & 1 & \cos \beta \\ \cos \gamma & \cos \beta & 1 \end{bmatrix}.$$

Then, A is positive semidefinite if and only if the following conditions are satisfied:

- i) $\alpha \leq \beta + \gamma$.
- ii) $\beta \leq \alpha + \gamma$.
- iii) $\gamma \leq \alpha + \beta$.
- iv) $\alpha + \beta + \gamma \leq 2\pi$.

Furthermore, A is positive definite if and only if all of these inequalities are strict. (Proof: See [149].)

Fact 8.8.16. Let $\lambda_1, \dots, \lambda_n \in \mathbb{C}$, assume that, for all $i = 1, \dots, n$, $\operatorname{Re} \lambda_i < 0$, and, for all $i, j = 1, \dots, n$, define $A \in \mathbb{C}^{n \times n}$ by

$$A_{(i,j)} \triangleq \frac{-1}{\overline{\lambda_i} + \lambda_j}.$$

Then, A is positive definite. (Proof: Note that $A = 2B \circ (1_{n \times n} - C)^{\circ-1}$, where $B_{(i,j)} = \frac{1}{(\overline{\lambda_i} - 1)(\lambda_j - 1)}$ and $C_{(i,j)} = \frac{(\overline{\lambda_i} + 1)(\lambda_j + 1)}{(\overline{\lambda_i} - 1)(\lambda_j - 1)}$. Then, note that B is positive semidefinite and that $(1_{n \times n} - C)^{\circ-1} = 1_{n \times n} + C + C^{\circ 2} + C^{\circ 3} + \dots$.) (Remark: A is the solution of a Lyapunov equation. See Fact 12.21.18 and Fact 12.21.19.) (Remark: A is a Cauchy matrix. See Fact 3.18.4, Fact 3.20.14, and Fact 3.20.15.) (Remark: A Cauchy matrix is also a Gram matrix defined in terms of the inner product of the functions $f_i(t) = e^{-\lambda_i t}$. See [201, p. 3].)

Fact 8.8.17. Let $\lambda_1, \dots, \lambda_n \in \text{OUD}$, and let $w_1, \dots, w_n \in \mathbb{C}$. Then, there exists a holomorphic function $\phi: \text{OUD} \mapsto \text{OUD}$ such that $\phi(\lambda_i) = w_i$ for all $i = 1, \dots, n$ if and only if $A \in \mathbb{C}^{n \times n}$ is positive semidefinite, where, for all $i, j = 1, \dots, n$,

$$A_{(i,j)} \triangleq \frac{1 - \overline{w_i}w_j}{1 - \overline{\lambda_i}\lambda_j}.$$

(Proof: See [985].) (Remark: A is a *Pick matrix*.)

Fact 8.8.18. Let $\alpha_0, \dots, \alpha_n > 0$, and define the tridiagonal matrix $A \in \mathbb{R}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} \alpha_0 + \alpha_1 & -\alpha_1 & 0 & 0 & \cdots & 0 \\ -\alpha_1 & \alpha_1 + \alpha_2 & -\alpha_2 & 0 & \cdots & 0 \\ 0 & -\alpha_2 & \alpha_2 + \alpha_3 & -\alpha_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & \alpha_{n-1} + \alpha_n \end{bmatrix}.$$

Then, A is positive definite. (Proof: For $k = 2, \dots, n$, the $k \times k$ leading principal subdeterminant of A is given by $\left[\sum_{i=0}^k \alpha_i^{-1}\right] \alpha_0 \alpha_1 \cdots \alpha_k$. See [146, p. 115].) (Remark: A is a stiffness matrix arising in structural analysis.) (Remark: See Fact 3.20.8.)

8.9 Facts on Identities and Inequalities for One Matrix

Fact 8.9.1. Let $n \leq 3$, let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, $|A|$ is positive semidefinite. (Proof: See [964].) (Remark: $|A|$ denotes the matrix whose entries are the absolute values of the entries of A .) (Remark: The result does not hold for $n \geq 4$. Let

$$A = \begin{bmatrix} 1 & \frac{1}{\sqrt{3}} & 0 & -\frac{1}{\sqrt{3}} \\ \frac{1}{\sqrt{3}} & 1 & \frac{1}{\sqrt{3}} & 0 \\ 0 & \frac{1}{\sqrt{3}} & 0 & \frac{1}{\sqrt{3}} \\ -\frac{1}{\sqrt{3}} & 0 & \frac{1}{\sqrt{3}} & 1 \end{bmatrix}.$$

Then, $\text{mspec}(A) = \{1 - \sqrt{6}/3, 1 - \sqrt{6}/3, 1 + \sqrt{6}/3, 1 + \sqrt{6}/3\}_{\text{ms}}$, whereas $\text{mspec}(|A|) = \{1, 1, 1 - \sqrt{12}/3, 1 + \sqrt{12}/3\}_{\text{ms}}$.)

Fact 8.9.2. Let $x \in \mathbb{F}^n$. Then,

$$xx^* \leq x^*xI.$$

Fact 8.9.3. Let $x \in \mathbb{F}^n$, assume that x is nonzero, and define $A \triangleq x^*xI - xx^*$. Then, A is positive semidefinite, $\text{mspec}(A) = \{x^*x, \dots, x^*x, 0\}_{\text{ms}}$, and $\text{rank } A = n - 1$.

Fact 8.9.4. Let $x, y \in \mathbb{F}^n$, assume that x and y are linearly independent, and define $A \triangleq (x^*x + y^*y)I - xx^* - yy^*$. Then, A is positive definite. Now, let $\mathbb{F} = \mathbb{R}$. Then,

$$\begin{aligned} \text{mspec}(A) = \{ & x^T x + y^T y, \dots, x^T x + y^T y, \\ & \frac{1}{2}(x^T x + y^T y) + \sqrt{\frac{1}{4}(x^T x - y^T y)^2 + (x^T y)^2}, \\ & \frac{1}{2}(x^T x + y^T y) - \sqrt{\frac{1}{4}(x^T x - y^T y)^2 + (x^T y)^2} \}_{\text{ms}}. \end{aligned}$$

(Proof: To show that A is positive definite, write $A = B + C$, where $B \triangleq x^* x I - x x^*$ and $C \triangleq y^* y I - y y^*$. Then, using Fact 8.9.3 it follows that $\mathcal{N}(B) = \text{span}\{x\}$ and $\mathcal{N}(C) = \text{span}\{y\}$. Now, it follows from Fact 8.7.3 that $\mathcal{N}(A) = \mathcal{N}(B) \cap \mathcal{N}(C) = \{0\}$. Therefore, A is nonsingular and thus positive definite. The expression for $\text{mspec}(A)$ follows from Fact 4.9.16.)

Fact 8.9.5. Let $x_1, \dots, x_n \in \mathbb{R}^3$, assume that $\text{span}\{x_1, \dots, x_n\} = \mathbb{R}^3$, and define $A \triangleq \sum_{i=1}^n (x_i^T x_i I - x_i x_i^T)$. Then, A is positive definite. Furthermore,

$$\lambda_1(A) < \lambda_2(A) + \lambda_3(A)$$

and

$$d_1(A) < d_2(A) + d_3(A).$$

(Proof: Suppose that $d_1(A) = A_{(1,1)}$. Then, $d_2(A) + d_3(A) - d_1(A) = 2 \sum_{i=1}^n x_{i(3)}^2 > 0$. Now, let $S \in \mathbb{R}^{3 \times 3}$ be such that $SAS^T = \sum_{i=1}^n (\hat{x}_i^T \hat{x}_i I - \hat{x}_i \hat{x}_i^T)$ is diagonal, where, for $i = 1, \dots, n$, $\hat{x}_i \triangleq Sx_i$. Then, for $i = 1, 2, 3$, $d_i(A) = \lambda_i(A)$.) (Remark: A is the inertia matrix for a rigid body consisting of n discrete particles. For a homogeneous continuum body $\mathcal{B}$ whose density is ρ , the inertia matrix is given by

$$I = \rho \iiint_{\mathcal{B}} (r^T r I - r r^T) dx dy dz,$$

where $r \triangleq \begin{bmatrix} x \\ y \\ z \end{bmatrix}$.) (Remark: The eigenvalues and diagonal entries of A represent the lengths of the sides of triangles. See Fact 1.11.17 and [1069, p. 220].)

Fact 8.9.6. Let $A \in \mathbb{F}^{2 \times 2}$, assume that A is positive semidefinite and nonzero, and define $B \in \mathbb{F}^{2 \times 2}$ by

$$B \triangleq \left(\text{tr } A + 2\sqrt{\det A} \right)^{-1/2} \left(A + \sqrt{\det A} I \right).$$

Then, $B = A^{1/2}$. (Proof: See [629, pp. 84, 266, 267].)

Fact 8.9.7. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then,

$$\text{rank } A = \nu_-(A) + \nu_+(A)$$

and

$$\text{def } A = \nu_0(A).$$

Fact 8.9.8. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and assume there exists $i \in \{1, \dots, n\}$ such that $A_{(i,i)} = 0$. Then, $\text{row}_i(A) = 0$ and $\text{col}_i(A) = 0$.

Fact 8.9.9. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, $A_{(i,i)} \geq 0$ for all $i = 1, \dots, n$, and $|A_{(i,j)}|^2 \leq A_{(i,i)} A_{(j,j)}$ for all $i, j = 1, \dots, n$.

Fact 8.9.10. Let $A \in \mathbb{F}^{n \times n}$. Then, $A \geq 0$ if and only if $A \geq -A$.

Fact 8.9.11. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, $A^2 \geq 0$.

Fact 8.9.12. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is skew Hermitian. Then, $A^2 \leq 0$.

Fact 8.9.13. Let $A \in \mathbb{F}^{n \times n}$, and let $\alpha > 0$. Then,

$$A^2 + A^{2*} \leq \alpha AA^* + \frac{1}{\alpha} A^*A.$$

Equality holds if and only if $\alpha A = A^*$.

Fact 8.9.14. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$(A - A^*)^2 \leq 0 \leq (A + A^*)^2 \leq 2(AA^* + A^*A).$$

Fact 8.9.15. Let $A \in \mathbb{F}^{n \times n}$, and let $\alpha > 0$. Then,

$$A + A^* \leq \alpha I + \alpha^{-1}AA^*.$$

Equality holds if and only if $A = \alpha I$.

Fact 8.9.16. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$2I \leq A + A^{-1}.$$

Equality holds if and only if $A = I$. Furthermore,

$$2n \leq \text{tr } A + \text{tr } A^{-1}.$$

Fact 8.9.17. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$(1_{1 \times n} A^{-1} 1_{n \times 1})^{-1} 1_{n \times n} \leq A.$$

(Proof: Set $B = 1_{n \times n}$ in Fact 8.21.14. See [1492].)

Fact 8.9.18. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then, $\begin{bmatrix} A & I \\ I & A^{-1} \end{bmatrix}$ is positive semidefinite.

Fact 8.9.19. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, $A^2 \leq A$ if and only if $0 \leq A \leq I$.

Fact 8.9.20. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, $\alpha I + A \geq 0$ if and only if $\alpha \geq -\lambda_{\min}(A)$. Furthermore,

$$A^2 + A + \frac{1}{4}I \geq 0.$$

Fact 8.9.21. Let $A \in \mathbb{F}^{n \times m}$. Then, $AA^* \leq I_n$ if and only if $A^*A \leq I_m$.

Fact 8.9.22. Let $A \in \mathbb{F}^{n \times n}$, and assume that either $AA^* \leq A^*A$ or $A^*A \leq AA^*$. Then, A is normal. (Proof: Use *ii*) of Corollary 8.4.10.)

Fact 8.9.23. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is a projector. Then,

$$0 \leq A \leq I.$$

Fact 8.9.24. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then,

$$\langle A^{-1} \rangle = \langle A^* \rangle^{-1}.$$

Fact 8.9.25. Let $A \in \mathbb{F}^{n \times m}$, and assume that A^*A is nonsingular. Then,

$$\langle A^* \rangle = A \langle A \rangle^{-1/2} A^*.$$

Fact 8.9.26. Let $A \in \mathbb{F}^{n \times n}$. Then, A is unitary if and only if there exists a nonsingular matrix $B \in \mathbb{F}^{n \times n}$ such that

$$A = \langle B^* \rangle^{-1/2} B.$$

If, in addition, A is real, then $\det B = \text{sign}(\det A)$. (Proof: For necessity, set $B = A$.) (Remark: See Fact 3.11.10.)

Fact 8.9.27. Let $A \in \mathbb{F}^{n \times n}$. Then, A is normal if and only if $\langle A \rangle = \langle A^* \rangle$. (Remark: See Fact 3.7.12.)

Fact 8.9.28. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$-\langle A \rangle - \langle A^* \rangle \leq A + A^* \leq \langle A \rangle + \langle A^* \rangle.$$

(Proof: See [886].)

Fact 8.9.29. Let $A \in \mathbb{F}^{n \times n}$, assume that A is normal, and let $\alpha, \beta \in (0, \infty)$. Then,

$$-\alpha \langle A \rangle - \beta \langle A^* \rangle \leq \langle \alpha A + \beta A^* \rangle \leq \alpha \langle A \rangle + \beta \langle A^* \rangle.$$

In particular,

$$-\langle A \rangle - \langle A^* \rangle \leq \langle A + A^* \rangle \leq \langle A \rangle + \langle A^* \rangle.$$

(Proof: See [886, 1494].) (Remark: See Fact 8.11.11.)

Fact 8.9.30. Let $A \in \mathbb{F}^{n \times n}$. The following statements hold:

- i) If $A \in \mathbb{F}^{n \times n}$ is positive definite, then $I + A$ is nonsingular and the matrices $I - B$ and $I + B$ are positive definite, where $B \triangleq (I + A)^{-1}(I - A)$.
- ii) If $I + A$ is nonsingular and the matrices $I - B$ and $I + B$ are positive definite, where $B \triangleq (I + A)^{-1}(I - A)$, then A is positive definite.

(Proof: See [463].) (Remark: For additional results on the Cayley transform, see Fact 3.11.28, Fact 3.11.29, Fact 3.11.30, Fact 3.19.12, and Fact 11.21.8.)

Fact 8.9.31. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\frac{1}{2j}(A - A^*)$ is positive definite. Then,

$$B \triangleq \left[\frac{1}{2}(A + A^*) \right]^{1/2} A^{-1} A^* \left[\frac{1}{2}(A + A^*) \right]^{-1/2}$$

is unitary. (Proof: See [466].) (Remark: A is *strictly dissipative* if $\frac{1}{2j}(A - A^*)$ is negative definite. A is strictly dissipative if and only if $-jA$ is dissipative. See [464, 465].) (Remark: $A^{-1}A^*$ is similar to a unitary matrix. See Fact 3.11.4.) (Remark: See Fact 8.13.11 and Fact 8.17.12.)

Fact 8.9.32. Let $A \in \mathbb{R}^{n \times n}$, assume that A is positive definite, assume that $A \leq I$, and define $(B_k)_{k=0}^\infty$ by $B_0 \triangleq 0$ and

$$B_{k+1} \triangleq B_k + \frac{1}{2}(A - B_k^2).$$

Then,

$$\lim_{k \rightarrow \infty} B_k = A^{1/2}.$$

(Proof: See [170, p. 181].) (Remark: See Fact 5.15.21.)

Fact 8.9.33. Let $A \in \mathbb{R}^{n \times n}$, assume that A is nonsingular, and define $(B_k)_{k=0}^\infty$ by $B_0 \triangleq A$ and

$$B_{k+1} \triangleq \frac{1}{2}(B_k + B_k^{-T}).$$

Then,

$$\lim_{k \rightarrow \infty} B_k = (AA^T)^{-1/2}A.$$

(Remark: The limit is unitary. See Fact 8.9.26. See [144, p. 224].)

Fact 8.9.34. Let $a, b \in \mathbb{R}$, and define the symmetric, Toeplitz matrix $A \in \mathbb{R}^{n \times n}$ by

$$A \triangleq aI_n + b1_{n \times n}.$$

Then, A is positive definite if and only if $a + nb > 0$ and $a > 0$. (Remark: See Fact 2.13.12 and Fact 4.10.15.)

Fact 8.9.35. Let $x_1, \dots, x_n \in \mathbb{R}^m$, and define

$$\bar{x} \triangleq \frac{1}{n} \sum_{j=1}^n x_j, \quad S \triangleq \frac{1}{n} \sum_{j=1}^n (x_j - \bar{x})(x_j - \bar{x})^T.$$

Then, for all $i = 1, \dots, n$,

$$(x_i - \bar{x})(x_i - \bar{x})^T \leq (n-1)S.$$

Furthermore, equality holds if and only if all of the elements of $\{x_1, \dots, x_n\} \setminus \{x_i\}$ are equal. (Proof: See [754, 1043, 1332].) (Remark: This result is an extension of the Laguerre-Samuelson inequality. See Fact 1.15.12.)

Fact 8.9.36. Let $x_1, \dots, x_n \in \mathbb{F}^n$, and define $A \in \mathbb{F}^{n \times n}$ by $A_{(i,j)} \triangleq x_i^* x_j$ for all $i, j = 1, \dots, n$, and $B \triangleq \begin{bmatrix} x_1 & \cdots & x_n \end{bmatrix}$. Then, $A = B^* B$. Consequently, A is positive semidefinite and $\text{rank } A = \text{rank } B$. Conversely, let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, there exist $x_1, \dots, x_n \in \mathbb{F}^n$ such that $A = B^* B$, where $B = \begin{bmatrix} x_1 & \cdots & x_n \end{bmatrix}$. (Proof: The converse is an immediate consequence of Corollary 5.4.5.) (Remark: A is the *Gram matrix* of $x_1, \dots, x_n$.)

Fact 8.9.37. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, there exists a matrix $B \in \mathbb{F}^{n \times n}$ such that B is lower triangular, B has nonnegative diagonal entries, and $A = BB^*$. If, in addition, A is positive definite, then B is unique and has positive diagonal entries. (Remark: This result is the *Cholesky decomposition*.)

Fact 8.9.38. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = m$. Then,

$$0 \leq A(A^*A)^{-1}A^* \leq I.$$

Fact 8.9.39. Let $A \in \mathbb{F}^{n \times m}$. Then, $I - A^*A$ is positive definite if and only if $I - AA^*$ is positive definite. In this case,

$$(I - A^*A)^{-1} = I + A^*(I - AA^*)^{-1}A.$$

Fact 8.9.40. Let $A \in \mathbb{F}^{n \times m}$, let α be a positive number, and define $A_\alpha \triangleq (\alpha I + A^*A)^{-1}A^*$. Then, the following statements are equivalent:

$$i) \quad AA_\alpha = A_\alpha A.$$

$$ii) \quad AA^* = A^*A.$$

Furthermore, the following statements are equivalent:

$$iii) \quad A_\alpha A^* = A^* A_\alpha.$$

$$iv) \quad AA^*A^2 = A^2A^*A.$$

(Proof: See [1299].) (Remark: A_α is a *regularized Tikhonov inverse*.)

Fact 8.9.41. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$A^{-1} \leq \frac{\alpha + \beta}{\alpha\beta}I - \frac{1}{\alpha\beta}A \leq \frac{(\alpha + \beta)^2}{4\alpha\beta}A^{-1},$$

where $\alpha \triangleq \lambda_{\max}(A)$ and $\beta \triangleq \lambda_{\min}(A)$. (Proof: See [972].)

Fact 8.9.42. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, the following statements hold:

$$i) \quad \text{If } \alpha \in [0, 1], \text{ then}$$

$$A^\alpha \leq \alpha A + (1 - \alpha)I.$$

$$ii) \quad \text{If } \alpha \in [0, 1] \text{ and } A \text{ is positive definite, then}$$

$$[\alpha A^{-1} + (1 - \alpha)I]^{-1} \leq A^\alpha \leq \alpha A + (1 - \alpha)I.$$

$$iii) \quad \text{If } \alpha \geq 1, \text{ then}$$

$$\alpha A + (1 - \alpha)I \leq A^\alpha.$$

$$iv) \quad \text{If } A \text{ is positive definite and either } \alpha \geq 1 \text{ or } \alpha \leq 0, \text{ then}$$

$$\alpha A + (1 - \alpha)I \leq A^\alpha \leq [\alpha A^{-1} + (1 - \alpha)I]^{-1}.$$

(Proof: See [530, pp. 122, 123].) (Remark: This result is a special case of the Young inequality. See Fact 1.9.2 and Fact 8.10.43.) (Remark: See Fact 8.12.26 and Fact 8.12.27.)

Fact 8.9.43. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$I - A^{-1} \leq \log A \leq A - I.$$

Furthermore, if $A \geq I$, then $\log A$ is positive semidefinite, and, if $A > I$, then $\log A$ is positive definite. (Proof: See Fact 1.9.22.)

8.10 Facts on Identities and Inequalities for Two or More Matrices

Fact 8.10.1. Let $\{A_i\}_{i=1}^\infty \subset \mathbf{H}^n$ and $\{B_i\}_{i=1}^\infty \subset \mathbf{H}^n$, assume that, for all $i \in \mathbb{P}$, $A_i \leq B_i$, and assume that $A \triangleq \lim_{i \rightarrow \infty} A_i$ and $B \triangleq \lim_{i \rightarrow \infty} B_i$ exist. Then, $A \leq B$.

Fact 8.10.2. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that $A \leq B$. Then, $\mathcal{R}(A) \subseteq \mathcal{R}(B)$ and $\text{rank } A \leq \text{rank } B$. Furthermore, $\mathcal{R}(A) = \mathcal{R}(B)$ if and only if $\text{rank } A = \text{rank } B$.

Fact 8.10.3. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, the following statements hold:

- i) $\lambda_{\min}(A) \leq \lambda_{\min}(B)$ if and only if $\lambda_{\min}(A)I \leq B$.
- ii) $\lambda_{\max}(A) \leq \lambda_{\max}(B)$ if and only if $A \leq \lambda_{\max}(B)I$.

Fact 8.10.4. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and consider the following conditions:

- i) $A \leq B$.
- ii) For all $i = 1, \dots, n$, $\lambda_i(A) \leq \lambda_i(B)$.
- iii) There exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that $A \leq SBS^*$.

Then, $i) \implies ii) \iff iii)$. (Remark: $i) \implies ii)$ is the monotonicity theorem given by Theorem 8.4.9.)

Fact 8.10.5. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, $0 < A \leq B$ if and only if $\text{sprad}(AB^{-1}) < 1$.

Fact 8.10.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$(A^{-1} + B^{-1})^{-1} = A(A + B)^{-1}B.$$

Fact 8.10.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$(A + B)^{-1} \leq \frac{1}{4}(A^{-1} + B^{-1}).$$

Equivalently,

$$A + B \leq AB^{-1}A + BA^{-1}B.$$

In both inequalities, equality holds if and only if $A = B$. (Proof: See [1490, p. 168].) (Remark: See Fact 1.10.4.)

Fact 8.10.8. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite, B is Hermitian, and $A + B$ is nonsingular. Then,

$$(A + B)^{-1} + (A + B)^{-1}B(A + B)^{-1} \leq A^{-1}.$$

If, in addition, B is nonsingular, the inequality is strict. (Proof: This inequality is equivalent to $BA^{-1}B \geq 0$. See [1050].)

Fact 8.10.9. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $\alpha \in [0, 1]$. Then,

$$\beta[\alpha A^{-1} + (1 - \alpha)B^{-1}] \leq [\alpha A + (1 - \alpha)B]^{-1},$$

where

$$\beta \triangleq \min_{\mu \in \text{mspec}(A^{-1}B)} \frac{4\mu}{(1 + \mu)^2}.$$

(Proof: See [1017].) (Remark: This result is a reverse form of a convex inequality.)

Fact 8.10.10. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times m}$, and assume that B is positive semidefinite. Then, $ABA^* = 0$ if and only if $AB = 0$.

Fact 8.10.11. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, AB is positive semidefinite if and only if AB is normal.

Fact 8.10.12. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that either *i*) A and B are positive semidefinite or *ii*) either A or B is positive definite. Then, AB is group invertible. (Proof: Use Theorem 8.3.2 and Theorem 8.3.5.)

Fact 8.10.13. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that A and $AB + BA$ are (positive semidefinite, positive definite). Then, B is (positive semidefinite, positive definite). (Proof: See [201, p. 8], [878, p. 120], or [1430]. Alternatively, the result follows from Corollary 11.9.4.)

Fact 8.10.14. Let $A, B, C \in \mathbb{F}^{n \times n}$, assume that A , B , and C are positive semidefinite, and assume that $A = B + C$. Then, the following statements are equivalent:

- i*) $\text{rank } A = \text{rank } B + \text{rank } C$.
- ii*) There exists $S \in \mathbb{F}^{m \times n}$ such that $\text{rank } S = m$, $\mathcal{R}(S) \cap \mathcal{N}(A) = \{0\}$, and either $B = AS^*(SAS^*)^{-1}SA$ or $C = AS^*(SAS^*)^{-1}SA$.

(Proof: See [285, 331].)

Fact 8.10.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian and nonsingular. Then, the following statements hold:

- i*) If every eigenvalue of AB is positive, then $\text{In } A = \text{In } B$.
- ii*) $\text{In } A - \text{In } B = \text{In}(A - B) + \text{In}(A^{-1} - B^{-1})$.
- iii*) If $\text{In } A = \text{In } B$ and $A \leq B$, then $B^{-1} \leq A^{-1}$.

(Proof: See [51, 109, 1047].) (Remark: The identity *ii*) is due to Styan. See [1047].) (Remark: An extension to singular A and B is given by Fact 8.20.14.)

Fact 8.10.16. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that $A \leq B$. Then, $A_{(i,i)} \leq B_{(i,i)}$ for all $i = 1, \dots, n$.

Fact 8.10.17. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that $A \leq B$. Then, $\text{sig } A \leq \text{sig } B$. (Proof: See [392, p. 148].)

Fact 8.10.18. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that $\langle A \rangle \leq B$. Then, either $A \leq B$ or $-A \leq B$. (Proof: See [1493].)

Fact 8.10.19. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite and B is positive definite. Then, $A \leq B$ if and only if $AB^{-1}A \leq A$.

Fact 8.10.20. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that $0 \leq A \leq B$. Then, there exists a matrix $S \in \mathbb{F}^{n \times n}$ such that $A = S^*BS$ and $S^*S \leq I$. (Proof: See [447, p. 269].)

Fact 8.10.21. Let $A, B, C, D \in \mathbb{F}^{n \times n}$, assume that A, B, C, D are positive semidefinite, and assume that $0 < D \leq C$ and $BCB \leq ADA$. Then, $B \leq A$. (Proof: See [84, 300].)

Fact 8.10.22. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then, there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that

$$\langle AB \rangle \leq \frac{1}{2}S(A^2 + B^2)S^*.$$

(Proof: See [90, 209].)

Fact 8.10.23. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, $ABA \leq B$ if and only if $AB = BA$. (Proof: See [1325].)

Fact 8.10.24. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite, $0 \leq A \leq I$, and B is positive definite. Then,

$$ABA \leq \frac{(\alpha + \beta)^2}{4\alpha\beta}B.$$

where $\alpha \triangleq \lambda_{\min}(B)$ and $\beta \triangleq \lambda_{\max}(B)$. (Proof: See [251].) (Remark: This inequality is related to Fact 1.16.6.)

Fact 8.10.25. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then,

$$(A + B)^{1/2} \leq A^{1/2} + B^{1/2}$$

if and only if $AB = BA$. (Proof: See [1317, p. 30].)

Fact 8.10.26. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that $0 \leq A \leq B$. Then,

$$(A + \frac{1}{4}A^2)^{1/2} \leq (B + \frac{1}{4}B^2)^{1/2}.$$

(Proof: See [1012].)

Fact 8.10.27. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $B \in \mathbb{F}^{l \times n}$. Then, BAB^* is positive definite if and only if $B(A + A^2)B^*$ is positive definite. (Proof: Diagonalize A using a unitary transformation and note that $BA^{1/2}$ and $B(A + A^2)^{1/2}$ have the same rank.)

Fact 8.10.28. Let $A, B, C \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and assume that B and C are positive semidefinite. Then,

$$2\operatorname{tr} \langle B^{1/2} C^{1/2} \rangle \leq \operatorname{tr}(AB + A^{-1}C).$$

Furthermore, there exists A such that equality holds if and only if $\operatorname{rank} B = \operatorname{rank} C = \operatorname{rank} B^{1/2} C^{1/2}$. (Proof: See [35, 494].) (Remark: A matrix A for which equality holds is given in [35].) (Remark: Applications to linear systems are given in [1442].)

Fact 8.10.29. Let $A_1, \dots, A_k \in \mathbb{F}^{n \times n}$, and assume that $A_1, \dots, A_k$ are positive definite. Then,

$$n^2 \left(\sum_{i=1}^k A_i \right)^{-1} \leq \sum_{i=1}^k A_i^{-1}.$$

(Remark: This result is an extension of Fact 1.15.37.)

Fact 8.10.30. Let $A_1, \dots, A_k \in \mathbb{F}^{n \times n}$, assume that $A_1, \dots, A_k$ are positive semidefinite, and let $p, q \in \mathbb{R}$ satisfy $1 \leq p \leq q$. Then,

$$\left(\frac{1}{k} \sum_{i=1}^k A_i^p \right)^{1/p} \leq \left(\frac{1}{k} \sum_{i=1}^k A_i^q \right)^{1/q}.$$

(Proof: See [193].)

Fact 8.10.31. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, let $S \in \mathbb{F}^{n \times n}$ be such that $SAS^* = \operatorname{diag}(\alpha_1, \dots, \alpha_n)$ and $SBS^* = \operatorname{diag}(\beta_1, \dots, \beta_n)$, and define

$$C_l \triangleq S^{-1} \operatorname{diag}(\min\{\alpha_1, \beta_1\}, \dots, \min\{\alpha_n, \beta_n\}) S^{-*}$$

and

$$C_u \triangleq S^{-1} \operatorname{diag}(\max\{\alpha_1, \beta_1\}, \dots, \max\{\alpha_n, \beta_n\}) S^{-*}.$$

Then, C_l and C_u are independent of the choice of S , and

$$C_l \leq A \leq C_u,$$

$$C_l \leq B \leq C_u.$$

(Proof: See [900].)

Fact 8.10.32. Let $A, B \in \mathbf{H}^{n \times n}$. Then, $\operatorname{glb}\{A, B\}$ exists in $\mathbf{H}^n$ with respect to the ordering “ $\leq$ ” if and only if either $A \leq B$ or $B \leq A$. (Proof: See [784].) (Remark: Let $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$. Then, $C = 0$ is a lower bound for $\{A, B\}$. Furthermore, $D = \begin{bmatrix} -1 & \sqrt{2} \\ \sqrt{2} & -1 \end{bmatrix}$, which has eigenvalues $-1 - \sqrt{2}$ and $-1 + \sqrt{2}$, is also a lower bound for $\{A, B\}$ but is not comparable with C .)

Fact 8.10.33. Let $A, B \in \mathbf{H}^{n \times n}$, and assume that A and B are positive semidefinite. Then, the following statements hold:

- i) $\{A, B\}$ does not necessarily have a least upper bound in $\mathbf{N}^n$.
- ii) If A and B are positive definite, then $\{A, B\}$ has a greatest lower bound in $\mathbf{N}^n$ if and only if A and B are comparable.

- iii) If A is a projector and $0 \leq B \leq I$, then $\{A, B\}$ has a greatest lower bound in $\mathbf{N}^n$.
- iv) If $A, B \in \mathbf{N}^n$ are projectors, then the greatest lower bound of $\{A, B\}$ in $\mathbf{N}^n$ is given by

$$\text{glb}\{A, B\} = 2A(A + B)^+B,$$

which is the projector onto $\mathcal{R}(A) \cap \mathcal{R}(B)$.

- v) $\text{glb}\{A, B\}$ exists in $\mathbf{N}^n$ if and only if $\text{glb}\{A, \text{glb}\{AA^+, BB^+\}\}$ and $\text{glb}\{B, \text{glb}\{AA^+, BB^+\}\}$ are comparable. In this case,

$$\text{glb}\{A, B\} = \min\{\text{glb}\{A, \text{glb}\{AA^+, BB^+\}\}, \text{glb}\{B, \text{glb}\{AA^+, BB^+\}\}\}.$$

- vi) $\text{glb}\{A, B\}$ exists if and only if $\text{sh}(A, B)$ and $\text{sh}(B, A)$ are comparable, where $\text{sh}(A, B) \triangleq \lim_{\alpha \rightarrow \infty} (\alpha B):A$. In this case,

$$\text{glb}\{A, B\} = \min\{\text{sh}(A, B), \text{sh}(B, A)\}.$$

(Proof: To prove *i*), let $A = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ and $B = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$, and suppose that Z is the least upper bound for A and B . Hence, $A \leq Z \leq I$ and $B \leq Z \leq I$, and thus $Z = I$. Next, note that $X \triangleq \begin{bmatrix} 4/3 & 2/3 \\ 2/3 & 4/3 \end{bmatrix}$ satisfies $A \leq X$ and $B \leq X$. However, it is not true that $Z \leq X$, which implies that $\{A, B\}$ does not have a least upper bound. See [239, p. 11]. Statement *ii*) is given in [441, 550, 1021]. Statements *iii*) and *v*) are given in [1021]. Statement *iv*) is given in [39]. The expression for the projector onto $\mathcal{R}(A) \cap \mathcal{R}(B)$ is given in Fact 6.4.41. Statement *vi*) is given in [50].) (Remark: The partially ordered cones $\mathbf{H}^n$ and $\mathbf{N}^n$ with the ordering “ $\leq$ ” are not lattices.) (Remark: $\text{sh}(A, B)$ is the shorted operator, see Fact 8.20.19. However, the usage here is more general since B need not be a projector. See [50].) (Remark: An alternative approach to showing that $\mathbf{N}^n$ is not a lattice is given in [900].) (Remark: The cone $\mathbf{N}$ is a partially ordered set under the spectral order, see Fact 8.10.35.)

Fact 8.10.34. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let p be a real number, and assume that either $p \in [1, 2]$ or A and B are positive definite and $p \in [-1, 0] \cup [1, 2]$. Then,

$$\left[\frac{1}{2}(A + B)\right]^p \leq \frac{1}{2}(A^p + B^p).$$

(Proof: See [854].)

Fact 8.10.35. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p, q \in \mathbb{R}$ satisfy $p \geq q \geq 0$. Then,

$$\left[\frac{1}{2}(A^q + B^q)\right]^{1/q} \leq \left[\frac{1}{2}(A^p + B^p)\right]^{1/p}.$$

Furthermore,

$$\mu(A, B) \triangleq \lim_{p \rightarrow \infty} \left[\frac{1}{2}(A^p + B^p)\right]^{1/p}$$

exists and satisfies

$$A \leq \mu(A, B), \quad B \leq \mu(A, B).$$

(Proof: See [171].) (Remark: $\mu(A, B)$ is the least upper bound of A and B with respect to the spectral order. See [54, 795] and Fact 8.19.4.)

Fact 8.10.36. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let $p \in (1, \infty)$, and let $\alpha \in [0, 1]$. Then,

$$\alpha^{1-1/p}A + (1 - \alpha)^{1-1/p}B \leq (A^p + B^p)^{1/p}.$$

(Proof: See [54].)

Fact 8.10.37. Let $A, B, C \in \mathbb{F}^{n \times n}$. Then,

$$A^*A + B^*B = (B + CA)^*(I + CC^*)^{-1}(B + CA) + (A - C^*B)(I + C^*C)^{-1}(A - C^*B).$$

(Proof: See [717].) (Remark: See Fact 8.13.29.)

Fact 8.10.38. Let $A \in \mathbb{F}^{n \times n}$, let $\alpha \in \mathbb{R}$, and assume that either A is nonsingular or $\alpha \geq 1$. Then,

$$(A^*A)^\alpha = A^*(AA^*)^{\alpha-1}A.$$

(Proof: Use the singular value decomposition.) (Remark: This result is given in [512, 526].)

Fact 8.10.39. Let $A, B \in \mathbb{F}^{n \times n}$, let $\alpha \in \mathbb{R}$, assume that A and B are positive semidefinite, and assume that either A and B are positive definite or $\alpha \geq 1$. Then,

$$(AB^2A)^\alpha = AB(BA^2B)^{\alpha-1}BA.$$

(Proof: Use Fact 8.10.38.)

Fact 8.10.40. Let $A, B, C \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, B is positive definite, and $B = C^*C$, and let $\alpha \in [0, 1]$. Then,

$$C^*(C^{-*}AC^{-1})^\alpha C \leq \alpha A + (1 - \alpha)B.$$

If, in addition, $\alpha \in (0, 1)$, then equality holds if and only if $A = B$. (Proof: See [995].)

Fact 8.10.41. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $p \in \mathbb{R}$. Furthermore, assume that either A and B are nonsingular or $p \geq 1$. Then,

$$(BAB^*)^p = BA^{1/2}(A^{1/2}B^*BA^{1/2})^{p-1}A^{1/2}B^*.$$

(Proof: See [526] or [530, p. 129].)

Fact 8.10.42. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $p \in \mathbb{R}$. Then,

$$(BAB)^p = BA^{1/2}(A^{1/2}B^2A^{1/2})^{p-1}A^{1/2}B.$$

(Proof: See [524, 674].)

Fact 8.10.43. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Furthermore, if A is positive definite, then define

$$A \# B \triangleq A^{1/2} \left(A^{-1/2} B A^{-1/2} \right)^{1/2} A^{1/2},$$

whereas, if A is singular, then define

$$A\#B \triangleq \lim_{\varepsilon \downarrow 0} (A + \varepsilon I)\#B.$$

Then, the following statements hold:

- i) $A\#B$ is positive semidefinite.
- ii) $A\#A = A$.
- iii) $A\#B = B\#A$.
- iv) $\mathcal{R}(A\#B) = \mathcal{R}(A) \cap \mathcal{R}(B)$.
- v) If $S \in \mathbb{F}^{m \times n}$ is right invertible, then $(SAS^*)\#(SBS^*) \leq S(A\#B)S^*$.
- vi) If $S \in \mathbb{F}^{n \times n}$ is nonsingular, then $(SAS^*)\#(SBS^*) = S(A\#B)S^*$.
- vii) If $C, D \in \mathbf{P}^n$, $A \leq C$, and $B \leq D$, then $A\#B \leq C\#D$.
- viii) If $C, D \in \mathbf{P}^n$, then

$$(A\#C) + (C\#D) \leq (A + B)\#(C + D).$$

- ix) If $A \leq B$, then

$$4A\#(B - A) = [A + A\#(4B - 3A)]\#[-A + A\#(4B - 3A)].$$

- x) If $\alpha \in [0, 1]$, then

$$\sqrt{\alpha}(A\#B) \pm \frac{1}{2}\sqrt{1 - \alpha}(A - B) \leq \frac{1}{2}(A + B).$$

- xi) $A\#B = \max\{X \in \mathbf{H}: \begin{bmatrix} A & X \\ X & B \end{bmatrix} \text{ is positive semidefinite}\}.$
- xii) Let $X \in \mathbb{F}^{n \times n}$, and assume that X is Hermitian and

$$\begin{bmatrix} A & X \\ X & B \end{bmatrix} \geq 0.$$

Then,

$$-A\#B \leq X \leq A\#B.$$

Furthermore, $\begin{bmatrix} A & A\#B \\ A\#B & B \end{bmatrix}$ and $\begin{bmatrix} A & -A\#B \\ -A\#B & B \end{bmatrix}$ are positive semidefinite.

- xiii) If $S \in \mathbb{F}^{n \times n}$ is unitary and $A^{1/2}SB^{1/2}$ is positive semidefinite, then $A\#B = A^{1/2}SB^{1/2}$.

Now, assume that A is positive definite. Then, the following statements hold:

- xiv) $(A\#B)A^{-1}(A\#B) = B$.
- xv) For all $\alpha \in \mathbb{R}$, $A\#B = A^{1-\alpha}(A^{\alpha-1}BA^{-\alpha})^{1/2}A^{\alpha}$.
- xvi) $A\#B = A(A^{-1}B)^{1/2} = (BA^{-1})^{1/2}A$.
- xvii) $A\#B = (A + B)[(A + B)^{-1}A(A + B)^{-1}B]^{1/2}$.

Now, assume that A and B are positive definite. Then, the following statements hold:

- xviii) $A\#B$ is positive definite.

xix) $S \triangleq (A^{-1/2}BA^{-1/2})^{1/2}A^{1/2}B^{-1/2}$ is unitary, and $A\#B = A^{1/2}SB^{1/2}$.

xx) $\det A\#B = \sqrt{(\det A)\det B}$.

xxi) $\det (A\#B)^2 = \det AB$.

xxii) $(A\#B)^{-1} = A^{-1}\#B^{-1}$.

xxiii) Let $A_0 \triangleq A$ and $B_0 \triangleq B$, and, for all $k \in \mathbb{N}$, define $A_{k+1} \triangleq 2(A_k^{-1} + B_k^{-1})^{-1}$ and $B_{k+1} \triangleq \frac{1}{2}(A_k + B_k)$. Then, for all $k \in \mathbb{N}$,

$$A_k \leq A_{k+1} \leq A\#B \leq B_{k+1} \leq B_k$$

and

$$\lim_{k \rightarrow \infty} A_k = \lim_{k \rightarrow \infty} B_k = A\#B.$$

xxiv) For all $\alpha \in (-1, 1)$, $\begin{bmatrix} A & \alpha A\#B \\ \alpha A\#B & B \end{bmatrix}$ is positive definite.

xxv) $\text{rank} \begin{bmatrix} A & A\#B \\ A\#B & B \end{bmatrix} = \text{rank} \begin{bmatrix} A & -A\#B \\ -A\#B & B \end{bmatrix} = n$.

Furthermore, the following statements hold:

xxvi) If $n = 2$, then

$$A\#B = \frac{\sqrt{\alpha\beta}}{\sqrt{\det(\alpha^{-1}A + \beta^{-1}B)}}(\alpha^{-1}A + \beta^{-1}B).$$

xxvii) If $0 < A \leq B$, then $\phi: [0, \infty) \mapsto \mathbf{P}^n$ defined by $\phi(p) \triangleq A^{-p}\#B^p$ is nondecreasing.

xxviii) If B is positive definite and $A \leq B$, then

$$A^2\#B^{-2} \leq A\#B^{-1} \leq I.$$

xxix) If A and B are positive semidefinite, then

$$(BA^2B)^{1/2} \leq B^{1/2}(B^{1/2}AB^{1/2})^{1/2}B^{1/2} \leq B^2.$$

Finally, let $X \in \mathbf{H}^n$. Then, the following statements are equivalent:

xxx) $\begin{bmatrix} A & X \\ X & B \end{bmatrix}$ is positive semidefinite.

xxxi) $XA^{-1}X \leq B$.

xxxii) $XB^{-1}X \leq A$.

xxxiii) $-A\#B \leq X \leq A\#B$.

(Proof: See [45, 486, 583, 877, 1314]. For *xiii*), *xix*), and *xxvi*), see [201, pp. 108, 109, 111]. For *xxvii*), see [46]. Statement *xxvii*) implies *xxviii*), which, in turn, implies *xxix*.) (Remark: The square roots in *xvi*) indicate a semisimple matrix with positive diagonal entries.) (Remark: $A\#B$ is the *geometric mean* of A and B . A related mean is defined in [486]. Alternative means and their differences are considered in [20]. Geometric means for an arbitrary number of positive-definite matrices are discussed in [57, 809, 1014, 1084].) (Remark: See Fact 12.23.4.) (Remark: Inverse problems are considered in [41].) (Remark: *xxix*) interpolates (8.6.6).) (Remark:

Compare statements *xiii*) and *xix*) with Fact 8.11.6.) (Remark: See Fact 10.10.4.) (Problem: For singular A and B , express $A\#B$ in terms of generalized inverses.)

Fact 8.10.44. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, the following statements are equivalent:

- i) $A \leq B$.
- ii) For all $t \geq 0$, $I \leq e^{-tA} \# e^{tB}$.
- iii) $\phi: [0, \infty) \mapsto \mathbf{P}^n$ defined by $\phi(t) \triangleq e^{-tA} \# e^{tB}$ is nondecreasing.

(Proof: See [46].)

Fact 8.10.45. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\alpha \in [0, 1]$. Furthermore, if A is positive definite, then define

$$A\#_{\alpha}B \triangleq A^{1/2} \left(A^{-1/2} B A^{-1/2} \right)^{\alpha} A^{1/2},$$

whereas, if A is singular, then define

$$A\#_{\alpha}B \triangleq \lim_{\varepsilon \downarrow 0} (A + \varepsilon I) \#_{\alpha} B.$$

Then, the following statements hold:

- i) $A\#_{\alpha}B = B\#_{1-\alpha}A$.
- ii) $(A\#_{\alpha}B)^{-1} = A^{-1} \#_{\alpha} B^{-1}$.

Fact 8.10.46. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $\alpha \in [0, 1]$. Then,

$$[\alpha A^{-1} + (1 - \alpha) B^{-1}]^{-1} \leq A^{1/2} \left(A^{-1/2} B A^{-1/2} \right)^{1-\alpha} A^{1/2} \leq \alpha A + (1 - \alpha) B,$$

or, equivalently,

$$[\alpha A^{-1} + (1 - \alpha) B^{-1}]^{-1} \leq A\#_{1-\alpha}B \leq \alpha A + (1 - \alpha) B,$$

or, equivalently,

$$[\alpha A + (1 - \alpha) B]^{-1} \leq A^{-1/2} \left(A^{-1/2} B A^{-1/2} \right)^{\alpha-1} A^{-1/2} \leq \alpha A^{-1} + (1 - \alpha) B^{-1}.$$

Consequently,

$$\text{tr} [\alpha A + (1 - \alpha) B]^{-1} \leq \text{tr} \left[A^{-1} \left(A^{-1/2} B A^{-1/2} \right)^{\alpha-1} \right] \leq \text{tr} [\alpha A^{-1} + (1 - \alpha) B^{-1}]$$

and

$$\frac{2\alpha\beta}{(\alpha+\beta)^2} (A + B) \leq 2(A^{-1} + B^{-1})^{-1} \leq A\#B \leq \frac{1}{2}(A + B) \leq \frac{(\alpha+\beta)^2}{2\alpha\beta} (A^{-1} + B^{-1})^{-1},$$

where

$$\alpha \triangleq \min\{\lambda_{\min}(A), \lambda_{\min}(B)\}$$

and

$$\beta \triangleq \max\{\lambda_{\max}(A), \lambda_{\max}(B)\}.$$

(Remark: The left-hand inequality in the first string of inequalities is the *Young inequality*. See [530, p. 122], Fact 1.10.21, and Fact 8.9.42. Setting $B = I$ yields

Fact 8.9.42. The fourth string of inequalities improves the fact that $\phi(A) = A^{-1}$ is convex as shown by *iv*) of Proposition 8.6.17. The last string of inequalities follows from the fourth string of inequalities with $\alpha = 1/2$ along with results given in [1283] and [1490, p. 174].) (Remark: Related inequalities are given by Fact 8.12.26 and Fact 8.12.27. See also Fact 8.20.18.)

Fact 8.10.47. Let $(x_i)_{i=1}^\infty \subset \mathbb{R}^n$, assume that $\sum_{i=1}^\infty x_i$ exists, and let $(A_i)_{i=1}^\infty \subset \mathbb{N}^n$ be such that $A_i \leq A_{i+1}$ for all $i \in \mathbb{P}$ and $\lim_{i \rightarrow \infty} \text{tr } A_i = \infty$. Then,

$$\lim_{k \rightarrow \infty} (\text{tr } A_k)^{-1} \sum_{i=1}^k A_i x_i = 0.$$

If, in addition A_i is positive definite for all $i \in \mathbb{P}$ and $\{\lambda_{\max}(A_i)/\lambda_{\min}(A_i)\}_{i=1}^\infty$ is bounded, then

$$\lim_{k \rightarrow \infty} A_k^{-1} \sum_{i=1}^k A_i x_i = 0.$$

(Proof: See [33].) (Remark: These identities are matrix versions of the *Kronecker lemma*.) (Remark: Extensions are given in [623].)

Fact 8.10.48. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, assume that $A \leq B$, and let $p \geq 1$. Then,

$$A^p \leq K(\lambda_{\min}(A), \lambda_{\min}(A), p) B^p \leq \left[\frac{\lambda_{\max}(A)}{\lambda_{\min}(A)} \right]^{p-1} B^p,$$

where

$$K(a, b, p) \triangleq \frac{a^p b - ab^p}{(p-1)(a-b)} \left[\frac{(p-1)(a^p - b^p)}{p(a^p b - ab^p)} \right]^p.$$

(Proof: See [249, 528] and [530, pp. 193, 194].) (Remark: $K(a, b, p)$ is the *Fan constant*.)

Fact 8.10.49. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive definite and B is positive semidefinite, and let $p \geq 1$. Then, there exist unitary matrices $U, V \in \mathbb{F}^{n \times n}$ such that

$$\frac{1}{K(\lambda_{\min}(A), \lambda_{\min}(A), p)} U(BAB)^p U^* \leq B^p A^p B^p \leq K(\lambda_{\min}(A), \lambda_{\min}(A), p) V(BAB)^p V^*,$$

where $K(a, b, p)$ is the Fan constant defined in Fact 8.10.48.) (Proof: See [249].) (Remark: See Fact 8.12.20, Fact 8.18.26, and Fact 9.9.17.)

Fact 8.10.50. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive definite, B is positive semidefinite, and $B \leq A$, and let $p \geq 1$ and $r \geq 1$. Then,

$$\left[A^{r/2} \left(A^{-1/2} B^p A^{-1/2} \right)^r A^{r/2} \right]^{1/p} \leq A^r.$$

In particular,

$$\left\langle A^{-1/2} B^p A^{1/2} \right\rangle^{2/p} \leq A^2.$$

(Proof: See [53].)

Fact 8.10.51. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite and B is positive semidefinite. Then, the following statements are equivalent:

- i) $B \leq A$.
- ii) For all $p, q, r, t \in \mathbb{R}$ such that $p \geq 1$, $r \geq 0$, $t \geq 0$, and $q \in [1, 2]$,

$$\left[A^{r/2} \left(A^{t/2} B^p A^{t/2} \right)^q A^{r/2} \right]^{\frac{r+t+1}{r+q+qp}} \leq A^{r+t+1}.$$

- iii) For all $p, q, r, \tau \in \mathbb{R}$ such that $p \geq 1$, $r \geq \tau$, $q \geq 1$, and $\tau \in [0, 1]$,

$$\left[A^{r/2} \left(A^{-\tau/2} B^p A^{-\tau/2} \right)^q A^{r/2} \right]^{\frac{r-\tau}{r-q\tau+qp}} \leq A^{r-\tau}.$$

- iv) For all $p, q, r, \tau \in \mathbb{R}$ be such that $p \geq 1$, $r \geq \tau$, $\tau \in [0, 1]$, and $q \geq 1$,

$$\left[A^{r/2} \left(A^{-\tau/2} B^p A^{-\tau/2} \right)^q A^{r/2} \right]^{\frac{r-\tau+1}{r-q\tau+qp}} \leq A^{r-\tau+1}.$$

In particular, if $B \leq A$, $p \geq 1$, and $r \geq 1$, then

$$\left[A^{r/2} \left(A^{-1/2} B^p A^{-1/2} \right)^r A^{r/2} \right]^{\frac{r-1}{pr}} \leq A^{r-1}.$$

(Proof: Condition *ii*) is given in [512], *iii*) appears in [531], and *iv*) appears in [512]. See also [513].) (Remark: Setting $q = r$ and $\tau = 1$ in *iv*) yields Fact 8.10.50.)

Fact 8.10.52. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then, the following statements are equivalent:

- i) $B \leq A$.
- ii) There exist $r \in [0, \infty)$, $p \in [1, \infty)$, and a nonnegative integer k such that $(k+1)(r+1) = p+r$ and

$$B^r \leq \left(B^{r/2} A^p B^{r/2} \right)^{\frac{1}{k+1}}.$$

- iii) There exist $r \in [0, \infty)$, $p \in [1, \infty)$, and a nonnegative integer k such that $(k+1)(r+1) = p+r$ and

$$\left(A^{r/2} B^p A^{r/2} \right)^{\frac{1}{k+1}} \leq A^r.$$

(Proof: See [914].) (Remark: See Fact 8.19.1.)

Fact 8.10.53. Each of the following functions $\phi: (0, \infty) \mapsto (0, \infty)$ yields an increasing function $\phi: \mathbf{P}^n \mapsto \mathbf{P}^n$:

- i) $\phi(x) = \frac{x^{p+1/2}}{x^{2p}+1}$, where $p \in [0, 1/2]$.
- ii) $\phi(x) = x(1+x) \log(1+1/x)$.
- iii) $\phi(x) = \frac{1}{(1+x) \log(1+1/x)}$.
- iv) $\phi(x) = \frac{x-1-\log x}{(\log x)^2}$.
- v) $\phi(x) = \frac{x(\log x)^2}{x-1-\log x}$.

- $vi) \phi(x) = \frac{x(x+2) \log(x+2)}{(x+1)^2}.$
- $vii) \phi(x) = \frac{x(x+1)}{(x+2) \log(x+2)}.$
- $viii) \phi(x) = \frac{(x^2-1) \log(1+x)}{x^2}.$
- $ix) \phi(x) = \frac{x(x-1)}{(x+1) \log(x+1)}.$
- $x) \phi(x) = \frac{(x-1)^2}{(x+1) \log x}.$
- $xi) \phi(x) = \frac{p-1}{p} \left(\frac{x^p-1}{x^{p-1}-1} \right), \text{ where } p \in [-1, 2].$
- $xii) \phi(x) = \frac{x-1}{\log x}.$
- $xiii) \phi(x) = \sqrt{x}.$
- $xiv) \phi(x) = \frac{2x}{x+1}.$
- $xv) \phi(x) = \frac{x-1}{x^p-1}, \text{ where } p \in (0, 1].$

(Proof: See [534, 1084]. To obtain $xii)$, $xiii)$, and $xiv)$, set $p = 1, 1/2, -1$, respectively, in $xi)$.)

Fact 8.10.54. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite, $A \leq B$, and $AB = BA$. Then, $A^2 \leq B^2$. (Proof: See [110].)

8.11 Facts on Identities and Inequalities for Partitioned Matrices

Fact 8.11.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, the following statements hold:

- $i) \begin{bmatrix} A & A \\ A & A \end{bmatrix} \text{ and } \begin{bmatrix} A & -A \\ -A & A \end{bmatrix} \text{ are positive semidefinite.}$
- $ii) \text{ If } \begin{bmatrix} \alpha & \beta \\ \beta & \gamma \end{bmatrix} \in \mathbb{F}^{2 \times 2} \text{ is positive semidefinite, then } \begin{bmatrix} \alpha A & \beta A \\ \beta A & \gamma A \end{bmatrix} \text{ is positive semidefinite.}$
- $iii) \text{ If } A \text{ and } \begin{bmatrix} \alpha & \beta \\ \beta & \gamma \end{bmatrix} \text{ are positive definite, then } \begin{bmatrix} \alpha A & \beta A \\ \beta A & \gamma A \end{bmatrix} \text{ is positive definite.}$

(Proof: Use Fact 7.4.16.)

Fact 8.11.2. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times m}$, assume that $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ is positive semidefinite, and assume that $\begin{bmatrix} \alpha & \beta \\ \beta & \gamma \end{bmatrix} \in \mathbb{F}^{2 \times 2}$ is positive semidefinite. Then, the following statements hold:

- $i) \begin{bmatrix} \alpha 1_{n \times n} & \beta 1_{n \times m} \\ \beta 1_{m \times n} & \gamma 1_{m \times m} \end{bmatrix} \text{ is positive semidefinite.}$
- $ii) \begin{bmatrix} \alpha A & \beta B \\ \beta B^* & \gamma C \end{bmatrix} \text{ is positive semidefinite.}$
- $iii) \text{ If } \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \text{ is positive definite and } \alpha \text{ and } \gamma \text{ are positive, then } \begin{bmatrix} \alpha A & \beta B \\ \beta B^* & \gamma C \end{bmatrix} \text{ is positive definite.}$

(Proof: To prove *i*), use Proposition 8.2.4. Statements *ii*) and *iii*) follow from Fact 8.21.12.)

Fact 8.11.3. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that A and B are partitioned identically as $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$ and $B = \begin{bmatrix} B_{11} & B_{12} \\ B_{12}^* & B_{22} \end{bmatrix}$. Then,

$$A_{22}|A + B_{22}|B \leq (A_{22} + B_{22})|(A + B).$$

Now, assume that A_{22} and B_{22} are positive definite. Then, equality holds if and only if $A_{12}A_{22}^{-1} = B_{12}B_{22}^{-1}$. (Proof: See [485, 1057].) (Remark: The first inequality, which follows from *xvii*) of Proposition 8.6.17, is an extension of Bergstrom's inequality, which corresponds to the case in which A_{11} is a scalar. See Fact 8.15.18.)

Fact 8.11.4. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, assume that A and B are partitioned identically as $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix}$ and $B = \begin{bmatrix} B_{11} & B_{12} \\ B_{12}^* & B_{22} \end{bmatrix}$, and assume that A_{11} and B_{11} are positive definite. Then,

$$(A_{12} + B_{12})^*(A_{11} + B_{11})^{-1}(A_{12} + B_{12}) \leq A_{12}^*A_{11}^{-1}A_{12} + B_{12}^*B_{11}^{-1}B_{12}$$

and

$$\begin{aligned} & \text{rank}[A_{12}^*A_{11}^{-1}A_{12} + B_{12}^*B_{11}^{-1}B_{12} - (A_{12} + B_{12})^*(A_{11} + B_{11})^{-1}(A_{12} + B_{12})] \\ &= \text{rank}(A_{12} - A_{11}B_{11}^{-1}B_{12}). \end{aligned}$$

Furthermore,

$$\frac{\det A}{\det A_{11}} + \frac{\det B}{\det B_{11}} \leq \frac{\det(A + B)}{\det(A_{11} + B_{11})} = \det[(A_{11} + B_{11})|(A + B)].$$

(Remark: The last inequality generalizes Fact 8.13.17.)

Fact 8.11.5. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$. Then, the following statements hold:

i) If $\mathcal{A}$ is positive semidefinite, then

$$0 \leq BC^+B^* \leq A.$$

ii) If $\mathcal{A}$ is positive definite, then C is positive definite and

$$0 \leq BC^{-1}B^* < A.$$

Now, assume that $n = m$. Then, the following statements hold:

iii) If $\mathcal{A}$ is positive semidefinite, then

$$-A - C \leq B + B^* \leq A + C.$$

iv) If $\mathcal{A}$ is positive definite, then

$$-A - C < B + B^* < A + C.$$

(Proof: The first two statements follow from Proposition 8.2.4. To prove the last

two statements, consider SAS^T , where $S \triangleq \begin{bmatrix} I & I \end{bmatrix}$ and $S \triangleq \begin{bmatrix} I & -I \end{bmatrix}$. (Remark: See Fact 8.21.40.)

Fact 8.11.6. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$. Then, $\mathcal{A}$ is positive semidefinite if and only if A and C are positive semidefinite and there exists a semicontractive matrix $S \in \mathbb{F}^{n \times m}$ such that

$$B = A^{1/2}SC^{1/2}.$$

(Proof: See [719].) (Remark: Compare this result with statements *xiii*) and *xix*) of Fact 8.10.43.)

Fact 8.11.7. Let $A, B, C \in \mathbb{F}^{n \times n}$, assume that $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{2n \times 2n}$ is positive semidefinite, and assume that $AB = BA$. Then,

$$B^*B \leq A^{1/2}CA^{1/2}.$$

(Proof: See [1492].)

Fact 8.11.8. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, $-A \leq B \leq A$ if and only if $\begin{bmatrix} A & B \\ B & A \end{bmatrix}$ is positive semidefinite. Furthermore, $-A < B < A$ if and only if $\begin{bmatrix} A & B \\ B & A \end{bmatrix}$ is positive definite. (Proof: Note that

$$\frac{1}{\sqrt{2}} \begin{bmatrix} I & -I \\ I & I \end{bmatrix} \begin{bmatrix} A & B \\ B & A \end{bmatrix} \frac{1}{\sqrt{2}} \begin{bmatrix} I & I \\ -I & I \end{bmatrix} = \begin{bmatrix} A - B & 0 \\ 0 & A + B \end{bmatrix}.)$$

Fact 8.11.9. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, assume that $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$ is positive semidefinite, and let $r \triangleq \text{rank } B$. Then, for all $k = 1, \dots, r$,

$$\prod_{i=1}^k \sigma_i(B) \leq \prod_{i=1}^k \max\{\lambda_i(A), \lambda_i(C)\}.$$

(Proof: See [1492].)

Fact 8.11.10. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$, and assume that $\mathcal{A}$ is positive definite. Then,

$$\text{tr } A^{-1} + \text{tr } C^{-1} \leq \text{tr } \mathcal{A}^{-1}.$$

Furthermore, B is nonzero if and only if

$$\text{tr } A^{-1} + \text{tr } C^{-1} < \text{tr } \mathcal{A}^{-1}.$$

(Proof: Use Proposition 8.2.5 or see [995].)

Fact 8.11.11. Let $A \in \mathbb{F}^{n \times m}$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} \langle A^* \rangle & A \\ A^* & \langle A \rangle \end{bmatrix}.$$

Then, $\mathcal{A}$ is positive semidefinite. If, in addition, $n = m$, then

$$-\langle A^* \rangle - \langle A \rangle \leq A + A^* \leq \langle A^* \rangle + \langle A \rangle.$$

(Proof: Use Fact 8.11.5.) (Remark: See Fact 8.9.29 and Fact 8.20.4.)

Fact 8.11.12. Let $A \in \mathbb{F}^{n \times n}$, assume that A is normal, and define

$$\mathcal{A} \triangleq \begin{bmatrix} \langle A \rangle & A \\ A^* & \langle A \rangle \end{bmatrix}.$$

Then, $\mathcal{A}$ is positive semidefinite. (Proof: See [711, p. 213].)

Fact 8.11.13. Let $A \in \mathbb{F}^{n \times n}$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} I & A \\ A^* & I \end{bmatrix}.$$

Then, $\mathcal{A}$ is (positive semidefinite, positive definite) if and only if A is (semicontractive, contractive).

Fact 8.11.14. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} A^*A & A^*B \\ B^*A & B^*B \end{bmatrix}.$$

Then, $\mathcal{A}$ is positive semidefinite, and

$$0 \leq A^*B(B^*B)^+B^*A \leq A^*A.$$

If $m = l$, then

$$-A^*A - B^*B \leq A^*B + B^*A \leq A^*A + B^*B.$$

If, in addition, $m = l = 1$ and $B^*B \neq 0$, then

$$|A^*B|^2 \leq A^*AB^*B.$$

(Remark: This result is the Cauchy-Schwarz inequality. See Fact 8.13.22.) (Remark: See Fact 8.21.41.)

Fact 8.11.15. Let $A, B \in \mathbb{F}^{n \times m}$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} I + A^*A & I - A^*B \\ I - B^*A & I + B^*B \end{bmatrix}$$

and

$$\mathcal{B} \triangleq \begin{bmatrix} I + A^*A & I + A^*B \\ I + B^*A & I + B^*B \end{bmatrix}.$$

Then, $\mathcal{A}$ and $\mathcal{B}$ are positive semidefinite,

$$0 \leq (I - A^*B)(I + B^*B)^{-1}(I - B^*A) \leq I + A^*A,$$

and

$$0 \leq (I + A^*B)(I + B^*B)^{-1}(I + B^*A) \leq I + A^*A.$$

(Remark: See Fact 8.13.25.)

Fact 8.11.16. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$I + AA^* = (A + B)(I + B^*B)^{-1}(A + B)^* + (I - AB^*)(I + BB^*)^{-1}(I - BA^*).$$

Therefore,

$$(A + B)(I + B^*B)^{-1}(A + B)^* \leq I + AA^*.$$

(Proof: Set $C = A$ in Fact 2.16.23. See also [1490, p. 185].)

Fact 8.11.17. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{n \times m}$, assume that A is positive semidefinite, and define

$$\mathcal{A} \triangleq \begin{bmatrix} A & AB \\ B^*A & B^*AB \end{bmatrix}.$$

Then,

$$\mathcal{A} = \begin{bmatrix} A^{1/2} \\ B^*A^{1/2} \end{bmatrix} \begin{bmatrix} A^{1/2} & A^{1/2}B \end{bmatrix},$$

and thus $\mathcal{A}$ is positive semidefinite. Furthermore,

$$0 \leq AB(B^*AB)^+B^*A \leq A.$$

Now, assume that $n = m$. Then,

$$-A - B^*AB \leq AB + B^*A \leq A + B^*AB.$$

Fact 8.11.18. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{n \times m}$, assume that A is positive definite, and define

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & B^*A^{-1}B \end{bmatrix}.$$

Then,

$$\mathcal{A} = \begin{bmatrix} A^{1/2} \\ B^*A^{-1/2} \end{bmatrix} \begin{bmatrix} A^{1/2} & A^{-1/2}B \end{bmatrix},$$

and thus $\mathcal{A}$ is positive semidefinite. Furthermore,

$$0 \leq B(B^*A^{-1}B)^+B^* \leq A.$$

Furthermore, if $\text{rank } B = m$, then

$$\text{rank}[A - B(B^*A^{-1}B)^{-1}B^*] = n - m.$$

Now, assume that $n = m$. Then,

$$-A - B^*A^{-1}B \leq B + B^* \leq A + B^*A^{-1}B.$$

(Proof: Use Fact 8.11.5.) (Remark: See Fact 8.21.42.) (Remark: The matrix $I - A^{-1/2}B(B^*A^{-1}B)^+B^*A^{-1/2}$ is a projector.)

Fact 8.11.19. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{n \times m}$, assume that A is positive definite, and define

$$\mathcal{A} \triangleq \begin{bmatrix} B^*AB & B^*B \\ B^*B & B^*A^{-1}B \end{bmatrix}.$$

Then,

$$\mathcal{A} = \begin{bmatrix} B^*A^{1/2} \\ B^*A^{-1/2} \end{bmatrix} \begin{bmatrix} A^{1/2}B & A^{-1/2}B \end{bmatrix},$$

and thus $\mathcal{A}$ is positive semidefinite. Furthermore,

$$0 \leq B^*B(B^*A^{-1}B)^+B^*B \leq B^*AB.$$

Now, assume that $n = m$. Then,

$$-B^*AB - B^*A^{-1}B \leq 2B^*B \leq B^*AB + B^*A^{-1}B.$$

(Proof: Use Fact 8.11.5.) (Remark: See Fact 8.13.23 and Fact 8.21.42.)

Fact 8.11.20. Let $A, B \in \mathbb{F}^{n \times m}$, let $\alpha, \beta \in (0, \infty)$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} \beta^{-1}I + \alpha A^*A & (A + B)^* \\ A + B & \alpha^{-1}I + \beta BB^* \end{bmatrix}.$$

Then,

$$\begin{aligned} \mathcal{A} &= \begin{bmatrix} \beta^{-1/2}I & \alpha^{1/2}A^* \\ \beta^{1/2}B & \alpha^{-1/2}I \end{bmatrix} \begin{bmatrix} \beta^{-1/2}I & \beta^{1/2}B^* \\ \alpha^{1/2}A & \alpha^{-1/2}I \end{bmatrix} \\ &= \begin{bmatrix} \alpha A^*A & A^* \\ A & \alpha^{-1}I \end{bmatrix} + \begin{bmatrix} \beta^{-1}I & B^* \\ B & \beta BB^* \end{bmatrix}, \end{aligned}$$

and thus $\mathcal{A}$ is positive semidefinite. Furthermore,

$$(A + B)^*(\alpha^{-1}I + \beta BB^*)^{-1}(A + B) \leq \beta^{-1}I + \alpha A^*A.$$

Now, assume that $n = m$. Then,

$$\begin{aligned} -(\beta^{-1/2} + \alpha^{-1/2})I - \alpha A^*A - \beta BB^* &\leq A + B + (A + B)^* \\ &\leq (\beta^{-1/2} + \alpha^{-1/2})I + \alpha A^*A + \beta BB^*. \end{aligned}$$

(Remark: See Fact 8.13.26 and Fact 8.21.43.)

Fact 8.11.21. Let $A, B \in \mathbb{F}^{n \times m}$, and assume that $I - A^*A$ and thus $I - AA^*$ are nonsingular. Then,

$$I - B^*B - (I - B^*A)(I - A^*A)^{-1}(I - A^*B) = -(A - B)^*(I - AA^*)^{-1}(A - B).$$

Now, assume that $I - A^*A$ is positive definite. Then,

$$I - B^*B \leq (I - B^*A)(I - A^*A)^{-1}(I - A^*B).$$

Now, assume that $I - B^*B$ is positive definite. Then, $I - A^*B$ is nonsingular. Next, define

$$\mathcal{A} \triangleq \begin{bmatrix} (I - A^*A)^{-1} & (I - B^*A)^{-1} \\ (I - A^*B)^{-1} & (I - B^*B)^{-1} \end{bmatrix}.$$

Then, $\mathcal{A}$ is positive semidefinite. Finally,

$$\begin{aligned} -(I - A^*A)^{-1} - (I - B^*B)^{-1} &\leq (I - B^*A)^{-1} + (I - A^*B)^{-1} \\ &\leq (I - A^*A)^{-1} + (I - B^*B)^{-1}. \end{aligned}$$

(Proof: For the first identity, set $D = -B^*$ and $C = -A^*$, and replace B with $-B$ in Fact 2.16.22. See [47, 1060]. The last statement follows from Fact 8.11.5.) (Remark: The identity is *Hua's matrix equality*. This result does not assume that either $I - A^*A$ or $I - B^*B$ is positive semidefinite. The inequality and Fact 8.13.25 constitute *Hua's inequalities*. See [1060, 1467].) (Remark: Extensions to the case

in which $I - A^*A$ is singular are considered in [1060].) (Remark: See Fact 8.9.39 and Fact 8.13.25.)

Fact 8.11.22. Let $A \in \mathbb{F}^{n \times n}$ be semicontractive, and define $B \in \mathbb{F}^{2n \times 2n}$ by

$$B \triangleq \begin{bmatrix} A & (I - AA^*)^{1/2} \\ (I - A^*A)^{1/2} & -A^* \end{bmatrix}.$$

Then, B is unitary. (Remark: See [508, p. 180].)

Fact 8.11.23. Let $A \in \mathbb{F}^{n \times m}$, and define $B \in \mathbb{F}^{(n+m) \times (n+m)}$ by

$$B \triangleq \begin{bmatrix} (I + A^*A)^{-1/2} & -A^*(I + AA^*)^{-1/2} \\ (I + AA^*)^{-1/2}A & (I + AA^*)^{-1/2} \end{bmatrix}.$$

Then, B is unitary and satisfies $A^* = \tilde{I}A\tilde{I}$, where $\tilde{I} \triangleq \text{diag}(I_m, -I_n)$. Furthermore, $\det B = 1$. (Remark: See [638].)

Fact 8.11.24. Let $A \in \mathbb{F}^{n \times m}$, assume that A is contractive, and define $B \in \mathbb{F}^{(n+m) \times (n+m)}$ by

$$B \triangleq \begin{bmatrix} (I - A^*A)^{-1/2} & A^*(I - AA^*)^{-1/2} \\ (I - AA^*)^{-1/2}A & (I - AA^*)^{-1/2} \end{bmatrix}.$$

Then, B is Hermitian and satisfies $A^*\tilde{I}A = \tilde{I}$, where $\tilde{I} \triangleq \text{diag}(I_m, -I_n)$. Furthermore, $\det B = 1$. (Remark: See [638].)

Fact 8.11.25. Let $X \in \mathbb{F}^{n \times m}$, and define $U \in \mathbb{F}^{(n+m) \times (n+m)}$ by

$$U \triangleq \begin{bmatrix} (I + X^*X)^{-1/2} & -X^*(I + XX^*)^{-1/2} \\ (I + XX^*)^{-1/2}X & (I + XX^*)^{-1/2} \end{bmatrix}.$$

Furthermore, let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, $D \in \mathbb{F}^{m \times m}$. Then, the following statements hold:

i) Assume that D is nonsingular, and let $X \triangleq D^{-1}C$. Then,

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} (A - BX)(I + X^*X)^{-1/2} & (B + AX^*)(I + XX^*)^{-1/2} \\ 0 & D(I + XX^*)^{1/2} \end{bmatrix} U.$$

ii) Assume that A is nonsingular and let $X \triangleq CA^{-1}$. Then,

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = U \begin{bmatrix} (I + X^*X)^{1/2}A & (I + X^*X)^{-1/2}(B + X^*D) \\ 0 & (I + XX^*)^{-1/2}(D - XB) \end{bmatrix}.$$

(Remark: See Proposition 2.8.3 and Proposition 2.8.4.) (Proof: See [638].)

Fact 8.11.26. Let $X \in \mathbb{F}^{n \times m}$, and define $U \in \mathbb{F}^{(n+m) \times (n+m)}$ by

$$U \triangleq \begin{bmatrix} (I - X^*X)^{-1/2} & X^*(I - XX^*)^{-1/2} \\ (I - XX^*)^{-1/2}X & (I - XX^*)^{-1/2} \end{bmatrix}.$$

Furthermore, let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{m \times n}$, $D \in \mathbb{F}^{m \times m}$. Then, the following statements hold:

- i) Assume that D is nonsingular, let $X \triangleq D^{-1}C$, and assume that $X^*X < I$. Then,

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = \begin{bmatrix} (A - BX)(I - X^*X)^{-1/2} & (B + AX^*)(I - XX^*)^{-1/2} \\ 0 & D(I - XX^*)^{1/2} \end{bmatrix} U.$$

- ii) Assume that A is nonsingular, let $X \triangleq CA^{-1}$, and assume that $X^*X < I$. Then,

$$\begin{bmatrix} A & B \\ C & D \end{bmatrix} = U \begin{bmatrix} (I - X^*X)^{1/2}A & (I - X^*X)^{-1/2}(B - X^*D) \\ 0 & (I - XX^*)^{-1/2}(D - XB) \end{bmatrix}.$$

(Proof: See [638].) (Remark: See Proposition 2.8.3 and Proposition 2.8.4.)

Fact 8.11.27. Let $A, B \in \mathbb{F}^{n \times m}$ and $C, D \in \mathbb{F}^{m \times m}$, assume that C and D are positive definite, and define

$$\mathcal{A} \triangleq \begin{bmatrix} AC^{-1}A^* + BD^{-1}B^* & A + B \\ (A + B)^* & C + D \end{bmatrix}.$$

Then, $\mathcal{A}$ is positive semidefinite, and

$$(A + B)(C + D)^{-1}(A + B)^* \leq AC^{-1}A^* + BD^{-1}B^*.$$

Now, assume that $n = m$. Then,

$$\begin{aligned} -AC^{-1}A^* - BD^{-1}B^* - C - D &\leq A + B + (A + B)^* \\ &\leq AC^{-1}A^* + BD^{-1}B^* + C + D. \end{aligned}$$

(Proof: See [658, 907] or [1098, p. 151].) (Remark: Replacing A, B, C, D by $\alpha B_1, (1 - \alpha)B_2, \alpha A_1, (1 - \alpha)A_2$ yields *xiv*) of Proposition 8.6.17.)

Fact 8.11.28. Let $A \in \mathbb{R}^{n \times n}$, assume that A is positive definite, and let $\mathcal{S} \subseteq \{1, \dots, n\}$. Then,

$$(A_{(\mathcal{S})})^{-1} \leq (A^{-1})_{(\mathcal{S})}.$$

(Proof: See [709, p. 474].) (Remark: Generalizations of this result are given in [328].)

Fact 8.11.29. Let $A_{ij} \in \mathbb{F}^{n_i \times n_j}$ for all $i, j = 1, \dots, k$, define

$$A \triangleq \begin{bmatrix} A_{11} & \cdots & A_{1k} \\ \vdots & \ddots & \vdots \\ A_{1k} & \cdots & A_{kk} \end{bmatrix},$$

and assume that A is square and positive definite. Furthermore, define

$$\hat{A} \triangleq \begin{bmatrix} \hat{A}_{11} & \cdots & \hat{A}_{1k} \\ \vdots & \ddots & \vdots \\ \hat{A}_{1k} & \cdots & \hat{A}_{kk} \end{bmatrix},$$

where $\hat{A}_{ij} = 1_{1 \times n_i} A_{ij} 1_{n_j \times 1}$ is the sum of the entries of A_{ij} for all $i, j = 1, \dots, k$. Then, $\hat{A}$ is positive definite. (Proof: $\hat{A} = BAB^T$, where the entries of $B \in \mathbb{R}^{k \times \sum_{i=1}^k n_i}$ are 0's and 1's. See [42].)

Fact 8.11.30. Let $A, D \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and assume that $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{n \times n}$ is positive semidefinite, C is positive definite, and D is positive definite. Then, $\begin{bmatrix} A+D & B \\ B^* & C \end{bmatrix}$ is positive definite.

Fact 8.11.31. Let $A \in \mathbb{F}^{(n+m+l) \times (n+m+l)}$, assume that A is positive semidefinite, and assume that A is of the form

$$A = \begin{bmatrix} A_{11} & A_{12} & 0 \\ A_{12}^* & A_{22} & A_{23} \\ 0 & A_{32}^* & A_{33} \end{bmatrix}.$$

Then, there exist positive-semidefinite matrices $B, C \in \mathbb{F}^{(n+m+l) \times (n+m+l)}$ such that $A = B + C$ and such that B and C have the form

$$B = \begin{bmatrix} B_{11} & B_{12} & 0 \\ B_{12}^* & B_{22} & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

and

$$C = \begin{bmatrix} 0 & 0 & 0 \\ 0 & C_{22} & C_{23} \\ 0 & C_{23}^* & C_{33} \end{bmatrix}.$$

(Proof: See [669].)

8.12 Facts on the Trace

Fact 8.12.1. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, let p and q be real numbers, and assume that $p \leq q$. Then,

$$\left(\frac{1}{n} \operatorname{tr} A^p\right)^{1/p} \leq \left(\frac{1}{n} \operatorname{tr} A^q\right)^{1/q}.$$

Furthermore,

$$\lim_{p \downarrow 0} \left(\frac{1}{n} \operatorname{tr} A^p\right)^{1/p} = \det A^{1/n}.$$

(Proof: Use Fact 1.15.30.)

Fact 8.12.2. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$n^2 \leq (\operatorname{tr} A) \operatorname{tr} A^{-1}.$$

Finally, equality holds if and only if $A = I_n$. (Remark: Bounds on $\text{tr } A^{-1}$ are given in [100, 307, 1052, 1132].)

Fact 8.12.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, the following statements hold:

i) Let $r \in [0, 1]$. Then, for all $k = 1, \dots, n$,

$$\sum_{i=k}^n \lambda_i^r(A) \leq \sum_{i=k}^n d_i^r(A).$$

In particular,

$$\text{tr } A^r \leq \sum_{i=1}^n A_{(i,i)}^r.$$

ii) Let $r \geq 1$. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k d_i^r(A) \leq \sum_{i=1}^k \lambda_i^r(A).$$

In particular,

$$\sum_{i=1}^n A_{(i,i)}^r \leq \text{tr } A^r.$$

iii) If either $r = 0$ or $r = 1$, then

$$\text{tr } A^r = \sum_{i=1}^n A_{(i,i)}^r.$$

iv) If $r \neq 0$ and $r \neq 1$, then

$$\text{tr } A^r = \sum_{i=1}^n A_{(i,i)}^r$$

if and only if A is diagonal.

(Proof: Use Fact 8.17.8 and Fact 2.21.8. See [946] and [948, p. 217].) (Remark: See Fact 8.17.8.)

Fact 8.12.4. Let $A \in \mathbb{F}^{n \times n}$, and let $p, q \in [0, \infty)$. Then,

$$\text{tr } (A^* p A^p)^q \leq \text{tr } (A^* A)^{pq}.$$

Furthermore, equality holds if and only if $\text{tr } A^* p A^p = \text{tr } (A^* A)^p$. (Proof: See [1208].)

Fact 8.12.5. Let $A \in \mathbb{F}^{n \times n}$, $p \in [2, \infty)$, and $q \in [1, \infty)$. Then, A is normal if and only if

$$\text{tr } (A^* p A^p)^q = \text{tr } (A^* A)^{pq}.$$

(Proof: See [1208].)

Fact 8.12.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that either A and B are Hermitian or A and B are skew Hermitian. Then, $\text{tr } AB$ is real. (Proof: $\text{tr } AB = \text{tr } A^* B^* = \text{tr } (BA)^* = \overline{\text{tr } BA} = \overline{\text{tr } AB}$. (Remark: See [1476] or [1490, p. 213].)

Fact 8.12.7. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and let $k \in \mathbb{N}$. Then, $\operatorname{tr} (AB)^k$ is real. (Proof: See [55].)

Fact 8.12.8. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then,

$$\operatorname{tr} AB \leq |\operatorname{tr} AB| \leq \sqrt{(\operatorname{tr} A^2) \operatorname{tr} B^2} \leq \frac{1}{2} \operatorname{tr} (A^2 + B^2).$$

The second inequality is an equality if and only if A and B are linearly dependent. The third inequality is an equality if and only if $\operatorname{tr} A^2 = \operatorname{tr} B^2$. All four terms are equal if and only if $A = B$. (Proof: Use the Cauchy-Schwarz inequality Corollary 9.3.9.) (Remark: See Fact 8.12.18.)

Fact 8.12.9. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that $-A \leq B \leq A$. Then,

$$\operatorname{tr} B^2 \leq \operatorname{tr} A^2.$$

(Proof: $0 \leq \operatorname{tr} [(A - B)(A + B)] = \operatorname{tr} A^2 - \operatorname{tr} B^2$. See [1318].) (Remark: For $0 \leq B \leq A$, this result is a special case of xxi) of Proposition 8.6.13.

Fact 8.12.10. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, $AB = 0$ if and only if $\operatorname{tr} AB = 0$.

Fact 8.12.11. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p, q \geq 1$ satisfy $1/p + 1/q = 1$. Then,

$$\operatorname{tr} AB \leq \operatorname{tr} \langle AB \rangle \leq (\operatorname{tr} A^p)^{1/p} (\operatorname{tr} B^q)^{1/q}.$$

Furthermore, equality holds for both inequalities if and only if A^{p-1} and B are linearly dependent. (Proof: See [946] and [948, pp. 219, 222].) (Remark: This result is a matrix version of Hölder's inequality.) (Remark: See Fact 8.12.12 and Fact 8.12.17.)

Fact 8.12.12. Let $A_1, \dots, A_m \in \mathbb{F}^{n \times n}$, assume that $A_1, \dots, A_m$ are positive semidefinite, and let $p_1, \dots, p_m \in [1, \infty)$ satisfy $\frac{1}{p_1} + \dots + \frac{1}{p_m} = 1$. Then,

$$\operatorname{tr} \langle A_1 \cdots A_m \rangle \leq \prod_{i=1}^m (\operatorname{tr} A_i^{p_i})^{1/p_i} \leq \operatorname{tr} \sum_{i=1}^m \frac{1}{p_i} A_i^{p_i}.$$

Furthermore, the following statements are equivalent:

- i) $\operatorname{tr} \langle A_1 \cdots A_m \rangle = \prod_{i=1}^m (\operatorname{tr} A_i^{p_i})^{1/p_i}$.
- ii) $\operatorname{tr} \langle A_1 \cdots A_m \rangle = \operatorname{tr} \sum_{i=1}^m \frac{1}{p_i} A_i^{p_i}$.
- iii) $A_1^{p_1} = \dots = A_m^{p_m}$.

(Proof: See [954].) (Remark: The first inequality is a matrix version of Hölder's inequality. The inequality involving the first and third terms is a matrix version of Young's inequality. See Fact 1.10.32 and Fact 1.15.31.)

Fact 8.12.13. Let $A_1, \dots, A_m \in \mathbb{F}^{n \times n}$, assume that $A_1, \dots, A_m$ are positive semidefinite, let $\alpha_1, \dots, \alpha_m$ be nonnegative numbers, and assume that $\sum_{i=1}^m \alpha_i \geq 1$.

Then,

$$\left| \operatorname{tr} \prod_{i=1}^m A_i^{\alpha_i} \right| \leq \prod_{i=1}^m (\operatorname{tr} A_i)^{\alpha_i}.$$

Furthermore, if $\sum_{i=1}^m \alpha_i = 1$, then equality holds if and only if $A_2, \dots, A_m$ are scalar multiples of A_1 , whereas, if $\sum_{i=1}^m \alpha_i > 1$, then equality holds if and only if $A_2, \dots, A_m$ are scalar multiples of A_1 and $\operatorname{rank} A_1 = 1$. (Proof: See [317].) (Remark: See Fact 8.12.11.)

Fact 8.12.14. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$|\operatorname{tr} AB|^2 \leq (\operatorname{tr} A^*A) \operatorname{tr} BB^*.$$

(Proof: See [1490, p. 25] or Corollary 9.3.9.) (Remark: See Fact 8.12.15.)

Fact 8.12.15. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$, and let $k \in \mathbb{N}$. Then,

$$|\operatorname{tr} (AB)^{2k}| \leq \operatorname{tr} (A^*ABB^*)^k \leq \operatorname{tr} (A^*A)^k (\operatorname{tr} BB^*)^k \leq [\operatorname{tr} (A^*A)^k] \operatorname{tr} (BB^*)^k.$$

In particular,

$$|\operatorname{tr} (AB)^2| \leq \operatorname{tr} A^*ABB^* \leq (\operatorname{tr} A^*A) \operatorname{tr} BB^*.$$

(Proof: See [1476] for the case $n = m$. If $n \neq m$, then A and B can be augmented with 0's.) (Problem: Show that

$$\left. \begin{array}{l} |\operatorname{tr} AB|^2 \\ |\operatorname{tr} (AB)^2| \end{array} \right\} \leq \operatorname{tr} A^*ABB^* \leq (\operatorname{tr} A^*A) \operatorname{tr} BB^*.$$

See Fact 8.12.14.)

Fact 8.12.16. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and let $k \geq 1$. Then,

$$\operatorname{tr} (A^2B^2)^k \leq (\operatorname{tr} A^2B^2)^k$$

and

$$\operatorname{tr} (AB)^{2k} \leq |\operatorname{tr} (AB)^{2k}| \leq \left\{ \begin{array}{l} \operatorname{tr} (A^2B^2)^k \\ \operatorname{tr} ((AB)^{2k}) \end{array} \right\} \leq \operatorname{tr} A^{2k}B^{2k}.$$

(Proof: Use Fact 8.12.15 and see [55, 1476].) (Remark: It follows from Fact 8.12.7 that $\operatorname{tr} (AB)^{2k}$ and $\operatorname{tr} (A^2B^2)^k$ are real.)

Fact 8.12.17. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\operatorname{tr} AB \leq \operatorname{tr} (AB^2A)^{1/2} = \operatorname{tr} \langle AB \rangle \leq \frac{1}{4} \operatorname{tr} (A+B)^2$$

and

$$\operatorname{tr} (AB)^2 \leq \operatorname{tr} A^2B^2 \leq \frac{1}{16} \operatorname{tr} (A+B)^4.$$

(Proof: See Fact 8.12.20 and Fact 9.9.18.)

Fact 8.12.18. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\begin{aligned}
 \operatorname{tr} AB &= \operatorname{tr} A^{1/2} B A^{1/2} \\
 &= \operatorname{tr} \left[\left(A^{1/2} B A^{1/2} \right)^{1/2} \left(A^{1/2} B A^{1/2} \right)^{1/2} \right] \\
 &\leq \left[\operatorname{tr} \left(A^{1/2} B A^{1/2} \right)^{1/2} \right]^2 \\
 &\leq (\operatorname{tr} A)(\operatorname{tr} B) \\
 &\leq \frac{1}{4} (\operatorname{tr} A + \operatorname{tr} B)^2 \\
 &\leq \frac{1}{2} [(\operatorname{tr} A)^2 + (\operatorname{tr} B)^2]
 \end{aligned}$$

and

$$\begin{aligned}
 \operatorname{tr} AB &\leq \sqrt{\operatorname{tr} A^2} \sqrt{\operatorname{tr} B^2} \\
 &\leq \frac{1}{4} \left(\sqrt{\operatorname{tr} A^2} + \sqrt{\operatorname{tr} B^2} \right)^2 \\
 &\leq \frac{1}{2} (\operatorname{tr} A^2 + \operatorname{tr} B^2) \\
 &\leq \frac{1}{2} [(\operatorname{tr} A)^2 + (\operatorname{tr} B)^2].
 \end{aligned}$$

(Remark: Use Fact 1.10.4.) (Remark: Note that

$$\operatorname{tr} \left(A^{1/2} B A^{1/2} \right)^{1/2} = \sum_{i=1}^n \lambda_i^{1/2}(AB).$$

The second inequality follows from Proposition 9.3.6 with $p = q = 2$, $r = 1$, and A and B replaced by $A^{1/2}$ and $B^{1/2}$.) (Remark: See Fact 2.12.16.)

Fact 8.12.19. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p \geq 1$. Then,

$$\operatorname{tr} AB \leq \operatorname{tr} (A^{p/2} B^p A^{p/2})^{1/p}.$$

(Proof: See [521].)

Fact 8.12.20. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p \geq 0$ and $r \geq 1$. Then,

$$\operatorname{tr} \left(A^{1/2} B A^{1/2} \right)^{pr} \leq \operatorname{tr} \left(A^{r/2} B^r A^{r/2} \right)^p.$$

In particular,

$$\operatorname{tr} \left(A^{1/2} B A^{1/2} \right)^{2p} \leq \operatorname{tr} (AB^2 A)^p$$

and

$$\operatorname{tr} AB \leq \operatorname{tr} (AB^2 A)^{1/2} = \operatorname{tr} \langle AB \rangle.$$

(Proof: Use Fact 8.18.20 and Fact 8.18.27.) (Remark: This result is the *Araki-Lieb-Thirring inequality*. See [69, 88] and [197, p. 258]. See Fact 8.10.49, Fact 8.18.26,

and Fact 9.9.17.) (Problem: Referring to Fact 8.12.18, compare the upper bounds

$$\operatorname{tr} AB \leq \begin{cases} \left[\operatorname{tr} (A^{1/2} B A^{1/2})^{1/2} \right]^2 \\ \sqrt{\operatorname{tr} A^2} \sqrt{\operatorname{tr} B^2} \\ \operatorname{tr} (AB^2 A)^{1/2}. \end{cases}$$

Fact 8.12.21. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $q \geq 0$ and $t \in [0, 1]$. Then,

$$\sigma_{\max}^{2tq}(A) \operatorname{tr} B^{tq} \leq \operatorname{tr} (A^t B^t A^t)^q \leq \operatorname{tr} (ABA)^{tq}.$$

(Proof: See [88].) (Remark: The right-hand inequality is equivalent to the Araki-Lieb-Thirring inequality, where $t = 1/r$ and $q = pr$. See Fact 8.12.20.)

Fact 8.12.22. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $k, m \in \mathbb{P}$, where $m \geq k$. Then,

$$\operatorname{tr} (A^k B^k)^m \leq \operatorname{tr} (A^m B^m)^k.$$

In particular,

$$\operatorname{tr} (AB)^m \leq \operatorname{tr} A^m B^m.$$

If, in addition, m is even, then

$$\operatorname{tr} (AB)^m \leq \operatorname{tr} (A^2 B^2)^{m/2} \leq \operatorname{tr} A^m B^m.$$

(Proof: Use Fact 8.18.20 and Fact 8.18.27.) (Remark: It follows from Fact 8.12.7 that $\operatorname{tr} (AB)^m$ is real.) (Remark: The result $\operatorname{tr} (AB)^m \leq \operatorname{tr} A^m B^m$ is the *Lieb-Thirring inequality*. See [197, p. 279]. The inequality $\operatorname{tr} (AB)^m \leq \operatorname{tr} (A^2 B^2)^{m/2}$ follows from Fact 8.12.20. See [1466, 1476].)

Fact 8.12.23. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p \geq r \geq 0$. Then,

$$\left[\operatorname{tr} (A^{1/2} B A^{1/2})^p \right]^{1/p} \leq \left[\operatorname{tr} (A^{1/2} B A^{1/2})^r \right]^{1/r}.$$

In particular,

$$\left[\operatorname{tr} (A^{1/2} B A^{1/2})^2 \right]^{1/2} \leq \operatorname{tr} AB \leq \begin{cases} \operatorname{tr} (AB^2 A)^{1/2} \\ \left[\operatorname{tr} (A^{1/2} B A^{1/2})^{1/2} \right]^2. \end{cases}$$

(Proof: The result follows from the power-sum inequality Fact 1.15.34. See [369].)

Fact 8.12.24. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, assume that $A \leq B$, and let $p, q \geq 0$. Then,

$$\operatorname{tr} A^p B^q \leq \operatorname{tr} B^{p+q}.$$

If, in addition, A and B are positive definite, then this inequality holds for all $p, q \in \mathbb{R}$ satisfying $q \geq -1$ and $p + q \geq 0$. (Proof: See [246].)

Fact 8.12.25. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, assume that $A \leq B$, let $f: [0, \infty) \mapsto [0, \infty)$, and assume that $f(0) = 0$, f is continuous, and f is increasing. Then,

$$\operatorname{tr} f(A) \leq \operatorname{tr} f(B).$$

Now, let $p > 1$ and $q \geq \max\{-1, -p/2\}$, and, if $q < 0$, assume that A is positive definite. Then,

$$\operatorname{tr} f(A^{q/2} B^p A^{q/2}) \leq \operatorname{tr} f(A^{p+q}).$$

(Proof: See [527].)

Fact 8.12.26. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\alpha \in [0, 1]$. Then,

$$\operatorname{tr} A^\alpha B^{1-\alpha} \leq (\operatorname{tr} A)^\alpha (\operatorname{tr} B)^{1-\alpha} \leq \operatorname{tr} [\alpha A + (1 - \alpha) B].$$

Furthermore, the first inequality is an equality if and only if A and B are linearly dependent, while the second inequality is an equality if and only if $A = B$. (Proof: Use Fact 8.12.11 or Fact 8.12.13 for the left-hand inequality and Fact 1.10.21 for the right-hand inequality.)

Fact 8.12.27. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $\alpha \in [0, 1]$. Then,

$$\left. \begin{array}{l} \operatorname{tr} A^{-\alpha} B^{\alpha-1} \\ \operatorname{tr} [\alpha A + (1 - \alpha) B]^{-1} \end{array} \right\} \leq (\operatorname{tr} A^{-1})^\alpha (\operatorname{tr} B^{-1})^{1-\alpha} \leq \operatorname{tr} [\alpha A^{-1} + (1 - \alpha) B^{-1}]$$

and

$$\operatorname{tr} [\alpha A + (1 - \alpha) B]^{-1} \leq \left\{ \begin{array}{l} (\operatorname{tr} A^{-1})^\alpha (\operatorname{tr} B^{-1})^{1-\alpha} \\ \operatorname{tr} [A^{-1} (A^{-1/2} B A^{-1/2})^{\alpha-1}] \end{array} \right\} \leq \operatorname{tr} [\alpha A^{-1} + (1 - \alpha) B^{-1}].$$

(Remark: In the first string of inequalities, the upper left inequality and right-hand inequality are equivalent to Fact 8.12.26. The lower left inequality is given by *xxviii*) of Proposition 8.6.17. The second string of inequalities combines the lower left inequality in the first string of inequalities with the third string of inequalities in Fact 8.10.46.) (Remark: These inequalities interpolate the convexity of $\phi(A) = \operatorname{tr} A^{-1}$. See Fact 1.10.21.)

Fact 8.12.28. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that B is positive semidefinite. Then,

$$|\operatorname{tr} AB| \leq \sigma_{\max}(A) \operatorname{tr} B.$$

(Proof: Use Proposition 8.4.13 and $\sigma_{\max}(A + A^*) \leq 2\sigma_{\max}(A)$.) (Remark: See Fact 5.12.4.)

Fact 8.12.29. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p \geq 1$. Then,

$$\operatorname{tr}(A^p + B^p) \leq \operatorname{tr}(A + B)^p \leq \left[(\operatorname{tr} A^p)^{1/p} + (\operatorname{tr} B^p)^{1/p} \right]^p.$$

Furthermore, the second inequality is an equality if and only if A and B are linearly independent. (Proof: See [246] and [946].) (Remark: The first inequality is the *Mc-*

Carthy inequality. The second inequality is a special case of the triangle inequality for the norm $\|\cdot\|_{\sigma p}$ and a matrix version of Minkowski's inequality.)

Fact 8.12.30. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let m be a positive integer, and define $p \in \mathbb{F}[s]$ by

$$p(s) = \operatorname{tr}(A + sB)^m.$$

Then, all of the coefficients of p are nonnegative. (Remark: This result is the *Bessis-Moussa-Villani trace conjecture*. See [687, 908] and Fact 8.12.31.)

Fact 8.12.31. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is Hermitian and B is positive semidefinite, and define

$$f(t) = e^{A+tB}.$$

Then, for all $k = 0, 1, \dots$ and $t \geq 0$,

$$(-1)^{k+1} f^{(k)}(t) \geq 0.$$

(Remark: This result is a consequence of the Bessis-Moussa-Villani trace conjecture. See [687, 908] and Fact 8.12.30.) (Remark: See Fact 8.14.18.)

Fact 8.12.32. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and let $f: \mathbb{R} \mapsto \mathbb{R}$. Then, the following statements hold:

i) If f is convex, then there exist unitary matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that

$$f\left[\frac{1}{2}(A+B)\right] \leq \frac{1}{2}[S_1\left(\frac{1}{2}[f(A)+f(B)]\right)S_1^* + S_2\left(\frac{1}{2}[f(A)+f(B)]\right)S_2^*].$$

ii) If f is convex and even, then there exist unitary matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that

$$f\left[\frac{1}{2}(A+B)\right] \leq \frac{1}{2}[S_1 f(A) S_1^* + S_2 f(B) S_2^*].$$

iii) If f is convex and increasing, then there exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that

$$f\left[\frac{1}{2}(A+B)\right] \leq S\left(\frac{1}{2}[f(A)+f(B)]\right)S^*.$$

iv) There exist unitary matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that

$$\langle A+B \rangle \leq S_1 \langle A \rangle S_1^* + S_2 \langle B \rangle S_2^*.$$

v) If f is convex, then

$$\operatorname{tr} f\left[\frac{1}{2}(A+B)\right] \leq \operatorname{tr} \frac{1}{2}[f(A)+f(B)].$$

(Proof: See [247, 248].) (Remark: Result v), which is a consequence of i), is *von Neumann's trace inequality*.) (Remark: See Fact 8.12.33.)

Fact 8.12.33. Let $f: \mathbb{R} \mapsto \mathbb{R}$, and assume that f is convex. Then, the following statements hold:

i) If $f(0) \leq 0$, $A \in \mathbb{F}^{n \times n}$ is Hermitian, and $S \in \mathbb{F}^{n \times m}$ is a contractive matrix, then

$$\operatorname{tr} f(S^*AS) \leq \operatorname{tr} S^*f(A)S.$$

ii) If $A_1, \dots, A_k \in \mathbb{F}^{n \times n}$ are Hermitian and $S_1, \dots, S_k \in \mathbb{F}^{n \times m}$ satisfy $\sum_{i=1}^k S_i^* S_i = I$, then

$$\operatorname{tr} f\left(\sum_{i=1}^k S_i^* A_i S_i\right) \leq \operatorname{tr} \sum_{i=1}^k S_i^* f(A_i) S_i.$$

iii) If $A \in \mathbb{F}^{n \times n}$ is Hermitian and $S \in \mathbb{F}^{n \times n}$ is a projector, then

$$\operatorname{tr} Sf(SAS)S \leq \operatorname{tr} Sf(A)S.$$

(Proof: See [248] and [1039, p. 36].) (Remark: Special cases are considered in [785].) (Remark: The first result is due to Brown and Kosaki, the second result is due to Hansen and Pedersen, and the third result is due to Berezin.) (Remark: The second result generalizes statement *v*) of Fact 8.12.32.)

Fact 8.12.34. Let $A, B \in \mathbb{F}^{n \times n}$, assume that B is positive semidefinite, and assume that $A^*A \leq B$. Then,

$$|\operatorname{tr} A| \leq \operatorname{tr} B^{1/2}.$$

(Proof: Corollary 8.6.11 with $r = 2$ implies that $(A^*A)^{1/2} \leq \operatorname{tr} B^{1/2}$. Letting $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, it follows from Fact 9.11.2 that $|\operatorname{tr} A| \leq \sum_{i=1}^n |\lambda_i| \leq \sum_{i=1}^n \sigma_i(A) = \operatorname{tr} (A^*A)^{1/2} \leq \operatorname{tr} B^{1/2}$. See [167].)

Fact 8.12.35. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive definite and B is positive semidefinite, let $\alpha \in [0, 1]$, and let $\beta \geq 0$. Then,

$$\operatorname{tr}(-BA^{-1}B + \beta B^\alpha) \leq \beta(1 - \frac{\alpha}{2}) \operatorname{tr} \left(\frac{\alpha\beta}{2} A \right)^{\alpha/(2-\alpha)}.$$

If, in addition, either A and B commute or B is a multiple of a projector, then

$$-BA^{-1}B + \beta B^\alpha \leq \beta(1 - \frac{\alpha}{2}) \left(\frac{\alpha\beta}{2} A \right)^{\alpha/(2-\alpha)}.$$

(Proof: See [634, 635].)

Fact 8.12.36. Let $A, P \in \mathbb{F}^{n \times n}$, $B, Q \in \mathbb{F}^{n \times m}$, and $C, R \in \mathbb{F}^{m \times m}$, and assume that $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix}, \begin{bmatrix} P & Q \\ Q^* & R \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ are positive semidefinite. Then,

$$|\operatorname{tr} BQ^*|^2 \leq (\operatorname{tr} AP)(\operatorname{tr} CR).$$

(Proof: See [886, 1494].)

Fact 8.12.37. Let $A, B \in \mathbb{F}^{n \times m}$, let $X \in \mathbb{F}^{n \times n}$, and assume that X is positive definite. Then,

$$|\operatorname{tr} A^*B|^2 \leq (\operatorname{tr} A^*XA)(\operatorname{tr} B^*X^{-1}A).$$

(Proof: Use Fact 8.12.36 with $\begin{bmatrix} X & I \\ I & X^{-1} \end{bmatrix}$ and $\begin{bmatrix} AA^* & AB^* \\ BA^* & BB^* \end{bmatrix}$. See [886, 1494].)

Fact 8.12.38. Let $A, B, C \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian and C is positive semidefinite. Then,

$$|\operatorname{tr} ABC^2 - \operatorname{tr} ACBC| \leq \frac{1}{4}[\lambda_1(A) - \lambda_n(A)][\lambda_1(B) - \lambda_n(B)] \operatorname{tr} C^2.$$

(Proof: See [250].)

Fact 8.12.39. Let $A_{11} \in \mathbb{R}^{n \times n}$, $A_{12} \in \mathbb{R}^{n \times m}$, and $A_{22} \in \mathbb{R}^{m \times m}$, define $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^T & A_{22} \end{bmatrix} \in \mathbb{R}^{(n+m) \times (n+m)}$, and assume that A is symmetric. Then, A is positive semidefinite if and only if, for all $B \in \mathbb{R}^{n \times m}$,

$$\operatorname{tr} BA_{12}^T \leq \operatorname{tr} \left(A_{11}^{1/2} B A_{22} B^T A_{11}^{1/2} \right)^{1/2}.$$

(Proof: See [167].)

Fact 8.12.40. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and assume that $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ is positive semidefinite. Then,

$$\operatorname{tr} B^* B \leq \sqrt{(\operatorname{tr} A^2)(\operatorname{tr} C^2)} \leq (\operatorname{tr} A)(\operatorname{tr} C).$$

(Proof: Use Fact 8.12.36 with $P = A$, $Q = B$, and $R = C$.) (Remark: The inequality involving the first and third terms is given in [1075].) (Remark: See Fact 8.12.41 for the case $n = m$.)

Fact 8.12.41. Let $A, B, C \in \mathbb{F}^{n \times n}$, and assume that $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{2n \times 2n}$ is positive semidefinite. Then,

$$|\operatorname{tr} B|^2 \leq (\operatorname{tr} A)(\operatorname{tr} C)$$

and

$$|\operatorname{tr} B^2| \leq \operatorname{tr} B^* B \leq \sqrt{(\operatorname{tr} A^2)(\operatorname{tr} C^2)} \leq (\operatorname{tr} A)(\operatorname{tr} C).$$

(Remark: The first result follows from Fact 8.12.42. In the second string, the first inequality is given by Fact 9.11.3, while the second inequality is given by Fact 8.12.40. The inequality $|\operatorname{tr} B^2| \leq \sqrt{(\operatorname{tr} A^2)(\operatorname{tr} C^2)}$ is given in [964].)

Fact 8.12.42. Let $A_{ij} \in \mathbb{F}^{n \times n}$ for all $i, j = 1, \dots, k$, define $A \in \mathbb{F}^{kn \times kn}$ by

$$A \triangleq \begin{bmatrix} A_{11} & \cdots & A_{1k} \\ \vdots & \ddots & \vdots \\ A_{1k}^* & \cdots & A_{kk} \end{bmatrix},$$

and assume that A is positive semidefinite. Then,

$$\begin{bmatrix} \operatorname{tr} A_{11} & \cdots & \operatorname{tr} A_{1k} \\ \vdots & \ddots & \vdots \\ \operatorname{tr} A_{1k}^* & \cdots & \operatorname{tr} A_{kk} \end{bmatrix} \geq 0$$

and

$$\begin{bmatrix} \operatorname{tr} A_{11}^2 & \cdots & \operatorname{tr} A_{1k}^* A_{1k} \\ \vdots & \ddots & \vdots \\ \operatorname{tr} A_{1k}^* A_{1k} & \cdots & \operatorname{tr} A_{kk}^2 \end{bmatrix} \geq 0.$$

(Proof: See [386, 964, 1075].) (Remark: See Fact 8.13.42.)

8.13 Facts on the Determinant

Fact 8.13.1. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\begin{aligned} \lambda_{\min}(A) &\leq \lambda_{\max}^{1/n}(A) \lambda_{\min}^{(n-1)/n}(A) \\ &\leq \lambda_n \\ &\leq \lambda_1 \\ &\leq \lambda_{\min}^{1/n}(A) \lambda_{\max}^{(n-1)/n}(A) \\ &\leq \lambda_{\max}(A) \end{aligned}$$

and

$$\begin{aligned} \lambda_{\min}^n(A) &\leq \lambda_{\max}(A) \lambda_{\min}^{n-1}(A) \\ &\leq \det A \\ &\leq \lambda_{\min}(A) \lambda_{\max}^{n-1}(A) \\ &\leq \lambda_{\max}^n(A). \end{aligned}$$

(Proof: Use Fact 5.11.29.)

Fact 8.13.2. Let $A \in \mathbb{F}^{n \times n}$, and assume that $A + A^*$ is positive semidefinite. Then,

$$\det \frac{1}{2}(A + A^*) \leq |\det A|.$$

Furthermore, if $A + A^*$ is positive definite, then equality holds if and only if A is Hermitian. (Proof: The inequality follows from Fact 5.11.25 and Fact 5.11.28.) (Remark: This result is the *Ostrowski-Taussky inequality*.) (Remark: See Fact 8.13.2.)

Fact 8.13.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that $A + A^*$ is positive semidefinite. Then,

$$[\det \frac{1}{2}(A + A^*)]^{2/n} + |\det \frac{1}{2}(A - A^*)|^{2/n} \leq |\det A|^{2/n}.$$

Furthermore, if $A + A^*$ is positive definite, then equality holds if and only if every eigenvalue of $(A + A^*)^{-1}(A - A^*)$ has the same absolute value. Finally, if $n \geq 2$, then

$$\det \frac{1}{2}(A + A^*) \leq \det \frac{1}{2}(A + A^*) + |\det \frac{1}{2}(A - A^*)| \leq |\det A|.$$

(Proof: See [466, 760]. To prove the last result, use Fact 1.10.30.) (Remark: Setting $A = 1 + j$ shows that the last result can fail for $n = 1$.) (Remark: $-A$ is semidissipative.) (Remark: The last result interpolates Fact 8.13.2.) (Remark: Extensions to the case in which $A + A^*$ is positive definite are considered in [1269].)

Fact 8.13.4. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then,

$$(\det A)^{2/n} + |\det(A + B)|^{2/n} \leq |\det(A + B)|^{2/n}.$$

Furthermore, if A is positive definite, then equality holds if and only if every eigenvalue of $A^{-1}B$ has the same absolute value. Finally, if $n \geq 2$, then

$$\det A \leq \det A + |\det B| \leq |\det(A + B)|.$$

(Remark: This result is a restatement of Fact 8.13.2 in terms of the Cartesian decomposition.)

Fact 8.13.5. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, assume that B is positive definite. Then,

$$\prod_{i=1}^n [\lambda_i^2(A) + \lambda_i^2(B)]^{1/2} \leq |\det(A + jB)| \leq \prod_{i=1}^n [\lambda_i^2(A) + \lambda_{n-i+1}^2(B)]^{1/2}.$$

(Proof: See [158].)

Fact 8.13.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite and B is skew Hermitian. Then,

$$\det A \leq |\det(A + B)|.$$

Furthermore, if A and B are real, then

$$\det A \leq \det(A + B).$$

Finally, if A is positive definite, then equality holds if and only if $B = 0$. (Proof: See [654, p. 447] and [1098, pp. 146, 163]. Now, suppose that A and B are real. If A is positive definite, then $A^{-1/2}BA^{-1/2}$ is skew symmetric, and thus $\det(A + B) = (\det A)\det(I + A^{-1/2}BA^{-1/2})$ is positive. If A is positive semidefinite, then a continuity argument implies that $\det(A + B)$ is nonnegative.) (Remark: Extensions of this result are given in [219].)

Fact 8.13.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite and B is Hermitian. Then,

$$\det(A + jB) = (\det A) \prod_{i=1}^n \left[1 + \sigma_i^2 \left(A^{-1/2}BA^{-1/2} \right) \right]^{1/2}.$$

(Proof: See [320].)

Fact 8.13.8. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$n + \operatorname{tr} \log A = n + \log \det A \leq n(\det A)^{1/n} \leq \operatorname{tr} A \leq (n \operatorname{tr} A^2)^{1/2},$$

with equality if and only if $A = I$. (Remark: The inequality

$$(\det A)^{1/n} \leq \frac{1}{n} \operatorname{tr} A$$

is a consequence of the arithmetic-mean–geometric-mean inequality.)

Fact 8.13.9. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that $A \leq B$. Then,

$$n \det A + \det B \leq \det(A + B).$$

(Proof: See [1098, pp. 154, 166].) (Remark: Under weaker conditions, Corollary 8.4.15 implies that $\det A + \det B \leq \det(A + B)$.)

Fact 8.13.10. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\det A + \det B + (2^n - 2)\sqrt{\det AB} \leq \det(A + B).$$

If, in addition, $B \leq A$, then

$$\det A + (2^n - 1)\det B \leq \det A + \det B + (2^n - 2)\sqrt{\det AB} \leq \det(A + B).$$

(Proof: See [1057] or [1184, p. 231].)

Fact 8.13.11. Let $A \in \mathbb{R}^{n \times n}$, and assume that $A + A^T$ is positive semidefinite. Then,

$$\left[\frac{1}{2}(A + A^T)\right]^A \leq \frac{1}{2}(A^A + A^{A^T}).$$

Now, assume that $A + A^T$ is positive definite. Then,

$$\left[\det \frac{1}{2}(A + A^T)\right] \left[\frac{1}{2}(A + A^T)\right]^{-1} \leq (\det A) \left[\frac{1}{2}(A^{-1} + A^{-T})\right].$$

Furthermore,

$$\left[\det \frac{1}{2}(A + A^T)\right] \left[\frac{1}{2}(A + A^T)\right]^{-1} < (\det A) \left[\frac{1}{2}(A^{-1} + A^{-T})\right]$$

if and only if $\text{rank}(A - A^T) \geq 4$. Finally, if $n \geq 4$ and $A - A^T$ is nonsingular, then

$$(\det A) \left[\frac{1}{2}(A^{-1} + A^{-T})\right] < \left[\det A - \det \frac{1}{2}(A - A^T)\right] \left[\frac{1}{2}(A + A^T)\right]^{-1}.$$

(Proof: See [465, 759].) (Remark: This result does not hold for complex matrices.)

(Remark: See Fact 8.9.31 and Fact 8.17.12.)

Fact 8.13.12. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is positive definite. Then,

$$\sum_{i=1}^n [\det A_{(\{1, \dots, i\})}]^{1/i} \leq (1 + \frac{1}{n})^n \text{tr } A < e \text{tr } A.$$

(Proof: See [29].)

Fact 8.13.13. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite and Toeplitz, and, for all $i = 1, \dots, n$, define $A_i \triangleq A_{(\{1, \dots, i\})} \in \mathbb{F}^{i \times i}$. Then,

$$(\det A)^{1/n} \leq (\det A_{n-1})^{1/(n-1)} \leq \dots \leq (\det A_2)^{1/2} \leq \det A_1.$$

Furthermore,

$$\frac{\det A}{\det A_{n-1}} \leq \frac{\det A_{n-1}}{\det A_{n-2}} \leq \dots \leq \frac{\det A_3}{\det A_2} \leq \frac{\det A_2}{\det A_1}.$$

(Proof: See [352] or [353, p. 682].)

Fact 8.13.14. Let $A, B \in \mathbb{F}^{n \times n}$, assume that B is Hermitian, and assume that $A^*BA < A + A^*$. Then, $\det A \neq 0$.

Fact 8.13.15. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $\alpha \in [0, 1]$. Then,

$$(\det A)^\alpha (\det B)^{1-\alpha} \leq \det[\alpha A + (1-\alpha)B].$$

Furthermore, equality holds if and only if $A = B$. (Proof: This inequality is a restatement of *xxviii*) of Proposition 8.6.17.) (Remark: This result is due to Bergstrom.) (Remark: $\alpha = 2$ yields $\sqrt{(\det A) \det B} \leq \det[\frac{1}{2}(A+B)]$.)

Fact 8.13.16. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, assume that either $A \leq B$ or $B \leq A$, and let $\alpha \in [0, 1]$. Then,

$$\det[\alpha A + (1-\alpha)B] \leq \alpha \det A + (1-\alpha) \det B.$$

(Proof: See [1406].)

Fact 8.13.17. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$\frac{\det A}{\det A_{[1;1]}} + \frac{\det B}{\det B_{[1;1]}} \leq \frac{\det(A+B)}{\det(A_{[1;1]} + B_{[1;1]})}.$$

(Proof: See [1098, p. 145].) (Remark: This inequality is a special case of *xli*) of Proposition 8.6.17.) (Remark: See Fact 8.11.4.)

Fact 8.13.18. Let $A_1, \dots, A_k \in \mathbb{F}^{n \times n}$, assume that $A_1, \dots, A_k$ are positive semidefinite, and let $\lambda_1, \dots, \lambda_k \in \mathbb{C}$. Then,

$$\det\left(\sum_{i=1}^k \lambda_i A_i\right) \leq \det\left(\sum_{i=1}^k |\lambda_i| A_i\right).$$

(Proof: See [1098, p. 144].)

Fact 8.13.19. Let $A, B, C \in \mathbb{R}^{n \times n}$, let $D \triangleq A + jB$, and assume that $CB + B^T C^T < D + D^*$. Then, $\det A \neq 0$.

Fact 8.13.20. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $m \in \mathbb{P}$. Then,

$$n^{1/m} (\det AB)^{1/n} \leq (\operatorname{tr} A^m B^m)^{1/m}.$$

(Proof: See [369].) (Remark: Assuming $\det B = 1$ and setting $m = 1$ yields Proposition 8.4.14.)

Fact 8.13.21. Let $A, B, C \in \mathbb{F}^{n \times n}$, define

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix},$$

and assume that $\mathcal{A}$ is positive semidefinite. Then,

$$|\det(B + B^*)| \leq \det(A + C).$$

If, in addition, $\mathcal{A}$ is positive definite, then

$$|\det(B + B^*)| < \det(A + C).$$

(Remark: Use Fact 8.11.5.)

Fact 8.13.22. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$|\det A^*B|^2 \leq (\det A^*A)(\det B^*B).$$

(Proof: Use Fact 8.11.14 or apply Fact 8.13.42 to $\begin{bmatrix} A^*A & B^*A \\ A^*B & B^*B \end{bmatrix}$.) (Remark: This result is a determinantal version of the Cauchy-Schwarz inequality.)

Fact 8.13.23. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and let $B \in \mathbb{F}^{m \times n}$, where $\text{rank } B = m$. Then,

$$(\det BB^*)^2 \leq (\det BAB^*) \det BA^{-1}B^*.$$

(Proof: Use Fact 8.11.19.)

Fact 8.13.24. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$|\det(A+B)|^2 + |\det(I-AB^*)|^2 \leq \det(I+AA^*)\det(I+B^*B)$$

and

$$|\det(A-B)|^2 + |\det(I+AB^*)|^2 \leq \det(I+AA^*)\det(I+B^*B).$$

Furthermore, the first inequality is an identity if and only if either $n = 1$, $A+B = 0$, or $AB^* = I$. (Proof: The result follows from Fact 8.11.16. See [1490, p. 184].)

Fact 8.13.25. Let $A, B \in \mathbb{F}^{n \times m}$, and assume that $I - A^*A$ and $I - B^*B$ are positive semidefinite. Then,

$$\begin{aligned} 0 &\leq \det(I - A^*A)\det(I - B^*B) \\ &\leq \begin{Bmatrix} |\det(I - A^*B)|^2 \\ |\det(I + A^*B)|^2 \end{Bmatrix} \\ &\leq \det(I + A^*A)\det(I + B^*B). \end{aligned}$$

Now, assume that $n = m$. Then,

$$\begin{aligned} 0 &\leq \det(I - A^*A)\det(I - B^*B) \\ &\leq |\det(I - A^*B)|^2 - |\det(A - B)|^2 \\ &\leq |\det(I - A^*B)|^2 \\ &\leq |\det(I - A^*B)|^2 + |\det(A + B)|^2 \\ &\leq \det(I + A^*A)\det(I + B^*B) \end{aligned}$$

and

$$\begin{aligned}
0 &\leq \det(I - A^*A)\det(I - B^*B) \\
&\leq |\det(I + A^*B)|^2 - |\det(A + B)|^2 \\
&\leq |\det(I + A^*B)|^2 \\
&\leq |\det(I + A^*B)|^2 + |\det(A - B)|^2 \\
&\leq \det(I + A^*A)\det(I + B^*B).
\end{aligned}$$

Finally,

$$\begin{bmatrix} \det[(I - A^*A)^{-1}] & \det[(I - A^*B)^{-1}] \\ \det[(I - B^*A)^{-1}] & \det[(I - B^*B)^{-1}] \end{bmatrix} \geq 0.$$

(Proof: The second inequality and Fact 8.11.21 are *Hua's inequalities*. See [47]. The third inequality follows from Fact 8.11.15. The first interpolation in the case $n = m$ is given in [1060].) (Remark: Generalizations of the last result are given in [1467].) (Remark: See Fact 8.11.21 and Fact 8.15.19.)

Fact 8.13.26. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\alpha, \beta \in (0, \infty)$. Then,

$$|\det(A + B)|^2 \leq \det(\beta^{-1}I + \alpha A^*A)\det(\alpha^{-1}I + \beta B B^*).$$

(Proof: Use Fact 8.11.20. See [1491].)

Fact 8.13.27. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{n \times m}$, and $D \in \mathbb{F}^{n \times l}$. Then,

$$|\det(AC^* + BD^*)|^2 \leq \det(AA^* + BB^*)\det(CC^* + DD^*).$$

(Proof: Use Fact 8.13.38 and $\mathcal{A}\mathcal{A}^* \geq 0$, where $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix}$.) (Remark: See Fact 2.14.22.)

Fact 8.13.28. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times m}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times m}$. Then,

$$|\det(A^*B + C^*D)|^2 \leq \det(A^*A + C^*C)\det(B^*B + D^*D).$$

(Proof: Use Fact 8.13.38 and $\mathcal{A}^*\mathcal{A} \geq 0$, where $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix}$.) (Remark: See Fact 2.14.18.)

Fact 8.13.29. Let $A, B, C \in \mathbb{F}^{n \times n}$. Then,

$$|\det(B + CA)|^2 \leq \det(A^*A + B^*B)\det(I + CC^*).$$

(Proof: See [717].) (Remark: See Fact 8.10.37.)

Fact 8.13.30. Let $A, B \in \mathbb{F}^{n \times m}$. Then, there exist unitary matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that

$$I + \langle A + B \rangle \leq S_1(I + \langle A \rangle)^{1/2}S_2(I + \langle B \rangle)S_2^*(I + \langle A \rangle)^{1/2}S_1^*.$$

Therefore,

$$\det(I + \langle A + B \rangle) \leq \det(I + \langle A \rangle)\det(I + \langle B \rangle).$$

(Proof: See [47, 1270].) (Remark: This result is due to Seiler and Simon.)

Fact 8.13.31. Let $A, B \in \mathbb{F}^{n \times n}$, assume that $A + A^* > 0$ and $B + B^* \geq 0$, and let $\alpha > 0$. Then, $\alpha I + AB$ is nonsingular and has no negative eigenvalues. Hence,

$$\det(\alpha I + AB) > 0.$$

(Proof: See [613].) (Remark: Equivalently, $-A$ is dissipative and $-B$ is semidissipative.) (Problem: Find a positive lower bound for $\det(\alpha I + AB)$ in terms of α , A , and B .)

Fact 8.13.32. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and define

$$\alpha \triangleq \frac{1}{n} \operatorname{tr} A$$

and

$$\beta \triangleq \frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n |A_{(i,j)}|.$$

Then,

$$|\det A| \leq (\alpha - \beta)^{n-1} [\alpha + (n-1)\beta].$$

Furthermore, if $A = aI_n + b1_{n \times n}$, where $a + nb > 0$ and $a > 0$, then $\alpha = a + b$, $\beta = b$, and equality holds. (Proof: See [1033].) (Remark: See Fact 2.13.12 and Fact 8.9.34.)

Fact 8.13.33. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and define

$$\beta \triangleq \frac{1}{n(n-1)} \sum_{\substack{i,j=1 \\ i \neq j}}^n \frac{|A_{(i,j)}|}{\sqrt{A_{(i,i)}A_{(j,j)}}}.$$

Then,

$$|\det A| \leq (1 - \beta)^{n-1} [1 + (n-1)\beta] \prod_{i=1}^n A_{(i,i)}.$$

(Proof: See [1033].) (Remark: This inequality strengthens Hadamard's inequality. See Fact 8.17.11. See also [412].)

Fact 8.13.34. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$|\det A| \leq \prod_{i=1}^n \left(\sum_{j=1}^n |A_{(i,j)}|^2 \right)^{1/2} = \prod_{i=1}^n \|\operatorname{row}_i(A)\|_2.$$

Furthermore, equality holds if and only if AA^* is diagonal. Now, let $\alpha > 0$ be such that, for all $i, j = 1, \dots, n$, $|A_{(i,j)}| \leq \alpha$. Then,

$$|\det A| \leq \alpha^n n^{n/2}.$$

If, in addition, at least one entry of A has absolute value less than α , then

$$|\det A| < \alpha^n n^{n/2}.$$

(Remark: Replace A with AA^* in Fact 8.17.11.) (Remark: This result is a direct consequence of Hadamard's inequality. See Fact 8.17.11.) (Remark: See Fact 2.13.14 and Fact 6.5.26.)

Fact 8.13.35. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, and assume that $\mathcal{A}$ is positive definite. Then,

$$\det \mathcal{A} = (\det A) \det(C - B^* A^{-1} B) \leq (\det A) \det C \leq \prod_{i=1}^{n+m} \mathcal{A}_{(i,i)}.$$

(Proof: The second inequality is obtained by successive application of the first inequality.) (Remark: $\det \mathcal{A} \leq (\det A) \det C$ is *Fischer's inequality*.)

Fact 8.13.36. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, assume that $\mathcal{A}$ is positive definite, let $k \triangleq \min\{m, n\}$, and, for $i = 1, \dots, n$, let $\lambda_i \triangleq \lambda_i(\mathcal{A})$. Then,

$$\prod_{i=1}^{n+m} \lambda_i \leq (\det A) \det C \leq \left(\prod_{i=k+1}^{n+m-k} \lambda_i \right) \prod_{i=1}^k \left[\frac{1}{2} (\lambda_i + \lambda_{n+m-i+1}) \right]^2.$$

(Proof: The left-hand inequality is given by Fact 8.13.35. The right-hand inequality is given in [1025].)

Fact 8.13.37. Let $A \in \mathbb{F}^{n \times n}$, and let $\mathcal{S} \subseteq \{1, \dots, n\}$. Then, the following statements hold:

i) If $\alpha \subset \{1, \dots, n\}$, then

$$\det A \leq [\det A_{(\alpha)}] \det A_{(\alpha^c)}.$$

ii) If $\alpha, \beta \subseteq \{1, \dots, n\}$, then

$$\det A_{(\alpha \cup \beta)} \leq \frac{[\det A_{(\alpha)}] \det A_{(\beta)}}{\det A_{(\alpha \cap \beta)}}.$$

iii) If $1 \leq k \leq n-1$, then

$$\left(\prod_{\{\alpha: \text{card}(\alpha)=k+1\}} \det A_{(\alpha)} \right)^{\binom{n-1}{k-1}} \leq \left(\prod_{\{\alpha: \text{card}(\alpha)=k\}} \det A_{(\alpha)} \right)^{\binom{n-1}{k}}.$$

(Proof: See [938].) (Remark: The first result is Fischer's inequality, see Fact 8.13.35. The second result is the *Hadamard-Fischer inequality*. The third result is *Szasz's inequality*. See [353, p. 680], [709, p. 479], and [938].) (Remark: See Fact 8.13.36.)

Fact 8.13.38. Let $A, B, C \in \mathbb{F}^{n \times n}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{2n \times 2n}$, and assume that $\mathcal{A}$ is positive semidefinite. Then,

$$0 \leq (\det A) \det C - |\det B|^2 \leq \det \mathcal{A} \leq (\det A) \det C.$$

Hence,

$$|\det B|^2 \leq (\det A) \det C.$$

Furthermore, $\mathcal{A}$ is positive definite if and only if

$$|\det B|^2 < (\det A) \det C.$$

(Proof: Assuming that A is positive definite, it follows that $0 \leq B^* A^{-1} B \leq C$, which implies that $|\det B|^2 / \det A \leq \det C$. Then, use continuity for the case in which A

is singular. For an alternative proof, see [1098, p. 142]. For the case in which A is positive definite, note that $0 \leq B^*A^{-1}B < C$, and thus $|\det B|^2/\det A < \det C$. (Remark: This result is due to Everitt.) (Remark: See Fact 8.13.42.) (Remark: When B is nonsquare, it is not necessarily true that $|\det(B^*B)|^2 < (\det A)\det C$. See [1492].)

Fact 8.13.39. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, and assume that $\mathcal{A}$ is positive semidefinite and A is positive definite. Then,

$$B^*A^{-1}B \leq \left[\frac{\lambda_{\max}(\mathcal{A}) - \lambda_{\min}(\mathcal{A})}{\lambda_{\max}(\mathcal{A}) + \lambda_{\min}(\mathcal{A})} \right]^2 C.$$

(Proof: See [886, 1494].)

Fact 8.13.40. Let $A, B, C \in \mathbb{F}^{n \times n}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{2n \times 2n}$, and assume that $\mathcal{A}$ is positive semidefinite. Then,

$$|\det B|^2 \leq \left[\frac{\lambda_{\max}(\mathcal{A}) - \lambda_{\min}(\mathcal{A})}{\lambda_{\max}(\mathcal{A}) + \lambda_{\min}(\mathcal{A})} \right]^{2n} (\det A)\det C.$$

Hence,

$$|\det B|^2 \leq \left[\frac{\lambda_{\max}(\mathcal{A}) - \lambda_{\min}(\mathcal{A})}{\lambda_{\max}(\mathcal{A}) + \lambda_{\min}(\mathcal{A})} \right]^2 (\det A)\det C.$$

Now, define $\hat{\mathcal{A}} \triangleq \begin{bmatrix} \det A & \det B \\ \det B^* & \det C \end{bmatrix} \in \mathbb{F}^{2 \times 2}$. Then,

$$|\det B|^2 \leq \left[\frac{\lambda_{\max}(\hat{\mathcal{A}}) - \lambda_{\min}(\hat{\mathcal{A}})}{\lambda_{\max}(\hat{\mathcal{A}}) + \lambda_{\min}(\hat{\mathcal{A}})} \right]^2 (\det A)\det C.$$

(Proof: See [886, 1494].) (Remark: The second and third bounds are not comparable. See [886, 1494].)

Fact 8.13.41. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, assume that $\mathcal{A}$ is positive semidefinite, and assume that A and C are positive definite. Then,

$$\det(A|\mathcal{A})\det(C|\mathcal{A}) \leq \det A.$$

(Proof: See [717].) (Remark: This result is the *reverse Fischer inequality*.)

Fact 8.13.42. Let $A_{ij} \in \mathbb{F}^{n \times n}$ for all $i, j = 1, \dots, k$, define

$$A \triangleq \begin{bmatrix} A_{11} & \cdots & A_{1k} \\ \vdots & \ddots & \vdots \\ A_{1k}^* & \cdots & A_{kk} \end{bmatrix},$$

assume that A is positive semidefinite, let $1 \leq k \leq n$, and define

$$\tilde{A}_k \triangleq \begin{bmatrix} A_{11}^{(k)} & \cdots & A_{1k}^{(k)} \\ \vdots & \ddots & \vdots \\ A_{1k}^{*(k)} & \cdots & A_{kk}^{(k)} \end{bmatrix}.$$

Then, $\tilde{A}_k$ is positive semidefinite. In particular,

$$\tilde{A}_n = \begin{bmatrix} \det A_{11} & \cdots & \det A_{1k} \\ \vdots & \ddots & \vdots \\ \det A_{1k}^* & \cdots & \det A_{kk} \end{bmatrix}$$

is positive semidefinite. Furthermore,

$$\det A \leq \det \tilde{A}.$$

Now, assume that A is positive definite. Then, $\det A = \det \tilde{A}$ if and only if, for all distinct $i, j = 1, \dots, k$, $A_{ij} = 0$. (Proof: The first statement is given in [386]. The inequality as well as the final statement are given in [1267].) (Remark: $B^{(k)}$ is the k th compound of B . See Fact 7.5.17.) (Remark: Note that every principal subdeterminant of $\tilde{A}_n$ is lower bounded by the determinant of a positive-semidefinite matrix. Hence, the inequality implies that $\tilde{A}_n$ is positive semidefinite.) (Remark: A weaker result is given in [388] and quoted in [961] in terms of elementary symmetric functions of the eigenvalues of each block.) (Remark: The example $A = \begin{bmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$ shows that $\tilde{A}$ can be positive definite while A is singular.) (Remark: The matrix whose (i, j) entry is $\det A_{ij}$ is a *determinantal compression* of A . See [387, 964, 1267].) (Remark: See Fact 8.12.42.)

8.14 Facts on Convex Sets and Convex Functions

Fact 8.14.1. Let $f: \mathbb{R}^n \mapsto \mathbb{R}^n$, and assume that f is convex. Then, for all $\alpha \in \mathbb{R}$, the sets $\{x \in \mathbb{R}^n: f(x) \leq \alpha\}$ and $\{x \in \mathbb{R}^n: f(x) < \alpha\}$ are convex. (Proof: See [495, p. 108].) (Remark: The converse is not true. Consider the function $f(x) = x^3$.

Fact 8.14.2. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let $\alpha \geq 0$, and define the set $\mathcal{S} \triangleq \{x \in \mathbb{F}^n: x^*Ax < \alpha\}$. Then, the following statements hold:

- i) $\mathcal{S}$ is open.
- ii) $\mathcal{S}$ is a blunt cone if and only if $\alpha = 0$.
- iii) $\mathcal{S}$ is nonempty if and only if either $\alpha > 0$ or $\lambda_{\min}(A) < 0$.
- iv) $\mathcal{S}$ is convex if and only if $A \geq 0$.
- v) $\mathcal{S}$ is convex and nonempty if and only if $\alpha > 0$ and $A \geq 0$.
- vi) The following statements are equivalent:
 - a) $\mathcal{S}$ is bounded.
 - b) $\mathcal{S}$ is convex and bounded.
 - c) $A > 0$.
- vii) The following statements are equivalent:
 - a) $\mathcal{S}$ is bounded and nonempty.
 - b) $\mathcal{S}$ is convex, bounded, and nonempty.

c) $\alpha > 0$ and $A > 0$.

Fact 8.14.3. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let $\alpha \geq 0$, and define the set $\mathcal{S} \triangleq \{x \in \mathbb{F}^n: x^*Ax \leq \alpha\}$. Then, the following statements hold:

- i) $\mathcal{S}$ is closed.
- ii) $0 \in \mathcal{S}$, and thus $\mathcal{S}$ is nonempty.
- iii) $\mathcal{S}$ is a pointed cone if and only if $\alpha = 0$ or $A \leq 0$.
- iv) $\mathcal{S}$ is convex if and only if $A \geq 0$.
- v) The following statements are equivalent:
 - a) $\mathcal{S}$ is bounded.
 - b) $\mathcal{S}$ is convex and bounded.
 - c) $A > 0$.

Fact 8.14.4. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let $\alpha \geq 0$, and define the set $\mathcal{S} \triangleq \{x \in \mathbb{F}^n: x^*Ax = \alpha\}$. Then, the following statements hold:

- i) $\mathcal{S}$ is closed.
- ii) $\mathcal{S}$ is nonempty if and only if either $\alpha = 0$ or $\lambda_{\max}(A) > 0$.
- iii) The following statements are equivalent:
 - a) $\mathcal{S}$ is a pointed cone.
 - b) $0 \in \mathcal{S}$.
 - c) $\alpha = 0$.
- iv) $\mathcal{S} = \{0\}$ if and only if $\alpha = 0$ and either $A > 0$ or $A < 0$.
- v) $\mathcal{S}$ is bounded if and only if either $A > 0$ or both $\alpha > 0$ and $A \leq 0$.
- vi) $\mathcal{S}$ is bounded and nonempty if and only if $A > 0$.
- vii) The following statements are equivalent:
 - a) $\mathcal{S}$ is convex.
 - b) $\mathcal{S}$ is convex and nonempty.
 - c) $\alpha = 0$ and either $A > 0$ or $A < 0$.
- viii) If $\alpha > 0$, then the following statements are equivalent:
 - a) $\mathcal{S}$ is nonempty.
 - b) $\mathcal{S}$ is not convex.
 - c) $\lambda_{\max}(A) > 0$.
- ix) The following statements are equivalent:
 - a) $\mathcal{S}$ is convex and bounded.
 - b) $\mathcal{S}$ is convex, bounded, and nonempty.
 - c) $\alpha = 0$ and $A > 0$.

Fact 8.14.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let $\alpha \geq 0$, and define the set $\mathcal{S} \triangleq \{x \in \mathbb{F}^n: x^*Ax \geq \alpha\}$. Then, the following statements hold:

- i) $\mathcal{S}$ is closed.
- ii) $\mathcal{S}$ is a pointed cone if and only if $\alpha = 0$.
- iii) $\mathcal{S}$ is nonempty if and only if either $\alpha = 0$ or $\lambda_{\max}(A) > 0$.
- iv) $\mathcal{S}$ is bounded if and only if $\mathcal{S} \subseteq \{0\}$.
- v) The following statements are equivalent:
 - a) $\mathcal{S}$ is bounded and nonempty.
 - b) $\mathcal{S} = \{0\}$.
 - c) $\alpha = 0$ and $A < 0$.
- vi) $\mathcal{S}$ is convex if and only if either $\mathcal{S}$ is empty or $\mathcal{S} = \mathbb{F}^n$.
- vii) $\mathcal{S}$ is convex and bounded if and only if $\mathcal{S}$ is empty.
- viii) The following statements are equivalent:
 - a) $\mathcal{S}$ is convex and nonempty.
 - b) $\mathcal{S} = \mathbb{F}^n$.
 - c) $\alpha = 0$ and $A \geq 0$.

Fact 8.14.6. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let $\alpha \geq 0$, and define the set $\mathcal{S} \triangleq \{x \in \mathbb{F}^n: x^*Ax > \alpha\}$. Then, the following statements hold:

- i) $\mathcal{S}$ is open.
- ii) $\mathcal{S}$ is a blunt cone if and only if $\alpha = 0$.
- iii) $\mathcal{S}$ is nonempty if and only if $\lambda_{\max}(A) > 0$.
- iv) The following statements are equivalent:
 - a) $\mathcal{S}$ is empty.
 - b) $\lambda_{\max}(A) \leq 0$.
 - c) $\mathcal{S}$ is bounded.
 - d) $\mathcal{S}$ is convex.

Fact 8.14.7. Let $A \in \mathbb{C}^{n \times n}$, and define the *numerical range* of A by

$$\Theta_1(A) \triangleq \{x^*Ax: x \in \mathbb{C}^n \text{ and } x^*x = 1\}$$

and the set

$$\Theta(A) \triangleq \{x^*Ax: x \in \mathbb{C}^n\}.$$

Then, the following statements hold:

- i) $\Theta_1(A)$ is a closed, bounded, convex subset of $\mathbb{C}$.
- ii) $\Theta(A) = \{0\} \cup \text{cone } \Theta_1(A)$.
- iii) $\Theta(A)$ is a pointed, closed, convex cone contained in $\mathbb{C}$.

- iv) If A is Hermitian, then $\Theta_1(A)$ is a closed, bounded interval contained in $\mathbb{R}$.
- v) If A is Hermitian, then $\Theta(A)$ is either $(-\infty, 0]$, $[0, \infty)$, or $\mathbb{R}$.
- vi) $\Theta_1(A)$ satisfies

$$\operatorname{co spec}(A) \subseteq \Theta_1(A) \subseteq \operatorname{co}\{\nu_1 + j\mu_1, \nu_1 + j\mu_n, \nu_n + j\mu_1, \nu_n + j\mu_n\},$$

where

$$\nu_1 \triangleq \lambda_{\max}\left[\frac{1}{2}(A + A^*)\right], \quad \nu_n \triangleq \lambda_{\min}\left[\frac{1}{2}(A + A^*)\right],$$

$$\mu_1 \triangleq \lambda_{\max}\left[\frac{1}{2j}(A - A^*)\right], \quad \mu_n \triangleq \lambda_{\min}\left[\frac{1}{2j}(A - A^*)\right].$$

- vii) If A is normal, then

$$\Theta_1(A) = \operatorname{co spec}(A).$$

- viii) If $n \leq 4$ and $\Theta_1(A) = \operatorname{co spec}(A)$, then A is normal.

- ix) $\Theta_1(A) = \operatorname{co spec}(A)$ if and only if either A is normal or there exist matrices $A_1 \in \mathbb{F}^{n_1 \times n_1}$ and $A_2 \in \mathbb{F}^{n_2 \times n_2}$ such that $n_1 + n_2 = n$, $\Theta_1(A_1) \subseteq \Theta_1(A_2)$, and A is unitarily similar to $\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}$.

(Proof: See [610] or [711, pp. 11, 52].) (Remark: $\Theta_1(A)$ is called the *field of values* in [711, p. 5].) (Remark: See Fact 4.10.24 and Fact 8.14.7.) (Remark: *viii*) is an example of the *quartic barrier*. See [351], Fact 8.15.37, and Fact 11.17.3.)

Fact 8.14.8. Let $A \in \mathbb{R}^{n \times n}$, and define the *real numerical range* of A by

$$\Psi_1(A) \triangleq \{x^T A x: x \in \mathbb{R}^n \text{ and } x^T x = 1\}$$

and the set

$$\Psi(A) \triangleq \{x^T A x: x \in \mathbb{R}^n\}.$$

Then, the following statements hold:

- i) $\Psi_1(A) = \Psi_1[\frac{1}{2}(A + A^T)]$.
- ii) $\Psi_1(A) = [\lambda_{\min}[\frac{1}{2}(A + A^T)], \lambda_{\max}[\frac{1}{2}(A + A^T)]]$.
- iii) If A is symmetric, then $\Psi_1(A) = [\lambda_{\min}(A), \lambda_{\max}(A)]$.
- iv) $\Psi(A) = \{0\} \cup \operatorname{cone} \Psi_1(A)$.
- v) $\Psi(A)$ is either $(-\infty, 0]$, $[0, \infty)$, or $\mathbb{R}$.
- vi) $\Psi_1(A) = \Theta_1(A)$ if and only if A is symmetric.

(Proof: See [711, p. 83].) (Remark: $\Theta_1(A)$ is defined in Fact 8.14.7.)

Fact 8.14.9. Let $A, B \in \mathbb{C}^{n \times n}$, assume that A and B are Hermitian, and define

$$\Theta_1(A, B) \triangleq \left\{ \begin{bmatrix} x^* A x \\ x^* B x \end{bmatrix} : x \in \mathbb{C}^n \text{ and } x^* x = 1 \right\} \subseteq \mathbb{R}^2.$$

Then, $\Theta_1(A, B)$ is convex. (Proof: See [1090].) (Remark: This result is an immediate consequence of Fact 8.14.7.)

Fact 8.14.10. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A and B are symmetric, and let α, β be real numbers. Then, the following statements are equivalent:

- i) There exists $x \in \mathbb{R}^n$ such that $x^T A x = \alpha$ and $x^T B x = \beta$.
- ii) There exists a positive-semidefinite matrix $X \in \mathbb{R}^{n \times n}$ such that $\text{tr} AX = \alpha$ and $\text{tr} BX = \beta$.

(Proof: See [153, p. 84].)

Fact 8.14.11. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A and B are symmetric, and define

$$\Psi_1(A, B) \triangleq \left\{ \begin{bmatrix} x^T A x \\ x^T B x \end{bmatrix} : x \in \mathbb{R}^n \text{ and } x^T x = 1 \right\} \subseteq \mathbb{R}^2$$

and

$$\Psi(A, B) \triangleq \left\{ \begin{bmatrix} x^T A x \\ x^T B x \end{bmatrix} : x \in \mathbb{R}^n \right\} \subseteq \mathbb{R}^2.$$

Then, $\Psi(A, B)$ is a pointed, convex cone. If, in addition, $n \geq 3$, then $\Psi_1(A, B)$ is convex. (Proof: See [153, pp. 84, 89] or [406, 1090].) (Remark: $\Psi(A, B) = [\text{cone } \Psi_1(A, B)] \cup \{[0]\}$.) (Remark: The set $\Psi(A, B)$ is not necessarily closed. See [406, 1063, 1064].)

Fact 8.14.12. Let $A, B \in \mathbb{R}^{n \times n}$, where $n \geq 2$, assume that A and B are symmetric, let $a, b \in \mathbb{R}^n$, let $a_0, b_0 \in \mathbb{R}$, assume that there exist real numbers α, β such that $\alpha A + \beta B > 0$, and define

$$\Psi(A, a, a_0, B, b, b_0) \triangleq \left\{ \begin{bmatrix} x^T A x + a^T x + a_0 \\ x^T B x + b^T x + b_0 \end{bmatrix} : x \in \mathbb{R}^n \right\} \subseteq \mathbb{R}^2.$$

Then, $\Psi(A, a, a_0, B, b, b_0)$ is closed and convex. (Proof: See [1090].)

Fact 8.14.13. Let $A, B, C \in \mathbb{R}^{n \times n}$, where $n \geq 3$, assume that A, B , and C are symmetric, and define

$$\Phi_1(A, B, C) \triangleq \left\{ \begin{bmatrix} x^T A x \\ x^T B x \\ x^T C x \end{bmatrix} : x \in \mathbb{R}^n \text{ and } x^T x = 1 \right\} \subseteq \mathbb{R}^3$$

and

$$\Phi(A, B, C) \triangleq \left\{ \begin{bmatrix} x^T A x \\ x^T B x \\ x^T C x \end{bmatrix} : x \in \mathbb{R}^n \right\} \subseteq \mathbb{R}^3.$$

Then, $\Phi_1(A, B, C)$ is convex and $\Phi(A, B, C)$ is a pointed, convex cone. (Proof: See [260, 1087, 1090].)

Fact 8.14.14. Let $A, B, C \in \mathbb{R}^{n \times n}$, where $n \geq 3$, assume that A, B , and C are symmetric, and define

$$\Phi(A, B, C) \triangleq \left\{ \begin{bmatrix} x^T A x \\ x^T B x \\ x^T C x \end{bmatrix} : x \in \mathbb{R}^n \right\} \subseteq \mathbb{R}^3.$$

Then, the following statements are equivalent:

- i) There exist real numbers α, β, γ such that $\alpha A + \beta B + \gamma C$ is positive definite.

- ii) $\Phi(A, B, C)$ is a pointed, one-sided, closed, convex cone, and, if $x \in \mathbb{R}^n$ satisfies $x^T A x = x^T B x = x^T C x = 0$, then $x = 0$.

(Proof: See [1090].)

Fact 8.14.15. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let $b \in \mathbb{F}^n$ and $c \in \mathbb{R}$, and define $f: \mathbb{F}^n \mapsto \mathbb{R}$ by

$$f(x) \triangleq x^* A x + \operatorname{Re}(b^* x) + c.$$

Then, the following statements hold:

- i) f is convex if and only if A is positive semidefinite.
 ii) f is strictly convex if and only if A is positive definite.

Now, assume that A is positive semidefinite. Then, f has a minimizer if and only if $b \in \mathcal{R}(A)$. In this case, the following statements hold.

- iii) The vector $x_0 \in \mathbb{F}^n$ is a minimizer of f if and only if x_0 satisfies $Ax_0 = -\frac{1}{2}b$.
 iv) $x_0 \in \mathbb{F}^m$ minimizes f if and only if there exists a vector $y \in \mathbb{F}^m$ such that

$$x_0 = -\frac{1}{2}A^+b + (I - A^+A)y.$$

- v) The minimum of f is given by

$$f(x_0) = c - x_0^* A x_0 = c - \frac{1}{4}b^* A^+ b.$$

- vi) If A is positive definite, then $x_0 = -\frac{1}{2}A^{-1}b$ is the unique minimizer of f , and the minimum of f is given by

$$f(x_0) = c - x_0^* A x_0 = c - \frac{1}{4}b^* A^{-1} b.$$

(Proof: Use Proposition 6.1.7 and note that, for every x_0 satisfying $Ax_0 = -\frac{1}{2}b$, it follows that

$$\begin{aligned} f(x_0) &= (x - x_0)^* A (x - x_0) + c - x_0^* A x_0 \\ &= (x - x_0)^* A (x - x_0) + c - \frac{1}{4}b^* A^+ b. \end{aligned}$$

(Remark: This result is the *quadratic minimization lemma*.) (Remark: See Fact 9.15.1.)

Fact 8.14.16. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and define $\phi: \mathbb{F}^{m \times n} \mapsto \mathbb{R}$ by $\phi(B) \triangleq \operatorname{tr} BAB^*$. Then, ϕ is strictly convex. (Proof: $\operatorname{tr}[\alpha(1 - \alpha)(B_1 - B_2)A(B_1 - B_2)^*] > 0$.)

Fact 8.14.17. Let $p, q \in \mathbb{R}$, and define $\phi: \mathbf{P}^n \times \mathbf{P}^n \rightarrow (0, \infty)$ by

$$\phi(A, B) \triangleq \operatorname{tr} A^p B^q.$$

Then, the following statements hold:

- i) If $p, q \in (0, 1)$ and $p + q \leq 1$, then $-\phi$ is convex.
 ii) If either $p, q \in [-1, 0)$ or $p \in [-1, 0)$, $q \in [1, 2]$, and $p + q \geq 1$, or $p \in [1, 2]$, $q \in [-1, 0]$, and $p + q \geq 1$, then ϕ is convex.

- iii) If p, q do not satisfy the hypotheses of either i) or ii), then neither ϕ nor $-\phi$ is convex.

(Proof: See [166].)

Fact 8.14.18. Let $B \in \mathbb{F}^{n \times n}$, assume that B is Hermitian, let $\alpha_1, \dots, \alpha_k \in (0, \infty)$, define $r \triangleq \sum_{i=1}^k \alpha_i$, assume that $r \leq 1$, let $q \in \mathbb{R}$, and define $\phi: \mathbf{P}^n \times \dots \times \mathbf{P}^n \rightarrow [0, \infty)$ by

$$\phi(A_1, \dots, A_k) \triangleq - \left[\text{tr } e^{B + \sum_{i=1}^k \alpha_i \log A_i} \right]^q.$$

If $q \in (0, 1/r]$, then ϕ is convex. Furthermore, if $q < 0$, then $-\phi$ is convex. (Proof: See [905, 933].) (Remark: See [989] and Fact 8.12.31.)

8.15 Facts on Quadratic Forms

Fact 8.15.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then,

$$\mathcal{N}(A) \subseteq \{x \in \mathbb{F}^n: x^*Ax = 0\}.$$

Furthermore,

$$\mathcal{N}(A) = \{x \in \mathbb{F}^n: x^*Ax = 0\}$$

if and only if either $A \geq 0$ or $A \leq 0$.

Fact 8.15.2. Let $x, y \in \mathbb{F}^n$. Then, $xx^* \leq yy^*$ if and only if there exists $\alpha \in \mathbb{R}$ such that $|\alpha| \in [0, 1]$ and $x = \alpha y$.

Fact 8.15.3. Let $x, y \in \mathbb{F}^n$. Then, $xy^* + yx^* \geq 0$ if and only if x and y are linearly dependent. (Proof: Evaluate the product of the nonzero eigenvalues of $xy^* + yx^*$, and use the Cauchy-Schwarz inequality $|x^*y|^2 \leq x^*xy^*y$.)

Fact 8.15.4. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, let $x \in \mathbb{F}^n$, and let $a \in [0, \infty)$. Then, the following statements are equivalent:

- i) $xx^* \leq aA$.
- ii) $x^*A^{-1}x \leq a$.
- iii) $\begin{bmatrix} A & x \\ x^* & a \end{bmatrix} \geq 0$.

(Proof: Use Fact 2.14.3 and Proposition 8.2.4. Note that, if $a = 0$, then $x = 0$.)

Fact 8.15.5. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, assume that $A + B$ is nonsingular, let $x, a, b \in \mathbb{F}^n$, and define $c \triangleq (A + B)^{-1}(Ax + Bb)$. Then,

$$(x-a)^*A(x-a) + (x-b)^*B(x-b) = (x-c)^*(A+B)(x-c) = (a-b)^*A(A+B)^{-1}B(a-b).$$

(Proof: See [1184, p. 278].)

Fact 8.15.6. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A is symmetric and B is skew symmetric, and let $x, y \in \mathbb{R}^n$. Then,

$$\begin{bmatrix} x \\ y \end{bmatrix}^T \begin{bmatrix} A & B \\ B^T & A \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = (x + jy)^*(A + jB)(x + jy).$$

(Remark: See Fact 4.10.26.)

Fact 8.15.7. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and let $x, y \in \mathbb{F}^n$. Then,

$$2\operatorname{Re} x^*y \leq x^*Ax + y^*A^{-1}y.$$

Furthermore, if $y = Ax$, then equality holds. Therefore,

$$x^*Ax = \max_{z \in \mathbb{F}^n} [2\operatorname{Re} x^*z - z^*Az].$$

(Proof: $(A^{1/2}x - A^{-1/2}y)^*(A^{1/2}x - A^{-1/2}y) \geq 0$.) (Remark: This result is due to Bellman. See [886, 1494].)

Fact 8.15.8. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and let $x, y \in \mathbb{F}^n$. Then,

$$|x^*y|^2 \leq (x^*Ax)(y^*A^{-1}y).$$

(Proof: Use Fact 8.11.14 with A replaced by $A^{1/2}x$ and B replaced by $A^{-1/2}y$.)

Fact 8.15.9. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and let $x \in \mathbb{F}^n$. Then,

$$(x^*x)^2 \leq (x^*Ax)(x^*A^{-1}x) \leq \frac{(\alpha + \beta)^2}{4\alpha\beta}(x^*x)^2,$$

where $\alpha \triangleq \lambda_{\min}(A)$ and $\beta \triangleq \lambda_{\max}(A)$. (Remark: The second inequality is the *Kantorovich inequality*. See Fact 1.15.36 and [22]. See also [927].)

Fact 8.15.10. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and let $x \in \mathbb{F}^n$. Then,

$$(x^*x)^{1/2}(x^*Ax)^{1/2} - x^*Ax \leq \frac{(\alpha - \beta)^2}{4(\alpha + \beta)}x^*x$$

and

$$(x^*x)(x^*A^2x) - (x^*Ax)^2 \leq \frac{1}{4}(\alpha - \beta)^2(x^*x)^2,$$

where $\alpha \triangleq \lambda_{\min}(A)$ and $\beta \triangleq \lambda_{\max}(A)$. (Proof: See [1079].) (Remark: Extensions of these results are given in [748, 1079].)

Fact 8.15.11. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, let $r \triangleq \operatorname{rank} A$, let $x \in \mathbb{F}^n$, and assume that $x \notin \mathcal{N}(A)$. Then,

$$\frac{x^*Ax}{x^*x} - \frac{x^*x}{x^*A^+x} \leq [\lambda_{\max}^{1/2}(A) - \lambda_r^{1/2}(A)]^2.$$

If, in addition, A is positive definite, then, for all nonzero $x \in \mathbb{F}^n$,

$$0 \leq \frac{x^*Ax}{x^*x} - \frac{x^*x}{x^*A^{-1}x} \leq [\lambda_{\max}^{1/2}(A) - \lambda_{\min}^{1/2}(A)]^2.$$

(Proof: See [1016, 1079]. The left-hand inequality in the last string of inequalities is given by Fact 8.15.9.)

Fact 8.15.12. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, let $y \in \mathbb{F}^n$, let $\alpha > 0$, and define $f: \mathbb{F}^n \mapsto \mathbb{R}$ by $f(x) \triangleq |x^*y|^2$. Then,

$$x_0 = \sqrt{\frac{\alpha}{y^*A^{-1}y}} A^{-1}y$$

minimizes $f(x)$ subject to $x^*Ax \leq \alpha$. Furthermore, $f(x_0) = \alpha y^*A^{-1}y$. (Proof: See [31].)

Fact 8.15.13. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $x \in \mathbb{F}^n$. Then,

$$(x^*A^2x)^2 \leq (x^*Ax)(x^*A^3x)$$

and

$$(x^*Ax)^2 \leq (x^*x)(x^*A^2x).$$

(Proof: Apply the Cauchy-Schwarz inequality Corollary 9.1.7.)

Fact 8.15.14. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $x \in \mathbb{F}^n$. If $\alpha \in [0, 1]$, then

$$x^*A^\alpha x \leq (x^*x)^{1-\alpha}(x^*Ax)^\alpha.$$

Furthermore, if $\alpha > 1$, then

$$(x^*Ax)^\alpha \leq (x^*x)^{\alpha-1}x^*A^\alpha x.$$

(Remark: The first inequality is the *Hölder-McCarthy inequality*, which is equivalent to the Young inequality. See Fact 8.9.42, Fact 8.10.43, [530, p. 125], and [532]. Matrix versions of the second inequality are given in [697].)

Fact 8.15.15. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, let $x \in \mathbb{F}^n$, and let $\alpha, \beta \in [1, \infty)$, where $\alpha \leq \beta$. Then,

$$(x^*A^\alpha x)^{1/\alpha} \leq (x^*A^\beta x)^{1/\beta}.$$

Now, assume that A is positive definite. Then,

$$x^*(\log A)x \leq \log x^*Ax \leq \frac{1}{\alpha} \log x^*A^\alpha x \leq \frac{1}{\beta} \log x^*A^\beta x.$$

(Proof: See [509].)

Fact 8.15.16. Let $A \in \mathbb{F}^{n \times n}$, $x, y \in \mathbb{F}^n$, and $\alpha \in (0, 1)$. Then,

$$|x^*Ay| \leq \|\langle A \rangle^\alpha x\|_2 \|\langle A^* \rangle^{1-\alpha} y\|_2.$$

Consequently,

$$|x^*Ay| \leq [x^*\langle A \rangle x]^{1/2} [y^*\langle A^* \rangle y]^{1/2}.$$

(Proof: See [775].)

Fact 8.15.17. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, assume that AB is Hermitian, and let $x \in \mathbb{F}^n$. Then,

$$|x^*ABx| \leq \text{sprad}(B)x^*Ax.$$

(Proof: See [911].) (Remark: This result is the sharpening by Halmos of Reid's inequality. Related results are given in [912].)

Fact 8.15.18. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $x \in \mathbb{F}^n$. Then,

$$x^*(A+B)^{-1}x \leq \frac{x^*A^{-1}xx^*B^{-1}x}{x^*A^{-1}x + x^*B^{-1}x} \leq \frac{1}{4}(x^*A^{-1}x + x^*B^{-1}x).$$

In particular,

$$\frac{1}{(A^{-1})_{(i,i)}} + \frac{1}{(B^{-1})_{(i,i)}} \leq \frac{1}{[(A+B)^{-1}]_{(i,i)}}.$$

(Proof: See [948, p. 201]. The right-hand inequality follows from Fact 1.10.4.)

(Remark: This result is *Bergstrom's inequality*.) (Remark: This result is a special case of Fact 8.11.3, which is a special case of *xvii*) of Proposition 8.6.17.)

Fact 8.15.19. Let $A, B \in \mathbb{F}^{n \times m}$, assume that $I - A^*A$ and $I - B^*B$ are positive semidefinite, and let $x \in \mathbb{C}^n$. Then,

$$x^*(I - A^*A)xx^*(I - B^*B)x \leq |x^*(I - A^*B)x|^2.$$

(Remark: This result is due to Marcus. See [1060].) (Remark: See Fact 8.13.25.)

Fact 8.15.20. Let $A, B \in \mathbb{R}^n$, and assume that A is Hermitian and B is positive definite. Then,

$$\lambda_{\max}(AB^{-1}) = \max\{\lambda \in \mathbb{R} : \det(A - \lambda B) = 0\} = \min_{x \in \mathbb{F}^n \setminus \{0\}} \frac{x^*Ax}{x^*Bx}.$$

(Proof: Use Lemma 8.4.3.)

Fact 8.15.21. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite and B is positive semidefinite. Then,

$$4(x^*x)(x^*Bx) < (x^*Ax)^2$$

for all nonzero $x \in \mathbb{F}^n$ if and only if there exists $\alpha > 0$ such that

$$\alpha I + \alpha^{-1}B < A.$$

In this case, $4B < A^2$, and hence $2B^{1/2} < A$. (Proof: Sufficiency follows from $\alpha x^*x + \alpha^{-1}x^*Bx < x^*Ax$. Necessity follows from Fact 8.15.22. The last result follows from $(A - 2\alpha I)^2 \geq 0$ or $2B^{1/2} \leq \alpha I + \alpha^{-1}B$.)

Fact 8.15.22. Let $A, B, C \in \mathbb{F}^{n \times n}$, assume that A, B, C are positive semidefinite, and assume that

$$4(x^*Cx)(x^*Bx) < (x^*Ax)^2$$

for all nonzero $x \in \mathbb{F}^n$. Then, there exists $\alpha > 0$ such that

$$\alpha C + \alpha^{-1}B < A.$$

(Proof: See [1083].)

Fact 8.15.23. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian and B is positive semidefinite. Then, $x^*Ax < 0$ for all $x \in \mathbb{F}^n$ such that $Bx = 0$ and $x \neq 0$ if and only if there exists $\alpha > 0$ such that $A < \alpha B$. (Proof: To prove necessity, suppose that, for every $\alpha > 0$, there exists a nonzero vector x such that $x^*Ax \geq \alpha x^*Bx$. Now, $Bx = 0$ implies that $x^*Ax \geq 0$. Sufficiency is immediate.)

Fact 8.15.24. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that A and B are Hermitian. Then, the following statements are equivalent:

- i) There exist $\alpha, \beta \in \mathbb{R}$ such that $\alpha A + \beta B$ is positive definite.
- ii) $\{x \in \mathbb{C}^n: x^*Ax = x^*Bx = 0\} = \{0\}$.

(Remark: This result is *Finsler's lemma*. See [83, 163, 866, 1340, 1352].) (Remark: See Fact 8.15.25, Fact 8.16.5, and Fact 8.16.6.)

Fact 8.15.25. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that A and B are symmetric. Then, the following statements are equivalent:

- i) There exist $\alpha, \beta \in \mathbb{R}$ such that $\alpha A + \beta B$ is positive definite.
- ii) Either $x^TAx > 0$ for all nonzero $x \in \{y \in \mathbb{F}^n: y^TB y = 0\}$ or $x^TAx < 0$ for all nonzero $x \in \{y \in \mathbb{F}^n: y^TB y = 0\}$.

Now, assume that $n \geq 3$. Then, the following statement is equivalent to i) and ii):

- iii) $\{x \in \mathbb{R}^n: x^TAx = x^TBx = 0\} = \{0\}$.

(Remark: This result is related to Finsler's lemma. See [83, 163, 1352].) (Remark: See Fact 8.15.24, Fact 8.16.5, and Fact 8.16.6.)

Fact 8.15.26. Let $A, B \in \mathbb{C}^{n \times n}$, assume that A and B are Hermitian, and assume that $x^*(A + jB)x$ is nonzero for all nonzero $x \in \mathbb{C}^n$. Then, there exists $t \in [0, \pi)$ such that $(\sin t)A + (\cos t)B$ is positive definite. (Proof: See [355] or [1230, p. 282].)

Fact 8.15.27. Let $A \in \mathbb{R}^{n \times n}$, assume that A is symmetric, and let $B \in \mathbb{R}^{n \times m}$. Then, the following statements are equivalent:

- i) $x^TAx > 0$ for all nonzero $x \in \mathcal{N}(B^T)$.

$$ii) \nu_+ \left(\begin{bmatrix} A & B \\ B^T & 0 \end{bmatrix} \right) = n.$$

Furthermore, the following statements are equivalent:

- iii) $x^TAx \geq 0$ for all $x \in \mathcal{N}(B^T)$.

$$iv) \nu_- \left(\begin{bmatrix} A & B \\ B^T & 0 \end{bmatrix} \right) = \text{rank } B.$$

(Proof: See [299, 945].) (Remark: See Fact 5.8.21 and Fact 8.15.28.)

Fact 8.15.28. Let $A \in \mathbb{R}^{n \times n}$, assume that A is symmetric, let $B \in \mathbb{R}^{n \times m}$, where $m \leq n$, and assume that $\begin{bmatrix} I_m & 0 \end{bmatrix} B$ is nonsingular. Then, the following

statements are equivalent:

- i) $x^T A x > 0$ for all nonzero $x \in \mathcal{N}(B^T)$.
- ii) For all $i = m+1, \dots, n$, the sign of the $i \times i$ leading principal subdeterminant of the matrix $\begin{bmatrix} 0 & B^T \\ B & A \end{bmatrix}$ is $(-1)^m$.

(Proof: See [94, p. 20], [936, p. 312], or [955].) (Remark: See Fact 8.15.27.)

Fact 8.15.29. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite and nonzero, let $x, y \in \mathbb{F}^n$, and assume that $x^* y = 0$. Then,

$$|x^* A y|^2 \leq \left[\frac{\lambda_{\max}(A) - \lambda_{\min}(A)}{\lambda_{\max}(A) + \lambda_{\min}(A)} \right]^2 (x^* A x)(y^* A y).$$

Furthermore, there exist vectors $x, y \in \mathbb{F}^n$ satisfying $x^* y = 0$ for which equality holds. (Proof: See [711, p. 443] or [886, 1494].) (Remark: This result is the *Wielandt inequality*.)

Fact 8.15.30. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, define $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$, and assume that A and C are positive semidefinite. Then, the following statements are equivalent:

- i) $\mathcal{A}$ is positive semidefinite.
- ii) $|x^* B y|^2 \leq (x^* A x)(y^* C y)$ for all $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$.
- iii) $2|x^* B y| \leq x^* A x + y^* C y$ for all $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$.

If, in addition, A and C are positive definite, then the following statement is equivalent to i)–iii):

$$\text{sprad}(B^* A^{-1} B C^{-1}) \leq 1.$$

Finally, if $\mathcal{A}$ is positive semidefinite and nonzero, then, for all $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$,

$$|x^* B y|^2 \leq \left[\frac{\lambda_{\max}(\mathcal{A}) - \lambda_{\min}(\mathcal{A})}{\lambda_{\max}(\mathcal{A}) + \lambda_{\min}(\mathcal{A})} \right]^2 (x^* A x)(y^* C y).$$

(Proof: See [709, p. 473] and [886, 1494].)

Fact 8.15.31. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let $x, y \in \mathbb{F}^n$, and assume that $x^* x = y^* y = 1$ and $x^* y = 0$. Then,

$$2|x^* A y| \leq \lambda_{\max}(A) - \lambda_{\min}(A).$$

Furthermore, there exist vectors $x, y \in \mathbb{F}^n$ satisfying $x^* x = y^* y = 1$ and $x^* y = 0$ for which equality holds. (Proof: See [886, 1494].) (Remark: $\lambda_{\max}(A) - \lambda_{\min}(A)$ is the *spread* of A . See Fact 9.9.30 and Fact 9.9.31.)

Fact 8.15.32. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is positive definite. Then,

$$\int_{\mathbb{R}^n} e^{-x^T A x} dx = \frac{\pi^{n/2}}{\sqrt{\det A}}.$$

Fact 8.15.33. Let $A \in \mathbb{R}^{n \times n}$, assume that A is positive definite, and define $f: \mathbb{R}^n \mapsto \mathbb{R}$ by

$$f(x) = \frac{e^{-\frac{1}{2}x^T A^{-1}x}}{(2\pi)^{n/2} \sqrt{\det A}}.$$

Then,

$$\begin{aligned} \int_{\mathbb{R}^n} f(x) \, dx &= 1, \\ \int_{\mathbb{R}^n} f(x) x x^T \, dx &= A, \end{aligned}$$

and

$$-\int_{\mathbb{R}^n} f(x) \log f(x) \, dx = \frac{1}{2} \log[(2\pi e)^n \det A].$$

(Proof: See [352] or use Fact 8.15.35.) (Remark: f is the multivariate normal density. The last expression is the *entropy*.)

Fact 8.15.34. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A and B are positive definite, and, for $k = 0, 1, 2, 3$, define

$$\mathcal{J}_k \triangleq \frac{1}{(2\pi)^{n/2} \sqrt{\det A}} \int_{\mathbb{R}^n} (x^T B x)^k e^{-\frac{1}{2}x^T A^{-1}x} \, dx.$$

Then,

$$\begin{aligned} \mathcal{J}_0 &= 1, \\ \mathcal{J}_1 &= \operatorname{tr} AB, \\ \mathcal{J}_2 &= (\operatorname{tr} AB)^2 + 2 \operatorname{tr} (AB)^2, \\ \mathcal{J}_3 &= (\operatorname{tr} AB)^3 + 6(\operatorname{tr} AB) [\operatorname{tr} (AB)^2] + 8 \operatorname{tr} (AB)^3. \end{aligned}$$

(Proof: See [1002, p. 80].) (Remark: These identities are *Lancaster's formulas*.)

Fact 8.15.35. Let $A \in \mathbb{R}^{n \times n}$, assume that A is positive definite, let $B \in \mathbb{R}^{n \times n}$, let $a, b \in \mathbb{R}^n$, and let $\alpha, \beta \in \mathbb{R}$. Then,

$$\begin{aligned} &\int_{\mathbb{R}^n} (x^T B x + b^T x + \beta) e^{-(x^T A x + a^T x + \alpha)} \, dx \\ &= \frac{\pi^{n/2}}{2\sqrt{\det A}} [2\beta + \operatorname{tr}(A^{-1}B) - b^T A^{-1}a + \frac{1}{2}a^T A^{-1}B A^{-1}a] e^{\frac{1}{4}a^T A^{-1}a - \alpha}. \end{aligned}$$

(Proof: See [654, p. 322].)

Fact 8.15.36. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a symmetric graph, where $\mathcal{X} = \{x_1, \dots, x_n\}$. Then, for all $z \in \mathbb{R}^n$, it follows that

$$z^T L z = \frac{1}{2} \sum (z_{(i)} - z_{(j)})^2,$$

where the sum is over the set $\{(i, j): (x_i, x_j) \in \mathcal{R}\}$. (Proof: See [269, pp. 29, 30] or [993].)

Fact 8.15.37. Let $n \leq 4$, let $A \in \mathbb{R}^{n \times n}$, assume that A is symmetric, and assume that, for all nonnegative vectors $x \in \mathbb{R}^n$, $x^T A x \geq 0$. Then, there exist $B, C \in \mathbb{R}^{n \times n}$ such that B is positive semidefinite, C is symmetric and nonnegative, and $A = B + C$. (Remark: The result does not hold for all $n > 5$. Hence, this result is an example of the *quartic barrier*. See [351], Fact 8.14.7, and Fact 11.17.3.) (Remark: A is *copositive*.)

8.16 Facts on Simultaneous Diagonalization

Fact 8.16.1. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian. Then, the following statements are equivalent:

- i) There exists a unitary matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and SBS^* are diagonal.
- ii) $AB = BA$.
- iii) AB and BA are Hermitian.

If, in addition, A is nonsingular, then the following condition is equivalent to i)–iii):

- iv) $A^{-1}B$ is Hermitian.

(Proof: See [174, p. 208], [447, pp. 188–190], or [709, p. 229].) (Remark: The equivalence of i) and ii) is given by Fact 5.17.7.)

Fact 8.16.2. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that A is nonsingular. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and SBS^* are diagonal if and only if $A^{-1}B$ is diagonalizable over $\mathbb{R}$. (Proof: See [709, p. 229] or [1098, p. 95].)

Fact 8.16.3. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are symmetric, and assume that A is nonsingular. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^T and SBS^T are diagonal if and only if $A^{-1}B$ is diagonalizable. (Proof: See [709, p. 229] and [1352].) (Remark: A and B are complex symmetric.)

Fact 8.16.4. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and SBS^* are diagonal if and only if there exists a positive-definite matrix $M \in \mathbb{F}^{n \times n}$ such that $AMB = BMA$. (Proof: See [83].)

Fact 8.16.5. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume there exist $\alpha, \beta \in \mathbb{R}$ such that $\alpha A + \beta B$ is positive definite. Then, there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and SBS^* are diagonal. (Proof: See [709, p. 465].) (Remark: This result extends a result due to Weierstrass. See [1352].) (Remark: Suppose that B is positive definite. Then, by necessity of Fact 8.16.2, it follows that $A^{-1}B$ is diagonalizable over $\mathbb{R}$, which proves iii) $\implies$ i) of Proposition 5.5.12.) (Remark: See Fact 8.16.6.)

Fact 8.16.6. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, assume that $\{x \in \mathbb{F}^n: x^* A x = x^* B x = 0\} = \{0\}$, and, if $\mathbb{F} = \mathbb{R}$, assume that $n \geq 3$. Then,

there exists a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that SAS^* and SBS^* are diagonal. (Proof: The result follows from Fact 5.17.9. See [950] or [1098, p. 96].) (Remark: For $\mathbb{F} = \mathbb{R}$, this result is due to Pesonen and Milnor. See [1352].) (Remark: See Fact 5.17.9, Fact 8.15.24, Fact 8.15.25, and Fact 8.16.5.)

8.17 Facts on Eigenvalues and Singular Values for One Matrix

Fact 8.17.1. Let $A = \begin{bmatrix} a & b \\ b & c \end{bmatrix} \in \mathbb{F}^{2 \times 2}$, assume that A is Hermitian, and let $\text{mspec}(A) = \{\lambda_1, \lambda_2\}_{\text{ms}}$. Then,

$$2|b| \leq \lambda_1 - \lambda_2.$$

Now, assume that A is positive semidefinite. Then,

$$\sqrt{2}|b| \leq \left(\sqrt{\lambda_1} - \sqrt{\lambda_2}\right)\sqrt{\lambda_1 + \lambda_2}.$$

If $c > 0$, then

$$\frac{|b|}{\sqrt{c}} \leq \sqrt{\lambda_1} - \sqrt{\lambda_2}.$$

If $a > 0$ and $c > 0$, then

$$\frac{|b|}{\sqrt{ac}} \leq \frac{\lambda_1 - \lambda_2}{\lambda_1 + \lambda_2}.$$

Finally, if A is positive definite, then

$$\frac{|b|}{a} \leq \frac{\lambda_1 - \lambda_2}{2\sqrt{\lambda_1\lambda_2}}$$

and

$$4|b| \leq \frac{\lambda_1^2 - \lambda_2^2}{\sqrt{\lambda_1\lambda_2}}.$$

(Proof: See [886, 1494].) (Remark: These inequalities are useful for deriving inequalities involving quadratic forms. See Fact 8.15.29 and Fact 8.15.30.)

Fact 8.17.2. Let $A \in \mathbb{F}^{n \times m}$. Then, for all $i = 1, \dots, \min\{n, m\}$,

$$\lambda_i(\langle A \rangle) = \sigma_i(A).$$

Hence,

$$\text{tr} \langle A \rangle = \sum_{i=1}^{\min\{n, m\}} \sigma_i(A).$$

Fact 8.17.3. Let $A \in \mathbb{F}^{n \times n}$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} \sigma_{\max}(A)I & A^* \\ A & \sigma_{\max}(A)I \end{bmatrix}.$$

Then, $\mathcal{A}$ is positive semidefinite. Furthermore,

$$\langle A + A^* \rangle \leq \left\{ \begin{array}{c} \langle A \rangle + \langle A^* \rangle \leq 2\sigma_{\max}(A)I \\ A^*A + I \end{array} \right\} \leq [\sigma_{\max}^2(A) + 1]I.$$

(Proof: See [1492].)

Fact 8.17.4. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $i = 1, \dots, n$,

$$-\sigma_i(A) \leq \lambda_i \left[\frac{1}{2}(A + A^*) \right] \leq \sigma_i(A).$$

Hence,

$$|\operatorname{tr} A| \leq \operatorname{tr} \langle A \rangle.$$

(Proof: See [1211].) (Remark: See Fact 5.11.25.)

Fact 8.17.5. Let $A \in \mathbb{F}^{n \times n}$, and let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_n$ are ordered such that $|\lambda_1| \geq \dots \geq |\lambda_n|$. If $p > 0$, then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k |\lambda_i|^p \leq \sum_{i=1}^k \sigma_i^p(A).$$

In particular, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k |\lambda_i| \leq \sum_{i=1}^k \sigma_i(A).$$

Hence,

$$|\operatorname{tr} A| \leq \sum_{i=1}^n |\lambda_i| \leq \sum_{i=1}^n \sigma_i(A) = \operatorname{tr} \langle A \rangle.$$

Furthermore, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k |\lambda_i|^2 \leq \sum_{i=1}^k \sigma_i^2(A).$$

Hence,

$$\operatorname{Re} \operatorname{tr} A^2 \leq |\operatorname{tr} A^2| \leq \sum_{i=1}^n |\lambda_i|^2 \leq \sum_{i=1}^n \sigma_i(A^2) = \operatorname{tr} \langle A^2 \rangle \leq \sum_{i=1}^n \sigma_i^2(A) = \operatorname{tr} A^* A.$$

(Proof: The result follows from Fact 5.11.28 and Fact 2.21.13. See [197, p. 42], [711, p. 176], or [1485, p. 19]. See Fact 9.13.17 for the inequality $\operatorname{tr} \langle A^2 \rangle = \operatorname{tr} (A^2 A^2)^{1/2} \leq \operatorname{tr} A^* A$.) Furthermore,

$$\sum_{i=1}^n |\lambda_i|^2 = \operatorname{tr} A^* A$$

if and only if A is normal. (Proof: See Fact 5.14.15.) Finally,

$$\sum_{i=1}^n \lambda_i^2 = \operatorname{tr} A^* A$$

if and only if A is Hermitian. (Proof: See Fact 3.7.13.) (Remark: The first result is *Weyl's inequalities*. The result $\sum_{i=1}^n |\lambda_i|^2 \leq \operatorname{tr} A^* A$ is *Schur's inequality*. See Fact 9.11.3.) (Problem: Determine when equality holds for the remaining inequalities.)

Fact 8.17.6. Let $A \in \mathbb{F}^{n \times n}$, let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_n$ are ordered such that $|\lambda_1| \geq \dots \geq |\lambda_n|$, and let $r > 0$. Then, for all $k = 1, \dots, n$,

$$\prod_{i=1}^k (1 + r|\lambda_i|) \leq \prod_{i=1}^k [1 + \sigma_i(A)].$$

(Proof: See [447, p. 222].)

Fact 8.17.7. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$|\operatorname{tr} A^2| \leq \begin{cases} \operatorname{tr} \langle A \rangle \langle A^* \rangle \\ \operatorname{tr} \langle A^2 \rangle \leq \operatorname{tr} \langle A \rangle^2 = \operatorname{tr} A^* A. \end{cases}$$

(Proof: For the upper inequality, see [886, 1494]. For the lower inequalities, use Fact 8.17.4 and Fact 9.11.3.) (Remark: See Fact 5.11.10, Fact 9.13.17, and Fact 9.13.18.)

Fact 8.17.8. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k d_i(A) \leq \sum_{i=1}^k \lambda_i(A)$$

with equality for $k = n$, that is,

$$\operatorname{tr} A = \sum_{i=1}^n d_i(A) = \sum_{i=1}^n \lambda_i(A).$$

That is, $[\lambda_1(A) \ \cdots \ \lambda_n(A)]^T$ strongly majorizes $[d_1(A) \ \cdots \ d_n(A)]^T$, and thus, for all $k = 1, \dots, n$,

$$\sum_{i=k}^n \lambda_i(A) \leq \sum_{i=k}^n d_i(A).$$

In particular,

$$\lambda_{\min}(A) \leq d_{\min}(A) \leq d_{\max}(A) \leq \lambda_{\max}(A).$$

Furthermore, the vector $[d_1(A) \ \cdots \ d_n(A)]^T$ is an element of the convex hull of the $n!$ vectors obtaining by permuting the components of $[\lambda_1(A) \ \cdots \ \lambda_n(A)]^T$. (Proof: See [197, p. 35], [709, p. 193], [971, p. 218], or [1485, p. 18]. The last statement follows from Fact 2.21.7.) (Remark: This result is *Schur's theorem*.) (Remark: See Fact 8.12.3.)

Fact 8.17.9. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, let k denote the number of positive diagonal entries of A , and let l denote the number of positive eigenvalues of A . Then,

$$\sum_{i=1}^k d_i^2(A) \leq \sum_{i=1}^l \lambda_i^2(A).$$

(Proof: Write $A = B + C$, where B is positive semidefinite, C is negative semidefinite, and $\operatorname{mspec}(A) = \operatorname{mspec}(B) \cup \operatorname{mspec}(C)$. Furthermore, without loss of gener-

ality, assume that $A_{(1,1)}, \dots, A_{(k,k)}$ are the positive diagonal entries of A . Then,

$$\begin{aligned} \sum_{i=1}^k d_i^2(A) &= \sum_{i=1}^k A_{(i,i)}^2 \leq \sum_{i=1}^k (A_{(i,i)} - C_{(i,i)})^2 \\ &= \sum_{i=1}^k B_{(i,i)}^2 \leq \sum_{i=1}^n B_{(i,i)}^2 \leq \operatorname{tr} B^2 = \sum_{i=1}^l \lambda_i^2(A). \end{aligned}$$

(Remark: This inequality can be written as

$$\operatorname{tr} (A + |A|)^{\circ 2} \leq \operatorname{tr} (A + \langle A \rangle)^2.$$

(Remark: This result is due to Y. Li.)

Fact 8.17.10. Let $x, y \in \mathbb{R}^n$, where $n \geq 2$. Then, the following statements are equivalent:

- i) y strongly majorizes by x .
- ii) x is an element of the convex hull of the vectors $y_1, \dots, y_{n!} \in \mathbb{R}^n$, where each of these $n!$ vectors is formed by permuting the components of y .
- iii) There exists a Hermitian matrix $A \in \mathbb{C}^{n \times n}$ such that $\begin{bmatrix} A_{(1,1)} & \cdots & A_{(n,n)} \end{bmatrix}^T = x$ and $\operatorname{mspec}(A) = \{y_{(1)}, \dots, y_{(n)}\}_{\operatorname{ms}}$.

(Remark: This result is the *Schur-Horn theorem*. Schur's theorem given by Fact 8.17.8 is $\text{iii}) \implies \text{i})$, while the result $\text{i}) \implies \text{iii})$ is due to [708]. The equivalence of $\text{ii})$ is given by Fact 2.21.7. The significance of this result is discussed in [153, 198, 262].)

(Remark: An equivalent version is given by Fact 3.11.19.)

Fact 8.17.11. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, for all $k = 1, \dots, n$,

$$\prod_{i=k}^n \lambda_i(A) \leq \prod_{i=k}^n d_i(A).$$

In particular,

$$\det A \leq \prod_{i=1}^n A_{(i,i)}.$$

Now, assume that A is positive definite. Then, equality holds if and only if A is diagonal. (Proof: See [530, pp. 21–24], [709, pp. 200, 477], or [1485, p. 18].)

(Remark: The case $k = 1$ is *Hadamard's inequality*.) (Remark: See Fact 8.13.34 and Fact 9.11.1.) (Remark: A strengthened version is given by Fact 8.13.33.) (Remark: A geometric interpretation is discussed in [539].)

Fact 8.17.12. Let $A \in \mathbb{F}^{n \times n}$, define $H \triangleq \frac{1}{2}(A + A^*)$ and $S \triangleq \frac{1}{2}(A - A^*)$, and assume that H is positive definite. Then, the following statements hold:

- i) A is nonsingular.
- ii) $\frac{1}{2}(A^{-1} + A^{-*}) = (H + S^*H^{-1}S)^{-1}$.
- iii) $\sigma_{\max}(A^{-1}) \leq \sigma_{\max}(H^{-1})$.
- iv) $\sigma_{\max}(A) \leq \sigma_{\max}(H + S^*H^{-1}S)$.

(Proof: See [978].) (Remark: See Fact 8.9.31 and Fact 8.13.11.)

Fact 8.17.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, $\{A_{(1,1)}, \dots, A_{(n,n)}\}_{\text{ms}} = \text{mspec}(A)$ if and only if A is diagonal. (Proof: Apply Fact 8.17.11 with $A + \beta I > 0$.)

Fact 8.17.14. Let $A \in \mathbb{F}^{n \times n}$. Then, $\begin{bmatrix} I & A \\ A^* & I \end{bmatrix}$ is positive semidefinite if and only if $\sigma_{\max}(A) \leq 1$. Furthermore, $\begin{bmatrix} I & A \\ A^* & I \end{bmatrix}$ is positive definite if and only if $\sigma_{\max}(A) < 1$. (Proof: Note that

$$\begin{bmatrix} I & A \\ A^* & I \end{bmatrix} = \begin{bmatrix} I & 0 \\ A^* & I \end{bmatrix} \begin{bmatrix} I & 0 \\ 0 & I - A^*A \end{bmatrix} \begin{bmatrix} I & A \\ 0 & I \end{bmatrix}.$$

Fact 8.17.15. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \lambda_i = \max\{\text{tr } S^*AS : S \in \mathbb{F}^{n \times k} \text{ and } S^*S = I_k\}$$

and

$$\sum_{i=n+1-k}^n \lambda_i = \min\{\text{tr } S^*AS : S \in \mathbb{F}^{n \times k} \text{ and } S^*S = I_k\}.$$

(Proof: See [709, p. 191].) (Remark: This result is the *minimum principle*.)

Fact 8.17.16. Let $A \in \mathbb{F}^{n \times n}$, assume that A is Hermitian, and let $S \in \mathbb{R}^{k \times n}$ satisfy $SS^* = I_k$. Then, for all $i = 1, \dots, k$,

$$\lambda_{i+n-k}(A) \leq \lambda_i(SAS^*) \leq \lambda_i(A).$$

Consequently,

$$\sum_{i=1}^k \lambda_{i+n-k}(A) \leq \text{tr } SAS^* \leq \sum_{i=1}^k \lambda_i(A)$$

and

$$\prod_{i=1}^k \lambda_{i+n-k}(A) \leq \det SAS^* \leq \prod_{i=1}^k \lambda_i(A).$$

(Proof: See [709, p. 190].) (Remark: This result is the *Poincaré separation theorem*.)

8.18 Facts on Eigenvalues and Singular Values for Two or More Matrices

Fact 8.18.1. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and assume that A and C are positive definite. Then, $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ is positive semidefinite if and only if

$$\sigma_{\max}(A^{-1/2}BC^{-1/2}) \leq 1.$$

Furthermore, $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ is positive definite if and only if

$$\sigma_{\max}(A^{-1/2}BC^{-1/2}) < 1.$$

(Proof: See [964].)

Fact 8.18.2. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, assume that A and C are positive definite, and assume that

$$\sigma_{\max}^2(B) \leq \sigma_{\min}(A)\sigma_{\min}(C).$$

Then, $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ is positive semidefinite. If, in addition,

$$\sigma_{\max}^2(B) < \sigma_{\min}(A)\sigma_{\min}(C),$$

then $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ is positive definite. (Proof: Note that

$$\begin{aligned} \sigma_{\max}^2(A^{-1/2}BC^{-1/2}) &\leq \lambda_{\max}(A^{-1/2}BC^{-1}B^*A^{-1/2}) \\ &\leq \sigma_{\max}(C^{-1})\lambda_{\max}(A^{-1/2}BB^*A^{-1/2}) \\ &\leq \frac{1}{\sigma_{\min}(C)}\lambda_{\max}(B^*A^{-1}B) \\ &\leq \frac{\sigma_{\max}(A^{-1})}{\sigma_{\min}(C)}\lambda_{\max}(B^*B) \\ &= \frac{1}{\sigma_{\min}(A)\sigma_{\min}(C)}\sigma_{\max}^2(B) \\ &\leq 1. \end{aligned}$$

The result now follows from Fact 8.18.1.)

Fact 8.18.3. Let $A, B \in \mathbb{F}^n$, assume that A and B are Hermitian, and define $\gamma \triangleq [\gamma_1 \cdots \gamma_n]$, where the components of γ are the components of $[\lambda_1(A) \cdots \lambda_n(A)] + [\lambda_n(B) \cdots \lambda_1(B)]$ arranged in decreasing order. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \gamma_i \leq \sum_{i=1}^k \lambda_i(A+B).$$

(Proof: The result follows from the Lidskii-Wielandt inequalities. See [197, p. 71] or [198, 380].) (Remark: This result provides an alternative lower bound for (8.6.12).)

Fact 8.18.4. Let $A, B \in \mathbf{H}^n$, let $k \in \{1, \dots, n\}$, and let $1 \leq i_1 \leq \cdots \leq i_k \leq n$. Then,

$$\sum_{j=1}^k \lambda_{i_j}(A) + \sum_{i=1}^k \lambda_{n-k+j}(B) \leq \sum_{j=1}^k \lambda_{i_j}(A+B) \leq \sum_{j=1}^k [\lambda_{i_j}(A) + \lambda_j(B)].$$

(Proof: See [1177, pp. 115, 116].)

Fact 8.18.5. Let $f: \mathbb{R} \mapsto \mathbb{R}$ be convex, define $f: \mathbf{H}^n \mapsto \mathbf{H}^n$ by (8.5.1), let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, for all $\alpha \in [0, 1]$,

$$[\alpha \lambda_1[f(A)] + (1-\alpha)\lambda_1[f(B)] \quad \cdots \quad \alpha \lambda_n[f(A)] + (1-\alpha)\lambda_n[f(B)]]$$

weakly majorizes

$$\begin{bmatrix} \lambda_1[f(\alpha A + (1 - \alpha)B)] & \cdots & \lambda_n[f(\alpha A + (1 - \alpha)B)] \end{bmatrix}.$$

If, in addition, f is either nonincreasing or nondecreasing, then, for all $i = 1, \dots, n$,

$$\lambda_i[f(\alpha A + (1 - \alpha)B)] \leq \alpha \lambda_i[f(A)] + (1 - \alpha) \lambda_i[f(B)].$$

(Proof: See [91].) (Remark: Convexity of $f: \mathbb{R} \mapsto \mathbb{R}$ does not imply convexity of $f: \mathbf{H}^n \mapsto \mathbf{H}^n$.)

Fact 8.18.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. If $r \in [0, 1]$, then

$$\begin{bmatrix} \lambda_1(A^r + B^r) & \cdots & \lambda_n(A^r + B^r) \end{bmatrix}$$

weakly majorizes

$$\begin{bmatrix} \lambda_1[(A + B)^r] & \cdots & \lambda_n[(A + B)^r] \end{bmatrix},$$

and, for all $i = 1, \dots, n$,

$$2^{1-r} \lambda_i[(A + B)^r] \leq \lambda_i(A^r + B^r).$$

If $r \geq 1$, then

$$\begin{bmatrix} \lambda_1[(A + B)^r] & \cdots & \lambda_n[(A + B)^r] \end{bmatrix}$$

weakly majorizes

$$\begin{bmatrix} \lambda_1(A^r + B^r) & \cdots & \lambda_n(A^r + B^r) \end{bmatrix},$$

and, for all $i = 1, \dots, n$,

$$\lambda_i(A^r + B^r) \leq 2^{r-1} \lambda_i[(A + B)^r].$$

(Proof: The result follows from Fact 8.18.5. See [58, 89, 91].)

Fact 8.18.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \sigma_i^2(A + jB) \leq \sum_{i=1}^k [\sigma_i^2(A) + \sigma_i^2(B)],$$

$$\sum_{i=1}^n \sigma_i^2(A + jB) = \sum_{i=1}^n [\sigma_i^2(A) + \sigma_i^2(B)],$$

$$\sum_{i=1}^k [\sigma_i^2(A + jB) + \sigma_{n-i}^2(A + jB)] \leq \sum_{i=1}^k [\sigma_i^2(A) + \sigma_i^2(B)],$$

$$\sum_{i=1}^n [\sigma_i^2(A + jB) + \sigma_{n-i}^2(A + jB)] = \sum_{i=1}^n [\sigma_i^2(A) + \sigma_i^2(B)],$$

and

$$\sum_{i=1}^k [\sigma_i^2(A) + \sigma_{n-i}^2(B)] \leq \sum_{i=1}^k \sigma_i^2(A + jB),$$

$$\sum_{i=1}^n [\sigma_i^2(A) + \sigma_{n-i}^2(B)] = \sum_{i=1}^n \sigma_i^2(A + jB).$$

(Proof: See [52, 320].) (Remark: The first identity is given by Fact 9.9.40.)

Fact 8.18.8. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, the following statements hold:

i) If $p \in [0, 1]$, then

$$\sigma_{\max}(A^p - B^p) \leq \sigma_{\max}^p(A - B).$$

ii) If $p \geq \sqrt{2}$, then

$$\sigma_{\max}(A^p - B^p) \leq p[\max\{\sigma_{\max}(A), \sigma_{\max}(B)\}]^{p-1} \sigma_{\max}(A - B).$$

iii) If a and b are positive numbers such that $aI \leq A \leq bI$ and $aI \leq B \leq bI$, then

$$\sigma_{\max}(A^p - B^p) \leq b[b^{p-2} + (p-1)a^{p-2}] \sigma_{\max}(A - B).$$

(Proof: See [206, 816].)

Fact 8.18.9. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, for all $i = 1, \dots, n$,

$$\sigma_i(A - B) \leq \sigma_i\left(\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix}\right).$$

(Proof: See [1255, 1483].)

Fact 8.18.10. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and assume that $\mathcal{A} \in \mathbb{F}^{(n+m) \times (n+m)}$ defined by

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$$

is positive semidefinite. Then, for all $i = 1, \dots, \min\{n, m\}$,

$$2\sigma_i(B) \leq \sigma_i(\mathcal{A}).$$

(Proof: See [215, 1255].)

Fact 8.18.11. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\max\{\sigma_{\max}^2(A), \sigma_{\max}^2(B)\} - \sigma_{\max}(AB) \leq \sigma_{\max}(A^*A - BB^*)$$

and

$$\sigma_{\max}(A^*A - BB^*) \leq \max\{\sigma_{\max}^2(A), \sigma_{\max}^2(B)\} - \min\{\sigma_{\min}^2(A), \sigma_{\min}^2(B)\}.$$

Furthermore,

$$\max\{\sigma_{\max}^2(A), \sigma_{\max}^2(B)\} + \min\{\sigma_{\min}^2(A), \sigma_{\min}^2(B)\} \leq \sigma_{\max}(A^*A + BB^*)$$

and

$$\sigma_{\max}(A^*A + BB^*) \leq \max\{\sigma_{\max}^2(A), \sigma_{\max}^2(B)\} + \sigma_{\max}(AB).$$

Now, assume that A and B are positive semidefinite. Then,

$$\max\{\lambda_{\max}(A), \lambda_{\max}(B)\} - \sigma_{\max}(A^{1/2}B^{1/2}) \leq \sigma_{\max}(A - B)$$

and

$$\sigma_{\max}(A - B) \leq \max\{\lambda_{\max}(A), \lambda_{\max}(B)\} - \min\{\lambda_{\min}(A), \lambda_{\min}(B)\}.$$

Furthermore,

$$\max\{\lambda_{\max}(A), \lambda_{\max}(B)\} + \min\{\lambda_{\min}(A), \lambda_{\min}(B)\} \leq \lambda_{\max}(A + B)$$

and

$$\lambda_{\max}(A + B) \leq \max\{\lambda_{\max}(A), \lambda_{\max}(B)\} + \sigma_{\max}(A^{1/2}B^{1/2}).$$

(Proof: See [824, 1486].) (Remark: See Fact 8.18.14 and Fact 9.13.8.)

Fact 8.18.12. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\begin{aligned} \max\{\sigma_{\max}(A), \sigma_{\max}(B)\} - \sigma_{\max}(A^{1/2}B^{1/2}) & \\ & \leq \sigma_{\max}(A - B) \\ & \leq \max\{\sigma_{\max}(A), \sigma_{\max}(B)\} \\ & \leq \sigma_{\max}(A + B) \\ & \leq \left\{ \begin{array}{c} \max\{\sigma_{\max}(A), \sigma_{\max}(B)\} + \sigma_{\max}(A^{1/2}B^{1/2}) \\ \sigma_{\max}(A) + \sigma_{\max}(B) \end{array} \right\} \\ & \leq 2 \max\{\sigma_{\max}(A), \sigma_{\max}(B)\}. \end{aligned}$$

(Proof: See [818, 824] and use Fact 8.18.13.) (Remark: See Fact 8.18.14.)

Fact 8.18.13. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite, and let $k \geq 1$. Then, for all $i = 1, \dots, n$,

$$2\sigma_i \left[A^{1/2}(A + B)^{k-1}B^{1/2} \right] \leq \lambda_i[(A + B)^k].$$

Hence,

$$2\sigma_{\max}(A^{1/2}B^{1/2}) \leq \lambda_{\max}(A + B)$$

and

$$\sigma_{\max}(A^{1/2}B^{1/2}) \leq \max\{\lambda_{\max}(A), \lambda_{\max}(B)\}.$$

(Proof: See Fact 8.18.11 and Fact 9.9.18.)

Fact 8.18.14. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\max\{\lambda_{\max}(A), \lambda_{\max}(B)\} - \sigma_{\max}(A^{1/2}B^{1/2}) \leq \sigma_{\max}(A - B)$$

and

$$\begin{aligned}
& \lambda_{\max}(A + B) \\
& \leq \frac{1}{2} \left[\lambda_{\max}(A) + \lambda_{\max}(B) + \sqrt{[\lambda_{\max}(A) - \lambda_{\max}(B)]^2 + 4\sigma_{\max}^2(A^{1/2}B^{1/2})} \right] \\
& \leq \begin{cases} \max\{\lambda_{\max}(A), \lambda_{\max}(B)\} + \sigma_{\max}(A^{1/2}B^{1/2}) \\ \lambda_{\max}(A) + \lambda_{\max}(B). \end{cases}
\end{aligned}$$

Furthermore,

$$\lambda_{\max}(A + B) = \lambda_{\max}(A) + \lambda_{\max}(B)$$

if and only if

$$\sigma_{\max}(A^{1/2}B^{1/2}) = \lambda_{\max}^{1/2}(A)\lambda_{\max}^{1/2}(B).$$

(Proof: See [818, 821, 824].) (Remark: See Fact 8.18.11, Fact 8.18.12, Fact 9.14.15, and Fact 9.9.46.) (Problem: Is $\sigma_{\max}(A - B) \leq \sigma_{\max}(A + B)$?)

Fact 8.18.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\sigma_{\max}(A^{1/2}B^{1/2}) \leq \sigma_{\max}^{1/2}(AB).$$

Equivalently,

$$\lambda_{\max}(A^{1/2}BA^{1/2}) \leq \lambda_{\max}^{1/2}(AB^2A).$$

Furthermore, $AB = 0$ if and only if $A^{1/2}B^{1/2} = 0$. (Proof: See [818] and [824].)

Fact 8.18.16. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\begin{aligned}
\operatorname{tr} AB & \leq \operatorname{tr} (AB^2A)^{1/2} \leq \frac{1}{4} \operatorname{tr} (A + B)^2, \\
\operatorname{tr} (AB)^2 & \leq \operatorname{tr} A^2B^2 \leq \frac{1}{16} \operatorname{tr} (A + B)^4,
\end{aligned}$$

and

$$\begin{aligned}
\sigma_{\max}(AB) & \leq \frac{1}{4} \sigma_{\max}[(A + B)^2] \\
& \leq \left\{ \begin{array}{l} \frac{1}{2} \sigma_{\max}(A^2 + B^2) \leq \frac{1}{2} \sigma_{\max}(A^2) + \frac{1}{2} \sigma_{\max}(B^2) \\ \frac{1}{4} \sigma_{\max}^2(A + B) \leq \frac{1}{4} [\sigma_{\max}(A) + \sigma_{\max}(B)]^2 \end{array} \right\} \\
& \leq \frac{1}{2} \sigma_{\max}^2(A) + \frac{1}{2} \sigma_{\max}^2(B).
\end{aligned}$$

(Proof: See Fact 9.9.18. The inequalities $\operatorname{tr} AB \leq \operatorname{tr} (AB^2A)^{1/2}$ and $\operatorname{tr} (AB)^2 \leq \operatorname{tr} A^2B^2$ follow from Fact 8.12.20.)

Fact 8.18.17. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and assume that B is positive definite. Then, for all $i, j, k \in \{1, \dots, n\}$ such that $j + k \leq i + 1$,

$$\lambda_i(AB) \leq \lambda_j(A)\lambda_k(B)$$

and

$$\lambda_{n-j+1}(A)\lambda_{n-k+1}(B) \leq \lambda_{n-i+1}(AB).$$

In particular, for all $i = 1, \dots, n$,

$$\lambda_i(A)\lambda_n(B) \leq \lambda_i(AB) \leq \lambda_i(A)\lambda_1(B).$$

(Proof: See [1177, pp. 126, 127].)

Fact 8.18.18. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and assume that B is Hermitian. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \lambda_i(A) \lambda_{n-i+1}(B) \leq \sum_{i=1}^k \lambda_i(AB)$$

and

$$\sum_{i=1}^k \lambda_{n-i+1}(AB) \leq \sum_{i=1}^k \lambda_i(A) \lambda_i(B).$$

In particular,

$$\sum_{i=1}^k \lambda_i(A) \lambda_{n-i+1}(B) \leq \operatorname{tr} AB \leq \sum_{i=1}^n \lambda_i(A) \lambda_i(B).$$

(Proof: See [838].) (Remark: See Fact 5.12.4, Fact 5.12.5, Fact 5.12.8, and Proposition 8.4.13.) (Remark: The upper and lower bounds for $\operatorname{tr} AB$ are related to Fact 1.16.4. See [200, p. 140].)

Fact 8.18.19. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let $\lambda_1(AB) \geq \dots \geq \lambda_n(AB) \geq 0$ denote the eigenvalues of AB , and let $1 \leq l_1 < \dots < l_k \leq n$. Then,

$$\sum_{i=1}^k \lambda_{l_i}(A) \lambda_{n-i+1}(B) \leq \sum_{i=1}^k \lambda_{l_i}(AB) \leq \sum_{i=1}^k \lambda_{l_i}(A) \lambda_i(B).$$

Furthermore,

$$\sum_{i=1}^k \lambda_{l_i}(A) \lambda_{n-l_i+1}(B) \leq \sum_{i=1}^k \lambda_i(AB).$$

In particular,

$$\sum_{i=1}^k \lambda_i(A) \lambda_{n-i+1}(B) \leq \sum_{i=1}^k \lambda_i(AB) \leq \sum_{i=1}^k \lambda_i(A) \lambda_i(B).$$

(Proof: See [1388].) (Remark: See Fact 8.18.22 and Fact 9.14.27.)

Fact 8.18.20. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. If $p \geq 1$, then

$$\sum_{i=1}^n \lambda_i^p(A) \lambda_{n-i+1}^p(B) \leq \operatorname{tr} \left(B^{1/2} A B^{1/2} \right)^p \leq \operatorname{tr} A^p B^p \leq \sum_{i=1}^n \lambda_i^p(A) \lambda_i^p(B).$$

If $0 \leq p \leq 1$, then

$$\sum_{i=1}^n \lambda_i^p(A) \lambda_{n-i+1}^p(B) \leq \operatorname{tr} A^p B^p \leq \operatorname{tr} \left(B^{1/2} A B^{1/2} \right)^p \leq \sum_{i=1}^n \lambda_i^p(A) \lambda_i^p(B).$$

Now, suppose that A and B are positive definite. If $p \leq -1$, then

$$\sum_{i=1}^n \lambda_i^p(A) \lambda_{n-i+1}^p(B) \leq \operatorname{tr} \left(B^{1/2} A B^{1/2} \right)^p \leq \operatorname{tr} A^p B^p \leq \sum_{i=1}^n \lambda_i^p(A) \lambda_i^p(B).$$

If $-1 \leq p \leq 0$, then

$$\sum_{i=1}^n \lambda_i^p(A) \lambda_{n-i+1}^p(B) \leq \operatorname{tr} A^p B^p \leq \operatorname{tr} \left(B^{1/2} A B^{1/2} \right)^p \leq \sum_{i=1}^n \lambda_i^p(A) \lambda_i^p(B).$$

(Proof: See [1389]. See also [278, 881, 909, 1392].) (Remark: See Fact 8.12.20. See Fact 8.12.15 for the indefinite case.)

Fact 8.18.21. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, for all $k = 1, \dots, n$,

$$\prod_{i=1}^k \lambda_i(AB) \leq \prod_{i=1}^k \sigma_i(AB) \leq \prod_{i=1}^k \lambda_i(A) \lambda_i(B)$$

with equality for $k = n$. Furthermore, for all $k = 1, \dots, n$,

$$\prod_{i=k}^n \lambda_i(A) \lambda_i(B) \leq \prod_{i=k}^n \sigma_i(AB) \leq \prod_{i=k}^n \lambda_i(AB)$$

with equality for $k = 1$. (Proof: Use Fact 5.11.28 and Fact 9.13.19.)

Fact 8.18.22. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let $\lambda_1(AB) \geq \dots \geq \lambda_n(AB) \geq 0$ denote the eigenvalues of AB , and let $1 \leq l_1 < \dots < l_k \leq n$. Then,

$$\prod_{i=1}^k \lambda_{l_i}(AB) \leq \prod_{i=1}^k \lambda_{l_i}(A) \lambda_{l_i}(B)$$

with equality for $k = n$. Furthermore,

$$\prod_{i=1}^k \lambda_{l_i}(A) \lambda_{n-l_i+1}(B) \leq \prod_{i=1}^k \lambda_i(AB)$$

with equality for $k = n$. In particular,

$$\prod_{i=1}^k \lambda_i(A) \lambda_{n-i+1}(B) \leq \prod_{i=1}^k \lambda_i(AB) \leq \prod_{i=1}^k \lambda_i(A) \lambda_i(B)$$

with equality for $k = n$. (Proof: See [1388].) (Remark: See Fact 8.18.19 and Fact 9.14.27.)

Fact 8.18.23. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $\lambda \in \operatorname{spec}(A)$. Then,

$$\frac{2}{n} \left[\frac{\lambda_{\min}^2(A) \lambda_{\min}^2(B)}{\lambda_{\min}^2(A) + \lambda_{\min}^2(B)} \right] < \lambda < \frac{n}{2} [\lambda_{\max}^2(A) + \lambda_{\max}^2(B)].$$

(Proof: See [729].)

Fact 8.18.24. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and define

$$k_A \triangleq \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)}, \quad k_B \triangleq \frac{\lambda_{\max}(B)}{\lambda_{\min}(B)},$$

and

$$\gamma \triangleq \frac{(\sqrt{k_A} + 1)^2}{\sqrt{k_A}} - \frac{k_B(\sqrt{k_A} - 1)^2}{\sqrt{k_A}}.$$

Then, if $\gamma < 0$, then

$$\frac{1}{2}\lambda_{\max}(A)\lambda_{\max}(B)\gamma \leq \lambda_{\min}(AB + BA) \leq \lambda_{\max}(AB + BA) \leq 2\lambda_{\max}(A)\lambda_{\max}(B),$$

whereas, if $\gamma > 0$, then

$$\frac{1}{2}\lambda_{\min}(A)\lambda_{\min}(B)\gamma \leq \lambda_{\min}(AB + BA) \leq \lambda_{\max}(AB + BA) \leq 2\lambda_{\max}(A)\lambda_{\max}(B).$$

Furthermore, if

$$\sqrt{k_A k_B} < 1 + \sqrt{k_A} + \sqrt{k_B},$$

then $AB + BA$ is positive definite. (Proof: See [1038].)

Fact 8.18.25. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive definite, assume that B is positive semidefinite, and let $\alpha > 0$ and $\beta > 0$ be such that $\alpha I \leq A \leq \beta I$. Then,

$$\sigma_{\max}(AB) \leq \frac{\alpha + \beta}{2\sqrt{\alpha\beta}} \text{sprad}(AB) \leq \frac{\alpha + \beta}{2\sqrt{\alpha\beta}} \sigma_{\max}(AB).$$

In particular,

$$\sigma_{\max}(A) \leq \frac{\alpha + \beta}{2\sqrt{\alpha\beta}} \text{sprad}(A) \leq \frac{\alpha + \beta}{2\sqrt{\alpha\beta}} \sigma_{\max}(A).$$

(Proof: See [1312].) (Remark: The left-hand inequality is tightest for $\alpha = \lambda_{\min}(A)$ and $\beta = \lambda_{\max}(A)$.) (Remark: This result is due to Bourin.)

Fact 8.18.26. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, the following statements hold:

i) If $q \in [0, 1]$, then

$$\sigma_{\max}(A^q B^q) \leq \sigma_{\max}^q(AB)$$

and

$$\sigma_{\max}(B^q A^q B^q) \leq \sigma_{\max}^q(BAB).$$

ii) If $q \in [0, 1]$, then

$$\lambda_{\max}(A^q B^q) \leq \lambda_{\max}^q(AB).$$

iii) If $q \geq 1$, then

$$\sigma_{\max}^q(AB) \leq \sigma_{\max}(A^q B^q).$$

iv) If $q \geq 1$, then

$$\lambda_{\max}^q(AB) \leq \lambda_{\max}(A^q B^q).$$

v) If $p \geq q > 0$, then

$$\sigma_{\max}^{1/q}(A^q B^q) \leq \sigma_{\max}^{1/p}(A^p B^p).$$

vi) If $p \geq q > 0$, then

$$\lambda_{\max}^{1/q}(A^q B^q) \leq \lambda_{\max}^{1/p}(A^p B^p).$$

(Proof: See [197, pp. 255–258] and [523].) (Remark: See Fact 8.10.49, Fact 8.12.20, Fact 9.9.16, and Fact 9.9.17.) (Remark: ii) is the *Cordes inequality*.)

Fact 8.18.27. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p \geq r \geq 0$. Then,

$$\begin{bmatrix} \lambda_1^{1/p}(A^p B^p) & \cdots & \lambda_n^{1/p}(A^p B^p) \end{bmatrix}$$

strongly log majorizes

$$\begin{bmatrix} \lambda_1^{1/r}(A^r B^r) & \cdots & \lambda_n^{1/r}(A^r B^r) \end{bmatrix}.$$

In fact, for all $q > 0$,

$$\det(A^q B^q)^{1/q} = (\det A) \det B.$$

(Proof: See [197, p. 257] or [1485, p. 20] and Fact 2.21.13.)

Fact 8.18.28. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, and assume that

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$$

is positive semidefinite. Then,

$$\begin{aligned} & \max\{\sigma_{\max}(A), \sigma_{\max}(B)\} \\ & \leq \sigma_{\max}(\mathcal{A}) \\ & \leq \frac{1}{2} \left[\sigma_{\max}(A) + \sigma_{\max}(B) + \sqrt{[\sigma_{\max}(A) - \sigma_{\max}(B)]^2 + 4\sigma_{\max}^2(C)} \right] \\ & \leq \sigma_{\max}(A) + \sigma_{\max}(B) \end{aligned}$$

and

$$\max\{\sigma_{\max}(A), \sigma_{\max}(B)\} \leq \sigma_{\max}(\mathcal{A}) \leq \max\{\sigma_{\max}(A), \sigma_{\max}(B)\} + \sigma_{\max}(C).$$

(Proof: See [719].) (Remark: See Fact 9.14.12.)

Fact 8.18.29. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$\begin{bmatrix} \lambda_1(\log A + \log B) & \cdots & \lambda_n(\log A + \log B) \end{bmatrix}$$

strongly log majorizes

$$\begin{bmatrix} \lambda_1(\log A^{1/2} B A^{1/2}) & \cdots & \lambda_n(\log A^{1/2} B A^{1/2}) \end{bmatrix}.$$

Consequently,

$$\log \det AB = \operatorname{tr}(\log A + \log B) = \operatorname{tr} \log A^{1/2} B A^{1/2} = \log \det A^{1/2} B A^{1/2}.$$

(Proof: See [90].)

Fact 8.18.30. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, the following statements hold:

- i) $\sigma_{\max}[\log(I + A)\log(I + B)] \leq \left(\log \left[1 + \sigma_{\max}^{1/2}(AB) \right] \right)^2$.
- ii) $\sigma_{\max}[\log(I + B)\log(I + A)\log(I + B)] \leq \left(\log \left[1 + \sigma_{\max}^{1/3}(BAB) \right] \right)^3$.
- iii) $\det[\log(I + A)\log(I + B)] \leq \det[\log(I + \langle AB \rangle^{1/2})]^2$.

$$iv) \det[\log(I+B)\log(I+A)\log(I+B)] \leq \det(\log[I+(BAB)^{1/3}])^3.$$

(Proof: See [1349].) (Remark: See Fact 11.16.6.)

Fact 8.18.31. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\sigma_{\max}[(I+A)^{-1}AB(I+B)^{-1}] \leq \frac{\sigma_{\max}(AB)}{\left[1 + \sigma_{\max}^{1/2}(AB)\right]^2}.$$

(Proof: See [1349].)

8.19 Facts on Alternative Partial Orderings

Fact 8.19.1. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then, the following statements are equivalent:

$$i) \log B \leq \log A.$$

ii) There exists $r \in (0, \infty)$ such that

$$B^r \leq \left(B^{r/2}A^rB^{r/2}\right)^{1/2}.$$

iii) There exists $r \in (0, \infty)$ such that

$$\left(A^{r/2}B^rA^{r/2}\right)^{1/2} \leq A^r.$$

iv) There exist $p, r \in (0, \infty)$ and a positive integer k such that $(k+1)r = p+r$ and

$$B^r \leq \left(B^{r/2}A^pB^{r/2}\right)^{\frac{1}{k+1}}.$$

v) There exist $p, r \in (0, \infty)$ and a positive integer k such that $(k+1)r = p+r$ and

$$\left(A^{r/2}B^pA^{r/2}\right)^{\frac{1}{k+1}} \leq A^r.$$

vi) For all $p, r \in [0, \infty)$,

$$B^r \leq \left(B^{r/2}A^pB^{r/2}\right)^{1/2}.$$

vii) For all $p, r \in [0, \infty)$,

$$\left(A^{r/2}B^pA^{r/2}\right)^{\frac{r}{r+p}} \leq A^r.$$

viii) For all $p, q, r, t \in \mathbb{R}$ such that $p \geq 0, r \geq 0, t \geq 0$, and $q \in [1, 2]$,

$$\left[A^{r/2}\left(A^{t/2}B^pA^{t/2}\right)^qA^{r/2}\right]^{\frac{r+t}{r+qt+qp}} \leq A^{r+t}.$$

(Remark: $\log B \leq \log A$ is the *chaotic order*. This order is weaker than the Löwner order since $A \leq B$ implies that $\log A \leq \log A$, but not vice versa.) (Proof: See [512, 914, 1471] and [530, pp. 139, 200].) (Remark: Additional conditions are given in [915].)

Fact 8.19.2. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and assume that $B \leq A$. Then, $\log B \leq \log A$. (Proof: Setting $\tau = 0$ and $q = 1$ in *iii*) of Fact 8.10.51 yields *iii*) of Fact 8.19.1.) (Remark: This result is *xviii*) of Proposition 8.6.13.)

Fact 8.19.3. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive definite and B is positive semidefinite, and let $\alpha > 0$. Then, the following statements are equivalent:

- i) $B^\alpha \leq A^\alpha$.
- ii) For all $p, q, r, \tau \in \mathbb{R}$ such that $p \geq \alpha$, $r \geq \tau$, $q \geq 1$, and $\tau \in [0, \alpha]$,

$$\left[A^{r/2} \left(A^{-\tau/2} B^p A^{-\tau/2} \right)^q A^{r/2} \right]^{\frac{r-\tau}{r-q\tau+qp}} \leq A^{r-\tau}.$$

(Proof: See [512].)

Fact 8.19.4. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite and B is positive semidefinite. Then, the following statements are equivalent:

- i) For all $k \in \mathbb{N}$, $B^k \leq A^k$.
- ii) For all $\alpha > 0$, $B^\alpha \leq A^\alpha$.
- iii) For all $p, r \in \mathbb{R}$ such that $p > r \geq 0$,

$$\left(A^{-r/2} B^p A^{-r/2} \right)^{\frac{2p-r}{p-r}} \leq A^{2p-r}.$$

- iv) For all $p, q, r, \tau \in \mathbb{R}$ such that $p \geq \tau$, $r \geq \tau$, $q \geq 1$, and $\tau \geq 0$,

$$\left[A^{r/2} \left(A^{-\tau/2} B^p A^{-\tau/2} \right)^q A^{r/2} \right]^{\frac{r-\tau}{r-q\tau+qp}} \leq A^{r-\tau}.$$

(Proof: See [531].) (Remark: A and B are related by the *spectral order*.)

Fact 8.19.5. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, if two of the following statements hold, then the remaining statement also holds:

- i) $A \stackrel{\text{rs}}{\leq} B$.
- ii) $A^2 \stackrel{\text{rs}}{\leq} B^2$.
- iii) $AB = BA$.

(Proof: See [110, 590, 591].) (Remark: The rank subtractivity partial ordering is defined in Fact 2.10.32.)

Fact 8.19.6. Let $A, B, C \in \mathbb{F}^{n \times n}$, and assume that A , B , and C are positive semidefinite. Then, the following statements hold:

- i) If $A^2 = AB$ and $B^2 = BA$, then $A = B$.
- ii) If $A^2 = AB$ and $B^2 = BC$, then $A^2 = AC$.

(Proof: Use Fact 2.10.33 and Fact 2.10.34.)

Fact 8.19.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite, and define

$$A \leq^* B$$

if and only if

$$A^2 = AB.$$

Then, “ $\leq^*$ ” is a partial ordering on $\mathbf{N}^{n \times n}$. (Proof: Use Fact 2.10.35 or Fact 8.19.6.) (Remark: The relation “ $\leq^*$ ” is the *star partial ordering*.)

Fact 8.19.8. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$A \leq^* B$$

if and only if

$$B^+ \leq^* A^+.$$

(Proof: See [646].) (Remark: The star partial ordering is defined in Fact 8.19.7.)

Fact 8.19.9. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, the following statements are equivalent:

- i) $A \leq^* B$.
- ii) $A \overset{\text{rs}}{\leq} B$ and $A^2 \overset{\text{rs}}{\leq} B^2$.

(Remark: See [601].) (Remark: The star partial ordering is defined in Fact 8.19.7.)

Fact 8.19.10. Let $A, B \in \mathbb{F}^{n \times m}$, and define

$$A \overset{\text{GL}}{\leq} B$$

if and only if the following conditions hold:

- i) $\langle A \rangle \leq \langle B \rangle$.
- ii) $\mathcal{R}(A^*) \subseteq \mathcal{R}(B^*)$.
- iii) $AB^* = \langle A \rangle \langle B \rangle$.

Then, “ $\overset{\text{GL}}{\leq}$ ” is a partial ordering on $\mathbb{F}^{n \times m}$. Furthermore, the following statements are equivalent:

- iv) $A \overset{\text{GL}}{\leq} B$.
- v) $A^* \overset{\text{GL}}{\leq} B^*$.
- vi) $\text{sprad}(B^+A) \leq 1$, $\mathcal{R}(A) \subseteq \mathcal{R}(B)$, $\mathcal{R}(A^*) \subseteq \mathcal{R}(B^*)$, and $AB^* = \langle A \rangle \langle B \rangle$.

Furthermore, if $A \overset{\text{rs}}{\leq} B$, then $A \overset{\text{GL}}{\leq} B$. Finally, if $A, B \in \mathbf{N}^n$, then $A \leq B$ if and only if $A \overset{\text{GL}}{\leq} B$. (Proof: See [655].) (Remark: The relation “ $\overset{\text{GL}}{\leq}$ ” is the *generalized Löwner partial ordering*. Remarkably, the Löwner, generalized Löwner, and star partial orderings are linked through the polar decomposition. See [655].)

8.20 Facts on Generalized Inverses

Fact 8.20.1. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $A + A^* \geq 0$.
- ii) $A^+ + A^{+*} \geq 0$.

If, in addition, A is group invertible, then the following statement is equivalent to i) and ii):

- iii) $A^\# + A^{\#*} \geq 0$.

(Proof: See [1329].)

Fact 8.20.2. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then, the following statements hold:

- i) $A^+ = A^D = A^\# \geq 0$.
- ii) $\text{rank } A = \text{rank } A^+$.
- iii) $A^{+1/2} \triangleq (A^{1/2})^+ = (A^+)^{1/2}$.
- iv) $A^{1/2} = A(A^+)^{1/2} = (A^+)^{1/2}A$.
- v) $AA^+ = A^{1/2}(A^{1/2})^+$.
- vi) $\begin{bmatrix} A & AA^+ \\ A^+A & A^+ \end{bmatrix}$ is positive semidefinite.
- vii) $A^+A + AA^+ \leq A + A^+$.
- viii) $A^+A \circ AA^+ \leq A \circ A^+$.

(Proof: See [1492] or Fact 8.11.5 and Fact 8.21.40 for vi)–viii).)

Fact 8.20.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then,

$$\text{rank } A \leq (\text{tr } A) \text{tr } A^+.$$

Furthermore, equality holds if and only if $\text{rank } A \leq 1$. (Proof: See [113].)

Fact 8.20.4. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\langle A^* \rangle = A \langle A \rangle^{+1/2} A^*.$$

(Remark: See Fact 8.11.11.)

Fact 8.20.5. Let $A \in \mathbb{F}^{n \times m}$, and define $S \in \mathbb{F}^{n \times n}$ by

$$S \triangleq \langle A \rangle + I_n - AA^+.$$

Then, S is positive definite, and

$$SAA^+S = \langle A \rangle AA^+ \langle A \rangle = AA^*.$$

(Proof: See [447, p. 432].) (Remark: This result provides an explicit congruence transformation for AA^+ and AA^* .) (Remark: See Fact 5.8.20.)

Fact 8.20.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$A = (A + B)(A + B)^+A.$$

Fact 8.20.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, the following statements are equivalent:

$$i) \quad A \stackrel{\text{rs}}{\leq} B.$$

$$ii) \quad \mathcal{R}(A) \subseteq \mathcal{R}(B) \text{ and } AB^+A = A.$$

(Proof: See [590, 591].) (Remark: See Fact 6.5.30.)

Fact 8.20.8. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, assume that $\nu_-(A) = \nu_-(B)$, and consider the following statements:

$$i) \quad A \stackrel{*}{\leq} B.$$

$$ii) \quad A \stackrel{\text{rs}}{\leq} B.$$

$$iii) \quad A \leq B.$$

$$iv) \quad \mathcal{R}(A) \subseteq \mathcal{R}(B) \text{ and } AB^+A \leq A.$$

Then, $i) \implies ii) \implies iii) \iff iv)$. If, in addition, A and B are positive semidefinite, then the following statement is equivalent to $iii)$ and $iv)$:

$$v) \quad \mathcal{R}(A) \subseteq \mathcal{R}(B) \text{ and } \text{sprad}(B^+A) \leq 1.$$

(Proof: $i) \implies ii)$ is given in [652]. See [110, 590, 601, 1223] and [1184, p. 229].)
(Remark: See Fact 8.20.7.)

Fact 8.20.9. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, the following statements are equivalent:

$$i) \quad A^2 \leq B^2.$$

$$ii) \quad \mathcal{R}(A) \subseteq \mathcal{R}(B) \text{ and } \sigma_{\max}(B^+A) \leq 1.$$

(Proof: See [601].)

Fact 8.20.10. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that $A \leq B$. Then, the following statements are equivalent:

$$i) \quad B^+ \leq A^+.$$

$$ii) \quad \text{rank } A = \text{rank } B.$$

$$iii) \quad \mathcal{R}(A) = \mathcal{R}(B).$$

Furthermore, the following statements are equivalent:

$$iv) \quad A^+ \leq B^+.$$

$$v) \quad A^2 = AB.$$

$$vi) \quad A^+ \stackrel{*}{\leq} B^+.$$

(Proof: See [646, 1003].)

Fact 8.20.11. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, if two of the following statements hold, then the remaining statement also holds:

$$i) \ A \leq B.$$

$$ii) \ B^+ \leq A^+.$$

$$iii) \ \text{rank } A = \text{rank } B.$$

(Proof: See [111, 1003, 1422, 1456].)

Fact 8.20.12. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, if two of the following statements hold, then the remaining statement also holds:

$$i) \ A \leq B.$$

$$ii) \ B^+ \leq A^+.$$

$$iii) \ \text{In } A = \text{In } B.$$

(Proof: See [109].)

Fact 8.20.13. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that $A \leq B$. Then,

$$0 \leq AA^+ \leq BB^+.$$

If, in addition, $\text{rank } A = \text{rank } B$, then

$$AA^+ = BB^+.$$

Fact 8.20.14. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that $\mathcal{R}(A) = \mathcal{R}(B)$. Then,

$$\text{In } A - \text{In } B = \text{In}(A - B) + \text{In}(A^+ - B^+).$$

(Proof: See [1047].) (Remark: See Fact 8.10.15.)

Fact 8.20.15. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that $A \leq B$. Then,

$$0 \leq AB^+A \leq A \leq A + B[(I - AA^+)B(I - AA^+)]^+B \leq B.$$

(Proof: See [646].)

Fact 8.20.16. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\text{spec}[(A + B)^+A] \subset [0, 1].$$

(Proof: Let C be positive definite and satisfy $B \leq C$. Then,

$$(A + C)^{-1/2}C(A + C)^{-1/2} \leq I.$$

The result now follows from Fact 8.20.17.)

Fact 8.20.17. Let $A, B, C \in \mathbb{F}^{n \times n}$, assume that A, B, C are positive semidefinite, and assume that $B \leq C$. Then, for all $i = 1, \dots, n$,

$$\lambda_i[(A+B)^+B] \leq \lambda_i[(A+C)^+C].$$

Consequently,

$$\text{tr}[(A+B)^+B] \leq \text{tr}[(A+C)^+C].$$

(Proof: See [1390].) (Remark: See Fact 8.20.16.)

Fact 8.20.18. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and define

$$A:B \triangleq A(A+B)^+B.$$

Then, the following statements hold:

- i) $A:B$ is positive semidefinite.
- ii) $A:B = \lim_{\varepsilon \downarrow 0} (A + \varepsilon I):(B + \varepsilon I)$.
- iii) $A:A = \frac{1}{2}A$.
- iv) $A:B = B:A = B - B(A+B)^+B = A - A(A+B)^+A$.
- v) $A:B \leq A$.
- vi) $A:B \leq B$.

$$vii) \quad A:B = - \begin{bmatrix} 0 & 0 & I \end{bmatrix} \begin{bmatrix} A & 0 & I \\ 0 & B & I \\ I & I & 0 \end{bmatrix}^+ \begin{bmatrix} 0 \\ 0 \\ I \end{bmatrix}.$$

viii) $A:B = (A^+ + B^+)^+$ if and only if $\mathcal{R}(A) = \mathcal{R}(B)$.

ix) $A(A+B)^+B = ACB$ for every (1)-inverse C of $A+B$.

x) $\text{tr}(A:B) \leq (\text{tr } B):(\text{tr } A)$.

xi) $\text{tr}(A:B) = (\text{tr } B):(\text{tr } A)$ if and only if there exists $\alpha \in [0, \infty)$ such that either $A = \alpha B$ or $B = \alpha A$.

xii) $\det(A:B) \leq (\det B):(\det A)$.

xiii) $\mathcal{R}(A:B) = \mathcal{R}(A) \cap \mathcal{R}(B)$.

xiv) $\mathcal{N}(A:B) = \mathcal{N}(A) + \mathcal{N}(B)$.

xv) $\text{rank}(A:B) = \text{rank } A + \text{rank } B - \text{rank}(A+B)$.

xvi) Let $S \in \mathbb{F}^{p \times n}$, and assume that S is right invertible. Then,

$$S(A:B)S^* \leq (SAS^*):(SBS^*).$$

xvii) Let $S \in \mathbb{F}^{n \times n}$, and assume that S is nonsingular. Then,

$$S(A:B)S^* = (SAS^*):(SBS^*).$$

xviii) For all positive numbers α, β ,

$$(\alpha^{-1}A):(\beta^{-1}B) \leq \alpha A + \beta B.$$

xi) Let $X \in \mathbb{F}^{n \times n}$, and assume that X is Hermitian and

$$\begin{bmatrix} A+B & A \\ A & A-X \end{bmatrix} \geq 0.$$

Then,

$$X \leq A:B.$$

Furthermore,

$$\begin{bmatrix} A+B & A \\ A & A-A:B \end{bmatrix} \geq 0.$$

xx) $\phi: \mathbf{N}^n \times \mathbf{N}^n \mapsto -\mathbf{N}^n$ defined by $\phi(A, B) \triangleq -A:B$ is convex.

xxi) If A and B are projectors, then $2(A:B)$ is the projector onto $\mathcal{R}(A) \cap \mathcal{R}(B)$.

xxii) If $A+B$ is positive definite, then

$$A:B = A(A+B)^{-1}B.$$

xxiii) $A\#B = [\frac{1}{2}(A+B)]\#[2(A:B)]$.

xxiv) If $C, D \in \mathbb{F}^{n \times n}$ are positive semidefinite, then

$$(A:B):C = A:(B:C)$$

and

$$A:C + B:D \leq (A+B):(C+D).$$

xxv) If $C, D \in \mathbb{F}^{n \times n}$ are positive semidefinite, $A \leq C$, and $B \leq D$, then

$$A:B \leq C:D.$$

xxvi) If A and B are positive definite, then

$$A:B = (A^{-1} + B^{-1})^{-1} \leq \frac{1}{2}(A\#B) \leq \frac{1}{4}(A+B).$$

xxvii) Let $x, y \in \mathbb{F}^n$. Then,

$$(x+y)^*(A:B)(x+y) \leq x^*Ax + y^*By.$$

xxviii) Let $x, y \in \mathbb{F}^n$. Then,

$$x^*(A:B)x \leq y^*Ay + (x-y)^*B(x-y).$$

xxix) Let $x \in \mathbb{F}^n$. Then,

$$x^*(A:B)x = \inf_{y \in \mathbb{F}^n} [y^*Ay + (x-y)^*B(x-y)].$$

xxx) Let $x \in \mathbb{F}^n$. Then,

$$x^*(A:B)x \leq (x^*Ax) : (x^*Bx).$$

(Proof: See [36, 37, 40, 583, 843, 1284], [1118, p. 189], and [1485, p. 9].) (Remark: $A:B$ is the *parallel sum* of A and B .) (Remark: See Fact 6.4.41 and Fact 6.4.42.) (Remark: A symmetric expression for the parallel sum of three or more positive-semidefinite matrices is given in [1284].)

Fact 8.20.19. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and assume that B is a projector. Then,

$$\text{sh}(A, B) \triangleq \min\{X \in \mathbb{N}^n : 0 \leq X \leq A \text{ and } \mathcal{R}(X) \subseteq \mathcal{R}(B)\}$$

exists. Furthermore,

$$\text{sh}(A, B) = A - AB_{\perp}(B_{\perp}AB_{\perp})^+B_{\perp}A.$$

That is,

$$\text{sh}(A, B) = A \begin{bmatrix} A & AB_{\perp} \\ B_{\perp}A & B_{\perp}AB_{\perp} \end{bmatrix}.$$

Finally,

$$\text{sh}(A, B) = \lim_{\alpha \rightarrow \infty} (\alpha B) : A.$$

(Proof: Existence of the minimum is proved in [40]. The expression for $\text{sh}(A, B)$ is given in [568]; a related expression involving the Schur complement is given in [36]. The last identity is shown in [40]. See also [50].) (Remark: $\text{sh}(A, B)$ is the *shorted operator*.)

Fact 8.20.20. Let $B \in \mathbb{R}^{m \times n}$, define

$$\mathcal{S} \triangleq \{A \in \mathbb{R}^{n \times n} : A \geq 0 \text{ and } \mathcal{R}(B^TBA) \subseteq \mathcal{R}(A)\},$$

and define $\phi : \mathcal{S} \mapsto -\mathbb{N}^m$ by $\phi(A) \triangleq -(BA^+B^T)^+$. Then, $\mathcal{S}$ is a convex cone, and ϕ is convex. (Proof: See [592].) (Remark: This result generalizes *xii*) of Proposition 8.6.17 in the case $r = p = 1$.)

Fact 8.20.21. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. If $(AB)^+ = B^+A^+$, then AB is range Hermitian. Furthermore, the following statements are equivalent:

- i) AB is range Hermitian.
- ii) $(AB)^{\#} = B^+A^+$.
- iii) $(AB)^+ = B^+A^+$.

(Proof: See [988].) (Remark: See Fact 6.4.28.)

Fact 8.20.22. Let $A \in \mathbb{F}^{n \times n}$ and $C \in \mathbb{F}^{m \times m}$, assume that A and C are positive semidefinite, let $B \in \mathbb{F}^{n \times m}$, and define $X \triangleq A^{+1/2}BC^{+1/2}$. Then, the following statements are equivalent:

- i) $\begin{bmatrix} A & B \\ B^* & C \end{bmatrix}$ is positive semidefinite.
- ii) $AA^+B = B$ and $X^*X \leq I_m$.
- iii) $BC^+C = B$ and $X^*X \leq I_m$.
- iv) $B = A^{1/2}XC^{1/2}$ and $X^*X \leq I_m$.

- v) There exists a matrix $Y \in \mathbb{F}^{n \times m}$ such that $B = A^{1/2}YC^{1/2}$ and $Y^*Y \leq I_m$.

(Proof: See [1485, p. 15].)

Fact 8.20.23. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, the following statements are equivalent:

- i) $A(A + B)^+B = 0$.
- ii) $B(A + B)^+A = 0$.
- iii) $A(A + B)^+A = A$.
- iv) $B(A + B)^+B = B$.
- v) $A(A + B)^+B + B(A + B)^+A = 0$.
- vi) $A(A + B)^+A + B(A + B)^+B = A + B$.
- vii) $\text{rank} \begin{bmatrix} A & B \end{bmatrix} = \text{rank } A + \text{rank } B$.
- viii) $\mathcal{R}(A) \cap \mathcal{R}(B) = \{0\}$.
- ix) $(A + B)^+ = [(I - BB^+)A(I - B^+B)]^+ + [(I - AA^+)B(I - A^+A)]^+.$

(Proof: See [1302].) (Remark: See Fact 6.4.32.)

8.21 Facts on the Kronecker and Schur Products

Fact 8.21.1. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and assume that every entry of A is nonzero. Then, $A^{\circ-1}$ is positive semidefinite if and only if $\text{rank } A = 1$. (Proof: See [889].)

Fact 8.21.2. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, assume that every entry of A is nonnegative, and let $\alpha \in [0, n - 2]$. Then, $A^{\circ\alpha}$ is positive semidefinite. (Proof: See [199, 491].) (Remark: In many cases, $A^{\circ\alpha}$ is positive semidefinite for all $\alpha \geq 0$. See Fact 8.8.5.)

Fact 8.21.3. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $k \geq 1$. If $r \in [0, 1]$, then

$$(A^r)^{\circ k} \leq (A^{\circ k})^r.$$

If $r \in [1, 2]$, then

$$(A^{\circ k})^r \leq (A^r)^{\circ k}.$$

If A is positive definite and $r \in [0, 1]$, then

$$(A^{\circ k})^{-r} \leq (A^{-r})^{\circ k}.$$

(Proof: See [1485, p. 8].)

Fact 8.21.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then,

$$(I \circ A)^2 \leq \frac{1}{2}(I \circ A^2 + A \circ A) \leq I \circ A^2$$

and

$$A \circ A \leq I \circ A^2.$$

Hence,

$$\sum_{i=1}^n A_{(i,i)}^2 \leq \sum_{i=1}^n \lambda_i^2(A).$$

Now, assume that A is positive definite. Then,

$$(A \circ A)^{-1} \leq A^{-1} \circ A^{-1}$$

and

$$(A \circ A^{-1})^{-1} \leq I \leq \left(A^{1/2} \circ A^{-1/2} \right)^2 \leq \frac{1}{2}(I + A \circ A^{-1}) \leq A \circ A^{-1}.$$

Furthermore,

$$(A \circ A^{-1})1_{n \times 1} = 1_{n \times 1}$$

and

$$1 \in \text{spec}(A \circ A^{-1}).$$

Next, let $\alpha \triangleq \lambda_{\min}(A)$ and $\beta \triangleq \lambda_{\max}(A)$. Then,

$$\frac{2\alpha\beta}{\alpha^2 + \beta^2} I \leq \frac{2\alpha\beta}{\alpha^2 + \beta^2} (A^2 \circ A^{-2})^{1/2} \leq \frac{\alpha\beta}{\alpha^2 + \beta^2} (I + A^2 \circ A^{-2}) \leq A \circ A^{-1}.$$

Define $\Phi(A) \triangleq A \circ A^{-1}$, and, for all $k \geq 1$, define

$$\Phi^{(k+1)}(A) \triangleq \Phi[\Phi^{(k)}(A)],$$

where $\Phi^{(1)}(A) \triangleq \Phi(A)$. Then, for all $k \geq 1$,

$$\Phi^{(k)}(A) \geq I$$

and

$$\lim_{k \rightarrow \infty} \Phi^{(k)}(A) = I.$$

(Proof: See [480, 772, 1383, 1384], [709, p. 475], and set $B = A^{-1}$ in Fact 8.21.31.)

(Remark: The convergence result also holds if A is an H -matrix [772]. $A \circ A^{-1}$ is the *relative gain array*.) (Remark: See Fact 8.21.38.)

Fact 8.21.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then, for all $i = 1, \dots, n$,

$$1 \leq A_{(i,i)}(A^{-1})_{(i,i)}.$$

Furthermore,

$$\max_{i=1, \dots, n} \sqrt{A_{(i,i)}(A^{-1})_{(i,i)} - 1} \leq \sum_{i=1}^n \sqrt{A_{(i,i)}(A^{-1})_{(i,i)} - 1}$$

and

$$\max_{i=1, \dots, n} \sqrt{A_{(i,i)}(A^{-1})_{(i,i)} - 1} \leq \sum_{i=1}^n \left[\sqrt{A_{(i,i)}(A^{-1})_{(i,i)} - 1} \right].$$

(Proof: See [482, p. 66-6].)

Fact 8.21.6. Let $\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, assume that $\mathcal{A}$ is positive definite, and partition $\mathcal{A}^{-1} = \begin{bmatrix} X & Y \\ Y^* & Z \end{bmatrix}$ conformably with $\mathcal{A}$. Then,

$$I \leq \begin{bmatrix} A \circ A^{-1} & 0 \\ 0 & Z \circ Z^{-1} \end{bmatrix} \leq \mathcal{A} \circ \mathcal{A}^{-1}$$

and

$$I \leq \begin{bmatrix} X \circ X^{-1} & 0 \\ 0 & C \circ C^{-1} \end{bmatrix} \leq \mathcal{A} \circ \mathcal{A}^{-1}.$$

(Proof: See [132].)

Fact 8.21.7. Let $A \in \mathbb{F}^{n \times n}$, let $p, q \in \mathbb{R}$, assume that A is positive semidefinite, and assume that either p and q are nonnegative or A is positive definite. Then,

$$A^{(p+q)/2} \circ A^{(p+q)/2} \leq A^p \circ A^q.$$

In particular,

$$I \leq A \circ A^{-1}.$$

(Proof: See [92].)

Fact 8.21.8. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and assume that $I_n \circ A = I_n$. Then,

$$\det A \leq \lambda_{\min}(A \circ \overline{A}).$$

(Proof: See [1408].)

Fact 8.21.9. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$-A^*A \circ I \leq A^* \circ A \leq A^*A \circ I.$$

(Proof: Use Fact 8.21.41 with $B = I$.)

Fact 8.21.10. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\langle A \circ A^* \rangle \leq \left\{ \begin{array}{c} A^*A \circ I \\ \langle A \rangle \circ \langle A^* \rangle \end{array} \right\} \leq \sigma_{\max}^2(A)I.$$

(Proof: See [1492] and Fact 8.21.22.)

Fact 8.21.11. Let $A \triangleq \begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ and $B \triangleq \begin{bmatrix} B_{11} & B_{12} \\ B_{12}^* & B_{22} \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$, and assume that A and B are positive semidefinite. Then,

$$(A_{11}|A) \circ (B_{11}|B) \leq (A_{11}|A) \circ B_{22} \leq (A_{11} \circ B_{11})|(A \circ B).$$

(Proof: See [896].)

Fact 8.21.12. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, $A \circ B$ is positive semidefinite. If, in addition, B is positive definite and $I \circ A$ is positive definite, then $A \circ B$ is positive definite. (Proof: By Fact 7.4.16, $A \otimes B$ is positive semidefinite, and the Schur product $A \circ B$ is a principal submatrix of the Kronecker product. If A is positive definite, use Fact 8.21.19 to obtain $\det(A \circ B) > 0$.) (Remark: The first result is *Schur's theorem*. The second result is *Schott's theorem*. See [925] and Fact 8.21.19.)

Fact 8.21.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then, there exist positive-definite matrices $B, C \in \mathbb{F}^{n \times n}$ such that $A = B \circ C$. (Remark: See [1098, pp. 154, 166].) (Remark: This result is due to Djokovic.)

Fact 8.21.14. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite and B is positive semidefinite. Then,

$$(1_{1 \times n} A^{-1} 1_{n \times 1})^{-1} B \leq A \circ B.$$

(Proof: See [484].) (Remark: Setting $B = 1_{n \times n}$ yields Fact 8.9.17.)

Fact 8.21.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$(1_{1 \times n} A^{-1} 1_{n \times 1} 1_{1 \times n} B^{-1} 1_{n \times 1})^{-1} 1_{n \times n} \leq A \circ B.$$

(Proof: See [1492].)

Fact 8.21.16. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive definite, let $B \in \mathbb{F}^{n \times n}$, and assume that B is positive semidefinite. Then,

$$\text{rank } B \leq \text{rank}(A \circ B) \leq \text{rank}(A \otimes B) = (\text{rank } A)(\text{rank } B).$$

(Remark: See Fact 7.4.23, Fact 7.6.6, and Fact 8.21.14.) (Remark: The first inequality is due to Djokovic. See [1098, pp. 154, 166].)

Fact 8.21.17. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. If $p \geq 1$, then

$$\text{tr } (A \circ B)^p \leq \text{tr } A^p \circ B^p.$$

If $0 \leq p \leq 1$, then

$$\text{tr } A^p \circ B^p \leq \text{tr } (A \circ B)^p.$$

Now, assume that A and B are positive definite. If $p \leq 0$, then

$$\text{tr } (A \circ B)^p \leq \text{tr } A^p \circ B^p.$$

(Proof: See [1392].)

Fact 8.21.18. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\lambda_{\min}(AB) \leq \lambda_{\min}(A \circ B).$$

Hence,

$$\lambda_{\min}(AB)I \leq \lambda_{\min}(A \circ B)I \leq A \circ B.$$

(Proof: See [765].) (Remark: This result interpolates the penultimate inequality in Fact 8.21.20.)

Fact 8.21.19. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\det AB \leq \left(\prod_{i=1}^n A_{(i,i)} \right) \det B \leq \det(A \circ B) \leq \prod_{i=1}^n A_{(i,i)} B_{(i,i)}.$$

Equivalently,

$$\det AB \leq [\det(I \circ A)] \det B \leq \det(A \circ B) \leq \prod_{i=1}^n A_{(i,i)} B_{(i,i)}.$$

Furthermore,

$$2 \det AB \leq \left(\prod_{i=1}^n A_{(i,i)} \right) \det B + \left(\prod_{i=1}^n B_{(i,i)} \right) \det A \leq \det(A \circ B) + (\det A) \det B.$$

Finally, the following statements hold:

- i) If $I \circ A$ and B are positive definite, then $A \circ B$ is positive definite.
- ii) If $I \circ A$ and B are positive definite and $\text{rank } A = 1$, then equality holds in the right-hand equality.
- iii) If A and B are positive definite, then equality holds in the right-hand equality if and only if B is diagonal.

(Proof: See [967, 1477] and [1184, p. 253].) (Remark: In the first string, the first and third inequalities follow from Hadamard's inequality Fact 8.17.11, while the second inequality is *Oppenheim's inequality*. See Fact 8.21.12.) (Remark: The right-hand inequality in the third string of inequalities is valid when A and B are M-matrices. See [44, 318].) (Problem: Compare the lower bounds $\det(A \# B)^2$ and $(\prod_{i=1}^n A_{(i,i)}) \det B$ for $\det(A \circ B)$. See Fact 8.21.20.)

Fact 8.21.20. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let $k \in \{1, \dots, n\}$, and let $r \in (0, 1]$. Then,

$$\prod_{i=k}^n \lambda_i(A) \lambda_i(B) \leq \prod_{i=k}^n \sigma_i(AB) \leq \prod_{i=k}^n \lambda_i(AB) \leq \prod_{i=k}^n \lambda_i^2(A \# B) \leq \prod_{i=k}^n \lambda_i(A \circ B)$$

and

$$\begin{aligned} \prod_{i=k}^n \lambda_i(A) \lambda_i(B) &\leq \prod_{i=k}^n \sigma_i(AB) \leq \prod_{i=k}^n \lambda_i(AB) \leq \prod_{i=k}^n \lambda_i^{1/r}(A^r B^r) \\ &\leq \prod_{i=k}^n e^{\lambda_i(\log A + \log B)} \leq \prod_{i=k}^n e^{\lambda_i[I \circ (\log A + \log B)]} \\ &\leq \prod_{i=k}^n \lambda_i^{1/r}(A^r \circ B^r) \leq \prod_{i=k}^n \lambda_i(A \circ B). \end{aligned}$$

Consequently,

$$\lambda_{\min}(AB)I \leq A \circ B$$

and

$$\det AB = \det(A \# B)^2 \leq \det(A \circ B).$$

(Proof: See [48, 480, 1382], [1485, p. 21], Fact 8.10.43, and Fact 8.18.21.)

Fact 8.21.21. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, let $k \in \{1, \dots, n\}$, and let $r > 0$. Then,

$$\prod_{i=k}^n \lambda_i^{-r}(A \circ B) \leq \prod_{i=k}^n \lambda_i^{-r}(AB).$$

(Proof: See [1381].)

Fact 8.21.22. Let $A, B \in \mathbb{F}^{n \times n}$, let $C, D \in \mathbb{F}^{m \times m}$, assume that A, B, C , and D are Hermitian, $A \leq B$, $C \leq D$, and that either A and C are positive semidefinite, A and D are positive semidefinite, or B and D are positive semidefinite. Then,

$$A \otimes C \leq B \otimes D.$$

If, in addition, $n = m$, then

$$A \circ C \leq B \circ D.$$

(Proof: See [43, 111].) (Problem: Under which conditions are these inequalities strict?)

Fact 8.21.23. Let $A, B, C, D \in \mathbb{F}^{n \times n}$, assume that A, B, C, D are positive semidefinite, and assume that $A \leq B$ and $C \leq D$. Then,

$$0 \leq A \otimes C \leq B \otimes D$$

and

$$0 \leq A \circ C \leq B \circ D.$$

(Proof: See Fact 8.21.22.)

Fact 8.21.24. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, $A \leq B$ if and only if $A \otimes A \leq B \otimes B$. (Proof: See [925].)

Fact 8.21.25. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, assume that $0 \leq A \leq B$, and let $k \geq 1$. Then,

$$A^{\circ k} \leq B^{\circ k}.$$

(Proof: $0 \leq (B - A) \circ (B + A)$ implies that $A \circ A \leq B \circ B$, that is, $A^{\circ 2} \leq B^{\circ 2}$.)

Fact 8.21.26. Let $A_1, \dots, A_k, B_1, \dots, B_k \in \mathbb{F}^{n \times n}$, and assume that $A_1, \dots, A_k, B_1, \dots, B_k$ are positive semidefinite. Then,

$$(A_1 + B_1) \otimes \cdots \otimes (A_k + B_k) \leq A_1 \otimes \cdots \otimes A_k + B_1 \otimes \cdots \otimes B_k.$$

(Proof: See [994, p. 143].)

Fact 8.21.27. Let $A_1, A_2, B_1, B_2 \in \mathbb{F}^{n \times n}$, assume that A_1, A_2, B_1, B_2 are positive semidefinite, assume that $0 \leq A_1 \leq B_1$ and $0 \leq A_2 \leq B_2$, and let $\alpha \in [0, 1]$. Then,

$$[\alpha A_1 + (1 - \alpha) B_1] \otimes [\alpha A_2 + (1 - \alpha) B_2] \leq \alpha (A_1 \otimes A_2) + (1 - \alpha) (B_1 \otimes B_2).$$

(Proof: See [1406].)

Fact 8.21.28. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, for all $i = 1, \dots, n$,

$$\lambda_n(A) \lambda_n(B) \leq \lambda_{i+n^2-n}(A \otimes B) \leq \lambda_i(A \circ B) \leq \lambda_i(A \otimes B) \leq \lambda_1(A) \lambda_1(B).$$

(Proof: The result follows from Proposition 7.3.1 and Theorem 8.4.5. For A, B positive semidefinite, the result is given in [962].)

Fact 8.21.29. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$, assume that A and B are positive semidefinite, let $r \in \mathbb{R}$, and assume that either A and B are positive

definite or r is positive. Then,

$$(A \otimes B)^r = A^r \otimes B^r.$$

(Proof: See [1019].)

Fact 8.21.30. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{k \times l}$. Then,

$$\langle A \otimes B \rangle = \langle A \rangle \otimes \langle B \rangle.$$

Fact 8.21.31. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. If $r \in [0, 1]$, then

$$A^r \circ B^r \leq (A \circ B)^r.$$

If $r \in [1, 2]$, then

$$(A \circ B)^r \leq A^r \circ B^r.$$

If A and B are positive definite and $r \in [0, 1]$, then

$$(A \circ B)^{-r} \leq A^{-r} \circ B^{-r}.$$

Therefore,

$$\begin{aligned} (A \circ B)^2 &\leq A^2 \circ B^2, \\ A \circ B &\leq (A^2 \circ B^2)^{1/2}, \\ A^{1/2} \circ B^{1/2} &\leq (A \circ B)^{1/2}. \end{aligned}$$

Furthermore,

$$A^2 \circ B^2 - \frac{1}{4}(\beta - \alpha)^2 I \leq (A \circ B)^2 \leq \frac{1}{2}[A^2 \circ B^2 + (AB)^{\circ 2}] \leq A^2 \circ B^2$$

and

$$A \circ B \leq (A^2 \circ B^2)^{1/2} \leq \frac{\alpha + \beta}{2\sqrt{\alpha\beta}} A \circ B,$$

where $\alpha \triangleq \lambda_{\min}(A \otimes B)$ and $\beta \triangleq \lambda_{\max}(A \otimes B)$. Hence,

$$\begin{aligned} A \circ B - \frac{1}{4}(\sqrt{\beta} - \sqrt{\alpha})^2 I &\leq (A^{1/2} \circ B^{1/2})^2 \\ &\leq \frac{1}{2} \left[A \circ B + (A^{1/2} B^{1/2})^{\circ 2} \right] \\ &\leq A \circ B \\ &\leq (A^2 \circ B^2)^{1/2} \\ &\leq \frac{\alpha + \beta}{2\sqrt{\alpha\beta}} A \circ B. \end{aligned}$$

(Proof: See [43, 1018, 1383], [709, p. 475], and [1485, p. 8].)

Fact 8.21.32. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, there exist unitary matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that

$$\langle A \circ B \rangle \leq \frac{1}{2}[S_1(\langle A \rangle \circ \langle B \rangle)S_1^* + S_2(\langle A \rangle \circ \langle B \rangle)S_2^*].$$

(Proof: See [90].)

Fact 8.21.33. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let k, l be nonzero integers such that $k \leq l$. Then,

$$(A^k \circ B^k)^{1/k} \leq (A^l \circ B^l)^{1/l}.$$

In particular,

$$(A^{-1} \circ B^{-1})^{-1} \leq A \circ B$$

and

$$(A \circ B)^{-1} \leq A^{-1} \circ B^{-1},$$

and, for all $k \geq 1$,

$$A \circ B \leq (A^k \circ B^k)^{1/k},$$

and

$$A^{1/k} \circ B^{1/k} \leq (A \circ B)^{1/k}.$$

Furthermore,

$$(A \circ B)^{-1} \leq A^{-1} \circ B^{-1} \leq \frac{(\alpha + \beta)^2}{4\alpha\beta} (A \circ B)^{-1},$$

where $\alpha \triangleq \lambda_{\min}(A \otimes B)$ and $\beta \triangleq \lambda_{\max}(A \otimes B)$. (Proof: See [1018].)

Fact 8.21.34. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite, B is positive semidefinite, and $I \circ B$ is positive definite. Then, for all $i = 1, \dots, n$,

$$[(A \circ B)^{-1}]_{(i,i)} \leq \frac{(A^{-1})_{(i,i)}}{B_{(i,i)}}.$$

Furthermore, if $\text{rank } B = 1$, then equality holds. (Proof: See [1477].)

Fact 8.21.35. Let $A, B \in \mathbb{F}^{n \times n}$. Then, A is positive semidefinite if and only if, for every positive-semidefinite matrix $B \in \mathbb{F}^{n \times n}$,

$$1_{1 \times n}(A \circ B)1_{n \times 1} \geq 0.$$

(Proof: See [709, p. 459].) (Remark: This result is *Fejer's theorem*.)

Fact 8.21.36. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$1_{1 \times n}[(A - B) \circ (A^{-1} - B^{-1})]1_{n \times 1} \leq 0.$$

Furthermore, equality holds if and only if $A = B$. (Proof: See [148, p. 8-8].)

Fact 8.21.37. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let $p, q \in \mathbb{R}$, and assume that one of the following conditions is satisfied:

- i) $p \leq q \leq -1$, and A and B are positive definite.
- ii) $p \leq -1 < 1 \leq q$, and A and B are positive definite.
- iii) $1 \leq p \leq q$.
- iv) $\frac{1}{2} \leq p \leq 1 \leq q$.
- v) $p \leq -1 \leq q \leq -\frac{1}{2}$, and A and B are positive definite.

Then,

$$(A^p \circ B^p)^{1/p} \leq (A^q \circ B^q)^{1/q}.$$

(Proof: See [1019]. Consider case *iii*). Since $p/q \leq 1$, it follows from Fact 8.21.31 that $A^p \circ B^p = (A^q)^{p/q} \circ (A^q)^{p/q} \leq (A^q \circ B^q)^{p/q}$. Then, use Corollary 8.6.11 with p replaced by $1/p$. See [1485, p. 8].) (Remark: See [92].)

Fact 8.21.38. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$2I \leq A \circ B^{-1} + B \circ A^{-1}.$$

(Proof: See [1383, 1492].) (Remark: Setting $B = A$ yields an inequality given by Fact 8.21.4.)

Fact 8.21.39. Let $A, B \in \mathbb{F}^{n \times m}$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} A^*A \circ B^*B & (A \circ B)^* \\ A \circ B & I \end{bmatrix}.$$

Then, $\mathcal{A}$ is positive semidefinite. Furthermore,

$$(A \circ B)^*(A \circ B) \leq \frac{1}{2}(A^*A \circ B^*B + A^*B \circ B^*A) \leq A^*A \circ B^*B.$$

(Proof: See [713, 1383, 1492].) (Remark: The inequality $(A \circ B)^*(A \circ B) \leq A^*A \circ B^*B$ is *Amemiya's inequality*. See [925].)

Fact 8.21.40. Let $A, B, C \in \mathbb{F}^{n \times n}$, define

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix},$$

and assume that $\mathcal{A}$ is positive semidefinite. Then,

$$-A \circ C \leq B \circ B^* \leq A \circ C$$

and

$$|\det(B \circ B^*)| \leq \det(A \circ C).$$

If, in addition, $\mathcal{A}$ is positive definite, then

$$-A \circ C < B \circ B^* < A \circ C$$

and

$$|\det(B \circ B^*)| < \det(A \circ C).$$

(Proof: See [1492].) (Remark: See Fact 8.11.5.)

Fact 8.21.41. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$-A^*A \circ B^*B \leq A^*B \circ B^*A \leq A^*A \circ B^*B$$

and

$$|\det(A^*B \circ B^*A)| \leq \det(A^*A \circ B^*B).$$

(Proof: Apply Fact 8.21.40 to $\begin{bmatrix} A^*A & A^*B \\ B^*A & B^*B \end{bmatrix}$.) (Remark: See Fact 8.11.14 and Fact 8.21.9.)

Fact 8.21.42. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$-A \circ B^*A^{-1}B \leq B \circ B^* \leq A \circ B^*A^{-1}B$$

and

$$|\det(B \circ B^*)| \leq \det(A \circ B^* A^{-1} B).$$

(Proof: Use Fact 8.11.19 and Fact 8.21.40.)

Fact 8.21.43. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\alpha, \beta \in (0, \infty)$.

$$\begin{aligned} -\left(\beta^{-1/2}I + \alpha A^*A\right) \circ \left(\alpha^{-1/2}I + \beta BB^*\right) &\leq (A + B) \circ (A + B)^* \\ &\leq \left(\beta^{-1/2}I + \alpha A^*A\right) \circ \left(\alpha^{-1/2}I + \beta BB^*\right). \end{aligned}$$

(Remark: See Fact 8.11.20.)

Fact 8.21.44. Let $A, B \in \mathbb{F}^{n \times m}$, and define

$$\mathcal{A} \triangleq \begin{bmatrix} A^*A \circ I & (A \circ B)^* \\ A \circ B & BB^* \circ I \end{bmatrix}.$$

Then, $\mathcal{A}$ is positive semidefinite. Now, assume that $n = m$. Then,

$$-A^*A \circ I - BB^* \circ I \leq A \circ B + (A \circ B)^* \leq A^*A \circ I + BB^* \circ I$$

and

$$-A^*A \circ BB^* \circ I \leq A \circ A^* \circ B \circ B^* \leq A^*A \circ BB^* \circ I.$$

(Remark: See Fact 8.21.40.)

Fact 8.21.45. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$A \circ B \leq \frac{1}{2}(A^2 + B^2) \circ I.$$

(Proof: Use Fact 8.21.44.)

Fact 8.21.46. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and define $e^{\circ A} \in \mathbb{F}^{n \times n}$ by $[e^{\circ A}]_{(i,j)} \triangleq e^{A_{(i,j)}}$. Then, $e^{\circ A}$ is positive semidefinite. (Proof: Note that $e^{\circ A} = 1_{n \times n} + \frac{1}{2}A \circ A + \frac{1}{3!}A \circ A \circ A + \cdots$, and use Fact 8.21.12. See [422, p. 10].)

Fact 8.21.47. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $p, q \in (0, \infty)$ satisfy $p \leq q$. Then,

$$I \circ (\log A + \log B) \leq \log(A^p \circ B^p)^{1/p} \leq \log(A^q \circ B^q)^{1/q}$$

and

$$I \circ (\log A + \log B) = \lim_{p \downarrow 0} \log(A^p \circ B^p)^{1/p}.$$

(Proof: See [1382].) (Remark: $\log(A^p \circ B^p)^{1/p} = \frac{1}{p} \log(A^p \circ B^p)$.)

Fact 8.21.48. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$I \circ (\log A + \log B) \leq \log(A \circ B).$$

(Proof: Set $p = 1$ in Fact 8.21.47. See [43] and [1485, p. 8].) (Remark: See Fact 11.14.21.)

Fact 8.21.49. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $C, D \in \mathbb{F}^{m \times n}$. Then,

$$(C \circ D)(A \circ B)^{-1}(C \circ D)^* \leq (CA^{-1}C^*) \circ (DB^{-1}D^*).$$

In particular,

$$(A \circ B)^{-1} \leq A^{-1} \circ B^{-1}$$

and

$$(C \circ D)(C \circ D)^* \leq (CC^*) \circ (DD^*).$$

(Proof: Form the Schur complement of the lower right block of the Schur product of the positive-semidefinite matrices $\begin{bmatrix} A & C^* \\ C & CA^{-1}C^* \end{bmatrix}$ and $\begin{bmatrix} B & D^* \\ D & DB^{-1}D^* \end{bmatrix}$. See [966, 1393], [1485, p. 13], or [1490, p. 198].)

Fact 8.21.50. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p, q \in (1, \infty)$ satisfy $1/p + 1/q = 1$. Then,

$$(A \circ B) + (C \circ D) \leq (A^p + C^p)^{1/p} \circ (B^q + D^q)^{1/q}.$$

(Proof: Use *xxiv*) of Proposition 8.6.17 with $r = 1/p$. See [1485, p. 10].) (Remark: Note the relationship between the *conjugate parameters* p, q and the *barycentric coordinates* $\alpha, 1 - \alpha$. See Fact 1.16.11.)

Fact 8.21.51. Let $A, B, C, D \in \mathbb{F}^{n \times n}$, assume that A, B, C , and D are positive definite. Then,

$$(A \# C) \circ (B \# D) \leq (A \circ B) \# (C \circ D).$$

Furthermore,

$$(A \# B) \circ (A \# B) \leq (A \circ B).$$

(Proof: See [92].)

8.22 Notes

The ordering $A \leq B$ is traditionally called the *Löwner ordering*. Proposition 8.2.4 is given in [14] and [846] with extensions in [167]. The proof of Proposition 8.2.7 is based on [264, p. 120], as suggested in [1249]. The proof given in [540, p. 307] is incomplete.

Theorem 8.3.4 is due to Newcomb [1035]. Proposition 8.4.13 is given in [699, 1022]. Special cases such as Fact 8.12.28 appear in numerous papers. The proofs of Lemma 8.4.4 and Theorem 8.4.5 are based on [1230]. Theorem 8.4.9 can also be obtained as a corollary of the *Fischer minimax theorem* given in [709, 971], which provides a geometric characterization of the eigenvalues of a symmetric matrix. Theorem 8.3.5 appears in [1118, p. 121]. Theorem 8.6.2 is given in [40]. Additional inequalities appear in [1007].

Functions that are nondecreasing on $\mathbf{P}^n$ are characterized by the theory of *monotone matrix functions* [197, 422]. See [1012] for a summary of the principal results.

The literature on convex maps is extensive. Result *xiv*) of Proposition 8.6.17 is due to Lieb and Ruskai [907]. Result *xxiv*) is the *Lieb concavity theorem*. See [197, p. 271] or [905]. Result *xxiv*) is due to Ando. Results *xlvi*) and *xlvi*) are due to Fan. Some extensions to strict convexity are considered in [971]. See also [43, 1024].

Products of positive-definite matrices are studied in [117, 118, 119, 121, 1458].

Essays on the legacy of Issai Schur appear in [780]. Schur complements are discussed in [288, 290, 658, 896, 922, 1057]. Majorization and eigenvalue inequalities for sums and products of matrices are discussed in [198].

Chapter Nine

Norms

Norms are used to quantify vectors and matrices, and they play a basic role in convergence analysis. This chapter introduces vector and matrix norms and their properties.

9.1 Vector Norms

For many applications it is useful to have a scalar measure of the magnitude of a vector x or a matrix A . *Norms* provide such measures.

Definition 9.1.1. A *norm* $\|\cdot\|$ on $\mathbb{F}^n$ is a function $\|\cdot\|: \mathbb{F}^n \mapsto [0, \infty)$ that satisfies the following conditions:

- i) $\|x\| \geq 0$ for all $x \in \mathbb{F}^n$.
- ii) $\|x\| = 0$ if and only if $x = 0$.
- iii) $\|\alpha x\| = |\alpha| \|x\|$ for all $\alpha \in \mathbb{F}$ and $x \in \mathbb{F}^n$.
- iv) $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in \mathbb{F}^n$.

Condition iv) is the *triangle inequality*.

A norm $\|\cdot\|$ on $\mathbb{F}^n$ is *monotone* if $|x| \leq |y|$ implies that $\|x\| \leq \|y\|$ for all $x, y \in \mathbb{F}^n$, while $\|\cdot\|$ is *absolute* if $\||x|\| = \|x\|$ for all $x \in \mathbb{F}^n$.

Proposition 9.1.2. Let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then, $\|\cdot\|$ is monotone if and only if $\|\cdot\|$ is absolute.

Proof. First, suppose that $\|\cdot\|$ is monotone. Let $x \in \mathbb{F}^n$, and define $y \triangleq |x|$. Then, $|y| = |x|$, and thus $|y| \leq |x|$ and $|x| \leq |y|$. Hence, $\|x\| \leq \|y\|$ and $\|y\| \leq \|x\|$, which implies that $\|x\| = \|y\|$. Thus, $\||x|\| = \|y\| = \|x\|$, which proves that $\|\cdot\|$ is absolute.

Conversely, suppose that $\|\cdot\|$ is absolute and, for convenience, let $n = 2$. Now, let $x, y \in \mathbb{F}^2$ be such that $|x| \leq |y|$. Then, there exist $\alpha_1, \alpha_2 \in [0, 1]$ and $\theta_1, \theta_2 \in \mathbb{R}$ such that $x_{(i)} = \alpha_i e^{j\theta_i} y_{(i)}$ for $i = 1, 2$. Since $\|\cdot\|$ is absolute, it follows

that

$$\begin{aligned}
 \|x\| &= \left\| \begin{bmatrix} \alpha_1 e^{j\theta_1} y_{(1)} \\ \alpha_2 e^{j\theta_2} y_{(2)} \end{bmatrix} \right\| \\
 &= \left\| \begin{bmatrix} \alpha_1 |y_{(1)}| \\ \alpha_2 |y_{(2)}| \end{bmatrix} \right\| \\
 &= \left\| \frac{1}{2}(1 - \alpha_1) \begin{bmatrix} -|y_{(1)}| \\ \alpha_2 |y_{(2)}| \end{bmatrix} + \frac{1}{2}(1 - \alpha_1) \begin{bmatrix} |y_{(1)}| \\ \alpha_2 |y_{(2)}| \end{bmatrix} + \alpha_1 \begin{bmatrix} |y_{(1)}| \\ \alpha_2 |y_{(2)}| \end{bmatrix} \right\| \\
 &\leq \left[\frac{1}{2}(1 - \alpha_1) + \frac{1}{2}(1 - \alpha_1) + \alpha_1 \right] \left\| \begin{bmatrix} |y_{(1)}| \\ \alpha_2 |y_{(2)}| \end{bmatrix} \right\| \\
 &= \left\| \begin{bmatrix} |y_{(1)}| \\ \alpha_2 |y_{(2)}| \end{bmatrix} \right\| \\
 &= \left\| \frac{1}{2}(1 - \alpha_2) \begin{bmatrix} |y_{(1)}| \\ -|y_{(2)}| \end{bmatrix} + \frac{1}{2}(1 - \alpha_2) \begin{bmatrix} |y_{(1)}| \\ |y_{(2)}| \end{bmatrix} + \alpha_2 \begin{bmatrix} |y_{(1)}| \\ |y_{(2)}| \end{bmatrix} \right\| \\
 &\leq \left\| \begin{bmatrix} |y_{(1)}| \\ |y_{(2)}| \end{bmatrix} \right\| \\
 &= \|y\|.
 \end{aligned}$$

Thus, $\|\cdot\|$ is monotone. $\square$

As we shall see, there are many different norms. For $x \in \mathbb{F}^n$, a useful class of norms consists of the *Hölder norms* defined by

$$\|x\|_p \triangleq \begin{cases} \left(\sum_{i=1}^n |x_{(i)}|^p \right)^{1/p}, & 1 \leq p < \infty, \\ \max_{i \in \{1, \dots, n\}} |x_{(i)}|, & p = \infty. \end{cases} \quad (9.1.1)$$

Note that, for all $x \in \mathbb{C}^n$ and $p \in [1, \infty]$, $\|\bar{x}\|_p = \|x\|_p$. These norms depend on *Minkowski's inequality* given by the following result.

Lemma 9.1.3. Let $p \in [1, \infty]$, and let $x, y \in \mathbb{F}^n$. Then,

$$\|x + y\|_p \leq \|x\|_p + \|y\|_p. \quad (9.1.2)$$

If $p = 1$, then equality holds if and only if, for all $i = 1, \dots, n$, there exists $\alpha_i \geq 0$ such that either $x_{(i)} = \alpha_i y_{(i)}$ or $y_{(i)} = \alpha_i x_{(i)}$. If $p \in (1, \infty)$, then equality holds if and only if there exists $\alpha \geq 0$ such that either $x = \alpha y$ or $y = \alpha x$.

Proof. See [162, 963] and Fact 1.16.25. $\square$

Proposition 9.1.4. Let $p \in [1, \infty]$. Then, $\|\cdot\|_p$ is a norm on $\mathbb{F}^n$.

For $p = 1$,

$$\|x\|_1 = \sum_{i=1}^n |x_{(i)}| \quad (9.1.3)$$

is the *absolute sum norm*; for $p = 2$,

$$\|x\|_2 = \left(\sum_{i=1}^n |x_{(i)}|^2 \right)^{1/2} = \sqrt{x^*x} \quad (9.1.4)$$

is the *Euclidean norm*; and, for $p = \infty$,

$$\|x\|_\infty = \max_{i \in \{1, \dots, n\}} |x_{(i)}| \quad (9.1.5)$$

is the *infinity norm*.

The Hölder norms satisfy the following monotonicity property, which is related to the power-sum inequality given by Fact 1.15.34.

Proposition 9.1.5. Let $1 \leq p \leq q \leq \infty$, and let $x \in \mathbb{F}^n$. Then,

$$\|x\|_\infty \leq \|x\|_q \leq \|x\|_p \leq \|x\|_1. \quad (9.1.6)$$

Assume, in addition, that $1 < p < q < \infty$. Then, x has at least two nonzero components if and only if

$$\|x\|_\infty < \|x\|_q < \|x\|_p < \|x\|_1. \quad (9.1.7)$$

Proof. If either $p = q$ or $x = 0$ or x has exactly one nonzero component, then $\|x\|_q = \|x\|_p$. Hence, to prove both (9.1.6) and (9.1.7), it suffices to prove (9.1.7) in the case that $1 < p < q < \infty$ and x has at least two nonzero components. Thus, let $n \geq 2$, let $x \in \mathbb{F}^n$ have at least two nonzero components, and define $f: [1, \infty) \rightarrow [0, \infty)$ by $f(\beta) \triangleq \|x\|_\beta$. Hence,

$$f'(\beta) = \frac{1}{\beta} \|x\|_\beta^{1-\beta} \sum_{i=1}^n \gamma_i,$$

where, for all $i = 1, \dots, n$,

$$\gamma_i \triangleq \begin{cases} |x_i|^\beta (\log |x_{(i)}| - \log \|x\|_\beta), & x_{(i)} \neq 0, \\ 0, & x_{(i)} = 0. \end{cases}$$

If $x_{(i)} \neq 0$, then $\log |x_{(i)}| < \log \|x\|_\beta$. It thus follows that $f'(\beta) < 0$, which implies that f is decreasing on $[1, \infty)$. Hence, (9.1.7) holds. $\square$

The following result is *Hölder's inequality*. For this result we interpret $1/\infty = 0$. Note that, for all $x, y \in \mathbb{F}^n$, $|x^T y| \leq |x|^T |y| = \|x \circ y\|_1$.

Proposition 9.1.6. Let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$, and let $x, y \in \mathbb{F}^n$. Then,

$$|x^T y| \leq \|x\|_p \|y\|_q. \quad (9.1.8)$$

Furthermore, equality holds if and only if $|x^T y| = |x|^T |y|$ and

$$\begin{cases} |x| \circ |y| = \|y\|_\infty |x|, & p = 1, \\ \|y\|_q^{1/p} |x|^{\circ 1/q} = \|x\|_p^{1/q} |y|^{\circ 1/p}, & 1 < p < \infty, \\ |x| \circ |y| = \|x\|_\infty |y|, & p = \infty. \end{cases} \quad (9.1.9)$$

Proof. See [273, p. 127], [709, p. 536], [800, p. 71], Fact 1.16.11, and Fact 1.16.12. $\square$

The case $p = q = 2$ is the *Cauchy-Schwarz inequality*.

Corollary 9.1.7. Let $x, y \in \mathbb{F}^n$. Then,

$$|x^T y| \leq \|x\|_2 \|y\|_2. \quad (9.1.10)$$

Furthermore, equality holds if and only if x and y are linearly dependent.

Proof. Suppose that $y \neq 0$, and define $M \triangleq \begin{bmatrix} \sqrt{y^* y} I & (y^* y)^{-1/2} y \\ y^* y I & 1 \end{bmatrix}$. Since $M^* M = \begin{bmatrix} y^* y I & y \\ y^* & 1 \end{bmatrix}$ is positive semidefinite, it follows from *iii*) of Proposition 8.2.4 that $yy^* \leq y^* y I$. Therefore, $x^* y y^* x \leq x^* x y^* y$, which is equivalent to (9.1.10) with x replaced by $\bar{x}$.

Now, suppose that x and y are linearly dependent. Then, there exists $\beta \in \mathbb{F}$ such that either $x = \beta y$ or $y = \beta x$. In both cases it follows that $|x^* y| = \|x\|_2 \|y\|_2$. Conversely, define $f: \mathbb{F}^n \times \mathbb{F}^n \rightarrow [0, \infty)$ by $f(\mu, \nu) \triangleq \mu^* \mu \nu^* \nu - |\mu^* \nu|^2$. Now, suppose that $f(x, y) = 0$ so that (x, y) minimizes f . Then, it follows that $f_\mu(x, y) = 0$, which implies that $y^* y x = y^* x y$. Hence, x and y are linearly dependent. $\square$

The norms $\|\cdot\|$ and $\|\cdot\|'$ on $\mathbb{F}^n$ are *equivalent* if there exist $\alpha, \beta > 0$ such that

$$\alpha \|x\| \leq \|x\|' \leq \beta \|x\| \quad (9.1.11)$$

for all $x \in \mathbb{F}^n$. Note that these inequalities can be written as

$$\frac{1}{\beta} \|x\|' \leq \|x\| \leq \frac{1}{\alpha} \|x\|'. \quad (9.1.12)$$

Hence, the word “equivalent” is justified.

The following result shows that every pair of norms on $\mathbb{F}^n$ is equivalent.

Theorem 9.1.8. Let $\|\cdot\|$ and $\|\cdot\|'$ be norms on $\mathbb{F}^n$. Then, $\|\cdot\|$ and $\|\cdot\|'$ are equivalent.

Proof. See [709, p. 272]. $\square$

9.2 Matrix Norms

One way to define norms for matrices is by viewing a matrix $A \in \mathbb{F}^{n \times m}$ as a vector in $\mathbb{F}^{nm}$, for example, as $\text{vec } A$.

Definition 9.2.1. A *norm* $\|\cdot\|$ on $\mathbb{F}^{n \times m}$ is a function $\|\cdot\|: \mathbb{F}^{n \times m} \mapsto [0, \infty)$ that satisfies the following conditions:

- i) $\|A\| \geq 0$ for all $A \in \mathbb{F}^{n \times m}$.
- ii) $\|A\| = 0$ if and only if $A = 0$.

- iii) $\|\alpha A\| = |\alpha| \|A\|$ for all $\alpha \in \mathbb{F}$ and $A \in \mathbb{F}^{n \times m}$.
 iv) $\|A + B\| \leq \|A\| + \|B\|$ for all $A, B \in \mathbb{F}^{n \times m}$.

If $\|\cdot\|$ is a norm on $\mathbb{F}^{nm}$, then $\|\cdot\|'$ defined by $\|A\|' \triangleq \|\text{vec } A\|$ is a norm on $\mathbb{F}^{n \times m}$. For example, Hölder norms can be defined for matrices by choosing $\|\cdot\| = \|\cdot\|_p$. Hence, for all $A \in \mathbb{F}^{n \times m}$, define

$$\|A\|_p \triangleq \begin{cases} \left(\sum_{i=1}^n \sum_{j=1}^m |A_{(i,j)}|^p \right)^{1/p}, & 1 \leq p < \infty, \\ \max_{\substack{i \in \{1, \dots, n\} \\ j \in \{1, \dots, m\}}} |A_{(i,j)}|, & p = \infty. \end{cases} \quad (9.2.1)$$

Note that the same symbol $\|\cdot\|_p$ is used to denote the Hölder norm for both vectors and matrices. This notation is consistent since, if $A \in \mathbb{F}^{n \times 1}$, then $\|A\|_p$ coincides with the vector Hölder norm. Furthermore, if $A \in \mathbb{F}^{n \times m}$ and $1 \leq p \leq \infty$, then

$$\|A\|_p = \|\text{vec } A\|_p. \quad (9.2.2)$$

It follows from (9.1.6) that, if $A \in \mathbb{F}^{n \times m}$ and $1 \leq p \leq q \leq \infty$, then

$$\|A\|_\infty \leq \|A\|_q \leq \|A\|_p \leq \|A\|_1. \quad (9.2.3)$$

If, in addition, $1 < p < q < \infty$ and A has at least two nonzero entries, then

$$\|A\|_\infty < \|A\|_q < \|A\|_p < \|A\|_1. \quad (9.2.4)$$

The Hölder norms in the cases $p = 1, 2, \infty$ are the most commonly used. Let $A \in \mathbb{F}^{n \times m}$. For $p = 2$ we define the *Frobenius norm* $\|\cdot\|_F$ by

$$\|A\|_F \triangleq \|A\|_2. \quad (9.2.5)$$

Since $\|A\|_2 = \|\text{vec } A\|_2$, it follows that

$$\|A\|_F = \|A\|_2 = \|\text{vec } A\|_2 = \|\text{vec } A\|_F. \quad (9.2.6)$$

It is easy to see that

$$\|A\|_F = \sqrt{\text{tr } A^* A}. \quad (9.2.7)$$

Let $\|\cdot\|$ be a norm on $\mathbb{F}^{n \times m}$. If $\|S_1 A S_2\| = \|A\|$ for all $A \in \mathbb{F}^{n \times m}$ and for all unitary matrices $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$, then $\|\cdot\|$ is *unitarily invariant*. Now, let $m = n$. If $\|A\| = \|A^*\|$ for all $A \in \mathbb{F}^{n \times n}$, then $\|\cdot\|$ is *self-adjoint*. If $\|I_n\| = 1$, then $\|\cdot\|$ is *normalized*. Note that the Frobenius norm is not normalized since $\|I_n\|_F = \sqrt{n}$. If $\|S A S^*\| = \|A\|$ for all $A \in \mathbb{F}^{n \times n}$ and for all unitary $S \in \mathbb{F}^{n \times n}$, then $\|\cdot\|$ is *weakly unitarily invariant*.

Matrix norms can be defined in terms of singular values. Let $\sigma_1(A) \geq \sigma_2(A) \geq \dots$ denote the singular values of $A \in \mathbb{F}^{n \times m}$. The following result gives a weak majorization condition for singular values.

Proposition 9.2.2. Let $A, B \in \mathbb{F}^{n \times m}$. Then, for all $k = 1, \dots, \min\{n, m\}$,

$$\sum_{i=1}^k [\sigma_i(A) - \sigma_i(B)] \leq \sum_{i=1}^k \sigma_i(A+B) \leq \sum_{i=1}^k [\sigma_i(A) + \sigma_i(B)]. \quad (9.2.8)$$

In particular,

$$\sigma_{\max}(A) - \sigma_{\max}(B) \leq \sigma_{\max}(A+B) \leq \sigma_{\max}(A) + \sigma_{\max}(B) \quad (9.2.9)$$

and

$$\operatorname{tr} \langle A \rangle - \operatorname{tr} \langle B \rangle \leq \operatorname{tr} \langle A+B \rangle \leq \operatorname{tr} \langle A \rangle + \operatorname{tr} \langle B \rangle. \quad (9.2.10)$$

Proof. Define $\mathcal{A}, \mathcal{B} \in \mathbf{H}^{n+m}$ by $\mathcal{A} \triangleq \begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix}$ and $\mathcal{B} \triangleq \begin{bmatrix} 0 & B \\ B^* & 0 \end{bmatrix}$. Then, Corollary 8.6.19 implies that, for all $k = 1, \dots, n+m$,

$$\sum_{i=1}^k \lambda_i(\mathcal{A} + \mathcal{B}) \leq \sum_{i=1}^k [\lambda_i(\mathcal{A}) + \lambda_i(\mathcal{B})].$$

Now, consider $k \leq \min\{n, m\}$. Then, it follows from Proposition 5.6.6 that, for all $i = 1, \dots, k$, $\lambda_i(\mathcal{A}) = \sigma_i(A)$. Setting $k = 1$ yields (9.2.9), while setting $k = \min\{n, m\}$ and using Fact 8.17.2 yields (9.2.10). $\square$

Proposition 9.2.3. Let $p \in [1, \infty]$, and let $A \in \mathbb{F}^{n \times m}$. Then, $\|\cdot\|_{\sigma p}$ defined by

$$\|A\|_{\sigma p} \triangleq \begin{cases} \left(\sum_{i=1}^{\min\{n, m\}} \sigma_i^p(A) \right)^{1/p}, & 1 \leq p < \infty, \\ \sigma_{\max}(A), & p = \infty, \end{cases} \quad (9.2.11)$$

is a norm on $\mathbb{F}^{n \times m}$.

Proof. Let $p \in [1, \infty]$. Then, it follows from Proposition 9.2.2 and Minkowski's inequality Fact 1.16.25 that

$$\begin{aligned} \|A+B\|_{\sigma p} &= \left(\sum_{i=1}^{\min\{n, m\}} \sigma_i^p(A+B) \right)^{1/p} \\ &\leq \left(\sum_{i=1}^{\min\{n, m\}} [\sigma_i(A) + \sigma_i(B)]^p \right)^{1/p} \\ &\leq \left(\sum_{i=1}^{\min\{n, m\}} \sigma_i^p(A) \right)^{1/p} + \left(\sum_{i=1}^{\min\{n, m\}} \sigma_i^p(B) \right)^{1/p} \\ &= \|A\|_{\sigma p} + \|B\|_{\sigma p}. \end{aligned} \quad \square$$

The norm $\|\cdot\|_{\sigma p}$ is a *Schatten norm*. Let $A \in \mathbb{F}^{n \times m}$. Then, for all $p \in [1, \infty)$,

$$\|A\|_{\sigma p} = (\operatorname{tr} \langle A^p \rangle)^{1/p}. \quad (9.2.12)$$

Special cases are

$$\|A\|_{\sigma_1} = \sigma_1(A) + \cdots + \sigma_{\min\{n,m\}}(A) = \operatorname{tr} \langle A \rangle, \quad (9.2.13)$$

$$\|A\|_{\sigma_2} = \left[\sigma_1^2(A) + \cdots + \sigma_{\min\{n,m\}}^2(A) \right]^{1/2} = (\operatorname{tr} A^*A)^{1/2} = \|A\|_F, \quad (9.2.14)$$

and

$$\|A\|_{\sigma_\infty} = \sigma_1(A) = \sigma_{\max}(A), \quad (9.2.15)$$

which are the *trace norm*, Frobenius norm, and *spectral norm*, respectively.

By applying Proposition 9.1.5 to the vector $[\sigma_1(A) \cdots \sigma_{\min\{n,m\}}(A)]^T$, we obtain the following result.

Proposition 9.2.4. Let $p, q \in [1, \infty)$, where $p \leq q$, and let $A \in \mathbb{F}^{n \times m}$. Then,

$$\|A\|_{\sigma_\infty} \leq \|A\|_{\sigma_q} \leq \|A\|_{\sigma_p} \leq \|A\|_{\sigma_1}. \quad (9.2.16)$$

Assume, in addition, that $1 < p < q < \infty$ and $\operatorname{rank} A \geq 2$. Then,

$$\|A\|_\infty < \|A\|_q < \|A\|_p < \|A\|_1. \quad (9.2.17)$$

The norms $\|\cdot\|_{\sigma_p}$ are not very interesting when applied to vectors. Let $x \in \mathbb{F}^n = \mathbb{F}^{n \times 1}$. Then, $\sigma_{\max}(x) = (x^*x)^{1/2} = \|x\|_2$, and, since $\operatorname{rank} x \leq 1$, it follows that, for all $p \in [1, \infty]$,

$$\|x\|_{\sigma_p} = \|x\|_2. \quad (9.2.18)$$

Proposition 9.2.5. Let $A \in \mathbb{F}^{n \times m}$. If $p \in (0, 2]$, then

$$\|A\|_{\sigma_p} \leq \|A\|_p. \quad (9.2.19)$$

If $p \geq 2$, then

$$\|A\|_p \leq \|A\|_{\sigma_p}. \quad (9.2.20)$$

Proof. See [1485, p. 50]. □

Proposition 9.2.6. Let $\|\cdot\|$ be a norm on $\mathbb{F}^{n \times n}$, and let $A \in \mathbb{F}^{n \times n}$. Then,

$$\operatorname{sprad}(A) = \lim_{k \rightarrow \infty} \|A^k\|^{1/k}. \quad (9.2.21)$$

Proof. See [709, p. 322]. □

9.3 Compatible Norms

The norms $\|\cdot\|$, $\|\cdot\|'$, and $\|\cdot\|''$ on $\mathbb{F}^{n \times l}$, $\mathbb{F}^{n \times m}$, and $\mathbb{F}^{m \times l}$, respectively, are *compatible* if, for all $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$,

$$\|AB\| \leq \|A\|' \|B\|''. \quad (9.3.1)$$

For $l = 1$, the norms $\|\cdot\|$, $\|\cdot\|'$, and $\|\cdot\|''$ on $\mathbb{F}^n$, $\mathbb{F}^{n \times m}$, and $\mathbb{F}^m$, respectively, are compatible if, for all $A \in \mathbb{F}^{n \times m}$ and $x \in \mathbb{F}^m$,

$$\|Ax\| \leq \|A\|' \|x\|''. \quad (9.3.2)$$

Furthermore, the norm $\|\cdot\|$ on $\mathbb{F}^n$ is *compatible* with the norm $\|\cdot\|'$ on $\mathbb{F}^{n \times n}$ if, for all $A \in \mathbb{F}^{n \times n}$ and $x \in \mathbb{F}^n$,

$$\|Ax\| \leq \|A\|' \|x\|. \quad (9.3.3)$$

Note that $\|I_n\|' \geq 1$. The norm $\|\cdot\|$ on $\mathbb{F}^{n \times n}$ is *submultiplicative* if, for all $A, B \in \mathbb{F}^{n \times n}$,

$$\|AB\| \leq \|A\| \|B\|. \quad (9.3.4)$$

Hence, the norm $\|\cdot\|$ on $\mathbb{F}^{n \times n}$ is submultiplicative if and only if $\|\cdot\|$, $\|\cdot\|'$, and $\|\cdot\|''$ are compatible. In this case, $\|I_n\| \geq 1$, while $\|\cdot\|$ is normalized if and only if $\|I_n\| = 1$.

Proposition 9.3.1. Let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$, and let $y \in \mathbb{F}^n$ be nonzero. Then, $\|x\|' \triangleq \|xy^*\|$ is a norm on $\mathbb{F}^n$, and $\|\cdot\|'$ is compatible with $\|\cdot\|$.

Proposition 9.3.2. Let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$, and let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{sprad}(A) \leq \|A\|. \quad (9.3.5)$$

Proof. Use Proposition 9.3.1 to construct a norm $\|\cdot\|'$ on $\mathbb{F}^n$ that is compatible with $\|\cdot\|$. Furthermore, let $A \in \mathbb{F}^{n \times n}$, let $\lambda \in \text{spec}(A)$, and let $x \in \mathbb{C}^n$ be an eigenvector of A associated with λ . Then, $Ax = \lambda x$ implies that $|\lambda| \|x\|' = \|Ax\|' \leq \|A\| \|x\|'$, and thus $|\lambda| \leq \|A\|$, which implies (9.3.5). Alternatively, under the additional assumption that $\|\cdot\|$ is submultiplicative, it follows from Proposition 9.2.6 that

$$\text{sprad}(A) = \lim_{k \rightarrow \infty} \|A^k\|^{1/k} \leq \lim_{k \rightarrow \infty} \|A\|^{k/k} = \|A\|. \quad \square$$

Proposition 9.3.3. Let $A \in \mathbb{F}^{n \times n}$, and let $\varepsilon > 0$. Then, there exists a submultiplicative norm $\|\cdot\|$ on $\mathbb{F}^{n \times n}$ such that

$$\text{sprad}(A) \leq \|A\| \leq \text{sprad}(A) + \varepsilon. \quad (9.3.6)$$

Proof. See [709, p. 297]. $\square$

Corollary 9.3.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{sprad}(A) < 1$. Then, there exists a submultiplicative norm $\|\cdot\|$ on $\mathbb{F}^{n \times n}$ such that $\|A\| < 1$.

We now identify some compatible norms. We begin with the Hölder norms.

Proposition 9.3.5. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. If $p \in [1, 2]$, then

$$\|AB\|_p \leq \|A\|_p \|B\|_p. \quad (9.3.7)$$

If $p \in [2, \infty]$ and q satisfies $1/p + 1/q = 1$, then

$$\|AB\|_p \leq \|A\|_p \|B\|_q \quad (9.3.8)$$

and

$$\|AB\|_p \leq \|A\|_q \|B\|_p. \quad (9.3.9)$$

Proof. First let $1 \leq p \leq 2$ so that $q \triangleq p/(p-1) \geq 2$. Using Hölder's inequality (9.1.8) and (9.1.6) with $p \leq q$ yields

$$\begin{aligned} \|AB\|_p &= \left(\sum_{i,j=1}^{n,l} |\text{row}_i(A) \text{col}_j(B)|^p \right)^{1/p} \\ &\leq \left(\sum_{i,j=1}^{n,l} \|\text{row}_i(A)\|_p^p \|\text{col}_j(B)\|_q^p \right)^{1/p} \\ &= \left(\sum_{i=1}^n \|\text{row}_i(A)\|_p^p \right)^{1/p} \left(\sum_{j=1}^l \|\text{col}_j(B)\|_q^p \right)^{1/p} \\ &\leq \left(\sum_{i=1}^n \|\text{row}_i(A)\|_p^p \right)^{1/p} \left(\sum_{j=1}^l \|\text{col}_j(B)\|_p^p \right)^{1/p} \\ &= \|A\|_p \|B\|_p. \end{aligned}$$

Next, let $2 \leq p \leq \infty$ so that $q \triangleq p/(p-1) \leq 2$. Using Hölder's inequality (9.1.8) and (9.1.6) with $q \leq p$ yields

$$\begin{aligned} \|AB\|_p &\leq \left(\sum_{i=1}^n \|\text{row}_i(A)\|_p^p \right)^{1/p} \left(\sum_{j=1}^l \|\text{col}_j(B)\|_q^p \right)^{1/p} \\ &\leq \left(\sum_{i=1}^n \|\text{row}_i(A)\|_p^p \right)^{1/p} \left(\sum_{j=1}^l \|\text{col}_j(B)\|_q^q \right)^{1/q} \\ &= \|A\|_p \|B\|_q. \end{aligned}$$

Similarly, it can be shown that (9.3.9) holds. $\square$

Proposition 9.3.6. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $p, q \in [1, \infty]$, define

$$r \triangleq \frac{1}{\frac{1}{p} + \frac{1}{q}},$$

and assume that $r \geq 1$. Then,

$$\|AB\|_{\sigma r} \leq \|A\|_{\sigma p} \|B\|_{\sigma q}. \quad (9.3.10)$$

In particular,

$$\|AB\|_{\sigma r} \leq \|A\|_{\sigma 2r} \|B\|_{\sigma 2r}. \quad (9.3.11)$$

Proof. Using Proposition 9.6.2 and Hölder's inequality with $1/(p/r) + 1/(q/r) = 1$, it follows that

$$\begin{aligned}
 \|AB\|_{\sigma r} &= \left(\sum_{i=1}^{\min\{n,m,l\}} \sigma_i^r(AB) \right)^{1/r} \\
 &\leq \left(\sum_{i=1}^{\min\{n,m,l\}} \sigma_i^r(A) \sigma_i^r(B) \right)^{1/r} \\
 &\leq \left[\left(\sum_{i=1}^{\min\{n,m,l\}} \sigma_i^p(A) \right)^{r/p} \left(\sum_{i=1}^{\min\{n,m,l\}} \sigma_i^q(B) \right)^{r/q} \right]^{1/r} \\
 &= \|A\|_{\sigma p} \|B\|_{\sigma q}. \quad \square
 \end{aligned}$$

Corollary 9.3.7. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\|AB\|_{\sigma\infty} \leq \|AB\|_{\sigma 2} \leq \left\{ \begin{array}{c} \|A\|_{\sigma\infty} \|B\|_{\sigma 2} \\ \|A\|_{\sigma 2} \|B\|_{\sigma\infty} \\ \|AB\|_{\sigma 1} \end{array} \right\} \leq \|A\|_{\sigma 2} \|B\|_{\sigma 2} \quad (9.3.12)$$

or, equivalently,

$$\sigma_{\max}(AB) \leq \|AB\|_{\mathbb{F}} \leq \left\{ \begin{array}{c} \sigma_{\max}(A) \|B\|_{\mathbb{F}} \\ \|A\|_{\mathbb{F}} \sigma_{\max}(B) \\ \text{tr} \langle AB \rangle \end{array} \right\} \leq \|A\|_{\mathbb{F}} \|B\|_{\mathbb{F}}. \quad (9.3.13)$$

Furthermore, for all $r \in [1, \infty]$,

$$\|AB\|_{\sigma 2r} \leq \|AB\|_{\sigma r} \leq \left\{ \begin{array}{c} \|A\|_{\sigma r} \sigma_{\max}(B) \\ \sigma_{\max}(A) \|B\|_{\sigma r} \\ \|A\|_{\sigma 2r} \|B\|_{\sigma 2r} \end{array} \right\} \leq \|A\|_{\sigma r} \|B\|_{\sigma r}. \quad (9.3.14)$$

In particular, setting $r = \infty$ yields

$$\sigma_{\max}(AB) \leq \sigma_{\max}(A) \sigma_{\max}(B). \quad (9.3.15)$$

Corollary 9.3.8. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\|AB\|_{\sigma 1} \leq \left\{ \begin{array}{c} \sigma_{\max}(A) \|B\|_{\sigma 1} \\ \|A\|_{\sigma 1} \sigma_{\max}(B). \end{array} \right. \quad (9.3.16)$$

Note that the inequality $\|AB\|_{\mathbb{F}} \leq \|A\|_{\mathbb{F}} \|B\|_{\mathbb{F}}$ in (9.3.13) is equivalent to (9.3.7) with $p = 2$ as well as (9.3.8) and (9.3.9) with $p = q = 2$.

The following result is the matrix version of the Cauchy-Schwarz inequality Corollary 9.1.7.

Corollary 9.3.9. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then,

$$|\operatorname{tr} A^*B| \leq \|A\|_{\mathbb{F}} \|B\|_{\mathbb{F}}. \quad (9.3.17)$$

Equality holds if and only if A and B^* are linearly dependent.

9.4 Induced Norms

In this section we consider the case in which there exists a nonzero vector $x \in \mathbb{F}^m$ such that (9.3.3) holds as an equality. This condition characterizes a special class of norms on $\mathbb{F}^{n \times n}$, namely, the *induced norms*.

Definition 9.4.1. Let $\|\cdot\|''$ and $\|\cdot\|$ be norms on $\mathbb{F}^m$ and $\mathbb{F}^n$, respectively. Then, $\|\cdot\|': \mathbb{F}^{n \times m} \mapsto \mathbb{F}$ defined by

$$\|A\|' = \max_{x \in \mathbb{F}^m \setminus \{0\}} \frac{\|Ax\|}{\|x\|''} \quad (9.4.1)$$

is an *induced norm* on $\mathbb{F}^{n \times m}$. In this case, $\|\cdot\|'$ is *induced by* $\|\cdot\|''$ and $\|\cdot\|$. If $m = n$ and $\|\cdot\|'' = \|\cdot\|$, then $\|\cdot\|'$ is *induced by* $\|\cdot\|$, and $\|\cdot\|'$ is an *equi-induced norm*.

The next result confirms that $\|\cdot\|'$ defined by (9.4.1) is a norm.

Theorem 9.4.2. Every induced norm is a norm. Furthermore, every equi-induced norm is normalized.

Proof. See [709, p. 293]. □

Let $A \in \mathbb{F}^{n \times m}$. It can be seen that (9.4.1) is equivalent to

$$\|A\|' = \max_{x \in \{y \in \mathbb{F}^m: \|y\|''=1\}} \|Ax\|. \quad (9.4.2)$$

Theorem 10.3.8 implies that the maximum in (9.4.2) exists. Since, for all $x \neq 0$,

$$\|A\|' = \max_{x \in \mathbb{F}^m \setminus \{0\}} \frac{\|Ax\|}{\|x\|''} \geq \frac{\|Ax\|}{\|x\|''}, \quad (9.4.3)$$

it follows that, for all $x \in \mathbb{F}^m$,

$$\|Ax\| \leq \|A\|' \|x\|'' \quad (9.4.4)$$

so that $\|\cdot\|$, $\|\cdot\|'$, and $\|\cdot\|''$ are compatible. If $m = n$ and $\|\cdot\|'' = \|\cdot\|$, then the norm $\|\cdot\|$ is compatible with the induced norm $\|\cdot\|'$. The next result shows that compatible norms can be obtained from induced norms.

Proposition 9.4.3. Let $\|\cdot\|$, $\|\cdot\|'$, and $\|\cdot\|''$ be norms on $\mathbb{F}^l$, $\mathbb{F}^m$, and $\mathbb{F}^n$, respectively. Furthermore, let $\|\cdot\|'''$ be the norm on $\mathbb{F}^{m \times l}$ induced by $\|\cdot\|$ and $\|\cdot\|'$, let $\|\cdot\|''''$ be the norm on $\mathbb{F}^{n \times m}$ induced by $\|\cdot\|'$ and $\|\cdot\|''$, and let $\|\cdot\|'''''$ be the norm on $\mathbb{F}^{n \times l}$ induced by $\|\cdot\|$ and $\|\cdot\|''$. If $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, then

$$\|AB\|''''' \leq \|A\|'''' \|B\|'''. \quad (9.4.5)$$

Proof. Note that, for all $x \in \mathbb{F}^l$, $\|Bx\|' \leq \|B\|''' \|x\|$, and, for all $y \in \mathbb{F}^m$, $\|Ay\|'' \leq \|A\|'''' \|y\|'$. Hence, for all $x \in \mathbb{F}^l$, it follows that

$$\|ABx\|'' \leq \|A\|'''' \|Bx\|' \leq \|A\|'''' \|B\|''' \|x\|,$$

which implies that

$$\|AB\|'''' = \max_{x \in \mathbb{F}^l \setminus \{0\}} \frac{\|ABx\|''}{\|x\|} \leq \|A\|'''' \|B\|'''. \quad \square$$

Corollary 9.4.4. Every equi-induced norm is submultiplicative.

The following result is a consequence of Corollary 9.4.4 and Proposition 9.3.2.

Corollary 9.4.5. Let $\|\cdot\|$ be an equi-induced norm on $\mathbb{F}^{n \times n}$, and let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{sprad}(A) \leq \|A\|. \quad (9.4.6)$$

By assigning $\|\cdot\|_p$ to $\mathbb{F}^m$ and $\|\cdot\|_q$ to $\mathbb{F}^n$, the *Hölder-induced norm* on $\mathbb{F}^{n \times m}$ is defined by

$$\|A\|_{q,p} \triangleq \max_{x \in \mathbb{F}^m \setminus \{0\}} \frac{\|Ax\|_q}{\|x\|_p}. \quad (9.4.7)$$

Proposition 9.4.6. Let $p, q, p', q' \in [1, \infty]$, where $p \leq p'$ and $q \leq q'$, and let $A \in \mathbb{F}^{n \times m}$. Then,

$$\|A\|_{q',p} \leq \|A\|_{q,p} \leq \|A\|_{q,p'}. \quad (9.4.8)$$

Proof. The result follows from Proposition 9.1.5. $\square$

A subtlety of induced norms is that the value of an induced norm may depend on the underlying field. In particular, the value of the induced norm of a real matrix A computed over the complex field may be different from the induced norm of A computed over the real field. Although the chosen field is usually not made explicit, we do so in special cases for clarity.

Proposition 9.4.7. Let $A \in \mathbb{R}^{n \times m}$, and let $\|A\|_{p,q,\mathbb{F}}$ denote the Hölder-induced norm of A evaluated over the field $\mathbb{F}$. Then,

$$\|A\|_{p,q,\mathbb{R}} \leq \|A\|_{p,q,\mathbb{C}}. \quad (9.4.9)$$

If $p \in [1, \infty]$, then

$$\|A\|_{p,1,\mathbb{R}} = \|A\|_{p,1,\mathbb{C}}. \quad (9.4.10)$$

Finally, if $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$, then

$$\|A\|_{\infty,p,\mathbb{R}} = \|A\|_{\infty,p,\mathbb{C}}. \quad (9.4.11)$$

Proof. See [690, p. 716]. $\square$

Example 9.4.8. Let $A = \begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$ and $x = \begin{bmatrix} x_1 & x_2 \end{bmatrix}^T$. Then, $\|Ax\|_1 = |x_1 - x_2| + |x_1 + x_2|$. Letting $x = \begin{bmatrix} 1 & j \end{bmatrix}^T$ so that $\|x\|_\infty = 1$, it follows that

$\|A\|_{1,\infty,\mathbb{C}} \geq 2\sqrt{2}$. On the other hand, $\|A\|_{1,\infty,\mathbb{R}} = 2$. Hence, in this case, the inequality (9.4.9) is strict. See [690, p. 716].

The following result gives explicit expressions for several Hölder-induced norms.

Proposition 9.4.9. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\|A\|_{2,2} = \sigma_{\max}(A). \quad (9.4.12)$$

If $p \in [1, \infty]$, then

$$\|A\|_{p,1} = \max_{i \in \{1, \dots, m\}} \|\text{col}_i(A)\|_p. \quad (9.4.13)$$

Finally, if $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$, then

$$\|A\|_{\infty,p} = \max_{i \in \{1, \dots, n\}} \|\text{row}_i(A)\|_q. \quad (9.4.14)$$

Proof. Since A^*A is Hermitian, it follows from Corollary 8.4.2 that, for all $x \in \mathbb{F}^m$,

$$x^*A^*Ax \leq \lambda_{\max}(A^*A)x^*x,$$

which implies that, for all $x \in \mathbb{F}^m$, $\|Ax\|_2 \leq \sigma_{\max}(A)\|x\|_2$, and thus $\|A\|_{2,2} \leq \sigma_{\max}(A)$. Now, let $x \in \mathbb{F}^{n \times n}$ be an eigenvector associated with $\lambda_{\max}(A^*A)$ so that $\|Ax\|_2 = \sigma_{\max}(A)\|x\|_2$, which implies that $\sigma_{\max}(A) \leq \|A\|_{2,2}$. Hence, (9.4.12) holds.

Next, note that, for all $x \in \mathbb{F}^m$,

$$\|Ax\|_p = \left\| \sum_{i=1}^m x_{(i)} \text{col}_i(A) \right\|_p \leq \sum_{i=1}^m |x_{(i)}| \|\text{col}_i(A)\|_p \leq \max_{i \in \{1, \dots, m\}} \|\text{col}_i(A)\|_p \|x\|_1,$$

and hence $\|A\|_{p,1} \leq \max_{i \in \{1, \dots, m\}} \|\text{col}_i(A)\|_p$. Next, let $j \in \{1, \dots, m\}$ be such that $\|\text{col}_j(A)\|_p = \max_{i \in \{1, \dots, m\}} \|\text{col}_i(A)\|_p$. Now, since $\|e_j\|_1 = 1$, it follows that $\|Ae_j\|_p = \|\text{col}_j(A)\|_p \|e_j\|_1$, which implies that

$$\max_{i \in \{1, \dots, m\}} \|\text{col}_i(A)\|_p = \|\text{col}_j(A)\|_p \leq \|A\|_{p,1},$$

and hence (9.4.13) holds.

Next, for all $x \in \mathbb{F}^m$, it follows from Hölder's inequality (9.1.8) that

$$\|Ax\|_{\infty} = \max_{i \in \{1, \dots, n\}} |\text{row}_i(A)x| \leq \max_{i \in \{1, \dots, n\}} \|\text{row}_i(A)\|_q \|x\|_p,$$

which implies that $\|A\|_{\infty,p} \leq \max_{i \in \{1, \dots, n\}} \|\text{row}_i(A)\|_q$. Next, let $j \in \{1, \dots, n\}$ be such that $\|\text{row}_j(A)\|_q = \max_{i \in \{1, \dots, n\}} \|\text{row}_i(A)\|_q$, and let nonzero $x \in \mathbb{F}^m$ be such that $|\text{row}_j(A)x| = \|\text{row}_j(A)\|_q \|x\|_p$. Hence,

$$\|Ax\|_{\infty} = \max_{i \in \{1, \dots, n\}} |\text{row}_i(A)x| \geq |\text{row}_j(A)x| = \|\text{row}_j(A)\|_q \|x\|_p,$$

which implies that

$$\max_{i \in \{1, \dots, n\}} \|\text{row}_i(A)\|_q = \|\text{row}_j(A)\|_q \leq \|A\|_{\infty,p},$$

and thus (9.4.14) holds. $\square$

Note that

$$\max_{i \in \{1, \dots, m\}} \|\text{col}_i(A)\|_2 = d_{\max}^{1/2}(A^*A) \quad (9.4.15)$$

and

$$\max_{i \in \{1, \dots, n\}} \|\text{row}_i(A)\|_2 = d_{\max}^{1/2}(AA^*). \quad (9.4.16)$$

Therefore, it follows from Proposition 9.4.9 that

$$\|A\|_{1,1} = \max_{i \in \{1, \dots, m\}} \|\text{col}_i(A)\|_1, \quad (9.4.17)$$

$$\|A\|_{2,1} = \max_{i \in \{1, \dots, m\}} \|\text{col}_i(A)\|_2 = d_{\max}^{1/2}(A^*A), \quad (9.4.18)$$

$$\|A\|_{\infty,1} = \|A\|_{\infty} = \max_{\substack{i \in \{1, \dots, n\} \\ j \in \{1, \dots, m\}}} |A_{(i,j)}|, \quad (9.4.19)$$

$$\|A\|_{\infty,2} = \max_{i \in \{1, \dots, n\}} \|\text{row}_i(A)\|_2 = d_{\max}^{1/2}(AA^*), \quad (9.4.20)$$

$$\|A\|_{\infty,\infty} = \max_{i \in \{1, \dots, n\}} \|\text{row}_i(A)\|_1. \quad (9.4.21)$$

For convenience, we define the *column norm*

$$\|A\|_{\text{col}} \triangleq \|A\|_{1,1} \quad (9.4.22)$$

and the *row norm*

$$\|A\|_{\text{row}} \triangleq \|A\|_{\infty,\infty}. \quad (9.4.23)$$

The following result follows from Corollary 9.4.5.

Corollary 9.4.10. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{sprad}(A) \leq \sigma_{\max}(A), \quad (9.4.24)$$

$$\text{sprad}(A) \leq \|A\|_{\text{col}}, \quad (9.4.25)$$

$$\text{sprad}(A) \leq \|A\|_{\text{row}}. \quad (9.4.26)$$

Proposition 9.4.11. Let $p, q \in [1, \infty]$ be such that $1/p + 1/q = 1$, and let $A \in \mathbb{F}^{n \times m}$. Then,

$$\|A\|_{q,p} \leq \|A\|_q. \quad (9.4.27)$$

Proof. For $p = 1$ and $q = \infty$, (9.4.27) follows from (9.4.19). For $q < \infty$ and $x \in \mathbb{F}^n$, it follows from Hölder's inequality (9.1.8) that

$$\begin{aligned}
\|Ax\|_q &= \left(\sum_{i=1}^n |\text{row}_i(A)x|^q \right)^{1/q} \leq \left(\sum_{i=1}^n \|\text{row}_i(A)\|_q^q \|x\|_p^q \right)^{1/q} \\
&= \left(\sum_{i=1}^n \sum_{j=1}^m |A_{(i,j)}|^q \right)^{1/q} \|x\|_p = \|A\|_q \|x\|_p,
\end{aligned}$$

which implies (9.4.27). $\square$

Next, we specialize Proposition 9.4.3 to the Hölder-induced norms.

Corollary 9.4.12. Let $p, q, r \in [1, \infty]$, and let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\|AB\|_{r,p} \leq \|A\|_{r,q} \|B\|_{q,p}. \quad (9.4.28)$$

In particular,

$$\|AB\|_{\text{col}} \leq \|A\|_{\text{col}} \|B\|_{\text{col}}, \quad (9.4.29)$$

$$\sigma_{\max}(AB) \leq \sigma_{\max}(A) \sigma_{\max}(B), \quad (9.4.30)$$

$$\|AB\|_{\text{row}} \leq \|A\|_{\text{row}} \|B\|_{\text{row}}, \quad (9.4.31)$$

$$\|AB\|_{\infty} \leq \|A\|_{\infty} \|B\|_{\text{col}}, \quad (9.4.32)$$

$$\|AB\|_{\infty} \leq \|A\|_{\text{row}} \|B\|_{\infty}, \quad (9.4.33)$$

$$d_{\max}^{1/2}(B^*A^*AB) \leq d_{\max}^{1/2}(A^*A) \|B\|_{\text{col}}, \quad (9.4.34)$$

$$d_{\max}^{1/2}(B^*A^*AB) \leq \sigma_{\max}(A) d_{\max}^{1/2}(B^*B), \quad (9.4.35)$$

$$d_{\max}^{1/2}(ABB^*A^*) \leq d_{\max}^{1/2}(AA^*) \sigma_{\max}(B), \quad (9.4.36)$$

$$d_{\max}^{1/2}(ABB^*A^*) \leq \|B\|_{\text{row}} d_{\max}^{1/2}(BB^*). \quad (9.4.37)$$

The following result is often useful.

Proposition 9.4.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{sprad}(A) < 1$. Then, there exists a submultiplicative norm $\|\cdot\|$ on $\mathbb{F}^{n \times n}$ such that $\|A\| < 1$. Furthermore, the series $\sum_{k=0}^{\infty} A^k$ converges absolutely, and

$$(I - A)^{-1} = \sum_{k=0}^{\infty} A^k. \quad (9.4.38)$$

Finally,

$$\frac{1}{1 + \|A\|} \leq \|(I - A)^{-1}\| \leq \frac{1}{1 - \|A\|} + \|I\| - 1. \quad (9.4.39)$$

If, in addition, $\|\cdot\|$ is normalized, then

$$\frac{1}{1 + \|A\|} \leq \|(I - A)^{-1}\| \leq \frac{1}{1 - \|A\|}. \quad (9.4.40)$$

Proof. Corollary 9.3.4 implies that there exists a submultiplicative norm $\|\cdot\|$ on $\mathbb{F}^{n \times n}$ such that $\|A\| < 1$. It thus follows that

$$\left\| \sum_{k=0}^{\infty} A^k \right\| \leq \sum_{k=0}^{\infty} \|A^k\| \leq \|I\| - 1 + \sum_{k=0}^{\infty} \|A\|^k = \frac{1}{1 - \|A\|} + \|I\| - 1,$$

which proves that the series $\sum_{k=0}^{\infty} A^k$ converges absolutely.

Next, we show that $I - A$ is nonsingular. If $I - A$ is singular, then there exists a nonzero vector $x \in \mathbb{C}^n$ such that $Ax = x$. Hence, $1 \in \text{spec}(A)$, which contradicts $\text{sprad}(A) < 1$. Next, to verify (9.4.38), note that

$$(I - A) \sum_{k=0}^{\infty} A^k = \sum_{k=0}^{\infty} A^k - \sum_{k=1}^{\infty} A^k = I + \sum_{k=1}^{\infty} A^k - \sum_{k=1}^{\infty} A^k = I,$$

which implies (9.4.38) and thus the right-hand inequality in (9.4.39). Furthermore,

$$\begin{aligned} 1 &\leq \|I\| \\ &= \|(I - A)(I - A)^{-1}\| \\ &\leq \|I - A\| \|(I - A)^{-1}\| \\ &\leq (1 + \|A\|) \|(I - A)^{-1}\|, \end{aligned}$$

which yields the left-hand inequality in (9.4.39). $\square$

9.5 Induced Lower Bound

We now consider a variation of the induced norm.

Definition 9.5.1. Let $\|\cdot\|$ and $\|\cdot\|'$ denote norms on $\mathbb{F}^m$ and $\mathbb{F}^n$, respectively, and let $A \in \mathbb{F}^{n \times m}$. Then, $\ell: \mathbb{F}^{n \times m} \mapsto \mathbb{R}$ defined by

$$\ell(A) \triangleq \begin{cases} \min_{y \in \mathcal{R}(A) \setminus \{0\}} \max_{x \in \{z \in \mathbb{F}^m: Az=y\}} \frac{\|y\|'}{\|x\|}, & A \neq 0, \\ 0, & A = 0, \end{cases} \quad (9.5.1)$$

is the *lower bound induced by $\|\cdot\|$ and $\|\cdot\|'$* . Equivalently,

$$\ell(A) \triangleq \begin{cases} \min_{x \in \mathbb{F}^m \setminus \mathcal{N}(A)} \max_{z \in \mathcal{N}(A)} \frac{\|Ax\|'}{\|x+z\|}, & A \neq 0, \\ 0, & A = 0. \end{cases} \quad (9.5.2)$$

Proposition 9.5.2. Let $\|\cdot\|$ and $\|\cdot\|'$ be norms on $\mathbb{F}^m$ and $\mathbb{F}^n$, respectively, let $\|\cdot\|''$ be the norm induced by $\|\cdot\|$ and $\|\cdot\|'$, let $\|\cdot\|'''$ be the norm induced by $\|\cdot\|'$ and $\|\cdot\|$, and let ℓ be the lower bound induced by $\|\cdot\|$ and $\|\cdot\|'$. Then, the following statements hold:

- i) $\ell(A)$ exists for all $A \in \mathbb{F}^{n \times m}$, that is, the minimum in (9.5.1) is attained.
- ii) If $A \in \mathbb{F}^{n \times m}$, then $\ell(A) = 0$ if and only if $A = 0$.

iii) For all $A \in \mathbb{F}^{n \times m}$ there exists a vector $x \in \mathbb{F}^m$ such that

$$\ell(A)\|x\| = \|Ax\|'. \quad (9.5.3)$$

iv) For all $A \in \mathbb{F}^{n \times m}$,

$$\ell(A) \leq \|A\|''. \quad (9.5.4)$$

v) If $A \neq 0$ and B is a (1)-inverse of A , then

$$1/\|B\|''' \leq \ell(A) \leq \|B\|'''. \quad (9.5.5)$$

vi) If $A, B \in \mathbb{F}^{n \times m}$ and either $\mathcal{R}(A) \subseteq \mathcal{R}(A+B)$ or $\mathcal{N}(A) \subseteq \mathcal{N}(A+B)$, then

$$\ell(A) - \|B\|''' \leq \ell(A+B). \quad (9.5.6)$$

vii) If $A, B \in \mathbb{F}^{n \times m}$ and either $\mathcal{R}(A+B) \subseteq \mathcal{R}(A)$ or $\mathcal{N}(A+B) \subseteq \mathcal{N}(A)$, then

$$\ell(A+B) \leq \ell(A) + \|B\|'''. \quad (9.5.7)$$

viii) If $n = m$ and $A \in \mathbb{F}^{n \times n}$ is nonsingular, then

$$\ell(A) = 1/\|A^{-1}\|'''. \quad (9.5.8)$$

Proof. See [582]. □

Proposition 9.5.3. Let $\|\cdot\|$, $\|\cdot\|'$, and $\|\cdot\|''$ be norms on $\mathbb{F}^l$, $\mathbb{F}^m$, and $\mathbb{F}^n$, respectively, let $\|\cdot\|'''$ denote the norm on $\mathbb{F}^{m \times l}$ induced by $\|\cdot\|$ and $\|\cdot\|'$, let $\|\cdot\|''''$ denote the norm on $\mathbb{F}^{n \times m}$ induced by $\|\cdot\|'$ and $\|\cdot\|''$, and let $\|\cdot\|'''''$ denote the norm on $\mathbb{F}^{n \times l}$ induced by $\|\cdot\|$ and $\|\cdot\|''$. If $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, then

$$\ell(A)\ell'(B) \leq \ell''(AB). \quad (9.5.9)$$

In addition, the following statements hold:

i) If either $\text{rank } B = \text{rank } AB$ or $\text{def } B = \text{def } AB$, then

$$\ell''(AB) \leq \|A\|''\ell(B). \quad (9.5.10)$$

ii) If $\text{rank } A = \text{rank } AB$, then

$$\ell''(AB) \leq \ell(A)\|B\|'''''. \quad (9.5.11)$$

iii) If $\text{rank } B = m$, then

$$\|A\|''\ell(B) \leq \|AB\|'''''. \quad (9.5.12)$$

iv) If $\text{rank } A = m$, then

$$\ell(A)\|B\|''''' \leq \|AB\|'''''. \quad (9.5.13)$$

Proof. See [582]. □

By assigning $\|\cdot\|_p$ to $\mathbb{F}^m$ and $\|\cdot\|_q$ to $\mathbb{F}^n$, the *Hölder-induced lower bound* on $\mathbb{F}^{n \times m}$ is defined by

$$\ell_{q,p}(A) \triangleq \begin{cases} \min_{y \in \mathcal{R}(A) \setminus \{0\}} \max_{x \in \{z \in \mathbb{F}^m: Az=y\}} \frac{\|y\|_q}{\|x\|_p}, & A \neq 0, \\ 0, & A = 0. \end{cases} \quad (9.5.14)$$

The following result shows that $\ell_{2,2}(A)$ is the smallest positive singular value of A .

Proposition 9.5.4. Let $A \in \mathbb{F}^{n \times m}$, assume that A is nonzero, and let $r \triangleq \text{rank } A$. Then,

$$\ell_{2,2}(A) = \sigma_r(A). \quad (9.5.15)$$

Proof. The result follows from the singular value decomposition. $\square$

Corollary 9.5.5. Let $A \in \mathbb{F}^{n \times m}$. If $n \leq m$ and A is right invertible, then

$$\ell_{2,2}(A) = \sigma_{\min}(A) = \sigma_n(A). \quad (9.5.16)$$

If $m \leq n$ and A is left invertible, then

$$\ell_{2,2}(A) = \sigma_{\min}(A) = \sigma_m(A). \quad (9.5.17)$$

Finally, if $n = m$ and A is nonsingular, then

$$\ell_{2,2}(A^{-1}) = \sigma_{\min}(A^{-1}) = \frac{1}{\sigma_{\max}(A)}. \quad (9.5.18)$$

Proof. Use Proposition 5.6.2 and Fact 6.3.29. $\square$

In contrast to the submultiplicativity condition (9.4.4) satisfied by the induced norm, the induced lower bound satisfies a supermultiplicativity condition. The following result is analogous to Proposition 9.4.3.

Proposition 9.5.6. Let $\|\cdot\|$, $\|\cdot\|'$, and $\|\cdot\|''$ be norms on $\mathbb{F}^l$, $\mathbb{F}^m$, and $\mathbb{F}^n$, respectively. Let $\ell(\cdot)$ be the lower bound induced by $\|\cdot\|$ and $\|\cdot\|'$, let $\ell'(\cdot)$ be the lower bound induced by $\|\cdot\|'$ and $\|\cdot\|''$, let $\ell''(\cdot)$ be the lower bound induced by $\|\cdot\|$ and $\|\cdot\|''$, let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, and assume that either A or B is right invertible. Then,

$$\ell'(A)\ell(B) \leq \ell''(AB). \quad (9.5.19)$$

Furthermore, if $1 \leq p, q, r \leq \infty$, then

$$\ell_{r,q}(A)\ell_{q,p}(B) \leq \ell_{r,p}(AB). \quad (9.5.20)$$

In particular,

$$\sigma_m(A)\sigma_l(B) \leq \sigma_l(AB). \quad (9.5.21)$$

Proof. See [582] and [867, pp. 369, 370]. $\square$

9.6 Singular Value Inequalities

Proposition 9.6.1. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, for all $i \in \{1, \dots, \min\{n, m\}\}$ and $j \in \{1, \dots, \min\{m, l\}\}$ such that $i + j \leq \min\{n, l\} + 1$,

$$\sigma_{i+j-1}(AB) \leq \sigma_i(A)\sigma_j(B). \quad (9.6.1)$$

In particular, for all $i = 1, \dots, \min\{n, m, l\}$,

$$\sigma_i(AB) \leq \sigma_{\max}(A)\sigma_i(B) \quad (9.6.2)$$

and

$$\sigma_i(AB) \leq \sigma_i(A)\sigma_{\max}(B). \quad (9.6.3)$$

Proof. See [711, p. 178]. $\square$

Proposition 9.6.2. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. If $r \geq 0$, then, for all $k = 1, \dots, \min\{n, m, l\}$,

$$\sum_{i=1}^k \sigma_i^r(AB) \leq \sum_{i=1}^k \sigma_i^r(A) \sigma_i^r(B). \quad (9.6.4)$$

In particular, for all $k = 1, \dots, \min\{n, m, l\}$,

$$\sum_{i=1}^k \sigma_i(AB) \leq \sum_{i=1}^k \sigma_i(A) \sigma_i(B). \quad (9.6.5)$$

If $r < 0$, $n = m = l$, and A and B are nonsingular, then

$$\sum_{i=1}^n \sigma_i^r(AB) \leq \sum_{i=1}^n \sigma_i^r(A) \sigma_i^r(B). \quad (9.6.6)$$

Proof. The first statement follows from Proposition 9.6.3 and Fact 2.21.9. For the case $r < 0$, use Fact 2.21.12. See [197, p. 94] or [711, p. 177]. $\square$

Proposition 9.6.3. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then, for all $k = 1, \dots, \min\{n, m, l\}$,

$$\prod_{i=1}^k \sigma_i(AB) \leq \prod_{i=1}^k \sigma_i(A) \sigma_i(B).$$

If, in addition, $n = m = l$, then

$$\prod_{i=1}^n \sigma_i(AB) = \prod_{i=1}^n \sigma_i(A) \sigma_i(B).$$

Proof. See [711, p. 172]. $\square$

Proposition 9.6.4. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. If $m \leq n$, then, for all $i = 1, \dots, \min\{n, m, l\}$,

$$\sigma_{\min}(A) \sigma_i(B) = \sigma_m(A) \sigma_i(B) \leq \sigma_i(AB). \quad (9.6.7)$$

If $m \leq l$, then, for all $i = 1, \dots, \min\{n, m, l\}$,

$$\sigma_i(A) \sigma_{\min}(B) = \sigma_i(A) \sigma_m(B) \leq \sigma_i(AB). \quad (9.6.8)$$

Proof. Corollary 8.4.2 implies that $\sigma_m^2(A) I_m = \lambda_{\min}(A^* A) I_m \leq A^* A$, which implies that $\sigma_m^2(A) B^* B \leq B^* A^* A B$. Hence, it follows from the monotonicity theorem Theorem 8.4.9 that, for all $i = 1, \dots, \min\{n, m, l\}$,

$$\sigma_m(A) \sigma_i(B) = \lambda_i[\sigma_m^2(A) B^* B]^{1/2} \leq \lambda_i^{1/2}(B^* A^* A B) = \sigma_i(AB),$$

which proves the left-hand inequality in (9.6.7). Similarly, for all $i = 1, \dots, \min\{n, m, l\}$,

$$\sigma_i(A) \sigma_m(B) = \lambda_i[\sigma_m^2(B) A A^*]^{1/2} \leq \lambda_i^{1/2}(A B B^* A^*) = \sigma_i(AB). \quad \square$$

Corollary 9.6.5. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\sigma_m(A)\sigma_{\min\{n,m,l\}}(B) \leq \sigma_{\min\{n,m,l\}}(AB) \leq \sigma_{\max}(A)\sigma_{\min\{n,m,l\}}(B), \quad (9.6.9)$$

$$\sigma_m(A)\sigma_{\max}(B) \leq \sigma_{\max}(AB) \leq \sigma_{\max}(A)\sigma_{\max}(B), \quad (9.6.10)$$

$$\sigma_{\min\{n,m,l\}}(A)\sigma_m(B) \leq \sigma_{\min\{n,m,l\}}(AB) \leq \sigma_{\min\{n,m,l\}}(A)\sigma_{\max}(B), \quad (9.6.11)$$

$$\sigma_{\max}(A)\sigma_m(B) \leq \sigma_{\max}(AB) \leq \sigma_{\max}(A)\sigma_{\max}(B). \quad (9.6.12)$$

Specializing Corollary 9.6.5 to the case in which A or B is square yields the following result.

Corollary 9.6.6. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{n \times l}$. Then, for all $i = 1, \dots, \min\{n, l\}$,

$$\sigma_{\min}(A)\sigma_i(B) \leq \sigma_i(AB) \leq \sigma_{\max}(A)\sigma_i(B). \quad (9.6.13)$$

In particular,

$$\sigma_{\min}(A)\sigma_{\max}(B) \leq \sigma_{\max}(AB) \leq \sigma_{\max}(A)\sigma_{\max}(B). \quad (9.6.14)$$

If $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times m}$, then, for all $i = 1, \dots, \min\{n, m\}$,

$$\sigma_i(A)\sigma_{\min}(B) \leq \sigma_i(AB) \leq \sigma_i(A)\sigma_{\max}(B). \quad (9.6.15)$$

In particular,

$$\sigma_{\max}(A)\sigma_{\min}(B) \leq \sigma_{\max}(AB) \leq \sigma_{\max}(A)\sigma_{\max}(B). \quad (9.6.16)$$

Corollary 9.6.7. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. If $m \leq n$, then

$$\sigma_{\min}(A)\|B\|_F = \sigma_m(A)\|B\|_F \leq \|AB\|_F. \quad (9.6.17)$$

If $m \leq l$, then

$$\|A\|_F\sigma_{\min}(B) = \|A\|_F\sigma_m(B) \leq \|AB\|_F. \quad (9.6.18)$$

Proposition 9.6.8. Let $A, B \in \mathbb{F}^{n \times m}$. Then, for all $i, j \in \{1, \dots, \min\{n, m\}\}$ such that $i + j \leq \min\{n, m\} + 1$,

$$\sigma_{i+j-1}(A+B) \leq \sigma_i(A) + \sigma_j(B) \quad (9.6.19)$$

and

$$\sigma_{i+j-1}(A) - \sigma_j(B) \leq \sigma_i(A+B). \quad (9.6.20)$$

Proof. See [711, p. 178]. □

Corollary 9.6.9. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\sigma_n(A) - \sigma_{\max}(B) \leq \sigma_n(A+B) \leq \sigma_n(A) + \sigma_{\max}(B). \quad (9.6.21)$$

If, in addition, $n = m$, then

$$\sigma_{\min}(A) - \sigma_{\max}(B) \leq \sigma_{\min}(A+B) \leq \sigma_{\min}(A) + \sigma_{\max}(B). \quad (9.6.22)$$

Proof. The result follows from Proposition 9.6.8. Alternatively, it follows from Lemma 8.4.3 and the Cauchy-Schwarz inequality Corollary 9.1.7 that, for all

nonzero $x \in \mathbb{F}^n$,

$$\begin{aligned}\lambda_{\min}[(A+B)(A+B)^*] &\leq \frac{x^*(AA^* + BB^* + AB^* + BA^*)x}{x^*x} \\ &= \frac{x^*AA^*x}{\|x\|_2^2} + \frac{x^*BB^*x}{\|x\|_2^2} + \operatorname{Re} \frac{2x^*AB^*x}{\|x\|_2^2} \\ &\leq \frac{x^*AA^*x}{\|x\|_2^2} + \sigma_{\max}^2(B) + 2 \frac{(x^*AA^*x)^{1/2}}{\|x\|_2} \sigma_{\max}(B).\end{aligned}$$

Minimizing with respect to x and using Lemma 8.4.3 yields

$$\begin{aligned}\sigma_n^2(A+B) &= \lambda_{\min}[(A+B)(A+B)^*] \\ &\leq \lambda_{\min}(AA^*) + \sigma_{\max}^2(B) + 2\lambda_{\min}^{1/2}(AA^*)\sigma_{\max}(B) \\ &= [\sigma_n(A) + \sigma_{\max}(B)]^2,\end{aligned}$$

which proves the right-hand inequality of (9.6.21). Finally, the left-hand inequality follows from the right-hand inequality with A and B replaced by $A+B$ and $-B$, respectively. $\square$

9.7 Facts on Vector Norms

Fact 9.7.1. Let $x, y \in \mathbb{F}^n$. Then, x and y are linearly dependent if and only if $|x|^{\circ 2}$ and $|y|^{\circ 2}$ are linearly dependent and $|x^*y| = |x|^T|y|$. (Remark: This equivalence clarifies the relationship between (9.1.9) with $p = 2$ and Corollary 9.1.7.)

Fact 9.7.2. Let $x, y \in \mathbb{F}^n$, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then,

$$\left| \|x\| - \|y\| \right| \leq \begin{cases} \|x+y\| \\ \|x-y\|. \end{cases}$$

Fact 9.7.3. Let $x, y \in \mathbb{F}^n$, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then, the following statements hold:

- i) If there exists $\beta \geq 0$ such that either $x = \beta y$ or $y = \beta x$, then $\|x+y\| = \|x\| + \|y\|$.
- ii) If $\|x+y\| = \|x\| + \|y\|$ and x and y are linearly dependent, then there exists $\beta \geq 0$ such that either $x = \beta y$ or $y = \beta x$.
- iii) If $\|x+y\|_2 = \|x\|_2 + \|y\|_2$, then there exists $\beta \geq 0$ such that either $x = \beta y$ or $y = \beta x$.

(Proof: For *iii*), use *v*) of Fact 9.7.4.) (Problem: Consider *iii*) with alternative norms.) (Problem: If x and y are linearly independent, then does it follow that $\|x+y\| < \|x\| + \|y\|$?)

Fact 9.7.4. Let $x, y, z \in \mathbb{F}^n$. Then, the following statements hold:

- i) $\frac{1}{2}(\|x+y\|_2^2 + \|x-y\|_2^2) = \|x\|_2^2 + \|y\|_2^2$.

ii) If x and y are nonzero, then

$$\frac{1}{2}(\|x\|_2 + \|y\|_2) \left\| \frac{x}{\|x\|_2} - \frac{y}{\|y\|_2} \right\|_2 \leq \|x - y\|_2.$$

iii) If x and y are nonzero, then

$$\left\| \frac{1}{\|x\|_2} x - \|x\|_2 y \right\|_2 = \left\| \frac{1}{\|y\|_2} y - \|y\|_2 x \right\|_2.$$

iv) If $\mathbb{F} = \mathbb{R}$, then

$$4x^T y = \|x + y\|_2^2 - \|x - y\|_2^2.$$

v) If $\mathbb{F} = \mathbb{C}$, then

$$4x^* y = \|x + y\|_2^2 - \|x - y\|_2^2 + j(\|x + jy\|_2^2 - \|x - jy\|_2^2).$$

vi) $\operatorname{Re} x^* y = \frac{1}{4}(\|x + y\|_2^2 - \|x - y\|_2^2) = \frac{1}{2}(\|x + y\|_2^2 - \|x\|_2^2 - \|y\|_2^2).$

vii) If $\mathbb{F} = \mathbb{C}$, then $\operatorname{Im} x^* y = \frac{j}{4}(\|x + jy\|_2^2 - \|x - jy\|_2^2).$

viii) $\|x + y\|_2 = \sqrt{\|x\|_2^2 + \|y\|_2^2 + 2\operatorname{Re} x^* y}.$

ix) $\|x - y\|_2 = \sqrt{\|x\|_2^2 + \|y\|_2^2 - 2\operatorname{Re} x^* y}.$

x) $\|x + y\|_2 \|x - y\|_2 \leq \|x\|_2^2 + \|y\|_2^2.$

xi) If $\|x + y\|_2 = \|x\|_2 + \|y\|_2$, then $\operatorname{Im} x^* y = 0$ and $\operatorname{Re} x^* y \geq 0.$

xii) $|x^* y| \leq \|x\|_2 \|y\|_2.$

xiii) If $\|x + y\|_2 \leq 2$, then

$$(1 - \|x\|_2^2)(1 - \|y\|_2^2) \leq |1 - \operatorname{Re} x^* y|^2.$$

xiv) For all nonzero $\alpha \in \mathbb{R}$,

$$\|x\|_2^2 \|y\|_2^2 - |x^* y|^2 \leq \alpha^{-2} \|\alpha y - x\|_2^2 \|x\|_2^2.$$

xv) If $\operatorname{Re} x^* y \neq 0$, then, for all nonzero $\alpha \in \mathbb{R}$,

$$\|x\|_2^2 \|y\|_2^2 - |x^* y|^2 \leq \alpha_0^{-2} \|\alpha_0 y - x\|_2^2 \|x\|_2^2 \leq \alpha^{-2} \|\alpha y - x\|_2^2 \|x\|_2^2,$$

where $\alpha_0 \triangleq x^* x / (\operatorname{Re} x^* y).$

xvi) x, y, z satisfy

$$\|x + y\|_2^2 + \|y + z\|_2^2 + \|z + x\|_2^2 = \|x\|_2^2 + \|y\|_2^2 + \|z\|_2^2 + \|x + y + z\|_2^2$$

and

$$\|x + y\|_2 + \|y + z\|_2 + \|z + x\|_2 \leq \|x\|_2 + \|y\|_2 + \|z\|_2 + \|x + y + z\|_2.$$

xvii) $|x^* z z^* y - \frac{1}{2} x^* y \|z\|_2^2| \leq \frac{1}{2} \|x\|_2 \|y\|_2 \|z\|_2^2.$

xviii) $|\operatorname{Re}(x^* z z^* y - \frac{1}{2} x^* y \|z\|_2^2)| \leq \frac{1}{2} \|z\|_2^2 \sqrt{\|x\|_2^2 \|y\|_2^2 - (\operatorname{Im} x^* y)^2}.$

xix) $|\operatorname{Im}(x^* z z^* y - \frac{1}{2} x^* y \|z\|_2^2)| \leq \frac{1}{2} \|z\|_2^2 \sqrt{\|x\|_2^2 \|y\|_2^2 - (\operatorname{Re} x^* y)^2}.$

Furthermore, the following statements are equivalent:

xx) $\|x - y\|_2 = \|x + y\|_2.$

$$xix) \|x + y\|_2^2 = \|x\|_2^2 + \|y\|_2^2.$$

$$xii) \operatorname{Re} x^*y = 0.$$

Now, let $x_1, \dots, x_k \in \mathbb{F}^n$, and assume that $x_i^*x_j = \delta_{ij}$ for all $i, j = 1, \dots, n$. Then, the following statement holds:

$$xiii) \sum_{i=1}^k |y^*x_i|^2 \leq \|y\|_2^2.$$

If, in addition, $k = n$, then the following statement holds:

$$xiv) \sum_{i=1}^n |y^*x_i|^2 = \|y\|_2^2.$$

(Remark: *i*) is the *parallelogram law*, which relates the diagonals and the sides of a parallelogram; *ii*) is the *Dunkl-Williams inequality*, which compares the distance between x and y with the distance between the projections of x and y onto the unit sphere (see [446], [1010, p. 515], and [1490, p. 28]); *iv*) and *v*) are the *polarization identity* (see [368, p. 54], [1030, p. 276], and Fact 1.18.2); *ix*) is the *cosine law* (see Fact 9.9.13 for a matrix version); *xiii*) is given in [1467] and implies Aczel's inequality given by Fact 1.16.19; *xv*) is given in [913]; *xvi*) is *Hlawka's identity* and *Hlawka's inequality* (see Fact 1.8.6, Fact 1.18.2, [1010, p. 521], and [1039, p. 100]); *xvii*) is *Buzano's inequality* (see [514] and Fact 1.17.2); *xviii*) and *xix*) are given in [1093]; the equivalence of *xi*) and *xii*) is the *Pythagorean theorem*; *xiii*) is *Bessel's inequality*; and *xiv*) is *Parseval's identity*. Note that *xiv*) implies *xiii*).) (Remark: Hlawka's inequality is called the *quadrilateral inequality* in [1202], which gives a geometric interpretation. In addition, [1202] provides an extension and geometric interpretation to the *polygonal inequalities*. See Fact 9.7.7.) (Remark: When $\mathbb{F} = \mathbb{R}$ and $n = 2$ the Euclidean norm of $\begin{bmatrix} x \\ y \end{bmatrix}$ is equivalent to the absolute value $|z| = |x + jy|$. See Fact 1.18.2.)

Fact 9.7.5. Let $x, y \in \mathbb{R}^3$, and let $\mathcal{S} \subset \mathbb{R}^3$ be the parallelogram with vertices $0, x, y$, and $x + y$. Then,

$$\operatorname{area}(\mathcal{S}) = \|x \times y\|_2.$$

(Remark: See Fact 2.20.13, Fact 2.20.14 and Fact 3.10.1.) (Remark: The parallelogram associated with the cross product can be interpreted as a bivector. See [605, 870] and [426, pp. 86–88].)

Fact 9.7.6. Let $x, y \in \mathbb{R}^n$, and assume that x and y are nonzero. Then,

$$\frac{x^T y}{\|x\|_2 \|y\|_2} (\|x\|_2 + \|y\|_2) \leq \|x + y\|_2 \leq \|x\|_2 + \|y\|_2.$$

Hence, if $x^T y = \|x\|_2 \|y\|_2$, then $\|x\|_2 + \|y\|_2 = \|x + y\|_2$. (Proof: See [1010, p. 517].) (Remark: This result is a *reverse triangle inequality*.) (Problem: Extend this result to complex vectors.)

Fact 9.7.7. Let $x_1, \dots, x_n \in \mathbb{F}^n$, and let $\alpha_1, \dots, \alpha_n$ be nonnegative numbers. Then,

$$\sum_{i=1}^n \alpha_i \left\| x_i - \sum_{j=1}^n \alpha_j x_j \right\|_2 \leq \sum_{i=1}^n \alpha_i \|x_i\|_2 + \left[\left(\sum_{i=1}^n \alpha_i \right) - 2 \right] \left\| \sum_{i=1}^n \alpha_i x_i \right\|_2.$$

In particular,

$$\sum_{i=1}^n \left\| \sum_{j=1, j \neq i}^n x_j \right\|_2 \leq \sum_{i=1}^n \|x_i\|_2 + (n-2) \left\| \sum_{i=1}^n x_i \right\|_2.$$

(Remark: The first inequality is the *generalized Hlawka inequality* or *polygonal inequalities*. The second inequality is the *Djokovic inequality*. See [1254] and Fact 9.7.4.)

Fact 9.7.8. Let $x, y \in \mathbb{R}^n$, let α and δ , be positive numbers, and let $p, q \in (0, \infty)$ satisfy $1/p + 1/q = 1$. Then,

$$\left(\frac{\alpha}{\alpha + \|y\|_2^q} \right)^{p-1} \delta^p \leq |\delta - x^T y|^p + \alpha^{p-1} \|x\|_2^p.$$

Equality holds if and only if $x = [\delta \|y\|_2^{q-2} / (\alpha + \|y\|_2^q)] y$. In particular,

$$\frac{\alpha \delta^2}{\alpha + \|y\|_2^2} \leq (\delta - x^T y)^2 + \alpha \|x\|_2^2.$$

Equality holds if and only if $x = [\delta / (\alpha + \|y\|_2^2)] y$. (Proof: See [1253].) (Remark: The first inequality is due to Pecaric. The case $p = q = 2$ is due to Dragomir and Yang. These results are generalizations of Hua's inequality. See Fact 1.15.13 and Fact 9.7.9.)

Fact 9.7.9. Let $x_1, \dots, x_n, y \in \mathbb{R}^n$, and let α and δ be positive numbers. Then,

$$\frac{\alpha}{\alpha + n} \|y\|_2^2 \leq \left\| y - \sum_{i=1}^n x_i \right\|_2^2 + \alpha \sum_{i=1}^n \|x_i\|_2^2.$$

Equality holds if and only if $x_1 = \dots = x_n = [1/(\alpha + n)] y$. (Proof: See [1253].) (Remark: This inequality, which is due to Dragomir and Yang, is a generalization of Hua's inequality. See Fact 1.15.13 and Fact 9.7.8.)

Fact 9.7.10. Let $x, y \in \mathbb{F}^n$, and assume that x and y are nonzero. Then,

$$\begin{aligned} \frac{\|x - y\|_2 - |\|x\|_2 - \|y\|_2|}{\min\{\|x\|_2, \|y\|_2\}} &\leq \left\| \frac{x}{\|x\|_2} - \frac{y}{\|y\|_2} \right\|_2 \\ &\leq \left\{ \begin{array}{l} \frac{\|x - y\|_2 + |\|x\|_2 - \|y\|_2|}{\max\{\|x\|_2, \|y\|_2\}} \\ \frac{2\|x - y\|_2}{\|x\|_2 + \|y\|_2} \end{array} \right\} \\ &\leq \left\{ \begin{array}{l} \frac{2\|x - y\|_2}{\max\{\|x\|_2, \|y\|_2\}} \\ \frac{2(\|x - y\|_2 + |\|x\|_2 - \|y\|_2|)}{\|x\|_2 + \|y\|_2} \end{array} \right\} \\ &\leq \frac{4\|x - y\|_2}{\|x\|_2 + \|y\|_2}. \end{aligned}$$

(Proof: See Fact 9.7.13 and [991].) (Remark: In the last string of inequalities, the first inequality is the *reverse Malignanda inequality*, the second and upper third terms constitute the *Malignanda inequality*, the second and lower third terms constitute the Dunkl-Williams inequality in an inner product space, the second and upper fourth terms constitute the *Massera-Schaffer inequality*.) (Remark: See Fact 1.18.5.)

Fact 9.7.11. Let $x, y \in \mathbb{F}^n$, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then, there exists a unique number $\alpha \in [1, 2]$ such that, for all $x, y \in \mathbb{F}^n$, at least one of which is nonzero,

$$\frac{2}{\alpha} \leq \frac{\|x+y\|^2 + \|x-y\|^2}{\|x\|^2 + \|y\|^2} \leq 2\alpha.$$

Furthermore, if $\|\cdot\| = \|\cdot\|_p$, then

$$\alpha = \begin{cases} 2^{(2-p)/p}, & 1 \leq p \leq 2, \\ 2^{(p-2)/p}, & p \geq 2. \end{cases}$$

(Proof: See [275, p. 258].) (Remark: This result is the *von Neumann-Jordan inequality*.) (Remark: When $p = 2$, it follows that $\alpha = 2$, and this result yields *i*) of Fact 9.7.4.)

Fact 9.7.12. Let $x, y \in \mathbb{F}^n$, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then,

$$\|x+y\| \leq \|x\| + \|y\| - \min\{\|x\|, \|y\|\} \left(2 - \left\| \frac{x}{\|x\|} + \frac{y}{\|y\|} \right\| \right) \leq \|x\| + \|y\|,$$

$$\|x-y\| \leq \|x\| + \|y\| - \min\{\|x\|, \|y\|\} \left(2 - \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \right) \leq \|x\| + \|y\|,$$

$$\|x\| + \|y\| - \max\{\|x\|, \|y\|\} \left(2 - \left\| \frac{x}{\|x\|} + \frac{y}{\|y\|} \right\| \right) \leq \|x+y\| \leq \|x\| + \|y\|,$$

and

$$\|x\| + \|y\| - \max\{\|x\|, \|y\|\} \left(2 - \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \right) \leq \|x-y\| \leq \|x\| + \|y\|.$$

(Proof: See [951].)

Fact 9.7.13. Let $x, y \in \mathbb{F}^n$, assume that x and y are nonzero, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then,

$$\begin{aligned} \frac{(\|x\| + \|y\|)(\|x+y\| - \|\|x\| - \|y\|\|)}{4 \min\{\|x\|, \|y\|\}} &\leq \frac{1}{4}(\|x\| + \|y\|) \left\| \frac{x}{\|x\|} + \frac{y}{\|y\|} \right\| \\ &\leq \frac{1}{2} \max\{\|x\|, \|y\|\} \left\| \frac{x}{\|x\|} + \frac{y}{\|y\|} \right\| \\ &\leq \frac{1}{2}(\|x+y\| + \max\{\|x\|, \|y\|\} - \|x\| - \|y\|) \\ &\leq \frac{1}{2}(\|x+y\| + \|\|x\| - \|y\|\|) \\ &\leq \|x+y\| \end{aligned}$$

and

$$\begin{aligned}
\frac{(\|x\| + \|y\|)(\|x - y\| - |\|x\| - \|y\||)}{4 \min\{\|x\|, \|y\|\}} &\leq \frac{1}{4}(\|x\| + \|y\|) \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \\
&\leq \frac{1}{2} \max\{\|x\|, \|y\|\} \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \\
&\leq \frac{1}{2}(\|x - y\| + \max\{\|x\|, \|y\|\} - \|x\| - \|y\|) \\
&\leq \frac{1}{2}(\|x - y\| + |\|x\| - \|y\||) \\
&\leq \|x - y\|.
\end{aligned}$$

Furthermore,

$$\begin{aligned}
\frac{\|x - y\| - |\|x\| - \|y\||}{\min\{\|x\|, \|y\|\}} &\leq \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\| \\
&\leq \frac{\|x - y\| + |\|x\| - \|y\||}{\max\{\|x\|, \|y\|\}} \\
&\leq \left\{ \begin{array}{c} \frac{2\|x - y\|}{\max\{\|x\|, \|y\|\}} \\ \frac{2(\|x - y\| + |\|x\| - \|y\||)}{\|x\| + \|y\|} \end{array} \right\} \\
&\leq \frac{4\|x - y\|}{\|x\| + \|y\|}.
\end{aligned}$$

(Proof: The result follows from Fact 9.7.12, [951, 991] and [1010, p. 516].) (Remark: In the last string of inequalities, the first inequality is the *reverse Maligranda inequality*, the second inequality is the *Maligranda inequality*, the second and upper fourth terms constitute the *Massera-Schaffer inequality*, and the second and fifth terms constitute the *Dunkl-Williams inequality*. See Fact 1.18.2 and Fact 9.7.4 for the case of the Euclidean norm.) (Remark: Extensions to more than two vectors are given in [794, 1078].)

Fact 9.7.14. Let $x, y \in \mathbb{F}^n$, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then,

$$\begin{aligned}
&\left. \begin{array}{l} \|x\|^2 + \|y\|^2 \\ 2\|x\|^2 - 4\|x\|\|y\| + 2\|y\|^2 \end{array} \right\} \leq \|x + y\|^2 + \|x - y\|^2 \\
&\leq 2\|x\|^2 + 4\|x\|\|y\| + 2\|y\|^2 \\
&\leq 4(\|x\|^2 + \|y\|^2).
\end{aligned}$$

(Proof: See [530, pp. 9, 10] and [1030, p. 278].)

Fact 9.7.15. Let $x, y \in \mathbb{F}^n$, let $\alpha \in [0, 1]$, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then,

$$\|x + y\| \leq \|\alpha x + (1 - \alpha)y\| + \|(1 - \alpha)x + \alpha y\| \leq \|x\| + \|y\|.$$

Fact 9.7.16. Let $x, y \in \mathbb{F}^n$, assume that x and y are nonzero, let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, and let $p \in \mathbb{R}$. Then, the following statements hold:

i) If $p \leq 0$, then

$$\| \|x\|^{p-1}x - \|y\|^{p-1}y \| \leq (2-p) \frac{\max\{\|x\|^p, \|y\|^p\}}{\max\{\|x\|, \|y\|\}} \|x - y\|.$$

ii) If $p \in [0, 1]$, then

$$\| \|x\|^{p-1}x - \|y\|^{p-1}y \| \leq (2-p) \frac{\|x - y\|}{[\max\{\|x\|, \|y\|\}]^{1-p}}.$$

iii) If $p \geq 1$, then

$$\| \|x\|^{p-1}x - \|y\|^{p-1}y \| \leq p[\max\{\|x\|, \|y\|\}]^{p-1} \|x - y\|.$$

(Proof: See [951].)

Fact 9.7.17. Let $x, y \in \mathbb{F}^n$, let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, assume that $\|x\| \neq \|y\|$, and let $p > 0$. Then,

$$\| \|x\| - \|y\| \| \leq \frac{\| \|x\|^p x - \|y\|^p y \|}{\| \|x\|^{p+1} - \|y\|^{p+1} \|} \| \|x\| - \|y\| \| \leq \|x - y\|.$$

(Proof: See [1010, p. 516].)

Fact 9.7.18. Let $x \in \mathbb{F}^n$, and let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$. Then,

$$\|x\|_2 \leq \sqrt{\|x\|_p \|x\|_q}.$$

Fact 9.7.19. Let $x, y \in \mathbb{F}^n$, let $p \in (0, 1]$, and define $\|\cdot\|_p$ as in (9.1.1). Then,

$$\|x\|_p + \|y\|_p \leq \|x + y\|_p.$$

(Remark: This result is a *reverse triangle inequality*.)

Fact 9.7.20. Let $x, y \in \mathbb{F}^n$, let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, let p and q be real numbers, and assume that $1 \leq p \leq q$. Then,

$$[\frac{1}{2}(\|x + \frac{1}{\sqrt{q-1}}y\|^q + \|x - \frac{1}{\sqrt{q-1}}y\|^q)]^{1/q} \leq [\frac{1}{2}(\|x + \frac{1}{\sqrt{p-1}}y\|^p + \|x - \frac{1}{\sqrt{p-1}}y\|^p)]^{1/p}.$$

(Proof: See [542, p. 207].) (Remark: This result is *Bonami's inequality*. See Fact 1.10.16.)

Fact 9.7.21. Let $x, y \in \mathbb{F}^{n \times n}$. If $p \in [1, 2]$, then

$$(\|x\|_p + \|y\|_p)^p + \| \|x\|_p - \|y\|_p \|^p \leq \|x + y\|_p^p + \|x - y\|_p^p$$

and

$$(\|x + y\|_p + \|x - y\|_p)^p + \| \|x + y\|_p - \|x - y\|_p \|^p \leq 2^p (\|x\|_p^p + \|y\|_p^p).$$

If $p \in [2, \infty]$, then

$$\|x + y\|_p^p + \|x - y\|_p^p \leq (\|x\|_p + \|y\|_{\sigma p})^p + \| \|x\|_p - \|y\|_p \|^p$$

and

$$2^p (\|x\|_p^p + \|y\|_p^p) \leq (\|x + y\|_p + \|x - y\|_p)^p + \| \|x + y\|_p - \|x - y\|_p \|^p.$$

(Proof: See [116, 906].) (Remark: These inequalities are versions of *Hanner's inequality*. These vector versions follow from inequalities on L_p by appropriate choice of measure.) (Remark: Matrix versions are given in Fact 9.9.36.)

Fact 9.7.22. Let $y \in \mathbb{F}^n$, let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, let $\|\cdot\|'$ be the norm on $\mathbb{F}^{n \times n}$ induced by $\|\cdot\|$, and define

$$\|y\|_D \triangleq \max_{x \in \{z \in \mathbb{F}^n: \|z\|=1\}} |y^*x|.$$

Then, $\|\cdot\|_D$ is a norm on $\mathbb{F}^n$. Furthermore,

$$\|y\| = \max_{x \in \{z \in \mathbb{F}^n: \|z\|_D=1\}} |y^*x|.$$

Hence, for all $x \in \mathbb{F}^n$,

$$|x^*y| \leq \|x\| \|y\|_D.$$

In addition,

$$\|xy^*\|' = \|x\| \|y\|_D.$$

Finally, let $p \in [1, \infty]$, and let $1/p + 1/q = 1$. Then,

$$\|\cdot\|_{pD} = \|\cdot\|_q.$$

Hence, for all $x \in \mathbb{F}^n$,

$$|x^*y| \leq \|x\|_p \|y\|_q$$

and

$$\|xy^*\|_{p,p} = \|x\|_p \|y\|_q.$$

(Proof: See [1230, p. 57].) (Remark: $\|\cdot\|_D$ is the *dual norm* of $\|\cdot\|$.)

Fact 9.7.23. Let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, and let $\alpha > 0$. Then, $f: \mathbb{F}^n \mapsto [0, \infty)$ defined by $f(x) = \|x\|$ is convex. Furthermore, $\{x \in \mathbb{F}^n: \|x\| \leq \alpha\}$ is symmetric, solid, convex, closed, and bounded. (Remark: See Fact 10.8.22.)

Fact 9.7.24. Let $x \in \mathbb{R}^n$, and let $\|\cdot\|$ be a norm on $\mathbb{R}^n$. Then, $x^T y > 0$ for all $y \in \mathbb{B}_{\|\cdot\|}(x) = \{z \in \mathbb{R}^n: \|z - x\| < \|x\|\}$.

Fact 9.7.25. Let $x, y \in \mathbb{R}^n$, assume that x and y are nonzero, assume that $x^T y = 0$, and let $\|\cdot\|$ be a norm on $\mathbb{R}^n$. Then, $\|x\| \leq \|x+y\|$. (Proof: If $\|x+y\| < \|x\|$, then $x+y \in \mathbb{B}_{\|\cdot\|}(0)$, and thus $y \in \mathbb{B}_{\|\cdot\|}(-x)$. By Fact 9.7.24, $x^T y < 0$.) (Remark: See [218, 901] for related results concerning matrices.)

Fact 9.7.26. Let $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$. Then,

$$\sigma_{\max}(xy^*) = \|xy^*\|_F = \|x\|_2 \|y\|_2$$

and

$$\sigma_{\max}(xx^*) = \|xx^*\|_F = \|x\|_2^2.$$

(Remark: See Fact 5.11.16.)

Fact 9.7.27. Let $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$. Then,

$$\|x \otimes y\|_2 = \|\text{vec}(x \otimes y^T)\|_2 = \|\text{vec}(yx^T)\|_2 = \|yx^T\|_2 = \|x\|_2 \|y\|_2.$$

Fact 9.7.28. Let $x \in \mathbb{F}^n$, and let $1 \leq p, q \leq \infty$. Then,

$$\|x\|_p = \|x\|_{p,q}.$$

Fact 9.7.29. Let $x \in \mathbb{F}^n$, and let $p, q \in [1, \infty)$, where $p \leq q$. Then,

$$\|x\|_q \leq \|x\|_p \leq n^{1/p-1/q} \|x\|_q.$$

(Proof: See [680], [681, p. 107].) (Remark: See Fact 1.15.5 and Fact 9.8.21.)

Fact 9.7.30. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$\|x\|_A \triangleq (x^*Ax)^{1/2}$$

is a norm on $\mathbb{F}^n$.

Fact 9.7.31. Let $\|\cdot\|$ and $\|\cdot\|'$ be norms on $\mathbb{F}^n$, and let $\alpha, \beta > 0$. Then, $\alpha\|\cdot\| + \beta\|\cdot\|'$ is also a norm on $\mathbb{F}^n$. Furthermore, $\max\{\|\cdot\|, \|\cdot\|'\}$ is a norm on $\mathbb{F}^n$. (Remark: $\min\{\|\cdot\|, \|\cdot\|'\}$ is not necessarily a norm.)

Fact 9.7.32. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then, $\|x\|' \triangleq \|Ax\|$ is a norm on $\mathbb{F}^n$.

Fact 9.7.33. Let $x \in \mathbb{F}^n$, and let $p \in [1, \infty]$. Then,

$$\|\overline{x}\|_p = \|x\|_p.$$

Fact 9.7.34. Let $x_1, \dots, x_k \in \mathbb{F}^n$, let $\alpha_1, \dots, \alpha_k$ be positive numbers, and assume that $\sum_{i=1}^k \alpha_i = 1$. Then,

$$|1_{1 \times n}(x_1 \circ \dots \circ x_k)| \leq \prod_{i=1}^k \|x_i\|_{1/\alpha_i}.$$

(Remark: This result is the *generalized Hölder inequality*. See [273, p. 128].)

9.8 Facts on Matrix Norms for One Matrix

Fact 9.8.1. Let $\mathcal{S} \subseteq \mathbb{F}^m$, assume that $\mathcal{S}$ is bounded, and let $A \in \mathbb{F}^{n \times m}$. Then, $A\mathcal{S}$ is bounded.

Fact 9.8.2. Let $A \in \mathbb{F}^{n \times n}$, assume that A is a idempotent, and assume that, for all $x \in \mathbb{F}^n$,

$$\|Ax\|_2 \leq \|x\|_2.$$

Then, A is a projector. (Proof: See [536, p. 42].)

Fact 9.8.3. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are projectors. Then, the following statements are equivalent:

i) $A \leq B$.

ii) For all $x \in \mathbb{F}^n$, $\|Ax\|_2 \leq \|Bx\|_2$.

- iii) $\mathcal{R}(A) \subseteq \mathcal{R}(A)$.
- iv) $AB = A$.
- v) $BA = A$.
- vi) $B - A$ is a projector.

(Proof: See [536, p. 43] and [1184, p. 24].) (Remark: See Fact 3.13.14 and Fact 3.13.17.)

Fact 9.8.4. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{sprad}(A) < 1$. Then, there exists a submultiplicative matrix norm $\|\cdot\|$ on $\mathbb{F}^{n \times n}$ such that $\|A\| < 1$. Furthermore,

$$\lim_{k \rightarrow \infty} A^k = 0.$$

Fact 9.8.5. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, and let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A^{-1}\| \geq \|I_n\|/\|A\|.$$

Fact 9.8.6. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonzero and idempotent, and let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A\| \geq 1.$$

Fact 9.8.7. Let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then, $\|\cdot\|$ is self-adjoint.

Fact 9.8.8. Let $A \in \mathbb{F}^{n \times m}$, let $\|\cdot\|$ be a norm on $\mathbb{F}^{n \times m}$, and define $\|A\|' \triangleq \|A^*\|$. Then, $\|\cdot\|'$ is a norm on $\mathbb{F}^{m \times n}$. If, in addition, $n = m$ and $\|\cdot\|$ is induced by $\|\cdot\|''$, then $\|\cdot\|'$ is induced by $\|\cdot\|''_D$. (Proof: See [709, p. 309] and Fact 9.8.10.) (Remark: See Fact 9.7.22 for the definition of the dual norm. $\|\cdot\|'$ is the *adjoint norm* of $\|\cdot\|$.) (Problem: Generalize this result to nonsquare matrices and norms that are not equi-induced.)

Fact 9.8.9. Let $1 \leq p \leq \infty$. Then, $\|\cdot\|_{\sigma_p}$ is unitarily invariant.

Fact 9.8.10. Let $A \in \mathbb{F}^{n \times m}$, and let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$. Then,

$$\|A^*\|_{p,p} = \|A\|_{q,q}.$$

In particular,

$$\|A^*\|_{\text{col}} = \|A\|_{\text{row}}.$$

(Proof: See Fact 9.8.8.)

Fact 9.8.11. Let $A \in \mathbb{F}^{n \times m}$, and let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$. Then,

$$\left\| \begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix} \right\|_{p,p} = \max\{\|A\|_{p,p}, \|A\|_{q,q}\}.$$

In particular,

$$\left\| \begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix} \right\|_{\text{col}} = \left\| \begin{bmatrix} 0 & A \\ A^* & 0 \end{bmatrix} \right\|_{\text{row}} = \max\{\|A\|_{\text{col}}, \|A\|_{\text{row}}\}.$$

Fact 9.8.12. Let $A \in \mathbb{F}^{n \times m}$. Then, the following inequalities hold:

- i) $\|A\|_{\text{F}} \leq \|A\|_1 \leq \sqrt{mn}\|A\|_{\text{F}}.$
- ii) $\|A\|_{\infty} \leq \|A\|_1 \leq mn\|A\|_{\infty}.$
- iii) $\|A\|_{\text{col}} \leq \|A\|_1 \leq m\|A\|_{\text{col}}.$
- iv) $\|A\|_{\text{row}} \leq \|A\|_1 \leq n\|A\|_{\text{row}}.$
- v) $\sigma_{\max}(A) \leq \|A\|_1 \leq \sqrt{mn \text{rank } A} \sigma_{\max}(A).$
- vi) $\|A\|_{\infty} \leq \|A\|_{\text{F}} \leq \sqrt{mn}\|A\|_{\infty}.$
- vii) $\frac{1}{\sqrt{n}}\|A\|_{\text{col}} \leq \|A\|_{\text{F}} \leq \sqrt{m}\|A\|_{\text{col}}.$
- viii) $\frac{1}{\sqrt{m}}\|A\|_{\text{row}} \leq \|A\|_{\text{F}} \leq \sqrt{n}\|A\|_{\text{row}}.$
- ix) $\sigma_{\max}(A) \leq \|A\|_{\text{F}} \leq \sqrt{\text{rank } A} \sigma_{\max}(A).$
- x) $\frac{1}{n}\|A\|_{\text{col}} \leq \|A\|_{\infty} \leq \|A\|_{\text{col}}.$
- xi) $\frac{1}{m}\|A\|_{\text{row}} \leq \|A\|_{\infty} \leq \|A\|_{\text{row}}.$
- xii) $\frac{1}{\sqrt{mn}}\sigma_{\max}(A) \leq \|A\|_{\infty} \leq \sigma_{\max}(A).$
- xiii) $\frac{1}{m}\|A\|_{\text{row}} \leq \|A\|_{\text{col}} \leq n\|A\|_{\text{row}}.$
- xiv) $\frac{1}{\sqrt{m}}\sigma_{\max}(A) \leq \|A\|_{\text{col}} \leq \sqrt{n}\sigma_{\max}(A).$
- xv) $\frac{1}{\sqrt{n}}\sigma_{\max}(A) \leq \|A\|_{\text{row}} \leq \sqrt{m}\sigma_{\max}(A).$

(Proof: See [709, p. 314] and [1501].) (Remark: See [681, p. 115] for matrices that attain these bounds.)

Fact 9.8.13. Let $A \in \mathbb{F}^{n \times m}$, and assume that A is normal. Then,

$$\frac{1}{\sqrt{mn}}\sigma_{\max}(A) \leq \|A\|_{\infty} \leq \text{sprad}(A) = \sigma_{\max}(A).$$

(Proof: Use Fact 5.14.15 and statement *xii*) of Fact 9.8.12.)

Fact 9.8.14. Let $A \in \mathbb{R}^{n \times n}$, assume that A is symmetric, and assume that every diagonal entry of A is zero. Then, the following conditions are equivalent:

- i) For all $x \in \mathbb{R}^n$ such that $1_{1 \times n}x = 0$, it follows that $x^T A x \leq 0$.
- ii) There exists a positive integer k and vectors $x_1, \dots, x_n \in \mathbb{R}^k$ such that, for all $i, j = 1, \dots, n$, $A_{(i,j)} = \|x_i - x_j\|_2^2$.

(Proof: See [18].) (Remark: This result is due to Schoenberg.) (Remark: A is a *Euclidean distance matrix*.)

Fact 9.8.15. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\|A^A\|_F \leq n^{(2-n)/2} \|A\|_F^{n-1}.$$

(Proof: See [1098, pp. 151, 165].)

Fact 9.8.16. Let $A \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ and $\|\cdot\|'$ be norms on $\mathbb{F}^n$, and define the induced norms

$$\|A\|'' \triangleq \max_{x \in \{y \in \mathbb{F}^m: \|y\|=1\}} \|Ax\|$$

and

$$\|A\|''' \triangleq \max_{x \in \{y \in \mathbb{F}^m: \|y\|'=1\}} \|Ax\|'.$$

Then,

$$\begin{aligned} \max_{A \in \{X \in \mathbb{F}^{n \times n}: X \neq 0\}} \frac{\|A\|''}{\|A\|'''} &= \max_{A \in \{X \in \mathbb{F}^{n \times n}: X \neq 0\}} \frac{\|A\|'''}{\|A\|''} \\ &= \max_{x \in \{y \in \mathbb{F}^n: y \neq 0\}} \frac{\|x\|}{\|x\|'} \max_{x \in \{y \in \mathbb{F}^n: y \neq 0\}} \frac{\|x\|'}{\|x\|}. \end{aligned}$$

(Proof: See [709, p. 303].) (Remark: This symmetry property is evident in Fact 9.8.12.)

Fact 9.8.17. Let $A \in \mathbb{F}^{n \times m}$, let $q, r \in [1, \infty]$, assume that $1 \leq q \leq r$, define

$$p \triangleq \frac{1}{\frac{1}{q} - \frac{1}{r}},$$

and assume that $p \geq 2$. Then,

$$\|A\|_p \leq \|A\|_{q,r}.$$

In particular,

$$\|A\|_\infty \leq \|A\|_{\infty, \infty}.$$

(Proof: See [476].) (Remark: This result is due to Hardy and Littlewood.)

Fact 9.8.18. Let $A \in \mathbb{R}^{n \times m}$. Then,

$$\begin{aligned} \left\| \begin{bmatrix} \|\text{row}_1(A)\|_2 \\ \vdots \\ \|\text{row}_n(A)\|_2 \end{bmatrix} \right\|_1 &\leq \sqrt{2} \|A\|_{1, \infty}, \\ \left\| \begin{bmatrix} \|\text{row}_1(A)\|_1 \\ \vdots \\ \|\text{row}_n(A)\|_1 \end{bmatrix} \right\|_2 &\leq \sqrt{2} \|A\|_{1, \infty}, \end{aligned}$$

$$\|A\|_{4/3}^{3/4} \leq \sqrt{2} \|A\|_{1, \infty}.$$

(Proof: See [542, p. 303].) (Remark: The first and third results are due to Littlewood, while the second result is due to Orlicz.)

Fact 9.8.19. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive semidefinite. Then,

$$\|A\|_{1,\infty} = \max_{x \in \{z \in \mathbb{F}^n: \|z\|_\infty=1\}} x^*Ax.$$

(Remark: This result is due to Tao. See [681, p. 116] and [1138].)

Fact 9.8.20. Let $A \in \mathbb{F}^{n \times n}$. If $p \in [1, 2]$, then

$$\|A\|_F \leq \|A\|_{\sigma p} \leq n^{1/p-1/2} \|A\|_F.$$

If $p \in [2, \infty]$, then

$$\|A\|_{\sigma p} \leq \|A\|_F \leq n^{1/2-1/p} \|A\|_{\sigma p}.$$

(Proof: See [200, p. 174].)

Fact 9.8.21. Let $A \in \mathbb{F}^{n \times n}$, and let $p, q \in [1, \infty]$. Then,

$$\|A\|_{p,p} \leq \begin{cases} n^{1/p-1/q} \|A\|_{q,q}, & p \leq q, \\ n^{1/q-1/p} \|A\|_{q,q}, & q \leq p. \end{cases}$$

Consequently,

$$n^{1/p-1} \|A\|_{\text{col}} \leq \|A\|_{p,p} \leq n^{1-1/p} \|A\|_{\text{col}},$$

$$n^{-|1/p-1/2|} \sigma_{\max}(A) \leq \|A\|_{p,p} \leq n^{|1/p-1/2|} \sigma_{\max}(A),$$

$$n^{-1/p} \|A\|_{\text{col}} \leq \|A\|_{p,p} \leq n^{1/p} \|A\|_{\text{row}}.$$

(Proof: See [680] and [681, p. 112].) (Remark: See Fact 9.7.29.) (Problem: Extend these inequalities to nonsquare matrices.)

Fact 9.8.22. Let $A \in \mathbb{F}^{n \times m}$, $p, q \in [1, \infty]$, and $\alpha \in [0, 1]$, and let $r \triangleq pq/[(1-\alpha)p + \alpha q]$. Then,

$$\|A\|_{r,r} \leq \|A\|_{p,p}^\alpha \|A\|_{q,q}^{1-\alpha}.$$

(Proof: See [680] or [681, p. 113].)

Fact 9.8.23. Let $A \in \mathbb{F}^{n \times m}$, and let $p \in [1, \infty]$. Then,

$$\|A\|_{p,p} \leq \|A\|_{\text{col}}^{1/p} \|A\|_{\text{row}}^{1-1/p}.$$

In particular,

$$\sigma_{\max}(A) \leq \sqrt{\|A\|_{\text{col}} \|A\|_{\text{row}}}.$$

(Proof: Set $\alpha = 1/p$, $p = 1$, and $q = \infty$ in Fact 9.8.22. See [681, p. 113]. To prove the special case $p = 2$ directly, note that $\lambda_{\max}(A^*A) \leq \|A^*A\|_{\text{col}} \leq \|A^*\|_{\text{col}} \|A\|_{\text{col}} = \|A\|_{\text{row}} \|A\|_{\text{col}}$.)

Fact 9.8.24. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\left. \begin{array}{l} \|A\|_{2,1} \\ \|A\|_{\infty,2} \end{array} \right\} \leq \sigma_{\max}(A).$$

(Proof: The result follows from Proposition 9.1.5.)

Fact 9.8.25. Let $A \in \mathbb{F}^{n \times m}$, and let $p \in [1, 2]$. Then,

$$\|A\|_{p,p} \leq \|A\|_{\text{col}}^{2/p-1} \sigma_{\max}^{2-2/p}(A).$$

(Proof: Let $\alpha = 2/p - 1$, $p = 1$, and $q = 2$ in Fact 9.8.22. See [681, p. 113].)

Fact 9.8.26. Let $A \in \mathbb{F}^{n \times n}$, and let $p \in [1, \infty]$. Then,

$$\|A\|_{p,p} \leq \| |A| \|_{p,p} \leq n^{\min\{1/p, 1-1/p\}} \|A\|_{p,p} \leq \sqrt{n} \|A\|_{p,p}.$$

(Remark: See [681, p. 117].)

Fact 9.8.27. Let $A \in \mathbb{F}^{n \times m}$, and let $p, q \in [1, \infty]$. Then,

$$\|\bar{A}\|_{q,p} = \|A\|_{q,p}.$$

Fact 9.8.28. Let $A \in \mathbb{F}^{n \times m}$, and let $p, q \in [1, \infty]$. Then,

$$\|A^*\|_{q,p} = \|A\|_{p/(p-1), q/(q-1)}.$$

Fact 9.8.29. Let $A \in \mathbb{F}^{n \times m}$, and let $p, q \in [1, \infty]$. Then,

$$\|A\|_{q,p} \leq \begin{cases} \|A\|_{p/(p-1)}, & 1/p + 1/q \leq 1, \\ \|A\|_q, & 1/p + 1/q \geq 1. \end{cases}$$

Fact 9.8.30. Let $A \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|\langle A \rangle\| = \|A\|.$$

Fact 9.8.31. Let $A, S \in \mathbb{F}^{n \times n}$, assume that S is nonsingular, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A\| \leq \frac{1}{2} \|SAS^{-1} + S^{-*}AS^*\|.$$

(Proof: See [61, 246].)

Fact 9.8.32. Let $A \in \mathbb{F}^{n \times n}$, assume that A is positive semidefinite, and let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A\|^{1/2} \leq \|A^{1/2}\|.$$

In particular,

$$\sigma_{\max}^{1/2}(A) = \sigma_{\max}(A^{1/2}).$$

Fact 9.8.33. Let $A_{11} \in \mathbb{F}^{n \times n}$, $A_{12} \in \mathbb{F}^{n \times m}$, and $A_{22} \in \mathbb{F}^{m \times m}$, assume that $\begin{bmatrix} A_{11} & A_{12} \\ A_{12}^* & A_{22} \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ is positive semidefinite, let $\|\cdot\|$ and $\|\cdot\|'$ be unitarily invariant norms on $\mathbb{F}^{n \times n}$ and $\mathbb{F}^{m \times m}$, respectively, and let $p > 0$. Then,

$$\|\langle A_{12} \rangle^p\|^2 \leq \|A_{11}^p\| \|A_{22}^p\|'.$$

(Proof: See [713].)

Fact 9.8.34. Let $A \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, let $\|\cdot\|_{\mathbb{D}}$ denote the dual norm on $\mathbb{F}^n$, and let $\|\cdot\|'$ denote the norm induced by $\|\cdot\|$ on $\mathbb{F}^{n \times n}$. Then,

$$\|A\|' = \max_{\substack{x, y \in \mathbb{F}^n \\ x, y \neq 0}} \frac{\operatorname{Re} y^* A x}{\|y\|_{\mathbb{D}} \|x\|}.$$

(Proof: See [681, p. 115].) (Remark: See Fact 9.7.22 for the definition of the dual norm.) (Problem: Generalize this result to obtain Fact 9.8.35 as a special case.)

Fact 9.8.35. Let $A \in \mathbb{F}^{n \times m}$, and let $p, q \in [1, \infty]$. Then,

$$\|A\|_{q,p} = \max_{\substack{x \in \mathbb{F}^m, y \in \mathbb{F}^n \\ x, y \neq 0}} \frac{|y^* A x|}{\|y\|_{q/(q-1)} \|x\|_p}.$$

Fact 9.8.36. Let $A \in \mathbb{F}^{n \times m}$, and let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$. Then,

$$\|A\|_{p,p} = \max_{\substack{x \in \mathbb{F}^m, y \in \mathbb{F}^n \\ x, y \neq 0}} \frac{|y^* A x|}{\|y\|_q \|x\|_p} = \max_{\substack{x \in \mathbb{F}^m, y \in \mathbb{F}^n \\ x, y \neq 0}} \frac{|y^* A x|}{\|y\|_{p/(p-1)} \|x\|_p}.$$

(Remark: See Fact 9.13.2 for the case $p = 2$.)

Fact 9.8.37. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is positive definite. Then,

$$\min_{x \in \mathbb{F}^n \setminus \{0\}} \frac{x^* A x}{\|A x\|_2 \|x\|_2} = \frac{2\sqrt{\alpha\beta}}{\alpha + \beta}$$

and

$$\min_{\alpha \geq 0} \sigma_{\max}(\alpha A - I) = \frac{\alpha - \beta}{\alpha + \beta},$$

where $\alpha \triangleq \lambda_{\max}(A)$ and $\beta \triangleq \lambda_{\min}(A)$. (Proof: See [609].) (Remark: These quantities are *antieigenvalues*.)

Fact 9.8.38. Let $A \in \mathbb{F}^{n \times n}$, and define

$$\operatorname{nrad}(A) \triangleq \max \{|x^* A x| : x \in \mathbb{C}^n \text{ and } x^* x \leq 1\}.$$

Then, the following statements hold:

- i) $\operatorname{nrad}(A) = \max\{|z| : z \in \Theta(A)\}.$
- ii) $\operatorname{sprad}(A) \leq \operatorname{nrad}(A) \leq \operatorname{nrad}(|A|) = \frac{1}{2} \operatorname{sprad}(|A| + |A|^T).$
- iii) $\frac{1}{2} \sigma_{\max}(A) \leq \operatorname{nrad}(A) \leq \frac{1}{2} [\sigma_{\max}(A) + \sigma_{\max}^{1/2}(A^2)] \leq \sigma_{\max}(A).$
- iv) If $A^2 = 0$, then $\operatorname{nrad}(A) = \sigma_{\max}(A).$
- v) If $\operatorname{nrad}(A) = \sigma_{\max}(A)$, then $\sigma_{\max}(A^2) = \sigma_{\max}^2(A).$
- vi) If A is normal, then $\operatorname{nrad}(A) = \operatorname{sprad}(A).$
- vii) $\operatorname{nrad}(A^k) \leq [\operatorname{nrad}(A)]^k$ for all $k \in \mathbb{N}.$
- viii) $\operatorname{nrad}(\cdot)$ is a weakly unitarily invariant norm on $\mathbb{F}^{n \times n}.$
- ix) $\operatorname{nrad}(\cdot)$ is not a submultiplicative norm on $\mathbb{F}^{n \times n}.$

- $x) \|\cdot\| \triangleq \alpha \text{nrad}(\cdot)$ is a submultiplicative norm on $\mathbb{F}^{n \times n}$ if and only if $\alpha \geq 4$.
- $xi) \text{nrad}(AB) \leq \text{nrad}(A)\text{nrad}(B)$ for all $A, B \in \mathbb{F}^{n \times n}$ such that A and B are normal.
- $xii) \text{nrad}(A \circ B) \leq \alpha \text{nrad}(A)\text{nrad}(B)$ for all $A, B \in \mathbb{F}^{n \times n}$ if and only if $\alpha \geq 2$.
- $xiii) \text{nrad}(A \oplus B) = \max\{\text{nrad}(A), \text{nrad}(B)\}$ for all $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$.

(Proof: See [709, p. 331] and [711, pp. 43, 44]. For *iii*), see [823].) (Remark: $\text{nrad}(A)$ is the *numerical radius* of A . $\Theta(A)$ is the numerical range. See Fact 8.14.7.) (Remark: $\text{nrad}(\cdot)$ is not submultiplicative. The example $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 0 & 2 \\ 2 & 0 \end{bmatrix}$, where B is normal, $\text{nrad}(A) = 1/2$, $\text{nrad}(B) = 2$, and $\text{nrad}(AB) = 2$, shows that *xi*) is not valid if only one of the matrices A and B is normal, which corrects [711, pp. 43, 73].) (Remark: *vii*) is the *power inequality*.)

Fact 9.8.39. Let $A \in \mathbb{F}^{n \times m}$, let $\gamma > \sigma_{\max}(A)$, and define $\beta \triangleq \sigma_{\max}(A)/\gamma$. Then,

$$\|A\|_F \leq \sqrt{-[\gamma^2/(2\pi)] \log \det(I - \gamma^{-2}A^*A)} \leq \beta^{-1} \sqrt{-\log(1 - \beta^2)} \|A\|_F.$$

(Proof: See [254].)

Fact 9.8.40. Let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then, $\|A\| = 1$ for all $A \in \mathbb{F}^{n \times n}$ such that $\text{rank } A = 1$ if and only if $\|E_{1,1}\| = 1$. (Proof: $\|A\| = \|E_{1,1}\| \sigma_{\max}(A)$.) (Remark: These equivalent normalizations are used in [1230, p. 74] and [197], respectively.)

Fact 9.8.41. Let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- $i) \sigma_{\max}(A) \leq \|A\|$ for all $A \in \mathbb{F}^{n \times n}$.
- $ii) \|\cdot\|$ is submultiplicative.
- $iii) \|A^2\| \leq \|A\|^2$ for all $A \in \mathbb{F}^{n \times n}$.
- $iv) \|A^k\| \leq \|A\|^k$ for all $k \geq 1$ and $A \in \mathbb{F}^{n \times n}$.
- $v) \|A \circ B\| \leq \|A\| \|B\|$ for all $A, B \in \mathbb{F}^{n \times n}$.
- $vi) \text{sprad}(A) \leq \|A\|$ for all $A \in \mathbb{F}^{n \times n}$.
- $vii) \|Ax\|_2 \leq \|A\| \|x\|_2$ for all $A \in \mathbb{F}^{n \times n}$ and $x \in \mathbb{F}^n$.
- $viii) \|A\|_{\infty} \leq \|A\|$ for all $A \in \mathbb{F}^{n \times n}$.
- $ix) \|E_{1,1}\| \geq 1$.
- $x) \sigma_{\max}(A) \leq \|A\|$ for all $A \in \mathbb{F}^{n \times n}$ such that $\text{rank } A = 1$.

(Proof: The equivalence of *i*)–*vii*) is given in [710] and [711, p. 211]. Since $\|A\| = \|E_{1,1}\| \sigma_{\max}(A)$ for all $A \in \mathbb{F}^{n \times n}$ such that $\text{rank } A = 1$, it follows that *vii*) and *viii*) are equivalent. To prove *ix*) $\implies$ *x*), let $A \in \mathbb{F}^{n \times n}$ satisfy $\text{rank } A = 1$. Then, $\|A\| = \sigma_{\max}(A) \|E_{1,1}\| \geq \sigma_{\max}(A)$. To show *x*) $\implies$ *ii*), define $\|\cdot\|' \triangleq \|E_{1,1}\|^{-1} \|\cdot\|$. Since $\|E_{1,1}\|' = 1$, it follows from [197, p. 94] that $\|\cdot\|'$ is submultiplicative. Since $\|E_{1,1}\|^{-1} \leq 1$, it follows that $\|\cdot\|$ is also submultiplicative. Alternatively,

$\|A\|' = \sigma_{\max}(A)$ for all $A \in \mathbb{F}^{n \times n}$ having rank 1. Then, Corollary 3.10 of [1230, p. 80] implies that $\|\cdot\|'$, and thus $\|\cdot\|$, is submultiplicative.)

Fact 9.8.42. Let $\Phi: \mathbb{F}^n \mapsto [0, \infty)$ satisfy the following conditions:

- i) If $x \neq 0$, then $\Phi(x) > 0$.
- ii) $\Phi(\alpha x) = |\alpha|\Phi(x)$ for all $\alpha \in \mathbb{R}$.
- iii) $\Phi(x + y) \leq \Phi(x) + \Phi(y)$ for all $x, y \in \mathbb{F}^n$.
- iv) If $A \in \mathbb{F}^{n \times n}$ is a permutation matrix, then $\Phi(Ax) = \Phi(x)$ for all $x \in \mathbb{F}^n$.
- v) $\Phi(|x|) = \Phi(x)$ for all $x \in \mathbb{F}^n$.

Furthermore, for $A \in \mathbb{F}^{n \times m}$, where $n \leq m$, define

$$\|A\| \triangleq \Phi[\sigma_1(A), \dots, \sigma_n(A)].$$

Then, $\|\cdot\|$ is a unitarily invariant norm on $\mathbb{F}^{n \times m}$. Conversely, if $\|\cdot\|$ is a unitarily invariant norm on $\mathbb{F}^{n \times m}$, where $n \leq m$, then $\Phi: \mathbb{F}^n \mapsto [0, \infty)$ defined by

$$\Phi(x) \triangleq \left\| \begin{bmatrix} x_{(1)} & \cdots & 0 & 0_{n \times (m-n)} \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & x_{(n)} & 0_{n \times (m-n)} \end{bmatrix} \right\|$$

satisfies i)–v). (Proof: See [1230, pp. 75, 76].) (Remark: Φ is a *symmetric gauge function*. This result is due to von Neumann. See Fact 2.21.14.)

Fact 9.8.43. Let $\|\cdot\|$ and $\|\cdot\|'$ denote norms on $\mathbb{F}^m$ and $\mathbb{F}^n$, respectively, and define $\hat{\ell}: \mathbb{F}^{n \times m} \mapsto \mathbb{R}$ by

$$\hat{\ell}(A) \triangleq \min_{x \in \mathbb{F}^m \setminus \{0\}} \frac{\|Ax\|'}{\|x\|},$$

or, equivalently,

$$\hat{\ell}(A) \triangleq \min_{x \in \{y \in \mathbb{F}^m: \|y\|=1\}} \|Ax\|'.$$

Then, for $A \in \mathbb{F}^{n \times m}$, the following statements hold:

- i) $\hat{\ell}(A) \geq 0$.
- ii) $\hat{\ell}(A) > 0$ if and only if $\text{rank } A = m$.
- iii) $\hat{\ell}(A) = \ell(A)$ if and only if either $A = 0$ or $\text{rank } A = m$.

(Proof: See [867, pp. 369, 370].) (Remark: $\hat{\ell}$ is a weaker version of ℓ .)

Fact 9.8.44. Let $A \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ be a normalized, submultiplicative norm on $\mathbb{F}^{n \times n}$, and assume that $\|I - A\| < 1$. Then, A is nonsingular. (Remark: See Fact 9.9.56.)

Fact 9.8.45. Let $\|\cdot\|$ be a normalized, submultiplicative norm on $\mathbb{F}^{n \times n}$. Then, $\|\cdot\|$ is equi-induced if and only if $\|A\| \leq \|A\|'$ for all $A \in \mathbb{F}^{n \times n}$ and for all normalized submultiplicative norms $\|\cdot\|'$ on $\mathbb{F}^{n \times n}$. (Proof: See [1234].) (Remark: As shown in [308, 383], not every normalized submultiplicative norm on $\mathbb{F}^{n \times n}$ is equi-induced or induced.)

9.9 Facts on Matrix Norms for Two or More Matrices

Fact 9.9.1. $\|\cdot\|'_\infty \triangleq n\|\cdot\|_\infty$ is submultiplicative on $\mathbb{F}^{n \times n}$. (Remark: It is not necessarily true that $\|AB\|_\infty \leq \|A\|_\infty\|B\|_\infty$. For example, let $A = B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$.)

Fact 9.9.2. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$. Then,

$$\|AB\|_\infty \leq m\|A\|_\infty\|B\|_\infty.$$

Furthermore, if $A = 1_{n \times m}$ and $B = 1_{m \times l}$, then $\|AB\|_\infty = m\|A\|_\infty\|B\|_\infty$.

Fact 9.9.3. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$. Then, $\|AB\| \leq \|A\|\|B\|$. Hence, if $\|A\| \leq 1$ and $\|B\| \leq 1$, then $\|AB\| \leq 1$. Finally, if either $\|A\| < 1$ or $\|B\| < 1$, then $\|AB\| < 1$. (Remark: $\text{sprad}(A) < 1$ and $\text{sprad}(B) < 1$ do not imply that $\text{sprad}(AB) < 1$. Let $A = B^T = \begin{bmatrix} 0 & 2 \\ 0 & 0 \end{bmatrix}$.)

Fact 9.9.4. Let $\|\cdot\|$ be a norm on $\mathbb{F}^{m \times m}$, and let

$$\delta > \sup \left\{ \frac{\|AB\|}{\|A\|\|B\|} : A, B \in \mathbb{F}^{m \times m}, A, B \neq 0 \right\}.$$

Then, $\|\cdot\|' \triangleq \delta\|\cdot\|$ is a submultiplicative norm on $\mathbb{F}^{m \times m}$. (Proof: See [709, p. 323].)

Fact 9.9.5. Let $A, B \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and assume that $A \leq B$. Then,

$$\|A\| \leq \|B\|.$$

(Proof: See [215].)

Fact 9.9.6. Let $A, B \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$, and assume that AB is normal. Then,

$$\|AB\| \leq \|BA\|.$$

(Proof: See [197, p. 253].)

Fact 9.9.7. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite and nonzero, and let $\|\cdot\|$ be a submultiplicative unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\frac{\|AB\|}{\|A\|\|B\|} \leq \frac{\|A+B\|}{\|A\| + \|B\|}$$

and

$$\frac{\|A \circ B\|}{\|A\|\|B\|} \leq \frac{\|A+B\|}{\|A\| + \|B\|}.$$

(Proof: See [675].) (Remark: See Fact 9.8.41.)

Fact 9.9.8. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$. Then, $\|\cdot\|' \triangleq 2\|\cdot\|$ is a submultiplicative norm on $\mathbb{F}^{n \times n}$ and satisfies

$$\|[A, B]\|' \leq \|A\|'\|B\|'.$$

Fact 9.9.9. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) There exist projectors $Q, P \in \mathbb{R}^{n \times n}$ such that $A = [P, Q]$.
- ii) $\sigma_{\max}(A) \leq 1/2$, A and $-A$ are unitarily similar, and A is skew Hermitian.

(Proof: See [903].) (Remark: Extensions are discussed in [984].) (Remark: See Fact 3.12.16 for the case of idempotent matrices.) (Remark: In the case $\mathbb{F} = \mathbb{R}$, the condition that A is skew symmetric implies that A and $-A$ are orthogonally similar. See Fact 5.9.10.)

Fact 9.9.10. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|AB\| \leq \sigma_{\max}(A)\|B\|$$

and

$$\|AB\| \leq \|A\|\sigma_{\max}(B).$$

Consequently, if $C \in \mathbb{F}^{n \times n}$, then

$$\|ABC\| \leq \sigma_{\max}(A)\|B\|\sigma_{\max}(C).$$

(Proof: See [820].)

Fact 9.9.11. Let $A, B \in \mathbb{F}^{n \times m}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{m \times m}$. If $p > 0$, then

$$\|\langle A^*B \rangle^p\|^2 \leq \|(A^*A)^p\| \|(B^*B)^p\|.$$

In particular,

$$\|(A^*BB^*A)^{1/4}\|^2 \leq \|\langle A \rangle\| \|\langle B \rangle\|$$

and

$$\|\langle A^*B \rangle\| = \|A^*B\|^2 \leq \|A^*A\| \|B^*B\|.$$

Furthermore,

$$\text{tr} \langle A^*B \rangle \leq \|A\|_{\text{F}} \|B\|_{\text{F}}$$

and

$$\left[\text{tr} (A^*BB^*A)^{1/4} \right]^2 \leq (\text{tr} \langle A \rangle)(\text{tr} \langle B \rangle).$$

(Proof: See [713] and use Fact 9.8.30.) (Problem: Noting Fact 9.12.1 and Fact 9.12.2, compare the lower bounds for $\|A\|_{\text{F}}\|B\|_{\text{F}}$ given by

$$\left. \begin{array}{l} \text{tr} \langle A^*B \rangle \\ |\text{tr} A^*B| \\ \sqrt{|\text{tr} (A^*B)^2|} \end{array} \right\} \leq \|A\|_{\text{F}} \|B\|_{\text{F}} \quad \left. \begin{array}{l} \\ \\ \sqrt{\text{tr} AA^*BB^*} \end{array} \right\}$$

Fact 9.9.12. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\begin{aligned}
(2\|A\|_F\|B\|_F)^{1/2} &\leq (\|A\|_F^2 + \|B\|_F^2)^{1/2} \\
&= \|(A^2 + B^2)^{1/2}\|_F \\
&\leq \|A + B\|_F \\
&\leq \sqrt{2}(\|A\|_F^2 + \|B\|_F^2)^{1/2}.
\end{aligned}$$

Fact 9.9.13. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\|A + B\|_F = \sqrt{\|A\|_F^2 + \|B\|_F^2 + 2\operatorname{tr} AB^*} \leq \|A\|_F + \|B\|_F.$$

In particular,

$$\|A - B\|_F = \sqrt{\|A\|_F^2 + \|B\|_F^2 - 2\operatorname{tr} AB^*}.$$

If, in addition, A is Hermitian and B is skew Hermitian, then $\operatorname{tr} AB^* = 0$, and thus

$$\|A + B\|_F^2 = \|A - B\|_F^2 = \|A\|_F^2 + \|B\|_F^2.$$

(Remark: The second identity is a matrix version of the cosine law given by *ix*) of Fact 9.7.4.)

Fact 9.9.14. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|AB\| \leq \frac{1}{4}\|(\langle A \rangle + \langle B^* \rangle)^2\|.$$

(Proof: See [212].)

Fact 9.9.15. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|AB\| \leq \frac{1}{4}\|(A + B)^2\|.$$

(Proof: See [212] or [1485, p. 77].) (Problem: Noting Fact 9.9.12, compare the lower bounds for $\|A + B\|_F$ given by

$$(2\|A\|_F\|B\|_F)^{1/2} \leq \|(A^2 + B^2)^{1/2}\|_F \leq \|A + B\|_F$$

and

$$2\|AB\|_F^{1/2} \leq \|(A + B)^2\|_F^{1/2} \leq \|A + B\|_F.)$$

Fact 9.9.16. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$, and let $p \in (0, \infty)$. If $p \in [0, 1]$, then

$$\|A^p B^p\| \leq \|AB\|^p.$$

If $p \in [1, \infty)$, then

$$\|AB\|^p \leq \|A^p B^p\|.$$

(Proof: See [203, 523].) (Remark: See Fact 8.18.26.)

Fact 9.9.17. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. If $p \in [0, 1]$, then

$$\|B^p A^p B^p\| \leq \|(BAB)^p\|.$$

Furthermore, if $p \geq 1$, then

$$\|(BAB)^p\| \leq \|B^p A^p B^p\|.$$

(Proof: See [69] and [197, p. 258].) (Remark: Extensions and a reverse inequality are given in Fact 8.10.49.) (Remark: See Fact 8.12.20 and Fact 8.18.26.)

Fact 9.9.18. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A^{1/2}B^{1/2}\| \leq \frac{1}{2}\|A + B\|.$$

Hence,

$$\|AB\| \leq \frac{1}{2}\|A^2 + B^2\|,$$

and thus

$$\|(A + B)^2\| \leq 2\|A^2 + B^2\|.$$

Consequently,

$$\|AB\| \leq \frac{1}{4}\|(A + B)^2\| \leq \frac{1}{2}\|A^2 + B^2\|.$$

(Proof: Let $p = 1/2$ and $X = I$ in Fact 9.9.49. The last inequality follows from Fact 9.9.15.) (Remark: See Fact 8.18.13.)

Fact 9.9.19. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let either $p = 1$ or $p \in [2, \infty]$. Then,

$$\|\langle AB \rangle^{1/2}\|_{\sigma p} \leq \frac{1}{2}\|A + B\|_{\sigma p}.$$

(Proof: See [90, 212].) (Remark: The inequality holds for all Q-norms. See [197].) (Remark: See Fact 8.18.13.)

Fact 9.9.20. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times l}$, and $p, q, q', r \in [1, \infty]$, and assume that $1/q + 1/q' = 1$. Then,

$$\|AB\|_p \leq \varepsilon_{pq}(n)\varepsilon_{pr}(l)\varepsilon_{q'r}(m)\|A\|_q\|B\|_r,$$

where

$$\varepsilon_{pq}(n) \triangleq \begin{cases} 1, & p \geq q, \\ n^{1/p-1/q}, & q \geq p. \end{cases}$$

Furthermore, there exist matrices $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$ such that equality holds. (Proof: See [564].) (Remark: Related results are given in [475, 476, 564, 565, 566, 828, 1313].)

Fact 9.9.21. Let $A, B \in \mathbb{C}^{n \times m}$. Then, there exist unitary matrices $S_1, S_2 \in \mathbb{C}^{m \times m}$ such that

$$\langle A + B \rangle \leq S_1 \langle A \rangle S_1^* + S_2 \langle B \rangle S_2^*.$$

(Remark: This result is a matrix version of the triangle inequality. See [47, 1271].)

Fact 9.9.22. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $p \in [1, \infty]$. Then,

$$\|A - B\|_{\sigma 2p}^2 \leq \|A^2 - B^2\|_{\sigma p}.$$

(Proof: See [813].) (Remark: The case $p = 1$ is due to Powers and Stormer.)

Fact 9.9.23. Let $A, B \in \mathbb{F}^{n \times n}$, and let $p \in [1, \infty]$. Then,

$$\|\langle A \rangle - \langle B \rangle\|_{\sigma p}^2 \leq \|A + B\|_{\sigma 2p} \|A - B\|_{\sigma 2p}.$$

(Proof: See [827].)

Fact 9.9.24. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\|\langle A \rangle - \langle B \rangle\|_{\sigma 1}^2 \leq 2\|A + B\|_{\sigma 1} \|A - B\|_{\sigma 1}.$$

(Proof: See [827].) (Remark: This result is due to Borchers and Kosaki. See [827].)

Fact 9.9.25. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\|\langle A \rangle - \langle B \rangle\|_F \leq \sqrt{2}\|A - B\|_F$$

and

$$\|\langle A \rangle - \langle B \rangle\|_F^2 + \|\langle A^* \rangle - \langle B^* \rangle\|_F^2 \leq 2\|A - B\|_F^2.$$

If, in addition, A and B are normal, then

$$\|\langle A \rangle - \langle B \rangle\|_F \leq \|A - B\|_F.$$

(Proof: See [47, 70, 812, 827] and [683, pp. 217, 218].)

Fact 9.9.26. Let $A, B \in \mathbb{R}^{n \times n}$. Then,

$$\|AB - BA\|_F \leq \sqrt{2}\|A\|_F \|B\|_F.$$

(Proof: See [242, 1385].) (Remark: The constant $\sqrt{2}$ holds for all n .) (Remark: Extensions to complex matrices are given in [243].)

Fact 9.9.27. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$\|AB - BA\|_F^2 + \|(A - B)^2\|_F^2 \leq \|A^2 - B^2\|_F^2.$$

(Proof: See [820].)

Fact 9.9.28. Let $A, B \in \mathbb{F}^{n \times n}$, let p be a positive number, and assume that either A is normal and $p \in [2, \infty]$, or A is Hermitian and $p \geq 1$. Then,

$$\|\langle A \rangle B - B \langle A \rangle\|_{\sigma p} \leq \|AB - BA\|_{\sigma p}.$$

(Proof: See [1].)

Fact 9.9.29. Let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$, and let $A, X, B \in \mathbb{F}^{n \times n}$. Then,

$$\|AX - XB\| \leq [\sigma_{\max}(A) + \sigma_{\max}(B)]\|X\|.$$

In particular,

$$\sigma_{\max}(AX - XA) \leq 2\sigma_{\max}(A)\sigma_{\max}(X).$$

Now, assume that A and B are positive semidefinite. Then,

$$\|AX - XB\| \leq \max\{\sigma_{\max}(A), \sigma_{\max}(B)\}\|X\|.$$

In particular,

$$\sigma_{\max}(AX - XA) \leq \sigma_{\max}(A)\sigma_{\max}(X).$$

Finally, assume that A and X are positive semidefinite. Then,

$$\|AX - XA\| \leq \frac{1}{2}\sigma_{\max}(A) \left\| \begin{bmatrix} X & 0 \\ 0 & X \end{bmatrix} \right\|.$$

In particular,

$$\sigma_{\max}(AX - XA) \leq \frac{1}{2}\sigma_{\max}(A)\sigma_{\max}(X).$$

(Proof: See [214].) (Remark: The first inequality is sharp since equality holds for $A = B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ and $X = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$.) (Remark: $\|\cdot\|$ can be extended to $\mathbb{F}^{2n \times 2n}$ by considering the n largest singular values of matrices in $\mathbb{F}^{2n \times 2n}$. For details, see [197, pp. 90, 98].)

Fact 9.9.30. Let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$, let $A, X \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian. Then,

$$\|AX - XA\| \leq [\lambda_{\max}(A) - \lambda_{\min}(A)]\|X\|.$$

(Proof: See [214].) (Remark: $\lambda_{\max}(A) - \lambda_{\min}(A)$ is the spread of A . See Fact 8.15.31 and Fact 9.9.31.)

Fact 9.9.31. Let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$, let $A, X \in \mathbb{F}^{n \times n}$, assume that A is normal, let $\text{spec}(A) = \{\lambda_1, \dots, \lambda_r\}$, and define

$$\text{spd}(A) \triangleq \max\{|\lambda_i(A) - \lambda_j(A)| : i, j = 1, \dots, r\}.$$

Then,

$$\|AX - XA\| \leq \sqrt{2}\text{spd}(A)\|X\|.$$

Furthermore, let $p \in [1, \infty]$. Then,

$$\|AX - XA\|_{\sigma p} \leq 2^{|2-p|/(2p)}\text{spd}(A)\|X\|_{\sigma p}.$$

In particular,

$$\|AX - XA\|_{\text{F}} \leq \text{spd}(A)\|X\|_{\text{F}}$$

and

$$\sigma_{\max}(AX - XA) \leq \sqrt{2}\text{spd}(A)\sigma_{\max}(X).$$

(Proof: See [214].) (Remark: $\text{spd}(A)$ is the spread of A . See Fact 8.15.31 and Fact 9.9.30.)

Fact 9.9.32. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\sigma_{\max}(\langle A \rangle - \langle B \rangle) \leq \frac{2}{\pi} \left[2 + \log \frac{\sigma_{\max}(A) + \sigma_{\max}(B)}{\sigma_{\max}(A - B)} \right] \sigma_{\max}(A - B).$$

(Remark: This result is due to Kato. See [827].)

Fact 9.9.33. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times l}$, and let $r = 1$ or $r = 2$. Then,

$$\|AB\|_{\sigma r} = \|A\|_{\sigma 2r} \|B\|_{\sigma 2r}$$

if and only if there exists $\alpha \geq 0$ such that $AA^* = \alpha B^*B$. Furthermore,

$$\|AB\|_{\infty} = \|A\|_{\infty} \|B\|_{\infty}$$

if and only if AA^* and B^*B have a common eigenvector associated with $\lambda_1(AA^*)$ and $\lambda_1(B^*B)$. (Proof: See [1442].)

Fact 9.9.34. Let $A, B \in \mathbb{F}^{n \times n}$. If $p \in (0, 2]$, then

$$2^{p-1}(\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p) \leq \|A + B\|_{\sigma p}^p + \|A - B\|_{\sigma p}^p \leq 2(\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p).$$

If $p \in [2, \infty)$, then

$$2(\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p) \leq \|A + B\|_{\sigma p}^p + \|A - B\|_{\sigma p}^p \leq 2^{p-1}(\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p).$$

If $p \in (1, 2]$ and $1/p + 1/q = 1$, then

$$\|A + B\|_{\sigma p}^q + \|A - B\|_{\sigma p}^q \leq 2(\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p)^{q/p}.$$

If $p \in [2, \infty)$ and $1/p + 1/q = 1$, then

$$2(\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p)^{q/p} \leq \|A + B\|_{\sigma p}^q + \|A - B\|_{\sigma p}^q.$$

(Proof: See [696].) (Remark: These inequalities are versions of the *Clarkson inequalities*. See Fact 1.18.2.) (Remark: See [696] for extensions to unitarily invariant norms. See [213] for additional extensions.)

Fact 9.9.35. Let $A, B \in \mathbb{C}^{n \times m}$. If $p \in [1, 2]$, then

$$[\|A\|^2 + (p-1)\|B\|^2]^{1/2} \leq [\tfrac{1}{2}(\|A + B\|^p + \|A - B\|^p)]^{1/p}.$$

If $p \in [2, \infty]$, then

$$[\tfrac{1}{2}(\|A + B\|^p + \|A - B\|^p)]^{1/p} \leq [\|A\|^2 + (p-1)\|B\|^2]^{1/2}.$$

(Proof: See [116, 164].) (Remark: This result is *Beckner's two-point inequality* or *optimal 2-uniform convexity*.)

Fact 9.9.36. Let $A, B \in \mathbb{F}^{n \times n}$. If either $p \in [1, 4/3]$ or both $p \in (4/3, 2]$ and $A + B$ and $A - B$ are positive semidefinite, then

$$(\|A\|_{\sigma p} + \|B\|_{\sigma p})^p + \||A\|_{\sigma p} - \|B\|_{\sigma p}\|^p \leq \|A + B\|_{\sigma p}^p + \|A - B\|_{\sigma p}^p.$$

Furthermore, if either $p \in [4, \infty]$ or both $p \in [2, 4)$ and A and B are positive semidefinite, then

$$\|A + B\|_{\sigma p}^p + \|A - B\|_{\sigma p}^p \leq (\|A\|_{\sigma p} + \|B\|_{\sigma p})^p + \||A\|_{\sigma p} - \|B\|_{\sigma p}\|^p.$$

(Proof: See [116, 811].) (Remark: These inequalities are versions of *Hanner's inequality*.) (Remark: Vector versions are given in Fact 9.7.21.)

Fact 9.9.37. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that A and B are Hermitian. If $p \in [1, 2]$, then

$$2^{1/2-1/p} \|(A^2 + B^2)^{1/2}\|_p \leq \|A + jB\|_{\sigma p} \leq \|(A^2 + B^2)^{1/2}\|_p$$

and

$$2^{1-2/p}(\|A\|_{\sigma p}^2 + \|B\|_{\sigma p}^2) \leq \|A + jB\|_{\sigma p}^2 \leq 2^{2/p-1}(\|A\|_{\sigma p}^2 + \|B\|_{\sigma p}^2).$$

Furthermore, if $p \in [2, \infty)$, then

$$\|(A^2 + B^2)^{1/2}\|_p \leq \|A + jB\|_{\sigma p} \leq 2^{1/2-1/p} \|(A^2 + B^2)^{1/2}\|_p$$

and

$$2^{2/p-1}(\|A\|_{\sigma p}^2 + \|B\|_{\sigma p}^2) \leq \|A + jB\|_{\sigma p}^2 \leq 2^{1-2/p}(\|A\|_{\sigma p}^2 + \|B\|_{\sigma p}^2).$$

(Proof: See [211].)

Fact 9.9.38. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that A and B are Hermitian. If $p \in [1, 2]$, then

$$2^{1-2/p}(\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p) \leq \|A + jB\|_{\sigma p}^p.$$

If $p \in [2, \infty]$, then

$$\|A + jB\|_{\sigma p}^p \leq 2^{1-2/p}(\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p).$$

In particular,

$$\|A + jB\|_{\mathbb{F}}^2 = \|A\|_{\mathbb{F}}^2 + \|B\|_{\mathbb{F}}^2 = \|(A^2 + B^2)^{1/2}\|_{\mathbb{F}}^2.$$

(Proof: See [211, 219].)

Fact 9.9.39. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that A is positive semidefinite and B is Hermitian. If $p \in [1, 2]$, then

$$\|A\|_{\sigma p}^2 + 2^{1-2/p}\|B\|_{\sigma p}^2 \leq \|A + jB\|_{\sigma p}^2.$$

If $p \in [2, \infty]$, then

$$\|A + jB\|_{\sigma p}^2 \leq \|A\|_{\sigma p}^2 + 2^{1-2/p}\|B\|_{\sigma p}^2.$$

In particular,

$$\|A\|_{\sigma 1}^2 + \frac{1}{2}\|B\|_{\sigma 1}^2 \leq \|A + jB\|_{\sigma 1}^2,$$

$$\|A + jB\|_{\mathbb{F}}^2 = \|A\|_{\mathbb{F}}^2 + \|B\|_{\mathbb{F}}^2,$$

and

$$\sigma_{\max}^2(A + jB) \leq \sigma_{\max}^2(A) + 2\sigma_{\max}^2(B).$$

In fact,

$$\|A\|_{\sigma 1}^2 + \|B\|_{\sigma 1}^2 \leq \|A + jB\|_{\sigma 1}^2.$$

(Proof: See [219].)

Fact 9.9.40. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that A and B are positive semidefinite. If $p \in [1, 2]$, then

$$\|A\|_{\sigma p}^2 + \|B\|_{\sigma p}^2 \leq \|A + jB\|_{\sigma p}^2.$$

If $p \in [2, \infty]$, then

$$\|A + jB\|_{\sigma p}^2 \leq \|A\|_{\sigma p}^2 + \|B\|_{\sigma p}^2.$$

Hence,

$$\|A\|_{\sigma 2}^2 + \|B\|_{\sigma 2}^2 = \|A + jB\|_{\sigma 2}^2.$$

In particular,

$$(\operatorname{tr} \langle A \rangle)^2 + (\operatorname{tr} \langle B \rangle)^2 \leq (\operatorname{tr} \langle A + jB \rangle)^2,$$

$$\sigma_{\max}^2(A + jB) \leq \sigma_{\max}^2(A) + \sigma_{\max}^2(A),$$

$$\|A + jB\|_{\mathbb{F}}^2 = \|A\|_{\mathbb{F}}^2 + \|B\|_{\mathbb{F}}^2.$$

(Proof: See [219].) (Remark: See Fact 8.18.7.)

Fact 9.9.41. Let $A \in \mathbb{F}^{n \times n}$, let $B \in \mathbb{F}^{n \times n}$, assume that B is Hermitian, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A - \frac{1}{2}(A + A^*)\| \leq \|A - B\|.$$

In particular,

$$\|A - \frac{1}{2}(A + A^*)\|_F \leq \|A - B\|_F$$

and

$$\sigma_{\max}[A - \frac{1}{2}(A + A^*)] \leq \sigma_{\max}(A - B).$$

(Proof: See [197, p. 275] and [1098, p. 150].)

Fact 9.9.42. Let $A, M, S, B \in \mathbb{F}^{n \times n}$, assume that $A = MS$, M is positive semidefinite, and S and B are unitary, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A - S\| \leq \|A - B\|.$$

In particular,

$$\|A - S\|_F \leq \|A - B\|_F.$$

(Proof: See [197, p. 276] and [1098, p. 150].) (Remark: $A = MS$ is the polar decomposition of A . See Corollary 5.6.5.)

Fact 9.9.43. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$, and let $k \in \mathbb{N}$. Then,

$$\|(A - B)^{2k+1}\| \leq 2^{2k} \|A^{2k+1} - B^{2k+1}\|.$$

(Proof: See [197, p. 294] or [758].)

Fact 9.9.44. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|\langle A \rangle - \langle B \rangle\| \leq \sqrt{2\|A + B\|\|A - B\|}.$$

(Proof: See [47].) (Remark: This result is due to Kosaki and Bhatia.)

Fact 9.9.45. Let $A, B \in \mathbb{F}^{n \times n}$, and let $p \geq 1$. Then,

$$\|\langle A \rangle - \langle B \rangle\|_{\sigma p} \leq \max\left\{2^{1/p-1/2}, 1\right\} \sqrt{\|A + B\|_{\sigma p} \|A - B\|_{\sigma p}}.$$

(Proof: See [47].) (Remark: This result is due to Kittaneh, Kosaki, and Bhatia.)

Fact 9.9.46. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{2n \times 2n}$. Then,

$$\left\| \begin{bmatrix} A + B & 0 \\ 0 & 0 \end{bmatrix} \right\| \leq \left\| \begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \right\| + \left\| \begin{bmatrix} A^{1/2}B^{1/2} & 0 \\ 0 & A^{1/2}B^{1/2} \end{bmatrix} \right\|.$$

In particular,

$$\sigma_{\max}(A + B) \leq \max\{\sigma_{\max}(A), \sigma_{\max}(B)\} + \sigma_{\max}(A^{1/2}B^{1/2})$$

and, for all $p \in [1, \infty)$,

$$\|A + B\|_{\sigma p} \leq (\|A\|_{\sigma p}^p + \|B\|_{\sigma p}^p)^{1/p} + 2^{1/p} \|A^{1/2}B^{1/2}\|_{\sigma p}.$$

(Proof: See [818, 821, 825].) (Remark: See Fact 9.14.15 for a tighter upper bound for $\sigma_{\max}(A + B)$.)

Fact 9.9.47. Let $A, X, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A^*XB\| \leq \frac{1}{2}\|AA^*X + XBB^*\|.$$

In particular,

$$\|A^*B\| \leq \frac{1}{2}\|AA^* + BB^*\|.$$

(Proof: See [61, 202, 209, 525, 815].) (Remark: The first result is *McIntosh's inequality*.) (Remark: See Fact 9.14.23.)

Fact 9.9.48. Let $A, X, B \in \mathbb{F}^{n \times n}$, assume that X is positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A^*XB + B^*XA\| \leq \|A^*XA + B^*XB\|.$$

In particular,

$$\|A^*B + B^*A\| \leq \|A^*A + B^*B\|.$$

(Proof: See [819].) (Remark: See [819] for extensions to the case in which X is not necessarily positive semidefinite.)

Fact 9.9.49. Let $A, X, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let $p \in [0, 1]$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A^pXB^{1-p} + A^{1-p}XB^p\| \leq \|AX + XB\|$$

and

$$\|A^pXB^{1-p} - A^{1-p}XB^p\| \leq |2p - 1|\|AX - XB\|.$$

(Proof: See [61, 203, 216, 510].) (Remark: These results are the *Heinz inequalities*.)

Fact 9.9.50. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is nonsingular and B is Hermitian, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|B\| \leq \frac{1}{2}\|ABA^{-1} + A^{-1}BA\|.$$

(Proof: See [347, 517].)

Fact 9.9.51. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. If $r \in [0, 1]$, then

$$\|A^r - B^r\| \leq \|\langle A - B \rangle^r\|.$$

Furthermore, if $r \in [1, \infty)$, then

$$\|\langle A - B \rangle^r\| \leq \|A^r - B^r\|.$$

In particular,

$$\|(A - B)^2\| \leq \|A^2 - B^2\|.$$

(Proof: See [197, pp. 293, 294] and [820].)

Fact 9.9.52. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$, and let $z \in \mathbb{F}$. Then,

$$\|A - |z|B\| \leq \|A + zB\| \leq \|A + |z|B\|.$$

In particular,

$$\|A - B\| \leq \|A + B\|.$$

(Proof: See [210].) (Remark: Extensions to weak log majorization are given in [1483].) (Remark: The special case $z = 1$ is given in [215].)

Fact 9.9.53. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. If $r \in [0, 1]$, then

$$\|(A + B)^r\| \leq \|A^r + B^r\|.$$

Furthermore, if $r \in [1, \infty)$, then

$$\|A^r + B^r\| \leq \|(A + B)^r\|.$$

In particular, if $k \geq 1$, then

$$\|A^k + B^k\| \leq \|(A + B)^k\|.$$

(Proof: See [58].)

Fact 9.9.54. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|\log(I + A) - \log(I + B)\| \leq \|\log(I + \langle A - B \rangle)\|$$

and

$$\|\log(I + A + B)\| \leq \|\log(I + A) + \log(I + B)\|.$$

(Proof: See [58] and [197, p. 293].) (Remark: See Fact 11.16.16.)

Fact 9.9.55. Let $A, X, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|(\log A)X - X(\log B)\| \leq \|A^{1/2}XB^{-1/2} - A^{-1/2}XB^{1/2}\|.$$

(Proof: See [216].) (Remark: See Fact 11.16.17.)

Fact 9.9.56. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, let $\|\cdot\|$ be a normalized submultiplicative norm on $\mathbb{F}^{n \times n}$, and assume that $\|A - B\| < 1/\|A^{-1}\|$. Then, B is nonsingular. (Remark: See Fact 9.8.44.)

Fact 9.9.57. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, let $\|\cdot\|$ be a normalized submultiplicative norm on $\mathbb{F}^{n \times n}$, let $\gamma > 0$, and assume that $\|A^{-1}\| < \gamma$ and $\|A - B\| < 1/\gamma$. Then, B is nonsingular,

$$\|B^{-1}\| \leq \frac{\gamma}{1 - \gamma\|B - A\|},$$

and

$$\|A^{-1} - B^{-1}\| \leq \gamma^2\|A - B\|.$$

(Proof: See [447, p. 148].) (Remark: See Fact 9.8.44.)

Fact 9.9.58. Let $A, B \in \mathbb{F}^{n \times n}$, let $\lambda \in \mathbb{C}$, assume that $\lambda I - A$ is nonsingular, let $\|\cdot\|$ be a normalized submultiplicative norm on $\mathbb{F}^{n \times n}$, let $\gamma > 0$, and assume that $\|(\lambda I - A)^{-1}\| < \gamma$ and $\|A - B\| < 1/\gamma$. Then, $\lambda I - B$ is nonsingular,

$$\|(\lambda I - B)^{-1}\| \leq \frac{\gamma}{1 - \gamma\|B - A\|},$$

and

$$\|(\lambda I - A)^{-1} - (\lambda I - B)^{-1}\| \leq \frac{\gamma^2\|A - B\|}{1 - \gamma\|A - B\|}.$$

(Proof: See [447, pp. 149, 150].) (Remark: See Fact 9.9.57.)

Fact 9.9.59. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and $A + B$ are nonsingular, and let $\|\cdot\|$ be a normalized submultiplicative norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A^{-1} - (A + B)^{-1}\| \leq \|A^{-1}\| \|(A + B)^{-1}\| \|B\|.$$

If, in addition, $\|A^{-1}B\| < 1$, then

$$\|A^{-1} + (A + B)^{-1}\| \leq \frac{\|A^{-1}\| \|A^{-1}B\|}{1 - \|A^{-1}B\|}.$$

Furthermore, if $\|A^{-1}B\| < 1$ and $\|B\| < 1/\|A^{-1}\|$, then

$$\|A^{-1} - (A + B)^{-1}\| \leq \frac{\|A^{-1}\|^2 \|B\|}{1 - \|A^{-1}\| \|B\|}.$$

Fact 9.9.60. Let $A \in \mathbb{F}^{n \times n}$, assume that A is nonsingular, let $E \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a normalized norm on $\mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} (A + E)^{-1} &= A^{-1}(I + EA^{-1})^{-1} \\ &= A^{-1} - A^{-1}EA^{-1} + O(\|E\|^2). \end{aligned}$$

Fact 9.9.61. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times k}$. Then,

$$\|A \otimes B\|_{\text{col}} = \|A\|_{\text{col}} \|B\|_{\text{col}},$$

$$\|A \otimes B\|_{\infty} = \|A\|_{\infty} \|B\|_{\infty},$$

$$\|A \otimes B\|_{\text{row}} = \|A\|_{\text{row}} \|B\|_{\text{row}}.$$

Furthermore, if $p \in [1, \infty]$, then

$$\|A \otimes B\|_p = \|A\|_p \|B\|_p.$$

Fact 9.9.62. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A \circ B\|^2 \leq \|A^*A\| \|B^*B\|.$$

(Proof: See [712].)

Fact 9.9.63. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are normal, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|A + B\| \leq \|\langle A \rangle + \langle B \rangle\|$$

and

$$\|A \circ B\| \leq \|\langle A \rangle \circ \langle B \rangle\|.$$

(Proof: See [90, 825] and [711, p. 213].)

Fact 9.9.64. Let $A \in \mathbb{R}^{n \times n}$, assume that A is nonsingular, let $b \in \mathbb{R}^n$, and let $\hat{x} \in \mathbb{R}^n$. Then,

$$\frac{1}{\kappa(A)} \frac{\|A\hat{x} - b\|}{\|b\|} \leq \frac{\|\hat{x} - A^{-1}b\|}{\|A^{-1}b\|} \leq \kappa(A) \frac{\|A\hat{x} - b\|}{\|b\|},$$

where $\kappa(A) \triangleq \|A\| \|A^{-1}\|$ and the vector and matrix norms are compatible. Equivalently, letting $\hat{b} \triangleq A\hat{x} - b$ and $x \triangleq A^{-1}b$, it follows that

$$\frac{1}{\kappa(A)} \frac{\|\hat{b}\|}{\|b\|} \leq \frac{\|\hat{x} - x\|}{\|x\|} \leq \kappa(A) \frac{\|\hat{b}\|}{\|b\|}.$$

(Remark: This result estimates the accuracy of an approximate solution $\hat{x}$ to $Ax = b$. $\kappa(A)$ is the *condition number* of A .) (Remark: See [1501].)

Fact 9.9.65. Let $A \in \mathbb{R}^{n \times n}$, assume that A is nonsingular, let $\hat{A} \in \mathbb{R}^{n \times n}$, assume that $\|A^{-1}\hat{A}\| < 1$, and let $b, \hat{b} \in \mathbb{R}^n$. Furthermore, let $x \in \mathbb{R}^n$ satisfy $Ax = b$, and let $\hat{x} \in \mathbb{R}^n$ satisfy $(A + \hat{A})\hat{x} = b + \hat{b}$. Then,

$$\frac{\|\hat{x} - x\|}{\|x\|} \leq \frac{\kappa(A)}{1 - \|A^{-1}\hat{A}\|} \left(\frac{\|\hat{b}\|}{\|b\|} + \frac{\|\hat{A}\|}{\|A\|} \right),$$

where $\kappa(A) \triangleq \|A\| \|A^{-1}\|$ and the vector and matrix norms are compatible. If, in addition, $\|A^{-1}\| \|\hat{A}\| < 1$, then

$$\frac{1}{\kappa(A) + 1} \frac{\|\hat{b} - \hat{A}x\|}{\|b\|} \leq \frac{\|\hat{x} - x\|}{\|x\|} \leq \frac{\kappa(A)}{1 - \|A^{-1}\hat{A}\|} \frac{\|\hat{b} - \hat{A}x\|}{\|b\|}.$$

(Proof: See [407, 408].)

Fact 9.9.66. Let $A, \hat{A} \in \mathbb{R}^{n \times n}$ satisfy $\|A^+\hat{A}\| < 1$, let $b \in \mathcal{R}(A)$, let $\hat{b} \in \mathbb{R}^n$, and assume that $b + \hat{b} \in \mathcal{R}(A + \hat{A})$. Furthermore, let $\hat{x} \in \mathbb{R}^n$ satisfy $(A + \hat{A})\hat{x} = b + \hat{b}$. Then, $x \triangleq A^+b + (I - A^+A)\hat{x}$ satisfies $Ax = b$ and

$$\frac{\|\hat{x} - x\|}{\|x\|} \leq \frac{\kappa(A)}{1 - \|A^+\hat{A}\|} \left(\frac{\|\hat{b}\|}{\|b\|} + \frac{\|\hat{A}\|}{\|A\|} \right),$$

where $\kappa(A) \triangleq \|A\| \|A^{-1}\|$ and the vector and matrix norms are compatible. (Proof: See [407].) (Remark: See [408] for a lower bound.)

9.10 Facts on Matrix Norms for Partitioned Matrices

Fact 9.10.1. Let $A \in \mathbb{F}^{n \times m}$ be the partitioned matrix

$$A = \begin{bmatrix} A_{11} & A_{12} & \cdots & A_{1k} \\ A_{21} & A_{22} & \cdots & A_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ A_{k1} & A_{k2} & \cdots & A_{kk} \end{bmatrix},$$

where $A_{ij} \in \mathbb{F}^{n_i \times n_j}$ for all $i, j = 1, \dots, k$. Furthermore, define $\mu(A) \in \mathbb{R}^{k \times k}$ by

$$\mu(A) \triangleq \begin{bmatrix} \sigma_{\max}(A_{11}) & \sigma_{\max}(A_{12}) & \cdots & \sigma_{\max}(A_{1k}) \\ \sigma_{\max}(A_{21}) & \sigma_{\max}(A_{22}) & \cdots & \sigma_{\max}(A_{2k}) \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_{\max}(A_{k1}) & \sigma_{\max}(A_{k2}) & \cdots & \sigma_{\max}(A_{kk}) \end{bmatrix}.$$

Finally, let $B \in \mathbb{F}^{n \times m}$ be partitioned conformally with A . Then, the following statements hold:

- i) For all $\alpha \in \mathbb{F}$, $\mu(\alpha A) \leq |\alpha| \mu(A)$.
- ii) $\mu(A + B) \leq \mu(A) + \mu(B)$.
- iii) $\mu(AB) \leq \mu(A) \mu(B)$.
- iv) $\text{sprad}(A) \leq \text{sprad}[\mu(A)]$.
- v) $\sigma_{\max}(A) \leq \sigma_{\max}[\mu(A)]$.

(Proof: See [400, 1055, 1205].) (Remark: $\mu(A)$ is a *matricial norm*.) (Remark: This result is a norm-compression inequality.)

Fact 9.10.2. Let $A \in \mathbb{F}^{n \times m}$ be the partitioned matrix

$$A = \begin{bmatrix} A_{11} & A_{12} & \cdots & A_{1k} \\ A_{21} & A_{22} & \cdots & A_{2k} \\ \vdots & \vdots & \ddots & \vdots \\ A_{k1} & A_{k2} & \cdots & A_{kk} \end{bmatrix},$$

where $A_{ij} \in \mathbb{F}^{n_i \times n_j}$ for all $i, j = 1, \dots, k$. Then, the following statements hold:

- i) If $p \in [1, 2]$, then

$$\sum_{i,j=1}^k \|A_{ij}\|_{\sigma p}^2 \leq \|A\|_{\sigma p}^2 \leq k^{4/p-2} \sum_{i,j=1}^k \|A_{ij}\|_{\sigma p}^2.$$

- ii) If $p \in [2, \infty]$, then

$$k^{4/p-2} \sum_{i,j=1}^k \|A_{ij}\|_{\sigma p}^2 \leq \|A\|_{\sigma p}^2 \leq \sum_{i,j=1}^k \|A_{ij}\|_{\sigma p}^2.$$

- iii) If $p \in [1, 2]$, then

$$\|A\|_{\sigma p}^p \leq \sum_{i,j=1}^k \|A_{ij}\|_{\sigma p}^p \leq k^{2-p} \|A\|_{\sigma p}^p.$$

iv) If $p \in [2, \infty)$, then

$$k^{2-p} \|A\|_{\sigma p}^p \leq \sum_{i,j=1}^k \|A_{ij}\|_{\sigma p}^p \leq \|A\|_{\sigma p}^p.$$

v) $\|A\|_{\sigma 2}^2 = \sum_{i,j=1}^k \|A_{ij}\|_{\sigma 2}^2.$

vii) For all $p \in [1, \infty)$,

$$\left(\sum_{i=1}^k \|A_{ii}\|_{\sigma p}^p \right)^{1/p} \leq \|A\|_{\sigma p}.$$

viii) For all $i = 1, \dots, k$,

$$\sigma_{\max}(A_{ii}) \leq \sigma_{\max}(A).$$

(Proof: See [129, 208].)

Fact 9.10.3. Let $A, B \in \mathbb{F}^{n \times n}$, and define $\mathcal{A} \in \mathbb{F}^{kn \times kn}$ by

$$\mathcal{A} \triangleq \begin{bmatrix} A & B & B & \cdots & B \\ B & A & B & \cdots & B \\ B & B & A & \ddots & B \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ B & B & B & \cdots & A \end{bmatrix}.$$

Then,

$$\sigma_{\max}(\mathcal{A}) = \max\{\sigma_{\max}(A + (k-1)B), \sigma_{\max}(A - B)\}.$$

Now, let $p \in [1, \infty)$. Then,

$$\|\mathcal{A}\|_{\sigma p} = (\|A + (k-1)B\|_{\sigma p}^p + (k-1)\|A - B\|_{\sigma p}^p)^{1/p}.$$

(Proof: See [129].)

Fact 9.10.4. Let $A \in \mathbb{F}^{n \times n}$, and define $\mathcal{A} \in \mathbb{F}^{kn \times kn}$ by

$$\mathcal{A} \triangleq \begin{bmatrix} A & A & A & \cdots & A \\ -A & A & A & \cdots & A \\ -A & -A & A & \ddots & A \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ -A & -A & -A & \cdots & A \end{bmatrix}.$$

Then,

$$\sigma_{\max}(\mathcal{A}) = \sqrt{\frac{2}{1 - \cos(\pi/k)}} \sigma_{\max}(A).$$

Furthermore, define $\mathcal{A}_0 \in \mathbb{F}^{kn \times kn}$ by

$$\mathcal{A}_0 \triangleq \begin{bmatrix} 0 & A & A & \cdots & A \\ -A & 0 & A & \cdots & A \\ -A & -A & 0 & \ddots & A \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ -A & -A & -A & \cdots & 0 \end{bmatrix}.$$

Then,

$$\sigma_{\max}(\mathcal{A}_0) = \sqrt{\frac{1 + \cos(\pi/k)}{1 - \cos(\pi/k)}} \sigma_{\max}(A).$$

(Proof: See [129].) (Remark: Extensions to Schatten norms are given in [129].)

Fact 9.10.5. Let $A, B, C, D \in \mathbb{F}^{n \times n}$. Then,

$$\frac{1}{2} \max\{\sigma_{\max}(A + B + C + D), \sigma_{\max}(A - B - C + D)\} \leq \sigma_{\max}\left(\begin{bmatrix} A & B \\ C & D \end{bmatrix}\right).$$

Now, let $p \in [1, \infty)$. Then,

$$\frac{1}{2}(\|A + B + C + D\|_{\sigma p}^p + \|A - B - C + D\|_{\sigma p}^p)^{1/p} \leq \left\| \begin{bmatrix} A & B \\ C & D \end{bmatrix} \right\|_{\sigma p}.$$

(Proof: See [129].)

Fact 9.10.6. Let $A, B, C \in \mathbb{F}^{n \times n}$, define

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix},$$

assume that $\mathcal{A}$ is positive semidefinite, let $p \in [1, \infty]$, and define

$$\mathcal{A}_0 \triangleq \begin{bmatrix} \|A\|_{\sigma p} & \|B\|_{\sigma p} \\ \|B\|_{\sigma p} & \|C\|_{\sigma p} \end{bmatrix}.$$

If $p \in [1, 2]$, then

$$\|\mathcal{A}_0\|_{\sigma p} \leq \|\mathcal{A}\|_{\sigma p}.$$

Furthermore, if $p \in [2, \infty]$, then

$$\|\mathcal{A}\|_{\sigma p} \leq \|\mathcal{A}_0\|_{\sigma p}.$$

Hence, if $p = 2$, then

$$\|\mathcal{A}_0\|_{\sigma p} = \|\mathcal{A}\|_{\sigma p}.$$

Finally, if $A = C$, B is Hermitian, and p is an integer, then

$$\|\mathcal{A}\|_{\sigma p}^p = \|A + B\|_{\sigma p}^p + \|A - B\|_{\sigma p}^p$$

and

$$\|\mathcal{A}_0\|_{\sigma p}^p = (\|A\|_{\sigma p} + \|B\|_{\sigma p})^p + \|\|A\|_{\sigma p} - \|B\|_{\sigma p}\|^p.$$

(Proof: See [810].) (Remark: This result is a norm-compression inequality.)

Fact 9.10.7. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$, define

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ B^* & C \end{bmatrix},$$

assume that $\mathcal{A}$ is positive semidefinite, and let $p \geq 1$. If $p \in [1, 2]$, then

$$\|\mathcal{A}\|_{\sigma p}^p \leq \|A\|_{\sigma p}^p + (2^p - 2)\|B\|_{\sigma p}^p + \|C\|_{\sigma p}^p.$$

Furthermore, if $p \geq 2$, then

$$\|\mathcal{A}\|_{\sigma p}^p + (2^p - 2)\|B\|_{\sigma p}^p + \|C\|_{\sigma p}^p \leq \|\mathcal{A}\|_{\sigma p}^p.$$

Finally, if $p = 2$, then

$$\|\mathcal{A}\|_{\sigma p}^p = \|A\|_{\sigma p}^p + (2^p - 2)\|B\|_{\sigma p}^p + \|C\|_{\sigma p}^p.$$

(Proof: See [86].)

Fact 9.10.8. Let $A \in \mathbb{F}^{n \times m}$ be the partitioned matrix

$$A = \begin{bmatrix} A_{11} & \cdots & A_{1k} \\ A_{21} & \cdots & A_{2k} \end{bmatrix},$$

where $A_{ij} \in \mathbb{F}^{n_i \times n_j}$ for all $i, j = 1, \dots, k$. Then, the following statements are conjectured to hold:

i) If $p \in [1, 2]$, then

$$\left\| \begin{bmatrix} \|A_{11}\|_{\sigma p} & \cdots & \|A_{1k}\|_{\sigma p} \\ \|A_{21}\|_{\sigma p} & \cdots & \|A_{2k}\|_{\sigma p} \end{bmatrix} \right\|_{\sigma p} \leq \|A\|_{\sigma p}.$$

ii) If $p \geq 2$, then

$$\|A\|_{\sigma p} \leq \left\| \begin{bmatrix} \|A_{11}\|_{\sigma p} & \cdots & \|A_{1k}\|_{\sigma p} \\ \|A_{21}\|_{\sigma p} & \cdots & \|A_{2k}\|_{\sigma p} \end{bmatrix} \right\|_{\sigma p}.$$

(Proof: See [87]. The result is true when all blocks have rank 1 or when $p \geq 4$.)

(Remark: This result is a norm-compression inequality.)

9.11 Facts on Matrix Norms and Eigenvalues Involving One Matrix

Fact 9.11.1. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$|\det A| \leq \prod_{i=1}^n \|\text{row}_i(A)\|_2$$

and

$$|\det A| \leq \prod_{i=1}^n \|\text{col}_i(A)\|_2.$$

(Proof: The result follows from Hadamard's inequality. See Fact 8.17.11.)

Fact 9.11.2. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\text{Re tr } A \leq |\text{tr } A| \leq \sum_{i=1}^n |\lambda_i| \leq \|A\|_{\sigma 1} = \text{tr } \langle A \rangle = \sum_{i=1}^n \sigma_i(A).$$

In addition, if A is normal, then

$$\|A\|_{\sigma 1} = \sum_{i=1}^n |\lambda_i|.$$

Finally, A is positive semidefinite if and only if

$$\|A\|_{\sigma 1} = \text{tr } A.$$

(Proof: See Fact 5.14.15 and Fact 9.13.19.) (Remark: See Fact 5.11.9 and Fact 5.14.15.) (Problem: Refine the second statement for necessity and sufficiency. See [742].)

Fact 9.11.3. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\begin{aligned} \text{Re tr } A^2 &\leq |\text{tr } A^2| \leq \sum_{i=1}^n |\lambda_i|^2 \leq \|A^2\|_{\sigma 1} = \text{tr } \langle A^2 \rangle = \sum_{i=1}^n \sigma_i(A^2) \\ &\leq \sum_{i=1}^n \sigma_i^2(A) = \text{tr } A^*A = \text{tr } \langle A \rangle^2 = \|A\|_{\sigma 2}^2 = \|A\|_{\text{F}}^2 \end{aligned}$$

and

$$\|A\|_{\text{F}}^2 - \sqrt{\frac{n^3-n}{12}} \| [A, A^*] \|_{\text{F}} \leq \sum_{i=1}^n |\lambda_i|^2 \leq \sqrt{\|A\|_{\text{F}}^4 - \frac{1}{2} \| [A, A^*] \|_{\text{F}}^2} \leq \|A\|_{\text{F}}^2.$$

Consequently, A is normal if and only if

$$\|A\|_{\text{F}}^2 = \sum_{i=1}^n |\lambda_i|^2.$$

Furthermore,

$$\sum_{i=1}^n |\lambda_i|^2 \leq \sqrt{\|A\|_{\text{F}}^4 - \frac{1}{4} (\text{tr } [A, A^*])^2} \leq \|A\|_{\text{F}}^2$$

and

$$\sum_{i=1}^n |\lambda_i|^2 \leq \sqrt{\|A\|_{\text{F}}^4 - \frac{n^2}{4} |\det [A, A^*]|^{2/n}} \leq \|A\|_{\text{F}}^2.$$

Finally, A is Hermitian if and only if

$$\|A\|_{\text{F}}^2 = \text{tr } A^2.$$

(Proof: Use Fact 8.17.5 and Fact 9.11.2. The lower bound involving the commutator is due to Henrici; the corresponding upper bound is given in [847]. The bounds in the penultimate statement are given in [847]. The last statement follows from Fact 3.7.13.) (Remark: $\text{tr } (A + A^*)^2 \geq 0$ and $\text{tr } (A - A^*)^2 \leq 0$ yield $|\text{tr } A^2| \leq \|A\|_{\text{F}}^2$.) (Remark: The result $\sum_{i=1}^n |\lambda_i|^2 \leq \|A\|_{\text{F}}^2$ is *Schur's inequality*. See Fact 8.17.5.) (Remark: See Fact 5.11.10, Fact 9.11.5, Fact 9.13.17, and Fact 9.13.20.) (Problem: Merge the first two strings.)

Fact 9.11.4. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$|\operatorname{tr} A^2| \leq (\operatorname{rank} A) \sqrt{\|A\|_F^4 - \frac{1}{2} \|[A, A^*]\|_F^2}.$$

(Proof: See [315].)

Fact 9.11.5. Let $A \in \mathbb{F}^{n \times n}$, let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, and define

$$\alpha \triangleq \sqrt{\left(\|A\|_F^2 - \frac{1}{n} |\operatorname{tr} A|^2\right)^2 - \frac{1}{2} \|[A, A^*]\|_F^2 + \frac{1}{n} |\operatorname{tr} A|^2}.$$

Then,

$$\begin{aligned} \sum_{i=1}^n |\lambda_i|^2 &\leq \alpha \leq \sqrt{\|A\|_F^4 - \frac{1}{2} \|[A, A^*]\|_F^2} \leq \|A\|_F^2, \\ \sum_{i=1}^n (\operatorname{Re} \lambda_i)^2 &\leq \frac{1}{2} (\alpha + \operatorname{Re} \operatorname{tr} A^2), \\ \sum_{i=1}^n (\operatorname{Im} \lambda_i)^2 &\leq \frac{1}{2} (\alpha - \operatorname{Re} \operatorname{tr} A^2). \end{aligned}$$

(Proof: See [732].) (Remark: The first string of inequalities interpolates the upper bound for $\sum_{i=1}^n |\lambda_i|^2$ in the second string of inequalities in Fact 9.11.3.)

Fact 9.11.6. Let $A \in \mathbb{F}^{n \times n}$, let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, and let $p \in (0, 2]$.

Then,

$$\sum_{i=1}^n |\lambda_i|^p \leq \sum_{i=1}^n \sigma_i^p(A) = \|A\|_{\sigma p}^p \leq \|A\|_p^p.$$

(Proof: The left-hand inequality, which holds for all $p > 0$, follows from Weyl's inequality in Fact 8.17.5. The right-hand inequality is given by Proposition 9.2.5.)

(Remark: This result is the *generalized Schur inequality*.) (Remark: The case of equality is discussed in [742] for $p \in [1, 2)$.)

Fact 9.11.7. Let $A \in \mathbb{F}^{n \times n}$, and let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\|A\|_F^2 - \sum_{i=1}^n |\lambda_i|^2 = 2 \left(\left\| \frac{1}{2j} (A - A^*) \right\|_F^2 - \sum_{i=1}^n |\operatorname{Im} \lambda_i|^2 \right).$$

(Proof: See Fact 5.11.22.) (Remark: This result is an extension of Browne's theorem.)

Fact 9.11.8. Let $A \in \mathbb{R}^{n \times n}$, and let $\lambda \in \operatorname{spec}(A)$. Then, the following inequalities hold:

- i) $|\lambda| \leq n \|A\|_\infty$.
- ii) $|\operatorname{Re} \lambda| \leq \frac{n}{2} \|A + A^T\|_\infty$.
- iii) $|\operatorname{Im} \lambda| \leq \frac{\sqrt{n^2 - n}}{2\sqrt{2}} \|A - A^T\|_\infty$.

(Proof: See [963, p. 140].) (Remark: i) and ii) are *Hirsch's theorems*, while iii) is *Bendixson's theorem*. See Fact 5.11.21.)

9.12 Facts on Matrix Norms and Eigenvalues Involving Two or More Matrices

Fact 9.12.1. Let $A, B \in \mathbb{F}^{n \times m}$, let $\text{mspec}(A^*B) = \{\lambda_1, \dots, \lambda_m\}_{\text{ms}}$, let $p, q \in [1, \infty]$ satisfy $1/p + 1/q = 1$, and define $r \triangleq \min\{m, n\}$. Then,

$$|\text{tr } A^*B| \leq \sum_{i=1}^m |\lambda_i| \leq \|A^*B\|_{\sigma 1} = \sum_{i=1}^m \sigma_i(A^*B) \leq \sum_{i=1}^r \sigma_i(A) \sigma_i(B) \leq \|A\|_{\sigma p} \|B\|_{\sigma q}.$$

In particular,

$$|\text{tr } A^*B| \leq \|A\|_{\text{F}} \|B\|_{\text{F}}.$$

(Proof: Use Proposition 9.6.2 and Fact 9.11.2. The last inequality in the string of inequalities is Hölder's inequality.) (Remark: See Fact 9.9.11.) (Remark: The result

$$|\text{tr } A^*B| \leq \sum_{i=1}^r \sigma_i(A) \sigma_i(B)$$

is *von Neumann's trace inequality*. See [250].)

Fact 9.12.2. Let $A, B \in \mathbb{F}^{n \times m}$, and let $\text{mspec}(A^*B) = \{\lambda_1, \dots, \lambda_m\}_{\text{ms}}$. Then,

$$|\text{tr } (A^*B)^2| \leq \sum_{i=1}^m |\lambda_i|^2 \leq \sum_{i=1}^m \sigma_i^2(A^*B) = \text{tr } AA^*BB^* = \|A^*B\|_{\text{F}}^2 \leq \|A\|_{\text{F}}^2 \|B\|_{\text{F}}^2.$$

(Proof: Use Fact 8.17.5.)

Fact 9.12.3. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and let $\text{mspec}(A + jB) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\sum_{i=1}^n |\text{Re } \lambda_i|^2 \leq \|A\|_{\text{F}}^2$$

and

$$\sum_{i=1}^n |\text{Im } \lambda_i|^2 \leq \|B\|_{\text{F}}^2.$$

(Proof: See [1098, p. 146].)

Fact 9.12.4. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and let $\|\cdot\|$ be a weakly unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} & \left\| \begin{bmatrix} \lambda_1(A) & & 0 \\ & \ddots & \\ 0 & & \lambda_n(A) \end{bmatrix} - \begin{bmatrix} \lambda_1(B) & & 0 \\ & \ddots & \\ 0 & & \lambda_n(B) \end{bmatrix} \right\| \leq \|A - B\| \\ & \leq \left\| \begin{bmatrix} \lambda_1(A) & & 0 \\ & \ddots & \\ 0 & & \lambda_n(A) \end{bmatrix} - \begin{bmatrix} \lambda_n(B) & & 0 \\ & \ddots & \\ 0 & & \lambda_1(B) \end{bmatrix} \right\|. \end{aligned}$$

In particular,

$$\max_{i \in \{1, \dots, n\}} |\lambda_i(A) - \lambda_i(B)| \leq \sigma_{\max}(A - B) \leq \max_{i \in \{1, \dots, n\}} |\lambda_i(A) - \lambda_{n-i+1}(B)|$$

and

$$\sum_{i=1}^n [\lambda_i(A) - \lambda_i(B)]^2 \leq \|A - B\|_F^2 \leq \sum_{i=1}^n [\lambda_i(A) - \lambda_{n-i+1}(B)]^2.$$

(Proof: See [47], [196, p. 38], [197, pp. 63, 69], [200, p. 38], [796, p. 126], [878, p. 134], [895], or [1230, p. 202].) (Remark: The first inequality is the *Lidskii-Mirsky-Wielandt theorem*. The result can be stated without norms using Fact 9.8.42. See [895].) (Remark: See Fact 9.14.29.) (Remark: The case in which A and B are normal is considered in Fact 9.12.8.)

Fact 9.12.5. Let $A, B \in \mathbb{F}^{n \times n}$, let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$ and $\text{mspec}(B) = \{\mu_1, \dots, \mu_n\}_{\text{ms}}$, and assume that A and B satisfy at least one of the following conditions:

- i) A and B are Hermitian.
- ii) A is Hermitian, and B is skew Hermitian.
- iii) A is skew Hermitian, and B is Hermitian.
- iv) A and B are unitary.
- v) There exist nonzero $\alpha, \beta \in \mathbb{C}$ such that αA and βB are unitary.
- vi) A , B , and $A - B$ are normal.

Then,

$$\min \sigma_{\max} \left(\begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix} - \begin{bmatrix} \mu_{\sigma(1)} & & 0 \\ & \ddots & \\ 0 & & \mu_{\sigma(n)} \end{bmatrix} \right) \leq \sigma_{\max}(A - B),$$

where the minimum is taken over all permutations σ of $\{1, \dots, n\}$. (Proof: See [200, pp. 52, 152].)

Fact 9.12.6. Let $A, B \in \mathbb{F}^{n \times n}$, let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$ and $\text{mspec}(B) = \{\mu_1, \dots, \mu_n\}_{\text{ms}}$, and assume that A is normal. Then,

$$\min \left\| \begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix} - \begin{bmatrix} \mu_{\sigma(1)} & & 0 \\ & \ddots & \\ 0 & & \mu_{\sigma(n)} \end{bmatrix} \right\|_F \leq \sqrt{n} \|A - B\|_F,$$

where the minimum is taken over all permutations σ of $\{1, \dots, n\}$. If, in addition, B is normal, then there exists $c \in (0, 2.9039)$ such that

$$\min \sigma_{\max} \left(\begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix} - \begin{bmatrix} \mu_{\sigma(1)} & & 0 \\ & \ddots & \\ 0 & & \mu_{\sigma(n)} \end{bmatrix} \right) \leq c \sigma_{\max}(A - B).$$

(Proof: See [200, pp. 152, 153, 173].) (Remark: Constants c for alternative Schatten norms are given in [200, p. 159].) (Remark: If, in addition, $A - B$ is normal, then

it follows from Fact 9.12.5 that the last inequality holds with $c = 1$.)

Fact 9.12.7. Let $A, B \in \mathbb{F}^{n \times n}$, let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$ and $\text{mspec}(B) = \{\mu_1, \dots, \mu_n\}_{\text{ms}}$, and assume that A is Hermitian. Then,

$$\min \left\| \begin{bmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{bmatrix} - \begin{bmatrix} \mu_{\sigma(1)} & & 0 \\ & \ddots & \\ 0 & & \mu_{\sigma(n)} \end{bmatrix} \right\|_{\text{F}} \leq \sqrt{2} \|A - B\|_{\text{F}},$$

where the minimum is taken over all permutations σ of $\{1, \dots, n\}$. (Proof: See [200, p. 174].)

Fact 9.12.8. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are normal, and let $\text{spec}(A) = \{\lambda_1, \dots, \lambda_q\}$ and $\text{spec}(B) = \{\mu_1, \dots, \mu_r\}$. Then,

$$\sigma_{\max}(A - B) \leq \max\{|\lambda_i - \lambda_j| : i = 1, \dots, q, j = 1, \dots, r\}.$$

(Proof: See [197, p. 164].) (Remark: The case in which A and B are Hermitian is considered in Fact 9.12.4.)

Fact 9.12.9. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are normal. Then, there exists a permutation σ of $1, \dots, n$ such that

$$\sum_{i=1}^n |\lambda_{\sigma(i)}(A) - \lambda_i(B)|^2 \leq \|A - B\|_{\text{F}}^2.$$

(Proof: See [709, p. 368] or [1098, pp. 160, 161].) (Remark: This inequality is the *Hoffman-Wielandt theorem*.) (Remark: The case in which A and B are Hermitian is considered in Fact 9.12.4.)

Fact 9.12.10. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is Hermitian and B is normal. Furthermore, let $\text{mspec}(B) = \{\lambda_1(B), \dots, \lambda_n(B)\}_{\text{ms}}$, where $\text{Re } \lambda_n(B) \leq \dots \leq \text{Re } \lambda_1(B)$. Then,

$$\sum_{i=1}^n |\lambda_i(A) - \lambda_i(B)|^2 \leq \|A - B\|_{\text{F}}^2.$$

(Proof: See [709, p. 370].) (Remark: This result is a special case of Fact 9.12.9.) (Remark: The left-hand side has the same value for all orderings that satisfy $\text{Re } \lambda_n(B) \leq \dots \leq \text{Re } \lambda_1(B)$.)

Fact 9.12.11. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be an induced norm on $\mathbb{F}^{n \times n}$. Then,

$$|\det A - \det B| \leq \begin{cases} \|A - B\| \frac{\|A\|^n - \|B\|^n}{\|A\| - \|B\|}, & \|A\| \neq \|B\|, \\ n \|A - B\| \|A\|^{n-1}, & \|A\| = \|B\|. \end{cases}$$

(Proof: See [505].) (Remark: See Fact 1.18.2.)

9.13 Facts on Matrix Norms and Singular Values for One Matrix

Fact 9.13.1. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\sigma_{\max}(A) = \max_{x \in \mathbb{F}^m \setminus \{0\}} \left(\frac{x^* A^* A x}{x^* x} \right)^{1/2},$$

and thus

$$\|Ax\|_2 \leq \sigma_{\max}(A) \|x\|_2.$$

Furthermore,

$$\lambda_{\min}^{1/2}(A^* A) = \min_{x \in \mathbb{F}^n \setminus \{0\}} \left(\frac{x^* A^* A x}{x^* x} \right)^{1/2},$$

and thus

$$\lambda_{\min}^{1/2}(A^* A) \|x\|_2 \leq \|Ax\|_2.$$

If, in addition, $m \leq n$, then

$$\sigma_m(A) = \min_{x \in \mathbb{F}^n \setminus \{0\}} \left(\frac{x^* A^* A x}{x^* x} \right)^{1/2},$$

and thus

$$\sigma_m(A) \|x\|_2 \leq \|Ax\|_2.$$

Finally, if $m = n$, then

$$\sigma_{\min}(A) = \min_{x \in \mathbb{F}^n \setminus \{0\}} \left(\frac{x^* A^* A x}{x^* x} \right)^{1/2},$$

and thus

$$\sigma_{\min}(A) \|x\|_2 \leq \|Ax\|_2.$$

(Proof: See Lemma 8.4.3.)

Fact 9.13.2. Let $A \in \mathbb{F}^{n \times m}$. Then,

$$\begin{aligned} \sigma_{\max}(A) &= \max\{|y^* A x| : x \in \mathbb{F}^m, y \in \mathbb{F}^n, \|x\|_2 = \|y\|_2 = 1\} \\ &= \max\{|y^* A x| : x \in \mathbb{F}^m, y \in \mathbb{F}^n, \|x\|_2 \leq 1, \|y\|_2 \leq 1\}. \end{aligned}$$

(Remark: See Fact 9.8.36.)

Fact 9.13.3. Let $x \in \mathbb{F}^n$ and $y \in \mathbb{F}^m$, and define $\mathcal{S} \triangleq \{A \in \mathbb{F}^{n \times m} : \sigma_{\max}(A) \leq 1\}$. Then,

$$\max_{A \in \mathcal{S}} x^* A y = \sqrt{x^* x y^* y}.$$

Fact 9.13.4. Let $\|\cdot\|$ be an equi-induced unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then, $\|\cdot\| = \sigma_{\max}(\cdot)$.

Fact 9.13.5. Let $\|\cdot\|$ be an equi-induced self-adjoint norm on $\mathbb{F}^{n \times n}$. Then, $\|\cdot\| = \sigma_{\max}(\cdot)$.

Fact 9.13.6. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\sigma_{\min}(A) - 1 \leq \sigma_{\min}(A + I) \leq \sigma_{\min}(A) + 1.$$

(Proof: Use Proposition 9.6.8.)

Fact 9.13.7. Let $A \in \mathbb{F}^{n \times n}$, assume that A is normal, and let $r \in \mathbb{N}$. Then,

$$\sigma_{\max}(A^r) = \sigma_{\max}^r(A).$$

(Remark: Matrices that are not normal might also satisfy these conditions. Consider $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$.)

Fact 9.13.8. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\sigma_{\max}^2(A) - \sigma_{\max}(A^2) \leq \sigma_{\max}(A^*A - AA^*) \leq \sigma_{\max}^2(A) - \sigma_{\min}^2(A)$$

and

$$\sigma_{\max}^2(A) + \sigma_{\min}^2(A) \leq \sigma_{\max}(A^*A + AA^*) \leq \sigma_{\max}^2(A) + \sigma_{\max}(A^2).$$

If $A^2 = 0$, then

$$\sigma_{\max}(A^*A - AA^*) = \sigma_{\max}^2(A).$$

(Proof: See [820, 824].) (Remark: See Fact 8.18.11.) (Remark: If A is normal, then it follows that $\sigma_{\max}^2(A) \leq \sigma_{\max}(A^2)$, although Fact 9.13.7 implies that equality holds.)

Fact 9.13.9. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $\text{sprad}(A) = \sigma_{\max}(A)$.
- ii) $\sigma_{\max}(A^i) = \sigma_{\max}^i(A)$ for all $i \in \mathbb{P}$.
- iii) $\sigma_{\max}(A^n) = \sigma_{\max}^n(A)$.

(Proof: See [493] and [711, p. 44].) (Remark: The result $\text{iii}) \implies \text{i})$ is due to Ptak.) (Remark: Additional conditions are given in [567].)

Fact 9.13.10. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\sigma_{\max}(A) \leq \sigma_{\max}(|A|) \leq \sqrt{\text{rank } A} \sigma_{\max}(A).$$

(Proof: See [681, p. 111].)

Fact 9.13.11. Let $A \in \mathbb{F}^{n \times n}$, and let $p \in [1, \infty)$ be an even integer. Then,

$$\|A\|_{\sigma p} \leq \| |A| \|_{\sigma p}.$$

In particular,

$$\|A\|_{\text{F}} \leq \| |A| \|_{\text{F}}$$

and

$$\sigma_{\max}(A) \leq \sigma_{\max}(|A|).$$

Finally, let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{C}^{n \times m}$. Then, $\|A\|_{\text{F}} = \| |A| \|_{\text{F}}$ for all $A \in \mathbb{C}^{n \times m}$ if and only if $\|\cdot\|$ is a constant multiple of $\|\cdot\|_{\text{F}}$. (Proof: See [712] and [730].)

Fact 9.13.12. Let $A \in \mathbb{R}^{n \times n}$, and assume that $r \triangleq \text{rank } A \geq 2$. If $r \text{tr } A^2 \leq (\text{tr } A)^2$, then

$$\sqrt{\frac{(\text{tr } A)^2 - \text{tr } A^2}{r(r-1)}} \leq \text{sprad}(A).$$

If $(\operatorname{tr} A)^2 \leq r \operatorname{tr} A^2$, then

$$\frac{|\operatorname{tr} A|}{r} + \sqrt{\frac{r \operatorname{tr} A^2 - (\operatorname{tr} A)^2}{r^2(r-1)}} \leq \operatorname{sprad}(A).$$

If $\operatorname{rank} A = 2$, then equality holds in both cases. Finally, if A is skew symmetric, then

$$\sqrt{\frac{3}{r(r-1)}} \|A\|_F \leq \operatorname{sprad}(A).$$

(Proof: See [718].)

Fact 9.13.13. Let $A \in \mathbb{R}^{n \times n}$. Then,

$$\sqrt{\frac{1}{2(n^2-n)}} (\|A\|_F^2 + \operatorname{tr} A^2) \leq \sigma_{\max}(A).$$

Furthermore, if $\|A\|_F \leq \operatorname{tr} A$, then

$$\sigma_{\max}(A) \leq \frac{1}{n} \operatorname{tr} A + \sqrt{\frac{n-1}{n} [\|A\|_F^2 - \frac{1}{n} (\operatorname{tr} A)^2]}.$$

(Proof: See [992], which considers the complex case.)

Fact 9.13.14. Let $A \in \mathbb{R}^{n \times n}$. Then, the polynomial $p \in \mathbb{R}[s]$ defined by

$$p(s) \triangleq s^n - \|A\|_F^2 s + (n-1) |\det A|^{2/(n-1)}$$

has either exactly one or exactly two positive roots $0 < \alpha \leq \beta$. Furthermore, α and β satisfy

$$\alpha^{(n-1)/2} \leq \sigma_{\min}(A) \leq \sigma_{\max}(A) \leq \beta^{(n-1)/2}.$$

(Proof: See [1139].)

Fact 9.13.15. Let $A \in \mathbb{R}^{n \times n}$, and, for all $k = 1, \dots, n$, define

$$\alpha_k \triangleq \sum_{\substack{j=1 \\ j \neq k}}^n |A_{(k,j)}|, \quad \beta_k \triangleq \sum_{\substack{i=1 \\ i \neq k}}^n |A_{(i,k)}|.$$

Then,

$$\min_{1 \leq k \leq n} \{|A_{(k,k)}| - \frac{1}{2}(\alpha_k + \beta_k)\} \leq \sigma_{\min}(A).$$

(Proof: See [764, 774].)

Fact 9.13.16. Let $A \in \mathbb{R}^{n \times n}$. Then,

$$\operatorname{tr} \langle A \rangle = \operatorname{tr} \langle A^* \rangle.$$

Fact 9.13.17. Let $A \in \mathbb{R}^{n \times n}$. Then, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \sigma_i(A^2) \leq \sum_{i=1}^k \sigma_i^2(A).$$

Hence,

$$\operatorname{tr} (A^{2*} A^2)^{1/2} \leq \operatorname{tr} A^* A,$$

that is,

$$\operatorname{tr} \langle A^2 \rangle \leq \operatorname{tr} \langle A \rangle^2.$$

(Proof: Let $B = A$ and $r = 1$ in Proposition 9.6.2. See also Fact 9.11.3.)

Fact 9.13.18. Let $A \in \mathbb{F}^{n \times n}$, and let k denote the number of nonzero eigenvalues of A . Then,

$$\left. \begin{aligned} |\operatorname{tr} A^2| &\leq \operatorname{tr} \langle A^2 \rangle \\ \operatorname{tr} \langle A \rangle \langle A^* \rangle & \\ \frac{1}{k} |\operatorname{tr} A|^2 & \end{aligned} \right\} \leq \operatorname{tr} \langle A \rangle^2.$$

(Proof: The upper bound for $|\operatorname{tr} A^2|$ is given by Fact 9.11.3. The upper bound for $\operatorname{tr} \langle A^2 \rangle$ is given by Fact 9.13.17. To prove the center inequality, let $A = S_1 D S_2$ denote the singular value decomposition of A . Then, $\operatorname{tr} \langle A \rangle \langle A^* \rangle = \operatorname{tr} S_3^* D S_3 D$, where $S_3 \triangleq S_1 S_2$, and $\operatorname{tr} A^* A = \operatorname{tr} D^2$. The result now follows using Fact 5.12.4. The remaining inequality is given by Fact 5.11.10.) (Remark: See Fact 5.11.10 and Fact 9.11.3.)

Fact 9.13.19. Let $A \in \mathbb{F}^{n \times n}$, and let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_n$ are ordered such that $|\lambda_1| \geq \dots \geq |\lambda_n|$. Then, for all $k = 1, \dots, n$,

$$\prod_{i=1}^k |\lambda_i|^2 \leq \prod_{i=1}^k \sigma_i(A^2) \leq \prod_{i=1}^k \sigma_i^2(A)$$

and

$$\prod_{i=1}^n |\lambda_i|^2 = \prod_{i=1}^n \sigma_i(A^2) = \prod_{i=1}^n \sigma_i^2(A) = |\det A|^2.$$

Furthermore, for all $k = 1, \dots, n$,

$$\left| \sum_{i=1}^k \lambda_i \right| \leq \sum_{i=1}^k |\lambda_i| \leq \sum_{i=1}^k \sigma_i(A),$$

and thus

$$|\operatorname{tr} A| \leq \sum_{i=1}^k |\lambda_i| \leq \operatorname{tr} \langle A \rangle.$$

(Proof: See [711, p. 172], and use Fact 5.11.28. For the last statement, use Fact 2.21.13.) (Remark: See Fact 5.11.28, Fact 8.18.21, and Fact 9.11.2.) (Remark: This result is due to Weyl.)

Fact 9.13.20. Let $A \in \mathbb{F}^{n \times n}$, and let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_n$ are ordered such that $|\lambda_n| \leq \dots \leq |\lambda_1|$, and let $p \geq 0$. Then, for all $k = 1, \dots, n$,

$$\left| \sum_{i=1}^k \lambda_i^p \right| \leq \sum_{i=1}^k |\lambda_i|^p \leq \sum_{i=1}^k \sigma_i^p(A).$$

(Proof: See [197, p. 42].) (Remark: This result is *Weyl's majorant theorem*.) (Remark: See Fact 9.11.3.)

Fact 9.13.21. Let $A \in \mathbb{F}^{n \times n}$, and define

$$\begin{aligned} r_i &\triangleq \sum_{j=1}^n |A_{(i,j)}|, & c_i &\triangleq \sum_{j=1}^n |A_{(j,i)}|, \\ r_{\min} &\triangleq \min_{i=1,\dots,n} \|r_i\|_2, & c_{\min} &\triangleq \min_{i=1,\dots,n} \|c_i\|_2, \\ \hat{r}_i &\triangleq \sum_{\substack{j=1 \\ j \neq i}}^n |A_{(i,j)}|, & \hat{c}_i &\triangleq \sum_{\substack{j=1 \\ j \neq i}}^n |A_{(j,i)}|, \end{aligned}$$

and

$$\alpha \triangleq \min_{i=1,\dots,n} (|A_{(i,i)}| - \hat{r}_i), \quad \beta \triangleq \min_{i=1,\dots,n} (|A_{(i,i)}| - \hat{c}_i).$$

Then, the following statements hold:

i) If $\alpha > 0$, then A is nonsingular and

$$\|A^{-1}\|_{\text{row}} < 1/\alpha.$$

ii) If $\beta > 0$, then A is nonsingular and

$$\|A^{-1}\|_{\text{col}} < 1/\beta.$$

iii) If $\alpha > 0$ and $\beta > 0$, then A is nonsingular, and

$$\sqrt{\alpha\beta} \leq \sigma_{\min}(A).$$

iv) $\sigma_{\min}(A)$ satisfies

$$\min_{i=1,\dots,n} \frac{1}{2} [2|A_{(i,i)}| - \hat{r}_i - \hat{c}_i] \leq \sigma_{\min}(A).$$

v) $\sigma_{\min}(A)$ satisfies

$$\min_{i=1,\dots,n} \frac{1}{2} \left[(4|A_{(i,i)}|^2 + [\hat{r}_i - \hat{c}_i]^2)^{1/2} - \hat{r}_i - \hat{c}_i \right] \leq \sigma_{\min}(A).$$

vi) $\sigma_{\min}(A)$ satisfies

$$\left(\frac{n-1}{n}\right)^{(n-1)/2} |\det A| \max \left\{ \frac{c_{\min}}{\prod_{i=1}^n c_i}, \frac{r_{\min}}{\prod_{i=1}^n r_i} \right\} \leq \sigma_{\min}(A).$$

(Proof: See Fact 9.8.23, [711, pp. 227, 231], and [707, 763, 1367].)

Fact 9.13.22. Let $A \in \mathbb{F}^{n \times n}$, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_n$ are ordered such that $|\lambda_n| \leq \dots \leq |\lambda_1|$. Then, for all $i = 1, \dots, n$,

$$\lim_{k \rightarrow \infty} \sigma_i^{1/k}(A^k) = |\lambda_i|.$$

In particular,

$$\lim_{k \rightarrow \infty} \sigma_{\max}^{1/k}(A^k) = \text{sprad}(A).$$

(Proof: See [711, p. 180].) (Remark: This identity is due to Yamamoto.) (Remark: The expression for $\text{sprad}(A)$ is a special case of Proposition 9.2.6.)

Fact 9.13.23. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonzero. Then,

$$\frac{1}{\sigma_{\max}(A)} = \min_{B \in \{X \in \mathbb{F}^{n \times n}: \det(I - AX) = 0\}} \sigma_{\max}(B).$$

Furthermore, there exists $B_0 \in \mathbb{F}^{n \times n}$ such that $\text{rank } B_0 = 1$, $\det(I - AB_0) = 0$, and

$$\frac{1}{\sigma_{\max}(A)} = \sigma_{\max}(B_0).$$

(Proof: If $\sigma_{\max}(B) < 1/\sigma_{\max}(A)$, then $\text{sprad}(AB) \leq \sigma_{\max}(AB) < 1$, and thus $I - AB$ is nonsingular. Hence,

$$\begin{aligned} \frac{1}{\sigma_{\max}(A)} &= \min_{B \in \{X \in \mathbb{F}^{n \times n} : \sigma_{\max}(X) \geq 1/\sigma_{\max}(A)\}} \sigma_{\max}(B) \\ &= \min_{B \in \{X \in \mathbb{F}^{n \times n} : \sigma_{\max}(X) < 1/\sigma_{\max}(A)\}^c} \sigma_{\max}(B) \\ &\leq \min_{B \in \{X \in \mathbb{F}^{n \times n} : \det(I - AX) = 0\}} \sigma_{\max}(B). \end{aligned}$$

Using the singular value decomposition, equality holds by constructing B_0 to have rank 1 and singular value $1/\sigma_{\max}(A)$. (Remark: This result is related to the *small-gain theorem*. See [1498, pp. 276, 277].)

9.14 Facts on Matrix Norms and Singular Values for Two or More Matrices

Fact 9.14.1. Let $a_1, \dots, a_n \in \mathbb{F}^n$ be linearly independent, and, for all $i = 1, \dots, n$, define

$$A_i \triangleq I - (a_i^* a_i)^{-1} a_i a_i^*.$$

Then,

$$\sigma_{\max}(A_n A_{n-1} \cdots A_1) < 1.$$

(Proof: Define $A \triangleq A_n A_{n-1} \cdots A_1$. Since $\sigma_{\max}(A_i) \leq 1$ for all $i = 1, \dots, n$, it follows that $\sigma_{\max}(A) \leq 1$. Suppose that $\sigma_{\max}(A) = 1$, and let $x \in \mathbb{F}^n$ satisfy $x^* x = 1$ and $\|Ax\|_2 = 1$. Then, for all $i = 1, \dots, n$, $\|A_i A_{i-1} \cdots A_1 x\|_2 = 1$. Consequently, $\|A_1 x\|_2 = 1$, which implies that $a_1^* x = 0$, and thus $A_1 x = x$. Hence, $\|A_i A_{i-1} \cdots A_2 x\|_2 = 1$. Repeating this argument implies that, for all $i = 1, \dots, n$, $a_i^* x = 0$. Since $a_1, \dots, a_n$ are linearly independent, it follows that $x = 0$, which is a contradiction.) (Remark: This result is due to Akers and Djokovic.)

Fact 9.14.2. Let $A_1, \dots, A_n \in \mathbb{F}^{n \times n}$, assume that, for all $i, j = 1, \dots, n$, $[A_i, A_j] = 0$, and assume that, for all $i = 1, \dots, n$, $\sigma_{\max}(A_i) = 1$ and $\text{sprad}(A_i) = 1$. Then,

$$\sigma_{\max}(A_n A_{n-1} \cdots A_1) < 1.$$

(Proof: See [1479].)

Fact 9.14.3. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then,

$$|\text{tr } AB| \leq \|AB\|_{\sigma_1} = \sum_{i=1}^r \sigma_i(AB) \leq \sum_{i=1}^r \sigma_i(A) \sigma_i(B).$$

(Proof: Use Proposition 9.6.2 and Fact 9.11.2.) (Remark: This result generalizes Fact 5.12.6.) (Remark: Sufficient conditions for equality are given in [1184, p. 107].)

Fact 9.14.4. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then,

$$|\operatorname{tr} AB| \leq \|AB\|_{\sigma 1} \leq \sigma_{\max}(A)\|B\|_{\sigma 1}.$$

(Proof: Use Corollary 9.3.8 and Fact 9.11.2.) (Remark: This result generalizes Fact 5.12.7.)

Fact 9.14.5. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{m \times n}$, and $p \in [1, \infty)$, and assume that AB is normal. Then,

$$\|AB\|_{\sigma p} \leq \|BA\|_{\sigma p}.$$

In particular,

$$\operatorname{tr} \langle AB \rangle \leq \operatorname{tr} \langle BA \rangle,$$

$$\|AB\|_F \leq \|BA\|_F,$$

$$\sigma_{\max}(AB) \leq \sigma_{\max}(BA).$$

(Proof: This result is due to Simon. See [246].)

Fact 9.14.6. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A is nonsingular, and assume that B is singular. Then,

$$\sigma_{\min}(A) \leq \sigma_{\max}(A - B).$$

Furthermore, if $\sigma_{\max}(A^{-1}) = \operatorname{sprad}(A^{-1})$, then there exists a singular matrix $C \in \mathbb{R}^{n \times n}$ such that $\sigma_{\max}(A - C) = \sigma_{\min}(A)$. (Proof: See [1098, p. 151].) (Remark: This result is due to Franck.)

Fact 9.14.7. Let $A \in \mathbb{C}^{n \times n}$, assume that A is nonsingular, let $\|\cdot\|$ and $\|\cdot\|'$ be norms on $\mathbb{C}^n$, let $\|\cdot\|''$ be the norm on $\mathbb{C}^{n \times n}$ induced by $\|\cdot\|$ and $\|\cdot\|'$, and let $\|\cdot\|'''$ be the norm on $\mathbb{C}^{n \times n}$ induced by $\|\cdot\|'$ and $\|\cdot\|$. Then,

$$\min\{\|B\|'' : B \in \mathbb{C}^{n \times n} \text{ and } A + B \text{ is nonsingular}\} = 1/\|A^{-1}\|'''.$$

In particular,

$$\min\{\|B\|_{\operatorname{col}} : B \in \mathbb{C}^{n \times n} \text{ and } A + B \text{ is singular}\} = 1/\|A^{-1}\|_{\operatorname{col}},$$

$$\min\{\sigma_{\max}(B) : B \in \mathbb{C}^{n \times n} \text{ and } A + B \text{ is singular}\} = \sigma_{\min}(A),$$

$$\min\{\|B\|_{\operatorname{row}} : B \in \mathbb{C}^{n \times n} \text{ and } A + B \text{ is singular}\} = 1/\|A^{-1}\|_{\operatorname{row}}.$$

(Proof: See [679] and [681, p. 111].) (Remark: This result is due to Gastinel. See [679].) (Remark: The result involving $\sigma_{\max}(B)$ is equivalent to the inequality in Fact 9.14.6.)

Fact 9.14.8. Let $A, B \in \mathbb{F}^{n \times m}$, and assume that $\operatorname{rank} A = \operatorname{rank} B$ and $\alpha \triangleq \sigma_{\max}(A^+) \sigma_{\max}(A - B) < 1$. Then,

$$\sigma_{\max}(B^+) < \frac{1}{1 - \alpha} \sigma_{\max}(A^+).$$

If, in addition, $n = m$, A and B are nonsingular, and $\sigma_{\max}(A - B) < \sigma_{\min}(A)$, then

$$\sigma_{\max}(B^{-1}) < \frac{\sigma_{\min}(A)}{\sigma_{\min}(A) - \sigma_{\max}(A - B)} \sigma_{\max}(A^{-1}).$$

(Proof: See [681, p. 400].)

Fact 9.14.9. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\sigma_{\max}(I - [A, B]) \geq 1.$$

(Proof: Since $\text{tr}[A, B] = 0$, it follows that there exists $\lambda \in \text{spec}(I - [A, B])$ such that $\text{Re } \lambda \geq 1$, and thus $|\lambda| \geq 1$. Hence, Corollary 9.4.5 implies that $\sigma_{\max}(I - [A, B]) \geq \text{sprad}(I - [A, B]) \geq |\lambda| \geq 1$.)

Fact 9.14.10. Let $A \in \mathbb{F}^{n \times m}$, and let $B \in \mathbb{F}^{k \times l}$ be a submatrix of A . Then, for all $i = 1, \dots, \min\{k, l\}$,

$$\sigma_i(B) \leq \sigma_i(A).$$

(Proof: Use Proposition 9.6.1.) (Remark: Sufficient conditions for singular value interlacing are given in [709, p. 419].)

Fact 9.14.11. Let

$$\mathcal{A} \triangleq \begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)},$$

assume that $\mathcal{A}$ is nonsingular, and define $\begin{bmatrix} E & F \\ G & H \end{bmatrix} \in \mathbb{F}^{(n+m) \times (n+m)}$ by

$$\begin{bmatrix} E & F \\ G & H \end{bmatrix} \triangleq \mathcal{A}^{-1}.$$

Then, the following statements hold:

i) For all $i = 1, \dots, \min\{n, m\} - 1$,

$$\frac{\sigma_{n-i}(A)}{\sigma_{\max}^2(\mathcal{A})} \leq \sigma_{m-i}(H) \leq \frac{\sigma_{n-i}(A)}{\sigma_{\min}^2(\mathcal{A})}.$$

ii) Assume that $n < m$. Then, for all $i = 1, \dots, m - n$,

$$\frac{1}{\sigma_{\max}(\mathcal{A})} \leq \sigma_i(H) \leq \frac{1}{\sigma_{\min}(\mathcal{A})}.$$

iii) Assume that $m < n$. Then, for all $i = 1, \dots, m - n$,

$$\sigma_{\min}(\mathcal{A}) \leq \sigma_i(H) \leq \sigma_{\max}(\mathcal{A}).$$

iv) Assume that $n = m$. Then, for all $i = 1, \dots, n$,

$$\frac{\sigma_i(A)}{\sigma_{\max}^2(\mathcal{A})} \leq \sigma_i(H) \leq \frac{\sigma_i(A)}{\sigma_{\min}^2(\mathcal{A})}.$$

v) Assume that $m < n$. Then,

$$\sigma_{\max}(H) \leq \frac{\sigma_{n-m+1}(A)}{\sigma_{\min}^2(\mathcal{A})}.$$

vi) Assume that $m < n$. Then, $H = 0$ if and only if $\text{def } A = m$.

Now, assume that $\mathcal{A}$ is unitary. Then, the following statements hold:

vii) If $n < m$, then

$$\sigma_i(D) = \begin{cases} 1, & 1 \leq i \leq m - n, \\ \sigma_{i-m+n}(A), & m - n < i \leq m. \end{cases}$$

viii) If $n = m$, then, for all $i = 1, \dots, n$,

$$\sigma_i(D) = \sigma_i(A).$$

ix) If $n \leq m$, then

$$|\det D| = \prod_{i=1}^m \sigma_i(D) = \prod_{i=1}^n \sigma_i(A) = |\det A|.$$

(Proof: See [575].) (Remark: Statement vi) is a special case of the nullity theorem given by Fact 2.11.20.) (Remark: Statement ix) follows from Fact 3.11.24 using Fact 5.11.28.)

Fact 9.14.12. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, $C \in \mathbb{F}^{k \times m}$, and $D \in \mathbb{F}^{k \times l}$. Then,

$$\sigma_{\max}\left(\begin{bmatrix} A & B \\ C & D \end{bmatrix}\right) \leq \sigma_{\max}\left(\begin{bmatrix} \sigma_{\max}(A) & \sigma_{\max}(B) \\ \sigma_{\max}(C) & \sigma_{\max}(D) \end{bmatrix}\right).$$

(Proof: See [719, 821].) (Remark: This result is due to Tomiyama.) (Remark: See Fact 8.18.28.)

Fact 9.14.13. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, and $C \in \mathbb{F}^{k \times m}$. Then, for all $X \in \mathbb{F}^{k \times l}$,

$$\max\left\{\sigma_{\max}\left(\begin{bmatrix} A & B \end{bmatrix}\right), \sigma_{\max}\left(\begin{bmatrix} A \\ C \end{bmatrix}\right)\right\} \leq \sigma_{\max}\left(\begin{bmatrix} A & B \\ C & X \end{bmatrix}\right).$$

Furthermore, there exists a matrix $X \in \mathbb{F}^{k \times l}$ such that equality holds. (Remark: This result is *Parrott's theorem*. See [366], [447, pp. 271, 272], and [1498, pp. 40–42].)

Fact 9.14.14. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{n \times l}$. Then,

$$\begin{aligned} \max\{\sigma_{\max}(A), \sigma_{\max}(B)\} &\leq \sigma_{\max}\left(\begin{bmatrix} A & B \end{bmatrix}\right) \\ &\leq [\sigma_{\max}^2(A) + \sigma_{\max}^2(B)]^{1/2} \\ &\leq \sqrt{2} \max\{\sigma_{\max}(A), \sigma_{\max}(B)\} \end{aligned}$$

and, if $n \leq \min\{m, l\}$,

$$[\sigma_n^2(A) + \sigma_n^2(B)]^{1/2} \leq \sigma_n\left(\begin{bmatrix} A & B \end{bmatrix}\right) \leq \begin{cases} [\sigma_n^2(A) + \sigma_{\max}^2(B)]^{1/2} \\ [\sigma_{\max}^2(A) + \sigma_n^2(B)]^{1/2}. \end{cases}$$

(Problem: Obtain analogous bounds for $\sigma_i(\begin{bmatrix} A & B \end{bmatrix})$.)

Fact 9.14.15. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} & \sigma_{\max}(A + B) \\ & \leq \frac{1}{2} \left[\sigma_{\max}(A) + \sigma_{\max}(B) \right. \\ & \quad \left. + \sqrt{[\sigma_{\max}(A) - \sigma_{\max}(B)]^2 + 4 \max\{\sigma_{\max}^2(\langle A \rangle^{1/2} \langle B \rangle^{1/2}), \sigma_{\max}^2(\langle A^* \rangle^{1/2} \langle B^* \rangle^{1/2})\}} \right] \\ & \leq \sigma_{\max}(A) + \sigma_{\max}(B). \end{aligned}$$

(Proof: See [821].) (Remark: See Fact 8.18.14.) (Remark: This result interpolates the triangle inequality for the maximum singular value.)

Fact 9.14.16. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\alpha > 0$. Then,

$$\sigma_{\max}(A + B) \leq [(1 + \alpha)\sigma_{\max}^2(A) + (1 + \alpha^{-1})\sigma_{\max}^2(B)]^{1/2}$$

and

$$\sigma_{\min}(A + B) \leq [(1 + \alpha)\sigma_{\min}^2(A) + (1 + \alpha^{-1})\sigma_{\max}^2(B)]^{1/2}.$$

Fact 9.14.17. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} \sigma_{\min}(A) - \sigma_{\max}(B) & \leq |\det(A + B)|^{1/n} \\ & \leq \prod_{i=1}^n |\sigma_i(A) + \sigma_{n-i+1}(B)|^{1/n} \\ & \leq \sigma_{\max}(A) + \sigma_{\max}(B). \end{aligned}$$

(Proof: See [721, p. 63] and [894].)

Fact 9.14.18. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $\sigma_{\max}(B) \leq \sigma_{\min}(A)$. Then,

$$\begin{aligned} 0 & \leq [\sigma_{\min}(A) - \sigma_{\max}(B)]^n \\ & \leq \prod_{i=1}^n |\sigma_i(A) - \sigma_{n-i+1}(B)| \\ & \leq |\det(A + B)| \\ & \leq \prod_{i=1}^n |\sigma_i(A) + \sigma_{n-i+1}(B)| \\ & \leq [\sigma_{\max}(A) + \sigma_{\max}(B)]^n. \end{aligned}$$

Hence, if $\sigma_{\max}(B) < \sigma_{\min}(A)$, then A is nonsingular and $A + \alpha B$ is nonsingular for all $-1 \leq \alpha \leq 1$. (Proof: See [894].) (Remark: See Fact 11.18.16.) (Remark: See Fact 5.12.12.)

Fact 9.14.19. Let $A, B \in \mathbb{F}^{n \times m}$. Then, the following statements are equivalent:

i) For all $k = 1, \dots, \min\{n, m\}$,

$$\sum_{i=1}^k \sigma_i(A) \leq \sum_{i=1}^k \sigma_i(B).$$

ii) For all unitarily invariant norms $\|\cdot\|$ on $\mathbb{F}^{n \times m}$, $\|A\| \leq \|B\|$.

(Proof: See [711, pp. 205, 206].) (Remark: This result is the *Fan dominance theorem*.)

Fact 9.14.20. Let $A, B \in \mathbb{F}^{n \times m}$. Then, for all $k = 1, \dots, \min\{n, m\}$,

$$\sum_{i=1}^k [\sigma_i(A) + \sigma_{\min\{n, m\}+1-i}(B)] \leq \sum_{i=1}^k \sigma_i(A+B) \leq \sum_{i=1}^k [\sigma_i(A) + \sigma_i(B)].$$

Furthermore, if either $\sigma_{\max}(A) < \sigma_{\min}(B)$ or $\sigma_{\max}(B) < \sigma_{\min}(A)$, then, for all $k = 1, \dots, \min\{n, m\}$,

$$\sum_{i=1}^k \sigma_i(A+B) \leq \sum_{i=1}^k |\sigma_i(A) - \sigma_{\min\{n, m\}+1-i}(B)|.$$

(Proof: See Proposition 9.2.2, [711, pp. 196, 197] and [894].)

Fact 9.14.21. Let $A, B \in \mathbb{F}^{n \times m}$, and let $\alpha \in [0, 1]$. Then, for all $i = 1, \dots, \min\{n, m\}$,

$$\sigma_i[\alpha A + (1 - \alpha)B] \leq \begin{cases} \sigma_i \left(\begin{bmatrix} A & 0 \\ 0 & B \end{bmatrix} \right) \\ \sigma_i \left(\begin{bmatrix} \sqrt{2\alpha}A & 0 \\ 0 & \sqrt{2(1-\alpha)}B \end{bmatrix} \right), \end{cases}$$

and

$$2\sigma_i(AB^*) \leq \sigma_i(\langle A \rangle^2 + \langle B \rangle^2).$$

Furthermore,

$$\langle \alpha A + (1 - \alpha)B \rangle^2 \leq \alpha \langle A \rangle^2 + (1 - \alpha) \langle B \rangle^2.$$

If, in addition, $n = m$, then, for all $i = 1, \dots, n$,

$$\frac{1}{2}\sigma_i(A + A^*) \leq \sigma_i \left(\begin{bmatrix} A & 0 \\ 0 & A \end{bmatrix} \right).$$

(Proof: See [698].) (Remark: See Fact 9.14.23.)

Fact 9.14.22. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times m}$, and let $p, q > 1$ satisfy $1/p + 1/q = 1$. Then, for all $i = 1, \dots, \min\{n, m, l\}$,

$$\sigma_i(AB^*) \leq \sigma_i \left(\frac{1}{p} \langle A \rangle^p + \frac{1}{q} \langle B \rangle^q \right).$$

Equivalently, there exists a unitary matrix $S \in \mathbb{F}^{m \times m}$ such that

$$\langle AB^* \rangle^{1/2} \leq S^* \left(\frac{1}{p} \langle A \rangle^p + \frac{1}{q} \langle B \rangle^q \right) S.$$

(Proof: See [47, 49, 694] or [1485, p. 28].) (Remark: This result is a matrix version of Young's inequality. See Fact 1.10.32.)

Fact 9.14.23. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times m}$. Then, for all $i = 1, \dots, \min\{n, m, l\}$,

$$\sigma_i(AB^*) \leq \frac{1}{2}\sigma_i(A^*A + B^*B).$$

(Proof: Set $p = q = 2$ in Fact 9.14.22. See [209].) (Remark: See Fact 9.9.47 and Fact 9.14.21.)

Fact 9.14.24. Let $A, B, C, D \in \mathbb{F}^{n \times m}$. Then, for all $i = 1, \dots, \min\{n, m\}$,

$$\sqrt{2}\sigma_i(\langle AB^* + CD^* \rangle) \leq \sigma_i\left(\begin{bmatrix} A & B \\ C & D \end{bmatrix}\right).$$

(Proof: See [693].)

Fact 9.14.25. Let $A, B, C, D, X \in \mathbb{F}^{n \times n}$, assume that A, B, C, D are positive semidefinite, and assume that $0 \leq A \leq C$ and $0 \leq B \leq D$. Then, for all $i = 1, \dots, n$,

$$\sigma_i(A^{1/2}XB^{1/2}) \leq \sigma_i(C^{1/2}XD^{1/2}).$$

(Proof: See [698, 816].)

Fact 9.14.26. Let $A_1, \dots, A_k \in \mathbb{F}^{n \times n}$, and let $l \in \{1, \dots, n\}$. Then,

$$\sum_{i=1}^l \sigma_i\left(\prod_{j=1}^k A_j\right) \leq \sum_{i=1}^l \prod_{j=1}^k \sigma_i(A_j).$$

(Proof: See [317].) (Remark: This result is a weak majorization relation.)

Fact 9.14.27. Let $A, B \in \mathbb{F}^{n \times m}$, and let $1 \leq l_1 < \dots < l_k \leq \min\{n, m\}$. Then,

$$\sum_{i=1}^k \sigma_{l_i}(A) \sigma_{n-i+1}(B) \leq \sum_{i=1}^k \sigma_{l_i}(AB) \leq \sum_{i=1}^k \sigma_{l_i}(A) \sigma_i(B)$$

and

$$\sum_{i=1}^k \sigma_{l_i}(A) \sigma_{n-l_i+1}(B) \leq \sum_{i=1}^k \sigma_i(AB).$$

In particular,

$$\sum_{i=1}^k \sigma_i(A) \sigma_{n-i+1}(B) \leq \sum_{i=1}^k \sigma_i(AB) \leq \sum_{i=1}^k \sigma_i(A) \sigma_i(B).$$

Furthermore,

$$\prod_{i=1}^k \sigma_{l_i}(AB) \leq \prod_{i=1}^k \sigma_{l_i}(A) \sigma_i(B)$$

with equality for $k = n$. Furthermore,

$$\prod_{i=1}^k \sigma_{l_i}(A) \sigma_{n-l_i+1}(B) \leq \prod_{i=1}^k \sigma_i(AB)$$

with equality for $k = n$. In particular,

$$\prod_{i=1}^k \sigma_i(A) \sigma_{n-i+1}(B) \leq \prod_{i=1}^k \sigma_i(AB) \leq \prod_{i=1}^k \sigma_i(A) \sigma_i(B)$$

with equality for $k = n$. (Proof: See [1388].) (Remark: See Fact 8.18.19 and Fact 8.18.22.) (Remark: The left-hand inequalities in the first and third strings are conjectures. See [1388].)

Fact 9.14.28. Let $A \in \mathbb{F}^{n \times m}$, let $k \geq 1$ satisfy $k < \text{rank } A$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times m}$. Then,

$$\min_{B \in \{X \in \mathbb{F}^{n \times n}: \text{rank } X \leq k\}} \|A - B\| = \|A - B_0\|,$$

where B_0 is formed by replacing $(\text{rank } A) - k$ smallest positive singular values in the singular value decomposition of A by 0's. Furthermore,

$$\sigma_{\max}(A - B_0) = \sigma_{k+1}(A)$$

and

$$\|A - B_0\|_F = \sqrt{\sum_{i=k+1}^r \sigma_i^2(A)}.$$

Furthermore, B_0 is the unique solution if and only if $\sigma_{k+1}(A) < \sigma_k(A)$. (Proof: The result follows from Fact 9.14.29 with $B_\sigma \triangleq \text{diag}[\sigma_1(A), \dots, \sigma_k(A), 0_{(n-k) \times (m-k)}]$, $S_1 = I_n$, and $S_2 = I_m$. See [569] and [1230, p. 208].) (Remark: This result is known as the *Schmidt-Mirsky theorem*. For the case of the Frobenius norm, the result is known as the *Eckart-Young theorem*. See [507] and [1230, p. 210].) (Remark: See Fact 9.15.4.)

Fact 9.14.29. Let $A, B \in \mathbb{F}^{n \times m}$, define $A_\sigma, B_\sigma \in \mathbb{F}^{n \times m}$ by

$$A_\sigma \triangleq \begin{bmatrix} \sigma_1(A) & & & \\ & \ddots & & \\ & & \sigma_r(A) & \\ & & & 0_{(n-r) \times (m-r)} \end{bmatrix},$$

where $r \triangleq \text{rank } A$, and

$$B_\sigma \triangleq \begin{bmatrix} \sigma_1(B) & & & \\ & \ddots & & \\ & & \sigma_l(B) & \\ & & & 0_{(n-l) \times (m-l)} \end{bmatrix},$$

where $l \triangleq \text{rank } B$, let $S_1 \in \mathbb{F}^{n \times n}$ and $S_2 \in \mathbb{F}^{m \times m}$ be unitary matrices, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times m}$. Then,

$$\|A_\sigma - B_\sigma\| \leq \|A - S_1 B S_2\| \leq \|A_\sigma + B_\sigma\|.$$

In particular,

$$\max_{i \in \{1, \dots, \max\{r, l\}\}} |\sigma_i(A) - \sigma_i(B)| \leq \sigma_{\max}(A - B) \leq \sigma_{\max}(A) + \sigma_{\max}(B).$$

(Proof: See [1390].) (Remark: In the case $S_1 = I_n$ and $S_2 = I_m$, the left-hand inequality is *Mirsky's theorem*. See [1230, p. 204].) (Remark: See Fact 9.12.4.)

Fact 9.14.30. Let $A, B \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = \text{rank } B$. Then,

$$\sigma_{\max}[AA^+(I - BB^+)] = \sigma_{\max}[BB^+(I - AA^+)]$$

$$\leq \min\{\sigma_{\max}(A^+), \sigma_{\max}(B^+)\}\sigma_{\max}(A - B).$$

(Proof: See [681, p. 400] and [1230, p. 141].)

Fact 9.14.31. Let $A, B \in \mathbb{F}^{n \times m}$. Then, for all $k = 1, \dots, \min\{n, m\}$,

$$\begin{aligned} \sum_{i=1}^k \sigma_i(A \circ B) &\leq \sum_{i=1}^k d_i^{1/2}(A^*A) d_i^{1/2}(BB^*) \\ &\leq \left\{ \begin{array}{l} \sum_{i=1}^k d_i^{1/2}(A^*A) \sigma_i(B) \\ \sum_{i=1}^k \sigma_i(A) d_i^{1/2}(BB^*) \end{array} \right\} \\ &\leq \sum_{i=1}^k \sigma_i(A) \sigma_i(B) \end{aligned}$$

and

$$\begin{aligned} \sum_{i=1}^k \sigma_i(A \circ B) &\leq \sum_{i=1}^k d_i^{1/2}(AA^*) d_i^{1/2}(B^*B) \\ &\leq \left\{ \begin{array}{l} \sum_{i=1}^k d_i^{1/2}(AA^*) \sigma_i(B) \\ \sum_{i=1}^k \sigma_i(A) d_i^{1/2}(B^*B) \end{array} \right\} \\ &\leq \sum_{i=1}^k \sigma_i(A) \sigma_i(B). \end{aligned}$$

In particular,

$$\sigma_{\max}(A \circ B) \leq \|A\|_{2,1} \|B\|_{\infty,2} \leq \left\{ \begin{array}{l} \|A\|_{2,1} \sigma_{\max}(B) \\ \sigma_{\max}(A) \|B\|_{\infty,2} \end{array} \right\} \leq \sigma_{\max}(A) \sigma_{\max}(B)$$

and

$$\sigma_{\max}(A \circ B) \leq \|A\|_{\infty,2} \|B\|_{2,1} \leq \left\{ \begin{array}{l} \|A\|_{\infty,2} \sigma_{\max}(B) \\ \sigma_{\max}(A) \|B\|_{2,1} \end{array} \right\} \leq \sigma_{\max}(A) \sigma_{\max}(B).$$

(Proof: See [56, 976, 1481] and [711, p. 334], and use Fact 2.21.2, Fact 8.17.8, and Fact 9.8.24.) (Remark: $d_i^{1/2}(A^*A)$ and $d_i^{1/2}(AA^*)$ are the i th largest Euclidean norms of the columns and rows of A , respectively.) (Remark: For related results, see [1345].) (Remark: The case of equality is discussed in [319].)

Fact 9.14.32. Let $A, B \in \mathbb{C}^{n \times m}$. Then,

$$\begin{aligned} \sum_{i=1}^n \sigma_i^2(A \circ B) &= \operatorname{tr}(A \circ B)(\overline{A \circ B})^T \\ &= \operatorname{tr}(A \circ \overline{A})(B \circ \overline{B})^T \\ &\leq \sum_{i=1}^n \sigma_i[(A \circ \overline{A})(B \circ \overline{B})^T] \\ &\leq \sum_{i=1}^n \sigma_i(A \circ \overline{A})\sigma_i(B \circ \overline{B}). \end{aligned}$$

(Proof: See [730].)

Fact 9.14.33. Let $A, B \in \mathbb{F}^{n \times m}$. Then,

$$\sigma_{\max}(A \circ B) \leq \sqrt{n}\|A\|_{\infty}\sigma_{\max}(B).$$

Now, assume that $n = m$ and that either A is positive semidefinite and B is Hermitian or A and B are nonnegative and symmetric. Then,

$$\sigma_{\max}(A \circ B) \leq \|A\|_{\infty}\sigma_{\max}(B).$$

Next, assume that A and B are real, let β denote the smallest positive entry of $|B|$, and assume that B is symmetric and positive semidefinite. Then,

$$\operatorname{sprad}(A \circ B) \leq \frac{\|A\|_{\infty}\|B\|_{\infty}}{\beta}\sigma_{\max}(B)$$

and

$$\operatorname{sprad}(B) \leq \operatorname{sprad}(|B|) \leq \frac{\|B\|_{\infty}}{\beta}\operatorname{sprad}(B).$$

(Proof: See [1080].)

Fact 9.14.34. Let $A, B \in \mathbb{F}^{n \times m}$, and let $p \in [1, \infty)$ be an even integer. Then,

$$\|A \circ B\|_{\sigma p}^2 \leq \|A \circ \overline{A}\|_{\sigma p}\|B \circ \overline{B}\|_{\sigma p}.$$

In particular,

$$\|A \circ B\|_{\mathbb{F}}^2 \leq \|A \circ \overline{A}\|_{\mathbb{F}}\|B \circ \overline{B}\|_{\mathbb{F}}$$

and

$$\sigma_{\max}^2(A \circ B) \leq \sigma_{\max}(A \circ \overline{A})\sigma_{\max}(B \circ \overline{B}).$$

Equality holds if $B = \overline{A}$. Furthermore,

$$\|A \circ A\|_{\sigma p} \leq \|A \circ \overline{A}\|_{\sigma p}.$$

In particular,

$$\|A \circ A\|_{\mathbb{F}} \leq \|A \circ \overline{A}\|_{\mathbb{F}}$$

and

$$\sigma_{\max}(A \circ A) \leq \sigma_{\max}(A \circ \overline{A}).$$

Now, assume that $n = m$. Then,

$$\|A \circ A^T\|_{\sigma p} \leq \|A \circ \overline{A}\|_{\sigma p}.$$

In particular,

$$\|A \circ A^T\|_F \leq \|A \circ \overline{A}\|_F$$

and

$$\sigma_{\max}(A \circ A^T) \leq \sigma_{\max}(A \circ \overline{A}).$$

Finally,

$$\|A \circ A^*\|_{\sigma p} \leq \|A \circ \overline{A}\|_{\sigma p}.$$

In particular,

$$\|A \circ A^*\|_F \leq \|A \circ \overline{A}\|_F$$

and

$$\sigma_{\max}(A \circ A^*) \leq \sigma_{\max}(A \circ \overline{A}).$$

(Proof: See [712, 1193].) (Remark: See Fact 7.6.16.)

Fact 9.14.35. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A and B are nonnegative, and let $\alpha \in [0, 1]$. Then,

$$\sigma_{\max}(A^{\circ\alpha} \circ B^{\circ(1-\alpha)}) \leq \sigma_{\max}^\alpha(A) \sigma_{\max}^{1-\alpha}(B).$$

In particular,

$$\sigma_{\max}(A^{\circ 1/2} \circ B^{\circ 1/2}) \leq \sqrt{\sigma_{\max}(A) \sigma_{\max}(B)}.$$

Finally,

$$\sigma_{\max}(A^{\circ 1/2} \circ A^{\circ 1/2T}) \leq \sigma_{\max}(A^{\circ\alpha} \circ A^{\circ(1-\alpha)T}) \leq \sigma_{\max}(A).$$

(Proof: See [1193].) (Remark: See Fact 7.6.17.)

Fact 9.14.36. Let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{C}^{n \times n}$, and let $A, X, B \in \mathbb{C}^{n \times n}$. Then,

$$\|A \circ X \circ B\| \leq \frac{1}{2}\sqrt{n}\|A \circ X \circ \overline{A} + B \circ X \circ \overline{B}\|$$

and

$$\|A \circ X \circ B\|^2 \leq n\|A \circ X \circ \overline{A}\|\|B \circ X \circ \overline{B}\|.$$

Furthermore,

$$\|A \circ X \circ B\|_F \leq \frac{1}{2}\|A \circ X \circ \overline{A} + B \circ X \circ \overline{B}\|_F.$$

(Proof: See [730].)

Fact 9.14.37. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{l \times k}$, and $p \in [1, \infty]$. Then,

$$\|A \otimes B\|_{\sigma p} = \|A\|_{\sigma p} \|B\|_{\sigma p}.$$

In particular,

$$\sigma_{\max}(A \otimes B) = \sigma_{\max}(A) \sigma_{\max}(B)$$

and

$$\|A \otimes B\|_F = \|A\|_F \|B\|_F.$$

(Proof: See [690, p. 722].)

9.15 Facts on Least Squares

Fact 9.15.1. Let $A \in \mathbb{F}^{n \times m}$ and $b \in \mathbb{F}^n$, and define

$$f(x) \triangleq (Ax - b)^*(Ax - b) = \|Ax - b\|_2^2,$$

where $x \in \mathbb{F}^m$. Then, f has a minimizer. Furthermore, $x \in \mathbb{F}^m$ minimizes f if and only if there exists a vector $y \in \mathbb{F}^m$ such that

$$x = A^+b + (I - A^+A)y.$$

In this case,

$$f(x) = b^*(I - AA^+)b.$$

Furthermore, if $y \in \mathbb{F}^m$ is such that $(I - A^+A)y$ is nonzero, then

$$\|A^+b\|_2 < \|A^+b + (I - A^+A)y\|_2 = \sqrt{\|A^+b\|_2^2 + \|(I - A^+A)y\|_2^2}.$$

Finally, A^+b is the unique minimizer of f if and only if A is left invertible. (Remark: The minimization of f is the *least squares problem*. See [15, 226, 1226]. Note that, unlike Proposition 6.1.7, consistency is not assumed.) (Remark: This result is a special case of Fact 8.14.15.)

Fact 9.15.2. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times l}$, and define

$$f(X) \triangleq \text{tr}[(AX - B)^*(AX - B)] = \|AX - B\|_F^2,$$

where $X \in \mathbb{F}^{m \times l}$. Then, $X = A^+B$ minimizes f . (Problem: Determine all minimizers.) (Problem: Consider $f(X) = \text{tr}[(AX - B)^*C(AX - B)]$, where $C \in \mathbb{F}^{n \times n}$ is positive definite.)

Fact 9.15.3. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times m}$, and define

$$f(X) \triangleq \text{tr}[(XA - B)^*(XA - B)] = \|XA - B\|_F^2,$$

where $X \in \mathbb{F}^{l \times n}$. Then, $X = BA^+$ minimizes f .

Fact 9.15.4. Let $A \in \mathbb{F}^{n \times m}$, $B \in \mathbb{F}^{n \times p}$, and $C \in \mathbb{F}^{q \times m}$, and let $k \geq 1$ satisfy $k < \text{rank } A$. Then,

$$\min_{X \in \{Y \in \mathbb{F}^{p \times q} : \text{rank } Y \leq k\}} \|A - BXC\|_F = \|A - BX_0C\|_F,$$

where $X_0 = B^+SC^+$ and S is formed by replacing all but the k largest singular values in the singular value decomposition of BB^+AC^+C by 0's. Furthermore, X_0 is a solution that minimizes $\|X\|_F$. Finally, X_0 is the unique solution if and only if either $\text{rank } BB^+AC^+C \leq k$ or both $k \leq \text{rank } BB^+AC^+C$ and $\sigma_{k+1}(BB^+AC^+C) < \sigma_k(BB^+AC^+C)$. (Proof: See [507].) (Remark: This result generalizes Fact 9.14.28.)

Fact 9.15.5. Let $A, B \in \mathbb{F}^{n \times m}$, and define

$$f(X) \triangleq \text{tr}[(AX - B)^*(AX - B)] = \|AX - B\|_F^2,$$

where $X \in \mathbb{F}^{m \times m}$ is unitary. Then, $X = S_1S_2$ minimizes f , where $S_1 \begin{bmatrix} \hat{B} & 0 \\ 0 & 0 \end{bmatrix} S_2$ is the singular value decomposition of A^*B . (Proof: See [144, p. 224]. See also [971, pp. 269, 270].)

Fact 9.15.6. Let $A, B \in \mathbb{R}^{n \times n}$, and define

$$f(X_1, X_2) \triangleq \operatorname{tr}[(X_1 A X_2 - B)^T (X_1 A X_2 - B)] = \|X_1 A X_2 - B\|_F^2,$$

where $X_1, X_2 \in \mathbb{R}^{n \times n}$ are orthogonal. Then, $(X_1, X_2) = (V_2^T U_1^T, V_1^T U_2^T)$ minimizes f , where $U_1 \begin{bmatrix} A & 0 \\ 0 & 0 \end{bmatrix} V_1$ is the singular value decomposition of A and $U_2 \begin{bmatrix} B & 0 \\ 0 & 0 \end{bmatrix} V_2$ is the singular value decomposition of B . (Proof: See [971, p. 270].) (Remark: This result is due to Kristof.) (Remark: See Fact 3.9.5.) (Problem: Extend this result to $\mathbb{C}$ and nonsquare matrices.)

9.16 Notes

The equivalence of absolute and monotone norms given by Proposition 9.1.2 is due to [155]. More general monotonicity conditions are considered in [768]. Induced lower bounds are treated in [867, pp. 369, 370]. See also [1230, pp. 33, 80]. The induced norms (9.4.13) and (9.4.14) are given in [310] and [681, p. 116]. Alternative norms for the convolution operator are given in [310, 1435]. Proposition 9.3.6 is given in [1127, p. 97]. Norm-related topics are discussed in [169]. Spectral perturbation theory in finite and infinite dimensions is treated in [796], where the emphasis is on the regularity of the spectrum as a function of the perturbation rather than on bounds for finite perturbations.

Chapter Ten

Functions of Matrices and Their Derivatives

The norms discussed in Chapter 9 provide the foundation for the development in this chapter of some basic results in topology and analysis.

10.1 Open Sets and Closed Sets

Let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, let $x \in \mathbb{F}^n$, and let $\varepsilon > 0$. Then, define the *open ball of radius ε centered at x* by

$$\mathbb{B}_\varepsilon(x) \triangleq \{y \in \mathbb{F}^n: \|x - y\| < \varepsilon\} \quad (10.1.1)$$

and the *sphere of radius ε centered at x* by

$$\mathbb{S}_\varepsilon(x) \triangleq \{y \in \mathbb{F}^n: \|x - y\| = \varepsilon\}. \quad (10.1.2)$$

Definition 10.1.1. Let $\mathcal{S} \subseteq \mathbb{F}^n$. The vector $x \in \mathcal{S}$ is an *interior point* of $\mathcal{S}$ if there exists $\varepsilon > 0$ such that $\mathbb{B}_\varepsilon(x) \subseteq \mathcal{S}$. The *interior* of $\mathcal{S}$ is the set

$$\text{int } \mathcal{S} \triangleq \{x \in \mathcal{S}: x \text{ is an interior point of } \mathcal{S}\}. \quad (10.1.3)$$

Finally, $\mathcal{S}$ is *open* if every element of $\mathcal{S}$ is an interior point, that is, if $\mathcal{S} = \text{int } \mathcal{S}$.

Definition 10.1.2. Let $\mathcal{S} \subseteq \mathcal{S}' \subseteq \mathbb{F}^n$. The vector $x \in \mathcal{S}$ is an *interior point of $\mathcal{S}$ relative to $\mathcal{S}'$* if there exists $\varepsilon > 0$ such that $\mathbb{B}_\varepsilon(x) \cap \mathcal{S}' \subseteq \mathcal{S}$ or, equivalently, $\mathbb{B}_\varepsilon(x) \cap \mathcal{S} = \mathbb{B}_\varepsilon(x) \cap \mathcal{S}'$. The *interior of $\mathcal{S}$ relative to $\mathcal{S}'$* is the set

$$\text{int}_{\mathcal{S}'} \mathcal{S} \triangleq \{x \in \mathcal{S}: x \text{ is an interior point of } \mathcal{S} \text{ relative to } \mathcal{S}'\}. \quad (10.1.4)$$

Finally, $\mathcal{S}$ is *open relative to $\mathcal{S}'$* if $\mathcal{S} = \text{int}_{\mathcal{S}'} \mathcal{S}$.

Definition 10.1.3. Let $\mathcal{S} \subseteq \mathbb{F}^n$. The vector $x \in \mathbb{F}^n$ is a *closure point* of $\mathcal{S}$ if, for all $\varepsilon > 0$, the set $\mathcal{S} \cap \mathbb{B}_\varepsilon(x)$ is not empty. The *closure* of $\mathcal{S}$ is the set

$$\text{cl } \mathcal{S} \triangleq \{x \in \mathbb{F}^n: x \text{ is a closure point of } \mathcal{S}\}. \quad (10.1.5)$$

Finally, the set $\mathcal{S}$ is *closed* if every closure point of $\mathcal{S}$ is an element of $\mathcal{S}$, that is, if $\mathcal{S} = \text{cl } \mathcal{S}$.

Definition 10.1.4. Let $\mathcal{S} \subseteq \mathcal{S}' \subseteq \mathbb{F}^n$. The vector $x \in \mathcal{S}'$ is a *closure point* of $\mathcal{S}$ relative to $\mathcal{S}'$ if, for all $\varepsilon > 0$, the set $\mathcal{S} \cap \mathbb{B}_\varepsilon(x)$ is not empty. The *closure* of $\mathcal{S}$ relative to $\mathcal{S}'$ is the set

$$\text{cl}_{\mathcal{S}'} \mathcal{S} \triangleq \{x \in \mathbb{F}^n: x \text{ is a closure point of } \mathcal{S} \text{ relative to } \mathcal{S}'\}. \quad (10.1.6)$$

Finally, $\mathcal{S}$ is *closed relative* to $\mathcal{S}'$ if $\mathcal{S} = \text{cl}_{\mathcal{S}'} \mathcal{S}$.

It follows from Theorem 9.1.8 on the equivalence of norms on $\mathbb{F}^n$ that these definitions are independent of the norm assigned to $\mathbb{F}^n$.

Let $\mathcal{S} \subseteq \mathcal{S}' \subseteq \mathbb{F}^n$. Then,

$$\text{cl}_{\mathcal{S}'} \mathcal{S} = (\text{cl } \mathcal{S}) \cap \mathcal{S}', \quad (10.1.7)$$

$$\text{int}_{\mathcal{S}'} \mathcal{S} = \mathcal{S}' \setminus \text{cl}(\mathcal{S}' \setminus \mathcal{S}), \quad (10.1.8)$$

and

$$\text{int } \mathcal{S} \subseteq \text{int}_{\mathcal{S}'} \mathcal{S} \subseteq \mathcal{S} \subseteq \text{cl}_{\mathcal{S}'} \mathcal{S} \subseteq \text{cl } \mathcal{S}. \quad (10.1.9)$$

The set $\mathcal{S}$ is *solid* if $\text{int } \mathcal{S}$ is not empty, while $\mathcal{S}$ is *completely solid* if $\text{cl int } \mathcal{S} = \text{cl } \mathcal{S}$. If $\mathcal{S}$ is completely solid, then $\mathcal{S}$ is solid. The *boundary* of $\mathcal{S}$ is the set

$$\text{bd } \mathcal{S} \triangleq \text{cl } \mathcal{S} \setminus \text{int } \mathcal{S}, \quad (10.1.10)$$

while the *boundary* of $\mathcal{S}$ relative to $\mathcal{S}'$ is the set

$$\text{bd}_{\mathcal{S}'} \mathcal{S} \triangleq \text{cl}_{\mathcal{S}'} \mathcal{S} \setminus \text{int}_{\mathcal{S}'} \mathcal{S}. \quad (10.1.11)$$

Note that the empty set is both open and closed, although it is not solid.

The set $\mathcal{S} \subset \mathbb{F}^n$ is *bounded* if there exists $\delta > 0$ such that, for all $x, y \in \mathcal{S}$,

$$\|x - y\| < \delta. \quad (10.1.12)$$

The set $\mathcal{S} \subset \mathbb{F}^n$ is *compact* if it is both closed and bounded.

10.2 Limits

Definition 10.2.1. The *sequence* $(x_1, x_2, \dots)$ is a tuple with a countably infinite number of components. We write $(x_i)_{i=1}^\infty$ for $(x_1, x_2, \dots)$.

Definition 10.2.2. The sequence $(\alpha_i)_{i=1}^\infty \subset \mathbb{F}$ *converges* to $\alpha \in \mathbb{F}$ if, for all $\varepsilon > 0$, there exists a positive integer $p \in \mathbb{P}$ such that $|\alpha_i - \alpha| < \varepsilon$ for all $i > p$. In this case, we write $\alpha = \lim_{i \rightarrow \infty} \alpha_i$ or $\alpha_i \rightarrow \alpha$ as $i \rightarrow \infty$, where $i \in \mathbb{P}$. Finally, the sequence $(\alpha_i)_{i=1}^\infty \subset \mathbb{F}$ *converges* if there exists $\alpha \in \mathbb{F}$ such that $(\alpha_i)_{i=1}^\infty$ converges to α .

Definition 10.2.3. The sequence $(x_i)_{i=1}^\infty \subset \mathbb{F}^n$ *converges* to $x \in \mathbb{F}^n$ if $\lim_{i \rightarrow \infty} \|x - x_i\| = 0$, where $\|\cdot\|$ is a norm on $\mathbb{F}^n$. In this case, we write $x = \lim_{i \rightarrow \infty} x_i$ or $x_i \rightarrow x$ as $i \rightarrow \infty$, where $i \in \mathbb{P}$. The sequence $(x_i)_{i=1}^\infty \subset \mathbb{F}^n$ *converges* if there exists $x \in \mathbb{F}^n$ such that $(x_i)_{i=1}^\infty$ converges to x . Similarly, $(A_i)_{i=1}^\infty \subset \mathbb{F}^{n \times m}$ *converges* to $A \in \mathbb{F}^{n \times m}$ if $\lim_{i \rightarrow \infty} \|A - A_i\| = 0$, where $\|\cdot\|$ is a norm on $\mathbb{F}^{n \times m}$. In this case, we write $A = \lim_{i \rightarrow \infty} A_i$ or $A_i \rightarrow A$ as $i \rightarrow \infty$, where $i \in \mathbb{P}$. Finally, the sequence

$(A_i)_{i=1}^\infty \subset \mathbb{F}^{n \times m}$ converges if there exists $A \in \mathbb{F}^{n \times m}$ such that $(A_i)_{i=1}^\infty$ converges to A .

It follows from Theorem 9.1.8 that convergence of a sequence is independent of the choice of norm.

Proposition 10.2.4. Let $\mathcal{S} \subseteq \mathbb{F}^n$. The vector $x \in \mathbb{F}^n$ is a closure point of $\mathcal{S}$ if and only if there exists a sequence $(x_i)_{i=1}^\infty \subseteq \mathcal{S}$ that converges to x .

Proof. Suppose that $x \in \mathbb{F}^n$ is a closure point of $\mathcal{S}$. Then, for all $i \in \mathbb{P}$, there exists a vector $x_i \in \mathcal{S}$ such that $\|x - x_i\| < 1/i$. Hence, $x - x_i \rightarrow 0$ as $i \rightarrow \infty$. Conversely, suppose that $(x_i)_{i=1}^\infty \subseteq \mathcal{S}$ is such that $x_i \rightarrow x$ as $i \rightarrow \infty$, and let $\varepsilon > 0$. Then, there exists a positive integer $p \in \mathbb{P}$ such that $\|x - x_i\| < \varepsilon$ for all $i > p$. Therefore, $x_{p+1} \in \mathcal{S} \cap \mathbb{B}_\varepsilon(x)$, and thus $\mathcal{S} \cap \mathbb{B}_\varepsilon(x)$ is not empty. Hence, x is a closure point of $\mathcal{S}$. $\square$

Theorem 10.2.5. Let $\mathcal{S} \subset \mathbb{F}^n$ be compact, and let $(x_i)_{i=1}^\infty \subseteq \mathcal{S}$. Then, there exists a subsequence $\{x_{i_j}\}_{j=1}^\infty$ of $(x_i)_{i=1}^\infty$ such that $\{x_{i_j}\}_{j=1}^\infty$ converges and $\lim_{j \rightarrow \infty} x_{i_j} \in \mathcal{S}$.

Proof. See [1030, p. 145]. $\square$

Next, we define convergence for the series $\sum_{i=1}^\infty x_i$ in terms of the *partial sums* $\sum_{i=1}^k x_i$.

Definition 10.2.6. Let $(x_i)_{i=1}^\infty \subset \mathbb{F}^n$, and let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then, the series $\sum_{i=1}^\infty x_i$ converges if $\{\sum_{i=1}^k x_i\}_{k=1}^\infty$ converges. Furthermore, $\sum_{i=1}^\infty x_i$ converges absolutely if the series $\sum_{i=1}^\infty \|x_i\|$ converges.

Proposition 10.2.7. Let $(x_i)_{i=1}^\infty \subset \mathbb{F}^n$, and assume that the series $\sum_{i=1}^\infty x_i$ converges absolutely. Then, the series $\sum_{i=1}^\infty x_i$ converges.

Definition 10.2.8. Let $(A_i)_{i=1}^\infty \subset \mathbb{F}^{n \times m}$, and let $\|\cdot\|$ be a norm on $\mathbb{F}^{n \times m}$. Then, the series $\sum_{i=1}^\infty A_i$ converges if $\{\sum_{i=1}^k A_i\}_{k=1}^\infty$ converges. Furthermore, $\sum_{i=1}^\infty A_i$ converges absolutely if the series $\sum_{i=1}^\infty \|A_i\|$ converges.

Proposition 10.2.9. Let $(A_i)_{i=1}^\infty \subset \mathbb{F}^{n \times m}$, and assume that the series $\sum_{i=1}^\infty A_i$ converges absolutely. Then, the series $\sum_{i=1}^\infty A_i$ converges.

10.3 Continuity

Definition 10.3.1. Let $\mathcal{D} \subseteq \mathbb{F}^m$, $f: \mathcal{D} \rightarrow \mathbb{F}^n$, and $x \in \mathcal{D}$. Then, f is *continuous* at x if, for every convergent sequence $(x_i)_{i=1}^\infty \subseteq \mathcal{D}$ such that $\lim_{i \rightarrow \infty} x_i = x$, it follows that $\lim_{i \rightarrow \infty} f(x_i) = f(x)$. Furthermore, let $\mathcal{D}_0 \subseteq \mathcal{D}$. Then, f is *continuous* on $\mathcal{D}_0$ if f is continuous at x for all $x \in \mathcal{D}_0$. Finally, f is *continuous* if it is continuous on $\mathcal{D}$.

Theorem 10.3.2. Let $\mathcal{D} \subseteq \mathbb{F}^n$ be convex, and let $f: \mathcal{D} \rightarrow \mathbb{F}$ be convex. Then, f is continuous on $\text{int}_{\text{aff } \mathcal{D}} \mathcal{D}$.

Proof. See [157, p. 81] and [1133, p. 82]. $\square$

Corollary 10.3.3. Let $A \in \mathbb{F}^{n \times m}$, and define $f: \mathbb{F}^m \rightarrow \mathbb{F}^n$ by $f(x) \triangleq Ax$. Then, f is continuous.

Proof. The result is a consequence of Theorem 10.3.2. Alternatively, let $x \in \mathbb{F}^m$, and let $(x_i)_{i=1}^\infty \subset \mathbb{F}^m$ be such that $x_i \rightarrow x$ as $i \rightarrow \infty$. Furthermore, let $\|\cdot\|$ and $\|\cdot\|'$ be compatible norms on $\mathbb{F}^m$ and $\mathbb{F}^{m \times n}$, respectively. Since $\|Ax - Ax_i\| \leq \|A\|' \|x - x_i\|$, it follows that $Ax_i \rightarrow Ax$ as $i \rightarrow \infty$. $\square$

Theorem 10.3.4. Let $\mathcal{D} \subseteq \mathbb{F}^m$, and let $f: \mathcal{D} \mapsto \mathbb{F}^n$. Then, the following statements are equivalent:

- i) f is continuous.
- ii) For all open $\mathcal{S} \subseteq \mathbb{F}^n$, the set $f^{-1}(\mathcal{S})$ is open relative to $\mathcal{D}$.
- iii) For all closed $\mathcal{S} \subseteq \mathbb{F}^n$, the set $f^{-1}(\mathcal{S})$ is closed relative to $\mathcal{D}$.

Proof. See [1030, pp. 87, 110]. $\square$

Corollary 10.3.5. Let $A \in \mathbb{F}^{n \times m}$ and $\mathcal{S} \subseteq \mathbb{F}^n$, and define $\mathcal{S}' \triangleq \{x \in \mathbb{F}^m: Ax \in \mathcal{S}\}$. If $\mathcal{S}$ is open, then $\mathcal{S}'$ is open. If $\mathcal{S}$ is closed, then $\mathcal{S}'$ is closed.

The following result is the *open mapping theorem*.

Theorem 10.3.6. Let $\mathcal{D} \subseteq \mathbb{F}^m$, let $A \in \mathbb{F}^{n \times m}$, assume that $\mathcal{D}$ is open, and assume that A is right invertible. Then, $A\mathcal{D}$ is open.

The following result is the *invariance of domain*.

Theorem 10.3.7. Let $\mathcal{D} \subseteq \mathbb{F}^n$, let $f: \mathcal{D} \mapsto \mathbb{F}^n$, assume that $\mathcal{D}$ is open, and assume that f is continuous and one-to-one. Then, $f(\mathcal{D})$ is open.

Proof. See [1217, p. 3]. $\square$

Theorem 10.3.8. Let $\mathcal{D} \subset \mathbb{F}^m$ be compact, and let $f: \mathcal{D} \mapsto \mathbb{F}^n$ be continuous. Then, $f(\mathcal{D})$ is compact.

Proof. See [1030, p. 146]. $\square$

The following corollary of Theorem 10.3.8 shows that a continuous real-valued function defined on a compact set has a minimizer.

Corollary 10.3.9. Let $\mathcal{D} \subset \mathbb{F}^m$ be compact, and let $f: \mathcal{D} \mapsto \mathbb{R}$ be continuous. Then, there exists $x_0 \in \mathcal{D}$ such that $f(x_0) \leq f(x)$ for all $x \in \mathcal{D}$.

The following result is the *Schauder fixed-point theorem*.

Theorem 10.3.10. Let $\mathcal{D} \subseteq \mathbb{F}^m$, assume that $\mathcal{D}$ is nonempty, closed, and convex, let $f: \mathcal{D} \rightarrow \mathcal{D}$, assume that f is continuous, and assume that $f(\mathcal{D})$ is bounded. Then, there exists $x \in \mathcal{D}$ such that $f(x) = x$.

Proof. See [1404, p. 167]. □

The following corollary for the case of a bounded domain is the *Brouwer fixed-point theorem*.

Corollary 10.3.11. Let $\mathcal{D} \subseteq \mathbb{F}^m$, assume that $\mathcal{D}$ is nonempty, compact, and convex, let $f: \mathcal{D} \rightarrow \mathcal{D}$, and assume that f is continuous. Then, there exists $x \in \mathcal{D}$ such that $f(x) = x$.

Proof. See [1404, p. 163]. □

Definition 10.3.12. Let $\mathcal{S} \subseteq \mathbb{F}^{n \times n}$. Then, $\mathcal{S}$ is *pathwise connected* if, for all $B_1, B_2 \in \mathcal{S}$, there exists a continuous function $f: [0, 1] \mapsto \mathcal{S}$ such that $f(0) = B_1$ and $f(1) = B_2$.

10.4 Derivatives

Let $\mathcal{D} \subseteq \mathbb{F}^m$, and let $x_0 \in \mathcal{D}$. Then, the *variational cone of $\mathcal{D}$ with respect to x_0* is the set

$$\text{vcone}(\mathcal{D}, x_0) \triangleq \{\xi \in \mathbb{F}^m: \text{ there exists } \alpha_0 > 0 \text{ such that } x_0 + \alpha\xi \in \mathcal{D}, \alpha \in [0, \alpha_0]\}. \quad (10.4.1)$$

Note that $\text{vcone}(\mathcal{D}, x_0)$ is a pointed cone, although it may consist of only the origin as can be seen from the example $x_0 = 0$ and

$$\mathcal{D} = \left\{x \in \mathbb{R}^2: 0 \leq x_{(1)} \leq 1, x_{(1)}^3 \leq x_{(2)} \leq x_{(1)}^2\right\}.$$

Now, let $\mathcal{D} \subseteq \mathbb{F}^m$ and $f: \mathcal{D} \rightarrow \mathbb{F}^n$. If $\xi \in \text{vcone}(\mathcal{D}, x_0)$, then the *one-sided directional differential of f at x_0 in the direction ξ* is defined by

$$D_+f(x_0; \xi) \triangleq \lim_{\alpha \downarrow 0} \frac{1}{\alpha} [f(x_0 + \alpha\xi) - f(x_0)] \quad (10.4.2)$$

if the limit exists. Similarly, if $\xi \in \text{vcone}(\mathcal{D}, x_0)$ and $-\xi \in \text{vcone}(\mathcal{D}, x_0)$, then the *two-sided directional differential $Df(x_0; \xi)$ of f at x_0 in the direction ξ* is defined by

$$Df(x_0; \xi) \triangleq \lim_{\alpha \rightarrow 0} \frac{1}{\alpha} [f(x_0 + \alpha\xi) - f(x_0)] \quad (10.4.3)$$

if the limit exists. If $\xi = e_i$ so that the direction ξ is one of the coordinate axes, then the *partial derivative of f with respect to $x_{(i)}$ at x_0* , denoted by $\frac{\partial f(x_0)}{\partial x_{(i)}}$, is given by

$$\frac{\partial f(x_0)}{\partial x_{(i)}} \triangleq \lim_{\alpha \rightarrow 0} \frac{1}{\alpha} [f(x_0 + \alpha e_i) - f(x_0)], \quad (10.4.4)$$

that is,

$$\frac{\partial f(x_0)}{\partial x_{(i)}} = Df(x_0; e_i), \quad (10.4.5)$$

when the two-sided directional differential $Df(x_0; e_i)$ exists.

Proposition 10.4.1. Let $\mathcal{D} \subseteq \mathbb{F}^m$ be a convex set, let $f: \mathcal{D} \rightarrow \mathbb{F}^n$ be convex, and let $x_0 \in \text{int } \mathcal{D}$. Then, $D_+f(x_0; \xi)$ exists for all $\xi \in \mathbb{F}^m$.

Proof. See [157, p. 83]. □

Note that $D_+f(x_0; \xi) = \pm\infty$ is possible if x_0 is an element of the boundary of $\mathcal{D}$. For example, consider the continuous function $f: [0, \infty) \mapsto \mathbb{R}$ given by $f(x) = 1 - \sqrt{x}$. In this case, $D_+f(x_0; \xi) = -\infty$ and thus does not exist.

Next, we consider a stronger form of differentiation.

Proposition 10.4.2. Let $\mathcal{D} \subseteq \mathbb{F}^m$ be solid and convex, let $f: \mathcal{D} \rightarrow \mathbb{F}^n$, and let $x_0 \in \mathcal{D}$. Then, there exists at most one matrix $F \in \mathbb{F}^{n \times m}$ satisfying

$$\lim_{\substack{x \rightarrow x_0 \\ x \in \mathcal{D} \setminus \{x_0\}}} \|x - x_0\|^{-1} [f(x) - f(x_0) - F(x - x_0)] = 0. \quad (10.4.6)$$

Proof. See [1404, p. 170]. □

In (10.4.6) the limit is taken over all sequences that are contained in $\mathcal{D}$, do not include x_0 , and converge to x_0 .

Definition 10.4.3. Let $\mathcal{D} \subseteq \mathbb{F}^m$ be solid and convex, let $f: \mathcal{D} \rightarrow \mathbb{F}^n$, let $x_0 \in \mathcal{D}$, and assume there exists a matrix $F \in \mathbb{F}^{n \times m}$ satisfying (10.4.6). Then, f is *differentiable at x_0* , and the matrix F is the (*Fréchet*) *derivative of f at x_0* . In this case, we write $f'(x_0) = F$ and

$$\lim_{\substack{x \rightarrow x_0 \\ x \in \mathcal{D} \setminus \{x_0\}}} \|x - x_0\|^{-1} [f(x) - f(x_0) - f'(x_0)(x - x_0)] = 0. \quad (10.4.7)$$

Note that Proposition 10.4.2 and Definition 10.4.3 do not require that x_0 lie in the interior of $\mathcal{D}$. We alternatively write $\frac{df(x_0)}{dx}$ for $f'(x_0)$.

Proposition 10.4.4. Let $\mathcal{D} \subseteq \mathbb{F}^m$ be solid and convex, let $f: \mathcal{D} \rightarrow \mathbb{F}^n$, let $x \in \mathcal{D}$, and assume that f is differentiable at x_0 . Then, f is continuous at x_0 .

Let $\mathcal{D} \subseteq \mathbb{F}^m$ be solid and convex, and let $f: \mathcal{D} \mapsto \mathbb{F}^n$. In terms of its scalar components, f can be written as $f = \begin{bmatrix} f_1 & \cdots & f_n \end{bmatrix}^T$, where $f_i: \mathcal{D} \mapsto \mathbb{F}$ for all $i = 1, \dots, n$ and $f(x) = \begin{bmatrix} f_1(x) & \cdots & f_n(x) \end{bmatrix}^T$ for all $x \in \mathcal{D}$. With this notation,

$f'(x_0)$ can be written as

$$f'(x_0) = \begin{bmatrix} f'_1(x_0) \\ \vdots \\ f'_n(x_0) \end{bmatrix}, \quad (10.4.8)$$

where $f'_i(x_0) \in \mathbb{F}^{1 \times m}$ is the *gradient* of f_i at x_0 and $f'(x_0)$ is the *Jacobian* of f at x_0 . Furthermore, if $x \in \text{int } \mathcal{D}$, then $f'(x_0)$ is related to the partial derivatives of f by

$$f'(x_0) = \left[\frac{\partial f(x_0)}{\partial x_{(1)}} \quad \cdots \quad \frac{\partial f(x_0)}{\partial x_{(m)}} \right], \quad (10.4.9)$$

where $\frac{\partial f(x_0)}{\partial x_{(i)}} \in \mathbb{F}^{n \times 1}$ for all $i = 1, \dots, m$. Note that the existence of the partial derivatives of f at x_0 does not imply that f is differentiable at x_0 , that is, $f'(x_0)$ given by (10.4.9) may not satisfy (10.4.7). Finally, note that the (i, j) entry of the $n \times m$ matrix $f'(x_0)$ is $\frac{\partial f_i(x_0)}{\partial x_{(j)}}$. For example, if $x \in \mathbb{F}^n$ and $A \in \mathbb{F}^{n \times n}$, then

$$\frac{d}{dx} Ax = A. \quad (10.4.10)$$

Let $\mathcal{D} \subseteq \mathbb{F}^m$ and $f: \mathcal{D} \mapsto \mathbb{F}^n$. If $f'(x)$ exists for all $x \in \mathcal{D}$ and $f': \mathcal{D} \mapsto \mathbb{F}^{n \times n}$ is continuous, then f is *continuously differentiable*, or C^1 . The *second derivative* of f at $x_0 \in \mathcal{D}$, denoted by $f''(x_0)$, is the derivative of $f': \mathcal{D} \mapsto \mathbb{F}^{n \times n}$ at $x_0 \in \mathcal{D}$. For $x_0 \in \mathcal{D}$ it can be seen that $f''(x_0): \mathbb{F}^m \times \mathbb{F}^m \mapsto \mathbb{F}^n$ is *bilinear*, that is, for all $\hat{\eta} \in \mathbb{F}^m$, the mapping $\eta \mapsto f''(x_0)(\eta, \hat{\eta})$ is linear and, for all $\eta \in \mathbb{F}^m$, the mapping $\hat{\eta} \mapsto f''(x_0)(\eta, \hat{\eta})$ is linear. Letting $f = [f_1 \quad \cdots \quad f_n]^T$, it follows that

$$f''(x_0)(\eta, \hat{\eta}) = \begin{bmatrix} \eta^T f''_1(x_0) \hat{\eta} \\ \vdots \\ \eta^T f''_n(x_0) \hat{\eta} \end{bmatrix}, \quad (10.4.11)$$

where, for all $i = 1, \dots, n$, the matrix $f''_i(x_0)$ is the $m \times m$ *Hessian* of f_i at x_0 . We write $f^{(2)}(x_0)$ for $f''(x_0)$ and $f^{(k)}(x_0)$ for the k th derivative of f at x_0 . f is C^k if $f^{(k)}(x)$ exists for all $x \in \mathcal{D}$ and $f^{(k)}$ is continuous on $\mathcal{D}$.

The following result is the *inverse function theorem*.

Theorem 10.4.5. Let $\mathcal{D} \subseteq \mathbb{F}^n$ be open, let $f: \mathcal{D} \mapsto \mathbb{F}^n$, and assume that f is C^k . Furthermore, let $x_0 \in \mathcal{D}$ be such that $\det f'(x_0) \neq 0$. Then, there exists an open set $\mathcal{N} \subset \mathbb{F}^n$ containing $f(x_0)$ and a C^k function $g: \mathcal{N} \mapsto \mathcal{D}$ such that $f[g(y)] = y$ for all $y \in \mathcal{N}$.

Let $S: [t_0, t_1] \mapsto \mathbb{F}^{n \times m}$, and assume that every entry of $S(t)$ is differentiable. Then, define $\dot{S}(t) \triangleq \frac{dS(t)}{dt} \in \mathbb{F}^{n \times m}$ for all $t \in [t_0, t_1]$ entrywise, that is, for all $i = 1, \dots, n$ and $j = 1, \dots, m$,

$$[\dot{S}(t)]_{(i,j)} \triangleq \frac{d}{dt} S_{(i,j)}(t). \quad (10.4.12)$$

If $t = t_0$ or $t = t_1$, then d^+/dt or d^-/dt (or just d/dt) denotes the right and left one-sided derivatives, respectively. Finally, define $\int_{t_0}^{t_1} S(t) dt$ entrywise, that is, for

all $i = 1, \dots, n$ and $j = 1, \dots, m$,

$$\left[\int_{t_0}^{t_1} S(t) dt \right]_{(i,j)} \triangleq \int_{t_0}^{t_1} [S(t)]_{(i,j)} dt. \quad (10.4.13)$$

10.5 Functions of a Matrix

Consider the function $f: \mathcal{D} \subseteq \mathbb{C} \mapsto \mathbb{C}$ defined by the power series

$$f(s) = \sum_{i=0}^{\infty} \beta_i s^i, \quad (10.5.1)$$

where $\beta_i \in \mathbb{C}$ for all $i \in \mathbb{N}$, and assume that this series converges for all $|s| < \gamma$. Then, for $A \in \mathbb{C}^{n \times n}$, we define

$$f(A) \triangleq \sum_{i=0}^{\infty} \beta_i A^i, \quad (10.5.2)$$

which converges for all $A \in \mathbb{C}^{n \times n}$ such that $\text{sprad}(A) < \gamma$. Now, assume that $A = SBS^{-1}$, where $S \in \mathbb{C}^{n \times n}$ is nonsingular, $B \in \mathbb{C}^{n \times n}$, and $\text{sprad}(B) < \gamma$. Then,

$$f(A) = Sf(B)S^{-1}. \quad (10.5.3)$$

If, in addition, $B = \text{diag}(J_1, \dots, J_r)$ is the Jordan form of A , then

$$f(A) = S \text{diag}[f(J_1), \dots, f(J_r)] S^{-1}. \quad (10.5.4)$$

Letting $J = \lambda I_k + N_k$ denote a $k \times k$ Jordan block, expanding and rearranging the infinite series $\sum_{i=0}^{\infty} \beta_i J^i$ shows that $f(J)$ is the $k \times k$ upper triangular Toeplitz matrix

$$\begin{aligned} f(J) &= f(\lambda)N_k + f'(\lambda)N_k + \frac{1}{2}f''(\lambda)N_k^2 + \dots + \frac{1}{(k-1)!}f^{(k-1)}(\lambda)N_k^{k-1} \\ &= \begin{bmatrix} f(\lambda) & f'(\lambda) & \frac{1}{2}f''(\lambda) & \dots & \frac{1}{(k-1)!}f^{(k-1)}(\lambda) \\ 0 & f(\lambda) & f'(\lambda) & \dots & \frac{1}{(k-2)!}f^{(k-2)}(\lambda) \\ 0 & 0 & f(\lambda) & \dots & \frac{1}{(k-3)!}f^{(k-3)}(\lambda) \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & f(\lambda) \end{bmatrix}. \end{aligned} \quad (10.5.5)$$

Next, we extend the definition $f(A)$ to functions $f: \mathcal{D} \subseteq \mathbb{C} \mapsto \mathbb{C}$ that are not necessarily of the form (10.5.1).

Definition 10.5.1. Let $f: \mathcal{D} \subseteq \mathbb{C} \mapsto \mathbb{C}$, let $A \in \mathbb{C}^{n \times n}$, where $\text{spec}(A) \subset \mathcal{D}$, and assume that, for all $\lambda_i \in \text{spec}(A)$, f is $k_i - 1$ times differentiable at λ_i , where $k_i \triangleq \text{ind}_A(\lambda_i)$ is the order of the largest Jordan block associated with λ_i as given by Theorem 5.3.3. Then, f is *defined* at A , and $f(A)$ is given by (10.5.3) and (10.5.4), where $f(J_i)$ is defined by (10.5.5) with $k = k_i$ and $\lambda = \lambda_i$.

Theorem 10.5.2. Let $A \in \mathbb{F}^{n \times n}$, let $\text{spec}(A) = \{\lambda_1, \dots, \lambda_r\}$, and, for $i = 1, \dots, r$, let $k_i \triangleq \text{ind}_A(\lambda_i)$. Furthermore, suppose that $f: \mathcal{D} \subseteq \mathbb{C} \mapsto \mathbb{C}$ is defined at A . Then, there exists a polynomial $p \in \mathbb{F}[s]$ such that $f(A) = p(A)$. Furthermore, there exists a unique polynomial p of minimal degree $\sum_{i=1}^r k_i$ satisfying $f(A) = p(A)$ and such that, for all $i = 1, \dots, r$ and $j = 0, 1, \dots, k_i - 1$,

$$f^{(j)}(\lambda_i) = p^{(j)}(\lambda_i). \quad (10.5.6)$$

This polynomial is given by

$$p(s) = \sum_{i=1}^r \left(\left[\prod_{\substack{j=1 \\ j \neq i}}^r (s - \lambda_j)^{n_j} \right] \sum_{k=0}^{k_i-1} \frac{1}{k!} \frac{d^k}{ds^k} \frac{f(s)}{\prod_{\substack{l=1 \\ l \neq i}}^r (s - \lambda_l)^{k_l}} \right) \bigg|_{s=\lambda_i} (s - \lambda_i)^k. \quad (10.5.7)$$

If, in addition, A is diagonalizable, then p is given by

$$p(s) = \sum_{i=1}^r f(\lambda_i) \prod_{\substack{j=1 \\ j \neq i}}^r \frac{s - \lambda_j}{\lambda_i - \lambda_j}. \quad (10.5.8)$$

Proof. See [359, pp. 263, 264]. □

The polynomial (10.5.7) is the *Lagrange-Hermite interpolation polynomial* for f .

The following result, which is known as the *identity theorem*, is a special case of Theorem 10.5.2.

Theorem 10.5.3. Let $A \in \mathbb{F}^{n \times n}$, let $\text{spec}(A) = \{\lambda_1, \dots, \lambda_r\}$, and, for $i = 1, \dots, r$, let $k_i \triangleq \text{ind}_A(\lambda_i)$. Furthermore, let $f: \mathcal{D} \subseteq \mathbb{C} \mapsto \mathbb{C}$ and $g: \mathcal{D} \subseteq \mathbb{C} \mapsto \mathbb{C}$ be analytic on a neighborhood of $\text{spec}(A)$. Then, $f(A) = g(A)$ if and only if, for all $i = 1, \dots, r$ and $j = 0, 1, \dots, k_i - 1$,

$$f^{(j)}(\lambda_i) = g^{(j)}(\lambda_i). \quad (10.5.9)$$

Corollary 10.5.4. Let $A \in \mathbb{F}^{n \times n}$, and let $f: \mathcal{D} \subset \mathbb{C} \rightarrow \mathbb{C}$ be analytic on a neighborhood of $\text{mspec}(A)$. Then,

$$\text{mspec}[f(A)] = f[\text{mspec}(A)]. \quad (10.5.10)$$

10.6 Matrix Square Root and Matrix Sign Functions

Theorem 10.6.1. Let $A \in \mathbb{C}^{n \times n}$, and assume that A is group invertible and has no eigenvalues in $(-\infty, 0)$. Then, there exists a unique matrix $B \in \mathbb{C}^{n \times n}$ such that $\text{spec}(B) \subset \text{ORHP} \cup \{0\}$ and such that $B^2 = A$. If, in addition, A is real, then B is real.

Proof. See [683, pp. 20, 31]. □

The matrix B given by Theorem 10.6.1 is the *principal square root* of A . This matrix is denoted by $A^{1/2}$. The existence of a square root that is not necessarily the principal square root is discussed in Fact 5.15.19.

The following result defines the *matrix sign function*.

Definition 10.6.2. Let $A \in \mathbb{C}^{n \times n}$, assume that A has no eigenvalues on the imaginary axis, and let

$$A = S \begin{bmatrix} J_1 & 0 \\ 0 & J_2 \end{bmatrix} S^{-1},$$

where $S \in \mathbb{C}^{n \times n}$ is nonsingular, $J_1 \in \mathbb{C}^{p \times p}$ and $J_2 \in \mathbb{C}^{q \times q}$ are in Jordan canonical form, and $\text{spec}(J_1) \subset \text{OLHP}$ and $\text{spec}(J_2) \subset \text{ORHP}$. Then, the *matrix sign* of A is defined by

$$\text{Sign}(A) \triangleq S \begin{bmatrix} -I_p & 0 \\ 0 & I_q \end{bmatrix} S^{-1}.$$

10.7 Matrix Derivatives

In this section we consider derivatives of differentiable scalar-valued functions with matrix arguments. Consider the linear function $f: \mathbb{F}^{m \times n} \mapsto \mathbb{F}$ given by $f(X) = \text{tr} AX$, where $A \in \mathbb{F}^{n \times m}$ and $X \in \mathbb{F}^{m \times n}$. In terms of vectors $x \in \mathbb{F}^{mn}$, we can define the linear function $\hat{f}(x) \triangleq (\text{vec } A)^T x$ so that $\hat{f}(\text{vec } X) = f(X) = (\text{vec } A)^T \text{vec } X$. Consequently, for all $Y \in \mathbb{F}^{m \times n}$, $f'(X_0)$ can be represented by $f'(X_0)Y = \text{tr } AY$.

These observations suggest that a convenient representation of the derivative $\frac{d}{dX}f(X)$ of a differentiable scalar-valued differentiable function $f(X)$ of a matrix argument $X \in \mathbb{F}^{m \times n}$ is the $n \times m$ matrix whose (i, j) entry is $\frac{\partial f(X)}{\partial X_{(j,i)}}$. Note the order of indices.

Proposition 10.7.1. Let $x \in \mathbb{F}^n$. Then, the following statements hold:

i) If $A \in \mathbb{F}^{n \times n}$, then

$$\frac{d}{dx} x^T A x = x^T (A + A^T). \quad (10.7.1)$$

ii) If $A \in \mathbb{F}^{n \times n}$ is symmetric, then

$$\frac{d}{dx} x^T A x = 2x^T A. \quad (10.7.2)$$

iii) If $A \in \mathbb{F}^{n \times n}$ is Hermitian, then

$$\frac{d}{dx} x^* A x = 2x^* A. \quad (10.7.3)$$

Proposition 10.7.2. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{l \times n}$. Then, the following statements hold:

i) For all $X \in \mathbb{F}^{m \times n}$,

$$\frac{d}{dX} \text{tr } AX = A. \quad (10.7.4)$$

ii) For all $X \in \mathbb{F}^{m \times l}$,

$$\frac{d}{dX} \operatorname{tr} AXB = BA. \quad (10.7.5)$$

iii) For all $X \in \mathbb{F}^{l \times m}$,

$$\frac{d}{dX} \operatorname{tr} AX^T B = A^T B^T. \quad (10.7.6)$$

iv) For all $X \in \mathbb{F}^{m \times l}$ and $k \geq 1$,

$$\frac{d}{dX} \operatorname{tr} (AXB)^k = kB(AXB)^{k-1}A. \quad (10.7.7)$$

v) For all $X \in \mathbb{F}^{m \times l}$,

$$\frac{d}{dX} \det AXB = B(AXB)^A A. \quad (10.7.8)$$

vi) For all $X \in \mathbb{F}^{m \times l}$ such that AXB is nonsingular,

$$\frac{d}{dX} \log \det AXB = B(AXB)^{-1}A. \quad (10.7.9)$$

Proposition 10.7.3. Let $A \in \mathbb{F}^{n \times m}$ and $B \in \mathbb{F}^{m \times n}$. Then, the following statements hold:

i) For all $X \in \mathbb{F}^{m \times m}$ and $k \geq 1$,

$$\frac{d}{dX} \operatorname{tr} AX^k B = \sum_{i=0}^{k-1} X^{k-1-i} B A X^i. \quad (10.7.10)$$

ii) For all nonsingular $X \in \mathbb{F}^{m \times m}$,

$$\frac{d}{dX} \operatorname{tr} AX^{-1} B = -X^{-1} B A X^{-1}. \quad (10.7.11)$$

iii) For all nonsingular $X \in \mathbb{F}^{m \times m}$,

$$\frac{d}{dX} \det AX^{-1} B = -X^{-1} B (A X^{-1} B)^A A X^{-1}. \quad (10.7.12)$$

iv) For all nonsingular $X \in \mathbb{F}^{m \times m}$,

$$\frac{d}{dX} \log \det AX^{-1} B = -X^{-1} B (A X^{-1} B)^{-1} A X^{-1}. \quad (10.7.13)$$

Proposition 10.7.4. The following statements hold:

i) Let $A, B \in \mathbb{F}^{n \times m}$. Then, for all $X \in \mathbb{F}^{m \times n}$,

$$\frac{d}{dX} \operatorname{tr} A X B X = A X B + B X A. \quad (10.7.14)$$

ii) Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then, for all $X \in \mathbb{F}^{n \times m}$,

$$\frac{d}{dX} \operatorname{tr} A X B X^T = B X^T A + B^T X^T A^T. \quad (10.7.15)$$

iii) Let $A \in \mathbb{F}^{k \times l}$, $B \in \mathbb{F}^{l \times m}$, $C \in \mathbb{F}^{n \times l}$, $D \in \mathbb{F}^{l \times l}$, and $E \in \mathbb{F}^{l \times k}$. Then, for all $X \in \mathbb{F}^{m \times n}$,

$$\frac{d}{dX} \operatorname{tr} A(D + BXC)^{-1} E = -C(D + BXC)^{-1} E A(D + BXC)^{-1} B. \quad (10.7.16)$$

iv) Let $A \in \mathbb{F}^{k \times l}$, $B \in \mathbb{F}^{l \times m}$, $C \in \mathbb{F}^{n \times l}$, $D \in \mathbb{F}^{l \times l}$, and $E \in \mathbb{F}^{l \times k}$. Then, for all $X \in \mathbb{F}^{n \times m}$,

$$\begin{aligned} & \frac{d}{dX} \operatorname{tr} A(D + BX^T C)^{-1} E \\ &= -B^T (D + BX^T C)^{-T} A^T E^T (D + BX^T C)^{-T} C^T. \end{aligned} \quad (10.7.17)$$

10.8 Facts Involving One Set

Fact 10.8.1. Let $x \in \mathbb{F}^n$, and let $\varepsilon > 0$. Then, $\mathbb{B}_\varepsilon(x)$ is completely solid and convex.

Fact 10.8.2. Let $\mathcal{S} \subset \mathbb{F}^n$, assume that $\mathcal{S}$ is bounded, let $\delta > 0$ satisfy $\|x - y\| < \delta$ for all $x, y \in \mathcal{S}$, and let $x_0 \in \mathcal{S}$. Then, $\mathcal{S} \subseteq \mathbb{B}_\delta(x_0)$.

Fact 10.8.3. Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then, $\operatorname{cl} \mathcal{S}$ is the smallest closed set containing $\mathcal{S}$, and $\operatorname{int} \mathcal{S}$ is the largest open set contained in $\mathcal{S}$.

Fact 10.8.4. Let $\mathcal{S} \subseteq \mathbb{F}^n$. If $\mathcal{S}$ is (open, closed), then $\mathcal{S}^\sim$ is (closed, open).

Fact 10.8.5. Let $\mathcal{S} \subseteq \mathcal{S}' \subseteq \mathbb{F}^n$. If $\mathcal{S}$ is (open relative to $\mathcal{S}'$, closed relative to $\mathcal{S}'$), then $\mathcal{S}' \setminus \mathcal{S}$ is (closed relative to $\mathcal{S}'$, open relative to $\mathcal{S}'$).

Fact 10.8.6. Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then,

$$(\operatorname{int} \mathcal{S})^\sim = \operatorname{cl}(\mathcal{S}^\sim)$$

and

$$\operatorname{bd} \mathcal{S} = \operatorname{bd} \mathcal{S}^\sim = (\operatorname{cl} \mathcal{S}) \cap (\operatorname{cl} \mathcal{S}^\sim) = [(\operatorname{int} \mathcal{S}) \cup \operatorname{int}(\mathcal{S}^\sim)]^\sim.$$

Hence, $\operatorname{bd} \mathcal{S}$ is closed.

Fact 10.8.7. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is either open or closed. Then, $\operatorname{int} \operatorname{bd} \mathcal{S}$ is empty. (Proof: See [68, p. 68].)

Fact 10.8.8. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is convex. Then, $\operatorname{cl} \mathcal{S}$, $\operatorname{int} \mathcal{S}$, and $\operatorname{int}_{\operatorname{aff} \mathcal{S}} \mathcal{S}$ are convex. (Proof: See [1133, p. 45] and [1134, p. 64].)

Fact 10.8.9. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is convex. Then, the following statements are equivalent:

- i) $\mathcal{S}$ is solid.
- ii) $\mathcal{S}$ is completely solid.
- iii) $\dim \mathcal{S} = n$.
- iv) $\operatorname{aff} \mathcal{S} = \mathbb{F}^n$.

Fact 10.8.10. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is solid. Then, $\operatorname{co} \mathcal{S}$ is completely solid.

Fact 10.8.11. Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then,

$$\text{cl } \mathcal{S} \subseteq \text{aff cl } \mathcal{S} = \text{aff } \mathcal{S}.$$

(Proof: See [239, p. 7].)

Fact 10.8.12. Let $k \leq n$, and let $x_1, \dots, x_k \in \mathbb{F}^n$. Then,

$$\text{int aff } \{x_1, \dots, x_k\} = \emptyset.$$

(Remark: See Fact 2.9.7.)

Fact 10.8.13. Let $\mathcal{S} \subseteq \mathbb{F}^n$. Then,

$$\text{co cl } \mathcal{S} \subseteq \text{cl co } \mathcal{S}.$$

Now, assume that $\mathcal{S}$ is either bounded or convex. Then,

$$\text{co cl } \mathcal{S} = \text{cl co } \mathcal{S}.$$

(Proof: Use Fact 10.8.8 and Fact 10.8.13.) (Remark: Although

$$\mathcal{S} = \left\{ x \in \mathbb{R}^2: x_{(1)}^2 x_{(2)}^2 = 1 \text{ for all } x_{(1)} > 0 \right\}$$

is closed, $\text{co } \mathcal{S}$ is not closed. Hence, $\text{co cl } \mathcal{S} \subset \text{cl co } \mathcal{S}$.)

Fact 10.8.14. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is open. Then, $\text{co } \mathcal{S}$ is open.

Fact 10.8.15. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is compact. Then, $\text{co } \mathcal{S}$ is compact.

Fact 10.8.16. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is solid. Then, $\dim \mathcal{S} = n$.

Fact 10.8.17. Let $\mathcal{S} \subseteq \mathbb{F}^m$, assume that $\mathcal{S}$ is solid, let $A \in \mathbb{F}^{n \times m}$, and assume that A is right invertible. Then, $A\mathcal{S}$ is solid. (Proof: Use Theorem 10.3.6.) (Remark: See Fact 2.10.4.)

Fact 10.8.18. $\mathbf{N}^n$ is a closed and completely solid subset of $\mathbb{F}^{n(n+1)/2}$. Furthermore,

$$\text{int } \mathbf{N}^n = \mathbf{P}^n.$$

Fact 10.8.19. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is convex. Then,

$$\text{int cl } \mathcal{S} = \text{int } \mathcal{S}.$$

Fact 10.8.20. Let $\mathcal{D} \subseteq \mathbb{F}^n$, and let x_0 belong to a solid, convex subset of $\mathcal{D}$. Then,

$$\dim \text{vcone}(\mathcal{D}, x_0) = n.$$

Fact 10.8.21. Let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is a subspace. Then, $\mathcal{S}$ is closed.

Fact 10.8.22. Let $\mathcal{S} \subseteq \mathbb{F}^n$, assume that $\mathcal{S}$ is symmetric, solid, convex, closed, and bounded, and, for all $x \in \mathbb{F}^n$, define

$$\|x\| \triangleq \min\{\alpha \geq 0: x \in \alpha\mathcal{S}\} = \max\{\alpha \geq 0: \alpha x \in \mathcal{S}\}.$$

Then, $\|\cdot\|$ is a norm on $\mathbb{F}^n$, and $\mathbb{B}_1(0) = \text{int } \mathcal{S}$. Conversely, let $\|\cdot\|$ be a norm on $\mathbb{F}^n$. Then, $\mathbb{B}_1(0)$ is convex, bounded, symmetric, and solid. (Proof: See [721, pp. 38, 39].) (Remark: In all cases, $\mathbb{B}_1(0)$ is defined with respect to $\|\cdot\|$. This result is due to Minkowski.) (Remark: See Fact 9.7.23.)

Fact 10.8.23. Let $\mathcal{S} \subseteq \mathbb{R}^m$, assume that $\mathcal{S}$ is nonempty, closed, and convex, and define $\mathcal{E} \subseteq \mathcal{S}$ by

$$\mathcal{E} \triangleq \{x \in \mathcal{S} : x \text{ is not a convex combination of two distinct elements of } \mathcal{S}\}.$$

Then, $\mathcal{E}$ is nonempty, closed, and convex, and

$$\mathcal{E} = \text{co } \mathcal{S}.$$

(Proof: See [447, pp. 482–484].) (Remark: $\mathcal{E}$ is the set of *extreme points* of $\mathcal{S}$.) (Remark: The last result is the *Krein-Milman theorem*.)

10.9 Facts Involving Two or More Sets

Fact 10.9.1. Let $\mathcal{S}_1 \subseteq \mathcal{S}_2 \subseteq \mathbb{F}^n$. Then,

$$\text{cl } \mathcal{S}_1 \subseteq \text{cl } \mathcal{S}_2$$

and

$$\text{int } \mathcal{S}_1 \subseteq \text{int } \mathcal{S}_2.$$

Fact 10.9.2. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$. Then, the following statements hold:

- i) $(\text{int } \mathcal{S}_1) \cap (\text{int } \mathcal{S}_2) = \text{int}(\mathcal{S}_1 \cap \mathcal{S}_2)$.
- ii) $(\text{int } \mathcal{S}_1) \cup (\text{int } \mathcal{S}_2) \subseteq \text{int}(\mathcal{S}_1 \cup \mathcal{S}_2)$.
- iii) $(\text{cl } \mathcal{S}_1) \cup (\text{cl } \mathcal{S}_2) = \text{cl}(\mathcal{S}_1 \cup \mathcal{S}_2)$.
- iv) $\text{bd}(\mathcal{S}_1 \cup \mathcal{S}_2) \subseteq (\text{bd } \mathcal{S}_1) \cup (\text{bd } \mathcal{S}_2)$.
- v) If $(\text{cl } \mathcal{S}_1) \cap (\text{cl } \mathcal{S}_2) = \emptyset$, then $\text{bd}(\mathcal{S}_1 \cup \mathcal{S}_2) = (\text{bd } \mathcal{S}_1) \cup (\text{bd } \mathcal{S}_2)$.

(Proof: See [68, p. 65].)

Fact 10.9.3. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$, assume that either $\mathcal{S}_1$ or $\mathcal{S}_2$ is closed, and assume that $\text{int } \mathcal{S}_1 = \text{int } \mathcal{S}_2 = \emptyset$. Then, $\text{int}(\mathcal{S}_1 \cup \mathcal{S}_2)$ is empty. (Proof: See [68, p. 69].) (Remark: $\text{int}(\mathcal{S}_1 \cup \mathcal{S}_2)$ is not necessarily empty if neither $\mathcal{S}_1$ nor $\mathcal{S}_2$ is closed. Consider the sets of rational and irrational numbers.)

Fact 10.9.4. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}_1$ is closed and $\mathcal{S}_2$ is compact. Then, $\mathcal{S}_1 + \mathcal{S}_2$ is closed. (Proof: See [442, p. 209].)

Fact 10.9.5. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are closed and compact. Then, $\mathcal{S}_1 + \mathcal{S}_2$ is closed and compact. (Proof: See [153, p. 34].)

Fact 10.9.6. Let $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3 \subseteq \mathbb{F}^n$, assume that $\mathcal{S}_1, \mathcal{S}_2$, and $\mathcal{S}_3$ are closed and convex, assume that $\mathcal{S}_1 \cap \mathcal{S}_2 \neq \emptyset$, $\mathcal{S}_2 \cap \mathcal{S}_3 \neq \emptyset$, and $\mathcal{S}_3 \cap \mathcal{S}_1 \neq \emptyset$, and assume that $\mathcal{S}_1 \cup \mathcal{S}_2 \cup \mathcal{S}_3$ is convex. Then, $\mathcal{S}_1 \cap \mathcal{S}_2 \cap \mathcal{S}_3 \neq \emptyset$. (Proof: See [153, p. 32].)

Fact 10.9.7. Let $\mathcal{S}_1, \mathcal{S}_2, \mathcal{S}_3 \subseteq \mathbb{F}^n$, assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are convex, $\mathcal{S}_2$ is closed, and $\mathcal{S}_3$ is bounded, and assume that $\mathcal{S}_1 + \mathcal{S}_3 \subseteq \mathcal{S}_2 + \mathcal{S}_3$. Then, $\mathcal{S}_1 \subseteq \mathcal{S}_2$. (Proof: See [239, p. 5].) (Remark: This result is due to Radstrom.)

Fact 10.9.8. Let $\mathcal{S} \subseteq \mathbb{F}^m$, assume that $\mathcal{S}$ is closed, let $A \in \mathbb{F}^{n \times m}$, and assume that A has full row rank. Then, $A\mathcal{S}$ is not necessarily closed. (Remark: See Theorem 10.3.6.)

Fact 10.9.9. Let $\mathcal{A}$ be a collection of open subsets of $\mathbb{R}^n$. Then, the union of all elements of $\mathcal{A}$ is open. If, in addition, $\mathcal{A}$ is finite, then the intersection of all elements of $\mathcal{A}$ is open. (Proof: See [68, p. 50].)

Fact 10.9.10. Let $\mathcal{A}$ be a collection of closed subsets of $\mathbb{R}^n$. Then, the intersection of all elements of $\mathcal{A}$ is closed. If, in addition, $\mathcal{A}$ is finite, then the union of all elements of $\mathcal{A}$ is closed. (Proof: See [68, p. 50].)

Fact 10.9.11. Let $\mathcal{A} = \{A_1, A_2, \dots\}$ be a collection of nonempty, closed subsets of $\mathbb{R}^n$ such that A_1 is bounded and such that, for all $i = 1, 2, \dots$, $A_{i+1} \subseteq A_i$. Then, $\bigcap_{i=1}^{\infty} A_i$ is closed and nonempty. (Proof: See [68, p. 56].) (Remark: This result is the *Cantor intersection theorem*.)

Fact 10.9.12. Let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, let $\mathcal{S} \subset \mathbb{F}^n$, assume that $\mathcal{S}$ is a subspace, let $y \in \mathbb{F}^n$, and define

$$\mu \triangleq \max_{x \in \{z \in \mathcal{S}: \|z\|=1\}} |y^*x|.$$

Then, there exists a vector $z \in \mathcal{S}^\perp$ such that

$$\max_{x \in \{z \in \mathbb{F}^n: \|z\|=1\}} |(y+z)^*x| = \mu.$$

(Proof: See [1230, p. 57].) (Remark: This result is a version of the *Hahn-Banach theorem*.) (Problem: Find a simple interpretation in $\mathbb{R}^2$.)

Fact 10.9.13. Let $\mathcal{S} \subset \mathbb{R}^n$, assume that $\mathcal{S}$ is a convex cone, let $x \in \mathbb{R}^n$, and assume that $x \notin \text{int } \mathcal{S}$. Then, there exists a nonzero vector $\lambda \in \mathbb{R}^n$ such that $\lambda^T x \leq 0$ and $\lambda^T z \geq 0$ for all $z \in \mathcal{S}$. (Remark: This result is a *separation theorem*. See [879, p. 37], [1096, p. 443], [1133, pp. 95–101], and [1235, pp. 96–100].)

Fact 10.9.14. Let $\mathcal{S}_1, \mathcal{S}_2 \subset \mathbb{R}^n$, and assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are convex. Then, the following statements are equivalent:

- i) There exist a nonzero vector $\lambda \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$ such that $\lambda^T x \leq \alpha$ for all $x \in \mathcal{S}_1$, $\lambda^T x \geq \alpha$ for all $x \in \mathcal{S}_2$, and either $\mathcal{S}_1$ or $\mathcal{S}_2$ is not contained in the affine hyperplane $\{x \in \mathbb{R}^n: \lambda^T x = \alpha\}$.
- ii) $\text{int}_{\text{aff } \mathcal{S}_1} \mathcal{S}_1$ and $\text{int}_{\text{aff } \mathcal{S}_2} \mathcal{S}_2$ are disjoint.

(Proof: See [180, p. 82].) (Remark: This result is a *proper separation theorem*.)

Fact 10.9.15. Let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, let $y \in \mathbb{F}^n$, let $\mathcal{S} \subseteq \mathbb{F}^n$, and assume that $\mathcal{S}$ is nonempty and closed. Then, there exists a vector $x_0 \in \mathcal{S}$ such that

$$\|y - x_0\| = \min_{x \in \mathcal{S}} \|y - x\|.$$

Now, assume that $\mathcal{S}$ is convex. Then, there exists a unique vector $x_0 \in \mathcal{S}$ such that

$$\|y - x_0\| = \min_{x \in \mathcal{S}} \|y - x\|.$$

In other words, there exists a vector $x_0 \in \mathcal{S}$ such that, for all $x \in \mathcal{S} \setminus \{x_0\}$,

$$\|y - x_0\| < \|y - x\|.$$

(Proof: See [447, pp. 470, 471].) (Remark: See Fact 10.9.17.)

Fact 10.9.16. Let $\|\cdot\|$ be a norm on $\mathbb{F}^n$, let $y_1, y_2 \in \mathbb{F}^n$, let $\mathcal{S} \subseteq \mathbb{F}^n$, assume that $\mathcal{S}$ is nonempty, closed, and convex, and let x_1 and x_2 denote the unique elements of $\mathcal{S}$ that are closest to y_1 and y_2 , respectively. Then,

$$\|x_1 - x_2\| \leq \|y_1 - y_2\|.$$

(Proof: See [447, pp. 474, 475].)

Fact 10.9.17. Let $\mathcal{S} \subseteq \mathbb{R}^n$, assume that $\mathcal{S}$ is a subspace, let $A \in \mathbb{F}^{n \times n}$ be the projector onto $\mathcal{S}$, and let $x \in \mathbb{F}^n$. Then,

$$\min_{y \in \mathcal{S}} \|x - y\|_2 = \|A_{\perp}x\|_2.$$

(Proof: See [536, p. 41] or [1230, p. 91].) (Remark: See Fact 10.9.15.)

Fact 10.9.18. Let $\mathcal{S}_1, \mathcal{S}_2 \subseteq \mathbb{R}^n$, assume that $\mathcal{S}_1$ and $\mathcal{S}_2$ are subspaces, let A_1 and A_2 be the projectors onto $\mathcal{S}_1$ and $\mathcal{S}_2$, respectively, and define

$$\text{dist}(\mathcal{S}_1, \mathcal{S}_2) \triangleq \max \left\{ \max_{\substack{x \in \mathcal{S}_1 \\ \|x\|=1}} \min_{y \in \mathcal{S}_2} \|x - y\|_2, \max_{\substack{y \in \mathcal{S}_2 \\ \|y\|=1}} \min_{x \in \mathcal{S}_1} \|x - y\|_2 \right\}.$$

Then,

$$\text{dist}(\mathcal{S}_1, \mathcal{S}_2) = \sigma_{\max}(A_1 - A_2).$$

If, in addition, $\dim \mathcal{S}_1 = \dim \mathcal{S}_2$, then

$$\text{dist}(\mathcal{S}_1, \mathcal{S}_2) = \sin \theta,$$

where θ is the minimal principal angle defined in Fact 5.11.39. (Proof: See [560, Chapter 13] and [1230, pp. 92, 93].) (Remark: If $\|\cdot\|$ is a norm on $\mathbb{F}^{n \times n}$, then

$$\text{dist}(\mathcal{S}_1, \mathcal{S}_2) \triangleq \|A_1 - A_2\|_2$$

defines a metric on the set of all subspaces of $\mathbb{F}^n$, yielding the *gap topology*.) (Remark: See Fact 5.12.17.)

10.10 Facts on Matrix Functions

Fact 10.10.1. Let $A \in \mathbb{C}^{n \times n}$, and assume that A is group invertible and has no eigenvalues in $(-\infty, 0)$. Then,

$$A^{1/2} = \frac{2}{\pi} A \int_0^\infty (t^2 I + A)^{-1} dt.$$

(Proof: See [683, p. 133].)

Fact 10.10.2. Let $A \in \mathbb{C}^{n \times n}$, and assume that A has no eigenvalues on the imaginary axis. Then, the following statements hold:

- i) $\text{Sign}(A)$ is involutory.
- ii) $A = \text{Sign}(A)$ if and only if A is involutory.
- iii) $[A, \text{Sign}(A)] = 0$.
- iv) $\text{Sign}(A) = \text{Sign}(A^{-1})$.
- v) If A is real, then $\text{Sign}(A)$ is real.
- vi) $\text{Sign}(A) = A(A^2)^{-1/2}$.
- vii) $\text{Sign}(A)$ is given by

$$\text{Sign}(A) = \frac{2}{\pi} A \int_0^\infty (t^2 I + A^2)^{-1} dt.$$

(Proof: See [683, pp. 39, 40 and Chapter 5] and [803].) (Remark: The square root in vi) is the principal square root.)

Fact 10.10.3. Let $A, B \in \mathbb{C}^{n \times n}$, assume that AB has no eigenvalues on the imaginary axis, and define $C \triangleq A(BA)^{-1/2}$. Then,

$$\text{Sign}\left(\begin{bmatrix} 0 & A \\ B & 0 \end{bmatrix}\right) = \begin{bmatrix} 0 & C \\ C^{-1} & 0 \end{bmatrix}.$$

If, in addition, A has no eigenvalues on the imaginary axis, then

$$\text{Sign}\left(\begin{bmatrix} 0 & A \\ I & 0 \end{bmatrix}\right) = \begin{bmatrix} 0 & A^{1/2} \\ A^{-1/2} & 0 \end{bmatrix}.$$

(Proof: See [683, p. 108].) (Remark: The square root is the principal square root.)

Fact 10.10.4. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that A and B are positive definite. Then,

$$\text{Sign}\left(\begin{bmatrix} 0 & B \\ A^{-1} & 0 \end{bmatrix}\right) = \begin{bmatrix} 0 & A\#B \\ (A\#B)^{-1} & 0 \end{bmatrix}.$$

(Proof: See [683, p. 131].) (Remark: The geometric mean is defined in Fact 8.10.43.)

10.11 Facts on Functions and Derivatives

Fact 10.11.1. Let $(x_i)_{i=1}^{\infty} \subset \mathbb{F}^n$. Then, $\lim_{i \rightarrow \infty} x_i = x$ if and only if $\lim_{i \rightarrow \infty} x_{i(j)} = x_{(j)}$ for all $j = 1, \dots, n$.

Fact 10.11.2. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \dots + a_0$, define $p_{\varepsilon_0, \dots, \varepsilon_{n-1}}(s) \triangleq s^n + (a_{n-1} + \varepsilon_{n-1})s^{n-1} + \dots + (a_1 + \varepsilon_1)s + a_0 + \varepsilon_0$, where $\varepsilon_0, \dots, \varepsilon_{n-1} \in \mathbb{R}$, let $\text{roots}(p) = \{\lambda_1, \dots, \lambda_r\}$, and, for all $i = 1, \dots, r$, let $\alpha_i \in \mathbb{R}$ satisfy $0 < \alpha_i < \max_{j \neq i} |\lambda_i - \lambda_j|$. Then, there exists $\varepsilon > 0$ such that, for all $\varepsilon_0, \dots, \varepsilon_{n-1}$ satisfying $|\varepsilon_i| < \varepsilon$, $i = 1, \dots, r$, the polynomial $p_{\varepsilon_0, \dots, \varepsilon_{n-1}}$ has exactly $\text{mult}_p(\lambda_i)$ roots in the disk $\{s \in \mathbb{C} : |s - \lambda_i| < \alpha_i\}$. (Proof: See [1005].) (Remark: This result shows that the roots of a polynomial are continuous functions of the coefficients.)

Fact 10.11.3. Let $p \in \mathbb{C}[s]$. Then,

$$\text{roots}(p') \subseteq \text{co roots}(p).$$

(Proof: See [447, p. 488].) (Remark: p' is the derivative of p .)

Fact 10.11.4. Let $\mathcal{S}_1 \subseteq \mathbb{F}^n$, assume that $\mathcal{S}_1$ is compact, let $\mathcal{S}_2 \subset \mathbb{F}^m$, let $f: \mathcal{S}_1 \times \mathcal{S}_2 \rightarrow \mathbb{R}$, and assume that f is continuous. Then, $g: \mathcal{S}_2 \rightarrow \mathbb{R}$ defined by $g(y) \triangleq \max_{x \in \mathcal{S}_1} f(x, y)$ is continuous. (Remark: A related result is given in [442, p. 208].)

Fact 10.11.5. Let $\mathcal{S} \subseteq \mathbb{F}^n$, assume that $\mathcal{S}$ is pathwise connected, let $f: \mathcal{S} \mapsto \mathbb{F}^n$, and assume that f is continuous. Then, $f(\mathcal{S})$ is pathwise connected. (Proof: See [1256, p. 65].)

Fact 10.11.6. Let $f: [0, \infty) \rightarrow \mathbb{R}$, assume that f is continuous, and assume that $\lim_{t \rightarrow \infty} f(t)$ exists. Then,

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t f(\tau) d\tau = \lim_{t \rightarrow \infty} f(t).$$

(Remark: The assumption that f is continuous can be weakened.)

Fact 10.11.7. Let $\mathcal{J} \subseteq \mathbb{R}$ be a finite or infinite interval, let $f: \mathcal{J} \rightarrow \mathbb{R}$, assume that f is continuous, and assume that, for all $x, y \in \mathcal{J}$, it follows that $f[\frac{1}{2}(x+y)] \leq \frac{1}{2}f(x+y)$. Then, f is convex. (Proof: See [1039, p. 10].) (Remark: This result is due to Jensen.) (Remark: See Fact 1.8.4.)

Fact 10.11.8. Let $A_0 \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ be a norm on $\mathbb{F}^{n \times n}$, and let $\varepsilon > 0$. Then, there exists $\delta > 0$ such that, if $A \in \mathbb{F}^{n \times n}$ and $\|A - A_0\| < \delta$, then

$$\text{dist}[\text{mspec}(A) - \text{mspec}(A_0)] < \varepsilon,$$

where

$$\text{dist}[\text{mspec}(A) - \text{mspec}(A_0)] \triangleq \min_{\sigma} \max_{i=1, \dots, n} |\lambda_{\sigma(i)}(A) - \lambda_i(A_0)|$$

and the minimum is taken over all permutations σ of $\{1, \dots, n\}$. (Proof: See [690, p. 399].)

Fact 10.11.9. Let $\mathcal{J} \subseteq \mathbb{R}$ be an interval, let $A: \mathcal{J} \mapsto \mathbb{F}^{n \times n}$, and assume that A is continuous. Then, for $i = 1, \dots, n$, there exist continuous functions $\lambda_i: \mathcal{J} \mapsto \mathbb{C}$ such that, for all $t \in \mathcal{J}$, $\text{mspec}(A(t)) = \{\lambda_1(t), \dots, \lambda_n(t)\}_{\text{ms}}$. (Proof: See [690, p. 399].) (Remark: The spectrum cannot always be continuously parameterized by more than one variable. See [690, p. 399].)

Fact 10.11.10. Let $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $g: \mathbb{R} \rightarrow \mathbb{R}$, and $h: \mathbb{R} \rightarrow \mathbb{R}$. Then, assuming each of the following integrals exists,

$$\frac{d}{d\alpha} \int_{g(\alpha)}^{h(\alpha)} f(t, \alpha) dt = f(h(\alpha), \alpha)h'(\alpha) - f(g(\alpha), \alpha)g'(\alpha) + \int_{g(\alpha)}^{h(\alpha)} \frac{\partial}{\partial \alpha} f(t, \alpha) dt.$$

(Remark: This identity is *Leibniz's rule*.)

Fact 10.11.11. Let $\mathcal{D} \subseteq \mathbb{R}^m$, assume that $\mathcal{D}$ is a convex set, and let $f: \mathcal{D} \rightarrow \mathbb{R}$. Then, f is convex if and only if the set $\{(x, y) \in \mathbb{R}^n \times \mathbb{R}: y \geq f(x)\}$ is convex.

Fact 10.11.12. Let $\mathcal{D} \subseteq \mathbb{R}^m$, assume that $\mathcal{D}$ is a convex set, let $f: \mathcal{D} \rightarrow \mathbb{R}$, and assume that f is convex. Then, f is continuous on $\text{int}_{\text{aff}} \mathcal{D}$.

Fact 10.11.13. Let $\mathcal{D} \subseteq \mathbb{R}^m$, assume that $\mathcal{D}$ is a convex set, let $f: \mathcal{D} \rightarrow \mathbb{R}$, and assume that f is convex. Then, $f^{-1}((-\infty, \alpha]) = \{x \in \mathcal{D}: f(x) \leq \alpha\}$ is convex.

Fact 10.11.14. Let $\mathcal{D} \subseteq \mathbb{R}^m$, assume that $\mathcal{D}$ is open and convex, let $f: \mathcal{D} \rightarrow \mathbb{R}$, and assume that f is C^1 on $\mathcal{D}$. Then, the following statements hold:

i) f is convex if and only if, for all $x, y \in \mathcal{D}$,

$$f(x) + (y - x)^T f'(x) \leq f(y).$$

ii) f is strictly convex if and only if, for all distinct $x, y \in \mathcal{D}$,

$$f(x) + (y - x)^T f'(x) < f(y).$$

(Remark: If f is not differentiable, then these inequalities can be stated in terms of directional differentials of f or the *subdifferential* of f . See [1039, pp. 29–31, 128–145].)

Fact 10.11.15. Let $f: \mathcal{D} \subseteq \mathbb{F}^m \mapsto \mathbb{F}^n$, and assume that $D_+ f(0; \xi)$ exists. Then, for all $\beta > 0$,

$$D_+ f(0; \beta \xi) = \beta D_+ f(0; \xi).$$

Fact 10.11.16. Define $f: \mathbb{R} \rightarrow \mathbb{R}$ by $f(x) \triangleq |x|$. Then, for all $\xi \in \mathbb{R}$,

$$D_+ f(0; \xi) = |\xi|.$$

Now, define $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$ by $f(x) \triangleq \sqrt{x^T x}$. Then, for all $\xi \in \mathbb{R}^n$,

$$D_+ f(0; \xi) = \sqrt{\xi^T \xi}.$$

Fact 10.11.17. Let $A, B \in \mathbb{F}^{n \times n}$. Then, for all $s \in \mathbb{F}$,

$$\frac{d}{ds}(A + sB)^2 = AB + BA + 2sB.$$

Hence,

$$\left. \frac{d}{ds}(A + sB)^2 \right|_{s=0} = AB + BA.$$

Furthermore, for all $k \geq 1$,

$$\left. \frac{d}{ds}(A + sB)^k \right|_{s=0} = \sum_{i=0}^{k-1} A^i B A^{k-1-i}.$$

Fact 10.11.18. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\mathcal{D} \triangleq \{s \in \mathbb{F} : \det(A + sB) \neq 0\}$. Then, for all $s \in \mathcal{D}$,

$$\frac{d}{ds}(A + sB)^{-1} = -(A + sB)^{-1}B(A + sB)^{-1}.$$

Hence, if A is nonsingular, then

$$\left. \frac{d}{ds}(A + sB)^{-1} \right|_{s=0} = -A^{-1}BA^{-1}.$$

Fact 10.11.19. Let $\mathcal{D} \subseteq \mathbb{F}$, let $A: \mathcal{D} \longrightarrow \mathbb{F}^{n \times n}$, and assume that A is differentiable. Then,

$$\frac{d}{ds} \det A(s) = \operatorname{tr} \left[A^A(s) \frac{d}{ds} A(s) \right] = \frac{1}{n-1} \operatorname{tr} \left[A(s) \frac{d}{ds} A^A(s) \right] = \sum_{i=1}^n \det A_i(s),$$

where $A_i(s)$ is obtained by differentiating the entries of the i th row of $A(s)$. If, in addition, $A(s)$ is nonsingular for all $s \in \mathcal{D}$, then

$$\frac{d}{ds} \log \det A(s) = \operatorname{tr} \left[A^{-1}(s) \frac{d}{ds} A(s) \right].$$

If $A(s)$ is positive definite for all $s \in \mathcal{D}$, then

$$\frac{d}{ds} \det A^{1/n}(s) = \frac{1}{n} [\det A^{1/n}(s)] \operatorname{tr} \left[A^{-1}(s) \frac{d}{ds} A(s) \right].$$

Finally, if $A(s)$ is nonsingular and has no negative eigenvalues for all $s \in \mathcal{D}$, then

$$\frac{d}{ds} \log^2 A(s) = 2 \operatorname{tr} \left[[\log A(s)] A^{-1}(s) \frac{d}{ds} A(s) \right]$$

and

$$\frac{d}{ds} \log A(s) = \int_0^1 [(A(s) - I)t + I]^{-1} \frac{d}{ds} A(s) [(A(s) - I)t + I]^{-1} dt.$$

(Proof: See [359, p. 267], [563], [1014], [1098, pp. 199, 212], [1129, p. 430], and [1183].) (Remark: See Fact 11.13.4.)

Fact 10.11.20. Let $\mathcal{D} \subseteq \mathbb{F}$, let $A: \mathcal{D} \rightarrow \mathbb{F}^{n \times n}$, assume that A is differentiable, and assume that $A(s)$ is nonsingular for all $s \in \mathcal{D}$. Then,

$$\frac{d}{ds}A^{-1}(s) = -A^{-1}(s) \left[\frac{d}{ds}A(s) \right] A^{-1}(s)$$

and

$$\operatorname{tr} \left[A^{-1}(s) \frac{d}{ds}A(s) \right] = -\operatorname{tr} \left[A(s) \frac{d}{ds}A^{-1}(s) \right].$$

(Proof: See [711, p. 491] and [1098, pp. 198, 212].)

Fact 10.11.21. Let $A, B \in \mathbb{F}^{n \times n}$. Then, for all $s \in \mathbb{F}$,

$$\frac{d}{ds} \det(A + sB) = \operatorname{tr}[B(A + sB)^A].$$

Hence,

$$\left. \frac{d}{ds} \det(A + sB) \right|_{s=0} = \operatorname{tr} BA^A = \sum_{i=1}^n \det[A \stackrel{i}{\leftarrow} \operatorname{col}_i(B)].$$

(Proof: Use Fact 10.11.19 and Fact 2.16.9.) (Remark: This result generalizes Lemma 4.4.8.)

Fact 10.11.22. Let $A \in \mathbb{F}^{n \times n}$, $r \in \mathbb{R}$, and $k \geq 1$. Then, for all $s \in \mathbb{C}$,

$$\frac{d^k}{ds^k} [\det(I + sA)]^r = (r \operatorname{tr} A)^k [\det(I + sA)]^r.$$

Hence,

$$\left. \frac{d^k}{ds^k} [\det(I + sA)]^r \right|_{s=0} = (r \operatorname{tr} A)^k.$$

Fact 10.11.23. Let $A \in \mathbb{R}^{n \times n}$, assume that A is symmetric, let $X \in \mathbb{R}^{m \times n}$, and assume that XAX^T is nonsingular. Then,

$$\left(\frac{d}{dX} \det XAX^T \right) = 2(\det XAX^T) A^T X^T (XAX^T)^{-1}.$$

(Proof: See [350].)

Fact 10.11.24. The following infinite series converge for $A \in \mathbb{F}^{n \times n}$ with the given bounds on $\operatorname{sprad}(A)$:

i) For all $A \in \mathbb{F}^{n \times n}$,

$$\sin A = A - \frac{1}{3!}A^3 + \frac{1}{5!}A^5 - \frac{1}{7!}A^7 + \cdots.$$

ii) For all $A \in \mathbb{F}^{n \times n}$,

$$\cos A = I - \frac{1}{2!}A^2 + \frac{1}{4!}A^4 - \frac{1}{6!}A^6 + \cdots.$$

iii) For all $A \in \mathbb{F}^{n \times n}$ such that $\operatorname{sprad}(A) < \pi/2$,

$$\tan A = A + \frac{1}{3}A^3 + \frac{2}{15}A^5 + \frac{17}{315}A^7 + \frac{62}{2835}A^9 + \cdots.$$

iv) For all $A \in \mathbb{F}^{n \times n}$ such that $\operatorname{sprad}(A) < 1$,

$$e^A = I + A + \frac{1}{2!}A^2 + \frac{1}{3!}A^3 + \frac{1}{4!}A^4 + \cdots.$$

v) For all $A \in \mathbb{F}^{n \times n}$ such that $\text{sprad}(A - I) < 1$,

$$\log A = -\left[I - A + \frac{1}{2}(I - A)^2 + \frac{1}{3}(I - A)^3 + \frac{1}{4}(I - A)^4 + \cdots\right].$$

vi) For all $A \in \mathbb{F}^{n \times n}$ such that $\text{sprad}(A) < 1$,

$$\log(I - A) = -(A + \frac{1}{2}A^2 + \frac{1}{3}A^3 + \frac{1}{4}A^4 + \cdots).$$

vii) For all $A \in \mathbb{F}^{n \times n}$ such that $\text{sprad}(A) < 1$,

$$\log(I + A) = A - \frac{1}{2}A^2 + \frac{1}{3}A^3 - \frac{1}{4}A^4 + \cdots.$$

viii) For all $A \in \mathbb{F}^{n \times n}$ such that $\text{spec}(A) \subset \text{ORHP}$,

$$\log A = \sum_{i=0}^{\infty} \frac{2}{2i+1} [(A - I)(A + I)^{-1}]^{2i+1}.$$

ix) For all $A \in \mathbb{F}^{n \times n}$,

$$\sinh A = \sin jA = A + \frac{1}{3!}A^3 + \frac{1}{5!}A^5 + \frac{1}{7!}A^7 + \cdots.$$

x) For all $A \in \mathbb{F}^{n \times n}$,

$$\cosh A = \cos jA = I + \frac{1}{2!}A^2 + \frac{1}{4!}A^4 + \frac{1}{6!}A^6 + \cdots.$$

xi) For all $A \in \mathbb{F}^{n \times n}$ such that $\text{sprad}(A) < \pi/2$,

$$\tanh A = \tan jA = A - \frac{1}{3}A^3 + \frac{2}{15}A^5 - \frac{17}{315}A^7 + \frac{62}{2835}A^9 - \cdots.$$

xii) Let $\alpha \in \mathbb{R}$. For all $A \in \mathbb{F}^{n \times n}$ such that $\text{sprad}(A) < 1$,

$$\begin{aligned} (I + A)^\alpha &= I + \alpha A + \frac{\alpha(\alpha-1)}{2!}A^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!}A^3 + \frac{1}{4}A^4 + \cdots \\ &= I + \binom{\alpha}{1}A + \binom{\alpha}{2}A^2 + \binom{\alpha}{3}A^3 + \binom{\alpha}{4}A^4 + \cdots. \end{aligned}$$

xiii) For all $A \in \mathbb{F}^{n \times n}$ such that $\text{sprad}(A) < 1$,

$$(I - A)^{-1} = I + A + A^2 + A^3 + A^4 + \cdots.$$

(Proof: See Fact 1.18.8.)

10.12 Notes

An introductory treatment of limits and continuity is given in [1030]. Fréchet and directional derivatives are discussed in [496], while differentiation of matrix functions is considered in [654, 948, 975, 1089, 1136, 1182]. In [1133, 1134] the set $\text{int}_{\text{aff } \mathcal{S}} \mathcal{S}$ is called the relative interior of $\mathcal{S}$. An extensive treatment of matrix functions is given in Chapter 6 of [711]; see also [716]. The identity theorem is discussed in [741]. The chain rule for matrix functions is considered in [948, 980]. Differentiation with respect to complex matrices is discussed in [776]. Extensive tables of derivatives of matrix functions are given in [374, pp. 586–593].

Chapter Eleven

The Matrix Exponential and Stability Theory

The matrix exponential function is fundamental to the study of linear ordinary differential equations. This chapter focuses on the properties of the matrix exponential as well as on stability theory.

11.1 Definition of the Matrix Exponential

The scalar initial value problem

$$\dot{x}(t) = ax(t), \quad (11.1.1)$$

$$x(0) = x_0, \quad (11.1.2)$$

where $t \in [0, \infty)$ and $a, x(t) \in \mathbb{R}$, has the solution

$$x(t) = e^{at}x_0, \quad (11.1.3)$$

where $t \in [0, \infty)$. We are interested in systems of linear differential equations of the form

$$\dot{x}(t) = Ax(t), \quad (11.1.4)$$

$$x(0) = x_0, \quad (11.1.5)$$

where $t \in [0, \infty)$, $x(t) \in \mathbb{R}^n$, and $A \in \mathbb{R}^{n \times n}$. Here $\dot{x}(t)$ denotes $\frac{dx(t)}{dt}$, where the derivative is one sided for $t = 0$ and two sided for $t > 0$. The solution of (11.1.4), (11.1.5) is given by

$$x(t) = e^{tA}x_0, \quad (11.1.6)$$

where $t \in [0, \infty)$ and e^{tA} is the *matrix exponential*. The following definition is based on (10.5.2).

Definition 11.1.1. Let $A \in \mathbb{F}^{n \times n}$. Then, the *matrix exponential* $e^A \in \mathbb{F}^{n \times n}$ or $\exp(A) \in \mathbb{F}^{n \times n}$ is the matrix

$$e^A \triangleq \sum_{k=0}^{\infty} \frac{1}{k!} A^k. \quad (11.1.7)$$

Note that $0! \triangleq 1$ and $e^{0_{n \times n}} = I_n$.

Proposition 11.1.2. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) The series (11.1.7) converges absolutely.
- ii) The series (11.1.7) converges to e^A .
- iii) Let $\|\cdot\|$ be a normalized submultiplicative norm on $\mathbb{F}^{n \times n}$. Then,

$$e^{-\|A\|} \leq \|e^A\| \leq e^{\|A\|}. \quad (11.1.8)$$

Proof. To prove i), let $\|\cdot\|$ be a normalized submultiplicative norm on $\mathbb{F}^{n \times n}$. Then, for all $k \geq 1$,

$$\sum_{i=0}^k \frac{1}{i!} \|A^i\| \leq \sum_{i=0}^k \frac{1}{i!} \|A\|^i \leq e^{\|A\|}.$$

Since the sequence $\{\sum_{i=0}^k \frac{1}{i!} \|A^i\|\}_{i=0}^\infty$ of partial sums is increasing and bounded, there exists $\alpha > 0$ such that the series $\sum_{i=0}^\infty \frac{1}{i!} \|A^i\|$ converges to α . Hence, the series $\sum_{i=0}^\infty \frac{1}{i!} A^i$ converges absolutely.

Next, ii) follows from i) using Proposition 10.2.9.

Next, we have

$$\|e^A\| = \left\| \sum_{i=0}^\infty \frac{1}{i!} A^i \right\| \leq \sum_{i=0}^\infty \frac{1}{i!} \|A^i\| \leq \sum_{i=0}^\infty \frac{1}{i!} \|A\|^i = e^{\|A\|},$$

which verifies (11.1.8). Finally, note that

$$1 \leq \|e^A\| \|e^{-A}\| \leq \|e^A\| e^{\|A\|},$$

and thus

$$e^{-\|A\|} \leq \|e^A\|. \quad \square$$

The following result generalizes the well-known scalar result.

Proposition 11.1.3. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$e^A = \lim_{k \rightarrow \infty} \left(I + \frac{1}{k} A\right)^k. \quad (11.1.9)$$

Proof. It follows from the binomial theorem that

$$\left(I + \frac{1}{k} A\right)^k = \sum_{i=0}^k \alpha_i(k) A^i,$$

where

$$\alpha_i(k) \triangleq \frac{1}{k^i} \binom{k}{i} = \frac{1}{k^i} \frac{k!}{i!(k-i)!}.$$

For all $i \in \mathbb{P}$, it follows that $\alpha_i(k) \rightarrow 1/i!$ as $k \rightarrow \infty$. Hence,

$$\lim_{k \rightarrow \infty} \left(I + \frac{1}{k} A\right)^k = \lim_{k \rightarrow \infty} \sum_{i=0}^k \alpha_i(k) A^i = \sum_{i=0}^\infty \frac{1}{i!} A^i = e^A. \quad \square$$

Proposition 11.1.4. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $t \in \mathbb{R}$,

$$e^{tA} - I = \int_0^t A e^{\tau A} d\tau \quad (11.1.10)$$

and

$$\frac{d}{dt} e^{tA} = A e^{tA}. \quad (11.1.11)$$

Proof. Note that

$$\int_0^t A e^{\tau A} d\tau = \int_0^t \sum_{k=0}^{\infty} \frac{1}{k!} \tau^k A^{k+1} d\tau = \sum_{k=0}^{\infty} \frac{1}{k!} \frac{t^{k+1}}{k+1} A^{k+1} = e^{tA} - I,$$

which yields (11.1.10), while differentiating (11.1.10) with respect to t yields (11.1.11). $\square$

Proposition 11.1.5. Let $A, B \in \mathbb{F}^{n \times n}$. Then, $AB = BA$ if and only if, for all $t \in [0, \infty)$,

$$e^{tA} e^{tB} = e^{t(A+B)}. \quad (11.1.12)$$

Proof. Suppose that $AB = BA$. By expanding e^{tA} , e^{tB} , and $e^{t(A+B)}$, it can be seen that the expansions of $e^{tA} e^{tB}$ and $e^{t(A+B)}$ are identical. Conversely, differentiating (11.1.12) twice with respect to t and setting $t = 0$ yields $AB = BA$. $\square$

Corollary 11.1.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $AB = BA$. Then,

$$e^A e^B = e^B e^A = e^{A+B}. \quad (11.1.13)$$

The converse of Corollary 11.1.6 is not true. For example, if $A \triangleq \begin{bmatrix} 0 & \pi \\ -\pi & 0 \end{bmatrix}$ and $B \triangleq \begin{bmatrix} 0 & (7+4\sqrt{3})\pi \\ (-7+4\sqrt{3})\pi & 0 \end{bmatrix}$, then $e^A = e^B = -I$ and $e^{A+B} = I$, although $AB \neq BA$. A partial converse is given by Fact 11.14.2.

Proposition 11.1.7. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$e^{A \otimes I_m} = e^A \otimes I_m, \quad (11.1.14)$$

$$e^{I_n \otimes B} = I_n \otimes e^B, \quad (11.1.15)$$

$$e^{A \oplus B} = e^A \otimes e^B. \quad (11.1.16)$$

Proof. Note that

$$\begin{aligned} e^{A \otimes I_m} &= I_{nm} + A \otimes I_m + \frac{1}{2!} (A \otimes I_m)^2 + \cdots \\ &= I_n \otimes I_m + A \otimes I_m + \frac{1}{2!} (A^2 \otimes I_m) + \cdots \\ &= (I_n + A + \frac{1}{2!} A^2 + \cdots) \otimes I_m \\ &= e^A \otimes I_m \end{aligned}$$

and similarly for (11.1.15). To prove (11.1.16), note that $(A \otimes I_m)(I_n \otimes B) = A \otimes B$ and $(I_n \otimes B)(A \otimes I_m) = A \otimes B$, which shows that $A \otimes I_m$ and $I_n \otimes B$ commute. Thus, by Corollary 11.1.6,

$$e^{A \oplus B} = e^{A \otimes I_m + I_n \otimes B} = e^{A \otimes I_m} e^{I_n \otimes B} = (e^A \otimes I_m)(I_n \otimes e^B) = e^A \otimes e^B. \quad \square$$

11.2 Structure of the Matrix Exponential

To elucidate the structure of the matrix exponential, recall that, by Theorem 4.6.1, every term A^k in (11.1.7) for $k > r \triangleq \deg \mu_A$ can be expressed as a linear combination of $I, A, \dots, A^{r-1}$. The following result provides an expression for e^{tA} in terms of $I, A, \dots, A^{r-1}$.

Proposition 11.2.1. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $t \in \mathbb{R}$,

$$e^{tA} = \frac{1}{2\pi j} \oint_{\mathcal{C}} (zI - A)^{-1} e^{tz} dz = \sum_{i=0}^{n-1} \psi_i(t) A^i, \quad (11.2.1)$$

where, for all $i = 0, \dots, n-1$, $\psi_i(t)$ is given by

$$\psi_i(t) \triangleq \frac{1}{2\pi j} \oint_{\mathcal{C}} \frac{\chi_A^{[i+1]}(z)}{\chi_A(z)} e^{tz} dz, \quad (11.2.2)$$

where $\mathcal{C}$ is a simple, closed contour in the complex plane enclosing $\text{spec}(A)$,

$$\chi_A(s) = s^n + \beta_{n-1}s^{n-1} + \dots + \beta_1s + \beta_0, \quad (11.2.3)$$

and the polynomials $\chi_A^{[1]}, \dots, \chi_A^{[n]}$ are defined by the recursion

$$s\chi_A^{[i+1]}(s) = \chi_A^{[i]}(s) - \beta_i, \quad i = 0, \dots, n-1,$$

where $\chi_A^{[0]} \triangleq \chi_A$ and $\chi_A^{[n]}(s) = 1$. Furthermore, for all $i = 0, \dots, n-1$ and $t \geq 0$, $\psi_i(t)$ satisfies

$$\psi_i^{(n)}(t) + \beta_{n-1}\psi_i^{(n-1)}(t) + \dots + \beta_1\psi_i'(t) + \beta_0\psi_i(t) = 0, \quad (11.2.4)$$

where, for all $i, j = 0, \dots, n-1$,

$$\psi_i^{(j)}(0) = \delta_{ij}. \quad (11.2.5)$$

Proof. See [569, p. 381], [888, 929], [1455, p. 31], and Fact 4.9.11. $\square$

The coefficient $\psi_i(t)$ of A^i in (11.2.1) can be further characterized in terms of the Laplace transform. Define

$$\hat{x}(s) \triangleq \mathcal{L}\{x(t)\} \triangleq \int_0^\infty e^{-st}x(t) dt. \quad (11.2.6)$$

Note that

$$\mathcal{L}\{\dot{x}(t)\} = s\hat{x}(s) - x(0) \quad (11.2.7)$$

and

$$\mathcal{L}\{\ddot{x}(t)\} = s^2\hat{x}(s) - sx(0) - \dot{x}(0). \quad (11.2.8)$$

The following result shows that the resolvent of A is the Laplace transform of the exponential of A . See (4.4.23).

Proposition 11.2.2. Let $A \in \mathbb{F}^{n \times n}$, and define $\psi_0, \dots, \psi_{n-1}$ as in Proposition 11.2.1. Then, for all $s \in \mathbb{C} \setminus \text{spec}(A)$,

$$\mathcal{L}\{e^{tA}\} = \int_0^\infty e^{-st} e^{tA} dt = (sI - A)^{-1}. \quad (11.2.9)$$

Furthermore, for all $i = 0, \dots, n-1$, the Laplace transform $\hat{\psi}_i(s)$ of $\psi_i(t)$ is given by

$$\hat{\psi}_i(s) = \frac{\chi_A^{[i+1]}(s)}{\chi_A(s)} \quad (11.2.10)$$

and

$$(sI - A)^{-1} = \sum_{i=0}^{n-1} \hat{\psi}_i(s) A^i. \quad (11.2.11)$$

Proof. Let $s \in \mathbb{C}$ satisfy $\text{Re } s > \text{spabs}(A)$ so that $A - sI$ is asymptotically stable. Thus, it follows from Lemma 11.9.2 that

$$\mathcal{L}\{e^{tA}\} = \int_0^\infty e^{-st} e^{tA} dt = \int_0^\infty e^{t(A-sI)} dt = (sI - A)^{-1}.$$

By analytic continuation, the expression $\mathcal{L}\{e^{tA}\}$ is given by (11.2.9) for all $s \in \mathbb{C} \setminus \text{spec}(A)$. $\square$

Comparing (11.2.11) with (4.4.23) yields

$$\sum_{i=0}^{n-1} \hat{\psi}_i(s) A^i = \frac{s^{n-1}}{\chi_A(s)} I + \frac{s^{n-2}}{\chi_A(s)} B_{n-2} + \dots + \frac{s}{\chi_A(s)} B_1 + B_0. \quad (11.2.12)$$

To further illustrate the structure of e^{tA} , where $A \in \mathbb{F}^{n \times n}$, let $A = SBS^{-1}$, where $B = \text{diag}(B_1, \dots, B_k)$ is the Jordan form of A . Hence, by Proposition 11.2.8,

$$e^{tA} = S e^{tB} S^{-1}, \quad (11.2.13)$$

where

$$e^{tB} = \text{diag}(e^{tB_1}, \dots, e^{tB_k}). \quad (11.2.14)$$

The structure of e^{tB} can thus be determined by considering the block $B_i \in \mathbb{F}^{\alpha_i \times \alpha_i}$, which, for all $i = 1, \dots, k$, has the form

$$B_i = \lambda_i I_{\alpha_i} + N_{\alpha_i}. \quad (11.2.15)$$

Since $\lambda_i I_{\alpha_i}$ and N_{α_i} commute, it follows from Proposition 11.1.5 that

$$e^{tB_i} = e^{t(\lambda_i I_{\alpha_i} + N_{\alpha_i})} = e^{\lambda_i t I_{\alpha_i}} e^{t N_{\alpha_i}} = e^{\lambda_i t} e^{t N_{\alpha_i}}. \quad (11.2.16)$$

Since $N_{\alpha_i}^{\alpha_i} = 0$, it follows that $e^{t N_{\alpha_i}}$ is a finite sum of powers of $t N_{\alpha_i}$. Specifically,

$$e^{t N_{\alpha_i}} = I_{\alpha_i} + t N_{\alpha_i} + \frac{1}{2} t^2 N_{\alpha_i}^2 + \dots + \frac{1}{(\alpha_i - 1)!} t^{\alpha_i - 1} N_{\alpha_i}^{\alpha_i - 1}, \quad (11.2.17)$$

and thus

$$e^{tN_{\alpha_i}} = \begin{bmatrix} 1 & t & \frac{t^2}{2} & \cdots & \frac{t^{\alpha_i-1}}{(\alpha_i-1)!} \\ 0 & 1 & t & \ddots & \frac{t^{\alpha_i-2}}{(\alpha_i-2)!} \\ 0 & 0 & 1 & \ddots & \frac{t^{\alpha_i-3}}{(\alpha_i-3)!} \\ \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{bmatrix}, \quad (11.2.18)$$

which is upper triangular and Toeplitz (see Fact 11.13.1). Alternatively, (11.2.18) follows from (10.5.5) with $f(s) = e^{st}$.

Note that (11.2.16) follows from (10.5.5) with $f(\lambda) = e^{\lambda t}$. Furthermore, every entry of e^{tB_i} is of the form $\frac{1}{r!} t^r e^{\lambda_i t}$, where $r \in \{0, \alpha_i - 1\}$ and λ_i is an eigenvalue of A . Reconstructing A by means of $A = SBS^{-1}$ shows that every entry of A is a linear combination of the entries of the blocks e^{tB_i} . If A is real, then e^{tA} is also real. Thus, the term $e^{\lambda_i t}$ for complex $\lambda_i = \nu_i + j\omega_i \in \text{spec}(A)$, where ν_i and ω_i are real, yields terms of the form $e^{\nu_i t} \cos \omega_i t$ and $e^{\nu_i t} \sin \omega_i t$.

The following result follows from (11.2.18) or Corollary 10.5.4.

Proposition 11.2.3. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\text{mspec}(e^A) = \{e^\lambda : \lambda \in \text{mspec}(A)\}_{\text{ms}}. \quad (11.2.19)$$

Proof. It can be seen that every diagonal entry of the Jordan form of e^A is of the form e^λ , where $\lambda \in \text{spec}(A)$. $\square$

Corollary 11.2.4. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\det e^A = e^{\text{tr } A}. \quad (11.2.20)$$

Corollary 11.2.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{tr } A = 0$. Then, $\det e^A = 1$.

Corollary 11.2.6. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) If e^A is unitary, then, $\text{spec}(A) \subset j\mathbb{R}$.
- ii) $\text{spec}(e^A)$ is real if and only if $\text{Im spec}(A) \subset \pi\mathbb{Z}$.

Proposition 11.2.7. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) A and e^A have the same number of Jordan blocks of corresponding sizes.
- ii) e^A is semisimple if and only if A is semisimple.
- iii) If $\mu \in \text{spec}(e^A)$, then

$$\text{am}_{\exp(A)}(\mu) = \sum_{\{\lambda \in \text{spec}(A) : e^\lambda = \mu\}} \text{am}_A(\lambda) \quad (11.2.21)$$

and

$$\text{gm}_{\exp(A)}(\mu) = \sum_{\{\lambda \in \text{spec}(A): e^\lambda = \mu\}} \text{gm}_A(\lambda). \quad (11.2.22)$$

- iv)* If e^A is simple, then A is simple.
- v)* If e^A is cyclic, then A is cyclic.
- vi)* e^A is a multiple of the identity if and only if A is semisimple and every pair of eigenvalues of A differs by an integer multiple of $2\pi j$.
- vii)* e^A is a real multiple of the identity if and only if A is semisimple, every pair of eigenvalues of A differs by an integer multiple of $2\pi j$, and the imaginary part of every eigenvalue of A is an integer multiple of πj .

Proof. To prove *i)*, note that, for all $t \neq 0$, $\text{def}(e^{tN_{\alpha_i}} - I_{\alpha_i}) = 1$, and thus the geometric multiplicity of (11.2.18) is 1. Since (11.2.18) has one distinct eigenvalue, it follows that (11.2.18) is cyclic. Hence, by Proposition 5.5.15, (11.2.18) is similar to a single Jordan block. Now, *i)* follows by setting $t = 1$ and applying this argument to each Jordan block of A . Statements *ii)*–*v)* follow by similar arguments.

To prove *vi)*, note that, for all $\lambda_i, \lambda_j \in \text{spec}(A)$, it follows that $e^{\lambda_i} = e^{\lambda_j}$. Furthermore, since A is semisimple, it follows from *ii)* that e^A is also semisimple. Since all of the eigenvalues of e^A are equal, it follows that e^A is a multiple of the identity. Finally, *viii)* is an immediate consequence of *vii)*. $\square$

Proposition 11.2.8. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i)* $(e^A)^T = e^{A^T}$.
- ii)* $(\overline{e^A}) = \overline{e^A}$.
- iii)* $(e^A)^* = e^{A^*}$.
- iv)* e^A is nonsingular, and $(e^A)^{-1} = e^{-A}$.
- v)* If $S \in \mathbb{F}^{n \times n}$ is nonsingular, then $e^{SAS^{-1}} = Se^AS^{-1}$.
- vi)* If $A = \text{diag}(A_1, \dots, A_k)$, where $A_i \in \mathbb{F}^{n_i \times n_i}$ for all $i = 1, \dots, k$, then $e^A = \text{diag}(e^{A_1}, \dots, e^{A_k})$.
- vii)* If A is Hermitian, then e^A is positive definite.
- viii)* e^A is Hermitian if and only if A is unitarily similar to a block-diagonal matrix $\text{diag}(A_1, \dots, A_k)$ such that, for all $i = 1, \dots, k$, e^{A_i} is a real multiple of the identity and, for all distinct $i, j = 1, \dots, k$, $\text{spec}(e^{A_i}) \neq \text{spec}(e^{A_j})$.

Furthermore, the following statements are equivalent:

- ix)* A is normal.
- x)* $\text{tr } e^{A^*} e^A = \text{tr } e^{A^* + A}$.
- xi)* $e^{A^*} e^A = e^{A^* + A}$.
- xii)* $e^A e^{A^*} = e^{A^*} e^A = e^{A^* + A}$.

xiii) A is unitarily similar to a block-diagonal matrix $\text{diag}(A_1, \dots, A_k)$ such that, for all $i = 1, \dots, k$, e^{A_i} is a multiple of the identity and, for all distinct $i, j = 1, \dots, k$, $\text{spec}(e^{A_i}) \neq \text{spec}(e^{A_j})$.

Finally, the following statements hold:

- xiv)* If A is normal, then e^A is normal.
- xv)* If e^A is normal and no pair of eigenvalues of A differ by an integer multiple of $2\pi j$, then A is normal.
- xvi)* A is skew Hermitian if and only if A is normal and e^A is unitary.
- xvii)* If $\mathbb{F} = \mathbb{R}$ and A is skew symmetric, then e^A is orthogonal and $\det e^A = 1$.
- xviii)* e^A is unitary if and only if A is unitarily similar to a block-diagonal matrix $\text{diag}(A_1, \dots, A_k)$ such that, for all $i = 1, \dots, k$, e^{A_i} is a unit-absolute-value multiple of the identity and, for all distinct $i, j = 1, \dots, k$, $\text{spec}(e^{A_i}) \neq \text{spec}(e^{A_j})$.
- xix)* If e^A is unitary, then either A is skew Hermitian or at least two eigenvalues of A differ by a nonzero integer multiple of $2\pi j$.

Proof. The equivalence of *ix)* and *x)* is given in [452, 1208], while the equivalence of *ix)* and *xii)* is given in [1172]. Note that $xii) \implies xi) \implies x)$. Statement *xiv)* follows from the fact that $ix) \implies xii)$. The equivalence of *ix)* and *xiii)* is given in [1468]; statement *xviii)* is analogous. To prove sufficiency in *xvi)*, note that $e^{A+A^*} = e^A e^{A^*} = e^A (e^A)^* = I = e^0$. Since $A + A^*$ is Hermitian, it follows from *iii)* of Proposition 11.2.9 that $A + A^* = 0$. To prove *xix)*, it follows from *xvii)* that, if every block A_i is scalar, then A is skew Hermitian, while, if at least one block A_i is not scalar, then A has at least two eigenvalues that differ by an integer multiple of $2\pi j$. $\square$

The converse of *ix)* is false. For example, the matrix $A \triangleq \begin{bmatrix} -2\pi & 4\pi \\ -2\pi & 2\pi \end{bmatrix}$ satisfies $e^A = I$ but is not normal. Likewise, $A = \begin{bmatrix} j\pi & 1 \\ 0 & -j\pi \end{bmatrix}$ satisfies $e^A = -I$ but is not normal. For both matrices, $e^{A^*} e^A = e^A e^{A^*} = I$, but $e^{A^*} e^A \neq e^{A^*+A}$, which is consistent with *xii)*. Both matrices have eigenvalues $\pm j\pi$.

Proposition 11.2.9. The following statements hold:

- i)* If $A, B \in \mathbb{F}^{n \times n}$ are similar, then e^A and e^B are similar.
- ii)* If $A, B \in \mathbb{F}^{n \times n}$ are unitarily similar, then e^A and e^B are unitarily similar.
- iii)* $B \in \mathbb{F}^{n \times n}$ is positive definite if and only if there exists a unique Hermitian matrix $A \in \mathbb{F}^{n \times n}$ such that $e^A = B$.
- iv)* $B \in \mathbb{F}^{n \times n}$ is Hermitian and nonsingular if and only if there exists a normal matrix $A \in \mathbb{C}^{n \times n}$ such that, for all $\lambda \in \text{spec}(A)$, $\text{Im } \lambda$ is an integer multiple of πj and $e^A = B$.
- v)* $B \in \mathbb{F}^{n \times n}$ is normal and nonsingular if and only if there exists a normal matrix $A \in \mathbb{F}^{n \times n}$ such that $e^A = B$.
- vi)* $B \in \mathbb{F}^{n \times n}$ is unitary if and only if there exists a normal matrix $A \in \mathbb{C}^{n \times n}$

such that $\text{mspec}(A) \subset j\mathbb{R}$ and $e^A = B$.

vii) $B \in \mathbb{F}^{n \times n}$ is unitary if and only if there exists a skew-Hermitian matrix $A \in \mathbb{C}^{n \times n}$ such that $e^A = B$.

viii) $B \in \mathbb{F}^{n \times n}$ is unitary if and only if there exists a Hermitian matrix $A \in \mathbb{F}^{n \times n}$ such that $e^{jA} = B$.

ix) $B \in \mathbb{R}^{n \times n}$ is orthogonal and $\det B = 1$ if and only if there exists a skew-symmetric matrix $A \in \mathbb{R}^{n \times n}$ such that $e^A = B$.

x) If A and B are normal and $e^A = e^B$, then $A + A^* = B + B^*$.

Proof. Statement *iii*) is given by Proposition 11.4.5. Statement *vii*) is given by *v*) of Proposition 11.6.7. To prove *x*), note that $e^{A+A^*} = e^{B+B^*}$, which, by *vii*) of Proposition 11.2.8, is positive definite. The result now follows from *iii*). $\square$

The converse of *i*) is false. For example, $A \triangleq \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ and $B \triangleq \begin{bmatrix} 0 & 2\pi \\ -2\pi & 0 \end{bmatrix}$ satisfy $e^A = e^B = I$, although A and B are not similar.

11.3 Explicit Expressions

In this section we present explicit expressions for the exponential of a general 2×2 real matrix A . Expressions are given in terms of both the entries of A and the eigenvalues of A .

Lemma 11.3.1. Let $A \triangleq \begin{bmatrix} a & b \\ 0 & d \end{bmatrix} \in \mathbb{C}^{2 \times 2}$. Then,

$$e^A = \begin{cases} e^a \begin{bmatrix} 1 & b \\ 0 & 1 \end{bmatrix}, & a = d, \\ \begin{bmatrix} e^a & b \frac{e^a - e^d}{a - d} \\ 0 & e^d \end{bmatrix}, & a \neq d. \end{cases} \quad (11.3.1)$$

The following result gives an expression for e^A in terms of the eigenvalues of A .

Proposition 11.3.2. Let $A \in \mathbb{C}^{2 \times 2}$, and let $\text{mspec}(A) = \{\lambda, \mu\}_{\text{ms}}$. Then,

$$e^A = \begin{cases} e^\lambda [(1 - \lambda)I + A], & \lambda = \mu, \\ \frac{\mu e^\lambda - \lambda e^\mu}{\mu - \lambda} I + \frac{e^\mu - e^\lambda}{\mu - \lambda} A, & \lambda \neq \mu. \end{cases} \quad (11.3.2)$$

Proof. The result follows from Theorem 10.5.2. Alternatively, suppose that $\lambda = \mu$. Then, there exists a nonsingular matrix $S \in \mathbb{C}^{2 \times 2}$ such that $A = S \begin{bmatrix} \lambda & \alpha \\ 0 & \lambda \end{bmatrix} S^{-1}$, where $\alpha \in \mathbb{C}$. Hence, $e^A = e^\lambda S \begin{bmatrix} 1 & \alpha \\ 0 & 1 \end{bmatrix} S^{-1} = e^\lambda [(1 - \lambda)I + A]$. Now, suppose that $\lambda \neq \mu$. Then, there exists a nonsingular matrix $S \in \mathbb{C}^{2 \times 2}$ such that $A = S \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix} S^{-1}$. Hence, $e^A = S \begin{bmatrix} e^\lambda & 0 \\ 0 & e^\mu \end{bmatrix} S^{-1}$. Then, the identity $\begin{bmatrix} e^\lambda & 0 \\ 0 & e^\mu \end{bmatrix} = \frac{\mu e^\lambda - \lambda e^\mu}{\mu - \lambda} I + \frac{e^\mu - e^\lambda}{\mu - \lambda} \begin{bmatrix} \lambda & 0 \\ 0 & \mu \end{bmatrix}$

yields the desired result. $\square$

Next, we give an expression for e^A in terms of the entries of $A \in \mathbb{R}^{2 \times 2}$.

Corollary 11.3.3. Let $A \triangleq \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbb{R}^{2 \times 2}$, and define $\gamma \triangleq (a-d)^2 + 4bc$ and $\delta \triangleq \frac{1}{2}|\gamma|^{1/2}$. Then,

$$e^A = \begin{cases} e^{\frac{a+d}{2}} \begin{bmatrix} \cos \delta + \frac{a-d}{2\delta} \sin \delta & \frac{b}{\delta} \sin \delta \\ \frac{c}{\delta} \sin \delta & \cos \delta - \frac{a-d}{2\delta} \sin \delta \end{bmatrix}, & \gamma < 0, \\ e^{\frac{a+d}{2}} \begin{bmatrix} 1 + \frac{a-d}{2} & b \\ c & 1 - \frac{a-d}{2} \end{bmatrix}, & \gamma = 0, \\ e^{\frac{a+d}{2}} \begin{bmatrix} \cosh \delta + \frac{a-d}{2\delta} \sinh \delta & \frac{b}{\delta} \sinh \delta \\ \frac{c}{\delta} \sinh \delta & \cosh \delta - \frac{a-d}{2\delta} \sinh \delta \end{bmatrix}, & \gamma > 0. \end{cases} \quad (11.3.3)$$

Proof. The eigenvalues of A are $\lambda \triangleq \frac{1}{2}(a+d-\sqrt{\gamma})$ and $\mu \triangleq \frac{1}{2}(a+d+\sqrt{\gamma})$. Hence, $\lambda = \mu$ if and only if $\gamma = 0$. The result now follows from Proposition 11.3.2. $\square$

Example 11.3.4. Let $A \triangleq \begin{bmatrix} \nu & \omega \\ -\omega & \nu \end{bmatrix} \in \mathbb{R}^{2 \times 2}$. Then,

$$e^{tA} = e^{\nu t} \begin{bmatrix} \cos \omega t & \sin \omega t \\ -\sin \omega t & \cos \omega t \end{bmatrix}. \quad (11.3.4)$$

On the other hand, if $A \triangleq \begin{bmatrix} \nu & \omega \\ \omega & -\nu \end{bmatrix}$, then

$$e^{tA} = \begin{bmatrix} \cosh \delta t + \frac{\nu}{\delta} \sinh \delta t & \frac{\omega}{\delta} \sinh \delta t \\ \frac{\omega}{\delta} \sinh \delta t & \cosh \delta t - \frac{\nu}{\delta} \sinh \delta t \end{bmatrix}, \quad (11.3.5)$$

where $\delta \triangleq \sqrt{\omega^2 + \nu^2}$.

Example 11.3.5. Let $\alpha \in \mathbb{F}$, and define $A \triangleq \begin{bmatrix} 0 & 1 \\ 0 & \alpha \end{bmatrix}$. Then,

$$e^{tA} = \begin{cases} \begin{bmatrix} 1 & \alpha^{-1}(e^{\alpha t} - 1) \\ 0 & e^{\alpha t} \end{bmatrix}, & \alpha \neq 0, \\ \begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix}, & \alpha = 0. \end{cases}$$

Example 11.3.6. Let $\theta \in \mathbb{R}$, and define $A \triangleq \begin{bmatrix} 0 & \theta \\ -\theta & 0 \end{bmatrix}$. Then,

$$e^A = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}.$$

Furthermore, define $B \triangleq \begin{bmatrix} 0 & \frac{\pi}{2} - \theta \\ -\frac{\pi}{2} + \theta & 0 \end{bmatrix}$. Then,

$$e^B = \begin{bmatrix} \sin \theta & \cos \theta \\ -\cos \theta & \sin \theta \end{bmatrix}.$$

Example 11.3.7. Consider the second-order mechanical vibration equation

$$m\ddot{q} + c\dot{q} + kq = 0, \quad (11.3.6)$$

where m is positive and c and k are nonnegative. Here m , c , and k denote mass, damping, and stiffness parameters, respectively. Equation (11.3.6) can be written in companion form as the system

$$\dot{x} = Ax, \quad (11.3.7)$$

where

$$x \triangleq \begin{bmatrix} q \\ \dot{q} \end{bmatrix}, \quad A \triangleq \begin{bmatrix} 0 & 1 \\ -k/m & -c/m \end{bmatrix}. \quad (11.3.8)$$

The inelastic case $k = 0$ is the simplest one since A is upper triangular. In this case,

$$e^{tA} = \begin{cases} \begin{bmatrix} 1 & t \\ 0 & 1 \end{bmatrix}, & k = c = 0, \\ \begin{bmatrix} 1 & \frac{m}{c}(1 - e^{-ct/m}) \\ 0 & e^{-ct/m} \end{bmatrix}, & k = 0, \ c > 0, \end{cases} \quad (11.3.9)$$

where $c = 0$ and $c > 0$ correspond to a rigid body and a damped rigid body, respectively.

Next, we consider the elastic case $c \geq 0$ and $k > 0$. In this case, we define

$$\omega_n \triangleq \sqrt{\frac{k}{m}}, \quad \zeta \triangleq \frac{c}{2\sqrt{mk}}, \quad (11.3.10)$$

where $\omega_n > 0$ denotes the (undamped) *natural frequency* of vibration and $\zeta \geq 0$ denotes the *damping ratio*. Now, A can be written as

$$A = \begin{bmatrix} 0 & 1 \\ -\omega_n^2 & -2\zeta\omega_n \end{bmatrix}, \quad (11.3.11)$$

and Corollary 11.3.3 yields

$$e^{tA} = \begin{cases} \begin{bmatrix} \cos \omega_n t & \frac{1}{\omega_n} \sin \omega_n t \\ -\omega_n \sin \omega_n t & \cos \omega_n t \end{bmatrix}, & \zeta = 0, \\ e^{-\zeta\omega_n t} \begin{bmatrix} \cos \omega_d t + \frac{\zeta}{\sqrt{1-\zeta^2}} \sin \omega_d t & \frac{1}{\omega_d} \sin \omega_d t \\ \frac{-\omega_d}{1-\zeta^2} \sin \omega_d t & \cos \omega_d t - \frac{\zeta}{\sqrt{1-\zeta^2}} \sin \omega_d t \end{bmatrix}, & 0 < \zeta < 1, \\ e^{-\omega_n t} \begin{bmatrix} 1 + \omega_n t & t \\ -\omega_n^2 t & 1 - \omega_n t \end{bmatrix}, & \zeta = 1, \\ e^{-\zeta\omega_n t} \begin{bmatrix} \cosh \omega_d t + \frac{\zeta}{\sqrt{\zeta^2-1}} \sinh \omega_d t & \frac{1}{\omega_d} \sinh \omega_d t \\ \frac{-\omega_d}{\zeta^2-1} \sinh \omega_d t & \cosh \omega_d t - \frac{\zeta}{\sqrt{\zeta^2-1}} \sinh \omega_d t \end{bmatrix}, & \zeta > 1, \end{cases} \quad (11.3.12)$$

where $\zeta = 0$, $0 < \zeta < 1$, $\zeta = 1$, and $\zeta > 1$ correspond to *undamped*, *underdamped*, *critically damped*, and *overdamped oscillators*, respectively, and where the *damped natural frequency* ω_d is the positive number

$$\omega_d \triangleq \begin{cases} \omega_n \sqrt{1 - \zeta^2}, & 0 < \zeta < 1, \\ \omega_n \sqrt{\zeta^2 - 1}, & \zeta > 1. \end{cases} \quad (11.3.13)$$

Note that m and k are not integers here.

11.4 Matrix Logarithms

Definition 11.4.1. Let $A \in \mathbb{F}^{n \times n}$. Then, $B \in \mathbb{F}^{n \times n}$ is a *logarithm* of A if $e^B = A$.

The following result shows that every complex, nonsingular matrix has a complex logarithm.

Proposition 11.4.2. Let $A \in \mathbb{C}^{n \times n}$. Then, there exists a matrix $B \in \mathbb{C}^{n \times n}$ such that $A = e^B$ if and only if A is nonsingular.

Proof. See [624, pp. 35, 60] or [711, p. 474]. □

Although the real number -1 does not have a real logarithm, the real matrix $B = \begin{bmatrix} 0 & \pi \\ -\pi & 0 \end{bmatrix}$ satisfies $e^B = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}$. These examples suggest that only certain real matrices have a real logarithm.

Proposition 11.4.3. Let $A \in \mathbb{R}^{n \times n}$. Then, there exists a matrix $B \in \mathbb{R}^{n \times n}$ such that $A = e^B$ if and only if A is nonsingular and, for every negative eigenvalue λ of A and for every positive integer k , the Jordan form of A has an even number of $k \times k$ blocks associated with λ .

Proof. See [711, p. 475]. □

Replacing A and B in Proposition 11.4.3 by e^A and A , respectively, yields the following result.

Corollary 11.4.4. Let $A \in \mathbb{R}^{n \times n}$. Then, for every negative eigenvalue λ of e^A and for every positive integer k , the Jordan form of e^A has an even number of $k \times k$ blocks associated with λ .

Since the matrix $A \triangleq \begin{bmatrix} -2\pi & 4\pi \\ -2\pi & 2\pi \end{bmatrix}$ satisfies $e^A = I$, it follows that a positive-definite matrix can have a logarithm that is not normal. However, the following result shows that every positive-definite matrix has exactly one Hermitian logarithm.

Proposition 11.4.5. The function $\exp: \mathbf{H}^n \mapsto \mathbf{P}^n$ is one-to-one and onto.

Let $A \in \mathbb{R}^{n \times n}$. If there exists a matrix $B \in \mathbb{R}^{n \times n}$ such that $A = e^B$, then Corollary 11.2.4 implies that $\det A = \det e^B = e^{\text{tr } B} > 0$. However, the converse is not true. Consider, for example, $A \triangleq \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix}$, which satisfies $\det A > 0$. However, Proposition 11.4.3 implies that there does not exist a matrix $B \in \mathbb{R}^{2 \times 2}$ such that $A = e^B$. On the other hand, note that $A = e^B e^C$, where $B \triangleq \begin{bmatrix} 0 & \pi \\ -\pi & 0 \end{bmatrix}$ and $C \triangleq \begin{bmatrix} 0 & 0 \\ 0 & \log 2 \end{bmatrix}$. While the product of two exponentials of real matrices has positive determinant, the following result shows that the converse is also true.

Proposition 11.4.6. Let $A \in \mathbb{R}^{n \times n}$. Then, there exist matrices $B, C \in \mathbb{R}^{n \times n}$ such that $A = e^B e^C$ if and only if $\det A > 0$.

Proof. Suppose that there exist $B, C \in \mathbb{R}^{n \times n}$ such that $A = e^B e^C$. Then, $\det A = (\det e^B)(\det e^C) > 0$. Conversely, suppose that $\det A > 0$. If A has no negative eigenvalues, then it follows from Proposition 11.4.3 that there exists $B \in \mathbb{R}^{n \times n}$ such that $A = e^B$. Hence, $A = e^B e^{0_{n \times n}}$. Now, suppose that A has at least one negative eigenvalue. Then, Theorem 5.3.5 on the real Jordan form implies that there exist a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ and matrices $A_1 \in \mathbb{R}^{n_1 \times n_1}$ and $A_2 \in \mathbb{R}^{n_2 \times n_2}$ such that $A = S \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} S^{-1}$, where every eigenvalue of A_1 is negative and where none of the eigenvalues of A_2 are negative. Since $\det A$ and $\det A_2$ are positive, it follows that n_1 is even. Now, write $A = S \begin{bmatrix} -I_{n_1} & 0 \\ 0 & I_{n_2} \end{bmatrix} \begin{bmatrix} -A_1 & 0 \\ 0 & A_2 \end{bmatrix} S^{-1}$. Since the eigenvalue -1 of $\begin{bmatrix} -I_{n_1} & 0 \\ 0 & I_{n_2} \end{bmatrix}$ appears in an even number of 1×1 Jordan blocks, it follows from Proposition 11.4.3 that there exists a matrix $\hat{B} \in \mathbb{R}^{n \times n}$ such that $\begin{bmatrix} -I_{n_1} & 0 \\ 0 & I_{n_2} \end{bmatrix} = e^{\hat{B}}$. Furthermore, since $\begin{bmatrix} -A_1 & 0 \\ 0 & A_2 \end{bmatrix}$ has no negative eigenvalues, it follows that there exists a matrix $\hat{C} \in \mathbb{R}^{n \times n}$ such that $\begin{bmatrix} -A_1 & 0 \\ 0 & A_2 \end{bmatrix} = e^{\hat{C}}$. Hence, $e^A = S e^{\hat{B}} e^{\hat{C}} S^{-1} = e^{S \hat{B} S^{-1}} e^{S \hat{C} S^{-1}}$. $\square$

Although $e^A e^B$ may be different from e^{A+B} , the following result, known as the *Baker-Campbell-Hausdorff series*, provides an expansion for a matrix function $C(t)$ that satisfies $e^{C(t)} = e^{tA} e^{tB}$.

Proposition 11.4.7. Let $A_1, \dots, A_l \in \mathbb{F}^{n \times n}$. Then, there exists $\varepsilon > 0$ such that, for all $t \in (-\varepsilon, \varepsilon)$,

$$e^{tA_1} \dots e^{tA_l} = e^{C(t)}, \quad (11.4.1)$$

where

$$C(t) \triangleq \sum_{i=1}^l t A_i + \sum_{1 \leq i < j \leq l} \frac{1}{2} t^2 [A_i, A_j] + O(t^3). \quad (11.4.2)$$

Proof. See [624, Chapter 3], [1162, p. 35], or [1366, p. 97]. $\square$

To illustrate (11.4.1), let $l = 2$, $A = A_1$, and $B = A_2$. Then, the first few terms of the series are given by

$$e^{tA} e^{tB} = e^{tA+tB+(t^2/2)[A,B]+(t^3/12)[[B,A],A+B]+\dots}. \quad (11.4.3)$$

The radius of convergence of this series is discussed in [379, 1037].

The following result is the *Lie-Trotter product formula*.

Corollary 11.4.8. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$e^{A+B} = \lim_{p \rightarrow \infty} \left[e^{\frac{1}{p}A} e^{\frac{1}{p}B} \right]^p. \quad (11.4.4)$$

Proof. Setting $l = 2$ and $t = 1/p$ in (11.4.1) yields, as $p \rightarrow \infty$,

$$\left[e^{\frac{1}{p}A} e^{\frac{1}{p}B} \right]^p = \left[e^{\frac{1}{p}(A+B)+O(1/p^2)} \right]^p = e^{A+B+O(1/p)} \rightarrow e^{A+B}. \quad \square$$

11.5 The Logarithm Function

Let $A \in \mathbb{F}^{n \times n}$ be positive definite so that $A = SBS^* \in \mathbb{F}^{n \times n}$, where $S \in \mathbb{F}^{n \times n}$ is unitary and $B \in \mathbb{R}^{n \times n}$ is diagonal with positive diagonal entries. In Section 8.5, $\log A$ is defined as $\log A = S(\log B)S^* \in \mathbf{H}^n$, where $(\log B)_{(i,i)} \triangleq \log B_{(i,i)}$. Since $\log A$ satisfies $A = e^{\log A}$, it follows that $\log A$ is a logarithm of A . The following result extends the definition of $\log A$ to arbitrary nonsingular matrices $A \in \mathbb{C}^{n \times n}$.

Theorem 11.5.1. Let $A \in \mathbb{C}^{n \times n}$. Then, the following statements hold:

i) If A is nonsingular, then the principal branch of the log function

$$\log: \mathbb{C} \setminus \{0\} \mapsto \{z: \operatorname{Re} z \neq 0 \text{ and } -\pi < \operatorname{Im} z \leq \pi\}$$

is defined at A .

ii) If A is nonsingular, then $\log A$ is a logarithm of A , that is, $e^{\log A} = A$.

iii) $\log e^A = A$ if and only if, for all $\lambda \in \operatorname{spec}(A)$, it follows that $|\operatorname{Im} \lambda| < \pi$.

iv) If A is nonsingular and $\operatorname{sprad}(A - I) \leq 1$, then $\log A$ is given by the series

$$\log A = \sum_{i=1}^{\infty} \frac{(-1)^{i+1}}{i} (A - I)^i, \quad (11.5.1)$$

which converges absolutely with respect to every submultiplicative norm $\|\cdot\|$ such that $\|A - I\| < 1$.

v) If $\operatorname{spec}(A) \subset \operatorname{ORHP}$, then $\log A$ is given by the series

$$\log A = \sum_{i=0}^{\infty} \frac{2}{2i+1} [(A - I)(A + I)^{-1}]^{2i+1}.$$

vi) If A has no eigenvalues in $(-\infty, 0]$, then

$$\log A = \int_0^1 (A - I)[t(A - I) + I]^{-1} dt.$$

vii) If A has no eigenvalues in $(-\infty, 0]$ and $\alpha \in [-1, 1]$, then

$$\log A^\alpha = \alpha \log A.$$

In particular,

$$\log A^{-1} = -\log A$$

and

$$\log A^{1/2} = \frac{1}{2}\log A.$$

viii) If A is real and $\text{spec}(A) \subset \text{ORHP}$, then $\log A$ is real.

ix) If A is real and nonsingular, then $\log A$ is real if and only if A is nonsingular and, for every negative eigenvalue λ of A and for every positive integer k , the Jordan form of A has an even number of $k \times k$ blocks associated with λ .

Now, let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{C}^{n \times n}$. Then, the following statements hold:

x) The function $\log$ is continuous on $\{X \in \mathbb{C}^{n \times n}: \|X - I\| < 1\}$.

xi) If $B \in \mathbb{C}^{n \times n}$ and $\|B\| < \log 2$, then $\|e^B - I\| < 1$ and $\log e^B = B$.

xii) $\exp: \mathbb{B}_{\log 2}(0) \mapsto \mathbb{F}^{n \times n}$ is one-to-one.

xiii) If $\|A - I\| < 1$, then

$$\|\log A\| \leq -\log(1 - \|A - I\|) \leq \frac{\|A - I\|}{1 - \|A - I\|}.$$

xiv) If $\|A - I\| < 2/3$, then

$$\|A - I\| \left[1 - \frac{\|A - I\|}{2(1 - \|A - I\|)} \right] \leq \|\log A\|.$$

xv) Assume that A is nonsingular, and let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$. Then,

$$\text{mspec}(\log A) = \{\log \lambda_1, \dots, \log \lambda_n\}_{\text{ms}}.$$

Proof. Statement *i)* follows from Definition 10.5.1 as well as the properties of the principal branch of the log function given by Fact 1.18.7. Statement *ii)* follows from the discussion in [711, p. 420].

Statement *iii)* is given in [683, p. 32].

Statements *iv)* and *v)* are given by Fact 10.11.24. See [624, pp. 34–35] and [683, p. 273].

Statement *vi)* is given in [683, p. 269].

Statement *vii)* is given in [683, p. 270].

Statement *ix)* follows from Proposition 11.4.3 and the discussion in [711, pp. 474–475].

Statements *x)* and *xi)* are proved in [624, pp. 34–35]. To prove the inequality in *xi)*, let $\|B\| < 2$, so that $e^{\|B\|} < 2$, and thus

$$\|e^B - I\| \leq \sum_{i=1}^{\infty} (i!)^{-1} \|B\|^i = e^{\|B\|} - 1 < 1.$$

To prove *xii*), let $B_1, B_2 \in \mathbb{B}_{\log 2}(0)$, and assume that $e^{B_1} = e^{B_2}$. Then, it follows from *ii*) that $B_1 = \log e^{B_1} = \log e^{B_2} = B_2$.

Finally, to prove *xiii*), let $\alpha \triangleq \|A - I\| < 1$. Then, it follows from (11.5.1) and *iv*) of Fact 1.18.7 that $\|\log A\| \leq \sum_{i=1}^{\infty} \alpha^i / i = -\log(1 - \alpha)$. For *xiv*), see [683, p. 647]. $\square$

For a nonsingular $A \in \mathbb{C}^{n \times n}$, the matrix $\log A$ given by Theorem 11.5.1 is the *principal logarithm*.

11.6 Lie Groups

Definition 11.6.1. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$, and assume that $\mathcal{S}$ is a group. Then, $\mathcal{S}$ is a *Lie group* if $\mathcal{S}$ is closed relative to $\text{GL}_{\mathbb{F}}(n)$.

Proposition 11.6.2. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$, and assume that $\mathcal{S}$ is a group. Then, $\mathcal{S}$ is a Lie group if and only if the limit of every convergent sequence in $\mathcal{S}$ is either an element of $\mathcal{S}$ or is singular.

The groups $\text{SL}_{\mathbb{F}}(n)$, $\text{U}(n)$, $\text{O}(n)$, $\text{SU}(n)$, $\text{SO}(n)$, $\text{U}(n, m)$, $\text{O}(n, m)$, $\text{SU}(n, m)$, $\text{SO}(n, m)$, $\text{Sp}_{\mathbb{F}}(n)$, $\text{Aff}_{\mathbb{F}}(n)$, $\text{SE}_{\mathbb{F}}(n)$, and $\text{Trans}_{\mathbb{F}}(n)$ defined in Proposition 3.3.6 are closed sets, and thus are Lie groups. Although the groups $\text{GL}_{\mathbb{F}}(n)$, $\text{PL}_{\mathbb{F}}(n)$, and $\text{UT}(n)$ (see Fact 3.21.5) are not closed sets, they are closed relative to $\text{GL}_{\mathbb{F}}(n)$, and thus they are Lie groups. Finally, the group $\mathcal{S} \subset \mathbb{C}^{2 \times 2}$ defined by

$$\mathcal{S} \triangleq \left\{ \begin{bmatrix} e^{jt} & 0 \\ 0 & e^{j\pi t} \end{bmatrix} : t \in \mathbb{R} \right\} \quad (11.6.1)$$

is not closed relative to $\text{GL}_{\mathbb{C}}(2)$, and thus is not a Lie group. For details, see [624, p. 4].

Proposition 11.6.3. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$, and assume that $\mathcal{S}$ is a Lie group. Furthermore, define

$$\mathcal{S}_0 \triangleq \{A \in \mathbb{F}^{n \times n} : e^{tA} \in \mathcal{S} \text{ for all } t \in \mathbb{R}\}. \quad (11.6.2)$$

Then, $\mathcal{S}_0$ is a Lie algebra.

Proof. See [624, pp. 39, 43, 44]. $\square$

The Lie algebra $\mathcal{S}_0$ defined by (11.6.2) is the *Lie algebra of $\mathcal{S}$* .

Proposition 11.6.4. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$, assume that $\mathcal{S}$ is a Lie group, and let $\mathcal{S}_0 \subseteq \mathbb{F}^{n \times n}$ be the Lie algebra of $\mathcal{S}$. Furthermore, let $S \in \mathcal{S}$ and $A \in \mathcal{S}_0$. Then, $SAS^{-1} \in \mathcal{S}_0$.

Proof. For all $t \in \mathbb{R}$, $e^{tA} \in \mathcal{S}$, and thus $e^{tSAS^{-1}} = Se^{tA}S^{-1} \in \mathcal{S}$. Hence, $SAS^{-1} \in \mathcal{S}_0$. $\square$

Proposition 11.6.5. The following statements hold:

- i) $\mathfrak{gl}_{\mathbb{F}}(n)$ is the Lie algebra of $\mathrm{GL}_{\mathbb{F}}(n)$.
- ii) $\mathfrak{gl}_{\mathbb{R}}(n) = \mathfrak{pl}_{\mathbb{R}}(n)$ is the Lie algebra of $\mathrm{PL}_{\mathbb{R}}(n)$.
- iii) $\mathfrak{pl}_{\mathbb{C}}(n)$ is the Lie algebra of $\mathrm{PL}_{\mathbb{C}}(n)$.
- iv) $\mathfrak{sl}_{\mathbb{F}}(n)$ is the Lie algebra of $\mathrm{SL}_{\mathbb{F}}(n)$.
- v) $\mathfrak{u}(n)$ is the Lie algebra of $\mathrm{U}(n)$.
- vi) $\mathfrak{so}(n)$ is the Lie algebra of $\mathrm{O}(n)$.
- vii) $\mathfrak{su}(n)$ is the Lie algebra of $\mathrm{SU}(n)$.
- viii) $\mathfrak{so}(n)$ is the Lie algebra of $\mathrm{SO}(n)$.
- ix) $\mathfrak{su}(n, m)$ is the Lie algebra of $\mathrm{U}(n, m)$.
- x) $\mathfrak{so}(n, m)$ is the Lie algebra of $\mathrm{O}(n, m)$.
- xi) $\mathfrak{su}(n, m)$ is the Lie algebra of $\mathrm{SU}(n, m)$.
- xii) $\mathfrak{so}(n, m)$ is the Lie algebra of $\mathrm{SO}(n, m)$.
- xiii) $\mathfrak{symp}_{\mathbb{F}}(2n)$ is the Lie algebra of $\mathrm{Symp}_{\mathbb{F}}(2n)$.
- xiv) $\mathfrak{osymp}_{\mathbb{F}}(2n)$ is the Lie algebra of $\mathrm{OSymp}_{\mathbb{F}}(2n)$.
- xv) $\mathfrak{aff}_{\mathbb{F}}(n)$ is the Lie algebra of $\mathrm{Aff}_{\mathbb{F}}(n)$.
- xvi) $\mathfrak{se}_{\mathbb{C}}(n)$ is the Lie algebra of $\mathrm{SE}_{\mathbb{C}}(n)$.
- xvii) $\mathfrak{se}_{\mathbb{R}}(n)$ is the Lie algebra of $\mathrm{SE}_{\mathbb{R}}(n)$.
- xviii) $\mathfrak{trans}_{\mathbb{F}}(n)$ is the Lie algebra of $\mathrm{Trans}_{\mathbb{F}}(n)$.

Proof. See [624, pp. 38–41]. $\square$

Proposition 11.6.6. Let $\mathcal{S} \subset \mathbb{F}^{n \times n}$, assume that $\mathcal{S}$ is a Lie group, and let $\mathcal{S}_0 \subseteq \mathbb{F}^{n \times n}$ be the Lie algebra of $\mathcal{S}$. Then, $\exp: \mathcal{S}_0 \mapsto \mathcal{S}$. Furthermore, if $\exp$ is onto, then $\mathcal{S}$ is pathwise connected.

Proof. Let $A \in \mathcal{S}_0$ so that $e^{tA} \in \mathcal{S}$ for all $t \in \mathbb{R}$. Hence, setting $t = 1$ implies that $\exp: \mathcal{S}_0 \mapsto \mathcal{S}$. Now, suppose that $\exp$ is onto, let $B \in \mathcal{S}$, and let $A \in \mathcal{S}_0$ be such that $e^A = B$. Then, $f(t) \triangleq e^{tA}$ satisfies $f(0) = I$ and $f(1) = B$, which implies that $\mathcal{S}$ is pathwise connected. $\square$

A Lie group can consist of multiple pathwise-connected components.

Proposition 11.6.7. Let $n \geq 1$. Then, the following functions are onto:

- i) $\exp: \mathfrak{gl}_{\mathbb{C}}(n) \mapsto \mathrm{GL}_{\mathbb{C}}(n)$.
- ii) $\exp: \mathfrak{gl}_{\mathbb{R}}(1) \mapsto \mathrm{PL}_{\mathbb{R}}(1)$.

- iii) $\exp: \mathfrak{pl}_{\mathbb{C}}(n) \mapsto \mathrm{PL}_{\mathbb{C}}(n)$.
- iv) $\exp: \mathfrak{sl}_{\mathbb{C}}(n) \mapsto \mathrm{SL}_{\mathbb{C}}(n)$.
- v) $\exp: \mathfrak{u}(n) \mapsto \mathrm{U}(n)$.
- vi) $\exp: \mathfrak{su}(n) \mapsto \mathrm{SU}(n)$.
- vii) $\exp: \mathfrak{so}(n) \mapsto \mathrm{SO}(n)$.

Furthermore, the following functions are not onto:

- viii) $\exp: \mathfrak{gl}_{\mathbb{R}}(n) \mapsto \mathrm{PL}_{\mathbb{R}}(n)$, where $n \geq 2$.
- ix) $\exp: \mathfrak{sl}_{\mathbb{R}}(n) \mapsto \mathrm{SL}_{\mathbb{R}}(n)$.
- x) $\exp: \mathfrak{so}(n) \mapsto \mathrm{O}(n)$.
- xi) $\exp: \mathrm{symp}_{\mathbb{R}}(2n) \mapsto \mathrm{Symp}_{\mathbb{R}}(2n)$.

Proof. Statement *i)* follows from Proposition 11.4.2, while *ii)* is immediate. Statements *iii)–vii)* can be verified by construction; see [1098, pp. 199, 212] for the proof of *v)* and *vii)*. The example $A \triangleq \begin{bmatrix} -1 & 0 \\ 0 & -2 \end{bmatrix}$ and Proposition 11.4.3 show that *viii)* is not onto. For $\lambda < 0$, $\lambda \neq -1$, Proposition 11.4.3 and the example $\begin{bmatrix} \lambda & 0 \\ 0 & 1/\lambda \end{bmatrix}$ given in [1162, p. 39] show that *ix)* is not onto. See also [103, pp. 84, 85]. Statement *viii)* shows that *x)* is not onto. For *xi)*, see [404]. $\square$

Proposition 11.6.8. The Lie groups $\mathrm{GL}_{\mathbb{C}}(n)$, $\mathrm{SL}_{\mathbb{F}}(n)$, $\mathrm{U}(n)$, $\mathrm{SU}(n)$, and $\mathrm{SO}(n)$ are pathwise connected. The Lie groups $\mathrm{GL}_{\mathbb{R}}(n)$, $\mathrm{O}(n)$, $\mathrm{O}(n, 1)$, and $\mathrm{SO}(n, 1)$ are not pathwise connected.

Proof. See [624, p. 15]. $\square$

Proposition 11.6.8 and *ix)* of Proposition 11.6.7 show that the converse of Proposition 11.6.6 does not hold, that is, pathwise connectedness does not imply that $\exp$ is onto. See [1162, p. 39].

11.7 Lyapunov Stability Theory

Consider the dynamical system

$$\dot{x}(t) = f[x(t)], \quad (11.7.1)$$

where $t \geq 0$, $x(t) \in \mathcal{D} \subseteq \mathbb{R}^n$, and $f: \mathcal{D} \rightarrow \mathbb{R}^n$ is continuous. We assume that, for all $x_0 \in \mathcal{D}$ and for all $T > 0$, there exists a unique C^1 solution $x: [0, T] \mapsto \mathcal{D}$ satisfying (11.7.1). If $x_e \in \mathcal{D}$ satisfies $f(x_e) = 0$, then $x(t) \equiv x_e$ is an *equilibrium* of (11.7.1). The following definition concerns the stability of an equilibrium of (11.7.1). Throughout this section, $\|\cdot\|$ denotes a norm on $\mathbb{R}^n$.

Definition 11.7.1. Let $x_e \in \mathcal{D}$ be an equilibrium of (11.7.1). Then, x_e is *Lyapunov stable* if, for all $\varepsilon > 0$, there exists $\delta > 0$ such that, if $\|x(0) - x_e\| < \delta$, then $\|x(t) - x_e\| < \varepsilon$ for all $t \geq 0$. Furthermore, x_e is *asymptotically stable* if it is Lyapunov stable and there exists $\varepsilon > 0$ such that, if $\|x(0) - x_e\| < \varepsilon$, then

$\lim_{t \rightarrow \infty} x(t) = x_e$. In addition, x_e is *globally asymptotically stable* if it is Lyapunov stable, $\mathcal{D} = \mathbb{R}^n$, and, for all $x(0) \in \mathbb{R}^n$, $\lim_{t \rightarrow \infty} x(t) = x_e$. Finally, x_e is *unstable* if it is not Lyapunov stable.

Note that, if $x_e \in \mathbb{R}^n$ is a globally asymptotically stable equilibrium, then x_e is the only equilibrium of (11.7.1).

The following result, known as *Lyapunov's direct method*, gives sufficient conditions for Lyapunov stability and asymptotic stability of an equilibrium of (11.7.1).

Theorem 11.7.2. Let $x_e \in \mathcal{D}$ be an equilibrium of the dynamical system (11.7.1), and assume there exists a C^1 function $V: \mathcal{D} \mapsto \mathbb{R}$ such that

$$V(x_e) = 0, \quad (11.7.2)$$

such that, for all $x \in \mathcal{D} \setminus \{x_e\}$,

$$V(x) > 0, \quad (11.7.3)$$

and such that, for all $x \in \mathcal{D}$,

$$V'(x)f(x) \leq 0. \quad (11.7.4)$$

Then, x_e is Lyapunov stable. If, in addition, for all $x \in \mathcal{D} \setminus \{x_e\}$,

$$V'(x)f(x) < 0, \quad (11.7.5)$$

then x_e is asymptotically stable. Finally, if $\mathcal{D} = \mathbb{R}^n$ and

$$\lim_{\|x\| \rightarrow \infty} V(x) = \infty, \quad (11.7.6)$$

then x_e is globally asymptotically stable.

Proof. For convenience, let $x_e = 0$. To prove Lyapunov stability, let $\varepsilon > 0$ be such that $\mathbb{B}_\varepsilon(0) \subseteq \mathcal{D}$. Since $\mathbb{S}_\varepsilon(0)$ is compact and $V(x)$ is continuous, it follows from Theorem 10.3.8 that $V[\mathbb{S}_\varepsilon(0)]$ is compact. Since $0 \notin \mathbb{S}_\varepsilon(0)$, $V(x) > 0$ for all $x \in \mathcal{D} \setminus \{0\}$, and $V[\mathbb{S}_\varepsilon(0)]$ is compact, it follows that $\alpha \triangleq \min V[\mathbb{S}_\varepsilon(0)]$ is positive. Next, since V is continuous, it follows that there exists $\delta \in (0, \varepsilon]$ such that $V(x) < \alpha$ for all $x \in \mathbb{B}_\delta(0)$. Now, let $x(t)$ for all $t \geq 0$ satisfy (11.7.1), where $\|x(0)\| < \delta$. Hence, $V[x(0)] < \alpha$. It thus follows from (11.7.4) that, for all $t \geq 0$,

$$V[x(t)] - V[x(0)] = \int_0^t V'[x(s)]f[x(s)] \, ds \leq 0,$$

and hence, for all $t \geq 0$,

$$V[x(t)] \leq V[x(0)] < \alpha.$$

Now, since $V(x) \geq \alpha$ for all $x \in \mathbb{S}_\varepsilon(0)$, it follows that $x(t) \notin \mathbb{S}_\varepsilon(0)$ for all $t \geq 0$. Hence, $\|x(t)\| < \varepsilon$ for all $t \geq 0$, which proves that $x_e = 0$ is Lyapunov stable.

To prove that $x_e = 0$ is asymptotically stable, let $\varepsilon > 0$ be such that $\mathbb{B}_\varepsilon(0) \subseteq \mathcal{D}$. Since (11.7.5) implies (11.7.4), it follows that there exists $\delta > 0$ such that, if $\|x(0)\| < \delta$, then $\|x(t)\| < \varepsilon$ for all $t \geq 0$. Furthermore, $\frac{d}{dt}V[x(t)] = V'[x(t)]f[x(t)] < 0$ for all $t \geq 0$, and thus $V[x(t)]$ is decreasing and bounded from below by zero. Now, suppose that $V[x(t)]$ does not converge to zero. Therefore, there exists $L > 0$

such that $V[x(t)] \geq L$ for all $t \geq 0$. Now, let $\delta' > 0$ be such that $V(x) < L$ for all $x \in \mathbb{B}_{\delta'}(0)$. Therefore, $\|x(t)\| \geq \delta'$ for all $t \geq 0$. Next, define $\gamma < 0$ by $\gamma \triangleq \max_{\delta' \leq \|x\| \leq \varepsilon} V'(x)f(x)$. Therefore, since $\|x(t)\| < \varepsilon$ for all $t \geq 0$, it follows that

$$V[x(t)] - V[x(0)] = \int_0^t V'[x(\tau)]f[x(\tau)] \, d\tau \leq \gamma t,$$

and hence

$$V(x(t)) \leq V[x(0)] + \gamma t.$$

However, $t > -V[x(0)]/\gamma$ implies that $V[x(t)] < 0$, which is a contradiction.

To prove that $x_e = 0$ is globally asymptotically stable, let $x(0) \in \mathbb{R}^n$, and let $\beta \triangleq V[x(0)]$. It follows from (11.7.6) that there exists $\varepsilon > 0$ such that $V(x) > \beta$ for all $x \in \mathbb{R}^n$ such that $\|x\| > \varepsilon$. Therefore, $\|x(0)\| \leq \varepsilon$, and, since $V[x(t)]$ is decreasing, it follows that $\|x(t)\| < \varepsilon$ for all $t > 0$. The remainder of the proof is identical to the proof of asymptotic stability. $\square$

11.8 Linear Stability Theory

We now specialize Definition 11.7.1 to the linear system

$$\dot{x}(t) = Ax(t), \tag{11.8.1}$$

where $t \geq 0$, $x(t) \in \mathbb{R}^n$, and $A \in \mathbb{R}^{n \times n}$. Note that $x_e = 0$ is an equilibrium of (11.8.1), and that $x_e \in \mathbb{R}^n$ is an equilibrium of (11.8.1) if and only if $x_e \in \mathcal{N}(A)$. Hence, if x_e is the globally asymptotically stable equilibrium of (11.8.1), then A is nonsingular and $x_e = 0$.

We consider three types of stability for the linear system (11.8.1). Unlike Definition 11.7.1, these definitions are stated in terms of the dynamics matrix rather than the equilibrium.

Definition 11.8.1. For $A \in \mathbb{F}^{n \times n}$, define the following classes of matrices:

- i) A is *Lyapunov stable* if $\text{spec}(A) \subset \text{CLHP}$ and, if $\lambda \in \text{spec}(A)$ and $\text{Re } \lambda = 0$, then λ is semisimple.
- ii) A is *semistable* if $\text{spec}(A) \subset \text{OLHP} \cup \{0\}$ and, if $0 \in \text{spec}(A)$, then 0 is semisimple.
- iii) A is *asymptotically stable* if $\text{spec}(A) \subset \text{OLHP}$.

The following result concerns Lyapunov stability, semistability, and asymptotic stability for (11.8.1).

Proposition 11.8.2. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

- i) $x_e = 0$ is a Lyapunov-stable equilibrium of (11.8.1).
- ii) At least one equilibrium of (11.8.1) is Lyapunov stable.

- iii) Every equilibrium of (11.8.1) is Lyapunov stable.
- iv) A is Lyapunov stable.
- v) For every initial condition $x(0) \in \mathbb{R}^n$, $x(t)$ is bounded for all $t \geq 0$.
- vi) $\|e^{tA}\|$ is bounded for all $t \geq 0$, where $\|\cdot\|$ is a norm on $\mathbb{R}^{n \times n}$.
- vii) For every initial condition $x(0) \in \mathbb{R}^n$, $e^{tA}x(0)$ is bounded for all $t \geq 0$.

The following statements are equivalent:

- viii) A is semistable.
- ix) $\lim_{t \rightarrow \infty} e^{tA}$ exists.
- x) For every initial condition $x(0)$, $\lim_{t \rightarrow \infty} x(t)$ exists.

In this case,

$$\lim_{t \rightarrow \infty} e^{tA} = I - AA^\# \quad (11.8.2)$$

The following statements are equivalent:

- xi) $x_e = 0$ is an asymptotically stable equilibrium of (11.8.1).
- xii) A is asymptotically stable.
- xiii) $\text{spabs}(A) < 0$.
- xiv) For every initial condition $x(0) \in \mathbb{R}^n$, $\lim_{t \rightarrow \infty} x(t) = 0$.
- xv) For every initial condition $x(0) \in \mathbb{R}^n$, $e^{tA}x(0) \rightarrow 0$ as $t \rightarrow \infty$.
- xvi) $e^{tA} \rightarrow 0$ as $t \rightarrow \infty$.

The following definition concerns the stability of a polynomial.

Definition 11.8.3. Let $p \in \mathbb{R}[s]$. Then, define the following terminology:

- i) p is *Lyapunov stable* if $\text{roots}(p) \subset \text{CLHP}$ and, if λ is an imaginary root of p , then $m_p(\lambda) = 1$.
- ii) p is *semistable* if $\text{roots}(p) \subset \text{OLHP} \cup \{0\}$ and, if $0 \in \text{roots}(p)$, then $m_p(0) = 1$.
- iii) p is *asymptotically stable* if $\text{roots}(p) \subset \text{OLHP}$.

For the following result, recall Definition 11.8.1.

Proposition 11.8.4. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) A is Lyapunov stable if and only if μ_A is Lyapunov stable.
- ii) A is semistable if and only if μ_A is semistable.

Furthermore, the following statements are equivalent:

- iii) A is asymptotically stable
- iv) μ_A is asymptotically stable.

v) χ_A is asymptotically stable.

Next, consider the factorization of the minimal polynomial μ_A of A given by

$$\mu_A = \mu_A^s \mu_A^u, \quad (11.8.3)$$

where μ_A^s and μ_A^u are monic polynomials such that

$$\text{roots}(\mu_A^s) \subset \text{OLHP} \quad (11.8.4)$$

and

$$\text{roots}(\mu_A^u) \subset \text{CRHP}. \quad (11.8.5)$$

Proposition 11.8.5. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix} S^{-1}, \quad (11.8.6)$$

where $A_1 \in \mathbb{R}^{r \times r}$ is asymptotically stable, $A_{12} \in \mathbb{R}^{r \times (n-r)}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ satisfies $\text{spec}(A_2) \subset \text{CRHP}$. Then,

$$\mu_A^s(A) = S \begin{bmatrix} 0 & C_{12s} \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1}, \quad (11.8.7)$$

where $C_{12s} \in \mathbb{R}^{r \times (n-r)}$ and $\mu_A^s(A_2)$ is nonsingular, and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & C_{12u} \\ 0 & 0 \end{bmatrix} S^{-1}, \quad (11.8.8)$$

where $C_{12u} \in \mathbb{R}^{r \times (n-r)}$ and $\mu_A^u(A_1)$ is nonsingular. Consequently,

$$\mathcal{N}[\mu_A^s(A)] = \mathcal{R}[\mu_A^u(A)] = \mathcal{R}\left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix}\right). \quad (11.8.9)$$

If, in addition, $A_{12} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} 0 & 0 \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1} \quad (11.8.10)$$

and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ 0 & 0 \end{bmatrix} S^{-1}. \quad (11.8.11)$$

Consequently,

$$\mathcal{R}[\mu_A^s(A)] = \mathcal{N}[\mu_A^u(A)] = \mathcal{R}\left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix}\right). \quad (11.8.12)$$

Corollary 11.8.6. Let $A \in \mathbb{R}^{n \times n}$. Then,

$$\mathcal{N}[\mu_A^s(A)] = \mathcal{R}[\mu_A^u(A)] \quad (11.8.13)$$

and

$$\mathcal{N}[\mu_A^u(A)] = \mathcal{R}[\mu_A^s(A)]. \quad (11.8.14)$$

Proof. It follows from Theorem 5.3.5 that there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (11.8.6) is satisfied, where $A_1 \in \mathbb{R}^{r \times r}$ is asymptotically stable, $A_{12} = 0$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ satisfies $\text{spec}(A_2) \subset \text{CRHP}$. The result now follows from Proposition 11.8.5. $\square$

In view of Corollary 11.8.6, we define the *asymptotically stable subspace* $\mathcal{S}_s(A)$ of A by

$$\mathcal{S}_s(A) \triangleq \mathcal{N}[\mu_A^s(A)] = \mathcal{R}[\mu_A^u(A)] \quad (11.8.15)$$

and the *unstable subspace* $\mathcal{S}_u(A)$ of A by

$$\mathcal{S}_u(A) \triangleq \mathcal{N}[\mu_A^u(A)] = \mathcal{R}[\mu_A^s(A)]. \quad (11.8.16)$$

Note that

$$\dim \mathcal{S}_s(A) = \text{def } \mu_A^s(A) = \text{rank } \mu_A^u(A) = \sum_{\substack{\lambda \in \text{spec}(A) \\ \text{Re } \lambda < 0}} \text{am}_A(\lambda) \quad (11.8.17)$$

and

$$\dim \mathcal{S}_u(A) = \text{def } \mu_A^u(A) = \text{rank } \mu_A^s(A) = \sum_{\substack{\lambda \in \text{spec}(A) \\ \text{Re } \lambda \geq 0}} \text{am}_A(\lambda). \quad (11.8.18)$$

Lemma 11.8.7. Let $A \in \mathbb{R}^{n \times n}$, assume that $\text{spec}(A) \subset \text{CRHP}$, let $x \in \mathbb{R}^n$, and assume that $\lim_{t \rightarrow \infty} e^{tA}x = 0$. Then, $x = 0$.

For the following result, note Proposition 11.8.2, Proposition 3.5.3, Fact 3.12.3, Fact 11.18.3, and Proposition 6.1.7.

Proposition 11.8.8. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) $\mathcal{S}_s(A) = \{x \in \mathbb{R}^n: \lim_{t \rightarrow \infty} e^{tA}x = 0\}$.
- ii) $\mu_A^s(A)$ and $\mu_A^u(A)$ are group invertible.
- iii) $P_s \triangleq I - \mu_A^s(A)[\mu_A^s(A)]^\#$ and $P_u \triangleq I - \mu_A^u(A)[\mu_A^u(A)]^\#$ are idempotent.
- iv) $P_s + P_u = I$.
- v) $P_{s\perp} = P_u$ and $P_{u\perp} = P_s$.
- vi) $\mathcal{S}_s(A) = \mathcal{R}(P_s) = \mathcal{N}(P_u)$.
- vii) $\mathcal{S}_u(A) = \mathcal{R}(P_u) = \mathcal{N}(P_s)$.
- viii) $\mathcal{S}_s(A)$ and $\mathcal{S}_u(A)$ are invariant subspaces of A .
- ix) $\mathcal{S}_s(A)$ and $\mathcal{S}_u(A)$ are complementary subspaces.
- x) P_s is the idempotent matrix onto $\mathcal{S}_s(A)$ along $\mathcal{S}_u(A)$.
- xi) P_u is the idempotent matrix onto $\mathcal{S}_u(A)$ along $\mathcal{S}_s(A)$.

Proof. To prove i), let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$ is asymptotically stable and $\text{spec}(A_2) \subset \text{CRHP}$. It then follows from Proposition 11.8.5 that

$$\mathcal{S}_s(A) = \mathcal{N}[\mu_A^s(A)] = \mathcal{R}\left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix}\right).$$

Furthermore,

$$e^{tA} = S \begin{bmatrix} e^{tA_1} & 0 \\ 0 & e^{tA_2} \end{bmatrix} S^{-1}.$$

To prove $\mathcal{S}_s(A) \subseteq \{z \in \mathbb{R}^n: \lim_{t \rightarrow \infty} e^{tA}z = 0\}$, let $x \triangleq S \begin{bmatrix} x_1 \\ 0 \end{bmatrix} \in \mathcal{S}_s(A)$, where $x_1 \in \mathbb{R}^r$. Then, $e^{tA}x = S \begin{bmatrix} e^{tA_1}x_1 \\ 0 \end{bmatrix} \rightarrow 0$ as $t \rightarrow \infty$. Hence, $x \in \{z \in \mathbb{R}^n: \lim_{t \rightarrow \infty} e^{tA}z = 0\}$.

Conversely, to prove $\{z \in \mathbb{R}^n: \lim_{t \rightarrow \infty} e^{tA}z = 0\} \subseteq \mathcal{S}_s(A)$, let $x \triangleq S \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in \mathbb{R}^n$ satisfy $\lim_{t \rightarrow \infty} e^{tA}x = 0$. Hence, $e^{tA_2}x_2 \rightarrow 0$ as $t \rightarrow \infty$. Since $\text{spec}(A_2) \subset \text{CRHP}$, it follows from Lemma 11.8.7 that $x_2 = 0$. Hence, $x \in \mathcal{R}(S \begin{bmatrix} I_r \\ 0 \end{bmatrix}) = \mathcal{S}_s(A)$.

The remaining statements follow directly from Proposition 11.8.5. $\square$

11.9 The Lyapunov Equation

In this section we specialize Theorem 11.7.2 to the linear system (11.8.1).

Corollary 11.9.1. Let $A \in \mathbb{R}^{n \times n}$, and assume there exist a positive-semidefinite matrix $R \in \mathbb{R}^{n \times n}$ and a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ satisfying

$$A^T P + PA + R = 0. \quad (11.9.1)$$

Then, A is Lyapunov stable. If, in addition, for all nonzero $\omega \in \mathbb{R}$,

$$\text{rank} \begin{bmatrix} j\omega I - A \\ R \end{bmatrix} = n, \quad (11.9.2)$$

then A is semistable. Finally, if R is positive definite, then A is asymptotically stable.

Proof. Define $V(x) \triangleq x^T P x$, which satisfies (11.7.2) with $x_e = 0$ and satisfies (11.7.3) for all nonzero $x \in \mathcal{D} = \mathbb{R}^n$. Furthermore, Theorem 11.7.2 implies that $V'(x)f(x) = 2x^T P A x = x^T (A^T P + PA)x = -x^T R x$, which satisfies (11.7.4) for all $x \in \mathbb{R}^n$. Thus, Theorem 11.7.2 implies that A is Lyapunov stable. If, in addition, R is positive definite, then (11.7.5) is satisfied for all $x \neq 0$, and thus A is asymptotically stable.

Alternatively, we now prove the first and third statements without using Theorem 11.7.2. Letting $\lambda \in \text{spec}(A)$, and letting $x \in \mathbb{C}^n$ be an associated eigenvector, it follows that $0 \geq -x^* R x = x^* (A^T P + PA)x = (\bar{\lambda} + \lambda)x^* P x$. Therefore, $\text{spec}(A) \subset \text{CLHP}$. Now, suppose that $j\omega \in \text{spec}(A)$, where $\omega \in \mathbb{R}$, and let $x \in \mathcal{N}[(j\omega I - A)^2]$. Defining $y \triangleq (j\omega I - A)x$, it follows that $(j\omega I - A)y = 0$, and thus $Ay = j\omega y$. Therefore, $-y^* R y = y^* (A^T P + PA)y = -j\omega y^* P y + j\omega y^* P y = 0$, and thus $Ry = 0$. Hence, $0 = x^* R y = -x^* (A^T P + PA)y = -x^* (A^T + j\omega I)P y = y^* P y$. Since P is positive definite, it follows that $y = 0$, that is, $(j\omega I - A)x = 0$. Therefore, $x \in \mathcal{N}(j\omega I - A)$. Now, Proposition 5.5.8 implies that $j\omega$ is semisimple. Therefore, A is Lyapunov stable.

Next, to prove that A is asymptotically stable, let $\lambda \in \text{spec}(A)$, and let $x \in \mathbb{C}^n$ be an associated eigenvector. Thus, $0 > -x^* R x = (\bar{\lambda} + \lambda)x^* P x$, which implies that A is asymptotically stable.

Finally, to prove that A is semistable, let $j\omega \in \text{spec}(A)$, where $\omega \in \mathbb{R}$ is nonzero, and let $x \in \mathbb{C}^n$ be an associated eigenvector. Then,

$$-x^*Rx = x^*(A^TP + PA)x = x^*[(j\omega I - A)^*P + P(j\omega I - A)]x = 0.$$

Therefore, $Rx = 0$, and thus

$$\begin{bmatrix} j\omega I - A \\ R \end{bmatrix} x = 0,$$

which implies that $x = 0$, which contradicts $x \neq 0$. Consequently, $j\omega \notin \text{spec}(A)$ for all nonzero $\omega \in \mathbb{R}$, and thus A is semistable. $\square$

Equation (11.9.1) is a *Lyapunov equation*. Converse results for Corollary 11.9.1 are given by Corollary 11.9.4, Proposition 11.9.6, Proposition 11.9.5, Proposition 11.9.6, and Proposition 12.8.3. The following lemma is useful for analyzing (11.9.1).

Lemma 11.9.2. Assume that $A \in \mathbb{F}^{n \times n}$ is asymptotically stable. Then,

$$\int_0^\infty e^{tA} dt = -A^{-1}. \quad (11.9.3)$$

Proof. Proposition 11.1.4 implies that $\int_0^t e^{\tau A} d\tau = A^{-1}(e^{tA} - I)$. Letting $t \rightarrow \infty$ yields (11.9.3). $\square$

The following result concerns Sylvester's equation. See Fact 5.10.21 and Proposition 7.2.4.

Proposition 11.9.3. Let $A, B, C \in \mathbb{R}^{n \times n}$. Then, there exists a unique matrix $X \in \mathbb{R}^{n \times n}$ satisfying

$$AX + XB + C = 0 \quad (11.9.4)$$

if and only if $B^T \oplus A$ is nonsingular. In this case, X is given by

$$X = -\text{vec}^{-1}[(B^T \oplus A)^{-1} \text{vec } C]. \quad (11.9.5)$$

If, in addition, $B^T \oplus A$ is asymptotically stable, then X is given by

$$X = \int_0^\infty e^{tA} C e^{tB} dt. \quad (11.9.6)$$

Proof. The first two statements follow from Proposition 7.2.4. If $B^T \oplus A$ is asymptotically stable, then it follows from (11.9.5) using Lemma 11.9.2 and Proposition 11.1.7 that

$$\begin{aligned}
X &= \int_0^\infty \text{vec}^{-1} \left(e^{t(B^T \oplus A)} \text{vec } C \right) dt = \int_0^\infty \text{vec}^{-1} \left(e^{tB^T} \otimes e^{tA} \right) \text{vec } C dt \\
&= \int_0^\infty \text{vec}^{-1} \text{vec} (e^{tA} C e^{tB}) dt = \int_0^\infty e^{tA} C e^{tB} dt. \quad \square
\end{aligned}$$

The following result provides a converse to Corollary 11.9.1 for the case of asymptotic stability.

Corollary 11.9.4. Let $A \in \mathbb{R}^{n \times n}$, and let $R \in \mathbb{R}^{n \times n}$. Then, there exists a unique matrix $P \in \mathbb{R}^{n \times n}$ satisfying (11.9.1) if and only if $A \oplus A$ is nonsingular. In this case, if R is symmetric, then P is symmetric. Now, assume that A is asymptotically stable. Then, $P \in \mathbf{S}^n$ is given by

$$P = \int_0^\infty e^{tA^T} R e^{tA} dt. \quad (11.9.7)$$

Finally, if R is (positive semidefinite, positive definite), then P is (positive semidefinite, positive definite).

Proof. First note that $A \oplus A$ is nonsingular if and only if $(A \oplus A)^T = A^T \oplus A^T$ is nonsingular. Now, the first statement follows from Proposition 11.9.3. To prove the second statement, note that $A^T(P - P^T) + (P - P^T)A = 0$, which implies that P is symmetric. Now, suppose that A is asymptotically stable. Then, Fact 11.18.33 implies that $A \oplus A$ is asymptotically stable. Consequently, (11.9.7) follows from (11.9.6). $\square$

The following results also include converse statements. We first consider asymptotic stability.

Proposition 11.9.5. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

- i) A is asymptotically stable.
- ii) For every positive-definite matrix $R \in \mathbb{R}^{n \times n}$ there exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that (11.9.1) is satisfied.
- iii) There exist a positive-definite matrix $R \in \mathbb{R}^{n \times n}$ and a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that (11.9.1) is satisfied.

Proof. The result $i) \implies ii)$ follows from Corollary 11.9.1. The implication $ii) \implies iii)$ is immediate. To prove $iii) \implies i)$, note that, since there exists a positive-semidefinite matrix P satisfying (11.9.1), it follows from Proposition 12.4.3 that (A, C) is observably asymptotically stable. Thus, there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that $A = S \begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix} S^{-1}$ and $C = \begin{bmatrix} C_1 & 0 \end{bmatrix} S^{-1}$, where (C_1, A_1) is observable and A_1 is asymptotically stable. Furthermore, since $(S^{-1}AS, CS)$ is detectable, it follows that A_2 is also asymptotically stable. Consequently, A is asymptotically stable. $\square$

Next, we consider the case of Lyapunov stability.

Proposition 11.9.6. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) If A is Lyapunov stable, then there exist a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ and a positive-semidefinite matrix $R \in \mathbb{R}^{n \times n}$ such that $\text{rank } R = \nu_-(A)$ and such that (11.9.1) is satisfied.
- ii) If there exist a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ and a positive-semidefinite matrix $R \in \mathbb{R}^{n \times n}$ such that (11.9.1) is satisfied, then A is Lyapunov stable.

Proof. To prove i), suppose that A is Lyapunov stable. Then, it follows from Theorem 5.3.5 and Definition 11.8.1 that there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that $A = S \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} S^{-1}$ is in real Jordan form, where $A_1 \in \mathbb{R}^{n_1 \times n_1}$, $A_2 \in \mathbb{R}^{n_2 \times n_2}$, $\text{spec}(A_1) \subset j\mathbb{R}$, A_1 is semisimple, and $\text{spec}(A_2) \subset \text{OLHP}$. Next, it follows from Fact 5.9.4 that there exists a nonsingular matrix $S_1 \in \mathbb{R}^{n_1 \times n_1}$ such that $A_1 = S_1 J_1 S_1^{-1}$, where $J_1 \in \mathbb{R}^{n_1 \times n_1}$ is skew symmetric. Then, it follows that $A_1^T P_1 + P_1 A_1 = S_1^{-T} (J_1 + J_1^T) S_1^{-1} = 0$, where $P_1 \triangleq S_1^{-T} S_1^{-1}$ is positive definite. Next, let $R_2 \in \mathbb{R}^{n_2 \times n_2}$ be positive definite, and let $P_2 \in \mathbb{R}^{n_2 \times n_2}$ be the positive-definite solution of $A_2^T P_2 + P_2 A_2 + R_2 = 0$. Hence, (11.9.1) is satisfied with $P \triangleq S^{-T} \begin{bmatrix} P_1 & 0 \\ 0 & P_2 \end{bmatrix} S^{-1}$ and $R \triangleq S^{-T} \begin{bmatrix} 0 & 0 \\ 0 & R_2 \end{bmatrix} S^{-1}$.

To prove ii), suppose there exist a positive-semidefinite matrix $R \in \mathbb{R}^{n \times n}$ and a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that (11.9.1) is satisfied. Let $\lambda \in \text{spec}(A)$, and let $x \in \mathbb{R}^n$ be an eigenvector of A associated with λ . It thus follows from (11.9.1) that $0 = x^* A^T P x + x^* P A x + x^* R x = (\lambda + \bar{\lambda}) x^* P x + x^* R x$. Therefore, $\text{Re } \lambda = -x^* R x / (2x^* P x)$, which shows that $\text{spec}(A) \subset \text{CLHP}$. Now, let $j\omega \in \text{spec}(A)$, and suppose that $x \in \mathbb{R}^n$ satisfies $(j\omega I - A)^2 x = 0$. Then, $(j\omega I - A)y = 0$, where $y = (j\omega I - A)x$. Computing $0 = y^* (A^T P + P A) y + y^* R y$ yields $y^* R y = 0$ and thus $R y = 0$. Therefore, $(A^T P + P A) y = 0$, and thus $y^* P y = (A^T + j\omega I) P y = 0$. Since P is positive definite, it follows that $y = (j\omega I - A)x = 0$. Therefore, $\mathcal{N}(j\omega I - A) = \mathcal{N}[(j\omega I - A)^2]$. Hence, it follows from Proposition 5.5.8 that $j\omega$ is semisimple. $\square$

Corollary 11.9.7. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) A is Lyapunov stable if and only if there exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that $A^T P + P A$ is negative semidefinite.
- ii) A is asymptotically stable if and only if there exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that $A^T P + P A$ is negative definite.

11.10 Discrete-Time Stability Theory

The theory of difference equations is concerned with the solutions of discrete-time dynamical systems of the form

$$x_{k+1} = f(x_k), \quad (11.10.1)$$

where $f: \mathbb{R}^n \rightarrow \mathbb{R}^n$, $k \in \mathbb{N}$, $x_k \in \mathbb{R}^n$, and x_0 is the initial condition. The solution $x_k \equiv x_e$ is an equilibrium of (11.10.1) if $x_e = f(x_e)$.

A linear discrete-time system has the form

$$x_{k+1} = Ax_k, \quad (11.10.2)$$

where $A \in \mathbb{R}^{n \times n}$. For $k \in \mathbb{N}$, x_k is given by

$$x_k = A^k x_0. \quad (11.10.3)$$

The behavior of the sequence $(x_k)_{k=0}^\infty$ is determined by the stability of A . To study the stability of discrete-time systems it is helpful to define the *open unit disk* (OUD) and the *closed unit disk* (CUD) by

$$\text{OUD} \triangleq \{x \in \mathbb{C}: |x| < 1\} \quad (11.10.4)$$

and

$$\text{CUD} \triangleq \{x \in \mathbb{C}: |x| \leq 1\}. \quad (11.10.5)$$

Definition 11.10.1. For $A \in \mathbb{F}^{n \times n}$, define the following classes of matrices:

- i) A is *discrete-time Lyapunov stable* if $\text{spec}(A) \subset \text{CUD}$ and, if $\lambda \in \text{spec}(A)$ and $|\lambda| = 1$, then λ is semisimple.
- ii) A is *discrete-time semistable* if $\text{spec}(A) \subset \text{OUD} \cup \{1\}$ and, if $1 \in \text{spec}(A)$, then 1 is semisimple.
- iii) A is *discrete-time asymptotically stable* if $\text{spec}(A) \subset \text{OUD}$.

Proposition 11.10.2. Let $A \in \mathbb{R}^{n \times n}$ and consider the linear discrete-time system (11.10.2). Then, the following statements are equivalent:

- i) A is discrete-time Lyapunov stable.
- ii) For every initial condition $x_0 \in \mathbb{R}^n$, the sequence $\{\|x_k\|\}_{k=1}^\infty$ is bounded, where $\|\cdot\|$ is a norm on $\mathbb{R}^n$.
- iii) For every initial condition $x_0 \in \mathbb{R}^n$, the sequence $\{\|A^k x_0\|\}_{k=1}^\infty$ is bounded, where $\|\cdot\|$ is a norm on $\mathbb{R}^n$.
- iv) The sequence $\{\|A^k\|\}_{k=1}^\infty$ is bounded, where $\|\cdot\|$ is a norm on $\mathbb{R}^{n \times n}$.

The following statements are equivalent:

- v) A is discrete-time semistable.
- vi) $\lim_{k \rightarrow \infty} A^k$ exists. In fact, $\lim_{k \rightarrow \infty} A^k = I - (I - A)(I - A)^\#$.
- vii) For every initial condition $x_0 \in \mathbb{R}^n$, $\lim_{k \rightarrow \infty} x_k$ exists.

The following statements are equivalent:

- viii) A is discrete-time asymptotically stable.
- ix) $\text{sprad}(A) < 1$.
- x) For every initial condition $x_0 \in \mathbb{R}^n$, $\lim_{k \rightarrow \infty} x_k = 0$.
- xi) For every initial condition $x_0 \in \mathbb{R}^n$, $A^k x_0 \rightarrow 0$ as $k \rightarrow \infty$.
- xii) $A^k \rightarrow 0$ as $k \rightarrow \infty$.

The following definition concerns the discrete-time stability of a polynomial.

Definition 11.10.3. Let $p \in \mathbb{R}[s]$. Then, define the following terminology:

- i) p is *discrete-time Lyapunov stable* if $\text{roots}(p) \subset \text{CUD}$ and, if λ is an imaginary root of p , then $m_p(\lambda) = 1$.
- ii) p is *discrete-time semistable* if $\text{roots}(p) \subset \text{OUD} \cup \{1\}$ and, if $1 \in \text{roots}(p)$, then $m_p(1) = 1$.
- iii) p is *discrete-time asymptotically stable* if $\text{roots}(p) \subset \text{OUD}$.

Proposition 11.10.4. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) A is discrete-time Lyapunov stable if and only if μ_A is discrete-time Lyapunov stable.
- ii) A is discrete-time semistable if and only if μ_A is discrete-time semistable.

Furthermore, the following statements are equivalent:

- iii) A is discrete-time asymptotically stable.
- iv) μ_A is discrete-time asymptotically stable.
- v) χ_A is discrete-time asymptotically stable.

We now consider the *discrete-time Lyapunov equation*

$$P = A^T P A + R = 0. \quad (11.10.6)$$

Proposition 11.10.5. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

- i) A is discrete-time asymptotically stable.
- ii) For every positive-definite matrix $R \in \mathbb{R}^{n \times n}$ there exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that (11.10.6) is satisfied.
- iii) There exist a positive-definite matrix $R \in \mathbb{R}^{n \times n}$ and a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that (11.10.6) is satisfied.

Proposition 11.10.6. Let $A \in \mathbb{R}^{n \times n}$. Then, A is discrete-time Lyapunov-stable if and only if there exist a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ and a positive-semidefinite matrix $R \in \mathbb{R}^{n \times n}$ such that (11.10.6) is satisfied.

11.11 Facts on Matrix Exponential Formulas

Fact 11.11.1. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) If $A^2 = 0$, then $e^{tA} = I + tA$.
- ii) If $A^2 = I$, then $e^{tA} = (\cosh t)I + (\sinh t)A$.
- iii) If $A^2 = -I$, then $e^{tA} = (\cos t)I + (\sin t)A$.

iv) If $A^2 = A$, then $e^{tA} = I + (e^t - 1)A$.

v) If $A^2 = -A$, then $e^{tA} = I + (1 - e^{-t})A$.

vi) If $\text{rank } A = 1$ and $\text{tr } A = 0$, then $e^{tA} = I + tA$.

vii) If $\text{rank } A = 1$ and $\text{tr } A \neq 0$, then $e^{tA} = I + \frac{e^{(\text{tr } A)t} - 1}{\text{tr } A} A$.

(Remark: See [1085].)

Fact 11.11.2. Let $A \triangleq \begin{bmatrix} 0 & I_n \\ I_n & 0 \end{bmatrix}$. Then,

$$e^{tA} = (\cosh t)I_{2n} + (\sinh t)A.$$

Furthermore,

$$e^{tJ_{2n}} = (\cos t)I_{2n} + (\sin t)J_{2n}.$$

Fact 11.11.3. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is skew symmetric. Then, $\{e^{\theta A} : \theta \in \mathbb{R}\} \subseteq \text{SO}(n)$ is a group. If, in addition, $n = 2$, then

$$\{e^{\theta J_2} : \theta \in \mathbb{R}\} = \text{SO}(2).$$

(Remark: Note that $e^{\theta J_2} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$. See Fact 3.11.6.)

Fact 11.11.4. Let $A \in \mathbb{R}^{n \times n}$, where

$$A \triangleq \begin{bmatrix} 0 & 1 & 0 & 0 & \cdots & 0 \\ 0 & 0 & 2 & 0 & \cdots & 0 \\ 0 & 0 & 0 & 3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \ddots & n-1 \\ 0 & 0 & 0 & 0 & \cdots & 0 \end{bmatrix}.$$

Then,

$$e^A = \begin{bmatrix} \binom{0}{0} & \binom{1}{0} & \binom{2}{0} & \binom{3}{0} & \cdots & \binom{n-1}{0} \\ 0 & \binom{1}{1} & \binom{2}{1} & \binom{3}{1} & \cdots & \binom{n-1}{1} \\ 0 & 0 & \binom{2}{2} & \binom{3}{2} & \cdots & \binom{n-1}{2} \\ \vdots & \vdots & \vdots & \ddots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \ddots & \binom{n-1}{n-2} \\ 0 & 0 & 0 & 0 & \cdots & \binom{n-1}{n-1} \end{bmatrix}.$$

Furthermore, if $k \geq n$, then

$$\sum_{i=1}^k i^{n-1} = \begin{bmatrix} 1^{n-1} & 2^{n-1} & \cdots & n^{n-1} \end{bmatrix} e^{-A} \begin{bmatrix} \binom{k}{1} \\ \vdots \\ \binom{k}{n} \end{bmatrix}.$$

(Proof: See [73].) (Remark: For related results, see [5], where A is called the *creation matrix*. See Fact 5.16.3.)

Fact 11.11.5. Let $A \in \mathbb{F}^{3 \times 3}$. If $\text{spec}(A) = \{\lambda\}$, then

$$e^{tA} = e^{\lambda t} \left[I + t(A - \lambda I) + \frac{1}{2}t^2(A - \lambda I)^2 \right].$$

If $\text{mspec}(A) = \{\lambda, \lambda, \mu\}_{\text{ms}}$, where $\mu \neq \lambda$, then

$$e^{tA} = e^{\lambda t} [I + t(A - \lambda I)] + \left[\frac{e^{\mu t} - e^{\lambda t}}{(\mu - \lambda)^2} - \frac{te^{\lambda t}}{\mu - \lambda} \right] (A - \lambda I)^2.$$

If $\text{spec}(A) = \{\lambda, \mu, \nu\}$, then

$$\begin{aligned} e^{tA} &= \frac{e^{\lambda t}}{(\lambda - \mu)(\lambda - \nu)} (A - \mu I)(A - \nu I) + \frac{e^{\mu t}}{(\mu - \lambda)(\mu - \nu)} (A - \lambda I)(A - \nu I) \\ &\quad + \frac{e^{\nu t}}{(\nu - \lambda)(\nu - \mu)} (A - \lambda I)(A - \mu I). \end{aligned}$$

(Proof: See [67].) (Remark: Additional expressions are given in [2, 175, 191, 321, 640, 1085, 1088].)

Fact 11.11.6. Let $x \in \mathbb{R}^3$, assume that x is nonzero, and define $\theta \triangleq \|x\|_2$. Then,

$$\begin{aligned} e^{K(x)} &= I + \frac{\sin \theta}{\theta} K(x) + \frac{1 - \cos \theta}{\theta^2} K^2(x) \\ &= I + \frac{\sin \theta}{\theta} K(x) + \frac{1}{2} \left[\frac{\sin(\frac{1}{2}\theta)}{\frac{1}{2}\theta} \right]^2 K^2(x) \\ &= (\cos \theta)I + \frac{\sin \theta}{\theta} K(x) + \frac{1 - \cos \theta}{\theta^2} xx^T. \end{aligned}$$

Furthermore,

$$e^{K(x)}x = x,$$

$$\text{spec}[e^{K(x)}] = \{1, e^{J\|x\|_2}, e^{-J\|x\|_2}\},$$

and

$$\text{tr } e^{K(x)} = 1 + 2\cos \theta = 1 + 2\cos \|x\|_2.$$

(Proof: The Cayley-Hamilton theorem or Fact 3.10.1 implies that $K^3(x) + \theta^2 K(x) = 0$. Then, every term $K^k(x)$ in the expansion of $e^{K(x)}$ can be expressed in terms of $K(x)$ or $K^2(x)$. Finally, Fact 3.10.1 implies that $\theta^2 I + K^2(x) = xx^T$.) (Remark: Fact 11.11.7 shows that, for all $z \in \mathbb{R}^3$, $e^{K(x)}z$ is the counterclockwise (right-hand-rule) rotation of z about the vector x through the angle θ , which is given by the Euclidean norm of x . In Fact 3.11.8, the cross product is used to construct the pivot vector x from a given pair of vectors having the same length.)

Fact 11.11.7. Let $x, y \in \mathbb{R}^3$, and assume that x and y are nonzero. Then, there exists a skew-symmetric matrix $A \in \mathbb{R}^{3 \times 3}$ such that $y = e^A x$ if and only if $x^T x = y^T y$. If $x \neq -y$, then one such matrix is $A = \theta K(z)$, where

$$z \triangleq \frac{1}{\|x \times y\|_2} x \times y$$

and

$$\theta \triangleq \cos^{-1} \left(\frac{x^T y}{\|x\|_2 \|y\|_2} \right).$$

If $x = -y$, then one such matrix is $A = \pi K(z)$, where $z \triangleq \|y\|_2^{-1} \nu \times y$ and $\nu \in \{y\}^\perp$ satisfies $\nu^T \nu = 1$. (Proof: This result follows from Fact 3.11.8 and Fact 11.11.6, which provide equivalent expressions for an orthogonal matrix that transforms a given vector into another given vector having the same length. This result thus provides a geometric interpretation for Fact 11.11.6.) (Remark: Note that z is the unit vector perpendicular to the plane containing x and y , where the direction of z is determined by the right-hand rule. An intuitive proof is to let x be the initial condition to the differential equation $\dot{w}(t) = K(z)w(t)$, that is, $w(0) = x$, where $t \in [0, \theta]$. Then, the derivative $\dot{w}(t)$ lies in the x, y plane and is perpendicular to $w(t)$ for all $t \in [0, \theta]$. Hence, $y = w(\theta)$.) (Remark: Since $\det e^A = e^{\text{tr} A} = 1$, it follows that every pair of vectors in $\mathbb{R}^3$ having the same Euclidean length are related by a *proper rotation*. See Fact 3.9.5 and Fact 3.14.4. This is a linear interpolation problem. See Fact 3.9.5, Fact 3.11.8, and [773].) (Remark: See Fact 3.11.31.) (Remark: Parameterizations of $\text{SO}(3)$ are considered in [1195, 1246].) (Problem: Extend this result to $\mathbb{R}^n$. See [135, 1164].)

Fact 11.11.8. Let $A \in \text{SO}(3)$, let $z \in \mathbb{R}^3$ be an eigenvector of A corresponding to the eigenvalue 1 of A , assume that $\|z\|_2 = 1$, assume that $\text{tr } A > -1$, and let $\theta \in (-\pi, \pi)$ satisfy $\text{tr } A = 1 + 2\cos \theta$. Then,

$$A = e^{\theta K(z)}.$$

(Remark: See Fact 5.11.2.)

Fact 11.11.9. Let $x, y \in \mathbb{R}^3$, and assume that x and y are nonzero. Then, $x^T x = y^T y$ if and only if

$$y = e^{\frac{\theta}{\|x \times y\|_2} (yx^T - xy^T)} x,$$

where

$$\theta \triangleq \cos^{-1} \left(\frac{x^T y}{\|x\|_2 \|y\|_2} \right).$$

(Proof: Use Fact 11.11.7.) (Remark: Note that $K(x \times y) = yx^T - xy^T$.)

Fact 11.11.10. Let $A \in \mathbb{R}^{3 \times 3}$, assume that $A \in \text{SO}(3)$ and $\text{tr } A > -1$, and let $\theta \in (-\pi, \pi)$ satisfy $\text{tr } A = 1 + 2\cos \theta$. Then,

$$\log A = \begin{cases} 0, & \theta = 0, \\ \frac{\theta}{2\sin \theta} (A - A^T), & \theta \neq 0. \end{cases}$$

(Proof: See [746, p. 364] and [1013].) (Remark: See Fact 11.15.10.)

Fact 11.11.11. Let $x \in \mathbb{R}^3$, assume that x is nonzero, and define $\theta \triangleq \|x\|_2$. Then,

$$K(x) = \frac{\theta}{2\sin \theta} [e^{K(x)} - e^{-K(x)}].$$

(Proof: Use Fact 11.11.10.) (Remark: See Fact 3.10.1.)

Fact 11.11.12. Let $A \in \text{SO}(3)$, let $x, y \in \mathbb{R}^3$, and assume that $x^T x = y^T y$. Then, $Ax = y$ if and only if, for all $t \in \mathbb{R}$,

$$Ae^{tK(x)}A^{-1} = e^{tK(y)}.$$

(Proof: See [887].)

Fact 11.11.13. Let $x, y, z \in \mathbb{R}^3$. Then, the following statements are equivalent:

i) For every $A \in \text{SO}(3)$, there exist $\alpha, \beta, \gamma \in \mathbb{R}$ such that

$$A = e^{\alpha K(x)} e^{\beta K(y)} e^{\gamma K(z)}.$$

ii) $y^T x = 0$ and $y^T z = 0$.

(Proof: See [887].) (Remark: This result is due to Davenport.) (Problem: Given $A \in \text{SO}(3)$, determine α, β, γ .)

Fact 11.11.14. Let $A \in \mathbb{R}^{4 \times 4}$, and assume that A is skew symmetric with $\text{mspec}(A) = \{j\omega, -j\omega, j\mu, -j\mu\}_{\text{ms}}$. If $\omega \neq \mu$, then

$$e^A = a_3 A^3 + a_2 A^2 + a_1 A + a_0 I,$$

where

$$\begin{aligned} a_3 &= (\omega^2 - \mu^2)^{-1} \left(\frac{1}{\mu} \sin \mu - \frac{1}{\omega} \sin \omega \right), \\ a_2 &= (\omega^2 - \mu^2)^{-1} (\cos \mu - \cos \omega), \\ a_1 &= (\omega^2 - \mu^2)^{-1} \left(\frac{\omega^2}{\mu} \sin \mu - \frac{\mu^2}{\omega} \sin \omega \right), \\ a_0 &= (\omega^2 - \mu^2)^{-1} (\omega^2 \cos \mu - \mu^2 \cos \omega). \end{aligned}$$

If $\omega = \mu$, then

$$e^A = (\cos \omega) I + \frac{\sin \omega}{\omega} A.$$

(Proof: See [607, p. 18] and [1088].) (Remark: There are typographical errors in [607, p. 18] and [1088].) (Remark: See Fact 4.9.20 and Fact 4.10.2.)

Fact 11.11.15. Let $a, b, c \in \mathbb{R}$, define the skew-symmetric matrix $A \in \mathbb{R}^{4 \times 4}$, by either

$$A \triangleq \begin{bmatrix} 0 & a & b & c \\ -a & 0 & -c & b \\ -b & c & 0 & -a \\ -c & -b & a & 0 \end{bmatrix}$$

or

$$A \triangleq \begin{bmatrix} 0 & a & b & c \\ -a & 0 & c & -b \\ -b & -c & 0 & a \\ -c & b & -a & 0 \end{bmatrix},$$

and define $\theta \triangleq \sqrt{a^2 + b^2 + c^2}$. Then,

$$\text{mspec}(A) = \{j\theta, -j\theta, j\theta, -j\theta\}_{\text{ms}}.$$

Furthermore,

$$A^k = \begin{cases} (-1)^{k/2} \theta^k I, & k \text{ even}, \\ (-1)^{(k-1)/2} \theta^{k-1} A, & k \text{ odd}, \end{cases}$$

and

$$e^A = (\cos \theta)I + \frac{\sin \theta}{\theta}A.$$

(Proof: See [1357].) (Remark: $(\sin 0)/0 = 1$.) (Remark: The skew-symmetric matrix A arises in the kinematic relationship between the angular velocity vector and quaternion (Euler-parameter) rates. See [152, p. 385].) (Remark: The two matrices A are similar. To show this, note that Fact 5.9.9 implies that A and $-A$ are similar. Then, apply the similarity transformation $S = \text{diag}(-1, 1, 1, 1)$.) (Remark: See Fact 4.9.20 and Fact 4.10.2.)

Fact 11.11.16. Let $x \in \mathbb{R}^3$, and define the skew-symmetric matrix $A \in \mathbb{R}^{4 \times 4}$ by

$$A = \begin{bmatrix} 0 & -x^T \\ x & -K(x) \end{bmatrix}.$$

Then, for all $t \in \mathbb{R}$,

$$e^{\frac{1}{2}tA} = \cos(\frac{1}{2}\|x\|t)I_4 + \frac{\sin(\frac{1}{2}\|x\|t)}{\|x\|}A.$$

(Proof: See [733, p. 34].) (Remark: The matrix $\frac{1}{2}A$ characterizes quaternion rates in terms of the angular velocity vector.)

Fact 11.11.17. Let $a, b \in \mathbb{R}^3$, define the skew-symmetric matrix $A \in \mathbb{R}^{4 \times 4}$ by

$$A = \begin{bmatrix} K(a) & b \\ -b^T & 0 \end{bmatrix},$$

and assume that $a^T b = 0$. Then,

$$e^A = I_4 + \frac{\sin \alpha}{\alpha}A + \frac{1 - \cos \alpha}{\alpha^2}A^2,$$

where $\alpha \triangleq \sqrt{a^T a + b^T b}$. (Proof: See [1334].) (Remark: See Fact 4.9.20 and Fact 4.10.2.)

Fact 11.11.18. Let $a, b \in \mathbb{R}^{n-1}$, define $A \in \mathbb{R}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} 0 & a^T \\ b & 0_{(n-1) \times (n-1)} \end{bmatrix},$$

and define $\alpha \triangleq \sqrt{|a^T b|}$. Then, the following statements hold:

i) If $a^T b < 0$, then

$$e^{tA} = I + \frac{\sin \alpha}{\alpha}A + \frac{1}{2} \left[\frac{\sin(\alpha/2)}{\alpha/2} \right]^2 A^2.$$

ii) If $a^T b = 0$, then

$$e^{tA} = I + A + \frac{1}{2}A^2.$$

iii) If $a^T b > 0$, then

$$e^{tA} = I + \frac{\sinh \alpha}{\alpha}A + \frac{1}{2} \left[\frac{\sinh(\alpha/2)}{\alpha/2} \right]^2 A^2.$$

(Proof: See [1480].)

11.12 Facts on the Matrix Sine and Cosine

Fact 11.12.1. Let $A \in \mathbb{C}^{n \times n}$, and define

$$\sin A \triangleq A - \frac{1}{3!}A^3 + \frac{1}{5!}A^5 - \frac{1}{7!}A^7 + \cdots$$

and

$$\cos A \triangleq I - \frac{1}{2!}A^2 + \frac{1}{4!}A^4 - \frac{1}{6!}A^6 + \cdots.$$

Then, the following statements hold:

- i) $\sin A = \frac{1}{2j}(e^{jA} - e^{-jA})$.
- ii) $\cos A = \frac{1}{2}(e^{jA} + e^{-jA})$.
- iii) $\sin^2 A + \cos^2 A = I$.
- iv) $\sin(2A) = 2(\sin A)\cos A$.
- v) $\cos(2A) = 2(\cos^2 A) - I$.
- vi) If A is real, then $\sin A = \operatorname{Re} e^{jA}$ and $\cos A = \operatorname{Re} e^{jA}$.
- vii) $\sin(A \oplus B) = (\sin A) \otimes \cos B - (\cos A) \otimes \sin B$.
- viii) $\cos(A \oplus B) = (\cos A) \otimes \cos B - (\sin A) \otimes \sin B$.
- ix) If A is involutory and k is an integer, then $\cos(k\pi A) = (-1)^k I$.

Furthermore, the following statements are equivalent:

- x) For all $t \in \mathbb{R}$, $\sin[(A + B)t] = \sin(tA)\cos(tB) + \cos(tA)\sin(tB)$.
- xi) For all $t \in \mathbb{R}$, $\cos[(A + B)t] = \cos(tA)\cos(tB) - \sin(tA)\sin(tB)$.
- xii) $AB = BA$.

(Proof: See [683, pp. 287, 288, 300].)

11.13 Facts on the Matrix Exponential for One Matrix

Fact 11.13.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is (lower triangular, upper triangular). Then, so is e^A . If, in addition, A is Toeplitz, then so is e^A . (Remark: See Fact 3.18.7.)

Fact 11.13.2. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$\operatorname{sprad}(e^A) = e^{\operatorname{spabs}(A)}.$$

Fact 11.13.3. Let $A \in \mathbb{R}^{n \times n}$, and let $X_0 \in \mathbb{R}^{n \times n}$. Then, the matrix differential equation

$$\begin{aligned}\dot{X}(t) &= AX(t), \\ X(0) &= X_0,\end{aligned}$$

where $t \geq 0$, has the unique solution

$$X(t) = e^{tA}X_0.$$

Fact 11.13.4. Let $A: [0, T] \mapsto \mathbb{R}^{n \times n}$, assume that A is continuous, and let $X_0 \in \mathbb{R}^{n \times n}$. Then, the matrix differential equation

$$\begin{aligned}\dot{X}(t) &= A(t)X(t), \\ X(0) &= X_0\end{aligned}$$

has a unique solution $X: [0, T] \mapsto \mathbb{R}^{n \times n}$. Furthermore, for all $t \in [0, T]$,

$$\det X(t) = e^{\int_0^t \operatorname{tr} A(\tau) d\tau} \det X_0.$$

Therefore, if X_0 is nonsingular, then $X(t)$ is nonsingular for all $t \in [0, T]$. If, in addition, for all $t_1, t_2 \in [0, T]$,

$$A(t_2) \int_{t_1}^{t_2} A(\tau) d\tau = \int_{t_1}^{t_2} A(\tau) d\tau A(t_2),$$

then, for all $t \in [0, T]$,

$$X(t) = e^{\int_0^t A(\tau) d\tau} X_0.$$

(Proof: It follows from Fact 10.11.19 that $(d/dt) \det X = \operatorname{tr}(X^A \dot{X}) = \operatorname{tr}(X^A AX) = \operatorname{tr}(XX^A A) = (\det X) \operatorname{tr} A$. This proof is given in [563]. See also [711, pp. 507, 508] and [1150, pp. 64–66].) (Remark: See Fact 11.13.4.) (Remark: The first result is *Jacobi's identity*.) (Remark: If the commutativity assumption does not hold, then the solution is given by the *Peano-Baker series*. See [1150, Chapter 3]. Alternative expressions for $X(t)$ are given by the Magnus, Fer, Baker-Campbell-Hausdorff-Dynkin, Wei-Norman, Goldberg, and Zassenhaus expansions. See [228, 443, 745, 746, 830, 949, 1056, 1244, 1274, 1414, 1415, 1419] and [621, pp. 118–120].)

Fact 11.13.5. Let $A: [0, T] \mapsto \mathbb{R}^{n \times n}$, assume that A is continuous, let $B: [0, T] \mapsto \mathbb{R}^{n \times m}$, assume that B is continuous, let $X: [0, T] \mapsto \mathbb{R}^{n \times n}$ satisfy the matrix differential equation

$$\begin{aligned}\dot{X}(t) &= A(t)X(t), \\ X(0) &= I,\end{aligned}$$

define

$$\Phi(t, \tau) \triangleq X(t)X^{-1}(\tau),$$

let $u: [0, T] \mapsto \mathbb{R}^m$, and assume that u is continuous. Then, the vector differential equation

$$\begin{aligned}\dot{x}(t) &= A(t)x(t) + B(t)u(t), \\ x(0) &= x_0\end{aligned}$$

has the unique solution

$$x(t) = X(t)x_0 + \int_0^t \Phi(t, \tau)B(\tau)u(\tau)d\tau.$$

(Remark: $\Phi(t, \tau)$ is the *state transition matrix*.)

Fact 11.13.6. Let $A \in \mathbb{R}^{n \times n}$, let $\lambda \in \text{spec}(A)$, and let $v \in \mathbb{C}^n$ be an eigenvector of A associated with λ . Then, for all $t \geq 0$,

$$x(t) \triangleq \text{Re}(e^{\lambda t} v)$$

satisfies $\dot{x}(t) = Ax(t)$. (Remark: $x(t)$ is an *eigensolution*.)

Fact 11.13.7. Let $A \in \mathbb{R}^{n \times n}$, let $\lambda \in \text{spec}(A)$, and let $(v_1, \dots, v_k) \in (\mathbb{C}^n)^k$ be a Jordan chain of A associated with λ . Then, for all $t \geq 0$ and all $\hat{k}$ such that $1 \leq \hat{k} \leq k$,

$$x(t) \triangleq \text{Re} \left[e^{\lambda t} \left(\frac{1}{(\hat{k}-1)!} t^{\hat{k}-1} v_1 + \dots + t v_{\hat{k}-1} + v_{\hat{k}} \right) \right]$$

satisfies $\dot{x}(t) = Ax(t)$. (Remark: See Fact 5.14.8 for the definition of a Jordan chain.) (Remark: $x(t)$ is a *generalized eigensolution*.) (Example: Let $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$, $\lambda = 0$, $\hat{k} = 2$, $v_1 = \begin{bmatrix} \beta \\ 0 \end{bmatrix}$, and $v_2 = \begin{bmatrix} 0 \\ \beta \end{bmatrix}$. Then, $x(t) = t v_1 + v_2 = \begin{bmatrix} \beta t \\ \beta \end{bmatrix}$ is a generalized eigensolution. Alternatively, choosing $\hat{k} = 1$ yields the eigensolution $x(t) = v_1 = \begin{bmatrix} \beta \\ 0 \end{bmatrix}$. Note that β represents velocity for the generalized eigensolution and position for the eigensolution. See [1062].)

Fact 11.13.8. Let $S: [t_0, t_1] \rightarrow \mathbb{R}^{n \times n}$ be differentiable. Then, for all $t \in [t_0, t_1]$,

$$\frac{d}{dt} S^2(t) = \dot{S}(t)S(t) + S(t)\dot{S}(t).$$

Let $S_1: [t_0, t_1] \rightarrow \mathbb{R}^{n \times m}$ and $S_2: [t_0, t_1] \rightarrow \mathbb{R}^{m \times l}$ be differentiable. Then, for all $t \in [t_0, t_1]$,

$$\frac{d}{dt} S_1(t)S_2(t) = \dot{S}_1(t)S_2(t) + S_1(t)\dot{S}_2(t).$$

Fact 11.13.9. Let $A \in \mathbb{F}^{n \times n}$, and define $A_1 \triangleq \frac{1}{2}(A + A^*)$ and $A_2 \triangleq \frac{1}{2}(A - A^*)$. Then, $A_1 A_2 = A_2 A_1$ if and only if A is normal. In this case, $e^{A_1} e^{A_2}$ is the polar decomposition of e^A . (Remark: See Fact 3.7.28.) (Problem: Obtain the polar decomposition of e^A when A is not normal.)

Fact 11.13.10. Let $A \in \mathbb{F}^{n \times m}$, and assume that $\text{rank } A = m$. Then,

$$A^+ = \int_0^\infty e^{-tA^*A} A^* dt.$$

Fact 11.13.11. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is nonsingular. Then,

$$A^{-1} = \int_0^\infty e^{-tA^*A} dt A^*.$$

Fact 11.13.12. Let $A \in \mathbb{F}^{n \times n}$, and let $k \triangleq \text{ind } A$. Then,

$$A^D = \int_0^\infty e^{-tA^k A^{(2k+1)*} A^{k+1}} dt A^k A^{(2k+1)*} A^k.$$

(Proof: See [570].)

Fact 11.13.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that $\text{ind } A = 1$. Then,

$$A^\# = \int_0^\infty e^{-tAA^{3*}A^2} dt AA^{3*}A.$$

(Proof: See Fact 11.13.12.)

Fact 11.13.14. Let $A \in \mathbb{F}^{n \times n}$, and let $k \triangleq \text{ind } A$. Then,

$$\int_0^t e^{\tau A} d\tau = A^D(e^{tA} - I) + (I - AA^D)(tI + \frac{1}{2!}t^2A + \cdots + \frac{1}{k!}t^kA^{k-1}).$$

If, in particular, A is group invertible, then

$$\int_0^t e^{\tau A} d\tau = A^\#(e^{tA} - I) + (I - AA^\#)t.$$

Fact 11.13.15. Let $A \in \mathbb{F}^{n \times n}$, let $\text{mspec}(A) = \{\lambda_1, \dots, \lambda_r, 0, \dots, 0\}_{\text{ms}}$, where $\lambda_1, \dots, \lambda_r$ are nonzero, and let $t > 0$. Then,

$$\det \int_0^t e^{\tau A} d\tau = t^{n-r} \prod_{i=1}^r \lambda_i^{-1} (e^{\lambda_i t} - 1).$$

Hence, $\det \int_0^t e^{\tau A} d\tau \neq 0$ if and only if, for every nonzero integer k , $2k\pi j/t \notin \text{spec}(A)$. Finally, $\det(e^{tA} - I) \neq 0$ if and only if $\det A \neq 0$ and $\det \int_0^t e^{\tau A} d\tau \neq 0$.

Fact 11.13.16. Let $A \in \mathbb{F}^{n \times n}$, and assume that there exists $\alpha \in \mathbb{R}$ such that $\text{spec}(A) \subset \{z \in \mathbb{C}: \alpha \leq \text{Im } z < 2\pi + \alpha\}$. Then, e^A is (diagonal, upper triangular, lower triangular) if and only if A is. (Proof: See [932].)

Fact 11.13.17. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) If A is unipotent, then the series (11.5.1) is finite, $\log A$ exists and is nilpotent, and $e^{\log A} = A$.
- ii) If A is nilpotent, then e^A is unipotent and $\log e^A = A$.

(Proof: See [624, p. 60].)

Fact 11.13.18. Let $B \in \mathbb{R}^{n \times n}$. Then, there exists a normal matrix $A \in \mathbb{R}^{n \times n}$ such that $B = e^A$ if and only if B is normal, nonsingular, and every negative eigenvalue of B has even algebraic multiplicity.

Fact 11.13.19. Let $C \in \mathbb{R}^{n \times n}$, assume that C is nonsingular, and let $k \geq 1$. Then, there exists a matrix $B \in \mathbb{R}^{n \times n}$ such that $C^{2k} = e^B$. (Proof: Use Proposition 11.4.3 with $A = C^2$, and note that every negative eigenvalue $-\alpha < 0$ of C^2 arises as the square of complex conjugate eigenvalues $\pm j\sqrt{\alpha}$ of C .)

11.14 Facts on the Matrix Exponential for Two or More Matrices

Fact 11.14.1. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{n \times m}$, and $C \in \mathbb{F}^{m \times m}$. Then,

$$e^{t \begin{bmatrix} A & B \\ 0 & C \end{bmatrix}} = \begin{bmatrix} e^{tA} & \int_0^t e^{(t-\tau)A} B e^{\tau C} d\tau \\ 0 & e^{tC} \end{bmatrix}.$$

Furthermore,

$$\int_0^t e^{\tau A} d\tau = \begin{bmatrix} I & 0 \end{bmatrix} e^{t \begin{bmatrix} A & I \\ 0 & 0 \end{bmatrix}} \begin{bmatrix} 0 \\ I \end{bmatrix}.$$

(Remark: The result can be extended to block- $k \times k$ matrices. See [1359]. For an application to sampled-data control, see [1053].)

Fact 11.14.2. Let $A, B \in \mathbb{F}^{n \times n}$, and consider the following conditions:

- i) $A = B$.
- ii) $e^A = e^B$.
- iii) $AB = BA$.
- iv) $Ae^B = e^B A$.
- v) $e^A e^B = e^B e^A$.
- vi) $e^A e^B = e^{A+B}$.
- vii) $e^A e^B = e^B e^A = e^{A+B}$.

Then, the following statements hold:

- viii) $iii) \implies iv) \implies v)$.
- ix) $iii) \implies vii)$.
- x) If $\text{spec}(A)$ is $2\pi j$ congruence free, then $ii) \implies iii) \implies iv) \iff v)$.
- xi) If $\text{spec}(A)$ and $\text{spec}(B)$ are $2\pi j$ congruence free, then $ii) \implies iii) \iff iv) \iff v)$.
- xii) If $\text{spec}(A + B)$ is $2\pi j$ congruence free, then $iii) \iff vii)$.
- xiii) If, for all $\lambda \in \text{spec}(A)$ and all $\mu \in \text{spec}(B)$, it follows that $(\lambda - \mu)/(2\pi j)$ is not a nonzero integer, then $ii) \implies i)$.
- xiv) If A and B are Hermitian, then $i) \iff ii) \implies iii) \iff iv) \iff v) \iff vi)$.

(Remark: The set $\mathcal{S} \subset \mathbb{C}$ is $2\pi j$ congruence free if no two elements of $\mathcal{S}$ differ by a nonzero integer multiple of $2\pi j$.) (Proof. See [629, pp. 88, 89, 270–272] and [1065, 1169, 1170, 1171, 1208, 1420, 1421]. The assumption of normality in operator versions of some of these statements in [1065, 1171] is not needed in the matrix case. Statement *xiii*) is given in [683, p. 32].) (Remark: The matrices $A \triangleq \begin{bmatrix} 0 & 1 \\ 0 & 2\pi j \end{bmatrix}$ and $B \triangleq \begin{bmatrix} 2\pi j & 0 \\ 0 & -2\pi j \end{bmatrix}$ do not commute but satisfy $e^A = e^B = e^{A+B} = I$. The same

statement holds for

$$A = 2\pi \begin{bmatrix} 0 & 0 & \sqrt{3}/2 \\ 0 & 0 & -1/2 \\ -\sqrt{3}/2 & 1/2 & 0 \end{bmatrix}, \quad B = 2\pi \begin{bmatrix} 0 & 0 & -\sqrt{3}/2 \\ 0 & 0 & -1/2 \\ \sqrt{3}/2 & 1/2 & 0 \end{bmatrix}.$$

Consequently, *vii*) does not imply *iii*.) (Problem: Does *vi*) imply *vii*)? Can *vii*) be replaced by *vi*) in *xii*)?)

Fact 11.14.3. Let $A, B \in \mathbb{R}^{n \times n}$. Then,

$$\frac{d}{dt} e^{A+tB} = \int_0^1 e^{\tau(A+tB)} B e^{(1-\tau)(A+tB)} d\tau.$$

Hence,

$$\text{Dexp}(A; B) = \left. \frac{d}{dt} e^{A+tB} \right|_{t=0} = \int_0^1 e^{\tau A} B e^{(1-\tau)A} d\tau.$$

Furthermore,

$$\frac{d}{dt} \text{tr } e^{A+tB} = \text{tr}(e^{A+tB} B).$$

Hence,

$$\left. \frac{d}{dt} \text{tr } e^{A+tB} \right|_{t=0} = \text{tr}(e^A B).$$

(Proof: See [170, p. 175], [442, p. 371], or [881, 977, 1027].)

Fact 11.14.4. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} \left. \frac{d}{dt} e^{A+tB} \right|_{t=0} &= \left(\frac{e^{\text{ad}_A} - I}{\text{ad}_A} \right) (B) e^A \\ &= e^A \left(\frac{I - e^{-\text{ad}_A}}{\text{ad}_A} \right) (B) \\ &= \sum_{k=0}^{\infty} \frac{1}{(k+1)!} \text{ad}_A^k (B) e^A. \end{aligned}$$

(Proof: The second and fourth expressions are given in [103, p. 49] and [746, p. 248], while the third expression appears in [1347]. See also [1366, pp. 107–110].)

(Remark: See Fact 2.18.6.)

Fact 11.14.5. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $e^A = e^B$. Then, the following statements hold:

- i*) If $|\lambda| < \pi$ for all $\lambda \in \text{spec}(A) \cup \text{spec}(B)$, then $A = B$.
- ii*) If $\lambda - \mu \neq 2k\pi j$ for all $\lambda \in \text{spec}(A)$, $\mu \in \text{spec}(B)$, and $k \in \mathbb{Z}$, then $[A, B] = 0$.
- iii*) If A is normal and $\sigma_{\max}(A) < \pi$, then $[A, B] = 0$.
- iv*) If A is normal and $\sigma_{\max}(A) = \pi$, then $[A^2, B] = 0$.

(Proof: See [1173, 1208] and [1366, p. 111].) (Remark: If $[A, B] = 0$, then $[A^2, B] = 0$.)

Fact 11.14.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are skew Hermitian. Then, $e^{tA}e^{tB}$ is unitary, and there exists a skew-Hermitian matrix $C(t)$ such that $e^{tA}e^{tB} = e^{C(t)}$. (Problem: Does (11.4.1) converge in this case? See [227, 458, 1123].)

Fact 11.14.7. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then,

$$\lim_{p \rightarrow 0} \left(e^{\frac{p}{2}A} e^{pB} e^{\frac{p}{2}A} \right)^{1/p} = e^{A+B}.$$

(Proof: See [53].) (Remark: This result is related to the Lie-Trotter formula given by Corollary 11.4.8. For extensions, see [9, 533].)

Fact 11.14.8. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then,

$$\lim_{p \rightarrow \infty} \left[\frac{1}{2}(e^{pA} + e^{pB}) \right]^{1/p} = e^{\frac{1}{2}(A+B)}.$$

(Proof: See [193].)

Fact 11.14.9. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\lim_{k \rightarrow \infty} \left[e^{\frac{1}{k}A} e^{\frac{1}{k}B} e^{-\frac{1}{k}A} e^{-\frac{1}{k}B} \right]^{k^2} = e^{[A,B]}.$$

Fact 11.14.10. Let $A \in \mathbb{F}^{n \times m}$, $X \in \mathbb{F}^{m \times l}$, and $B \in \mathbb{F}^{l \times n}$. Then,

$$\frac{d}{dX} \operatorname{tr} e^{AXB} = B e^{AXB} A.$$

Fact 11.14.11. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\left. \frac{d}{dt} e^{tA} e^{tB} e^{-tA} e^{-tB} \right|_{t=0} = 0$$

and

$$\left. \frac{d}{dt} e^{\sqrt{t}A} e^{\sqrt{t}B} e^{-\sqrt{t}A} e^{-\sqrt{t}B} \right|_{t=0} = AB - BA.$$

Fact 11.14.12. Let $A, B, C \in \mathbb{F}^{n \times n}$, assume there exists $\beta \in \mathbb{F}$ such that $[A, B] = \beta B + C$, and assume that $[A, C] = [B, C] = 0$. Then,

$$e^{A+B} = e^A e^{\phi(\beta)B} e^{\psi(\beta)C},$$

where

$$\phi(\beta) \triangleq \begin{cases} \frac{1}{\beta}(1 - e^{-\beta}), & \beta \neq 0, \\ 1, & \beta = 0, \end{cases}$$

and

$$\psi(\beta) \triangleq \begin{cases} \frac{1}{\beta^2}(1 - \beta - e^{-\beta}), & \beta \neq 0, \\ -\frac{1}{2}, & \beta = 0. \end{cases}$$

(Proof: See [556, 1264].)

Fact 11.14.13. Let $A, B \in \mathbb{F}^{n \times n}$, and assume there exist $\alpha, \beta \in \mathbb{F}$ such that $[A, B] = \alpha A + \beta B$. Then,

$$e^{t(A+B)} = e^{\phi(t)A} e^{\psi(t)B},$$

where

$$\phi(t) \triangleq \begin{cases} t, & \alpha = \beta = 0, \\ \alpha^{-1} \log(1 + \alpha t), & \alpha = \beta \neq 0, 1 + \alpha t > 0, \\ \int_0^t \frac{\alpha - \beta}{\alpha e^{(\alpha - \beta)\tau} - \beta} d\tau, & \alpha \neq \beta, \end{cases}$$

and

$$\psi(t) \triangleq \int_0^t e^{-\beta \phi(\tau)} d\tau.$$

(Proof: See [1265].)

Fact 11.14.14. Let $A, B \in \mathbb{F}^{n \times n}$, and assume there exists nonzero $\beta \in \mathbb{F}$ such that $[A, B] = \alpha B$. Then, for all $t > 0$,

$$e^{t(A+B)} = e^{tA} e^{[(1-e^{-\alpha t})/\alpha]B}.$$

(Proof: Apply Fact 11.14.12 with $[tA, tB] = \alpha t(tB)$ and $\beta = \alpha t$.)

Fact 11.14.15. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $[[A, B], A] = 0$ and $[[A, B], B] = 0$. Then, for all $t \in \mathbb{R}$,

$$e^{tA} e^{tB} = e^{tA + tB + (t^2/2)[A, B]}.$$

In particular,

$$e^A e^B = e^{A+B + \frac{1}{2}[A, B]} = e^{A+B} e^{\frac{1}{2}[A, B]} = e^{\frac{1}{2}[A, B]} e^{A+B}$$

and

$$e^B e^{2A} e^B = e^{2A+2B}.$$

(Proof: See [624, pp. 64–66] and [1431].)

Fact 11.14.16. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that $[A, B] = B^2$. Then,

$$e^{A+B} = e^A (I + B).$$

Fact 11.14.17. Let $A, B \in \mathbb{F}^{n \times n}$. Then, for all $t \in [0, \infty)$,

$$e^{t(A+B)} = e^{tA} e^{tB} + \sum_{k=2}^{\infty} C_k t^k,$$

where, for all $k \in \mathbb{N}$,

$$C_{k+1} \triangleq \frac{1}{k+1} ([A+B]C_k + [B, D_k]), \quad C_0 \triangleq 0,$$

and

$$D_{k+1} \triangleq \frac{1}{k+1} (AD_k + D_k B), \quad D_0 \triangleq I.$$

(Proof: See [1125].)

Fact 11.14.18. Let $A, B \in \mathbb{F}^{n \times n}$. Then, for all $t \in [0, \infty)$,

$$e^{t(A+B)} = e^{tA} e^{tB} e^{tC_2} e^{tC_3} \dots,$$

where

$$C_2 \triangleq -\frac{1}{2}[A, B], \quad C_3 \triangleq \frac{1}{3}[B, [A, B]] + \frac{1}{6}[A, [A, B]].$$

(Remark: This result is the *Zassenhaus product formula*. See [683, p. 236] and [1176].) (Remark: Higher order terms are given in [1176].) (Remark: Conditions for convergence do not seem to be available.)

Fact 11.14.19. Let $A \in \mathbb{R}^{2n \times 2n}$, and assume that A is symplectic and discrete-time Lyapunov stable. Then, $\text{spec}(A) \subset \{s \in \mathbb{C} : |s| = 1\}$, $\text{am}_A(1)$ and $\text{am}_A(-1)$ are even, A is semisimple, and there exists a Hamiltonian matrix $B \in \mathbb{R}^{2n \times 2n}$ such that $A = e^B$. (Proof: Since A is symplectic and discrete-time Lyapunov stable, it follows that the spectrum of A is a subset of the unit circle and A is semisimple. Therefore, the only negative eigenvalue that A can have is -1 . Since all nonreal eigenvalues appear in complex conjugate pairs and A has even order, and since, by Fact 3.19.10, $\det A = 1$, it follows that the eigenvalues -1 and 1 (if present) have even algebraic multiplicity. The fact that A has a Hamiltonian logarithm now follows from Theorem 2.6 of [404].) (Remark: See *xiii*) of Proposition 11.6.5.)

Fact 11.14.20. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A is positive definite, and assume that B is positive semidefinite. Then,

$$A + B \leq A^{1/2} e^{A^{-1/2} B A^{-1/2}} A^{1/2}.$$

Hence,

$$\frac{\det(A+B)}{\det A} \leq e^{\text{tr } A^{-1} B}.$$

Furthermore, for each inequality, equality holds if and only if $B = 0$. (Proof: For positive-semidefinite A it follows that $e^A \leq I + A$.)

Fact 11.14.21. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then,

$$I \circ (A + B) \leq \log(e^A \circ e^B).$$

(Proof: See [43, 1485].) (Remark: See Fact 8.21.48.)

Fact 11.14.22. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, assume that $A \leq B$, let $\alpha, \beta \in \mathbb{R}$, assume that either $\alpha I \leq A \leq \beta I$ or $\alpha I \leq B \leq \beta I$, and let $t > 0$. Then,

$$e^{tA} \leq S(t, e^{\beta-\alpha}) e^{tB},$$

where, for $t > 0$ and $h > 0$,

$$S(t, h) \triangleq \begin{cases} \frac{(h^t - 1)h^{t/(h^t-1)}}{et \log h}, & h \neq 1, \\ 1, & h = 1. \end{cases}$$

(Proof: See [518].) (Remark: $S(t, h)$ is Specht's ratio. See Fact 1.10.22 and Fact 1.15.19.)

Fact 11.14.23. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, let $\alpha, \beta \in \mathbb{R}$, assume that $\alpha I \leq A \leq \beta I$ and $\alpha I \leq B \leq \beta I$, and let $t > 0$. Then,

$$\begin{aligned} \frac{1}{S(1, e^{\beta-\alpha})S^{1/t}(t, e^{\beta-\alpha})} [\alpha e^{tA} + (1-\alpha)e^{tB}]^{1/t} \\ \leq e^{\alpha A + (1-\alpha)B} \\ \leq S(1, e^{\beta-\alpha}) [\alpha e^{tA} + (1-\alpha)e^{tB}]^{1/t}, \end{aligned}$$

where $S(t, h)$ is defined in Fact 11.14.22. (Proof: See [518].)

Fact 11.14.24. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$\log \det A = \operatorname{tr} \log A$$

and

$$\log \det AB = \operatorname{tr}(\log A + \log B).$$

Fact 11.14.25. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive definite. Then,

$$\operatorname{tr}(A - B) \leq \operatorname{tr}[A(\log A - \log B)]$$

and

$$(\log \operatorname{tr} A - \log \operatorname{tr} B) \operatorname{tr} A \leq \operatorname{tr}[A(\log A - \log B)].$$

(Proof: See [159] and [197, p. 281].) (Remark: The first inequality is *Klein's inequality*. See [201, p. 118].) (Remark: The second inequality is equivalent to the thermodynamic inequality. See Fact 11.14.31.) (Remark: $\operatorname{tr}[A(\log A - \log B)]$ is the *relative entropy of Umegaki*.)

Fact 11.14.26. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and define

$$\mu(A, B) \triangleq e^{\frac{1}{2}(\log A + \log B)}.$$

Then, the following statements hold:

- i) $\mu(A, A^{-1}) = I$.
- ii) $\mu(A, B) = \mu(B, A)$.
- iii) If $AB = BA$, then $\mu(A, B) = AB$.

(Proof: See [74].) (Remark: With multiplication defined by μ , the set of $n \times n$ positive-definite matrices is a commutative Lie group. See [74].)

Fact 11.14.27. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive definite, and let $p > 0$. Then,

$$\frac{1}{p} \operatorname{tr}[A \log(B^{p/2} A^p B^{p/2})] \leq \operatorname{tr}[A(\log A + \log B)] \leq \frac{1}{p} \operatorname{tr}[A \log(A^p B^p A^p)].$$

Furthermore,

$$\lim_{p \downarrow 0} \frac{1}{p} \operatorname{tr}[A \log(B^{p/2} A^p B^{p/2})] = \operatorname{tr}[A(\log A + \log B)] = \lim_{p \downarrow 0} \frac{1}{p} \operatorname{tr}[A \log(A^p B^p A^p)].$$

(Proof: See [53, 160, 533, 674].) (Remark: This inequality has applications to quantum information theory.)

Fact 11.14.28. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, let $q \geq p > 0$, let $h \triangleq \lambda_{\max}(e^A)/\lambda_{\min}(e^B)$, and define

$$S(1, h) \triangleq \frac{(h-1)h^{1/(h-1)}}{e \log h}.$$

Then, there exist unitary matrices $U, V \in \mathbb{F}^{n \times n}$ such that

$$\frac{1}{S(1, h)} U e^{A+B} U^* \leq e^{\frac{1}{2}A} e^B e^{\frac{1}{2}A} \leq S(1, h) V e^{A+B} V^*.$$

Furthermore,

$$\operatorname{tr} e^{A+B} \leq \operatorname{tr} e^A e^B \leq S(1, h) \operatorname{tr} e^{A+B},$$

$$\operatorname{tr} (e^{pA} \# e^{pB})^{2/p} \leq \operatorname{tr} e^{A+B} \leq \operatorname{tr} (e^{\frac{p}{2}B} e^{pA} e^{\frac{p}{2}B})^{1/p} \leq \operatorname{tr} (e^{\frac{q}{2}B} e^{qA} e^{\frac{q}{2}B})^{1/q},$$

$$\operatorname{tr} e^{A+B} = \lim_{p \downarrow 0} \operatorname{tr} (e^{\frac{p}{2}B} e^{pA} e^{\frac{p}{2}B})^{1/p},$$

$$e^{A+B} = \lim_{p \downarrow 0} (e^{pA} \# e^{pB})^{2/p}.$$

Moreover, $\operatorname{tr} e^{A+B} = \operatorname{tr} e^A e^B$ if and only if $AB = BA$. Furthermore, for all $i = 1, \dots, n$,

$$\frac{1}{S(1, h)} \lambda_i(e^{A+B}) \leq \lambda_i(e^A e^B) \leq S(1, h) \lambda_i(e^{A+B}).$$

Finally, let $\alpha \in [0, 1]$. Then,

$$\lim_{p \downarrow 0} (e^{pA} \#_{\alpha} e^{pB})^{1/p} = e^{(1-\alpha)A + \alpha B}$$

and

$$\operatorname{tr} (e^{pA} \#_{\alpha} e^{pB})^{1/p} \leq \operatorname{tr} e^{(1-\alpha)A + \alpha B}.$$

(Proof: See [252].) (Remark: The left-hand inequality in the second string of inequalities is the *Golden-Thompson inequality*. See Fact 11.16.4.) (Remark: Since $S(1, h) > 1$ for all $h > 1$, the left-hand inequality in the first string of inequalities does not imply the Golden-Thompson inequality.) (Remark: For $i = 1$, the stronger eigenvalue inequality $\lambda_{\max}(e^{A+B}) \leq \lambda_{\max}(e^A e^B)$ holds. See Fact 11.16.4.) (Remark: $S(1, h)$ is Specht's ratio given by Fact 11.14.22.) (Remark: The generalized geometric mean is defined in Fact 8.10.45.)

Fact 11.14.29. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then,

$$(\operatorname{tr} e^A) e^{\operatorname{tr}(e^A B) / \operatorname{tr} e^A} \leq \operatorname{tr} e^{A+B}.$$

(Proof: See [159].) (Remark: This result is the *Peierls-Bogoliubov inequality*.) (Remark: This inequality is equivalent to the thermodynamic inequality. See Fact 11.14.31.)

Fact 11.14.30. Let $A, B, C \in \mathbb{F}^{n \times n}$, and assume that A, B , and C are positive definite. Then,

$$\operatorname{tr} e^{\log A - \log B + \log C} \leq \operatorname{tr} \int_0^{\infty} A(B + xI)^{-1} C(B + xI)^{-1} dx.$$

(Proof: See [905, 933].) (Remark: $-\log B$ is correct.) (Remark: $\operatorname{tr} e^{A+B+C} \leq |\operatorname{tr} e^A e^B e^C|$ is not necessarily true.)

Fact 11.14.31. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A is positive definite, $\operatorname{tr} A = 1$, and B is Hermitian. Then,

$$\operatorname{tr} AB \leq \operatorname{tr}(A \log A) + \log \operatorname{tr} e^B.$$

Furthermore, equality holds if and only if

$$A = (\operatorname{tr} e^B)^{-1} e^B.$$

(Proof: See [159].) (Remark: This result is the *thermodynamic inequality*. Equivalent forms are given by Fact 11.14.25 and Fact 11.14.29.)

Fact 11.14.32. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then,

$$\|A - B\|_F \leq \|\log(e^{-\frac{1}{2}A} e^B e^{\frac{1}{2}A})\|_F.$$

(Proof: See [201, p. 203].) (Remark: This result has a distance interpretation in terms of geodesics. See [201, p. 203] and [207, 1013, 1014].)

Fact 11.14.33. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are skew Hermitian. Then, there exist unitary matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that

$$e^A e^B = e^{S_1 A S_1^{-1} + S_2 B S_2^{-1}}.$$

(Proof: See [1210, 1272, 1273].)

Fact 11.14.34. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are Hermitian. Then, there exist unitary matrices $S_1, S_2 \in \mathbb{F}^{n \times n}$ such that

$$e^{\frac{1}{2}A} e^B e^{\frac{1}{2}A} = e^{S_1 A S_1^{-1} + S_2 B S_2^{-1}}.$$

(Proof: See [1209, 1210, 1272, 1273].) (Problem: Determine the relationship between this result and Fact 11.14.33.)

Fact 11.14.35. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and assume that $B \leq A$. Furthermore, let $p, q, r, t \in \mathbb{R}$, and assume that $r \geq t \geq 0$, $p \geq 0$, $p + q \geq 0$, and $p + q + r > 0$. Then,

$$\left[e^{\frac{r}{2}A} e^{qA+pB} e^{\frac{r}{2}A} \right]^{t/(p+q+r)} \leq e^{tA}.$$

(Proof: See [1350].)

Fact 11.14.36. Let $A \in \mathbb{F}^{n \times n}$ and $B \in \mathbb{F}^{m \times m}$. Then,

$$\operatorname{tr} e^{A \oplus B} = (\operatorname{tr} e^A)(\operatorname{tr} e^B).$$

Fact 11.14.37. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, and $C \in \mathbb{F}^{l \times l}$. Then,

$$e^{A \oplus B \oplus C} = e^A \otimes e^B \otimes e^C.$$

Fact 11.14.38. Let $A \in \mathbb{F}^{n \times n}$, $B \in \mathbb{F}^{m \times m}$, $C \in \mathbb{F}^{k \times k}$, and $D \in \mathbb{F}^{l \times l}$. Then,

$$\operatorname{tr} e^{A \otimes I \otimes B \otimes I + I \otimes C \otimes I \otimes D} = \operatorname{tr} e^{A \otimes B} \operatorname{tr} e^{C \otimes D}.$$

(Proof: By Fact 7.4.29, a similarity transformation involving the Kronecker permutation matrix can be used to reorder the inner two terms. See [1220].)

Fact 11.14.39. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that A and B are positive definite. Then, $A \# B$ is the unique positive-definite solution X of the matrix equation

$$\log(A^{-1}X) + \log(B^{-1}X) = 0.$$

(Proof: See [1014].)

11.15 Facts on the Matrix Exponential and Eigenvalues, Singular Values, and Norms for One Matrix

Fact 11.15.1. Let $A \in \mathbb{F}^{n \times n}$, assume that e^A is positive definite, and assume that $\sigma_{\max}(A) < 2\pi$. Then, A is Hermitian. (Proof: See [851, 1172].)

Fact 11.15.2. Let $A \in \mathbb{F}^{n \times n}$, and define $f: [0, \infty) \mapsto (0, \infty)$ by $f(t) \triangleq \sigma_{\max}(e^{At})$. Then,

$$f'(0) = \frac{1}{2}\lambda_{\max}(A + A^*).$$

Hence, there exists $\varepsilon > 0$ such that $f(t) \triangleq \sigma_{\max}(e^{tA})$ is decreasing on $[0, \varepsilon)$ if and only if A is dissipative. (Proof: The result follows from *iii*) of Fact 11.15.7. See [1402].) (Remark: The derivative is one sided.)

Fact 11.15.3. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $t \geq 0$,

$$\frac{d}{dt} \|e^{tA}\|_F^2 = \operatorname{tr} e^{tA}(A + A^*)e^{tA*}.$$

Hence, if A is dissipative, then $f(t) \triangleq \|e^{tA}\|_F$ is decreasing on $[0, \infty)$. (Proof: See [1402].)

Fact 11.15.4. Let $A \in \mathbb{F}^{n \times n}$. Then,

$$|\operatorname{tr} e^{2A}| \leq \operatorname{tr} e^A e^{A*} \leq \operatorname{tr} e^{A+A*} \leq \left[n \operatorname{tr} e^{2(A+A*)} \right]^{1/2} \leq \frac{n}{2} + \frac{1}{2} \operatorname{tr} e^{2(A+A*)}.$$

In addition, $\operatorname{tr} e^A e^{A*} = \operatorname{tr} e^{A+A*}$ if and only if A is normal. (Proof: See [184], [711, p. 515], and [1208].) (Remark: $\operatorname{tr} e^A e^{A*} \leq \operatorname{tr} e^{A+A*}$ is *Bernstein's inequality*. See [47].) (Remark: See Fact 3.7.12.)

Fact 11.15.5. Let $A \in \mathbb{F}^{n \times n}$. Then, for all $k = 1, \dots, n$,

$$\prod_{i=1}^k \sigma_i(e^A) \leq \prod_{i=1}^k \lambda_i \left[e^{\frac{1}{2}(A+A*)} \right] = \prod_{i=1}^k e^{\lambda_i[\frac{1}{2}(A+A*)]} \leq \prod_{i=1}^k e^{\sigma_i(A)}.$$

Furthermore, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \sigma_i(e^A) \leq \sum_{i=1}^k \lambda_i \left[e^{\frac{1}{2}(A+A*)} \right] = \sum_{i=1}^k e^{\lambda_i[\frac{1}{2}(A+A*)]} \leq \sum_{i=1}^k e^{\sigma_i(A)}.$$

In particular,

$$\sigma_{\max}(e^A) \leq \lambda_{\max}\left[e^{\frac{1}{2}(A+A^*)}\right] = e^{\frac{1}{2}\lambda_{\max}(A+A^*)} \leq e^{\sigma_{\max}(A)}$$

or, equivalently,

$$\lambda_{\max}(e^A e^{A^*}) \leq \lambda_{\max}(e^{A+A^*}) = e^{\lambda_{\max}(A+A^*)} \leq e^{2\sigma_{\max}(A)}.$$

Furthermore,

$$|\det e^A| = |e^{\operatorname{tr} A}| \leq e^{|\operatorname{tr} A|} \leq e^{\operatorname{tr} \langle A \rangle}$$

and

$$\operatorname{tr} \langle e^A \rangle \leq \sum_{i=1}^n e^{\sigma_i(A)}.$$

(Proof: See [1211], Fact 2.21.13, Fact 8.17.4, and Fact 8.17.5.)

Fact 11.15.6. Let $A \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|e^A e^{A^*}\| \leq \|e^{A+A^*}\|.$$

In particular,

$$\lambda_{\max}(e^A e^{A^*}) \leq \lambda_{\max}(e^{A+A^*})$$

and

$$\operatorname{tr} e^A e^{A^*} \leq \operatorname{tr} e^{A+A^*}.$$

(Proof: See [342].)

Fact 11.15.7. Let $A, B \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ be the norm on $\mathbb{F}^{n \times n}$ induced by the norm $\|\cdot\|'$ on $\mathbb{F}^n$, let $\operatorname{mspec}(A) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, and define

$$\mu(A) \triangleq \lim_{\varepsilon \downarrow 0} \frac{\|I + \varepsilon A\| - 1}{\varepsilon}.$$

Then, the following statements hold:

- i) $\mu(A) = D_+(f; I)$, where $f: \mathbb{F}^{n \times n} \mapsto \mathbb{R}$ is defined by $f(A) \triangleq \|A\|$.
- ii) $\mu(A) = \lim_{t \downarrow 0} t^{-1} \log \|e^{tA}\| = \sup_{t > 0} t^{-1} \log \|e^{tA}\|$.
- iii) $\mu(A) = \left. \frac{d^+}{dt} \|e^{tA}\| \right|_{t=0} = \left. \frac{d^+}{dt} \log \|e^{tA}\| \right|_{t=0}$.
- iv) $\mu(I) = 1$, $\mu(-I) = -1$, and $\mu(0) = 0$.
- v) $\operatorname{spabs}(A) = \lim_{t \rightarrow \infty} t^{-1} \log \|e^{tA}\| = \inf_{t > 0} t^{-1} \log \|e^{tA}\|$.
- vi) For all $i = 1, \dots, n$,

$$-\|A\| \leq -\mu(-A) \leq \operatorname{Re} \lambda_i \leq \operatorname{spabs}(A) \leq \mu(A) \leq \|A\|.$$

vii) For all $\alpha \in \mathbb{R}$, $\mu(\alpha A) = |\alpha| \mu[(\operatorname{sign} \alpha)A]$.

viii) For all $\alpha \in \mathbb{F}$, $\mu(A + \alpha I) = \mu(A) + \operatorname{Re} \alpha$.

ix) $\max\{\mu(A) - \mu(-B), -\mu(-A) + \mu(B)\} \leq \mu(A + B) \leq \mu(A) + \mu(B)$.

x) $\mu: \mathbb{F}^{n \times n} \mapsto \mathbb{R}$ is convex.

xi) $|\mu(A) - \mu(B)| \leq \max\{|\mu(A - B)|, |\mu(B - A)|\} \leq \|A - B\|$.

xii) For all $x \in \mathbb{F}^n$, $\max\{-\mu(-A), -\mu(A)\}\|x\|' \leq \|Ax\|'$.

xiii) If A is nonsingular, then $\max\{-\mu(-A), -\mu(A)\} \leq 1/\|A^{-1}\|$.

xiv) For all $t \geq 0$ and all $i = 1, \dots, n$,

$$e^{-\|A\|t} \leq e^{-\mu(-A)t} \leq e^{(\operatorname{Re} \lambda_i)t} \leq e^{\operatorname{spabs}(A)t} \leq \|e^{tA}\| \leq e^{\mu(A)t} \leq e^{\|A\|t}.$$

xv) $\mu(A) = \min\{\beta \in \mathbb{R} : \|e^{tA}\| \leq e^{\beta t} \text{ for all } t \geq 0\}$.

xvi) If $\|\cdot\|' = \|\cdot\|_1$, and thus $\|\cdot\| = \|\cdot\|_{\text{col}}$, then

$$\mu(A) = \max_{j \in \{1, \dots, n\}} \left(\operatorname{Re} A_{(j,j)} + \sum_{\substack{i=1 \\ i \neq j}}^n |A_{(i,j)}| \right).$$

xvii) If $\|\cdot\|' = \|\cdot\|_2$ and thus $\|\cdot\| = \sigma_{\max}(\cdot)$, then

$$\mu(A) = \lambda_{\max}[\tfrac{1}{2}(A + A^*)].$$

xviii) If $\|\cdot\|' = \|\cdot\|_{\infty}$, and thus $\|\cdot\| = \|\cdot\|_{\text{row}}$, then

$$\mu(A) = \max_{i \in \{1, \dots, n\}} \left(\operatorname{Re} A_{(i,i)} + \sum_{\substack{j=1 \\ j \neq i}}^n |A_{(i,j)}| \right).$$

(Proof: See [399, 402, 1067, 1245], [690, pp. 653–655], and [1316, p. 150].) (Remark: $\mu(\cdot)$ is the *matrix measure* or *logarithmic derivative* or *initial growth rate*. For applications, see [690] and [1380]. See Fact 11.18.11 for the logarithmic derivative of an asymptotically stable matrix.) (Remark: The directional differential $D_+f(A; I)$ is defined in (10.4.2).) (Remark: *vi*) and *xvii*) yield Fact 5.11.24.) (Remark: Higher order logarithmic derivatives are studied in [205].)

Fact 11.15.8. Let $A \in \mathbb{F}^{n \times n}$, let $\beta > \operatorname{spabs}(A)$, let $\gamma \geq 1$, and let $\|\cdot\|$ be a normalized, submultiplicative norm on $\mathbb{F}^{n \times n}$. Then, for all $t \geq 0$,

$$\|e^{tA}\| \leq \gamma e^{\beta t}$$

if and only if, for all $k \geq 1$ and $\alpha > \beta$,

$$\|(\alpha I - A)^{-k}\| \leq \frac{\gamma}{(\alpha - \beta)^k}.$$

(Remark: This result is a consequence of the *Hille-Yosida theorem*. See [361, pp. 26] and [690, p. 672].)

Fact 11.15.9. Let $A \in \mathbb{R}^{n \times n}$, let $\beta \in \mathbb{R}$, and assume there exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that

$$A^T P + PA \leq 2\beta P.$$

Then, for all $t \geq 0$,

$$\sigma_{\max}(e^{tA}) \leq \sqrt{\sigma_{\max}(P)/\sigma_{\min}(P)} e^{\beta t}.$$

(Remark: See [690, p. 665].) (Remark: See Fact 11.18.9.)

Fact 11.15.10. Let $A \in \text{SO}(3)$. Then,

$$\theta \triangleq 2 \cos^{-1} \left(\frac{1}{2} \sqrt{1 + \text{tr } A} \right).$$

Then,

$$\theta = \sigma_{\max}(\log A) = \frac{1}{\sqrt{2}} \|\log A\|_F.$$

(Remark: See Fact 3.11.10 and Fact 11.11.10.) (Remark: θ is a Riemannian metric giving the length of the shortest geodesic curve on $\text{SO}(3)$ between A and I . See [1013].)

11.16 Facts on the Matrix Exponential and Eigenvalues, Singular Values, and Norms for Two or More Matrices

Fact 11.16.1. Let $A, B \in \mathbb{F}^{n \times n}$. Then,

$$\begin{aligned} |\text{tr } e^{A+B}| &\leq \text{tr } e^{\frac{1}{2}(A+B)} e^{\frac{1}{2}(A+B)^*} \\ &\leq \text{tr } e^{\frac{1}{2}(A+A^*+B+B^*)} \\ &\leq \text{tr } e^{\frac{1}{2}(A+A^*)} e^{\frac{1}{2}(B+B^*)} \\ &\leq \left(\text{tr } e^{A+A^*} \right)^{1/2} \left(\text{tr } e^{B+B^*} \right)^{1/2} \\ &\leq \frac{1}{2} \text{tr} \left(e^{A+A^*} + e^{B+B^*} \right) \end{aligned}$$

and

$$\left\{ \text{tr } e^A e^B \right\} \leq \frac{1}{2} \text{tr} \left(e^A e^{A^*} + e^B e^{B^*} \right) \leq \frac{1}{2} \text{tr} \left(e^{A+A^*} + e^{B+B^*} \right).$$

(Proof: See [184, 343, 1075] and [711, p. 514].)

Fact 11.16.2. Let $A, B \in \mathbb{F}^{n \times n}$. Then, for all $p > 0$,

$$\sigma_{\max} \left[e^{A+B} - \left(e^{\frac{1}{p}A} e^{\frac{1}{p}B} \right)^p \right] \leq \frac{1}{2p} \sigma_{\max}([A, B]) e^{\sigma_{\max}(A) + \sigma_{\max}(B)}.$$

(Proof: See [683, p. 237] and [1015].) (Remark: See Corollary 10.8.8 and Fact 11.16.3.)

Fact 11.16.3. Let $A \in \mathbb{F}^{n \times n}$, and define $A_H \triangleq \frac{1}{2}(A+A^*)$ and $A_S \triangleq \frac{1}{2}(A-A^*)$. Then, for all $p > 0$,

$$\sigma_{\max} \left[e^A - \left(e^{\frac{1}{p}A_H} e^{\frac{1}{p}A_S} \right)^p \right] \leq \frac{1}{4p} \sigma_{\max}([A^*, A]) e^{\frac{1}{2} \lambda_{\max}(A+A^*)}.$$

(Proof: See [1015].) (Remark: See Fact 10.8.8.)

Fact 11.16.4. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|e^{A+B}\| \leq \left\| e^{\frac{1}{2}A} e^{B} e^{\frac{1}{2}A} \right\| \leq \|e^A e^B\|.$$

If, in addition, $p > 0$, then

$$\|e^{A+B}\| \leq \left\| e^{\frac{p}{2}A} e^B e^{\frac{p}{2}A} \right\|^{1/p}$$

and

$$\|e^{A+B}\| = \lim_{p \downarrow 0} \left\| e^{\frac{p}{2}A} e^B e^{\frac{p}{2}A} \right\|^{1/p}.$$

Furthermore, for all $k = 1, \dots, n$,

$$\prod_{i=1}^k \lambda_i(e^{A+B}) \leq \prod_{i=1}^k \lambda_i(e^A e^B) \leq \prod_{i=1}^k \sigma_i(e^A e^B)$$

with equality for $k = n$, that is,

$$\prod_{i=1}^n \lambda_i(e^{A+B}) = \prod_{i=1}^n \lambda_i(e^A e^B) = \prod_{i=1}^n \sigma_i(e^A e^B) = \det(e^A e^B).$$

In fact,

$$\begin{aligned} \det(e^{A+B}) &= \prod_{i=1}^n \lambda_i(e^{A+B}) \\ &= \prod_{i=1}^n e^{\lambda_i(A+B)} \\ &= e^{\operatorname{tr}(A+B)} \\ &= e^{(\operatorname{tr} A) + (\operatorname{tr} B)} \\ &= e^{\operatorname{tr} A} e^{\operatorname{tr} B} \\ &= \det(e^A) \det(e^B) \\ &= \det(e^A e^B) \\ &= \prod_{i=1}^n \sigma_i(e^A e^B). \end{aligned}$$

Furthermore, for all $k = 1, \dots, n$,

$$\sum_{i=1}^k \lambda_i(e^{A+B}) \leq \sum_{i=1}^k \lambda_i(e^A e^B) \leq \sum_{i=1}^k \sigma_i(e^A e^B).$$

In particular,

$$\lambda_{\max}(e^{A+B}) \leq \lambda_{\max}(e^A e^B) \leq \sigma_{\max}(e^A e^B),$$

$$\operatorname{tr} e^{A+B} \leq \operatorname{tr} e^A e^B \leq \operatorname{tr} \langle e^A e^B \rangle,$$

and, for all $p > 0$,

$$\operatorname{tr} e^{A+B} \leq \operatorname{tr} (e^{\frac{p}{2}A} e^B e^{\frac{p}{2}A}).$$

Finally, $\operatorname{tr} e^{A+B} = \operatorname{tr} e^A e^B$ if and only if A and B commute. (Proof: See [53], [197, p. 261], Fact 5.11.28, Fact 2.21.13, and Fact 9.11.2. For the last statement, see [1208].) (Remark: Note that $\det(e^{A+B}) = \det(e^A) \det(e^B)$ even though e^{A+B} and $e^A e^B$ may not be equal. See [683, p. 265] or [711, p. 442].) (Remark: $\operatorname{tr} e^{A+B} \leq \operatorname{tr} e^A e^B$ is the Golden-Thompson inequality. See Fact 11.14.28.) (Remark: $\|e^{A+B}\| \leq$

$\|e^{\frac{1}{2}A}e^Be^{\frac{1}{2}A}\|$ is *Segal's inequality*. See [47].) (Problem: Compare the upper bound $\text{tr} \langle e^A e^B \rangle$ for $\text{tr} e^A e^B$ with the upper bound $S(1, h) \text{tr} e^{A+B}$ given by Fact 11.14.28.)

Fact 11.16.5. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, let $q, p > 0$, where $q \leq p$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\left\| \left(e^{\frac{q}{2}A} e^{qB} e^{\frac{q}{2}A} \right)^{1/q} \right\| \leq \left\| \left(e^{\frac{p}{2}A} e^{pB} e^{\frac{p}{2}A} \right)^{1/p} \right\|.$$

(Proof: See [53].)

Fact 11.16.6. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then,

$$e^{\sigma_{\max}^{1/2}(AB)} - 1 \leq \sigma_{\max}^{1/2}[(e^A - I)(e^B - I)]$$

and

$$e^{\sigma_{\max}^{1/3}(BAB)} - 1 \leq \sigma_{\max}^{1/3}[(e^B - I)(e^A - I)(e^B - I)].$$

(Proof: See [1349].) (Remark: See Fact 8.18.30.)

Fact 11.16.7. Let $A, B \in \mathbb{F}^{n \times n}$, and let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$. Then, for all $t \geq 0$,

$$\|e^{tA} - e^{tB}\| \leq e^{\|A\|t} (e^{\|A-B\|t} - 1).$$

Fact 11.16.8. Let $A, B \in \mathbb{F}^{n \times n}$, and let $t \geq 0$. Then,

$$e^{t(A+B)} = e^{tA} + \int_0^t e^{(t-\tau)A} B e^{\tau(A+B)} d\tau.$$

(Proof: See [683, p. 238].)

Fact 11.16.9. Let $A, B \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ be a normalized submultiplicative norm on $\mathbb{F}^{n \times n}$, and let $t \geq 0$. Then,

$$\|e^{tA} - e^{tB}\| \leq t\|A - B\|e^{t \max\{\|A\|, \|B\|\}}.$$

(Proof: See [683, p. 265].)

Fact 11.16.10. Let $A, B \in \mathbb{R}^{n \times n}$, and assume that A is normal. Then, for all $t \geq 0$,

$$\sigma_{\max}(e^{tA} - e^{tB}) \leq \sigma_{\max}(e^{tA}) [e^{\sigma_{\max}(A-B)t} - 1].$$

(Proof: See [1420].)

Fact 11.16.11. Let $A \in \mathbb{F}^{n \times n}$, let $\|\cdot\|$ be an induced norm on $\mathbb{F}^{n \times n}$, and let $\alpha > 0$ and $\beta \in \mathbb{R}$ be such that, for all $t \geq 0$,

$$\|e^{tA}\| \leq \alpha e^{\beta t}.$$

Then, for all $B \in \mathbb{F}^{n \times n}$ and $t \geq 0$,

$$\|e^{t(A+B)}\| \leq \alpha e^{(\beta+\alpha\|B\|)t}.$$

(Proof: See [690, p. 406].)

Fact 11.16.12. Let $A, B \in \mathbb{C}^{n \times n}$, assume that A and B are idempotent, assume that $A \neq B$, and let $\|\cdot\|$ be a norm on $\mathbb{C}^{n \times n}$. Then,

$$\|e^{JA} - e^{JB}\| = |e^J - 1| \|A - B\| < \|A - B\|.$$

(Proof: See [1028].) (Remark: $|e^J - 1| \approx 0.96$.)

Fact 11.16.13. Let $A, B \in \mathbb{C}^{n \times n}$, assume that A and B are Hermitian, let $X \in \mathbb{C}^{n \times n}$, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{C}^{n \times n}$. Then,

$$\|e^{JA}X - Xe^{JB}\| \leq \|AX - XB\|.$$

(Proof: See [1028].) (Remark: This result is a matrix version of x) of Fact 1.18.6.)

Fact 11.16.14. Let $A \in \mathbb{F}^{n \times n}$, and, for all $i = 1, \dots, n$, define $f_i: [0, \infty) \mapsto \mathbb{R}$ by $f_i(t) \triangleq \log \sigma_i(e^{tA})$. Then, A is normal if and only if, for all $i = 1, \dots, n$, f_i is convex. (Proof: See [93] and [452].) (Remark: The statement in [93] that convexity holds on $\mathbb{R}$ is erroneous. A counterexample is $A \triangleq \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ for which $\log \sigma_1(e^{tA}) = |t|$ and $\log \sigma_2(e^{tA}) = -|t|$.)

Fact 11.16.15. Let $A \in \mathbb{F}^{n \times n}$, and, for nonzero $x \in \mathbb{F}^n$, define $f_x: \mathbb{R} \mapsto \mathbb{R}$ by $f_x(t) \triangleq \log \sigma_{\max}(e^{tA}x)$. Then, A is normal if and only if, for all nonzero $x \in \mathbb{F}^n$, f_x is convex. (Proof: See [93].) (Remark: This result is due to Friedland.)

Fact 11.16.16. Let $A, B \in \mathbb{F}^{n \times n}$, assume that A and B are positive semidefinite, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|e^{(A-B)} - I\| \leq \|e^A - e^B\|$$

and

$$\|e^A + e^B\| \leq \|e^{A+B} + I\|.$$

(Proof: See [58] and [197, p. 294].) (Remark: See Fact 9.9.54.)

Fact 11.16.17. Let $A, X, B \in \mathbb{F}^{n \times n}$, assume that A and B are Hermitian, and let $\|\cdot\|$ be a unitarily invariant norm on $\mathbb{F}^{n \times n}$. Then,

$$\|AX - XB\| \leq \|e^{\frac{1}{2}A}Xe^{-\frac{1}{2}B} - e^{-\frac{1}{2}B}Xe^{\frac{1}{2}A}\|.$$

(Proof: See [216].) (Remark: See Fact 9.9.55.)

11.17 Facts on Stable Polynomials

Fact 11.17.1. Let $a_1, \dots, a_n$ be nonzero real numbers, let

$$\Delta \triangleq \{i \in \{1, \dots, n-1\} : \frac{a_{i+1}}{a_i} < 0\},$$

let $b_1, \dots, b_n$ be real numbers satisfying $b_1 < \dots < b_n$, define $f: (0, \infty) \mapsto \mathbb{R}$ by

$$f(x) = a_n x^{b_n} + \dots + a_1 x^{b_1},$$

and define

$$\mathfrak{S} \triangleq \{x \in (0, \infty) : f(x) = 0\}.$$

Furthermore, for all $x \in \mathfrak{S}$, define the multiplicity of x to be the positive integer m such that $f(x) = f'(x) = \cdots = f^{(m-1)}(x) = 0$ and $f^{(m)}(x) \neq 0$, and let $\mathfrak{S}'$ denote the multiset consisting of all elements of $\mathfrak{S}$ counting multiplicity. Then,

$$\text{card}(\mathfrak{S}') \leq \text{card}(\Delta).$$

If, in addition, $b_1, \dots, b_n$ are nonnegative integers, then $\text{card}(\Delta) - \text{card}(\mathfrak{S}')$ is even. (Proof: See [839, 1400].) (Remark: This result is the *Descartes rule of signs*.)

Fact 11.17.2. Let $p \in \mathbb{R}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$. If p is asymptotically stable, then $a_0, \dots, a_{n-1}$ are positive. Now, assume that $a_0, \dots, a_{n-1}$ are positive. Then, the following statements hold:

i) If $n = 1$ or $n = 2$, then p is asymptotically stable.

ii) If $n = 3$, then p is asymptotically stable if and only if

$$a_0 < a_1 a_2.$$

iii) If $n = 4$, then p is asymptotically stable if and only if

$$a_1^2 + a_0 a_3^2 < a_1 a_2 a_3.$$

iv) If $n = 5$, then p is asymptotically stable if and only if

$$\begin{aligned} a_2 &< a_3 a_4, \\ a_2^2 + a_1 a_4^2 &< a_0 a_4 + a_2 a_3 a_4, \\ a_0^2 + a_1 a_2^2 + a_1^2 a_4^2 + a_0 a_3^2 a_4 &< a_0 a_2 a_3 + 2a_0 a_1 a_4 + a_1 a_2 a_3 a_4. \end{aligned}$$

(Remark: These results are special cases of the *Routh criterion*, which provides stability criteria for polynomials of arbitrary degree n . See [301].)

Fact 11.17.3. Let $\varepsilon \in [0, 1]$, let $n \in \{2, 3, 4\}$, let $p_\varepsilon \in \mathbb{R}[s]$, where $p_\varepsilon(s) = s^n + a_{n-1}s^{n-1} + \cdots + \varepsilon a_0$, and assume that p_1 is asymptotically stable. Then, for all $\varepsilon \in (0, 1]$, p_ε is asymptotically stable. Furthermore, $p_0(s)/s$ is asymptotically stable. (Remark: The result does not hold for $n = 5$. A counterexample is $p(s) = s^5 + 2s^4 + 3s^3 + 5s^2 + 2s + 2.5\varepsilon$, which is asymptotically stable if and only if $\varepsilon \in (4/5, 1]$. This result is another instance of the quartic barrier. See [351], Fact 8.14.7, and Fact 8.15.37.)

Fact 11.17.4. Let $p \in \mathbb{R}[s]$ be monic, and define $q(s) \triangleq s^n p(1/s)$, where $n \triangleq \deg p$. Then, p is asymptotically stable if and only if q is asymptotically stable. (Remark: See Fact 4.8.1 and Fact 11.17.5.)

Fact 11.17.5. Let $p \in \mathbb{R}[s]$ be monic, and assume that p is semistable. Then, $q(s) \triangleq p(s)/s$ and $\hat{q}(s) \triangleq s^n p(1/s)$ are asymptotically stable. (Remark: See Fact 4.8.1 and Fact 11.17.4.)

Fact 11.17.6. Let $p, q \in \mathbb{R}[s]$, assume that p is even, assume that q is odd, and assume that every coefficient of $p+q$ is positive. Then, $p+q$ is asymptotically stable

if and only if every root of p and every root of q is imaginary, and the roots of p and the roots of q are interlaced on the imaginary axis. (Proof: See [221, 301, 705].) (Remark: This result is the *Hermite-Biehler* or *interlacing theorem*.) (Example: $s^2 + 2s + 5 = (s^2 + 5) + 2s$.)

Fact 11.17.7. Let $p \in \mathbb{R}[s]$ be asymptotically stable, and let $p(s) = \beta_n s^n + \beta_{n-1} s^{n-1} + \cdots + \beta_1 s + \beta_0$, where $\beta_n > 0$. Then, for all $i = 1, \dots, n-2$,

$$\beta_{i-1} \beta_{i+2} < \beta_i \beta_{i+1}.$$

(Remark: This result is a necessary condition for asymptotic stability, which can be used to show that a given polynomial with positive coefficients is unstable.)

(Remark: This result is due to Xie. See [1474]. For alternative conditions, see [221, p. 68].)

Fact 11.17.8. Let $n \in \mathbb{P}$ be even, let $m \triangleq n/2$, let $p \in \mathbb{R}[s]$, where $p(s) = \beta_n s^n + \beta_{n-1} s^{n-1} + \cdots + \beta_1 s + \beta_0$ and $\beta_n > 0$, and assume that p is asymptotically stable. Then, for all $i = 1, \dots, m-1$,

$$\binom{m}{i} \beta_0^{(m-i)/m} \beta_n^{i/m} \leq \beta_{2i}.$$

(Remark: This result is a necessary condition for asymptotic stability, which can be used to show that a given polynomial with positive coefficients is unstable.)

(Remark: This result is due to Borobia and Dormido. See [1474, 1475] for extensions to polynomials of odd degree.)

Fact 11.17.9. Let $p, q \in \mathbb{R}[s]$, where $p(s) = \alpha_n s^n + \alpha_{n-1} s^{n-1} + \cdots + \alpha_1 s + \alpha_0$ and $q(s) = \beta_m s^m + \beta_{m-1} s^{m-1} + \cdots + \beta_1 s + \beta_0$. If p and q are (Lyapunov, asymptotically) stable, then $r(s) \triangleq \alpha_l \beta_l s^l + \alpha_{l-1} \beta_{l-1} s^{l-1} + \cdots + \alpha_1 \beta_1 s + \alpha_0 \beta_0$, where $l \triangleq \min\{m, n\}$, is (Lyapunov, asymptotically) stable. (Proof: See [543].) (Remark: The polynomial r is the *Schur product* of p and q . See [82, 762].)

Fact 11.17.10. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is diagonalizable over $\mathbb{R}$. Then, χ_A has all positive coefficients if and only if A is asymptotically stable. (Proof: Sufficiency follows from Fact 11.17.2. For necessity, note that all of the roots of χ_A are real and that $\chi_A(\lambda) > 0$ for all $\lambda \geq 0$. Hence, $\text{roots}(\chi_A) \subset (-\infty, 0)$.)

Fact 11.17.11. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

- i) $\chi_{A \oplus A}$ has all positive coefficients.
- ii) $\chi_{A \oplus A}$ is asymptotically stable.
- iii) $A \oplus A$ is asymptotically stable.
- iv) A is asymptotically stable.

(Proof: If A is not asymptotically stable, then Fact 11.18.32 implies that $A \oplus A$ has a nonnegative eigenvalue λ . Since $\chi_{A \oplus A}(\lambda) = 0$, it follows that $\chi_{A \oplus A}$ cannot have all positive coefficients. See [519, Theorem 5].) (Remark: A similar method of proof is used in Proposition 8.2.7.)

Fact 11.17.12. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

i) χ_A and $\chi_{A^{(2,1)}}$ have all positive coefficients.

ii) A is asymptotically stable.

(Proof: See [1243].) (Remark: The additive compound $A^{(2,1)}$ is defined in Fact 7.5.17.)

Fact 11.17.13. For $i = 1, \dots, n-1$, let $a_i, b_i \in \mathbb{R}$ satisfy $0 < a_i \leq b_i$, define $\phi_1, \phi_2, \psi_1, \psi_2 \in \mathbb{R}[s]$ by

$$\begin{aligned}\phi_1(s) &= b_n s^n + a_{n-2} s^{n-2} + b_{n-4} s^{n-4} + \dots, \\ \phi_2(s) &= a_n s^n + b_{n-2} s^{n-2} + a_{n-4} s^{n-4} + \dots, \\ \psi_1(s) &= b_{n-1} s^{n-1} + a_{n-3} s^{n-3} + b_{n-5} s^{n-5} + \dots, \\ \psi_2(s) &= a_{n-1} s^{n-1} + b_{n-3} s^{n-3} + a_{n-5} s^{n-5} + \dots,\end{aligned}$$

assume that $\phi_1 + \psi_1$, $\phi_1 + \psi_2$, $\phi_2 + \psi_1$, and $\phi_2 + \psi_2$ are asymptotically stable, let $p \in \mathbb{R}[s]$, where $p(s) = \beta_n s^n + \beta_{n-1} s^{n-1} + \dots + \beta_1 s + \beta_0$, and assume that, for all $i = 1, \dots, n$, $a_i \leq \beta_i \leq b_i$. Then, p is asymptotically stable. (Proof: See [447, pp. 466, 467].) (Remark: This result is *Kharitonov's theorem*.)

11.18 Facts on Stable Matrices

Fact 11.18.1. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is semistable. Then, A is Lyapunov stable.

Fact 11.18.2. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is Lyapunov stable. Then, A is group invertible.

Fact 11.18.3. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is semistable. Then, A is group invertible.

Fact 11.18.4. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are similar. Then, A is (Lyapunov stable, semistable, asymptotically stable, discrete-time Lyapunov stable, discrete-time semistable, discrete-time asymptotically stable) if and only if B is.

Fact 11.18.5. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is semistable. Then,

$$\lim_{t \rightarrow \infty} e^{tA} = I - AA^\#,$$

and thus

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t e^{\tau A} d\tau = I - AA^\#.$$

(Remark: See Fact 10.11.6, Fact 11.18.1, and Fact 11.18.2.)

Fact 11.18.6. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is Lyapunov stable. Then,

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t e^{\tau A} d\tau = I - AA^\#.$$

(Remark: See Fact 11.18.2.)

Fact 11.18.7. Let $A, B \in \mathbb{F}^{n \times n}$. Then, $\lim_{\alpha \rightarrow \infty} e^{A+\alpha B}$ exists if and only if B is semistable. In this case,

$$\lim_{\alpha \rightarrow \infty} e^{A+\alpha B} = e^{(I-BB^\#)A}(I-BB^\#) = (I-BB^\#)e^{A(I-BB^\#)}.$$

(Proof: See [284].)

Fact 11.18.8. Let $A \in \mathbb{F}^{n \times n}$, assume that A is asymptotically stable, let $\beta > \text{spabs}(A)$, and let $\|\cdot\|$ be a submultiplicative norm on $\mathbb{F}^{n \times n}$. Then, there exists $\gamma > 0$ such that, for all $t \geq 0$,

$$\|e^{tA}\| \leq \gamma e^{\beta t}.$$

(Remark: See [558, pp. 201–206] and [786].)

Fact 11.18.9. Let $A \in \mathbb{R}^{n \times n}$, assume that A is asymptotically stable, let $\beta \in (\text{spabs}(A), 0)$, let $P \in \mathbb{R}^{n \times n}$ be positive definite and satisfy

$$A^T P + PA \leq 2\beta P,$$

and let $\|\cdot\|$ be a normalized, submultiplicative norm on $\mathbb{R}^{n \times n}$. Then, for all $t \geq 0$,

$$\|e^{tA}\| \leq \sqrt{\|P\|\|P^{-1}\|} e^{\beta t}.$$

(Remark: See [689].) (Remark: See Fact 11.15.9.)

Fact 11.18.10. Let $A \in \mathbb{F}^{n \times n}$, assume that A is asymptotically stable, let $R \in \mathbb{F}^{n \times n}$, assume that R is positive definite, and let $P \in \mathbb{F}^{n \times n}$ be the positive-definite solution of $A^*P + PA + R = 0$. Then,

$$\sigma_{\max}(e^{tA}) \leq \sqrt{\frac{\sigma_{\max}(P)}{\sigma_{\min}(P)}} e^{-t\lambda_{\min}(RP^{-1})/2}$$

and

$$\|e^{tA}\|_F \leq \sqrt{\|P\|_F\|P^{-1}\|_F} e^{-t\lambda_{\min}(RP^{-1})/2}.$$

If, in addition, $A + A^*$ is negative definite, then

$$\|e^{tA}\|_F \leq e^{-t\lambda_{\min}(-A-A^*)/2}.$$

(Proof: See [952].)

Fact 11.18.11. Let $A \in \mathbb{R}^{n \times n}$, assume that A is asymptotically stable, let $R \in \mathbb{R}^{n \times n}$, assume that R is positive definite, and let $P \in \mathbb{R}^{n \times n}$ be the positive-definite solution of $A^T P + PA + R = 0$. Furthermore, define the vector norm $\|x\|' \triangleq \sqrt{x^T P x}$ on $\mathbb{R}^n$, let $\|\cdot\|$ denote the induced norm on $\mathbb{R}^{n \times n}$, and let $\mu(\cdot)$ denote the corresponding logarithmic derivative. Then,

$$\mu(A) = -\lambda_{\min}(RP^{-1})/2.$$

Consequently,

$$\|e^{tA}\| \leq e^{-t\lambda_{\min}(RP^{-1})/2}.$$

(Proof: See [728] and use *xiv*) of Fact 11.15.7.) (Remark: See Fact 11.15.7 for the definition and properties of the logarithmic derivative.)

Fact 11.18.12. Let $A \in \mathbb{F}^{n \times n}$. Then, A is similar to a skew-Hermitian matrix if and only if there exists a positive-definite matrix $P \in \mathbb{F}^{n \times n}$ such that $A^*P + PA = 0$. (Remark: See Fact 5.9.4.)

Fact 11.18.13. Let $A \in \mathbb{R}^{n \times n}$. Then, A and A^2 are asymptotically stable if and only if, for all $\lambda \in \text{spec}(A)$, there exist $r > 0$ and $\theta \in (\frac{\pi}{2}, \frac{3\pi}{4}) \cup (\frac{5\pi}{4}, \frac{3\pi}{2})$ such that $\lambda = re^{j\theta}$.

Fact 11.18.14. Let $A \in \mathbb{R}^{n \times n}$. Then, A is group invertible and $2k\pi j \notin \text{spec}(A)$ for all $k \geq 1$ if and only if

$$AA^\# = (e^A - I)(e^A - I)^\#.$$

In particular, if A is semistable, then this identity holds. (Proof: Use *ii*) of Fact 11.21.10 and *ix*) of Proposition 11.8.2.)

Fact 11.18.15. Let $A \in \mathbb{F}^{n \times n}$. Then, A is asymptotically stable if and only if A^{-1} is asymptotically stable. Hence, $e^{tA} \rightarrow 0$ as $t \rightarrow \infty$ if and only if $e^{tA^{-1}} \rightarrow 0$ as $t \rightarrow \infty$.

Fact 11.18.16. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A is asymptotically stable, and assume that $\sigma_{\max}(B \oplus B) < \sigma_{\min}(A \oplus A)$. Then, $A + B$ is asymptotically stable. (Proof: Since $A \oplus A$ is nonsingular, Fact 9.14.18 implies that $A \oplus A + \alpha(B \oplus B) = (A + \alpha B) \oplus (A + \alpha B)$ is nonsingular for all $0 \leq \alpha \leq 1$. Now, suppose that $A + B$ is not asymptotically stable. Then, there exists $\alpha_0 \in (0, 1]$ such that $A + \alpha_0 B$ has an imaginary eigenvalue, and thus $(A + \alpha_0 B) \oplus (A + \alpha_0 B) = A \oplus A + \alpha_0(B \oplus B)$ is singular, which is a contradiction.) (Remark: This result provides a suboptimal solution of a nearness problem. See [679, Section 7] and Fact 9.14.18.)

Fact 11.18.17. Let $A \in \mathbb{C}^{n \times n}$, assume that A is asymptotically stable, let $\|\cdot\|$ denote either $\sigma_{\max}(\cdot)$ or $\|\cdot\|_F$, and define

$$\beta(A) \triangleq \{\|B\|: B \in \mathbb{C}^{n \times n} \text{ and } A + B \text{ is not asymptotically stable}\}.$$

Then,

$$\begin{aligned} \frac{1}{2}\sigma_{\min}(A \otimes A) &\leq \beta(A) \\ &= \min_{\gamma \in \mathbb{R}} \sigma_{\min}(A + \gamma jI) \\ &\leq \min\{\text{spabs}(A), \sigma_{\min}(A), \frac{1}{2}\sigma_{\max}(A + A^*)\}. \end{aligned}$$

Furthermore, let $R \in \mathbb{F}^{n \times n}$, assume that R is positive definite, and let $P \in \mathbb{F}^{n \times n}$ be the positive-definite solution of $A^*P + PA + R = 0$. Then,

$$\frac{1}{2}\sigma_{\min}(R)/\|P\| \leq \beta(A).$$

If, in addition, $A + A^*$ is negative definite, then

$$-\frac{1}{2}\lambda_{\min}(A + A^*) \leq \beta(A).$$

(Proof: See [679, 1360].) (Remark: The analogous problem for real matrices and real perturbations is discussed in [1108].)

Fact 11.18.18. Let $A \in \mathbb{F}^{n \times n}$, assume that A is asymptotically stable, let $V \in \mathbb{F}^{n \times n}$, assume that V is positive definite, and let $Q \in \mathbb{R}^n$ be the positive-definite solution of $AQ + QA^* + V = 0$. Then, for all $t \geq 0$,

$$\|e^{tA}\|_F^2 = \operatorname{tr} e^{tA} e^{tA^*} \leq \kappa(Q) \operatorname{tr} e^{-tS^{-1}VS^{-*}} \leq \kappa(Q) \operatorname{tr} e^{-[t/\sigma_{\max}(Q)]V},$$

where $S \in \mathbb{F}^{n \times n}$ satisfies $Q = SS^*$ and $\kappa(Q) \triangleq \sigma_{\max}(Q)/\sigma_{\min}(Q)$. If, in particular, A satisfies $AQ + QA^* + I = 0$, then

$$\|e^{tA}\|_F^2 \leq n\kappa(Q)e^{-t/\sigma_{\max}(Q)}.$$

(Proof: See [1468].) (Remark: Fact 11.15.4 yields $e^{tA}e^{tA^*} \leq e^{t(A+A^*)}$. However, this bound is poor when $A + A^*$ is not asymptotically stable. See [185].) (Remark: See Fact 11.18.19.)

Fact 11.18.19. Let $A \in \mathbb{F}^{n \times n}$, assume that A is asymptotically stable, let $V \in \mathbb{F}^{n \times n}$, assume that V is positive definite, and let $Q \in \mathbb{R}^n$ be the positive-definite solution of $AQ + QA^* + I = 0$. Then, for all $t \geq 0$,

$$\sigma_{\max}^2(e^{tA}) \leq \kappa(Q)e^{-t/\sigma_{\max}(Q)},$$

where $\kappa(Q) \triangleq \sigma_{\max}(Q)/\sigma_{\min}(Q)$. (Proof: See references in [1377, 1378].) (Remark: Since $\|e^{tA}\|_F \leq \sqrt{n}\sigma_{\max}(e^{tA})$, it follows that this inequality implies the last inequality in Fact 11.18.18.)

Fact 11.18.20. Let $A \in \mathbb{R}^{n \times n}$, and assume that every entry of $A \in \mathbb{R}^{n \times n}$ is positive. Then, A is unstable. (Proof: See Fact 4.11.5.)

Fact 11.18.21. Let $A \in \mathbb{R}^{n \times n}$. Then, A is asymptotically stable if and only if there exist matrices $B, C \in \mathbb{R}^{n \times n}$ such that B is positive definite, C is dissipative, and $A = BC$. (Proof: $A = P^{-1}(-A^TP - R)$.) (Remark: To reverse the order of factors, consider A^T .)

Fact 11.18.22. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements hold:

- i) All of the real eigenvalues of A are positive if and only if A is the product of two dissipative matrices.
- ii) A is nonsingular and $A \neq \alpha I$ for all $\alpha < 0$ if and only if A is the product of two asymptotically stable matrices.
- iii) A is nonsingular if and only if A is the product of three or fewer asymptotically stable matrices.

(Proof: See [126, 1459].)

Fact 11.18.23. Let $p \in \mathbb{R}[s]$, where $p(s) = s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_1s + \beta_0$ and $\beta_0, \dots, \beta_n > 0$. Furthermore, define $A \in \mathbb{R}^{n \times n}$ by

$$A \triangleq \begin{bmatrix} \beta_{n-1} & \beta_{n-3} & \beta_{n-5} & \beta_{n-7} & \cdots & \cdots & 0 \\ 1 & \beta_{n-2} & \beta_{n-4} & \beta_{n-6} & \cdots & \cdots & 0 \\ 0 & \beta_{n-1} & \beta_{n-3} & \beta_{n-5} & \cdots & \cdots & 0 \\ 0 & 1 & \beta_{n-2} & \beta_{n-4} & \cdots & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & \cdots & \beta_1 & 0 \\ 0 & 0 & 0 & \cdots & \cdots & \beta_2 & \beta_0 \end{bmatrix}.$$

If p is Lyapunov stable, then every subdeterminant of A is nonnegative. (Remark: A is *totally nonnegative*.) Furthermore, p is asymptotically stable if and only if every leading principal subdeterminant of A is positive. (Proof: See [82].) (Remark: The second statement is due to Hurwitz.) (Remark: The diagonal entries of A are $\beta_{n-1}, \dots, \beta_0$.) (Problem: Show that this condition for stability is equivalent to the condition given in [481, p. 183] in terms of an alternative matrix $\hat{A}$.)

Fact 11.18.24. Let $A \in \mathbb{R}^{n \times n}$, assume that A is tridiagonal, and assume that $A_{(i,i)} > 0$ for all $i = 1, \dots, n$ and $A_{(i,i+1)}A_{(i+1,i)} > 0$ for all $i = 1, \dots, n-1$. Then, A is asymptotically stable. (Proof: See [287].) (Remark: This result is due to Barnett and Storey.)

Fact 11.18.25. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is cyclic. Then, there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that $A_S \triangleq SAS^{-1}$ is given by the tridiagonal matrix

$$A_S = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ -\alpha_n & 0 & 1 & \cdots & 0 & 0 \\ 0 & -\alpha_{n-1} & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & -\alpha_2 & -\alpha_1 \end{bmatrix},$$

where $\alpha_1, \dots, \alpha_n$ are real numbers. If $\alpha_1\alpha_2 \cdots \alpha_n \neq 0$, then the number of eigenvalues of A in the OLHP is equal to the number of positive elements in $\{\alpha_1, \alpha_1\alpha_2, \dots, \alpha_1\alpha_2 \cdots \alpha_n\}_{\text{ms}}$. Furthermore, $A_S^T P + P A_S + R = 0$, where

$$P \triangleq \text{diag}(\alpha_1\alpha_2 \cdots \alpha_n, \alpha_1\alpha_2 \cdots \alpha_{n-1}, \dots, \alpha_1\alpha_2, \alpha_1)$$

and

$$R \triangleq \text{diag}(0, \dots, 0, 2\alpha_1^2).$$

Finally, A_S is asymptotically stable if and only if $\alpha_1, \dots, \alpha_n > 0$. (Remark: A_S is in *Schwarz form*.) (Proof: See [146, pp. 52, 95].) (Remark: See Fact 11.18.27 and Fact 11.18.26.)

Fact 11.18.26. Let $\alpha_1, \dots, \alpha_n$ be real numbers, and define $A \in \mathbb{R}^{n \times n}$ by

$$A = \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 & 0 \\ -\alpha_n & 0 & 1 & \cdots & 0 & 0 \\ 0 & -\alpha_{n-1} & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \cdots & -\alpha_2 & \alpha_1 \end{bmatrix}.$$

Then, $\text{spec}(A) \subset \text{ORHP}$ if and only if $\alpha_1, \dots, \alpha_n > 0$. (Proof: See [711, p. 111].) (Remark: Note the absence of the minus sign in the (n, n) entry compared to the matrix in Fact 11.18.25. This minus sign changes the sign of all eigenvalues of A .)

Fact 11.18.27. Let $\alpha_1, \alpha_2, \alpha_3 > 0$, and define $A_R, P, R \in \mathbb{R}^{3 \times 3}$ by the tridiagonal matrix

$$A_R \triangleq \begin{bmatrix} -\alpha_1 & \alpha_2^{1/2} & 0 \\ -\alpha_2^{1/2} & 0 & \alpha_3^{1/2} \\ 0 & -\alpha_3^{1/2} & 0 \end{bmatrix}$$

and the diagonal matrices

$$P \triangleq I, \quad R \triangleq \text{diag}(2\alpha_1, 0, 0).$$

Then, $A_R^T P + P A_R + R = 0$. (Remark: The matrix A_R is in *Routh form*. The Routh form A_R and the Schwarz form A_S are related by $A_R = S_{RS} A_S S_{RS}^{-1}$, where

$$S_{RS} \triangleq \begin{bmatrix} 0 & 0 & \alpha_1^{1/2} \\ 0 & -(\alpha_1 \alpha_2)^{1/2} & 0 \\ (\alpha_1 \alpha_2 \alpha_3)^{1/2} & 0 & 0 \end{bmatrix}.)$$

(Remark: See Fact 11.18.25.)

Fact 11.18.28. Let $\alpha_1, \alpha_2, \alpha_3 > 0$, and define $A_C, P, R \in \mathbb{R}^{3 \times 3}$ by the tridiagonal matrix

$$A_C \triangleq \begin{bmatrix} 0 & 1/a_3 & 0 \\ -1/a_2 & 0 & 1/a_2 \\ 0 & -1/a_1 & -1/a_1 \end{bmatrix}$$

and the diagonal matrices

$$P \triangleq \text{diag}(a_3, a_2, a_1), \quad R \triangleq \text{diag}(0, 0, 2),$$

where $a_1 \triangleq 1/\alpha_1$, $a_2 \triangleq \alpha_1/\alpha_2$, and $a_3 \triangleq \alpha_2/(\alpha_1 \alpha_3)$. Then, $A_C^T P + P A_C + R = 0$. (Remark: The matrix A_C is in *Chen form*.) The Schwarz form A_S and the Chen form A_C are related by $A_S = S_{SC} A_C S_{SC}^{-1}$, where

$$S_{SC} \triangleq \begin{bmatrix} 1/(\alpha_1 \alpha_3) & 0 & 0 \\ 0 & 1/\alpha_2 & 0 \\ 0 & 0 & 1/\alpha_1 \end{bmatrix}.)$$

(Proof: See [313, p. 346].) (Remark: The Schwarz, Routh, and Chen forms provide the basis for the Routh criterion. See [32, 268, 313, 1073].) (Remark: A circuit interpretation of the Chen form is given in [965].)

Fact 11.18.29. Let $\alpha_1, \dots, \alpha_n > 0$ and $\beta_1, \dots, \beta_n > 0$, and define $A \in \mathbb{R}^{n \times n}$ by

$$A = \begin{bmatrix} -\alpha_1 & 0 & \cdots & 0 & -\beta_1 \\ \beta_2 & -\alpha_2 & \cdots & 0 & 0 \\ \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & \ddots & -\alpha_{n-1} & 0 \\ 0 & 0 & \cdots & \beta_n & -\alpha_n \end{bmatrix}.$$

Then,

$$\chi_A(s) = (s + \alpha_1)(s + \alpha_2) \cdots (s + \alpha_n) + \beta_1 \beta_2 \cdots \beta_n.$$

Furthermore, if

$$(\cos \pi/n)^n < \frac{\alpha_1 \cdots \alpha_n}{\beta_1 \cdots \beta_n},$$

then A is asymptotically stable. (Remark: If $n = 2$, then A is asymptotically stable for all positive $\alpha_1, \beta_1, \alpha_2, \beta_2$.) (Proof: See [1213].) (Remark: This result is the *secant condition*.)

Fact 11.18.30. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is asymptotically stable.
- ii) There exist a negative-definite matrix $B \in \mathbb{F}^{n \times n}$, a skew-Hermitian matrix $C \in \mathbb{F}^{n \times n}$, and a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A = B + SCS^{-1}$.
- iii) There exist a negative-definite matrix $B \in \mathbb{F}^{n \times n}$, a skew-Hermitian matrix $C \in \mathbb{F}^{n \times n}$, and a nonsingular matrix $S \in \mathbb{F}^{n \times n}$ such that $A = S(B+C)S^{-1}$.

(Proof: See [370].)

Fact 11.18.31. Let $A \in \mathbb{R}^{n \times n}$, and let $k \geq 2$. Then, there exist asymptotically stable matrices $A_1, \dots, A_k \in \mathbb{R}^{n \times n}$ such that $A = \sum_{i=1}^k A_i$ if and only if $\operatorname{tr} A < 0$. (Proof: See [747].)

Fact 11.18.32. Let $A \in \mathbb{R}^{n \times n}$. Then, A is (Lyapunov stable, semistable, asymptotically stable) if and only if $A \oplus A$ is. (Proof: Use Fact 7.5.7 and the fact that $\operatorname{vec}(e^{tA} V e^{tA^*}) = e^{t(A \oplus \overline{A})} \operatorname{vec} V$.)

Fact 11.18.33. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{m \times m}$. Then, the following statements hold:

- i) If A and B are (Lyapunov stable, semistable, asymptotically stable), then so is $A \oplus B$.
- ii) If $A \oplus B$ is (Lyapunov stable, semistable, asymptotically stable), then so is either A or B .

(Proof: Use Fact 7.5.7.)

Fact 11.18.34. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is asymptotically stable. Then,

$$(A \oplus A)^{-1} = \int_{-\infty}^{\infty} (j\omega I - A)^{-1} \otimes (j\omega I - A)^{-1} d\omega$$

and

$$\int_{-\infty}^{\infty} (\omega^2 I + A^2) d\omega = -\pi A^{-1}.$$

(Proof: Use $(j\omega I - A)^{-1} + (-j\omega I - A)^{-1} = -2A(\omega^2 I + A^2)^{-1}$.)

Fact 11.18.35. Let $A \in \mathbb{R}^{2 \times 2}$. Then, A is asymptotically stable if and only if $\text{tr } A < 0$ and $\det A > 0$.

Fact 11.18.36. Let $A \in \mathbb{C}^{n \times n}$. Then, there exists a unique asymptotically stable matrix $B \in \mathbb{C}^{n \times n}$ such that $B^2 = -A$. (Remark: This result is stated in [1231]. The uniqueness of the square root for complex matrices that have no eigenvalues in $(-\infty, 0]$ is implicitly assumed in [1232].) (Remark: See Fact 5.15.19.)

Fact 11.18.37. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) If A is semidissipative, then A is Lyapunov stable.
- ii) If A is dissipative, then A is asymptotically stable.
- iii) If A is Lyapunov stable and normal, then A is semidissipative.
- iv) If A is asymptotically stable and normal, then A is dissipative.
- v) If A is discrete-time Lyapunov stable and normal, then A is semicontractive.

Fact 11.18.38. Let $M \in \mathbb{R}^{r \times r}$, assume that M is positive definite, let $C, K \in \mathbb{R}^{r \times r}$, assume that C and K are positive semidefinite, and consider the equation

$$M\ddot{q} + C\dot{q} + Kq = 0.$$

Furthermore, define

$$A \triangleq \begin{bmatrix} 0 & I \\ -M^{-1}K & -M^{-1}C \end{bmatrix}.$$

Then, the following statements hold:

- i) A is Lyapunov stable if and only if $C + K$ is positive definite.
- ii) A is Lyapunov stable if and only if $\text{rank} \begin{bmatrix} C \\ K \end{bmatrix} = r$.
- iii) A is semistable if and only if $(M^{-1}K, C)$ is observable.
- iv) A is asymptotically stable if and only if A is semistable and K is positive definite.

(Proof: See [186].) (Remark: See Fact 5.12.21.)

11.19 Facts on Almost Nonnegative Matrices

Fact 11.19.1. Let $A \in \mathbb{R}^{n \times n}$. Then, e^{tA} is nonnegative for all $t \geq 0$ if and only if A is almost nonnegative. (Proof: Let $\alpha > 0$ be such that $\alpha I + A$ is nonnegative, and consider $e^{t(\alpha I + A)}$. See [181, p. 74], [182, p. 146], [190, 365], or [1197, p. 37].)

Fact 11.19.2. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is almost nonnegative. Then, e^{tA} is positive for all $t > 0$ if and only if A is irreducible. (Proof: See [1184, p. 208].)

Fact 11.19.3. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, and assume that A is almost nonnegative. Then, the following statements are equivalent:

- i) There exist $\alpha \in (0, \infty)$ and $B \in \mathbb{R}^{n \times n}$ such that $A = B - \alpha I$, B is nonnegative, and $\text{sprad}(B) \leq \alpha$.
- ii) $\text{spec}(A) \subset \text{OLHP} \cup \{0\}$.
- iii) $\text{spec}(A) \subset \text{CLHP}$.
- iv) If $\lambda \in \text{spec}(A)$ is real, then $\lambda \leq 0$.
- v) Every principal subdeterminant of $-A$ is nonnegative.
- vi) For every diagonal, positive-definite matrix $B \in \mathbb{R}^{n \times n}$, it follows that $A - B$ is nonsingular.

(Example: $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$.) (Remark: A is an N -matrix if A is almost nonnegative and i)–vi) hold.) (Remark: This result follows from Fact 4.11.6.)

Fact 11.19.4. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, and assume that A is almost nonnegative. Then, the following conditions are equivalent:

- i) A is a group-invertible N -matrix.
- ii) A is a Lyapunov-stable N -matrix.
- iii) A is a semistable N -matrix.
- iv) A is Lyapunov stable.
- v) A is semistable.
- vi) A is an N -matrix, and there exist $\alpha \in (0, \infty)$ and a nonnegative matrix $B \in \mathbb{R}^{n \times n}$ such that $A = B - \alpha I$ and $\alpha^{-1}B$ is discrete-time semistable.
- vii) There exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that $A^T P + PA$ is negative semidefinite.

Furthermore, consider the following statements:

- viii) There exists a positive vector $p \in \mathbb{R}^n$ such that $-Ap$ is nonnegative.
- ix) There exists a nonzero nonnegative vector $p \in \mathbb{R}^n$ such that $-Ap$ is nonnegative.

Then, $\text{viii}) \implies [i\text{--}vii]) \implies ix)$. (Proof: See [182, pp. 152–155] and [183]. The statement $[i\text{--}vii]) \implies ix)$ is given by Fact 4.11.10.) (Remark: The converse of

$viii) \implies [i)-vii])$ does not hold. For example, $A = \begin{bmatrix} 0 & 1 \\ 0 & -1 \end{bmatrix}$ is almost negative and semistable, but there does not exist a positive vector $p \in \mathbb{R}^2$ such that $-Ap$ is nonnegative. However, note that $viii)$ holds for A^T , but not for $\text{diag}(A, A^T)$ or its transpose.) (Remark: A discrete-time semistable matrix is called *semiconvergent* in [182, p. 152].) (Remark: The last statement follows from the fact that the function $V(x) = p^T x$ is a Lyapunov function for the system $\dot{x} = -Ax$ for $x \in [0, \infty)^n$ with Lyapunov derivative $\dot{V}(x) = -A^T p$. See [187, 615].)

Fact 11.19.5. Let $A \in \mathbb{R}^{n \times n}$, where $n \geq 2$, and assume that A is almost nonnegative. Then, the following conditions are equivalent:

- i) A is a nonsingular N-matrix.
- ii) A is asymptotically stable.
- iii) A is an asymptotically stable N-matrix.
- iv) There exist $\alpha \in (0, \infty)$ and a nonnegative matrix $B \in \mathbb{R}^{n \times n}$ such that $A = B - \alpha I$ and $\text{sprad}(B) < \alpha$.
- v) If $\lambda \in \text{spec}(A)$ is real, then $\lambda < 0$.
- vi) If $B \in \mathbb{R}^{n \times n}$ is nonnegative and diagonal, then $A - B$ is nonsingular.
- vii) Every principal subdeterminant of $-A$ is positive.
- viii) Every leading principal subdeterminant of $-A$ is positive.
- ix) For all $i = 1, \dots, n$, the sign of the i th leading principal subdeterminant of A is $(-1)^i$.
- x) For all $k \in \{1, \dots, n\}$, the sum of all $k \times k$ principal subdeterminants of $-A$ is positive.
- xi) There exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that $A^T P + PA$ is negative definite.
- xii) There exists a positive vector $p \in \mathbb{R}^n$ such that $-Ap$ is positive.
- xiii) There exists a nonnegative vector $p \in \mathbb{R}^n$ such that $-Ap$ is positive.
- xiv) If $p \in \mathbb{R}^n$ and $-Ap$ is nonnegative, then $p \geq 0$ is nonnegative.
- xv) For every nonnegative vector $y \in \mathbb{R}^n$, there exists a unique nonnegative vector $x \in \mathbb{R}^n$ such that $Ax = -y$.
- xvi) A is nonsingular and $-A^{-1}$ is nonnegative.

(Proof: See [181, pp. 134–140] or [711, pp. 114–116].) (Remark: $-A$ is a nonsingular M-matrix. See Fact 4.11.6.)

Fact 11.19.6. For $i, j = 1, \dots, n$, let $\sigma_{ij} \in [0, \infty)$, and define $A \in \mathbb{R}^{n \times n}$ by $A_{(i,j)} \triangleq \sigma_{ij}$ for all $i \neq j$ and $A_{(i,i)} \triangleq -\sum_{j=1}^n \sigma_{ij}$. Then, the following statements hold:

- i) A is almost nonnegative.
- ii) $-A1_{n \times 1} = \begin{bmatrix} \sigma_{11} & \dots & \sigma_{nn} \end{bmatrix}^T$ is nonnegative.

- iii) $\text{spec}(A) \subset \text{OLHP} \cup \{0\}$.
- iv) A is an N-matrix.
- v) A is a group-invertible N-matrix.
- vi) A is a Lyapunov-stable N-matrix.
- vii) A is a semistable N-matrix.

If, in addition, $\sigma_{11}, \dots, \sigma_{nn}$ are positive, then A is a nonsingular N-matrix. (Proof: It follows from the Gershgorin circle theorem given by Fact 4.10.16 that every eigenvalue λ of A is an element of a disk in $\mathbb{C}$ centered at $-\sum_{j=1}^n \sigma_{ij} \leq 0$ and with radius $\sum_{j=1, j \neq i}^n \sigma_{ij}$. Hence, if $\sigma_{ii} = 0$, then either $\lambda = 0$ or $\text{Re } \lambda < 0$, whereas, if $\sigma_{ii} > 0$, then $\text{Re } \lambda \leq \sigma_{ii} < 0$. Thus, iii) holds. Statements iv)–vii) follow from ii) and Fact 11.19.4. The last statement follows from the Gershgorin circle theorem.) (Remark: A^T is a *compartmental matrix*. See [190, 617, 1387].) (Problem: Determine necessary and sufficient conditions on the parameters σ_{ij} such that A is a nonsingular N-matrix.)

Fact 11.19.7. Let $\mathcal{G} = (\mathcal{X}, \mathcal{R})$ be a graph, where $\mathcal{X} = \{x_1, \dots, x_n\}$, and let $L \in \mathbb{R}^{n \times n}$ denote either the in-Laplacian or the out-Laplacian of $\mathcal{G}$. Then, the following statements hold:

- i) $-L$ is semistable.
- ii) $\lim_{t \rightarrow \infty} e^{-Lt}$ exists.

(Remark: Use Fact 11.19.6.) (Remark: The spectrum of the Laplacian is discussed in [7].)

Fact 11.19.8. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is asymptotically stable. Then, at least one of the following statements holds:

- i) All of the diagonal entries of A are negative.
- ii) At least one diagonal entry of A is negative and at least one off-diagonal entry of A is negative.

(Proof: See [506].) (Remark: *sign stability* is discussed in [751].)

11.20 Facts on Discrete-Time-Stable Polynomials

Fact 11.20.1. Let $p \in \mathbb{R}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \dots + a_0$. Then, the following statements hold:

- i) If $n = 1$, then p is discrete-time asymptotically stable if and only if $|a_0| < 1$.
- ii) If $n = 2$, then p is discrete-time asymptotically stable if and only if $|a_0| < 1$ and $|a_1| < 1 + a_0$.
- iii) If $n = 3$, then p is discrete-time asymptotically stable if and only if $|a_0| < 1$, $|a_0 + a_2| < |1 + a_1|$, and $|a_1 - a_0 a_2| < 1 - a_0^2$.

(Remark: These results are the *Schur-Cohn criterion*. See [136, p. 185]. Conditions

for polynomials of arbitrary degree n follow from the *Jury test*. See [313, 782].) (Remark: For $n = 3$, an alternative form is given in [690, p. 355].)

Fact 11.20.2. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, and define $\hat{p} \in \mathbb{C}[s]$ by

$$\hat{p}(s) \triangleq z^{n-1} + \frac{a_{n-1} - a_0 \bar{a}_1}{1 - |a_0|^2} z^{n-1} + \frac{a_{n-2} - a_0 \bar{a}_2}{1 - |a_0|^2} z^{n-2} + \cdots + \frac{a_1 - a_0 \bar{a}_{n-1}}{1 - |a_0|^2}.$$

Then, p is discrete-time asymptotically stable if and only if $|a_0| < 1$ and $\hat{p}$ is discrete-time asymptotically stable. (Proof: See [690, p. 354].)

Fact 11.20.3. Let $p \in \mathbb{R}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$. Then, the following statements hold:

- i) If $a_0 \leq \cdots \leq a_{n-1} \leq 1$, then $\text{roots}(p) \subset \{z \in \mathbb{C} : |z| \leq 1 + |a_0| - a_0\}$.
- ii) If $0 < a_0 \leq \cdots \leq a_{n-1} \leq 1$, then $\text{roots}(p) \subset \text{CUD}$.
- iii) If $0 < a_0 < \cdots < a_{n-1} < 1$, then p is discrete-time asymptotically stable.

(Proof: For i), see [1189]. For ii), see [1004, p. 272]. For iii), use Fact 11.20.2. See [690, p. 355].) (Remark: If there exists $r > 0$ such that $0 < ra_0 < \cdots < r^{n-1}a_{n-1} < r^n$, then $\text{roots}(p) \subset \{z \in \mathbb{C} : |z| \leq r\}$.) (Remark: Statement ii) is the *Enestrom-Keakeya theorem*.)

Fact 11.20.4. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, assume that $a_0, \dots, a_{n-1}$ are nonzero, and let $\lambda \in \text{roots}(p)$. Then,

$$|\lambda| \leq \max\{2|a_{n-1}|, 2|a_{n-2}/a_{n-1}|, \dots, 2|a_1/a_2|, |a_0/a_1|\}.$$

(Remark: This result is due to Bourbaki. See [1005].)

Fact 11.20.5. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, assume that $a_0, \dots, a_{n-1}$ are nonzero, and let $\lambda \in \text{roots}(p)$. Then,

$$|\lambda| \leq \sum_{i=1}^{n-1} |a_i|^{1/(n-i)}$$

and

$$|\lambda + \frac{1}{2}a_{n-1}| \leq \frac{1}{2}|a_{n-1}| + \sum_{i=0}^{n-2} |a_i|^{1/(n-i)}.$$

(Remark: These results are due to Walsh. See [1005].)

Fact 11.20.6. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, and let $\lambda \in \text{roots}(p)$. Then,

$$\frac{|a_0|}{|a_0| + \max\{|a_1|, \dots, |a_{n-1}|, 1\}} < |\lambda| \leq \max\{|a_0|, 1 + |a_1|, \dots, 1 + |a_{n-1}|\}.$$

(Proof: The lower bound is proved in [1005], while the upper bound is proved in [401].) (Remark: The upper bound is *Cauchy's estimate*.) (Remark: The weaker upper bound

$$|\lambda| < 1 + \max_{i=0, \dots, n-1} |a_i|$$

is given in [136, p. 184] and [1005].)

Fact 11.20.7. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, and let $\lambda \in \text{roots}(p)$. Then,

$$|\lambda| \leq \frac{1}{2}(1 + |a_{n-1}|) + \sqrt{\max_{i=0, \dots, n-2} |a_i| + \frac{1}{4}(1 - |a_{n-1}|)^2},$$

$$|\lambda| \leq \max\{2, |a_0| + |a_{n-1}|, |a_1| + |a_{n-1}|, \dots, |a_{n-2}| + |a_{n-1}|\},$$

$$|\lambda| \leq \sqrt{2 + \max_{i=0, \dots, n-2} |a_i|^2 + |a_{n-1}|^2}.$$

(Proof: See [401].) (Remark: The first inequality is due to Joyal, Labelle, and Rahman. See [1005].)

Fact 11.20.8. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, assume that $a_0, \dots, a_{n-1}$ are nonzero, define

$$\alpha \triangleq \max \left\{ \left| \frac{a_0}{a_1} \right|, \left| \frac{a_1}{a_2} \right|, \dots, \left| \frac{a_{n-2}}{a_{n-1}} \right| \right\}$$

and

$$\beta \triangleq \max \left\{ \left| \frac{a_1}{a_2} \right|, \left| \frac{a_2}{a_3} \right|, \dots, \left| \frac{a_{n-2}}{a_{n-1}} \right| \right\},$$

and let $\lambda \in \text{roots}(p)$. Then,

$$|\lambda| \leq \frac{1}{2}(\beta + |a_{n-1}|) + \sqrt{\alpha|a_{n-1}| + \frac{1}{4}(\beta - |a_{n-1}|)^2},$$

$$|\lambda| \leq |a_{n-1}| + \alpha,$$

$$|\lambda| \leq \max \left\{ \left| \frac{a_0}{a_1} \right|, 2\beta, 2|a_{n-1}| \right\},$$

$$|\lambda| \leq 2 \max_{i=1, \dots, n-1} |a_i|^{1/(n-i)},$$

$$|\lambda| \leq \sqrt{2|a_{n-1}|^2 + \alpha^2 + \beta^2}.$$

(Proof: See [401, 918].) (Remark: The third inequality is *Kojima's bound*, while the fourth inequality is *Fujiwara's bound*.)

Fact 11.20.9. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, define $\alpha \triangleq 1 + \sum_{i=0}^{n-1} |a_i|^2$, and let $\lambda \in \text{roots}(p)$. Then,

$$|\lambda| \leq \frac{1}{n}|a_{n-1}| + \sqrt{\frac{n}{n-1} \left(n-1 + \sum_{i=0}^{n-1} |a_i|^2 - \frac{1}{n}|a_{n-1}|^2 \right)},$$

$$|\lambda| \leq \frac{1}{2} \left(|a_{n-1}| + 1 + \sqrt{(|a_{n-1}| - 1)^2 + 4 \sqrt{\sum_{i=0}^{n-2} |a_i|^2}} \right),$$

$$|\lambda| \leq \frac{1}{2} \left(|a_{n-1}| + \cos \frac{\pi}{n} + \sqrt{(|a_{n-1}| - \cos \frac{\pi}{n})^2 + (|a_{n-2}| + 1)^2 + \sum_{i=0}^{n-3} |a_i|^2} \right),$$

$$|\lambda| \leq \cos \frac{\pi}{n+1} + \frac{1}{2} \left(|a_{n-1}| + \sqrt{\sum_{i=0}^{n-1} |a_i|^2} \right),$$

and

$$\sqrt{\frac{1}{2}(\alpha - \sqrt{\alpha^2 - 4|a_0|^2})} \leq |\lambda| \leq \sqrt{\frac{1}{2}(\alpha + \sqrt{\alpha^2 - 4|a_0|^2})}.$$

Furthermore,

$$|\operatorname{Re} \lambda| \leq \frac{1}{2} \left(|\operatorname{Re} a_{n-1}| + \cos \frac{\pi}{n} + \sqrt{(|\operatorname{Re} a_{n-1}| - \cos \frac{\pi}{n})^2 + (|a_{n-2}| + 1)^2 + \sum_{i=0}^{n-3} |a_i|^2} \right)$$

and

$$|\operatorname{Im} \lambda| \leq \frac{1}{2} \left(|\operatorname{Im} a_{n-1}| + \cos \frac{\pi}{n} + \sqrt{(|\operatorname{Im} a_{n-1}| - \cos \frac{\pi}{n})^2 + (|a_{n-2}| + 1)^2 + \sum_{i=0}^{n-3} |a_i|^2} \right).$$

(Proof: See [514, 822, 826, 918].) (Remark: The first bound is due to Linden (see [826]), the fourth bound is due to Fujii and Kubo, and the upper bound in the fifth result, which follows from Fact 5.11.21 and Fact 5.11.30, is due to Parodi, see also [802, 817].) (Remark: The Parodi bound is a refinement of the Carmichael-Mason Bound. See Fact 11.20.10.)

Fact 11.20.10. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, let $r, q \in (1, \infty)$, assume that $1/r + 1/q = 1$, define $\alpha \triangleq (\sum_{i=0}^{n-1} |a_i|^r)^{1/r}$, and let $\lambda \in \operatorname{roots}(p)$. Then,

$$|\lambda| \leq (1 + \alpha^q)^{1/q}.$$

In particular, if $r = q = 2$, then

$$|\lambda| \leq \sqrt{1 + |a_{n-1}|^2 + \cdots + |a_0|^2}.$$

(Proof: See [918, 1005].) (Remark: Letting $r \rightarrow \infty$ yields the upper bound in Fact 11.20.6.) (Remark: The result for $r = q = 2$ is due to Carmichael and Mason.)

Fact 11.20.11. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \cdots + a_0$, let $\operatorname{mroots}(p) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}}$, and let $r > 0$ be the unique positive root of $\hat{p}(s) \triangleq s^n - |a_{n-1}|s^{n-1} - \cdots - |a_0|$. Then,

$$r(\sqrt[n]{2} - 1) \leq \max_{i=1, \dots, n} |\lambda_i| \leq r.$$

Furthermore,

$$r(\sqrt[n]{2} - 1) \leq \frac{1}{n} \sum_{i=1}^n |\lambda_i| < r.$$

Finally, the third inequality is an equality if and only if $\lambda_1 = \dots = \lambda_n$. (Remark: The first inequality is due to Cohn, the second inequality is due to Cauchy, and the third and fourth inequalities are due to Berwald. See [1005] and [1004, p. 245].)

Fact 11.20.12. Let $p \in \mathbb{C}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \dots + a_0$, define $\alpha \triangleq 1 + \sum_{i=0}^{n-1} |a_i|^2$, and let $\lambda \in \text{roots}(p)$. Then,

$$\sqrt{\frac{1}{2}(\alpha - \sqrt{\alpha^2 - 4|a_0|^2})} \leq |\lambda| \leq \sqrt{\frac{1}{2}(\alpha + \sqrt{\alpha^2 - 4|a_0|^2})}.$$

(Proof: See [823]. The result follows from Fact 5.11.29 and Fact 5.11.30.)

Fact 11.20.13. Let $p \in \mathbb{R}[s]$, where $p(s) = s^n + a_{n-1}s^{n-1} + \dots + a_0$, assume that $a_0, \dots, a_{n-1}$ are nonnegative, and let $x_1, \dots, x_m \in [0, \infty)$. Then,

$$p(\sqrt[n]{x_1 \cdots x_m}) \leq \sqrt[n]{p(x_1) \cdots p(x_m)}.$$

(Proof: See [1040].) (Remark: This result, which is due to Mihet, extends a result of Huygens for the case $p(x) = x + 1$.)

11.21 Facts on Discrete-Time-Stable Matrices

Fact 11.21.1. Let $A \in \mathbb{R}^{2 \times 2}$. Then, A is discrete-time asymptotically stable if and only if $|\text{tr } A| < 1 + \det A$ and $|\det A| < 1$.

Fact 11.21.2. Let $A \in \mathbb{F}^{n \times n}$. Then, A is discrete-time (Lyapunov stable, semistable, asymptotically stable) if and only if A^2 is.

Fact 11.21.3. Let $A \in \mathbb{R}^{n \times n}$, and let $\chi_A(s) = s^n + a_{n-1}s^{n-1} + \dots + a_1s + a_0$. Then, for all $k \geq 0$,

$$A^k = x_1(k)I + x_2(k)A + \dots + x_n(k)A^{n-1},$$

where, for all $i = 1, \dots, n$ and all $k \geq 0$, $x_i: \mathbb{N} \mapsto \mathbb{R}$ satisfies

$$x_i(k+n) + a_{n-1}x_i(k+n-1) + \dots + a_1x_i(k+1) + a_0x_i(k) = 0,$$

with, for all $i, j = 1, \dots, n$, the initial conditions

$$x_i(j-1) = \delta_{ij}.$$

(Proof: See [853].)

Fact 11.21.4. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) If A is semicontractive, then A is discrete-time Lyapunov stable.
- ii) If A is contractive, then A is discrete-time asymptotically stable.

- iii) If A is discrete-time Lyapunov stable and normal, then A is semicontractive.
- iv) If A is discrete-time asymptotically stable and normal, then A is contractive.

(Problem: Prove these results by using Fact 11.15.6.)

Fact 11.21.5. Let $A \in \mathbb{F}^{n \times n}$. Then, A is discrete-time (Lyapunov stable, semistable, asymptotically stable) if and only if $A \otimes A$ is. (Proof: Use Fact 7.4.15.)

Fact 11.21.6. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{m \times m}$. Then, the following statements hold:

- i) If A and B are discrete-time (Lyapunov stable, semistable, asymptotically stable), then $A \otimes B$ is discrete-time (Lyapunov stable, semistable, asymptotically stable).
- ii) If $A \otimes B$ is discrete-time (Lyapunov stable, semistable, asymptotically stable), then either A or B is discrete-time (Lyapunov stable, semistable, asymptotically stable).

(Proof: Use Fact 7.4.15.)

Fact 11.21.7. Let $A \in \mathbb{R}^{n \times n}$, and assume that A is (Lyapunov stable, semistable, asymptotically stable). Then, e^A is discrete-time (Lyapunov stable, semistable, asymptotically stable). (Problem: If $B \in \mathbb{R}^{n \times n}$ is discrete-time (Lyapunov stable, semistable, asymptotically stable), when does there exist a (Lyapunov-stable, semistable, asymptotically stable) matrix $A \in \mathbb{R}^{n \times n}$ such that $B = e^A$? See Proposition 11.4.3.)

Fact 11.21.8. The following statements hold:

- i) If $A \in \mathbb{R}^{n \times n}$ is discrete-time asymptotically stable, then $B \triangleq (A+I)^{-1}(A-I)$ is asymptotically stable.
- ii) If $B \in \mathbb{R}^{n \times n}$ is asymptotically stable, then $A \triangleq (I+B)(I-B)^{-1}$ is discrete-time asymptotically stable.
- iii) If $A \in \mathbb{R}^{n \times n}$ is discrete-time asymptotically stable, then there exists a unique asymptotically stable matrix $B \in \mathbb{R}^{n \times n}$ such that $A = (I+B)(I-B)^{-1}$. In fact, $B = (A+I)^{-1}(A-I)$.
- iv) If $B \in \mathbb{R}^{n \times n}$ is asymptotically stable, then there exists a unique discrete-time asymptotically stable matrix $A \in \mathbb{R}^{n \times n}$ such that $B = (A+I)^{-1}(A-I)$. In fact, $A = (I+B)(I-B)^{-1}$.

(Proof: See [657].) (Remark: For additional results on the Cayley transform, see Fact 3.11.29, Fact 3.11.28, Fact 3.11.30, Fact 3.19.12, and Fact 8.9.30.) (Problem: Obtain analogous results for Lyapunov-stable and semistable matrices.)

Fact 11.21.9. Let $\begin{bmatrix} P_1 & P_{12} \\ P_{12}^T & P_2 \end{bmatrix} \in \mathbb{R}^{2n \times 2n}$ be positive definite, where $P_1, P_{12}, P_2 \in \mathbb{R}^{n \times n}$. If $P_1 \geq P_2$, then $A \triangleq P_1^{-1}P_{12}^T$ is discrete-time asymptotically stable, while,

if $P_2 \geq P_1$, then $A \triangleq P_2^{-1}P_{12}$ is discrete-time asymptotically stable. (Proof: If $P_1 \geq P_2$, then $P_1 - P_{12}P_1^{-1}P_1P_1^{-1}P_{12}^T \geq P_1 - P_{12}P_2^{-2}P_{12}^T > 0$. See [334].)

Fact 11.21.10. Let $A \in \mathbb{R}^{n \times n}$, and let $\|\cdot\|$ be a norm on $\mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) A is discrete-time Lyapunov stable if and only if $\{\|A^k\|\}_{k=0}^\infty$ is bounded.
- ii) A is discrete-time semistable if and only if $A_\infty \triangleq \lim_{k \rightarrow \infty} A^k$ exists.
- iii) Assume that A is discrete-time semistable. Then, $A_\infty \triangleq I - (A - I)(A - I)^\#$ is idempotent and $\text{rank } A_\infty = \text{amult}_A(1)$. If, in addition, $\text{rank } A = 1$, then, for every eigenvector x of A associated with the eigenvalue 1, there exists $y \in \mathbb{F}^n$ such that $y^*x = 1$ and $A_\infty = xy^*$.
- iv) A is discrete-time asymptotically stable if and only if $\lim_{k \rightarrow \infty} A^k = 0$.

(Remark: A proof of ii) is given in [998, p. 640]. See Fact 11.21.14.)

Fact 11.21.11. Let $A \in \mathbb{F}^{n \times n}$. Then, A is discrete-time Lyapunov stable if and only if

$$A_\infty \triangleq \lim_{k \rightarrow \infty} \frac{1}{k} \sum_{i=0}^{k-1} A^i$$

exists. In this case,

$$A_\infty = I - (A - I)(A - I)^\#.$$

(Proof: See [998, p. 633].) (Remark: A is *Cesaro summable*.) (Remark: See Fact 6.3.34.)

Fact 11.21.12. Let $A \in \mathbb{F}^{n \times n}$. Then, A is discrete-time asymptotically stable if and only if

$$\lim_{k \rightarrow \infty} A^k = 0.$$

In this case,

$$(I - A)^{-1} = \sum_{i=1}^{\infty} A^i,$$

where the series converges absolutely.

Fact 11.21.13. Let $A \in \mathbb{F}^{n \times n}$, and assume that A is unitary. Then, A is discrete-time Lyapunov stable.

Fact 11.21.14. Let $A, B \in \mathbb{R}^{n \times n}$, assume that A is discrete-time semistable, and let $A_\infty \triangleq \lim_{k \rightarrow \infty} A^k$. Then,

$$\lim_{k \rightarrow \infty} \left(A + \frac{1}{k}B\right)^k = A_\infty e^{A_\infty B A_\infty}.$$

(Proof: See [233, 1429].) (Remark: If A is idempotent, then $A_\infty = A$. The existence of A_∞ is guaranteed by Fact 11.21.10, which also implies that A_∞ is idempotent.)

Fact 11.21.15. Let $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) A is discrete-time Lyapunov stable if and only if there exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that $P - A^T P A$ is positive semidefinite.

- ii) A is discrete-time asymptotically stable if and only if there exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ such that $P - A^T P A$ is positive definite.

(Remark: The *discrete-time Lyapunov equation* or the *Stein equation* is $P = A^T P A + R$.)

Fact 11.21.16. Let $(A_k)_{k=0}^\infty \subset \mathbb{R}^{n \times n}$ and, for $k \in \mathbb{N}$, consider the discrete-time, time-varying system

$$x_{k+1} = A_k x_k.$$

Furthermore, assume there exist real numbers $\beta \in (0, 1)$, $\gamma > 0$, and $\varepsilon > 0$ such that, for all $k \in \mathbb{N}$,

$$\text{sprad}(A_k) < \beta,$$

$$\|A_k\| < \gamma,$$

$$\|A_{k+1} - A_k\| < \varepsilon,$$

where $\|\cdot\|$ is a norm on $\mathbb{R}^{n \times n}$. Then, $x_k \rightarrow 0$ as $k \rightarrow \infty$. (Proof: See [642, pp. 170–173].) (Remark: This result arises from the theory of *infinite matrix products*. See [76, 230, 231, 375, 608, 704, 861].)

Fact 11.21.17. Let $A \in \mathbb{F}^{n \times n}$, and define

$$r(A) \triangleq \sup_{\{z \in \mathbb{C}: |z| > 1\}} \frac{|z| - 1}{\sigma_{\min}(zI - A)}.$$

Then,

$$r(A) \leq \sup_{k \geq 0} \sigma_{\max}(A^k) \leq \text{ner}(A).$$

Hence, if A is discrete-time Lyapunov stable, then $r(A)$ is finite. (Proof: See [1413].) (Remark: This result is the *Kreiss matrix theorem*.) (Remark: The constant en is the best possible. See [1413].)

Fact 11.21.18. Let $p \in \mathbb{R}[s]$, and assume that p is discrete-time semistable. Then, $C(p)$ is discrete-time semistable, and there exists $v \in \mathbb{R}^n$ such that

$$\lim_{k \rightarrow \infty} C^k(p) = 1_{n \times 1} v^T.$$

(Proof: Since $C(p)$ is a companion form matrix, it follows from Proposition 11.10.4 that its minimal polynomial is p . Hence, $C(p)$ is discrete-time semistable. Now, it follows from Proposition 11.10.2 that $\lim_{k \rightarrow \infty} C^k(p)$ exists, and thus the state x_k of the difference equation $x_{k+1} = C(p)x_k$ converges for all initial conditions x_0 . The structure of $C(p)$ shows that all components of $\lim_{k \rightarrow \infty} x_k$ converge to the same value. Hence, all rows of $\lim_{k \rightarrow \infty} C^k(p)$ are equal.)

11.22 Facts on Lie Groups

Fact 11.22.1. The groups $\text{UT}(n)$, $\text{UT}_+(n)$, $\text{UT}_{\pm 1}(n)$, $\text{SUT}(n)$, and $\{I_n\}$ are Lie groups. Furthermore, $\text{ut}(n)$ is the Lie algebra of $\text{UT}(n)$, $\text{sut}(n)$ is the Lie algebra of $\text{SUT}(n)$, and $\{0_{n \times n}\}$ is the Lie algebra of $\{I_n\}$. (Remark: See Fact 3.21.4 and Fact 3.21.5.) (Problem: Determine the Lie algebras of $\text{UT}_+(n)$ and $\text{UT}_{\pm 1}(n)$.)

11.23 Facts on Subspace Decomposition

Fact 11.23.1. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$ is asymptotically stable, $A_{12} \in \mathbb{R}^{r \times (n-r)}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$. Then,

$$\mu_A^s(A) = S \begin{bmatrix} 0 & B_{12s} \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

where $B_{12s} \in \mathbb{R}^{r \times (n-r)}$, and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & B_{12u} \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

where $B_{12u} \in \mathbb{R}^{r \times (n-r)}$ and $\mu_A^u(A_1)$ is nonsingular. Consequently,

$$\mathcal{R}\left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix}\right) \subseteq \mathcal{S}_s(A).$$

If, in addition, $A_{12} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} 0 & 0 \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

$$\mathcal{S}_u(A) \subseteq \mathcal{R}\left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix}\right).$$

(Proof: The result follows from Fact 4.10.12.)

Fact 11.23.2. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$, $A_{12} \in \mathbb{R}^{r \times (n-r)}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ satisfies $\text{spec}(A_2) \subset \text{CRHP}$. Then,

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & C_{12s} \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

where $C_{12s} \in \mathbb{R}^{r \times (n-r)}$ and $\mu_A^s(A_2)$ is nonsingular, and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & C_{12u} \\ 0 & 0 \end{bmatrix} S^{-1},$$

where $C_{12u} \in \mathbb{R}^{r \times (n-r)}$. Consequently,

$$\mathcal{S}_s(A) \subseteq \mathcal{R}\left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix}\right).$$

If, in addition, $A_{12} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ 0 & 0 \end{bmatrix} S^{-1},$$

$$\mathcal{R}\left(S \begin{bmatrix} 0 & \\ & I_{n-r} \end{bmatrix}\right) \subseteq \mathcal{S}_u(A).$$

Fact 11.23.3. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$ satisfies $\text{spec}(A_1) \subset \text{CRHP}$, $A_{12} \in \mathbb{R}^{r \times (n-r)}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$. Then,

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & B_{12s} \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

where $\mu_A^s(A_1)$ is nonsingular and $B_{12s} \in \mathbb{R}^{r \times (n-r)}$, and

$$\mu_A^u(A) = S \begin{bmatrix} 0 & B_{12u} \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

where $B_{12u} \in \mathbb{R}^{r \times (n-r)}$. Consequently,

$$\mathcal{R}\left(S \begin{bmatrix} I_r & \\ & 0 \end{bmatrix}\right) \subseteq \mathcal{S}_u(A).$$

If, in addition, $A_{12} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

$$\mu_A^u(A) = S \begin{bmatrix} 0 & 0 \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

$$\mathcal{S}_s(A) \subseteq \mathcal{R}\left(S \begin{bmatrix} 0 & \\ & I_{n-r} \end{bmatrix}\right).$$

Fact 11.23.4. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$, $A_{12} \in \mathbb{R}^{r \times (n-r)}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ is asymptotically stable. Then,

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & C_{12s} \\ 0 & 0 \end{bmatrix} S^{-1},$$

where $C_{12s} \in \mathbb{R}^{r \times (n-r)}$, and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & C_{12u} \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

where $\mu_A^u(A_2)$ is nonsingular and $C_{12u} \in \mathbb{R}^{r \times (n-r)}$. Consequently,

$$\mathcal{S}_u(A) \subseteq \mathcal{R} \left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix} \right).$$

If, in addition, $A_{12} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ 0 & 0 \end{bmatrix} S^{-1},$$

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

$$\mathcal{R} \left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix} \right) \subseteq \mathcal{S}_s(A).$$

Fact 11.23.5. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$ satisfies $\text{spec}(A_1) \subset \text{CRHP}$, $A_{12} \in \mathbb{R}^{r \times (n-r)}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ is asymptotically stable. Then,

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & C_{12s} \\ 0 & 0 \end{bmatrix} S^{-1},$$

where $C_{12s} \in \mathbb{R}^{r \times (n-r)}$ and $\mu_A^s(A_1)$ is nonsingular, and

$$\mu_A^u(A) = S \begin{bmatrix} 0 & C_{12u} \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

where $C_{12u} \in \mathbb{R}^{r \times (n-r)}$ and $\mu_A^u(A_2)$ is nonsingular. Consequently,

$$\mathcal{S}_u(A) = \mathcal{R} \left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix} \right).$$

If, in addition, $A_{12} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ 0 & 0 \end{bmatrix} S^{-1}$$

and

$$\mu_A^u(A) = S \begin{bmatrix} 0 & 0 \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

Consequently,

$$\mathcal{S}_s(A) = \mathcal{R} \left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix} \right).$$

Fact 11.23.6. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$ is asymptotically stable, $A_{21} \in \mathbb{R}^{(n-r) \times r}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$. Then,

$$\mu_A^s(A) = S \begin{bmatrix} 0 & 0 \\ B_{21s} & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

where $B_{21s} \in \mathbb{R}^{(n-r) \times r}$, and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ B_{21u} & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

where $B_{21u} \in \mathbb{R}^{(n-r) \times r}$ and $\mu_A^u(A_1)$ is nonsingular. Consequently,

$$\mathcal{S}_u(A) \subseteq \mathcal{R}\left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix}\right).$$

If, in addition, $A_{21} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} 0 & 0 \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

$$\mathcal{R}\left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix}\right) \subseteq \mathcal{S}_s(A).$$

Fact 11.23.7. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$, $A_{21} \in \mathbb{R}^{(n-r) \times r}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ satisfies $\text{spec}(A_2) \subset \text{CRHP}$. Then,

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ C_{21s} & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

where $C_{21s} \in \mathbb{R}^{(n-r) \times r}$ and $\mu_A^s(A_2)$ is nonsingular, and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ C_{21u} & 0 \end{bmatrix} S^{-1},$$

where $C_{21u} \in \mathbb{R}^{(n-r) \times r}$. Consequently,

$$\mathcal{R}\left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix}\right) \subseteq \mathcal{S}_u(A).$$

If, in addition, $A_{21} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ 0 & 0 \end{bmatrix} S^{-1},$$

$$\mathfrak{S}_s(A) \subseteq \mathcal{R} \left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix} \right).$$

Fact 11.23.8. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$ is asymptotically stable, $A_{21} \in \mathbb{R}^{(n-r) \times r}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ satisfies $\text{spec}(A_2) \subset \text{CRHP}$. Then,

$$\mu_A^s(A) = S \begin{bmatrix} 0 & 0 \\ C_{21s} & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

where $C_{21s} \in \mathbb{R}^{n-r \times r}$ and $\mu_A^s(A_2)$ is nonsingular, and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ C_{21u} & 0 \end{bmatrix} S^{-1},$$

where $C_{21u} \in \mathbb{R}^{(n-r) \times r}$ and $\mu_A^u(A_1)$ is nonsingular. Consequently,

$$\mathfrak{S}_u(A) = \mathcal{R} \left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix} \right).$$

If, in addition, $A_{21} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} 0 & 0 \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1}$$

and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ 0 & 0 \end{bmatrix} S^{-1}.$$

Consequently,

$$\mathfrak{S}_s(A) = \mathcal{R} \left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix} \right).$$

Fact 11.23.9. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$, $A_{21} \in \mathbb{R}^{(n-r) \times r}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$ is asymptotically stable. Then,

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ B_{21s} & 0 \end{bmatrix} S^{-1},$$

where $B_{21s} \in \mathbb{R}^{(n-r) \times r}$, and

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ B_{21u} & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

where $B_{21u} \in \mathbb{R}^{(n-r) \times r}$ and $\mu_A^u(A_2)$ is nonsingular. Consequently,

$$\mathcal{R}\left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix}\right) \subseteq \mathcal{S}(A).$$

If, in addition, $A_{21} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ 0 & 0 \end{bmatrix} S^{-1},$$

$$\mu_A^u(A) = S \begin{bmatrix} \mu_A^u(A_1) & 0 \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

$$\mathcal{S}_u(A) \subseteq \mathcal{R}\left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix}\right).$$

Fact 11.23.10. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$ satisfies $\text{spec}(A_1) \subset \text{CRHP}$, $A_{21} \in \mathbb{R}^{(n-r) \times r}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$. Then,

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ C_{12s} & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

where $C_{12s} \in \mathbb{R}^{(n-r) \times r}$ and $\mu_A^s(A_1)$ is nonsingular, and

$$\mu_A^u(A) = S \begin{bmatrix} 0 & 0 \\ C_{21u} & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

where $C_{21u} \in \mathbb{R}^{(n-r) \times r}$. Consequently,

$$\mathcal{S}_s(A) \subseteq \mathcal{R}\left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix}\right).$$

If, in addition, $A_{21} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

$$\mu_A^u(A) = S \begin{bmatrix} 0 & 0 \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

$$\mathcal{R}\left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix}\right) \subseteq \mathcal{S}_u(A).$$

Fact 11.23.11. Let $A \in \mathbb{R}^{n \times n}$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \in \mathbb{R}^{r \times r}$ satisfies $\text{spec}(A_1) \subset \text{CRHP}$, $A_{21} \in \mathbb{R}^{(n-r) \times r}$, and $A_2 \in \mathbb{R}^{(n-r) \times (n-r)}$.

$\mathbb{R}^{(n-r) \times (n-r)}$ is asymptotically stable. Then,

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ C_{21s} & 0 \end{bmatrix} S^{-1},$$

where $C_{21s} \in \mathbb{R}^{(n-r) \times r}$ and $\mu_A^s(A_1)$ is nonsingular, and

$$\mu_A^u(A) = S \begin{bmatrix} 0 & 0 \\ C_{21u} & \mu_A^u(A_2) \end{bmatrix} S^{-1},$$

where $C_{21u} \in \mathbb{R}^{(n-r) \times r}$ and $\mu_A^u(A_2)$ is nonsingular. Consequently,

$$\mathfrak{S}_s(A) = \mathcal{R} \left(S \begin{bmatrix} 0 \\ I_{n-r} \end{bmatrix} \right).$$

If, in addition, $A_{21} = 0$, then

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & 0 \\ 0 & 0 \end{bmatrix} S^{-1}$$

and

$$\mu_A^u(A) = S \begin{bmatrix} 0 & 0 \\ 0 & \mu_A^u(A_2) \end{bmatrix} S^{-1}.$$

Consequently,

$$\mathfrak{S}_u(A) = \mathcal{R} \left(S \begin{bmatrix} I_r \\ 0 \end{bmatrix} \right).$$

11.24 Notes

The Laplace transform (11.2.10) is given in [1201, p. 34]. Computational methods are discussed in [683, 1015]. An arithmetic-mean-geometric-mean iteration for computing the matrix exponential and matrix logarithm is given in [1232].

The exponential function plays a central role in the theory of Lie groups, see [168, 295, 624, 724, 740, 1162, 1366]. Applications to robotics and kinematics are given in [986, 1026, 1070]. Additional applications are discussed in [294].

The real logarithm is discussed in [360, 664, 1048, 1102]. The multiplicity and properties of logarithms are discussed in [462].

An asymptotically stable polynomial is traditionally called *Hurwitz*. Semistability is defined in [283] and developed in [186, 195]. Stability theory is treated in [620, 885, 1094] and [541, Chapter XV]. Solutions of the Lyapunov equation under weak conditions are considered in [1207]. Structured solutions of the Lyapunov equation are discussed in [793].

Chapter Twelve

Linear Systems and Control Theory

This chapter considers linear state space systems with inputs and outputs. These systems are considered in both the time domain and frequency (Laplace) domain. Some basic results in control theory are also presented.

12.1 State Space and Transfer Function Models

Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$, and, for $t \geq t_0$, consider the *state equation*

$$\dot{x}(t) = Ax(t) + Bu(t), \quad (12.1.1)$$

with the *initial condition*

$$x(t_0) = x_0. \quad (12.1.2)$$

In (12.1.1), $x(t) \in \mathbb{R}^n$ is the *state*, and $u(t) \in \mathbb{R}^m$ is the *input*.

The following result give the solution of (12.1.1) known as the *variation of constants formula*.

Proposition 12.1.1. For $t \geq t_0$ the state $x(t)$ of the dynamical equation (12.1.1) with initial condition (12.1.2) is given by

$$x(t) = e^{(t-t_0)A}x_0 + \int_{t_0}^t e^{(t-\tau)A}Bu(\tau) \, d\tau. \quad (12.1.3)$$

Proof. Multiplying (12.1.1) by e^{-tA} yields

$$e^{-tA}[\dot{x}(t) - Ax(t)] = e^{-tA}Bu(t),$$

which is equivalent to

$$\frac{d}{dt}[e^{-tA}x(t)] = e^{-tA}Bu(t).$$

Integrating over $[t_0, t]$ yields

$$e^{-tA}x(t) = e^{-t_0A}x(t_0) + \int_{t_0}^t e^{-\tau A}Bu(\tau) \, d\tau.$$

Now, multiplying by e^{tA} yields (12.1.3).

Alternatively, let $x(t)$ be given by (12.1.3). Then, it follows from Leibniz's rule Fact 10.11.10 that

$$\begin{aligned}\dot{x}(t) &= \frac{d}{dt}e^{(t-t_0)A}x_0 + \frac{d}{dt}\int_{t_0}^t e^{(t-\tau)A}Bu(\tau) d\tau \\ &= Ae^{(t-t_0)A}x_0 + \int_{t_0}^t Ae^{(t-\tau)A}Bu(\tau) d\tau + Bu(t) \\ &= Ax(t) + Bu(t).\end{aligned}\quad \square$$

For convenience, we can reset the clock and assume without loss of generality that $t_0 = 0$. In this case, $x(t)$ for all $t \geq 0$ is given by

$$x(t) = e^{tA}x_0 + \int_0^t e^{(t-\tau)A}Bu(\tau) d\tau. \quad (12.1.4)$$

If $u(t) = 0$ for all $t \geq 0$, then, for all $t \geq 0$, $x(t)$ is given by

$$x(t) = e^{tA}x_0. \quad (12.1.5)$$

Now, let $u(t) = \delta(t)v$, where $\delta(t)$ is the *unit impulse* at $t = 0$ and $v \in \mathbb{R}^m$. Then, for all $t \geq 0$, $x(t)$ is given by

$$x(t) = e^{tA}x_0 + e^{tA}Bv. \quad (12.1.6)$$

Let $a < b$. Then, $\delta(t)$, which has physical dimensions of 1/time, satisfies

$$\int_a^b \delta(\tau) d\tau = \begin{cases} 0, & a > 0 \text{ or } b \leq 0, \\ 1, & a \leq 0 < b. \end{cases} \quad (12.1.7)$$

More generally, if $g: \mathcal{D} \rightarrow \mathbb{R}^n$, where $[a, b] \subseteq \mathcal{D} \subseteq \mathbb{R}$, $t_0 \in \mathcal{D}$, and g is continuous at t_0 , then

$$\int_a^b \delta(\tau - t_0)g(\tau) d\tau = \begin{cases} 0, & a > t_0 \text{ or } b \leq t_0, \\ g(t_0), & a \leq t_0 < b. \end{cases} \quad (12.1.8)$$

Alternatively, let the input $u(t)$ be constant or a *step function*, that is, $u(t) = v$ for all $t \geq 0$, where $v \in \mathbb{R}^m$. Then, by a change of variable of integration, it follows that, for all $t \geq 0$,

$$x(t) = e^{tA}x_0 + \int_0^t e^{\tau A} d\tau Bv. \quad (12.1.9)$$

Using Fact 11.13.14, (12.1.9) can be written for all $t \geq 0$ as

$$x(t) = e^{tA}x_0 + \left[A^D(e^{tA} - I) + (I - AA^D) \sum_{i=1}^{\text{ind } A} (i!)^{-1} t^i A^{i-1} \right] Bv. \quad (12.1.10)$$

If A is group invertible, then, for all $t \geq 0$, (12.1.10) becomes

$$x(t) = e^{tA}x_0 + [A^\#(e^{tA} - I) + t(I - AA^\#)]Bv. \quad (12.1.11)$$

If, in addition, A is nonsingular, then, for all $t \geq 0$, (12.1.11) becomes

$$x(t) = e^{tA}x_0 + A^{-1}(e^{tA} - I)Bv. \quad (12.1.12)$$

Next, consider the *output equation*

$$y(t) = Cx(t) + Du(t), \quad (12.1.13)$$

where $t \geq 0$, $y(t) \in \mathbb{R}^l$ is the *output*, $C \in \mathbb{R}^{l \times n}$, and $D \in \mathbb{R}^{l \times m}$. Then, for all $t \geq 0$, the *total response* is

$$y(t) = Ce^{tA}x_0 + \int_0^t Ce^{(t-\tau)A}Bu(\tau) d\tau + Du(t). \quad (12.1.14)$$

If $u(t) = 0$ for all $t \geq 0$, then the *free response* is given by

$$y(t) = Ce^{tA}x_0, \quad (12.1.15)$$

while, if $x_0 = 0$, then the *forced response* is given by

$$y(t) = \int_0^t Ce^{(t-\tau)A}Bu(\tau) d\tau + Du(t). \quad (12.1.16)$$

Setting $u(t) = \delta(t)v$ yields, for all $t > 0$, the total response

$$y(t) = Ce^{tA}x_0 + H(t)v, \quad (12.1.17)$$

where, for all $t \geq 0$, the *impulse response function* $H(t)$ is defined by

$$H(t) \triangleq Ce^{tA}B + \delta(t)D. \quad (12.1.18)$$

The corresponding forced response is the *impulse response*

$$y(t) = H(t)v = Ce^{tA}Bv + \delta(t)Dv. \quad (12.1.19)$$

Alternatively, if $u(t) = v$ for all $t \geq 0$, then the total response is

$$y(t) = Ce^{tA}x_0 + \int_0^t Ce^{\tau A} d\tau Bv + Dv, \quad (12.1.20)$$

and the forced response is the *step response*

$$y(t) = \int_0^t H(\tau) d\tau v = \int_0^t Ce^{\tau A} d\tau Bv + Dv. \quad (12.1.21)$$

In general, the forced response can be written as

$$y(t) = \int_0^t H(t-\tau)u(\tau) d\tau. \quad (12.1.22)$$

Setting $u(t) = \delta(t)v$ yields (12.1.20) by noting that

$$\int_0^t \delta(t - \tau) \delta(\tau) d\tau = \delta(t). \quad (12.1.23)$$

Proposition 12.1.2. Let $D = 0$ and $m = 1$, and assume that $x_0 = Bv$. Then, the free response and the impulse response are equal and given by

$$y(t) = Ce^{tA}x_0 = Ce^{tA}Bv. \quad (12.1.24)$$

12.2 Laplace Transform Analysis

Now, consider the linear system

$$\dot{x}(t) = Ax(t) + Bu(t), \quad (12.2.1)$$

$$y(t) = Cx(t) + Du(t), \quad (12.2.2)$$

with state $x(t) \in \mathbb{R}^n$, input $u(t) \in \mathbb{R}^m$, and output $y(t) \in \mathbb{R}^l$, where $t \geq 0$ and $x(0) = x_0$. Taking Laplace transforms yields

$$s\hat{x}(s) - x_0 = A\hat{x}(s) + B\hat{u}(s), \quad (12.2.3)$$

$$\hat{y}(s) = C\hat{x}(s) + D\hat{u}(s), \quad (12.2.4)$$

where

$$\hat{x}(s) \triangleq \mathcal{L}\{x(t)\} = \int_0^\infty e^{-st}x(t) dt, \quad (12.2.5)$$

$$\hat{u}(s) \triangleq \mathcal{L}\{u(t)\}, \quad (12.2.6)$$

and

$$\hat{y}(s) \triangleq \mathcal{L}\{y(t)\}. \quad (12.2.7)$$

Hence,

$$\hat{x}(s) = (sI - A)^{-1}x_0 + (sI - A)^{-1}B\hat{u}(s), \quad (12.2.8)$$

and thus

$$\hat{y}(s) = C(sI - A)^{-1}x_0 + [C(sI - A)^{-1}B + D]\hat{u}(s). \quad (12.2.9)$$

We can also obtain (12.2.9) from the time-domain expression for $y(t)$ given by (12.1.14). Using Proposition 11.2.2, it follows from (12.1.14) that

$$\begin{aligned} \hat{y}(s) &= \mathcal{L}\{Ce^{tA}x_0\} + \mathcal{L}\left\{\int_0^t Ce^{(t-\tau)A}Bu(\tau) d\tau\right\} + D\hat{u}(s) \\ &= C\mathcal{L}\{e^{tA}\}x_0 + C\mathcal{L}\{e^{tA}\}B\hat{u}(s) + D\hat{u}(s) \\ &= C(sI - A)^{-1}x_0 + [C(sI - A)^{-1}B + D]\hat{u}(s), \end{aligned} \quad (12.2.10)$$

which coincides with (12.2.9). We define

$$G(s) \triangleq C(sI - A)^{-1}B + D. \quad (12.2.11)$$

Note that $G \in \mathbb{R}^{l \times m}(s)$, that is, by Definition 4.7.2, G is a rational transfer function. Since $\mathcal{L}\{\delta(t)\} = 1$, it follows that

$$G(s) = \mathcal{L}\{H(t)\}. \quad (12.2.12)$$

Using (4.7.2), G can be written as

$$G(s) = \frac{1}{\chi_A(s)} C(sI - A)^A B + D. \quad (12.2.13)$$

It follows from (4.7.3) that G is a proper rational transfer function. Furthermore, G is a strictly proper rational transfer function if and only if $D = 0$, whereas G is an exactly proper rational transfer function if and only if $D \neq 0$. Finally, if A is nonsingular, then

$$G(0) = -CA^{-1}B + D. \quad (12.2.14)$$

Let $A \in \mathbb{R}^{n \times n}$. If $|s| > \text{sprad}(A)$, then Proposition 9.4.13 implies that

$$(sI - A)^{-1} = \frac{1}{s} \left(I - \frac{1}{s} A \right)^{-1} = \sum_{k=0}^{\infty} \frac{1}{s^{k+1}} A^k, \quad (12.2.15)$$

where the series is absolutely convergent, and thus

$$\begin{aligned} G(s) &= D + \frac{1}{s} CB + \frac{1}{s^2} CAB + \cdots \\ &= \sum_{k=0}^{\infty} \frac{1}{s^k} H_k, \end{aligned} \quad (12.2.16)$$

where, for $k \geq 0$, the *Markov parameter* $H_k \in \mathbb{R}^{l \times m}$ is defined by

$$H_k \triangleq \begin{cases} D, & k = 0, \\ CA^{k-1}B, & k \geq 1. \end{cases} \quad (12.2.17)$$

It follows from (12.2.15) that $\lim_{s \rightarrow \infty} (sI - A)^{-1} = 0$, and thus

$$\lim_{s \rightarrow \infty} G(s) = D. \quad (12.2.18)$$

Finally, it follows from Definition 4.7.3 that

$$\text{reldeg } G = \min\{k \geq 0: H_k \neq 0\}. \quad (12.2.19)$$

12.3 The Unobservable Subspace and Observability

Let $A \in \mathbb{R}^{n \times n}$ and $C \in \mathbb{R}^{l \times n}$, and, for $t \geq 0$, consider the linear system

$$\dot{x}(t) = Ax(t), \quad (12.3.1)$$

$$x(0) = x_0, \quad (12.3.2)$$

$$y(t) = Cx(t). \quad (12.3.3)$$

Definition 12.3.1. The *unobservable subspace* $\mathcal{U}_{t_f}(A, C)$ of (A, C) at time $t_f > 0$ is the subspace

$$\mathcal{U}_{t_f}(A, C) \triangleq \{x_0 \in \mathbb{R}^n: y(t) = 0 \text{ for all } t \in [0, t_f]\}. \quad (12.3.4)$$

Let $t_f > 0$. Then, Definition 12.3.1 states that $x_0 \in \mathcal{U}_{t_f}(A, C)$ if and only if $y(t) = 0$ for all $t \in [0, t_f]$. Since $y(t) = 0$ for all $t \in [0, t_f]$ is the free response corresponding to $x_0 = 0$, it follows that $0 \in \mathcal{U}_{t_f}(A, C)$. Now, suppose there exists a nonzero vector $x_0 \in \mathcal{U}_{t_f}(A, C)$. Then, with $x(0) = x_0$, the free response is given by $y(t) = 0$ for all $t \in [0, t_f]$, and thus x_0 cannot be determined from knowledge of $y(t)$ for all $t \in [0, t_f]$.

The following result provides explicit expressions for $\mathcal{U}_{t_f}(A, C)$.

Lemma 12.3.2. Let $t_f > 0$. Then, the following subspaces are equal:

- i) $\mathcal{U}_{t_f}(A, C)$.
- ii) $\bigcap_{t \in [0, t_f]} \mathcal{N}(Ce^{tA})$.
- iii) $\bigcap_{i=0}^{n-1} \mathcal{N}(CA^i)$.
- iv) $\mathcal{N}\left(\begin{bmatrix} C \\ CA \\ \vdots \\ CA^{n-1} \end{bmatrix}\right)$.
- v) $\mathcal{N}\left(\int_0^{t_f} e^{tA^T} C^T C e^{tA} dt\right)$.

If, in addition, $\lim_{t_f \rightarrow \infty} \int_0^{t_f} e^{tA^T} C^T C e^{tA} dt$ exists, then the following subspace is equal to i)–v):

$$vi) \mathcal{N}\left(\int_0^\infty e^{tA^T} C^T C e^{tA} dt\right).$$

Proof. The proof is dual to the proof of Lemma 12.6.2. □

Lemma 12.3.2 shows that $\mathcal{U}_{t_f}(A, C)$ is independent of t_f . We thus write $\mathcal{U}(A, C)$ for $\mathcal{U}_{t_f}(A, C)$, and call $\mathcal{U}(A, C)$ the *unobservable subspace* of (A, C) . (A, C) is *observable* if $\mathcal{U}(A, C) = \{0\}$. For convenience, define the $nl \times n$ *observability matrix*

$$\mathcal{O}(A, C) \triangleq \begin{bmatrix} C \\ CA \\ \vdots \\ CA^{n-1} \end{bmatrix} \quad (12.3.5)$$

so that

$$\mathcal{U}(A, C) = \mathcal{N}[\mathcal{O}(A, C)]. \quad (12.3.6)$$

Define

$$p \triangleq n - \dim \mathcal{U}(A, C) = n - \text{def } \mathcal{O}(A, C). \quad (12.3.7)$$

Corollary 12.3.3. For all $t_f > 0$,

$$p = \dim \mathcal{U}(A, C)^\perp = \text{rank } \mathcal{O}(A, C) = \text{rank} \int_0^{t_f} e^{tA^T} C^T C e^{tA} dt. \quad (12.3.8)$$

If, in addition, $\lim_{t_f \rightarrow \infty} \int_0^{t_f} e^{tA^T} C^T C e^{tA} dt$ exists, then

$$p = \text{rank} \int_0^\infty e^{tA^T} C^T C e^{tA} dt. \quad (12.3.9)$$

Corollary 12.3.4. $\mathcal{U}(A, C)$ is an invariant subspace of A .

The following result shows that the unobservable subspace $\mathcal{U}(A, C)$ is unchanged by output injection

$$\dot{x}(t) = Ax(t) + Fy(t). \quad (12.3.10)$$

Proposition 12.3.5. Let $F \in \mathbb{R}^{n \times l}$. Then,

$$\mathcal{U}(A + FC, C) = \mathcal{U}(A, C). \quad (12.3.11)$$

In particular, (A, C) is observable if and only if $(A + FC, C)$ is observable.

Proof. The proof is dual to the proof of Proposition 12.6.5. $\square$

Let $\tilde{\mathcal{U}}(A, C) \subseteq \mathbb{R}^n$ be a subspace that is complementary to $\mathcal{U}(A, C)$. Then, $\tilde{\mathcal{U}}(A, C)$ is an *observable subspace* in the sense that, if $x_0 = x'_0 + x''_0$, where $x'_0 \in \tilde{\mathcal{U}}(A, C)$ is nonzero and $x''_0 \in \mathcal{U}(A, C)$, then it is possible to determine x'_0 from knowledge of $y(t)$ for $t \in [0, t_f]$. Using Proposition 3.5.3, let $\mathcal{P} \in \mathbb{R}^{n \times n}$ be the unique idempotent matrix such that $\mathcal{R}(\mathcal{P}) = \tilde{\mathcal{U}}(A, C)$ and $\mathcal{N}(\mathcal{P}) = \mathcal{U}(A, C)$. Then, $x'_0 = \mathcal{P}x_0$. The following result constructs $\mathcal{P}$ and provides an expression for x'_0 in terms of $y(t)$ for $\tilde{\mathcal{U}}(A, C) \triangleq \mathcal{U}(A, C)^\perp$. In this case, $\mathcal{P}$ is a projector.

Lemma 12.3.6. Let $t_f > 0$, and define $\mathcal{P} \in \mathbb{R}^{n \times n}$ by

$$\mathcal{P} \triangleq \left(\int_0^{t_f} e^{tA^T} C^T C e^{tA} dt \right)^+ \int_0^{t_f} e^{tA^T} C^T C e^{tA} dt. \quad (12.3.12)$$

Then, $\mathcal{P}$ is the projector onto $\mathcal{U}(A, C)^\perp$, and $\mathcal{P}_\perp$ is the projector onto $\mathcal{U}(A, C)$. Hence,

$$\mathcal{R}(\mathcal{P}) = \mathcal{N}(\mathcal{P}_\perp) = \mathcal{U}(A, C)^\perp, \quad (12.3.13)$$

$$\mathcal{N}(\mathcal{P}) = \mathcal{R}(\mathcal{P}_\perp) = \mathcal{U}(A, C), \quad (12.3.14)$$

$$\text{rank } \mathcal{P} = \text{def } \mathcal{P}_\perp = \dim \mathcal{U}(A, C)^\perp = p, \quad (12.3.15)$$

$$\text{def } \mathcal{P} = \text{rank } \mathcal{P}_\perp = \dim \mathcal{U}(A, C) = n - p. \quad (12.3.16)$$

If $x_0 = x'_0 + x''_0$, where $x'_0 \in \mathcal{U}(A, C)^\perp$ and $x''_0 \in \mathcal{U}(A, C)$, then

$$x'_0 = \mathcal{P}x_0 = \left(\int_0^{t_f} e^{tA^T} C^T C e^{tA} dt \right)^+ \int_0^{t_f} e^{tA^T} C^T y(t) dt. \quad (12.3.17)$$

Finally, (A, C) is observable if and only if $\mathcal{P} = I_n$. In this case, for all $x_0 \in \mathbb{R}^n$,

$$x_0 = \left(\int_0^{t_f} e^{tA^T} C^T C e^{tA} dt \right)^{-1} \int_0^{t_f} e^{tA^T} C^T y(t) dt. \quad (12.3.18)$$

Lemma 12.3.7. Let $\alpha \in \mathbb{R}$. Then,

$$\mathcal{U}(A + \alpha I, C) = \mathcal{U}(A, C). \quad (12.3.19)$$

The following result uses a coordinate transformation to characterize the observable dynamics of a system.

Theorem 12.3.8. There exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that

$$A = S \begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix} S^{-1}, \quad C = [C_1 \quad 0] S^{-1}, \quad (12.3.20)$$

where $A_1 \in \mathbb{R}^{p \times p}$, $C_1 \in \mathbb{R}^{l \times p}$, and (A_1, C_1) is observable.

Proof. The proof is dual to the proof of Theorem 12.6.8. $\square$

Proposition 12.3.9. Let $S \in \mathbb{R}^{n \times n}$, and assume that S is orthogonal. Then, the following conditions are equivalent:

- i) A and C have the form (12.3.20), where $A_1 \in \mathbb{R}^{p \times p}$, $C_1 \in \mathbb{R}^{l \times p}$, and (A_1, C_1) is observable.
- ii) $\mathcal{U}(A, C) = \mathcal{R}(S \begin{bmatrix} 0 \\ I_{n-p} \end{bmatrix})$.
- iii) $\mathcal{U}(A, C)^\perp = \mathcal{R}(S \begin{bmatrix} I_p \\ 0 \end{bmatrix})$.
- iv) $\mathcal{P} = S \begin{bmatrix} I_p & 0 \\ 0 & 0 \end{bmatrix} S^T$.

Proposition 12.3.10. Let $S \in \mathbb{R}^{n \times n}$, and assume that S is nonsingular. Then, the following conditions are equivalent:

- i) A and C have the form (12.3.20), where $A_1 \in \mathbb{R}^{p \times p}$, $C_1 \in \mathbb{R}^{l \times p}$, and (A_1, C_1) is observable.
- ii) $\mathcal{U}(A, C) = \mathcal{R}(S \begin{bmatrix} 0 \\ I_{n-p} \end{bmatrix})$.
- iii) $\mathcal{U}(A, C)^\perp = \mathcal{R}(S^{-T} \begin{bmatrix} I_p \\ 0 \end{bmatrix})$.

Definition 12.3.11. Let $S \in \mathbb{R}^{n \times n}$, assume that S is nonsingular, and let A and C have the form (12.3.20), where $A_1 \in \mathbb{R}^{p \times p}$, $C_1 \in \mathbb{R}^{l \times p}$, and (A_1, C_1) is observable. Then, the *unobservable spectrum* of (A, C) is $\text{spec}(A_2)$, while the *unobservable*

multispectrum of (A, C) is $\text{mspec}(A_2)$. Furthermore, $\lambda \in \mathbb{C}$ is an *unobservable eigenvalue* of (A, C) if $\lambda \in \text{spec}(A_2)$.

Definition 12.3.12. The *observability pencil* $\mathcal{O}_{A,C}(s)$ is the pencil

$$\mathcal{O}_{A,C} = P \begin{bmatrix} A \\ -C \end{bmatrix}, [I]_0, \quad (12.3.21)$$

that is,

$$\mathcal{O}_{A,C}(s) = \begin{bmatrix} sI - A \\ C \end{bmatrix}. \quad (12.3.22)$$

Proposition 12.3.13. Let $\lambda \in \text{spec}(A)$. Then, λ is an unobservable eigenvalue of (A, C) if and only if

$$\text{rank} \begin{bmatrix} \lambda I - A \\ C \end{bmatrix} < n. \quad (12.3.23)$$

Proof. The proof is dual to the proof of Proposition 12.6.13. $\square$

Proposition 12.3.14. Let $\lambda \in \text{mspec}(A)$ and $F \in \mathbb{R}^{n \times m}$. Then, λ is an unobservable eigenvalue of (A, C) if and only if λ is an unobservable eigenvalue of $(A + FC, C)$.

Proof. The proof is dual to the proof of Proposition 12.6.14. $\square$

Proposition 12.3.15. Assume that (A, C) is observable. Then, the Smith form of $\mathcal{O}_{A,C}$ is $\begin{bmatrix} I_n \\ 0_{l \times n} \end{bmatrix}$.

Proof. The proof is dual to the proof of Proposition 12.6.15. $\square$

Proposition 12.3.16. Let $p_1, \dots, p_{n-p}$ be the similarity invariants of A_2 , where, for all $i = 1, \dots, n-p-1$, p_i divides p_{i+1} . Then, there exist unimodular matrices $S_1 \in \mathbb{R}^{(n+l) \times (n+l)}[s]$ and $S_2 \in \mathbb{R}^{n \times n}[s]$ and such that, for all $s \in \mathbb{C}$,

$$\begin{bmatrix} sI - A \\ C \end{bmatrix} = S_1(s) \begin{bmatrix} I_p \\ p_1(s) \\ \vdots \\ p_{n-p}(s) \\ 0_{l \times n} \end{bmatrix} S_2(s). \quad (12.3.24)$$

Consequently,

$$\text{Sz zeros}(\mathcal{O}_{A,C}) = \bigcup_{i=1}^{n-p} \text{roots}(p_i) = \text{roots}(\chi_{A_2}) = \text{spec}(A_2) \quad (12.3.25)$$

and

$$\text{mSz zeros}(\mathcal{O}_{A,C}) = \bigcup_{i=1}^{n-p} \text{mroots}(p_i) = \text{mroots}(\chi_{A_2}) = \text{mspec}(A_2). \quad (12.3.26)$$

Proof. The proof is dual to the proof of Proposition 12.6.16. $\square$

Proposition 12.3.17. Let $s \in \mathbb{C}$. Then,

$$\mathcal{O}(A, C) \subseteq \operatorname{Re} \mathcal{R} \left(\begin{bmatrix} sI - A \\ C \end{bmatrix} \right). \quad (12.3.27)$$

Proof. The proof is dual to the proof of Proposition 12.6.17. $\square$

The next result characterizes observability in several equivalent ways.

Theorem 12.3.18. The following statements are equivalent:

- i) (A, C) is observable.
- ii) There exists $t > 0$ such that $\int_0^t e^{\tau A^T} C^T C e^{\tau A} d\tau$ is positive definite.
- iii) $\int_0^t e^{\tau A^T} C^T C e^{\tau A} d\tau$ is positive definite for all $t > 0$.
- iv) $\operatorname{rank} \mathcal{O}(A, C) = n$.
- v) Every eigenvalue of (A, C) is observable.

If, in addition, $\lim_{t \rightarrow \infty} \int_0^t e^{\tau A^T} C^T C e^{\tau A} d\tau$ exists, then the following condition is equivalent to i)–v):

- vi) $\int_0^\infty e^{tA^T} C^T C e^{tA} dt$ is positive definite.

Proof. The proof is dual to the proof of Theorem 12.6.18. $\square$

The following result implies that arbitrary eigenvalue placement is possible for (12.3.10) when (A, C) is observable.

Proposition 12.3.19. The pair (A, C) is observable if and only if, for every polynomial $p \in \mathbb{R}[s]$ such that $\deg p = n$, there exists a matrix $F \in \mathbb{R}^{m \times n}$ such that $\operatorname{mspec}(A + FC) = \operatorname{mroots}(p)$.

Proof. The proof is dual to the proof of Proposition 12.6.19. $\square$

12.4 Observable Asymptotic Stability

Let $A \in \mathbb{R}^{n \times n}$ and $C \in \mathbb{R}^{l \times n}$, and define $p \triangleq n - \dim \mathcal{U}(A, C)$.

Definition 12.4.1. (A, C) is *observably asymptotically stable* if

$$\mathcal{S}_u(A) \subseteq \mathcal{U}(A, C). \quad (12.4.1)$$

Proposition 12.4.2. Let $F \in \mathbb{R}^{n \times l}$. Then, (A, C) is observably asymptotically stable if and only if $(A + FC, C)$ is observably asymptotically stable.

Proposition 12.4.3. The following statements are equivalent:

- i) (A, C) is observably asymptotically stable.
- ii) There exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that (12.3.20) holds,

where $A_1 \in \mathbb{R}^{p \times p}$ is asymptotically stable and $C_1 \in \mathbb{R}^{l \times p}$.

iii) There exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.3.20) holds, where $A_1 \in \mathbb{R}^{p \times p}$ is asymptotically stable and $C_1 \in \mathbb{R}^{l \times p}$.

iv) $\lim_{t \rightarrow \infty} C e^{tA} = 0$.

v) The positive-semidefinite matrix $P \in \mathbb{R}^{n \times n}$ defined by

$$P \triangleq \int_0^\infty e^{tA^T} C^T C e^{tA} dt \quad (12.4.2)$$

exists.

vi) There exists a positive-semidefinite matrix $P \in \mathbb{R}^{n \times n}$ satisfying

$$A^T P + P A + C^T C = 0. \quad (12.4.3)$$

In this case, the positive-semidefinite matrix $P \in \mathbb{R}^{n \times n}$ defined by (12.4.2) satisfies (12.4.3).

Proof. The proof is dual to the proof of Proposition 12.7.3. $\square$

The matrix P defined by (12.4.2) is the *observability Gramian*, and (12.4.3) is the *observation Lyapunov equation*.

Proposition 12.4.4. Assume that (A, C) is observably asymptotically stable, let $P \in \mathbb{R}^{n \times n}$ be the positive-semidefinite matrix defined by (12.4.2), and define $\mathcal{P} \in \mathbb{R}^{n \times n}$ by (12.3.12). Then, the following statements hold:

i) $PP^+ = \mathcal{P}$.

ii) $\mathcal{R}(P) = \mathcal{R}(\mathcal{P}) = \mathcal{U}(A, C)^\perp$.

iii) $\mathcal{N}(P) = \mathcal{N}(\mathcal{P}) = \mathcal{U}(A, C)$.

iv) $\text{rank } P = \text{rank } \mathcal{P} = p$.

v) P is the only positive-semidefinite solution of (12.4.3) whose rank is p .

Proof. The proof is dual to the proof of Proposition 12.7.4. $\square$

Proposition 12.4.5. Assume that (A, C) is observably asymptotically stable, let $P \in \mathbb{R}^{n \times n}$ be the positive-semidefinite matrix defined by (12.4.2), and let $\hat{P} \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

i) $\hat{P}$ is positive semidefinite and satisfies (12.4.3).

ii) There exists a positive-semidefinite matrix $P_0 \in \mathbb{R}^{n \times n}$ such that $\hat{P} = P + P_0$ and $A^T P_0 + P_0 A = 0$.

In this case,

$$\text{rank } \hat{P} = p + \text{rank } P_0 \quad (12.4.4)$$

and

$$\text{rank } P_0 \leq \sum_{\substack{\lambda \in \text{spec}(A) \\ \lambda \in j\mathbb{R}}} \text{gmult}_A(\lambda). \quad (12.4.5)$$

Proof. The proof is dual to the proof of Proposition 12.7.5. $\square$

Proposition 12.4.6. The following statements are equivalent:

- i) (A, C) is observably asymptotically stable, every imaginary eigenvalue of A is semisimple, and A has no ORHP eigenvalues.
- ii) (12.4.3) has a positive-definite solution $P \in \mathbb{R}^{n \times n}$.

Proof. The proof is dual to the proof of Proposition 12.7.6. $\square$

Proposition 12.4.7. The following statements are equivalent:

- i) (A, C) is observably asymptotically stable, and A has no imaginary eigenvalues.
- ii) (12.4.3) has exactly one positive-semidefinite solution $P \in \mathbb{R}^{n \times n}$.

In this case, $P \in \mathbb{R}^{n \times n}$ is given by (12.4.2) and satisfies $\text{rank } P = p$.

Proof. The proof is dual to the proof of Proposition 12.7.7. $\square$

Corollary 12.4.8. Assume that A is asymptotically stable. Then, the positive-semidefinite matrix $P \in \mathbb{R}^{n \times n}$ defined by (12.4.2) is the unique solution of (12.4.3) and satisfies $\text{rank } P = p$.

Proof. The proof is dual to the proof of Corollary 12.7.4. $\square$

Proposition 12.4.9. The following statements are equivalent:

- i) (A, C) is observable, and A is asymptotically stable.
- ii) (12.4.3) has exactly one positive-semidefinite solution $P \in \mathbb{R}^{n \times n}$, and P is positive definite.

In this case, $P \in \mathbb{R}^{n \times n}$ is given by (12.4.2).

Proof. The proof is dual to the proof of Proposition 12.7.9. $\square$

Corollary 12.4.10. Assume that A is asymptotically stable. Then, the positive-semidefinite matrix $P \in \mathbb{R}^{n \times n}$ defined by (12.4.2) exists. Furthermore, P is positive definite if and only if (A, C) is observable.

12.5 Detectability

Let $A \in \mathbb{R}^{n \times n}$ and $C \in \mathbb{R}^{l \times n}$, and define $p \triangleq n - \dim \mathcal{U}(A, C)$.

Definition 12.5.1. (A, C) is *detectable* if

$$\mathcal{U}(A, C) \subseteq \mathcal{S}_s(A). \quad (12.5.1)$$

Proposition 12.5.2. Let $F \in \mathbb{R}^{n \times l}$. Then, (A, C) is detectable if and only if $(A + FC, C)$ is detectable.

Proposition 12.5.3. The following statements are equivalent:

- i) A is asymptotically stable.
- ii) (A, C) is detectable and observably asymptotically stable.

Proof. The proof is dual to the proof of Proposition 12.8.3. □

Proposition 12.5.4. The following statements are equivalent:

- i) (A, C) is detectable.
- ii) There exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that (12.3.20) holds, where $A_1 \in \mathbb{R}^{p \times p}$, $C_1 \in \mathbb{R}^{l \times p}$, (A_1, C_1) is observable, and $A_2 \in \mathbb{R}^{(n-p) \times (n-p)}$ is asymptotically stable.
- iii) There exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.3.20) holds, where $A_1 \in \mathbb{R}^{p \times p}$, $C_1 \in \mathbb{R}^{l \times p}$, (A_1, C_1) is observable, and $A_2 \in \mathbb{R}^{(n-p) \times (n-p)}$ is asymptotically stable.
- iv) Every CRHP eigenvalue of (A, C) is observable.

Proof. The proof is dual to the proof of Proposition 12.8.4. □

Proposition 12.5.5. The following statements are equivalent:

- i) (A, C) is observably asymptotically stable and detectable.
- ii) A is asymptotically stable.

Proof. The proof is dual to the proof of Proposition 12.8.5. □

Corollary 12.5.6. The following statements are equivalent:

- i) There exists a positive-semidefinite matrix $P \in \mathbb{R}^{n \times n}$ satisfying (12.4.3), and (A, C) is detectable.
- ii) A is asymptotically stable.

Proof. The proof is dual to the proof of Proposition 12.8.6. □

12.6 The Controllable Subspace and Controllability

Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$, and, for $t \geq 0$, consider the linear system

$$\dot{x}(t) = Ax(t) + Bu(t), \quad (12.6.1)$$

$$x(0) = 0. \quad (12.6.2)$$

Definition 12.6.1. The *controllable subspace* $\mathcal{C}_{t_f}(A, B)$ of (A, B) at time $t_f > 0$ is the subspace

$\mathcal{C}_{t_f}(A, B) \triangleq \{x_f \in \mathbb{R}^n: \text{ there exists a continuous control } u: [0, t_f] \mapsto \mathbb{R}^m$
 such that the solution $x(\cdot)$ of (12.6.1), (12.6.2) satisfies $x(t_f) = x_f\}$. (12.6.3)

Let $t_f > 0$. Then, Definition 12.6.1 states that $x_f \in \mathcal{C}_{t_f}(A, B)$ if and only if there exists a continuous control $u: [0, t_f] \mapsto \mathbb{R}^m$ such that

$$x_f = \int_0^{t_f} e^{(t_f-t)A} B u(t) dt. \quad (12.6.4)$$

The following result provides explicit expressions for $\mathcal{C}_{t_f}(A, B)$.

Lemma 12.6.2. Let $t_f > 0$. Then, the following subspaces are equal:

- i) $\mathcal{C}_{t_f}(A, B)$.
- ii) $\left[\bigcap_{t \in [0, t_f]} \mathcal{N}(B^T e^{tA^T}) \right]^\perp$.
- iii) $\left[\bigcap_{i=0}^{n-1} \mathcal{N}(B^T A^i T) \right]^\perp$.
- iv) $\mathcal{R} \left(\begin{bmatrix} B & AB & \cdots & A^{n-1}B \end{bmatrix} \right)$.
- v) $\mathcal{R} \left(\int_0^{t_f} e^{tA} B B^T e^{tA^T} dt \right)$.

If, in addition, $\lim_{t_f \rightarrow \infty} \int_0^{t_f} e^{tA} B B^T e^{tA^T} dt$ exists, then the following subspace is equal to i)–v):

$$vi) \mathcal{R} \left(\int_0^\infty e^{tA} B B^T e^{tA^T} dt \right).$$

Proof. To prove that i) $\subseteq$ ii), let $\eta \in \bigcap_{t \in [0, t_f]} \mathcal{N}(B^T e^{tA^T})$ so that $\eta^T e^{tA} B = 0$ for all $t \in [0, t_f]$. Now, let $u: [0, t_f] \mapsto \mathbb{R}^m$ be continuous. Then, $\eta^T \int_0^{t_f} e^{(t_f-t)A} B u(t) dt = 0$, which implies that $\eta \in \mathcal{C}_{t_f}(A, B)^\perp$.

To prove that ii) $\subseteq$ iii), let $\eta \in \bigcap_{i=0}^{n-1} \mathcal{N}(B^T A^i T)$ so that $\eta^T A^i B = 0$ for all $i = 0, 1, \dots, n-1$. It follows from the Cayley-Hamilton theorem Theorem 4.4.7 that $\eta^T A^i B = 0$ for all $i \geq 0$. Now, let $t \in [0, t_f]$. Then, $\eta^T e^{tA} B = \sum_{i=0}^\infty t^i (i!)^{-1} \eta^T A^i B = 0$, and thus $\eta \in \mathcal{N}(B^T e^{tA^T})$.

To show that iii) $\subseteq$ iv), let $\eta \in \mathcal{R} \left(\begin{bmatrix} B & AB & \cdots & A^{n-1}B \end{bmatrix} \right)^\perp$. Then, $\eta \in \mathcal{N} \left(\begin{bmatrix} B & AB & \cdots & A^{n-1}B \end{bmatrix}^T \right)$, which implies that $\eta^T A^i B = 0$ for all $i = 0, 1, \dots, n-1$.

To prove that iv) $\subseteq$ v), let $\eta \in \mathcal{N} \left(\int_0^{t_f} e^{tA} B B^T e^{tA^T} dt \right)$. Then,

$$\eta^T \int_0^{t_f} e^{tA} B B^T e^{tA^T} dt \eta = 0,$$

which implies that $\eta^T e^{tA} B = 0$ for all $t \in [0, t_f]$. Differentiating with respect to t and setting $t = 0$ implies that $\eta^T A^i B = 0$ for all $i = 0, 1, \dots, n-1$. Hence, $\eta \in \mathcal{R}([B \ AB \ \dots \ A^{n-1}B])^\perp$.

To prove that $v) \subseteq i)$, let $\eta \in \mathcal{C}_{t_f}(A, B)^\perp$. Then, $\eta^T \int_0^{t_f} e^{(t_f-t)A} B u(t) dt = 0$ for all continuous $u: [0, t_f] \mapsto \mathbb{R}^m$. Letting $u(t) = B^T e^{(t_f-t)A^T} \eta^T$, implies that $\eta^T \int_0^{t_f} e^{tA} B B^T e^{tA^T} dt \eta = 0$, and thus $\eta \in \mathcal{N}(\int_0^{t_f} e^{tA} B B^T e^{tA^T} dt)$. $\square$

Lemma 12.6.2 shows that $\mathcal{C}_{t_f}(A, B)$ is independent of t_f . We thus write $\mathcal{C}(A, B)$ for $\mathcal{C}_{t_f}(A, B)$, and call $\mathcal{C}(A, B)$ the *controllable subspace* of (A, B) . (A, B) is *controllable* if $\mathcal{C}(A, B) = \mathbb{R}^n$. For convenience, define the $m \times nm$ *controllability matrix*

$$\mathcal{K}(A, B) \triangleq [B \ AB \ \dots \ A^{n-1}B] \quad (12.6.5)$$

so that

$$\mathcal{C}(A, B) = \mathcal{R}[\mathcal{K}(A, B)]. \quad (12.6.6)$$

Define

$$q \triangleq \dim \mathcal{C}(A, B) = \text{rank } \mathcal{K}(A, B). \quad (12.6.7)$$

Corollary 12.6.3. For all $t_f > 0$,

$$q = \dim \mathcal{C}(A, B) = \text{rank } \mathcal{K}(A, B) = \text{rank} \int_0^{t_f} e^{tA} B B^T e^{tA^T} dt. \quad (12.6.8)$$

If, in addition, $\lim_{t_f \rightarrow \infty} \int_0^{t_f} e^{tA} B B^T e^{tA^T} dt$ exists, then

$$q = \text{rank} \int_0^\infty e^{tA} B B^T e^{tA^T} dt. \quad (12.6.9)$$

Corollary 12.6.4. $\mathcal{C}(A, B)$ is an invariant subspace of A .

The following result shows that the controllable subspace $\mathcal{C}(A, B)$ is unchanged by full-state feedback $u(t) = Kx(t) + v(t)$.

Proposition 12.6.5. Let $K \in \mathbb{R}^{m \times n}$. Then,

$$\mathcal{C}(A + BK, B) = \mathcal{C}(A, B). \quad (12.6.10)$$

In particular, (A, B) is controllable if and only if $(A + BK, B)$ is controllable.

Proof. Note that

$$\begin{aligned} \mathcal{C}(A + BK, B) &= \mathcal{R}[\mathcal{K}(A + BK, B)] \\ &= \mathcal{R}([B \ AB + BKB \ A^2B + ABKB + BKAB + BKBKB \ \dots]) \\ &= \mathcal{R}[\mathcal{K}(A, B)] = \mathcal{C}(A, B). \end{aligned} \quad \square$$

Let $\tilde{\mathcal{C}}(A, B) \subseteq \mathbb{R}^n$ be a subspace that is complementary to $\mathcal{C}(A, B)$. Then, $\tilde{\mathcal{C}}(A, B)$ is an *uncontrollable subspace* in the sense that, if $x_f = x'_f + x''_f \in \mathbb{R}^n$, where $x'_f \in \mathcal{C}(A, B)$ and $x''_f \in \tilde{\mathcal{C}}(A, B)$ is nonzero, then there exists a continuous control $u: [0, t_f] \rightarrow \mathbb{R}^m$ such that $x(t_f) = x'_f$, but there exists no continuous control such that $x(t_f) = x_f$. Using Proposition 3.5.3, let $\mathcal{Q} \in \mathbb{R}^{n \times n}$ be the unique idempotent matrix such that $\mathcal{R}(\mathcal{Q}) = \mathcal{C}(A, B)$ and $\mathcal{N}(\mathcal{Q}) = \tilde{\mathcal{C}}(A, B)$. Then, $x'_f = \mathcal{Q}x_f$. The following result constructs $\mathcal{Q}$ and a continuous control $u(\cdot)$ that yields $x(t_f) = x'_f$ for $\tilde{\mathcal{C}}(A, B) \triangleq \mathcal{C}(A, B)^\perp$. In this case, $\mathcal{Q}$ is a projector.

Lemma 12.6.6. Let $t_f > 0$, and define $\mathcal{Q} \in \mathbb{R}^{n \times n}$ by

$$\mathcal{Q} \triangleq \left(\int_0^{t_f} e^{tA} B B^T e^{tA^T} dt \right)^+ \int_0^{t_f} e^{tA} B B^T e^{tA^T} dt. \quad (12.6.11)$$

Then, $\mathcal{Q}$ is the projector onto $\mathcal{C}(A, B)$, and $\mathcal{Q}_\perp$ is the projector onto $\mathcal{C}(A, B)^\perp$. Hence,

$$\mathcal{R}(\mathcal{Q}) = \mathcal{N}(\mathcal{Q}_\perp) = \mathcal{C}(A, B), \quad (12.6.12)$$

$$\mathcal{N}(\mathcal{Q}) = \mathcal{R}(\mathcal{Q}) = \mathcal{C}(A, B)^\perp, \quad (12.6.13)$$

$$\text{rank } \mathcal{Q} = \text{def } \mathcal{Q}_\perp = \dim \mathcal{C}(A, B) = q, \quad (12.6.14)$$

$$\text{def } \mathcal{Q} = \text{rank } \mathcal{Q}_\perp = \dim \mathcal{C}(A, B)^\perp = n - q. \quad (12.6.15)$$

Now, define $u: [0, t_f] \mapsto \mathbb{R}^m$ by

$$u(t) \triangleq B^T e^{(t_f-t)A^T} \left(\int_0^{t_f} e^{\tau A} B B^T e^{\tau A^T} d\tau \right)^+ x_f. \quad (12.6.16)$$

If $x_f = x'_f + x''_f$, where $x'_f \in \mathcal{C}(A, B)$ and $x''_f \in \mathcal{C}(A, B)^\perp$, then

$$x'_f = \mathcal{Q}x_f = \int_0^{t_f} e^{(t_f-t)A} B u(t) dt. \quad (12.6.17)$$

Finally, (A, B) is controllable if and only if $\mathcal{Q} = I_n$. In this case, for all $x_f \in \mathbb{R}^n$,

$$x_f = \int_0^{t_f} e^{(t_f-t)A} B u(t) dt, \quad (12.6.18)$$

where $u: [0, t_f] \mapsto \mathbb{R}^m$ is given by

$$u(t) = B^T e^{(t_f-t)A^T} \left(\int_0^{t_f} e^{\tau A} B B^T e^{\tau A^T} d\tau \right)^{-1} x_f. \quad (12.6.19)$$

Lemma 12.6.7. Let $\alpha \in \mathbb{R}$. Then,

$$\mathcal{C}(A + \alpha I, B) = \mathcal{C}(A, B). \quad (12.6.20)$$

The following result uses a coordinate transformation to characterize the controllable dynamics of (12.6.1).

Theorem 12.6.8. There exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that

$$A = S \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix} S^{-1}, \quad B = S \begin{bmatrix} B_1 \\ 0 \end{bmatrix}, \quad (12.6.21)$$

where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, and (A_1, B_1) is controllable.

Proof. Let $\alpha < 0$ be such that $A_\alpha \triangleq A + \alpha I$ is asymptotically stable, and let $Q \in \mathbb{R}^{n \times n}$ be the positive-semidefinite solution of

$$A_\alpha Q + Q A_\alpha^T + B B^T = 0 \quad (12.6.22)$$

given by

$$Q = \int_0^\infty e^{t A_\alpha} B B^T e^{t A_\alpha^T} dt.$$

It now follows from Lemma 12.6.2 and Lemma 12.6.7 that

$$\mathcal{R}(Q) = \mathcal{R}[\mathcal{C}(A_\alpha, B)] = \mathcal{R}[\mathcal{C}(A, B)].$$

Hence,

$$\text{rank } Q = \dim \mathcal{C}(A_\alpha, B) = \dim \mathcal{C}(A, B) = q.$$

Next, let $S \in \mathbb{R}^{n \times n}$ be an orthogonal matrix such that $Q = S \begin{bmatrix} Q_1 & 0 \\ 0 & 0 \end{bmatrix} S^T$, where $Q_1 \in \mathbb{R}^{q \times q}$ is positive definite. Writing $A_\alpha = S \begin{bmatrix} \hat{A}_1 & \hat{A}_{12} \\ \hat{A}_{21} & \hat{A}_2 \end{bmatrix} S^{-1}$ and $B = S \begin{bmatrix} B_1 \\ B_2 \end{bmatrix}$, where $\hat{A}_1 \in \mathbb{R}^{q \times q}$ and $B_1 \in \mathbb{R}^{q \times m}$, it follows from (12.6.22) that

$$\begin{aligned} \hat{A}_1 Q_1 + Q_1 \hat{A}_1^T + B_1 B_1^T &= 0, \\ \hat{A}_{21} Q_1 + B_2 B_1^T &= 0, \\ B_2 B_2^T &= 0. \end{aligned}$$

Therefore, $B_2 = 0$ and $\hat{A}_{21} = 0$, and thus

$$A_\alpha = S \begin{bmatrix} \hat{A}_1 & \hat{A}_{12} \\ 0 & \hat{A}_2 \end{bmatrix} S^{-1}, \quad B = S \begin{bmatrix} B_1 \\ 0 \end{bmatrix}.$$

Furthermore,

$$A = S \begin{bmatrix} \hat{A}_1 & \hat{A}_{12} \\ 0 & \hat{A}_2 \end{bmatrix} S^{-1} - \alpha I = S \left(\begin{bmatrix} \hat{A}_1 & \hat{A}_{12} \\ 0 & \hat{A}_2 \end{bmatrix} - \alpha I \right) S^{-1}.$$

Hence,

$$A = S \begin{bmatrix} A_1 & A_{12} \\ 0 & A_2 \end{bmatrix} S^{-1},$$

where $A_1 \triangleq \hat{A}_1 - \alpha I_q$, $A_{12} \triangleq \hat{A}_{12}$, and $A_2 \triangleq \hat{A}_2 - \alpha I_{n-q}$. □

Proposition 12.6.9. Let $S \in \mathbb{R}^{n \times n}$, and assume that S is orthogonal. Then, the following conditions are equivalent:

- i) A and B have the form (12.6.21), where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, and (A_1, B_1) is controllable.
- ii) $\mathcal{C}(A, B) = \mathcal{R}(S \begin{bmatrix} I_q \\ 0 \end{bmatrix})$.

$$iii) \mathcal{C}(A, B)^\perp = \mathcal{R}(S \begin{bmatrix} 0 \\ I_{n-q} \end{bmatrix}).$$

$$iv) \mathcal{Q} = S \begin{bmatrix} I_q & 0 \\ 0 & 0 \end{bmatrix} S^T.$$

Proposition 12.6.10. Let $S \in \mathbb{R}^{n \times n}$, and assume that S is nonsingular. Then, the following conditions are equivalent:

- i) A and B have the form (12.6.21), where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, and (A_1, B_1) is controllable.
- ii) $\mathcal{C}(A, B) = \mathcal{R}(S \begin{bmatrix} I_q \\ 0 \end{bmatrix})$.
- iii) $\mathcal{C}(A, B)^\perp = \mathcal{R}(S^{-T} \begin{bmatrix} 0 \\ I_{n-q} \end{bmatrix})$.

Definition 12.6.11. Let $S \in \mathbb{R}^{n \times n}$, assume that S is nonsingular, and let A and B have the form (12.6.21), where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, and (A_1, B_1) is controllable. Then, the *uncontrollable spectrum* of (A, B) is $\text{spec}(A_2)$, while the *uncontrollable multispectrum* of (A, B) is $\text{mspec}(A_2)$. Furthermore, $\lambda \in \mathbb{C}$ is an *uncontrollable eigenvalue* of (A, B) if $\lambda \in \text{spec}(A_2)$.

Definition 12.6.12. The *controllability pencil* $\mathcal{C}_{A,B}(s)$ is the pencil

$$\mathcal{C}_{A,B} = P_{[A \ -B], [I \ 0]}, \quad (12.6.23)$$

that is,

$$\mathcal{C}_{A,B}(s) = \begin{bmatrix} sI - A & B \end{bmatrix}. \quad (12.6.24)$$

Proposition 12.6.13. Let $\lambda \in \text{spec}(A)$. Then, λ is an uncontrollable eigenvalue of (A, B) if and only if

$$\text{rank} \begin{bmatrix} \lambda I - A & B \end{bmatrix} < n. \quad (12.6.25)$$

Proof. Since (A_1, B_1) is controllable, it follows from (12.6.21) that

$$\begin{aligned} \text{rank} \begin{bmatrix} \lambda I - A & B \end{bmatrix} &= \text{rank} \begin{bmatrix} \lambda I - A_1 & A_{12} & B_1 \\ 0 & \lambda I - A_2 & 0 \end{bmatrix} \\ &= \text{rank} \begin{bmatrix} \lambda I - A_1 & B_1 \end{bmatrix} + \text{rank}(\lambda I - A_2) \\ &= q + \text{rank}(\lambda I - A_2). \end{aligned}$$

Hence, $\text{rank} \begin{bmatrix} \lambda I - A & B \end{bmatrix} < n$ if and only if $\text{rank}(\lambda I - A_2) < n - q$, that is, if and only if $\lambda \in \text{spec}(A_2)$. $\square$

Proposition 12.6.14. Let $\lambda \in \text{mspec}(A)$ and $K \in \mathbb{R}^{n \times m}$. Then, λ is an uncontrollable eigenvalue of (A, B) if and only if λ is an uncontrollable eigenvalue of $(A + BK, B)$.

Proof. In the notation of Theorem 12.6.8, partition $B_1 = \begin{bmatrix} B_{11} & B_{12} \end{bmatrix}$, where $B_{11} \in \mathbb{R}^{q \times m}$, and partition $K = \begin{bmatrix} K_1 \\ K_2 \end{bmatrix}$, where $K_1 \in \mathbb{R}^{q \times m}$. Then,

$$A + BK = \begin{bmatrix} A_1 + B_{11}K_1 & A_{12} + B_{12}K_2 \\ 0 & A_2 \end{bmatrix}.$$

Consequently, the uncontrollable spectrum of $A + BK$ is $\text{spec}(A_2)$.

Proposition 12.6.15. Assume that (A, B) is controllable. Then, the Smith form of $\mathcal{C}_{A,B}$ is $\begin{bmatrix} I_n & 0_{n \times m} \end{bmatrix}$.

Proof. First, note that, if $\lambda \in \mathbb{C}$ is not an eigenvalue of A , then $n = \text{rank}(\lambda I - A) = \text{rank} \begin{bmatrix} \lambda I - A & B \end{bmatrix} = \text{rank } \mathcal{C}_{A,B}(\lambda)$. Therefore, $\text{rank } \mathcal{C}_{A,B} = n$, and thus $\mathcal{C}_{A,B}$ has n Smith polynomials. Furthermore, since (A, B) is controllable, it follows that (A, B) has no uncontrollable eigenvalues. Therefore, it follows from Proposition 12.6.13 that, for all $\lambda \in \text{spec}(A)$, $\text{rank} \begin{bmatrix} \lambda I - A & B \end{bmatrix} = n$. Consequently, $\text{rank } \mathcal{C}_{A,B}(\lambda) = n$ for all $\lambda \in \mathbb{C}$. Thus, every Smith polynomial $\mathcal{C}_{A,B}$ is 1. $\square$

Proposition 12.6.16. Let $p_1, \dots, p_{n-q}$ be the similarity invariants of A_2 , where, for all $i = 1, \dots, n - q - 1$, p_i divides p_{i+1} . Then, there exist unimodular matrices $S_1 \in \mathbb{R}^{n \times n}[s]$ and $S_2 \in \mathbb{R}^{(n+m) \times (n+m)}[s]$ such that, for all $s \in \mathbb{C}$,

$$\begin{bmatrix} sI - A & B \end{bmatrix} = S_1(s) \begin{bmatrix} I_q & & & \\ & p_1(s) & & \\ & & \ddots & \\ & & & p_{n-q}(s) \end{bmatrix} \begin{bmatrix} 0_{n \times m} \\ S_2(s) \end{bmatrix}. \quad (12.6.26)$$

Consequently,

$$\text{Szeros}(\mathcal{C}_{A,B}) = \bigcup_{i=1}^{n-q} \text{roots}(p_i) = \text{roots}(\chi_{A_2}) = \text{spec}(A_2) \quad (12.6.27)$$

and

$$\text{mSzeros}(\mathcal{C}_{A,B}) = \bigcup_{i=1}^{n-q} \text{mroots}(p_i) = \text{mroots}(\chi_{A_2}) = \text{mspec}(A_2). \quad (12.6.28)$$

Proof. Let $S \in \mathbb{R}^{n \times n}$ be as in Theorem 12.6.8, let $\hat{S}_1 \in \mathbb{R}^{q \times q}[s]$ and $\hat{S}_2 \in \mathbb{R}^{(q+m) \times (q+m)}[s]$ be unimodular matrices such that

$$\hat{S}_1(s) \begin{bmatrix} sI_q - A_1 & B_1 \end{bmatrix} \hat{S}_2(s) = \begin{bmatrix} I_q & 0_{q \times m} \end{bmatrix},$$

and let $\hat{S}_3, \hat{S}_4 \in \mathbb{R}^{(n-q) \times (n-q)}$ be unimodular matrices such that

$$\hat{S}_3(s)(sI - A_2)\hat{S}_4(s) = \hat{P}(s),$$

where $\hat{P} \triangleq \text{diag}(p_1, \dots, p_{n-q})$. Then,

$$\begin{aligned} \begin{bmatrix} sI - A & B \end{bmatrix} &= S \begin{bmatrix} \hat{S}_1^{-1}(s) & 0 \\ 0 & \hat{S}_3^{-1}(s) \end{bmatrix} \begin{bmatrix} I_q & 0 & 0_{q \times m} \\ 0 & \hat{P}(s) & 0 \end{bmatrix} \\ &\times \begin{bmatrix} I_q & 0 & -\hat{S}_1(s)A_{12} \\ 0 & 0 & \hat{S}_4^{-1}(s) \\ 0 & I_m & 0 \end{bmatrix} \begin{bmatrix} \hat{S}_2^{-1}(s) & 0 \\ 0 & I_{n-q} \end{bmatrix} \begin{bmatrix} I_q & 0 & 0_{q \times m} \\ 0 & 0 & I_m \\ 0 & I_{n-q} & 0 \end{bmatrix} \begin{bmatrix} S^{-1} & 0 \\ 0 & I_m \end{bmatrix}. \quad \square \end{aligned}$$

Proposition 12.6.17. Let $s \in \mathbb{C}$. Then,

$$\mathcal{C}(A, B) \subseteq \text{Re } \mathcal{R}(\begin{bmatrix} sI - A & B \end{bmatrix}). \quad (12.6.29)$$

Proof. Using Proposition 12.6.9 and the notation in the proof of Proposition 12.6.16, it follows that, for all $s \in \mathbb{C}$,

$$\mathcal{C}(A, B) = \mathcal{R}(S \begin{bmatrix} I_q \\ 0 \end{bmatrix}) \subseteq \mathcal{R}\left(S \begin{bmatrix} \hat{S}_1^{-1}(s) & 0 \\ 0 & \hat{S}_3^{-1}(s)\hat{P}(s) \end{bmatrix}\right) = \mathcal{R}(\begin{bmatrix} sI - A & B \end{bmatrix}). \quad \square$$

The next result characterizes controllability in several equivalent ways.

Theorem 12.6.18. The following statements are equivalent:

- i) (A, B) is controllable.
- ii) There exists $t > 0$ such that $\int_0^t e^{\tau A} B B^T e^{\tau A^T} d\tau$ is positive definite.
- iii) $\int_0^t e^{\tau A} B B^T e^{\tau A^T} d\tau$ is positive definite for all $t > 0$.
- iv) $\text{rank } \mathcal{K}(A, B) = n$.
- v) Every eigenvalue of (A, B) is controllable.

If, in addition, $\lim_{t \rightarrow \infty} \int_0^t e^{\tau A} B B^T e^{\tau A^T} d\tau$ exists, then the following condition is equivalent to i)–v):

- vi) $\int_0^\infty e^{tA} B B^T e^{tA^T} dt$ is positive definite.

Proof. The equivalence of i)–iv) follows from Lemma 12.6.2.

To prove iv) $\implies$ v), suppose that v) does not hold, that is, there exist $\lambda \in \text{spec}(A)$ and a nonzero vector $x \in \mathbb{C}^n$ such that $x^* A = \lambda x^*$ and $x^* B = 0$. It thus follows that $x^* A B = \lambda x^* B = 0$. Similarly, $x^* A^i B = 0$ for all $i = 0, 1, \dots, n-1$. Hence, $(\text{Re } x)^T \mathcal{K}(A, B) = 0$ and $(\text{Im } x)^T \mathcal{K}(A, B) = 0$. Since $\text{Re } x$ and $\text{Im } x$ are not both zero, it follows that $\dim \mathcal{C}(A, B) < n$.

Conversely, to show that v) implies iv), suppose that $\text{rank } \mathcal{K}(A, B) < n$. Then, there exists a nonzero vector $x \in \mathbb{R}^n$ such that $x^T A^i B = 0$ for all $i = 0, \dots, n-1$. Now, let $p \in \mathbb{R}[s]$ be a nonzero polynomial of minimal degree such that $x^T p(A) = 0$. Note that p is not a constant polynomial and that $x^T \mu_A(A) = 0$. Thus, $1 \leq \deg p \leq \deg \mu_A$. Now, let $\lambda \in \mathbb{C}$ be such that $p(\lambda) = 0$, and let $q \in \mathbb{R}[s]$ be such that $p(s) = q(s)(s - \lambda)$ for all $s \in \mathbb{C}$. Since $\deg q < \deg p$, it follows that $x^T q(A) \neq 0$.

Therefore, $\eta \triangleq q(A)x$ is nonzero. Furthermore, $\eta^T(A - \lambda I) = x^T p(A) = 0$. Since $x^T A^i B = 0$ for all $i = 0, \dots, n-1$, it follows that $\eta^T B = x^T q(A)B = 0$. Consequently, $v)$ does not hold. $\square$

The following result implies that arbitrary eigenvalue placement is possible for (12.6.1) when (A, B) is controllable.

Proposition 12.6.19. The pair (A, B) is controllable if and only if, for every polynomial $p \in \mathbb{R}[s]$ such that $\deg p = n$, there exists a matrix $K \in \mathbb{R}^{m \times n}$ such that $\text{mspec}(A + BK) = \text{mroots}(p)$.

Proof. For the case $m = 1$ let $A_c \triangleq C(\chi_A)$ and $B_c \triangleq e_n$ as in (12.9.5). Then, Proposition 12.9.3 implies that $\mathcal{K}(A_c, B_c)$ is nonsingular, while Corollary 12.9.9 implies that $A_c = S^{-1}AS$ and $B_c = S^{-1}B$. Now, let $\text{mroots}(p) = \{\lambda_1, \dots, \lambda_n\}_{\text{ms}} \subset \mathbb{C}$. Letting $K \triangleq e_n^T [C(p) - A_c] S^{-1}$ it follows that

$$\begin{aligned} A + BK &= S(A_c + B_c K S) S^{-1} \\ &= S(A_c + E_{n,n}[C(p) - A_c]) S^{-1} \\ &= SC(p) S^{-1}. \end{aligned}$$

The case $m > 1$ requires the multivariable controllable canonical form. See [1150, p. 248]. $\square$

12.7 Controllable Asymptotic Stability

Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$, and define $q \triangleq \dim \mathcal{C}(A, C)$.

Definition 12.7.1. (A, B) is *controllably asymptotically stable* if

$$\mathcal{C}(A, B) \subseteq \mathcal{S}_s(A). \quad (12.7.1)$$

Proposition 12.7.2. Let $K \in \mathbb{R}^{m \times n}$. Then, (A, B) is controllably asymptotically stable if and only if $(A + BK, B)$ is controllably asymptotically stable.

Proposition 12.7.3. The following statements are equivalent:

- i) (A, B) is controllably asymptotically stable.
- ii) There exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) holds, where $A_1 \in \mathbb{R}^{q \times q}$ is asymptotically stable and $B_1 \in \mathbb{R}^{q \times m}$.
- iii) There exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) holds, where $A_1 \in \mathbb{R}^{q \times q}$ is asymptotically stable and $B_1 \in \mathbb{R}^{q \times m}$.
- iv) $\lim_{t \rightarrow \infty} e^{tA} B = 0$.
- v) The positive-semidefinite matrix $Q \in \mathbb{R}^{n \times n}$ defined by

$$Q \triangleq \int_0^\infty e^{tA} B B^T e^{tA^T} dt \quad (12.7.2)$$

exists.

vi) There exists a positive-semidefinite matrix $Q \in \mathbb{R}^{n \times n}$ satisfying

$$AQ + QA^T + BB^T = 0. \quad (12.7.3)$$

In this case, the positive-semidefinite matrix $Q \in \mathbb{R}^{n \times n}$ defined by (12.7.2) satisfies (12.7.3).

Proof. To prove *i) $\implies$ ii)*, assume that (A, B) is controllably asymptotically stable so that $\mathcal{C}(A, B) \subseteq \mathcal{S}_s(A) = \mathcal{N}[\mu_A^s(A)] = \mathcal{R}[\mu_A^u(A)]$. Using Theorem 12.6.8, it follows that there exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) is satisfied, where $A_1 \in \mathbb{R}^{q \times q}$ and (A_1, B_1) is controllable. Thus, $\mathcal{R}(S \begin{bmatrix} I_q \\ 0 \end{bmatrix}) = \mathcal{C}(A, B) \subseteq \mathcal{R}[\mu_A^s(A)]$.

Next, note that

$$\mu_A^s(A) = S \begin{bmatrix} \mu_{A_1}^s(A_1) & B_{12s} \\ 0 & \mu_{A_1}^s(A_2) \end{bmatrix} S^{-1},$$

where $B_{12s} \in \mathbb{R}^{q \times (n-q)}$, and suppose that A_1 is not asymptotically stable with CRHP eigenvalue λ . Then, $\lambda \notin \text{roots}(\mu_A^s)$, and thus $\mu_A^s(A_1) \neq 0$. Let $x_1 \in \mathbb{R}^{n-q}$ satisfy $\mu_A^s(A_1)x_1 \neq 0$. Then,

$$\begin{bmatrix} x_1 \\ 0 \end{bmatrix} \in \mathcal{R}\left(S \begin{bmatrix} I_q \\ 0 \end{bmatrix}\right) = \mathcal{C}(A, B)$$

and

$$\mu_A^s(A)S \begin{bmatrix} x_1 \\ 0 \end{bmatrix} = S \begin{bmatrix} \mu_{A_1}^s(A_1)x_1 \\ 0 \end{bmatrix},$$

and thus $\begin{bmatrix} x_1 \\ 0 \end{bmatrix} \notin \mathcal{N}[\mu_A^s(A)] = \mathcal{S}_s(A)$, which implies that $\mathcal{C}(A, B)$ is not contained in $\mathcal{S}_s(A)$. Hence, A_1 is asymptotically stable.

To prove *iii) $\implies$ iv)*, assume there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) holds, where $A_1 \in \mathbb{R}^{k \times k}$ is asymptotically stable and $B_1 \in \mathbb{R}^{k \times m}$. Thus, $e^{tA}B = \begin{bmatrix} e^{tA_1}B_1 \\ 0 \end{bmatrix} S \rightarrow 0$ as $t \rightarrow \infty$.

Next, to prove that *iv)* implies *v)*, assume that $e^{tA}B \rightarrow 0$ as $t \rightarrow \infty$. Then, every entry of $e^{tA}B$ involves exponentials of t , where the coefficients of t have negative real part. Hence, so does every entry of $e^{tA}BB^Te^{tA^T}$, which implies that $\int_0^\infty e^{tA}BB^Te^{tA^T} dt$ exists.

To prove *v) $\implies$ vi)*, note that, since $Q = \int_0^\infty e^{tA}BB^Te^{tA^T} dt$ exists, it follows that $e^{tA}BB^Te^{tA^T} \rightarrow 0$ as $t \rightarrow \infty$. Thus,

$$\begin{aligned}
 AQ + QA^T &= \int_0^\infty \left[Ae^{tA}BB^Te^{tA^T} + e^{tA}BB^Te^{tA^T}A \right] dt \\
 &= \int_0^\infty \frac{d}{dt} e^{tA}BB^Te^{tA^T} dt \\
 &= \lim_{t \rightarrow \infty} e^{tA}BB^Te^{tA^T} - BB^T = -BB^T,
 \end{aligned}$$

which shows that Q satisfies (12.4.3).

To prove $vi) \implies i)$, suppose there exists a positive-semidefinite matrix $Q \in \mathbb{R}^{n \times n}$ satisfying (12.7.3). Then,

$$\begin{aligned}
 \int_0^t e^{tA}BB^Te^{tA^T} d\tau &= - \int_0^t e^{\tau A} (AQ + QA^T) e^{tA^T} d\tau = - \int_0^t \frac{d}{d\tau} e^{\tau A} QA^T d\tau \\
 &= Q - e^{tA}Qe^{tA^T} \leq Q.
 \end{aligned}$$

Next, it follows from Theorem 12.6.8 that there exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) is satisfied, where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, and (A_1, B_1) is controllable. Consequently, we have

$$\begin{aligned}
 \int_0^t e^{\tau A_1} B_1 B_1^T e^{\tau A_1^T} d\tau &= \begin{bmatrix} I & 0 \end{bmatrix} S \int_0^t e^{\tau A} BB^T e^{\tau A^T} d\tau S^T \begin{bmatrix} I \\ 0 \end{bmatrix} \\
 &\leq \begin{bmatrix} I & 0 \end{bmatrix} S Q S^T \begin{bmatrix} I \\ 0 \end{bmatrix}.
 \end{aligned}$$

Thus, it follows from Proposition 8.6.3 that $Q_1 \triangleq \int_0^\infty e^{tA_1} B_1 B_1^T e^{tA_1^T} dt$ exists. Since (A_1, B_1) is controllable, it follows from $vi)$ of Theorem 12.6.18 that Q_1 is positive definite.

Now, let λ be an eigenvalue of A_1^T , and let $x_1 \in \mathbb{C}^n$ be an associated eigenvector. Consequently, $\alpha \triangleq x_1^* Q_1 x_1$ is positive, and

$$\alpha = x_1^* \int_0^\infty e^{\bar{\lambda} t} BB_1^T e^{\lambda t} dt x_1 = x_1^* B_1 B_1^T x_1 \int_0^\infty e^{2(\operatorname{Re} \lambda)t} dt.$$

Hence, $\int_0^\infty e^{2(\operatorname{Re} \lambda)t} dt = \alpha / x_1^* B_1 B_1^T x_1$ exists, and thus $\operatorname{Re} \lambda < 0$. Consequently, A_1 is asymptotically stable, and thus $\mathcal{C}(A, B) \subseteq \mathcal{S}_s(A)$, that is, (A, B) is controllably asymptotically stable. $\square$

The matrix $Q \in \mathbb{R}^{n \times n}$ defined by (12.7.2) is the *controllability Gramian*, and (12.7.3) is the *control Lyapunov equation*.

Proposition 12.7.4. Assume that (A, B) is controllably asymptotically stable, let $Q \in \mathbb{R}^{n \times n}$ be the positive-semidefinite matrix defined by (12.7.2), and define $\mathcal{Q} \in \mathbb{R}^{n \times n}$ by (12.6.11). Then, the following statements hold:

$$i) \quad QQ^+ = \mathcal{Q}.$$

- ii) $\mathcal{R}(Q) = \mathcal{R}(\mathcal{Q}) = \mathcal{C}(A, B)$.
- iii) $\mathcal{N}(Q) = \mathcal{N}(\mathcal{Q}) = \mathcal{C}(A, B)^\perp$.
- iv) $\text{rank } Q = \text{rank } \mathcal{Q} = q$.
- v) Q is the only positive-semidefinite solution of (12.7.3) whose rank is q .

Proof. See [1207] for the proof of v). □

Proposition 12.7.5. Assume that (A, B) is controllably asymptotically stable, let $Q \in \mathbb{R}^{n \times n}$ be the positive-semidefinite matrix defined by (12.7.2), and let $\hat{Q} \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

- i) $\hat{Q}$ is positive semidefinite and satisfies (12.7.3).
- ii) There exists a positive-semidefinite matrix $Q_0 \in \mathbb{R}^{n \times n}$ such that $\hat{Q} = Q + Q_0$ and $AQ_0 + Q_0A^T = 0$.

In this case,

$$\text{rank } \hat{Q} = q + \text{rank } Q_0 \quad (12.7.4)$$

and

$$\text{rank } Q_0 \leq \sum_{\substack{\lambda \in \text{spec}(A) \\ \lambda \in j\mathbb{R}}} \text{gmult}_A(\lambda). \quad (12.7.5)$$

Proof. See [1207]. □

Proposition 12.7.6. The following statements are equivalent:

- i) (A, B) is controllably asymptotically stable, every imaginary eigenvalue of A is semisimple, and A has no ORHP eigenvalues.
- ii) (12.7.3) has a positive-definite solution $Q \in \mathbb{R}^{n \times n}$.

Proof. See [1207]. □

Proposition 12.7.7. The following statements are equivalent:

- i) (A, B) is controllably asymptotically stable, and A has no imaginary eigenvalues.
- ii) (12.7.3) has exactly one positive-semidefinite solution $Q \in \mathbb{R}^{n \times n}$.

In this case, $Q \in \mathbb{R}^{n \times n}$ is given by (12.7.2) and satisfies $\text{rank } Q = q$.

Proof. See [1207]. □

Corollary 12.7.8. Assume that A is asymptotically stable. Then, the positive-semidefinite matrix $Q \in \mathbb{R}^{n \times n}$ defined by (12.7.2) is the unique solution of (12.7.3) and satisfies $\text{rank } Q = q$.

Proof. See [1207]. □

Proposition 12.7.9. The following statements are equivalent:

- i) (A, B) is controllable, and A is asymptotically stable.
- ii) (12.7.3) has exactly one positive-semidefinite solution $Q \in \mathbb{R}^{n \times n}$, and Q is positive definite.

In this case, $Q \in \mathbb{R}^{n \times n}$ is given by (12.7.2).

Proof. See [1207]. □

Corollary 12.7.10. Assume that A is asymptotically stable. Then, the positive-semidefinite matrix $Q \in \mathbb{R}^{n \times n}$ defined by (12.7.2) exists. Furthermore, Q is positive definite if and only if (A, B) is controllable.

12.8 Stabilizability

Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$, and define $q \triangleq \dim \mathcal{C}(A, B)$.

Definition 12.8.1. (A, B) is *stabilizable* if

$$\mathcal{S}_u(A) \subseteq \mathcal{C}(A, B). \quad (12.8.1)$$

Proposition 12.8.2. Let $K \in \mathbb{R}^{m \times n}$. Then, (A, B) is stabilizable if and only if $(A + BK, B)$ is stabilizable.

Proposition 12.8.3. The following statements are equivalent:

- i) A is asymptotically stable.
- ii) (A, B) is stabilizable and controllably asymptotically stable.

Proof. Suppose that A is asymptotically stable. Then, $\mathcal{S}_u(A) = \{0\}$, and $\mathcal{S}_s(A) = \mathbb{R}^n$. Thus, $\mathcal{S}_u(A) \subseteq \mathcal{C}(A, B)$, and $\mathcal{C}(A, B) \subseteq \mathcal{S}_s(A)$. Conversely, assume that (A, B) is stabilizable and controllably asymptotically stable. Then, $\mathcal{S}_u(A) \subseteq \mathcal{C}(A, B) \subseteq \mathcal{S}_s(A)$, and thus $\mathcal{S}_u(A) = \{0\}$. □

Proposition 12.8.4. The following statements are equivalent:

- i) (A, B) is stabilizable.
- ii) There exists an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) holds, where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, (A_1, B_1) is controllable, and $A_2 \in \mathbb{R}^{(n-q) \times (n-q)}$ is asymptotically stable.
- iii) There exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) holds, where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, (A_1, B_1) is controllable, and $A_2 \in \mathbb{R}^{(n-q) \times (n-q)}$ is asymptotically stable.
- iv) Every CRHP eigenvalue of (A, B) is controllable.

Proof. To prove i) $\implies$ ii), assume that (A, B) is stabilizable so that $\mathcal{S}_u(A) = \mathcal{N}[\mu_A^u(A)] = \mathcal{R}[\mu_A^s(A)] \subseteq \mathcal{C}(A, B)$. Using Theorem 12.6.8, it follows that there exists

an orthogonal matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) is satisfied, where $A_1 \in \mathbb{R}^{q \times q}$ and (A_1, B_1) is controllable. Thus, $\mathcal{R}[\mu_A^s(A)] \subseteq \mathcal{C}(A, B) = \mathcal{R}(S \begin{bmatrix} I_q \\ 0 \end{bmatrix})$.

Next, note that

$$\mu_A^s(A) = S \begin{bmatrix} \mu_A^s(A_1) & B_{12s} \\ 0 & \mu_A^s(A_2) \end{bmatrix} S^{-1},$$

where $B_{12s} \in \mathbb{R}^{q \times (n-q)}$, and suppose that A_2 is not asymptotically stable with CRHP eigenvalue λ . Then, $\lambda \notin \text{roots}(\mu_A^s)$, and thus $\mu_A^s(A_2) \neq 0$. Let $x_2 \in \mathbb{R}^{n-q}$ satisfy $\mu_A^s(A_2)x_2 \neq 0$. Then,

$$\mu_A^s(A)S \begin{bmatrix} 0 \\ x_2 \end{bmatrix} = S \begin{bmatrix} B_{12s}x_2 \\ \mu_A^s(A_2)x_2 \end{bmatrix} \notin \mathcal{R}\left(S \begin{bmatrix} I_q \\ 0 \end{bmatrix}\right) = \mathcal{C}(A, B),$$

which implies that $\mathcal{S}_u(A)$ is not contained in $\mathcal{C}(A, B)$. Hence, A_2 is asymptotically stable.

The statement *ii*) implies *iii*) is immediate.

To prove *iii*) $\implies$ *iv*), let $\lambda \in \text{spec}(A)$ be a CRHP eigenvalue of A . Since A_2 is asymptotically stable, it follows that $\lambda \notin \text{spec}(A_2)$. Consequently, Proposition 12.6.13 implies that λ is not an uncontrollable eigenvalue of (A, B) , and thus λ is a controllable eigenvalue of (A, B) .

To prove *iv*) $\implies$ *i*), let $S \in \mathbb{R}^{n \times n}$ be nonsingular and such that A and B have the form (12.6.21), where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, and (A_1, B_1) is controllable. Since every CRHP eigenvalue of (A, B) is controllable, it follows from Proposition 12.6.13 that A_2 is asymptotically stable. From Fact 11.23.4 it follows that $\mathcal{S}_u(A) \subseteq \mathcal{R}(S \begin{bmatrix} I_q \\ 0 \end{bmatrix}) = \mathcal{C}(A, B)$, which implies that (A, B) is stabilizable. $\square$

Proposition 12.8.5. The following statements are equivalent:

- i*) (A, B) is controllably asymptotically stable and stabilizable.
- ii*) A is asymptotically stable.

Proof. Since (A, B) is stabilizable, it follows from Proposition 12.5.4 that there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.6.21) holds, where $A_1 \in \mathbb{R}^{q \times q}$, $B_1 \in \mathbb{R}^{q \times m}$, (A_1, B_1) is controllable, and $A_2 \in \mathbb{R}^{(n-q) \times (n-q)}$ is asymptotically stable. Then,

$$\int_0^\infty e^{tA} B B^T e^{tA^T} dt = S \begin{bmatrix} \int_0^\infty e^{tA_1} B_1 B_1^T e^{tA_1^T} dt & 0 \\ 0 & 0 \end{bmatrix} S^{-1}.$$

Since the integral on the left-hand side exists by assumption, the integral on the right-hand side also exists. Since (A_1, B_1) is controllable, it follows from *vii*) of Theorem 12.6.18 that $Q_1 \triangleq \int_0^\infty e^{tA_1} B_1 B_1^T e^{tA_1^T} dt$ is positive definite.

Now, let λ be an eigenvalue of A_1^T , and let $x_1 \in \mathbb{C}^q$ be an associated eigenvector. Consequently, $\alpha \triangleq x_1^* Q_1 x_1$ is positive, and

$$\alpha = x_1^* \int_0^\infty e^{\bar{\lambda}t} B_1 B_1^T e^{\lambda t} dt x_1 = x_1^* B_1 B_1^T x_1 \int_0^\infty e^{2(\operatorname{Re} \lambda)t} dt.$$

Hence, $\int_0^\infty e^{2(\operatorname{Re} \lambda)t} dt$ exists, and thus $\operatorname{Re} \lambda < 0$. Consequently, A_1 is asymptotically stable, and thus A is asymptotically stable. $\square$

Corollary 12.8.6. The following statements are equivalent:

- i) There exists a positive-semidefinite matrix $Q \in \mathbb{R}^{n \times n}$ satisfying (12.7.3), and (A, B) is stabilizable.
- ii) A is asymptotically stable.

Proof. The result follows from Proposition 12.7.3 and Proposition 12.8.5. $\square$

12.9 Realization Theory

Given a proper rational transfer function G we wish to determine (A, B, C, D) such that (12.2.11) holds. The following terminology is convenient.

Definition 12.9.1. Let $G \in \mathbb{R}^{l \times m}(s)$. If $l = m = 1$, then G is a *single-input/single-output (SISO)* rational transfer function; if $l = 1$ and $m > 1$, then G is a *multiple-input/single-output (MISO)* rational transfer function; if $l > 1$ and $m = 1$, then G is a *single-input/multiple-output (SIMO)* rational transfer function; and, if $l > 1$ or $m > 1$, then G is a *multiple-input/multiple output (MIMO)* rational transfer function.

Definition 12.9.2. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, and assume that $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, $C \in \mathbb{R}^{l \times n}$, and $D \in \mathbb{R}^{l \times m}$ satisfy $G(s) = C(sI - A)^{-1}B + D$. Then, $\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ is a *realization* of G , which is written as

$$G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]. \quad (12.9.1)$$

The *order* of the realization (12.9.1) is the order of A . Finally, the realization (12.9.1) is *controllable and observable* if (A, B) is controllable and (A, C) is observable.

Suppose that $n = 0$. Then, A , B , and C are empty matrices, and $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$ is given by

$$G(s) = 0_{l \times 0}(sI_{0 \times 0} - 0_{0 \times 0})^{-1}0_{0 \times m} + D = 0_{l \times m} + D = D. \quad (12.9.2)$$

Therefore, the order of the realization $\left[\begin{array}{c|c} 0_{0 \times 0} & 0_{0 \times m} \\ \hline 0_{l \times 0} & D \end{array} \right]$ is zero.

Although the realization (12.9.1) is not unique, the matrix D is unique and is given by

$$D = G(\infty). \quad (12.9.3)$$

Furthermore, note that $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ if and only if $G - D \sim \left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$. Therefore, it suffices to construct realizations for strictly proper transfer functions.

The following result shows that every strictly proper, SISO rational transfer function G has a realization. In fact, two realizations are the *controllable canonical form* $G \sim \left[\begin{array}{c|c} A_c & B_c \\ \hline C_c & 0 \end{array} \right]$ and the *observable canonical form* $G \sim \left[\begin{array}{c|c} A_o & B_o \\ \hline C_o & 0 \end{array} \right]$. If G is exactly proper, then a realization can be obtained for $G - G(\infty)$.

Proposition 12.9.3. Let $G \in \mathbb{R}_{\text{prop}}(s)$ be the SISO strictly proper rational transfer function

$$G(s) = \frac{\alpha_{n-1}s^{n-1} + \alpha_{n-2}s^{n-2} + \cdots + \alpha_1s + \alpha_0}{s^n + \beta_{n-1}s^{n-1} + \cdots + \beta_1s + \beta_0}. \quad (12.9.4)$$

Then, $G \sim \left[\begin{array}{c|c} A_c & B_c \\ \hline C_c & 0 \end{array} \right]$, where A_c, B_c, C_c are defined by

$$A_c \triangleq \begin{bmatrix} 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \\ -\beta_0 & -\beta_1 & -\beta_2 & \cdots & -\beta_{n-1} \end{bmatrix}, \quad B_c \triangleq \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix}, \quad (12.9.5)$$

$$C_c \triangleq [\alpha_0 \quad \alpha_1 \quad \alpha_2 \quad \cdots \quad \alpha_{n-1}], \quad (12.9.6)$$

and $G \sim \left[\begin{array}{c|c} A_o & B_o \\ \hline C_o & 0 \end{array} \right]$, where A_o, B_o, C_o are defined by

$$A_o \triangleq \begin{bmatrix} 0 & 0 & \cdots & 0 & -\beta_0 \\ 1 & 0 & \cdots & 0 & -\beta_1 \\ 0 & 1 & \cdots & 0 & -\beta_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 1 & -\beta_{n-1} \end{bmatrix}, \quad B_o \triangleq \begin{bmatrix} \alpha_0 \\ \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_{n-1} \end{bmatrix}, \quad (12.9.7)$$

$$C_o \triangleq [0 \quad 0 \quad \cdots \quad 0 \quad 1]. \quad (12.9.8)$$

Furthermore, (A_c, B_c) is controllable, and (A_o, C_o) is observable. Finally, the following statements are equivalent:

- i) The numerator and denominator of G given in (12.9.4) are coprime.
- ii) (A_c, C_c) is observable.
- iii) (A_c, B_c, C_c) is controllable and observable.
- iv) (A_o, B_o) is controllable.
- v) (A_o, B_o, C_o) is controllable and observable.

Proof. The realizations can be verified directly. Furthermore, note that

$$\mathcal{K}(A_c, B_c) = \mathcal{O}(A_o, C_o) = \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & 0 & \ddots & 1 & -\beta_{n-1} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & 1 & \ddots & -\beta_3 & -\beta_2 \\ 0 & 1 & -\beta_{n-1} & \cdots & -\beta_2 & -\beta_1 \\ 1 & -\beta_{n-1} & -\beta_{n-2} & \cdots & -\beta_1 & -\beta_0 \end{bmatrix}.$$

It follows from Fact 2.13.8 that $\det \mathcal{K}(A_c, B_c) = \det \mathcal{O}(A_o, C_o) = (-1)^{\lfloor n/2 \rfloor}$, which implies that (A_c, B_c) is controllable and (A_o, C_o) is observable.

To prove the last statement, let $p, q \in \mathbb{R}[s]$ denote the numerator and denominator, respectively, of G in (12.9.4). Then, for $n = 2$,

$$\mathcal{K}(A_o, B_o) = \mathcal{O}^T(A_c, C_c) = B(p, q) \hat{I} \begin{bmatrix} 1 & -\beta_1 \\ 0 & 1 \end{bmatrix},$$

where $B(p, q)$ is the Bezout matrix of p and q . It follows from *ix*) of Fact 4.8.6 that $B(p, q)$ is nonsingular if and only if p and q are coprime. $\square$

The following result shows that every proper rational transfer function has a realization.

Theorem 12.9.4. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$. Then, there exist $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, $C \in \mathbb{R}^{l \times n}$, and $D \in \mathbb{R}^{l \times m}$ such that $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$.

Proof. By Proposition 12.9.3, every entry $G_{(i,j)}$ of G has a realization $G_{(i,j)} \sim \left[\begin{array}{c|c} A_{ij} & B_{ij} \\ \hline C_{ij} & D_{ij} \end{array} \right]$. Combining these realizations yields a realization of G . $\square$

Proposition 12.9.5. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$ have the n th-order realization $\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$, let $S \in \mathbb{R}^{n \times n}$, and assume that S is nonsingular. Then,

$$G \sim \left[\begin{array}{c|c} SAS^{-1} & SB \\ \hline CS^{-1} & D \end{array} \right]. \quad (12.9.9)$$

If, in addition, $\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ is controllable and observable, then so is $\left[\begin{array}{c|c} SAS^{-1} & SB \\ \hline CS^{-1} & D \end{array} \right]$.

Definition 12.9.6. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, and let $\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ and $\left[\begin{array}{c|c} \hat{A} & \hat{B} \\ \hline \hat{C} & \hat{D} \end{array} \right]$ be n th-order realizations of G . Then, $\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ and $\left[\begin{array}{c|c} \hat{A} & \hat{B} \\ \hline \hat{C} & \hat{D} \end{array} \right]$ are *equivalent* if there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that $\hat{A} = SAS^{-1}$, $\hat{B} = SB$, and $\hat{C} = CS^{-1}$.

The following result shows that the Markov parameters of a rational transfer function are independent of the realization.

Proposition 12.9.7. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, and assume that $G \sim \left[\frac{A}{C} \middle| \frac{B}{D} \right]$, where $A \in \mathbb{R}^{n \times n}$, and $G \sim \left[\frac{\hat{A}}{\hat{C}} \middle| \frac{\hat{B}}{\hat{D}} \right]$, where $A \in \mathbb{R}^{\hat{n} \times \hat{n}}$. Then, $D = \hat{D}$, and, for all $k \geq 0$, $CA^k B = \hat{C}\hat{A}^k\hat{B}$.

Proposition 12.9.8. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, assume that G has the n th-order realizations $\left[\frac{A_1}{C_1} \middle| \frac{B_1}{D} \right]$ and $\left[\frac{A_2}{C_2} \middle| \frac{B_2}{D} \right]$, and assume that both of these realizations are controllable and observable. Then, these realizations are equivalent. Furthermore, there exists a unique matrix $S \in \mathbb{R}^{n \times n}$ such that

$$\left[\frac{A_2}{C_2} \middle| \frac{B_2}{D} \right] = \left[\frac{SA_1S^{-1}}{C_1S^{-1}} \middle| \frac{SB_1}{D} \right]. \quad (12.9.10)$$

In fact,

$$S = (\mathcal{O}_2^T \mathcal{O}_2)^{-1} \mathcal{O}_2^T \mathcal{O}_1, \quad S^{-1} = \mathcal{K}_1 \mathcal{K}_2^T (\mathcal{K}_2 \mathcal{K}_2^T)^{-1}, \quad (12.9.11)$$

where, for $i = 1, 2$, $\mathcal{K}_i \triangleq \mathcal{K}(A_i, B_i)$ and $\mathcal{O}_i \triangleq \mathcal{O}(A_i, C_i)$.

Proof. By Proposition 12.9.7, the realizations $\left[\frac{A_1}{C_1} \middle| \frac{B_1}{D} \right]$ and $\left[\frac{A_2}{C_2} \middle| \frac{B_2}{D} \right]$ generate the same Markov parameters. Hence, $\mathcal{O}_1 A_1 \mathcal{K}_1 = \mathcal{O}_2 A_2 \mathcal{K}_2$, $\mathcal{O}_1 B_1 = \mathcal{O}_2 B_2$, and $C_1 \mathcal{K}_1 = C_2 \mathcal{K}_2$. Since $\left[\frac{A_2}{C_2} \middle| \frac{B_2}{D} \right]$ is controllable and observable, it follows that the $n \times n$ matrices $\mathcal{K}_2 \mathcal{K}_2^T$ and $\mathcal{O}_2^T \mathcal{O}_2$ are nonsingular. Consequently, $A_2 = SA_1 S^{-1}$, $B_2 = SB_1$, and $C_2 = C_1 S^{-1}$.

To prove uniqueness, assume there exists a matrix $\hat{S} \in \mathbb{R}^{n \times n}$ such that $A_2 = \hat{S} A_1 \hat{S}^{-1}$, $B_2 = \hat{S} B_1$, and $C_2 = C_1 \hat{S}^{-1}$. Then, it follows that $\mathcal{O}_1 \hat{S} = \mathcal{O}_2$. Since $\mathcal{O}_1 S = \mathcal{O}_2$, it follows that $\mathcal{O}_1 (S - \hat{S}) = 0$. Consequently, $S = \hat{S}$. $\square$

Corollary 12.9.9. Let $G \in \mathbb{R}_{\text{prop}}(s)$ be given by (12.9.4), assume that G has the n th-order controllable and observable realization $\left[\frac{A}{C} \middle| \frac{B}{0} \right]$, and define A_c, B_c, C_c by (12.9.5), (12.9.6) and A_o, B_o, C_o by (12.9.7), (12.9.8). Furthermore, define $S_c \triangleq [\mathcal{O}(A, B)]^{-1} \mathcal{O}(A_c, B_c)$. Then,

$$S_c^{-1} = \mathcal{K}(A, B) [\mathcal{K}(A_c, B_c)]^{-1} \quad (12.9.12)$$

and

$$\left[\frac{S_c A S_c^{-1}}{C S_c^{-1}} \middle| \frac{S_c B}{0} \right] = \left[\frac{A_c}{C_c} \middle| \frac{B_c}{0} \right]. \quad (12.9.13)$$

Furthermore, define $S_o \triangleq [\mathcal{O}(A, B)]^{-1} \mathcal{O}(A_o, B_o)$. Then,

$$S_o^{-1} = \mathcal{K}(A, B) [\mathcal{K}(A_o, B_o)]^{-1} \quad (12.9.14)$$

and

$$\left[\frac{S_o A S_o^{-1}}{C S_o^{-1}} \middle| \frac{S_o B}{0} \right] = \left[\frac{A_o}{C_o} \middle| \frac{B_o}{0} \right]. \quad (12.9.15)$$

The following result, known as the *Kalman decomposition*, is useful for constructing controllable and observable realizations.

Proposition 12.9.10. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. Then, there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that

$$A = S \begin{bmatrix} A_1 & 0 & A_{13} & 0 \\ A_{21} & A_2 & A_{23} & A_{24} \\ 0 & 0 & A_3 & 0 \\ 0 & 0 & A_{43} & A_4 \end{bmatrix} S^{-1}, \quad B = S \begin{bmatrix} B_1 \\ B_2 \\ 0 \\ 0 \end{bmatrix}, \quad (12.9.16)$$

$$C = [C_1 \quad 0 \quad C_3 \quad 0] S^{-1}, \quad (12.9.17)$$

where, for $i = 1, \dots, 4$, $A_i \in \mathbb{R}^{n_i \times n_i}$, $(\left[\begin{array}{c|c} A_1 & 0 \\ \hline A_{21} & A_2 \end{array} \right], \left[\begin{array}{c} B_1 \\ B_2 \end{array} \right])$ is controllable, and $(\left[\begin{array}{c|c} A_1 & A_{13} \\ \hline 0 & A_3 \end{array} \right], [C_1 \ C_3])$ is observable. Furthermore, the following statements hold:

- i) (A, B) is stabilizable if and only if A_3 and A_4 are asymptotically stable.
- ii) (A, B) is controllable if and only if A_3 and A_4 are empty.
- iii) (A, C) is detectable if and only if A_2 and A_4 are asymptotically stable.
- iv) (A, C) is observable if and only if A_2 and A_4 are empty.
- v) $G \sim \left[\begin{array}{c|c} A_1 & B_1 \\ \hline C_1 & D \end{array} \right]$.
- vi) The realization $\left[\begin{array}{c|c} A_1 & B_1 \\ \hline C_1 & D \end{array} \right]$ is controllable and observable.

Proof. Let $\alpha \leq 0$ be such that $A + \alpha I$ is asymptotically stable, and let $Q \in \mathbb{R}^{n \times n}$ and $P \in \mathbb{R}^{n \times n}$ denote the controllability and observability Gramians of the system $(A + \alpha I, B, C)$. Then, Theorem 8.3.4 implies that there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that

$$Q = S \begin{bmatrix} Q_1 & & 0 \\ & Q_2 & \\ 0 & & 0 \end{bmatrix} S^T, \quad P = S^{-T} \begin{bmatrix} P_1 & & 0 \\ & 0 & \\ 0 & & P_2 \end{bmatrix} S^{-1},$$

where Q_1 and P_1 are the same order, and where Q_1, Q_2, P_1 , and P_2 are positive definite and diagonal. The form of SAS^{-1} , SB , and CS^{-1} given by (12.9.17) now follows from (12.7.3) and (12.4.3) with A replaced by $A + \alpha I$, where, as in the proof of Theorem 12.6.8, $SAS^{-1} = S(A + \alpha I)S^{-1} - \alpha I$. Finally, statements i)–v) are immediate, while it can be verified directly that $\left[\begin{array}{c|c} A_1 & B_1 \\ \hline C_1 & D_1 \end{array} \right]$ is a realization of G . $\square$

Note that the uncontrollable multispectrum of (A, B) is given by $\text{mspec}(A_3) \cup \text{mspec}(A_4)$, while the unobservable multispectrum of (A, C) is given by $\text{mspec}(A_2) \cup \text{mspec}(A_4)$. Likewise, the *uncontrollable-unobservable multispectrum* of (A, B, C) is given by $\text{mspec}(A_4)$.

Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$. Then, define the *i-step observability matrix* $\mathcal{O}_i(A, C) \in$

$\mathbb{R}^{il \times n}$ by

$$\mathcal{O}_i(A, C) \triangleq \begin{bmatrix} C \\ CA \\ \vdots \\ CA^{i-1} \end{bmatrix} \quad (12.9.18)$$

and the j -step controllability matrix $\mathcal{K}_j(A, B) \in \mathbb{R}^{n \times jm}$ by

$$\mathcal{K}_j(A, B) \triangleq [B \quad AB \quad \dots \quad A^{j-1}B]. \quad (12.9.19)$$

Note that $\mathcal{O}(A, C) = \mathcal{O}_n(A, C)$ and $\mathcal{K}(A, B) = \mathcal{K}_n(A, B)$. Furthermore, define the Markov block-Hankel matrix $\mathcal{H}_{i,j,k}(G) \in \mathbb{R}^{il \times jm}$ of G by

$$\mathcal{H}_{i,j,k}(G) \triangleq \mathcal{O}_i(A, C)A^k\mathcal{K}_j(A, B). \quad (12.9.20)$$

Note that $\mathcal{H}_{i,j,k}(G)$ is the block-Hankel matrix of Markov parameters given by

$$\begin{aligned} \mathcal{H}_{i,j,k}(G) &= \begin{bmatrix} CA^k B & CA^{k+1}B & CA^{k+2}B & \dots & CA^{k+j-1}B \\ CA^{k+1}B & CA^{k+2}B & \ddots & \ddots & \ddots \\ CA^{k+2}B & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ CA^{k+i-1}B & \ddots & \ddots & \ddots & CA^{k+j+i-2}B \end{bmatrix} \\ &= \begin{bmatrix} H_{k+1} & H_{k+2} & H_{k+3} & \dots & H_{k+j} \\ H_{k+2} & H_{k+3} & \ddots & \ddots & \ddots \\ H_{k+3} & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \\ H_{k+i} & \ddots & \ddots & \ddots & H_{k+j+i-1} \end{bmatrix}. \end{aligned} \quad (12.9.21)$$

Note that

$$\mathcal{H}_{i,j,0}(G) = \mathcal{O}_i(A, C)\mathcal{K}_j(A, B) \quad (12.9.22)$$

and

$$\mathcal{H}_{i,j,1}(G) = \mathcal{O}_i(A, C)A\mathcal{K}_j(A, B). \quad (12.9.23)$$

Furthermore, define

$$\mathcal{H}(G) \triangleq \mathcal{H}_{n,n,0}(G) = \mathcal{O}(A, C)\mathcal{K}(A, B). \quad (12.9.24)$$

The following result provides a MIMO extension of Fact 4.8.8.

Proposition 12.9.11. Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$, where $A \in \mathbb{R}^{n \times n}$. Then, the following statements are equivalent:

- i) The realization $\left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$ is controllable and observable.
- ii) $\text{rank } \mathcal{H}(G) = n$.
- iii) For all $i, j \geq n$, $\text{rank } \mathcal{H}_{i,j,0}(G) = n$.
- iv) There exist $i, j \geq n$ such that $\text{rank } \mathcal{H}_{i,j,0}(G) = n$.

Proof. The equivalence of ii), iii), and iv) follows from Fact 2.11.7. To prove i) $\implies$ ii), note that, since the $n \times n$ matrices $\mathcal{O}^T(A, C)\mathcal{O}(A, C)$ and $\mathcal{K}(A, B)\mathcal{K}^T(A, B)$ are positive definite, it follows that

$$n = \text{rank } \mathcal{O}^T(A, C)\mathcal{O}(A, C)\mathcal{K}(A, B)\mathcal{K}^T(A, B) \leq \text{rank } \mathcal{H}(G) \leq n.$$

Conversely, $n = \text{rank } \mathcal{H}(G) \leq \min\{\text{rank } \mathcal{O}(A, C), \text{rank } \mathcal{K}(A, B)\} \leq n$. $\square$

Proposition 12.9.12. Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$, where $A \in \mathbb{R}^{n \times n}$, assume that $\left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$ is controllable and observable, and let $i, j \geq 1$ be such that $\text{rank } \mathcal{O}_i(A, C) = \text{rank } \mathcal{K}_j(A, B) = n$. Then,

$$A = \mathcal{O}_i^+(A, C)\mathcal{H}_{i,j,1}(G)\mathcal{K}_j^+(A, B), \quad (12.9.25)$$

$$B = \mathcal{K}_j(A, B) \begin{bmatrix} I_m \\ 0_{(j-1)n \times m} \end{bmatrix}, \quad (12.9.26)$$

$$C = \begin{bmatrix} I_l & 0_{l \times (i-1)l} \end{bmatrix} \mathcal{O}_i(A, C). \quad (12.9.27)$$

Proposition 12.9.13. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, let $i, j \geq 1$, define $n \triangleq \text{rank } \mathcal{H}_{i,j,0}(G)$, and let $L \in \mathbb{R}^{l \times n}$ and $R \in \mathbb{R}^{n \times jm}$ be such that $\mathcal{H}_{i,j,0}(G) = LR$. Then, the realization

$$G \sim \left[\begin{array}{c|c} L^+ \mathcal{H}_{i,j,1}(G) R^+ & R \begin{bmatrix} I_m \\ 0_{(j-1)n \times m} \end{bmatrix} \\ \hline \begin{bmatrix} I_l & 0_{l \times (i-1)l} \end{bmatrix} L & 0 \end{array} \right] \quad (12.9.28)$$

is controllable and observable.

A rational transfer function $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$ can have realizations of different orders. For example, letting

$$A = 1, \quad B = 1, \quad C = 1, \quad D = 0$$

and

$$\hat{A} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad \hat{B} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}, \quad \hat{C} = \begin{bmatrix} 1 & 0 \end{bmatrix}, \quad \hat{D} = 0,$$

it follows that

$$G(s) = C(sI - A)^{-1}B + D = \hat{C}(sI - \hat{A})^{-1}\hat{B} + \hat{D} = \frac{1}{s-1}.$$

Generally, it is desirable to find realizations whose order is as small as possible.

Definition 12.9.14. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, and assume that $G \sim \left[\frac{A}{C} \middle| \frac{B}{D} \right]$. Then, $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ is a *minimal realization* of G if its order is less than or equal to the order of every realization of G . In this case, we write

$$G \stackrel{\min}{\sim} \left[\frac{A}{C} \middle| \frac{B}{D} \right]. \quad (12.9.29)$$

Note that the minimality of a realization is independent of D .

The following result shows that the controllable and observable realization $\left[\frac{A_1}{C_1} \middle| \frac{B_1}{D_1} \right]$ of G in Proposition 12.9.10 is, in fact, minimal.

Corollary 12.9.15. Let $G \in \mathbb{R}^{l \times m}(s)$, and assume that $G \sim \left[\frac{A}{C} \middle| \frac{B}{D} \right]$. Then, $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ is minimal if and only if it is controllable and observable.

Proof. To prove necessity, suppose that $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ is either not controllable or not observable. Then, Proposition 12.9.10 can be used to construct a realization of G of order less than n . Hence, $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ is not minimal.

To prove sufficiency, assume that $A \in \mathbb{R}^{n \times n}$, and assume that $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ is not minimal. Hence, G has a minimal realization $\left[\frac{\hat{A}}{\hat{C}} \middle| \frac{\hat{B}}{\hat{D}} \right]$ of order $\hat{n} < n$. Since the Markov parameters of G are independent of the realization, it follows from Proposition 12.9.11 that $\text{rank } \mathcal{H}(G) = \hat{n} < n$. However, since $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ is observable and controllable, it follows from Proposition 12.9.11 that $\text{rank } \mathcal{H}(G) = n$, which is a contradiction. $\square$

Theorem 12.9.16. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\frac{A}{C} \middle| \frac{B}{D} \right]$ and $A \in \mathbb{R}^{n \times n}$. Then,

$$\text{poles}(G) \subseteq \text{spec}(A) \quad (12.9.30)$$

and

$$\text{mpoles}(G) \subseteq \text{mspec}(A). \quad (12.9.31)$$

Furthermore, the following statements are equivalent:

- i) $G \stackrel{\min}{\sim} \left[\frac{A}{C} \middle| \frac{B}{D} \right]$.
- ii) $\text{Mcdeg}(G) = n$.
- iii) $\text{mpoles}(G) = \text{mspec}(A)$.

Proof. See [1150, p. 319]. $\square$

Definition 12.9.17. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \stackrel{\min}{\sim} \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. Then, G is (asymptotically stable, semistable, Lyapunov stable) if A is.

Proposition 12.9.18. Let $G = p/q \in \mathbb{R}_{\text{prop}}(s)$, where $p, q \in \mathbb{R}[s]$, and assume that p and q are coprime. Then, G is (asymptotically stable, semistable, Lyapunov stable) if and only if q is.

Proposition 12.9.19. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$. Then, G is (asymptotically stable, semistable, Lyapunov stable) if and only if every entry of G is.

Definition 12.9.20. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \stackrel{\min}{\sim} \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ and A is asymptotically stable. Then, the realization $\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ is *balanced* if the controllability and observability Gramians (12.7.2) and (12.4.2) are diagonal and equal.

Proposition 12.9.21. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \stackrel{\min}{\sim} \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ and A is asymptotically stable. Then, there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that the realization $G \sim \left[\begin{array}{c|c} SAS^{-1} & SB \\ \hline CS^{-1} & D \end{array} \right]$ is balanced.

Proof. It follows from Corollary 8.3.7 that there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that SQS^T and $S^{-T}PS^{-1}$ are diagonal, where Q and P are the controllability and observability Gramians (12.7.2) and (12.4.2). Hence, the realization $\left[\begin{array}{c|c} SAS^{-1} & SB \\ \hline CS^{-1} & D \end{array} \right]$ is balanced. $\square$

12.10 Zeros

In Section 4.7 the Smith-McMillan decomposition is used to define transmission zeros and blocking zeros of a transfer function $G(s)$. We now define the invariant zeros of a realization of $G(s)$ and relate these zeros to the transmission zeros. These zeros are related to the Smith zeros of a polynomial matrix as well as the spectrum of a pencil.

Definition 12.10.1. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. Then, the *Rosenbrock system matrix* $\mathcal{Z} \in \mathbb{R}^{(n+l) \times (n+m)}[s]$ is the polynomial matrix

$$\mathcal{Z}(s) \triangleq \begin{bmatrix} sI - A & B \\ C & -D \end{bmatrix}. \quad (12.10.1)$$

Furthermore, $z \in \mathbb{C}$ is an *invariant zero* of the realization $\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ if

$$\text{rank } \mathcal{Z}(z) < \text{rank } \mathcal{Z}. \quad (12.10.2)$$

Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$ and $A \in \mathbb{R}^{n \times n}$, and note that $\mathcal{Z}$ is the pencil

$$\mathcal{Z}(s) = P \begin{bmatrix} A & -B \\ -C & D \end{bmatrix} \begin{bmatrix} I_n & 0 \\ 0 & 0 \end{bmatrix} (s) \quad (12.10.3)$$

$$= s \begin{bmatrix} I_n & 0 \\ 0 & 0 \end{bmatrix} - \begin{bmatrix} A & -B \\ -C & D \end{bmatrix}. \quad (12.10.4)$$

Thus,

$$\text{Szeros}(\mathcal{Z}) = \text{spec} \left(\begin{bmatrix} A & -B \\ -C & D \end{bmatrix}, \begin{bmatrix} I_n & 0 \\ 0 & 0 \end{bmatrix} \right) \quad (12.10.5)$$

and

$$\text{mSzeros}(\mathcal{Z}) = \text{mspec} \left(\begin{bmatrix} A & -B \\ -C & D \end{bmatrix}, \begin{bmatrix} I_n & 0 \\ 0 & 0 \end{bmatrix} \right). \quad (12.10.6)$$

Hence, we define the set of invariant zeros of $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ by

$$\text{izeros} \left(\left[\frac{A}{C} \middle| \frac{B}{D} \right] \right) \triangleq \text{Szeros}(\mathcal{Z})$$

and the multiset of invariant zeros of $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ by

$$\text{mizeros} \left(\left[\frac{A}{C} \middle| \frac{B}{D} \right] \right) \triangleq \text{mSzeros}(\mathcal{Z}).$$

Note that $P \begin{bmatrix} A & -B \\ -C & D \end{bmatrix} \begin{bmatrix} I_n & 0 \\ 0 & 0 \end{bmatrix}$ is regular if and only if $\text{rank } \mathcal{Z} = n + \min\{l, m\}$.

The following result shows that a strictly proper transfer function with full-state observation or full-state actuation has no invariant zeros.

Proposition 12.10.2. Let $G \in \mathbb{R}^{l \times m}(s)$, where $G \sim \left[\frac{A}{C} \middle| \frac{B}{0} \right]$ and $A \in \mathbb{R}^{n \times n}$. Then, the following statements hold:

- i) If $m = n$ and B is nonsingular, then $\text{rank } \mathcal{Z} = n + \text{rank } C$ and $\left[\frac{A}{C} \middle| \frac{B}{0} \right]$ has no invariant zeros.
- ii) If $l = n$ and C is nonsingular, then $\text{rank } \mathcal{Z} = n + \text{rank } B$ and $\left[\frac{A}{C} \middle| \frac{B}{0} \right]$ has no invariant zeros.
- iii) If $m = n$ and B is nonsingular, then $P \begin{bmatrix} I_n & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A & -B \\ -C & 0 \end{bmatrix}$ is regular if and only if $\text{rank } C = \min\{l, n\}$.
- iv) If $l = n$ and C is nonsingular, then $P \begin{bmatrix} I_n & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} A & -B \\ -C & 0 \end{bmatrix}$ is regular if and only if $\text{rank } B = \min\{m, n\}$.

It is useful to note that, for all $s \notin \text{spec}(A)$,

$$\mathcal{Z}(s) = \begin{bmatrix} I & 0 \\ C(sI - A)^{-1} & I \end{bmatrix} \begin{bmatrix} sI - A & B \\ 0 & -G(s) \end{bmatrix} \quad (12.10.7)$$

$$= \begin{bmatrix} sI - A & 0 \\ C & -G(s) \end{bmatrix} \begin{bmatrix} I & (sI - A)^{-1}B \\ 0 & I \end{bmatrix}. \quad (12.10.8)$$

Proposition 12.10.3. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. If $s \notin \text{spec}(A)$, then

$$\text{rank } \mathcal{Z}(s) = n + \text{rank } G(s). \quad (12.10.9)$$

Furthermore,

$$\text{rank } \mathcal{Z} = n + \text{rank } G. \quad (12.10.10)$$

Proof. For $s \notin \text{spec}(A)$, (12.10.9) follows from (12.10.7). Therefore, it follows from Proposition 4.3.6 and Proposition 4.7.8 that

$$\begin{aligned} \text{rank } \mathcal{Z} &= \max_{s \in \mathbb{C}} \text{rank } \mathcal{Z}(s) \\ &= \max_{s \in \mathbb{C} \setminus \text{spec}(A)} \text{rank } \mathcal{Z}(s) \\ &= n + \max_{s \in \mathbb{C} \setminus \text{spec}(A)} \text{rank } G(s) \\ &= n + \text{rank } G. \end{aligned} \quad \square$$

Note that the realization in Proposition 12.10.3 is not assumed to be minimal. Therefore, $P \left[\begin{array}{c|c} A & -B \\ \hline -C & D \end{array} \right] \left[\begin{array}{c} I_n \\ 0 \end{array} \right]$ is (regular, singular) for one realization of G if and only if it is (regular, singular) for every realization of G . In fact, the following result shows that $P \left[\begin{array}{c|c} A & -B \\ \hline -C & D \end{array} \right] \left[\begin{array}{c} I_n \\ 0 \end{array} \right]$ is regular if and only if G has full rank.

Corollary 12.10.4. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. Then, $P \left[\begin{array}{c|c} A & -B \\ \hline -C & D \end{array} \right] \left[\begin{array}{c} I_n \\ 0 \end{array} \right]$ is regular if and only if $\text{rank } G = \min\{l, m\}$.

In the SISO case, it follows from (12.10.7) and (12.10.8) that, for all $s \in \mathbb{C} \setminus \text{spec}(A)$,

$$\det \mathcal{Z}(s) = -[\det(sI - A)]G(s). \quad (12.10.11)$$

Consequently, for all $s \in \mathbb{C}$,

$$\det \mathcal{Z}(s) = -C(sI - A)^A B - \det(sI - A)D. \quad (12.10.12)$$

The identity (12.10.12) also follows from Fact 2.14.2.

In particular, if $s \in \text{spec}(A)$, then

$$\det \mathcal{Z}(s) = -C(sI - A)^A B. \quad (12.10.13)$$

If, in addition, $n \geq 2$ and $\text{rank}(sI - A) \leq n - 2$, then it follows from Fact 2.16.8 that $(sI - A)^A = 0$, and thus

$$\det \mathcal{Z}(s) = 0. \quad (12.10.14)$$

Alternatively, in the case $n = 1$, it follows that, for all $s \in \mathbb{C}$, $(sI - A)^A = 1$, and thus, for all $s \in \mathbb{C}$,

$$\det \mathcal{Z}(s) = -CB - (sI - A)D. \quad (12.10.15)$$

Next, it follows from (12.10.11) and (12.10.12) that

$$G(s) = \frac{C(sI - A)^A B + \det(sI - A)D}{\det(sI - A)} \quad (12.10.16)$$

$$= \frac{-\det \mathcal{Z}(s)}{\det(sI - A)}. \quad (12.10.17)$$

Consequently, $G \neq 0$ if and only if $\det \mathcal{Z} \neq 0$.

We now have the following result for scalar transfer functions.

Corollary 12.10.5. Let $G \in \mathbb{R}_{\text{prop}}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. Then, the following statements are equivalent:

- i) $P \left[\begin{array}{c|c} A & -B \\ \hline -C & D \end{array} \right], \left[\begin{array}{c|c} I_n & 0 \\ \hline 0 & 0 \end{array} \right]$ is regular.
- ii) $G \neq 0$.
- iii) $\text{rank } G = 1$.
- iv) $\det \mathcal{Z} \neq 0$.
- v) $\text{rank } \mathcal{Z} = n + 1$.
- vi) $C(sI - A)^A B + \det(sI - A)D$ is not the zero polynomial.

In this case,

$$\text{mizeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right) = \text{mroots}(\det \mathcal{Z}) \quad (12.10.18)$$

and

$$\text{mizeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right) = \text{mtzeros}(G) \cup [\text{mspec}(A) \setminus \text{mpoles}(G)]. \quad (12.10.19)$$

If, in addition, $G \stackrel{\min}{\sim} \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$, then

$$\text{mizeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right) = \text{mtzeros}(G). \quad (12.10.20)$$

Now, suppose that G is square, that is, $l = m$. Then, it follows from (12.10.7) and (12.10.8) that, for all $s \in \mathbb{C} \setminus \text{spec}(A)$,

$$\det \mathcal{Z}(s) = (-1)^l \det(sI - A) \det G(s), \quad (12.10.21)$$

and thus

$$\det G(s) = \frac{(-1)^l \det \mathcal{Z}(s)}{\det(sI - A)}. \quad (12.10.22)$$

Furthermore, for all $s \in \mathbb{C}$,

$$[\det(sI - A)]^{l-1} \det \mathcal{Z}(s) = (-1)^l \det [C(sI - A)^A B + \det(sI - A)D]. \quad (12.10.23)$$

Hence, for all $s \in \text{spec}(A)$, it follows that

$$\det [C(sI - A)^A B] = 0. \quad (12.10.24)$$

We thus have the following result for square transfer functions G that satisfy $\det G \neq 0$.

Corollary 12.10.6. Let $G \in \mathbb{R}_{\text{prop}}^{l \times l}(s)$, where $G \sim \left[\frac{A}{C} \middle| \frac{B}{D} \right]$. Then, the following statements are equivalent:

- i) $P_{\left[\frac{A}{C} \middle| \frac{B}{D} \right], \left[\begin{smallmatrix} I_n & 0 \\ 0 & 0 \end{smallmatrix} \right]}$ is regular.
- ii) $\det G \neq 0$.
- iii) $\text{rank } G = l$.
- iv) $\det \mathcal{Z} \neq 0$.
- v) $\text{rank } \mathcal{Z} = n + l$.
- vi) $\det[C(sI - A)^A B + \det(sI - A)D]$ is not the zero polynomial.

In this case,

$$\text{mizeros}\left(\left[\frac{A}{C} \middle| \frac{B}{D}\right]\right) = \text{mroots}(\det \mathcal{Z}), \quad (12.10.25)$$

$$\text{mizeros}\left(\left[\frac{A}{C} \middle| \frac{B}{D}\right]\right) = \text{mtzeros}(G) \cup [\text{mspec}(A) \setminus \text{mpoles}(G)], \quad (12.10.26)$$

and

$$\text{izeros}\left(\left[\frac{A}{C} \middle| \frac{B}{D}\right]\right) = \text{tzeros}(G) \cup [\text{spec}(A) \setminus \text{poles}(G)]. \quad (12.10.27)$$

If, in addition, $G \stackrel{\min}{\sim} \left[\frac{A}{C} \middle| \frac{B}{D} \right]$, then

$$\text{mizeros}\left(\left[\frac{A}{C} \middle| \frac{B}{D}\right]\right) = \text{mtzeros}(G). \quad (12.10.28)$$

Example 12.10.7. Consider $G \in \mathbb{R}^{2 \times 2}(s)$ defined by

$$G(s) \triangleq \begin{bmatrix} \frac{s-1}{s+1} & 0 \\ 0 & \frac{s+1}{s-1} \end{bmatrix}. \quad (12.10.29)$$

Then, the Smith-McMillan form of G is given by

$$G(s) \triangleq S_1(s) \begin{bmatrix} \frac{1}{s^2-1} & 0 \\ 0 & s^2-1 \end{bmatrix} S_2(s), \quad (12.10.30)$$

where $S_1, S_2 \in \mathbb{R}^{2 \times 2}[s]$ are the unimodular matrices

$$S_1(s) \triangleq \begin{bmatrix} (s-1)^2 & -1 \\ -\frac{1}{4}(s+1)^2(s-2) & \frac{1}{4}(s+2) \end{bmatrix} \quad (12.10.31)$$

and

$$S_2(s) \triangleq \begin{bmatrix} \frac{1}{4}(s-1)^2(s+2) & (s+1)^2 \\ \frac{1}{4}(s-2) & 1 \end{bmatrix}. \quad (12.10.32)$$

Thus, $\text{mpoles}(G) = \text{mtzeros}(G) = \{1, -1\}$. Furthermore, a minimal realization of G is given by

$$G \stackrel{\min}{\sim} \left[\begin{array}{cc|cc} -1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 \\ \hline -2 & 0 & 1 & 0 \\ 0 & 2 & 0 & 1 \end{array} \right]. \quad (12.10.33)$$

Finally, note that $\det \mathcal{Z}(s) = (-1)^2 \det(sI - A) \det G = s^2 - 1$, which confirms (12.10.28).

Theorem 12.10.8. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. Then,

$$\text{izeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right) \setminus \text{spec}(A) \subseteq \text{tzeros}(G) \quad (12.10.34)$$

and

$$\text{tzeros}(G) \setminus \text{poles}(G) \subseteq \text{izeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right). \quad (12.10.35)$$

If, in addition, $G \stackrel{\min}{\sim} \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$, then

$$\text{izeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right) \setminus \text{poles}(G) = \text{tzeros}(G) \setminus \text{poles}(G). \quad (12.10.36)$$

Proof. To prove (12.10.34), let $z \in \text{izeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right) \setminus \text{spec}(A)$. Since $z \notin \text{spec}(A)$ it follows from Theorem 12.9.16 that $z \notin \text{poles}(G)$. It now follows from Proposition 12.10.3 that $n + \text{rank } G(z) = \text{rank } \mathcal{Z}(z) < \text{rank } \mathcal{Z} = n + \text{rank } G$, which implies that $\text{rank } G(z) < \text{rank } G$. Thus, $z \in \text{tzeros}(G)$.

To prove (12.10.35), let $z \in \text{tzeros}(G) \setminus \text{poles}(G)$. Then, it follows from Proposition 12.10.3 that $\text{rank } \mathcal{Z}(z) = n + \text{rank } G(z) < n + \text{rank } G = \text{rank } \mathcal{Z}$, which implies that $z \in \text{izeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right)$. The last statement follows from (12.10.34), (12.10.35), and Theorem 12.9.16. $\square$

The following result is a stronger form of Theorem 12.10.8.

Theorem 12.10.9. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$, let $S \in \mathbb{R}^{n \times n}$, assume that S is nonsingular, and let A , B , and C have the form (12.9.16), (12.9.17), where $\left(\begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix}, \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} \right)$ is controllable and $\left(\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}, \begin{bmatrix} C_1 & C_3 \end{bmatrix} \right)$ is observable. Then,

$$\text{mtzeros}(G) = \text{mizeros} \left(\left[\begin{array}{c|c} A_1 & B_1 \\ \hline C_1 & D \end{array} \right] \right) \quad (12.10.37)$$

and

$$\text{mizeros} \left(\left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right] \right) = \text{mspec}(A_2) \cup \text{mspec}(A_3) \cup \text{mspec}(A_4) \cup \text{mtzeros}(G). \quad (12.10.38)$$

Proof. Defining $\mathcal{Z}$ by (12.10.1), note that, in the notation of Proposition 12.9.10, $\mathcal{Z}$ has the same Smith form as

$$\tilde{\mathcal{Z}} \triangleq \begin{bmatrix} sI - A_4 & -A_{43} & 0 & 0 & 0 \\ 0 & sI - A_3 & 0 & 0 & 0 \\ -A_{24} & -A_{23} & sI - A_2 & -A_{21} & B_2 \\ 0 & -A_{13} & 0 & sI - A_1 & B_1 \\ 0 & C_3 & 0 & C_1 & -D \end{bmatrix}.$$

Hence, it follows from Proposition 12.10.3 that $\text{rank } \mathcal{Z} = \text{rank } \tilde{\mathcal{Z}} = n + r$, where $r \triangleq \text{rank } G$. Let $\tilde{p}_1, \dots, \tilde{p}_{n+r}$ be the Smith polynomials of $\tilde{\mathcal{Z}}$. Then, since $\tilde{p}_{n+r}$ is the monic greatest common divisor of all $(n + r) \times (n + r)$ subdeterminants of $\tilde{\mathcal{Z}}$, it follows that $\tilde{p}_{n+r} = \chi_{A_1} \chi_{A_2} \chi_{A_3} p_r$, where p_r is the r th Smith polynomial of $\begin{bmatrix} sI - A_1 & B_1 \\ C_1 & -D \end{bmatrix}$. Therefore,

$$\text{mSzeros}(\mathcal{Z}) = \text{mspec}(A_2) \cup \text{mspec}(A_3) \cup \text{mspec}(A_4) \cup \text{mSzeros} \left(\begin{bmatrix} sI - A_1 & B_1 \\ C_1 & -D \end{bmatrix} \right).$$

Next, using the Smith-McMillan decomposition Theorem 4.7.5, it follows that there exist unimodular matrices $S_1 \in \mathbb{R}^{l \times l}[s]$ and $S_2 \in \mathbb{R}^{m \times m}[s]$ such that $G = S_1 D_0^{-1} N_0 S_2$, where

$$D_0 \triangleq \begin{bmatrix} q_1 & & 0 \\ & \ddots & \\ & & q_r \\ 0 & & I_{l-r} \end{bmatrix}, \quad N_0 \triangleq \begin{bmatrix} p_1 & & 0 \\ & \ddots & \\ & & p_r \\ 0 & & 0_{(l-r) \times (m-r)} \end{bmatrix}.$$

Now, define the polynomial matrix $\hat{\mathcal{Z}} \in \mathbb{R}^{(n+l) \times (n+m)}[s]$ by

$$\hat{\mathcal{Z}} \triangleq \begin{bmatrix} I_{n-l} & 0_{(n-l) \times l} & 0_{(n-l) \times m} \\ 0_{l \times (n-l)} & D_0 & N_0 S_2 \\ 0_{l \times (n-l)} & S_1 & 0_{l \times m} \end{bmatrix}.$$

Since S_1 is unimodular, it follows that the Smith form $\mathcal{S}$ of $\hat{\mathcal{Z}}$ is given by

$$\mathcal{S} = \begin{bmatrix} I_n & 0_{n \times m} \\ 0_{l \times n} & N_0 \end{bmatrix}.$$

Consequently, $\text{mSzeros}(\hat{\mathcal{Z}}) = \text{mSzeros}(\mathcal{S}) = \text{mtzeros}(G)$.

Next, note that

$$\text{rank} \begin{bmatrix} I_{n-l} & 0_{(n-l) \times l} & 0_{(n-l) \times m} \\ 0_{l \times (n-l)} & D_0 & N_0 S_2 \end{bmatrix} = \text{rank} \begin{bmatrix} I_{n-l} & 0_{(n-l) \times l} \\ 0_{l \times (n-l)} & D_0 \\ 0_{l \times (n-l)} & S_1 \end{bmatrix} = n$$

and that

$$G = \begin{bmatrix} 0_{l \times (n-l)} & S_1 & 0_{l \times m} \end{bmatrix} \begin{bmatrix} I_{n-l} & 0_{(n-l) \times l} \\ 0_{l \times (n-l)} & D_0 \end{bmatrix}^{-1} \begin{bmatrix} 0_{(n-l) \times m} \\ N_0 S_2 \end{bmatrix}.$$

Furthermore, $G \stackrel{\min}{\sim} \left[\frac{A_1}{C_1} \middle| \frac{B_1}{D} \right]$, Consequently, $\hat{\mathcal{Z}}$ and $\begin{bmatrix} sI - A_1 & B_1 \\ C_1 & D \end{bmatrix}$ have no decoupling zeros [1144, pp. 64–70], and it thus follows from Theorem 3.1 of [1144, p.

106] that $\hat{\mathcal{Z}}$ and $\begin{bmatrix} sI - A_1 & B_1 \\ C_1 & -D \end{bmatrix}$ have the same Smith form. Thus,

$$\text{mSzeros} \left(\begin{bmatrix} sI - A_1 & B_1 \\ C_1 & -D \end{bmatrix} \right) = \text{mSzeros}(\hat{\mathcal{Z}}) = \text{mtzeros}(G).$$

Consequently,

$$\text{mizeros} \left(\left[\frac{A_1}{C_1} \middle| \frac{B_1}{D} \right] \right) = \text{mSzeros} \left(\begin{bmatrix} sI - A_1 & B_1 \\ C_1 & -D \end{bmatrix} \right) = \text{mtzeros}(G),$$

which proves (12.10.37).

Finally, to prove (12.10.34) note that

$$\begin{aligned} \text{mizeros} \left(\left[\frac{A}{C} \middle| \frac{B}{D} \right] \right) &= \text{mSzeros}(\mathcal{Z}) \\ &= \text{mspec}(A_2) \cup \text{mspec}(A_3) \cup \text{mspec}(A_4) \cup \text{mSzeros} \left(\begin{bmatrix} sI - A_1 & B_1 \\ -C_1 & -D \end{bmatrix} \right) \\ &= \text{mspec}(A_2) \cup \text{mspec}(A_3) \cup \text{mspec}(A_4) \cup \text{mtzeros}(G). \end{aligned} \quad \square$$

Proposition 12.10.10. Equivalent realizations have the same invariant zeros. Furthermore, invariant zeros are not changed by full-state feedback.

Proof. Let $u = Kx + v$, which leads to the rational transfer function

$$G_K \sim \left[\frac{A + BK}{C + DK} \middle| \frac{B}{D} \right]. \quad (12.10.39)$$

Since

$$\begin{bmatrix} zI - (A + BK) & B \\ C + DK & -D \end{bmatrix} = \begin{bmatrix} zI - A & B \\ C & -D \end{bmatrix} \begin{bmatrix} I & 0 \\ -K & I \end{bmatrix}, \quad (12.10.40)$$

it follows that $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ and $\left[\frac{A + BK}{C + DK} \middle| \frac{B}{D} \right]$ have the same invariant zeros. $\square$

The following result provides an interpretation of condition *i*) of Theorem 12.17.9.

Proposition 12.10.11. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\frac{A}{C} \middle| \frac{B}{D} \right]$, and assume that $R \triangleq D^T D$ is positive definite. Then, the following statements hold:

i) $\text{rank } \mathcal{Z} = n + m$.

ii) $z \in \mathbb{C}$ is an invariant zero of $\left[\frac{A}{C} \middle| \frac{B}{D} \right]$ if and only if z is an unobservable eigenvalue of $(A - BR^{-1}D^T C, [I - DR^{-1}D^T]C)$.

Proof. To prove *i*), assume that $\text{rank } \mathcal{Z} < n + m$. Then, for every $s \in \mathbb{C}$, there exists a nonzero vector $\begin{bmatrix} x \\ y \end{bmatrix} \in \mathcal{N}[\mathcal{Z}(s)]$, that is,

$$\begin{bmatrix} sI - A & B \\ C & -D \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = 0.$$

Consequently, $Cx - Dy = 0$, which implies that $D^T Cx - Ry = 0$, and thus $y = R^{-1}D^T Cx$. Furthermore, since $(sI - A + BR^{-1}D^T C)x = 0$, choosing $s \notin$

$\text{spec}(A - BR^{-1}D^TC)$ yields $x = 0$, and thus $y = 0$, which is a contradiction.

To prove *ii*), note that z is an invariant zero of $\left[\frac{A}{C} \middle| \frac{B}{D}\right]$ if and only if $\text{rank } \mathcal{Z}(z) < n + m$, which holds if and only if there exists a nonzero vector $\begin{bmatrix} x \\ y \end{bmatrix} \in \mathcal{N}[\mathcal{Z}(z)]$. This condition is equivalent to

$$\begin{bmatrix} sI - A + BR^{-1}D^TC \\ (I - DR^{-1}D^T)C \end{bmatrix} x = 0,$$

where $x \neq 0$. This last condition is equivalent to the fact that z is an unobservable eigenvalue of $(A - BR^{-1}D^TC, [I - DR^{-1}D^T]C)$. $\square$

Corollary 12.10.12. Assume that $R \triangleq D^TD$ is positive definite, and assume that $(A - BR^{-1}D^TC, [I - DR^{-1}D^T]C)$ is observable. Then, $\left[\frac{A}{C} \middle| \frac{B}{D}\right]$ has no invariant zeros.

12.11 H_2 System Norm

Consider the system

$$\dot{x}(t) = Ax(t) + Bu(t), \quad (12.11.1)$$

$$y(t) = Cx(t), \quad (12.11.2)$$

where $A \in \mathbb{R}^{n \times n}$ is asymptotically stable, $B \in \mathbb{R}^{n \times m}$, and $C \in \mathbb{R}^{l \times n}$. Then, for all $t \geq 0$, the impulse response function defined by (12.1.18) is given by

$$H(t) = Ce^{tA}B. \quad (12.11.3)$$

The L_2 norm of $H(\cdot)$ is given by

$$\|H\|_{L_2} \triangleq \left(\int_0^\infty \|H(t)\|_F^2 dt \right)^{1/2}. \quad (12.11.4)$$

The following result provides expressions for $\|H(\cdot)\|_{L_2}$ in terms of the controllability and observability Gramians.

Theorem 12.11.1. Assume that A is asymptotically stable. Then, the L_2 norm of H is given by

$$\|H\|_{L_2}^2 = \text{tr } CQC^T = \text{tr } B^TPB, \quad (12.11.5)$$

where $Q, P \in \mathbb{R}^{n \times n}$ satisfy

$$AQ + QA^T + BB^T = 0, \quad (12.11.6)$$

$$A^TP + PA + C^TC = 0. \quad (12.11.7)$$

Proof. Note that

$$\|H\|_{L_2}^2 = \int_0^\infty \text{tr } Ce^{tA}BB^Te^{tA^T}C^T dt = \text{tr } CQC^T,$$

where Q satisfies (12.11.6). The dual expression (12.11.7) follows in a similar manner or by noting that

$$\begin{aligned}\operatorname{tr} CQC^T &= \operatorname{tr} C^T CQ = -\operatorname{tr} (A^T P + PA)Q \\ &= -\operatorname{tr} (AQ + QA^T)P = \operatorname{tr} BB^T P = \operatorname{tr} B^T PB.\end{aligned}\quad \square$$

For the following definition, note that

$$\|G(s)\|_F = [\operatorname{tr} G(s)G^*(s)]^{1/2}. \quad (12.11.8)$$

Definition 12.11.2. The H_2 norm of $G \in \mathbb{R}^{l \times m}(s)$ is the nonnegative number

$$\|G\|_{H_2} \triangleq \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} \|G(j\omega)\|_F^2 d\omega \right)^{1/2}. \quad (12.11.9)$$

The following result is *Parseval's theorem*, which relates the L_2 norm of the impulse response function to the H_2 norm of its transform.

Theorem 12.11.3. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{smallmatrix} A & B \\ C & 0 \end{smallmatrix} \right]$, and assume that $A \in \mathbb{R}^{n \times n}$ is asymptotically stable. Then,

$$\int_0^{\infty} H(t)H^T(t) dt = \frac{1}{2\pi} \int_{-\infty}^{\infty} G(j\omega)G^*(j\omega) d\omega. \quad (12.11.10)$$

Therefore,

$$\|H\|_{L_2} = \|G\|_{H_2}. \quad (12.11.11)$$

Proof. First note that

$$G(s) = \mathcal{L}\{H(t)\} = \int_0^{\infty} H(t)e^{-st} dt$$

and that

$$H(t) = \frac{1}{2\pi} \int_{-\infty}^{\infty} G(j\omega)e^{j\omega t} d\omega.$$

Hence,

$$\begin{aligned}\int_0^{\infty} H(t)H^T(t)e^{-st} dt &= \int_0^{\infty} \left(\frac{1}{2\pi} \int_{-\infty}^{\infty} G(j\omega)e^{j\omega t} d\omega \right) H^T(t)e^{-st} dt \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} G(j\omega) \left(\int_0^{\infty} H^T(t)e^{-(s-j\omega)t} dt \right) d\omega \\ &= \frac{1}{2\pi} \int_{-\infty}^{\infty} G(j\omega)G^T(s-j\omega) d\omega.\end{aligned}$$

Setting $s = 0$ yields (12.11.7), while taking the trace of (12.11.10) yields (12.11.11). $\square$

Corollary 12.11.4. Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$, where $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$, and assume that $A \in \mathbb{R}^{n \times n}$ is asymptotically stable. Then,

$$\|G\|_{\text{H}_2}^2 = \|H\|_{\text{L}_2}^2 = \text{tr } CQC^T = \text{tr } B^T P B, \quad (12.11.12)$$

where $Q, P \in \mathbb{R}^{n \times n}$ satisfy (12.11.6) and (12.11.7), respectively.

The following corollary of Theorem 12.11.3 provides a frequency domain expression for the solution of the Lyapunov equation.

Corollary 12.11.5. Let $A \in \mathbb{R}^{n \times n}$, assume that A is asymptotically stable, let $B \in \mathbb{R}^{n \times m}$, and define $Q \in \mathbb{R}^{n \times n}$ by

$$Q = \frac{1}{2\pi} \int_{-\infty}^{\infty} (j\omega I - A)^{-1} B B^T (j\omega I - A)^{-*} d\omega. \quad (12.11.13)$$

Then, Q satisfies

$$AQ + QA^T + BB^T = 0. \quad (12.11.14)$$

Proof. The result follows directly from Theorem 12.11.3 with $H(t) = e^{tA}B$ and $G(s) = (sI - A)^{-1}B$. Alternatively, it follows from (12.11.14) that

$$\int_{-\infty}^{\infty} (j\omega I - A)^{-1} d\omega Q + Q \int_{-\infty}^{\infty} (j\omega I - A)^{-*} d\omega = \int_{-\infty}^{\infty} (j\omega I - A)^{-1} B B^T (j\omega I - A)^{-*} d\omega.$$

Assuming that A is diagonalizable with eigenvalues $\lambda_i = -\sigma_i + j\omega_i$, it follows that

$$\int_{-\infty}^{\infty} \frac{d\omega}{j\omega - \lambda_i} = \int_{-\infty}^{\infty} \frac{\sigma_i - j\omega}{\sigma_i^2 + \omega^2} d\omega = \frac{\sigma_i \pi}{|\sigma_i|} - j \lim_{r \rightarrow \infty} \int_{-r}^r \frac{\omega}{\sigma_i^2 + \omega^2} d\omega = \pi,$$

which implies that

$$\int_{-\infty}^{\infty} (j\omega I - A)^{-1} d\omega = \pi I_n,$$

which yields (12.11.13). See [309] for a proof of the general case. $\square$

Proposition 12.11.6. Let $G_1, G_2 \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$ be asymptotically stable rational transfer functions. Then,

$$\|G_1 + G_2\|_{\text{H}_2} \leq \|G_1\|_{\text{H}_2} + \|G_2\|_{\text{H}_2}. \quad (12.11.15)$$

Proof. Let $G_1 \stackrel{\text{min}}{\sim} \left[\begin{array}{c|c} A_1 & B_1 \\ \hline C_1 & 0 \end{array} \right]$ and $G_2 \stackrel{\text{min}}{\sim} \left[\begin{array}{c|c} A_2 & B_2 \\ \hline C_2 & 0 \end{array} \right]$, where $A_1 \in \mathbb{R}^{n_1 \times n_1}$ and $A_2 \in \mathbb{R}^{n_2 \times n_2}$. It follows from Proposition 12.13.2 that

$$G_1 + G_2 \sim \left[\begin{array}{cc|c} A_1 & 0 & B_1 \\ 0 & A_2 & B_2 \\ \hline C_1 & C_2 & 0 \end{array} \right].$$

Now, Theorem 12.11.3 implies that $\|G_1\|_{H_2} = \sqrt{\text{tr } C_1 Q_1 C_1^T}$ and $\|G_2\|_{H_2} = \sqrt{\text{tr } C_2 Q_2 C_2^T}$, where $Q_1 \in \mathbb{R}^{n_1 \times n_1}$ and $Q_2 \in \mathbb{R}^{n_2 \times n_2}$ are the unique positive-definite matrices satisfying $A_1 Q_1 + Q_1 A_1^T + B_1 B_1^T = 0$ and $A_2 Q_2 + Q_2 A_2^T + B_2 B_2^T = 0$. Furthermore,

$$\|G_1 + G_2\|_{H_2}^2 = \text{tr} \begin{bmatrix} C_1 & C_2 \end{bmatrix} Q \begin{bmatrix} C_1^T \\ C_2^T \end{bmatrix},$$

where $Q \in \mathbb{R}^{(n_1+n_2) \times (n_1+n_2)}$ is the unique, positive-semidefinite matrix satisfying

$$\begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix} Q + Q \begin{bmatrix} A_1 & 0 \\ 0 & A_2 \end{bmatrix}^T + \begin{bmatrix} B_1 \\ B_2 \end{bmatrix} \begin{bmatrix} B_1 \\ B_2 \end{bmatrix}^T = 0.$$

It can be seen that $Q = \begin{bmatrix} Q_1 & Q_{12} \\ Q_{12}^T & Q_2 \end{bmatrix}$, where Q_1 and Q_2 are as given above and where Q_{12} satisfies $A_1 Q_{12} + Q_{12} A_2^T + B_1 B_2^T = 0$. Now, using the Cauchy-Schwarz inequality (9.3.17) and *iii*) of Proposition 8.2.4, it follows that

$$\begin{aligned} \|G_1 + G_2\|_{H_2}^2 &= \text{tr}(C_1 Q_1 C_1^T + C_2 Q_2 C_2^T + C_2 Q_{12}^T C_1^T + C_1 Q_{12} C_2^T) \\ &= \|G_1\|_{H_2}^2 + \|G_2\|_{H_2}^2 + 2 \text{tr } C_1 Q_{12} Q_2^{-1/2} Q_2^{1/2} C_2^T \\ &\leq \|G_1\|_{H_2}^2 + \|G_2\|_{H_2}^2 + 2 \text{tr}(C_1 Q_{12} Q_2^{-1} Q_{12}^T C_1^T) \text{tr}(C_2 Q_2 C_2^T) \\ &\leq \|G_1\|_{H_2}^2 + \|G_2\|_{H_2}^2 + 2 \text{tr}(C_1 Q_1 C_1^T) \text{tr}(C_2 Q_2 C_2^T) \\ &= (\|G_1\|_{H_2} + \|G_2\|_{H_2})^2. \end{aligned} \quad \square$$

12.12 Harmonic Steady-State Response

The following result concerns the response of a linear system to a harmonic input.

Theorem 12.12.1. For $t \geq 0$, consider the linear system

$$\dot{x}(t) = Ax(t) + Bu(t), \quad (12.12.1)$$

with harmonic input

$$u(t) = \text{Re } u_0 e^{j\omega_0 t}, \quad (12.12.2)$$

where $u_0 \in \mathbb{C}^m$ and $\omega_0 \in \mathbb{R}$ is such that $j\omega_0 \notin \text{spec}(A)$. Then, $x(t)$ is given by

$$x(t) = e^{tA}(x(0) - \text{Re}[(j\omega_0 I - A)^{-1}Bu_0]) + \text{Re}[(j\omega_0 I - A)^{-1}Bu_0 e^{j\omega_0 t}]. \quad (12.12.3)$$

Proof. We have

$$\begin{aligned}
 x(t) &= e^{tA}x(0) + \int_0^t e^{(t-\tau)A} B \operatorname{Re}(u_0 e^{j\omega_0 \tau}) d\tau \\
 &= e^{tA}x(0) + e^{tA} \operatorname{Re} \left[\int_0^t e^{-\tau A} e^{j\omega_0 \tau} d\tau B u_0 \right] \\
 &= e^{tA}x(0) + e^{tA} \operatorname{Re} \left[\int_0^t e^{\tau(j\omega_0 I - A)} d\tau B u_0 \right] \\
 &= e^{tA}x(0) + e^{tA} \operatorname{Re} \left[(j\omega_0 I - A)^{-1} (e^{t(j\omega_0 I - A)} - I) B u_0 \right] \\
 &= e^{tA}x(0) + \operatorname{Re} [(j\omega_0 I - A)^{-1} (e^{j\omega_0 t I} - e^{tA}) B u_0] \\
 &= e^{tA}x(0) + \operatorname{Re} [(j\omega_0 I - A)^{-1} (-e^{tA}) B u_0] + \operatorname{Re} [(j\omega_0 I - A)^{-1} e^{j\omega_0 t} B u_0] \\
 &= e^{tA} (x(0) - \operatorname{Re} [(j\omega_0 I - A)^{-1} B u_0]) + \operatorname{Re} [(j\omega_0 I - A)^{-1} B u_0 e^{j\omega_0 t}]. \quad \square
 \end{aligned}$$

Theorem 12.12.1 shows that the total response $y(t)$ of the linear system $G \sim \begin{bmatrix} A & B \\ C & 0 \end{bmatrix}$ to a harmonic input can be written as $y(t) = y_{\text{trans}}(t) + y_{\text{hss}}(t)$, where the transient component

$$y_{\text{trans}}(t) \triangleq C e^{tA} (x(0) - \operatorname{Re} [(j\omega_0 I - A)^{-1} B u_0]) \quad (12.12.4)$$

depends on the initial condition and the input, and the harmonic steady-state component

$$y_{\text{hss}}(t) = \operatorname{Re} [G(j\omega_0) u_0 e^{j\omega_0 t}] \quad (12.12.5)$$

depends only on the input.

If A is asymptotically stable, then $\lim_{t \rightarrow \infty} y_{\text{trans}}(t) = 0$, and thus $y(t)$ approaches its harmonic steady-state component $y_{\text{hss}}(t)$ for large t . Since the harmonic steady-state component is sinusoidal, it follows that $y(t)$ does not converge in the usual sense.

Finally, if A is semistable, then it follows from *vii*) of Proposition 11.8.2 that

$$\lim_{t \rightarrow \infty} y_{\text{trans}}(t) = C(I - AA^\#)(x(0) - \operatorname{Re} [(j\omega_0 I - A)^{-1} B u_0]), \quad (12.12.6)$$

which represents a constant offset to the harmonic steady-state component.

In the SISO case, let $u_0 \triangleq a_0(\sin \phi_0 + j \cos \phi_0)$, and consider the input

$$u(t) = a_0 \sin(\omega_0 t + \phi_0) = \operatorname{Re} u_0 e^{j\omega_0 t}. \quad (12.12.7)$$

Then, writing $G(j\omega_0) = \operatorname{Re} M e^{j\theta}$, it follows that

$$y_{\text{hss}}(t) = a_0 M \sin(\omega_0 t + \phi_0 + \theta). \quad (12.12.8)$$

12.13 System Interconnections

Let $G \in \mathbb{R}_{\text{prop}}^{l \times m}(s)$. We define the *parahermitian conjugate* $G^\sim$ of G by

$$G^\sim \triangleq G^T(-s). \quad (12.13.1)$$

The following result provides realizations for G^T , $G^\sim$, and G^{-1} .

Proposition 12.13.1. Let $G_{\text{prop}}^{l \times m}(s)$, and assume that $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. Then,

$$G^T \sim \left[\begin{array}{c|c} A^T & C^T \\ \hline B^T & D^T \end{array} \right] \quad (12.13.2)$$

and

$$G^\sim \sim \left[\begin{array}{c|c} -A^T & -C^T \\ \hline B^T & D^T \end{array} \right]. \quad (12.13.3)$$

Furthermore, if G is square and D is nonsingular, then

$$G^{-1} \sim \left[\begin{array}{c|c} A - BD^{-1}C & BD^{-1} \\ \hline -D^{-1}C & D^{-1} \end{array} \right]. \quad (12.13.4)$$

Proof. Since $y = Gu$, it follows that G^{-1} satisfies $u = G^{-1}y$. Since $\dot{x} = Ax + Bu$ and $y = Cx + Du$, it follows that $u = -D^{-1}Cx + D^{-1}y$, and thus $\dot{x} = Ax + B(-D^{-1}Cx + D^{-1}y) = (A - BD^{-1}C)x + BD^{-1}y$. $\square$

Note that, if $G \in \mathbb{R}_{\text{prop}}(s)$ and $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$, then $G \sim \left[\begin{array}{c|c} A^T & B^T \\ \hline C^T & D \end{array} \right]$.

Let $G_1 \in \mathbb{R}_{\text{prop}}^{l_1 \times m_1}(s)$ and $G_2 \in \mathbb{R}_{\text{prop}}^{l_2 \times m_2}(s)$. If $m_2 = l_1$, then the *cascade interconnection* of G_1 and G_2 shown in Figure 12.13.1 is the product G_2G_1 , while the *parallel interconnection* shown in Figure 12.13.2 is the sum $G_1 + G_2$. Note that G_2G_1 is defined only if $m_2 = l_1$, whereas $G_1 + G_2$ requires that $m_1 = m_2$ and $l_1 = l_2$.




Figure 12.13.1
Cascade Interconnection of Linear Systems

Proposition 12.13.2. Let $G_1 \sim \left[\begin{array}{c|c} A_1 & B_1 \\ \hline C_1 & D_1 \end{array} \right]$ and $G_2 \sim \left[\begin{array}{c|c} A_2 & B_2 \\ \hline C_2 & D_2 \end{array} \right]$. Then,

$$G_2G_1 \sim \left[\begin{array}{cc|c} A_1 & 0 & B_1 \\ B_2C_1 & A_2 & B_2D_1 \\ \hline D_2C_1 & C_2 & D_2D_1 \end{array} \right] \quad (12.13.5)$$




Figure 12.13.2
Parallel Interconnection of Linear Systems

and

$$G_1 + G_2 \sim \left[\begin{array}{cc|c} A_1 & 0 & B_1 \\ 0 & A_2 & B_2 \\ \hline C_1 & C_2 & D_1 + D_2 \end{array} \right]. \quad (12.13.6)$$

Proof. Consider the state space equations

$$\begin{aligned} \dot{x}_1 &= A_1 x_1 + B_1 u_1, & \dot{x}_2 &= A_2 x_2 + B_2 u_2, \\ y_1 &= C_1 x_1 + D_1 u_1, & y_2 &= C_2 x_2 + D_2 u_2. \end{aligned}$$

Since $u_2 = y_1$, it follows that

$$\begin{aligned} \dot{x}_2 &= A_2 x_2 + B_2 C_1 x_1 + B_2 D_1 u_1, \\ y_2 &= C_2 x_2 + D_2 C_1 x_1 + D_2 D_1 u_1, \end{aligned}$$

and thus

$$\begin{aligned} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} &= \begin{bmatrix} A_1 & 0 \\ B_2 C_1 & A_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} B_1 \\ B_2 D_1 \end{bmatrix} u_1, \\ y_2 &= \begin{bmatrix} D_2 C_1 & C_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + D_2 D_1 u_1, \end{aligned}$$

which yields the realization (12.13.5) of $G_2 G_1$. The realization (12.13.6) for $G_1 + G_2$ can be obtained by similar techniques. $\square$

It is sometimes useful to combine transfer functions by concatenating them into row, column, or block-diagonal transfer functions.

Proposition 12.13.3. Let $G_1 \sim \left[\begin{array}{c|c} A_1 & B_1 \\ \hline C_1 & D_1 \end{array} \right]$ and $G_2 \sim \left[\begin{array}{c|c} A_2 & B_2 \\ \hline C_2 & D_2 \end{array} \right]$. Then,

$$\begin{bmatrix} G_1 & G_2 \end{bmatrix} \sim \left[\begin{array}{cc|cc} A_1 & 0 & B_1 & 0 \\ 0 & A_2 & 0 & B_2 \\ \hline C_1 & C_2 & D_1 & D_2 \end{array} \right], \quad (12.13.7)$$

$$\begin{bmatrix} G_1 \\ G_2 \end{bmatrix} \sim \left[\begin{array}{cc|c} A_1 & 0 & B_1 \\ 0 & A_2 & B_2 \\ \hline C_1 & 0 & D_1 \\ 0 & C_2 & D_2 \end{array} \right], \quad (12.13.8)$$

$$\begin{bmatrix} G_1 & 0 \\ 0 & G_2 \end{bmatrix} \sim \left[\begin{array}{cc|cc} A_1 & 0 & B_1 & 0 \\ 0 & A_2 & 0 & B_2 \\ \hline C_1 & 0 & D_1 & 0 \\ 0 & C_2 & 0 & D_2 \end{array} \right]. \quad (12.13.9)$$

Next, we interconnect a pair of systems G_1, G_2 by means of feedback as shown in Figure 12.13.3. It can be seen that u and y are related by

$$\hat{y} = (I + G_1 G_2)^{-1} G_1 \hat{u} \quad (12.13.10)$$

or

$$\hat{y} = G_1 (I + G_2 G_1)^{-1} \hat{u}. \quad (12.13.11)$$

The equivalence of (12.13.10) and (12.13.11) follows from the push-through identity given by Fact 2.16.16,

$$(I + G_1 G_2)^{-1} G_1 = G_1 (I + G_2 G_1)^{-1}. \quad (12.13.12)$$

A realization of this rational transfer function is given by the following result.




Figure 12.13.3
Feedback Interconnection of Linear Systems

Proposition 12.13.4. Let $G_1 \sim \left[\begin{array}{c|c} A_1 & B_1 \\ \hline C_1 & D_1 \end{array} \right]$ and $G_2 \sim \left[\begin{array}{c|c} A_2 & B_2 \\ \hline C_2 & D_2 \end{array} \right]$, and assume that $\det(I + D_1 D_2) \neq 0$. Then,

$$\begin{aligned} & [I + G_1 G_2]^{-1} G_1 \\ & \sim \left[\begin{array}{cc|c} A_1 - B_1(I + D_2 D_1)^{-1} D_2 C_1 & -B_1(I + D_2 D_1)^{-1} C_2 & B_1(I + D_2 D_1)^{-1} \\ B_2(I + D_1 D_2)^{-1} C_1 & A_2 - B_2(I + D_1 D_2)^{-1} D_1 C_2 & B_2(I + D_1 D_2)^{-1} D_1 \\ \hline (I + D_1 D_2)^{-1} C_1 & -(I + D_1 D_2)^{-1} D_1 C_2 & (I + D_1 D_2)^{-1} D_1 \end{array} \right]. \end{aligned} \quad (12.13.13)$$

12.14 Standard Control Problem

The standard control problem shown in Figure 12.14.1 involves four distinct signals, namely, an *exogenous input* w , a *control input* u , a *performance variable* z , and a *feedback signal* y . This system can be written as

$$\begin{bmatrix} \hat{z}(s) \\ \hat{y}(s) \end{bmatrix} = \tilde{g}(s) \begin{bmatrix} \hat{w}(s) \\ \hat{u}(s) \end{bmatrix}, \quad (12.14.1)$$

where $\mathcal{G}(s)$ is partitioned as

$$\mathcal{G} \triangleq \begin{bmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{bmatrix} \quad (12.14.2)$$

with the realization

$$\mathcal{G} \sim \left[\begin{array}{c|cc} A & D_1 & B \\ \hline E_1 & E_0 & E_2 \\ C & D_2 & D \end{array} \right], \quad (12.14.3)$$

which represents the equations

$$\dot{x} = Ax + D_1w + Bu, \quad (12.14.4)$$

$$z = E_1x + E_0w + E_2u, \quad (12.14.5)$$

$$y = Cx + D_2w + Du. \quad (12.14.6)$$

Consequently,

$$\mathcal{G}(s) = \begin{bmatrix} E_1(sI - A)^{-1}D_1 + E_0 & E_1(sI - A)^{-1}B + E_2 \\ C(sI - A)^{-1}D_1 + D_2 & C(sI - A)^{-1}B + D \end{bmatrix}, \quad (12.14.7)$$

which shows that G_{11} , G_{12} , G_{21} , and G_{22} have the realizations

$$G_{11} \sim \left[\begin{array}{c|c} A & D_1 \\ \hline E_1 & E_0 \end{array} \right], \quad G_{12} \sim \left[\begin{array}{c|c} A & B \\ \hline E_1 & E_2 \end{array} \right], \quad (12.14.8)$$

$$G_{21} \sim \left[\begin{array}{c|c} A & D_1 \\ \hline C & D_2 \end{array} \right], \quad G_{22} \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]. \quad (12.14.9)$$




Figure 12.14.1
Standard Control Problem

Letting G_c denote a feedback controller with realization

$$G_c \sim \left[\begin{array}{c|c} A_c & B_c \\ \hline C_c & D_c \end{array} \right], \quad (12.14.10)$$

we interconnect G and G_c according to

$$\hat{u}(s) = G_c(s)\hat{y}(s). \quad (12.14.11)$$

The resulting rational transfer function $\tilde{\mathcal{G}}$ satisfying $\hat{z}(s) = \tilde{\mathcal{G}}(s)\hat{w}(s)$ is thus given by

$$\tilde{\mathcal{G}} = G_{11} + G_{12}G_c(I - G_{22}G_c)^{-1}G_{21} \quad (12.14.12)$$

or

$$\tilde{\mathcal{G}} = G_{11} + G_{12}(I - G_cG_{22})^{-1}G_cG_{21}. \quad (12.14.13)$$

A realization of $\tilde{\mathcal{G}}$ is given by the following result.

Proposition 12.14.1. Let $\mathcal{G}$ and G_c have the realizations (12.14.3) and (12.14.10), and assume that $\det(I - DD_c) \neq 0$. Then,

$$\tilde{\mathcal{G}} \sim \left[\begin{array}{cc|c} A + BD_c(I - DD_c)^{-1}C & B(I - D_cD)^{-1}C_c & D_1 + BD_c(I + DD_c)^{-1}D_2 \\ B_c(I - DD_c)^{-1}C & A_c + B_c(I - DD_c)^{-1}DC_c & B_c(I - DD_c)^{-1}D_2 \\ \hline E_1 + E_2D_c(I - DD_c)^{-1}C & E_2(I - D_cD)^{-1}C_c & E_0 + E_2D_c(I - DD_c)^{-1}D_2 \end{array} \right]. \quad (12.14.14)$$

The realization (12.14.14) can be simplified when $DD_c = 0$. For example, if $D = 0$, then

$$\tilde{\mathcal{G}} \sim \left[\begin{array}{cc|c} A + BD_cC & BC_c & D_1 + BD_cD_2 \\ B_cC & A_c & B_cD_2 \\ \hline E_1 + E_2D_cC & E_2C_c & E_0 + E_2D_cD_2 \end{array} \right], \quad (12.14.15)$$

whereas, if $D_c = 0$, then

$$\tilde{\mathcal{G}} \sim \left[\begin{array}{cc|c} A & BC_c & D_1 \\ B_cC & A_c + B_cDC_c & B_cD_2 \\ \hline E_1 & E_2C_c & E_0 \end{array} \right]. \quad (12.14.16)$$

Finally, if both $D = 0$ and $D_c = 0$, then

$$\tilde{\mathcal{G}} \sim \left[\begin{array}{cc|c} A & BC_c & D_1 \\ B_cC & A_c & B_cD_2 \\ \hline E_1 & E_2C_c & E_0 \end{array} \right]. \quad (12.14.17)$$

The feedback interconnection shown in Figure 12.14.1 forms the basis for the *standard control problem* in feedback control. For this problem the signal w is an exogenous signal representing a command or a disturbance, while the signal z is the *performance variable*, that is, the variable whose behavior reflects the performance of the closed-loop system. The performance variable may or may not be physically measured. The *controlled input* or the *control* u is the output of the feedback controller G_c , while the *measurement* signal y serves as the input to the *feedback controller* G_c . The standard control problem is the following: Given knowledge of w , determine G_c to minimize a performance criterion $J(G_c)$.

12.15 Linear-Quadratic Control

Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$, and consider the system

$$\dot{x}(t) = Ax(t) + Bu(t), \quad (12.15.1)$$

$$x(0) = x_0, \quad (12.15.2)$$

where $t \geq 0$. Furthermore, let $K \in \mathbb{R}^{m \times n}$, and consider the full-state-feedback control law

$$u(t) = Kx(t). \quad (12.15.3)$$

The objective of the *linear-quadratic control problem* is to minimize the *quadratic performance measure*

$$J(K, x_0) = \int_0^\infty [x^T(t)R_1x(t) + 2x^T(t)R_{12}u(t) + u^T(t)R_2u(t)] dt, \quad (12.15.4)$$

where $R_1 \in \mathbb{R}^{n \times n}$, $R_{12} \in \mathbb{R}^{n \times m}$, and $R_2 \in \mathbb{R}^{m \times m}$. We assume that $\begin{bmatrix} R_1 & R_{12} \\ R_{12}^T & R_2 \end{bmatrix}$ is positive semidefinite and R_2 is positive definite.

The performance measure (12.15.4) indicates the desire to maintain the state vector $x(t)$ close to the zero equilibrium without an excessive expenditure of control effort. Specifically, the term $x^T(t)R_1x(t)$ is a measure of the deviation of the state $x(t)$ from the zero state, where the $n \times n$ positive-semidefinite matrix R_1 determines how much weighting is associated with each component of the state. Likewise, the $m \times m$ positive-definite matrix R_2 weights the magnitude of the control input. Finally, the cross-weighting term R_{12} arises naturally when additional filters are used to shape the system response or in specialized applications.

Using (12.15.1) and (12.15.3), the closed-loop dynamic system can be written as

$$\dot{x}(t) = (A + BK)x(t) \quad (12.15.5)$$

so that

$$x(t) = e^{t\tilde{A}}x_0, \quad (12.15.6)$$

where $\tilde{A} \triangleq A + BK$. Thus, the performance measure (12.15.4) becomes

$$\begin{aligned} J(K, x_0) &= \int_0^\infty x^T(t)\tilde{R}x(t) dt = \int_0^\infty x_0^T e^{t\tilde{A}^T} \tilde{R} e^{t\tilde{A}} x_0 dt \\ &= \text{tr } x_0^T \int_0^\infty e^{t\tilde{A}^T} \tilde{R} e^{t\tilde{A}} dt x_0 = \text{tr } \int_0^\infty e^{t\tilde{A}^T} \tilde{R} e^{t\tilde{A}} dt x_0 x_0^T, \end{aligned} \quad (12.15.7)$$

where

$$\tilde{R} \triangleq R_1 + R_{12}K + K^T R_{12}^T + K^T R_2 K. \quad (12.15.8)$$

Now, consider the standard control problem with plant

$$\mathcal{G} \sim \left[\begin{array}{c|cc} A & D_1 & B \\ \hline E_1 & 0 & E_2 \\ I_n & 0 & 0 \end{array} \right] \quad (12.15.9)$$

and full-state feedback $u = Kx$. Then, the closed-loop transfer function is given by

$$\tilde{\mathcal{G}} \sim \left[\begin{array}{c|c} A + BK & D_1 \\ \hline E_1 + E_2K & 0 \end{array} \right]. \quad (12.15.10)$$

The following result shows that the quadratic performance measure (12.15.4) is equal to the H_2 norm of a transfer function.

Proposition 12.15.1. Assume that $D_1 = x_0$ and

$$\begin{bmatrix} R_1 & R_{12} \\ R_{12}^T & R_2 \end{bmatrix} = \begin{bmatrix} E_1^T \\ E_2^T \end{bmatrix} \begin{bmatrix} E_1 & E_2 \end{bmatrix}, \quad (12.15.11)$$

and let $\tilde{\mathcal{G}}$ be given by (12.15.10). Then,

$$J(K, x_0) = \|\tilde{\mathcal{G}}\|_{H_2}^2. \quad (12.15.12)$$

Proof. The result follows from Proposition 12.1.2. $\square$

For the following development, we assume that (12.15.11) holds so that R_1 , R_{12} , and R_2 are given by

$$R_1 = E_1^T E_1, \quad R_{12} = E_1^T E_2, \quad R_2 = E_2^T E_2. \quad (12.15.13)$$

To develop necessary conditions for the linear-quadratic control problem, we restrict K to the set of stabilizing gains

$$\mathcal{S} \triangleq \{K \in \mathbb{R}^{m \times n}: A + BK \text{ is asymptotically stable}\}. \quad (12.15.14)$$

Obviously, $\mathcal{S}$ is nonempty if and only if (A, B) is stabilizable. The following result gives necessary conditions that characterize a stabilizing solution K of the linear-quadratic control problem.

Theorem 12.15.2. Assume that (A, B) is stabilizable, assume that $K \in \mathcal{S}$ solves the linear-quadratic control problem, and assume that $(A + BK, D_1)$ is controllable. Then, there exists an $n \times n$ positive-semidefinite matrix P such that K is given by

$$K = -R_2^{-1}(B^T P + R_{12}^T) \quad (12.15.15)$$

and such that P satisfies

$$\hat{A}_R^T P + P \hat{A}_R + \hat{R}_1 - P B R_2^{-1} B^T P = 0, \quad (12.15.16)$$

where

$$\hat{A}_R \triangleq A - B R_2^{-1} R_{12}^T \quad (12.15.17)$$

and

$$\hat{R}_1 \triangleq R_1 - R_{12}R_2^{-1}R_{12}^T. \quad (12.15.18)$$

Furthermore, the minimal cost is given by

$$J(K) = \text{tr } PV, \quad (12.15.19)$$

where $V \triangleq D_1D_1^T$.

Proof. Since $K \in \mathcal{S}$, it follows that $\tilde{A}$ is asymptotically stable. It then follows that $J(K)$ is given by (12.15.19), where $P \triangleq \int_0^\infty e^{t\tilde{A}^T}\tilde{R}e^{t\tilde{A}}dt$ is positive semidefinite and satisfies the Lyapunov equation

$$\tilde{A}^TP + P\tilde{A} + \tilde{R} = 0. \quad (12.15.20)$$

Note that (12.15.20) can be written as

$$(A + BK)^TP + P(A + BK) + R_1 + R_{12}K + K^TR_{12}^T + K^TR_2K = 0. \quad (12.15.21)$$

To optimize (12.15.19) subject to the constraint (12.15.20) over the open set $\mathcal{S}$, form the Lagrangian

$$\mathcal{L}(K, P, Q, \lambda_0) \triangleq \text{tr} \left[\lambda_0 PV + Q \left(\tilde{A}^TP + P\tilde{A} + \tilde{R} \right) \right], \quad (12.15.22)$$

where the Lagrange multipliers $\lambda_0 \geq 0$ and $Q \in \mathbb{R}^{n \times n}$ are not both zero. Note that the $n \times n$ Lagrange multiplier Q accounts for the $n \times n$ constraint equation (12.15.20).

The necessary condition $\partial\mathcal{L}/\partial P = 0$ implies

$$\tilde{A}Q + Q\tilde{A}^T + \lambda_0 V = 0. \quad (12.15.23)$$

Since $\tilde{A}$ is asymptotically stable, it follows from Proposition 11.9.3 that, for all $\lambda_0 \geq 0$, (12.15.23) has a unique solution Q and, furthermore, Q is positive semidefinite. In particular, if $\lambda_0 = 0$, then $Q = 0$. Since λ_0 and Q are not both zero, we can set $\lambda_0 = 1$ so that (12.15.23) becomes

$$\tilde{A}Q + Q\tilde{A}^T + V = 0. \quad (12.15.24)$$

Since $(\tilde{A}, D_1)$ is controllable, it follows from Corollary 12.7.10 that Q is positive definite.

Next, evaluating $\partial\mathcal{L}/\partial K = 0$ yields

$$R_2KQ + (B^TP + R_{12}^T)Q = 0. \quad (12.15.25)$$

Since Q is positive definite, it follows from (12.15.25) that (12.15.15) is satisfied. Furthermore, using (12.15.15), it follows that (12.15.20) is equivalent to (12.15.16). $\square$

With K given by (12.15.15) the closed-loop dynamics matrix $\tilde{A} = A + BK$ is given by

$$\tilde{A} = A - BR_2^{-1}(B^TP + R_{12}^T), \quad (12.15.26)$$

where P is the solution of the *Riccati equation* (12.15.16).

12.16 Solutions of the Riccati Equation

For convenience in the following development, we assume that $R_{12} = 0$. With this assumption, the gain K given by (12.15.15) becomes

$$K = -R_2^{-1}B^TP. \quad (12.16.1)$$

Defining

$$\Sigma \triangleq BR_2^{-1}B^T, \quad (12.16.2)$$

(12.15.26) becomes

$$\tilde{A} = A - \Sigma P, \quad (12.16.3)$$

while the Riccati equation (12.15.16) can be written as

$$A^TP + PA + R_1 - P\Sigma P = 0. \quad (12.16.4)$$

Note that (12.16.4) has the alternative representation

$$(A - \Sigma P)^TP + P(A - \Sigma P) + R_1 + P\Sigma P = 0, \quad (12.16.5)$$

which is equivalent to the Lyapunov equation

$$\tilde{A}^TP + P\tilde{A} + \tilde{R} = 0, \quad (12.16.6)$$

where

$$\tilde{R} \triangleq R_1 + P\Sigma P. \quad (12.16.7)$$

By comparing (12.15.16) and (12.16.4), it can be seen that the linear-quadratic control problems with (A, B, R_1, R_{12}, R_2) and $(\hat{A}_R, B, \hat{R}_1, 0, R_2)$ are equivalent. Hence, there is no loss of generality in assuming that $R_{12} = 0$ in the following development, where A and R_1 take the place of $\hat{A}_R$ and $\hat{R}_1$, respectively.

To motivate the subsequent development, the following examples demonstrate the existence of solutions under various assumptions on (A, B, E_1) . In the following four examples, (A, B) is not stabilizable.

Example 12.16.1. Let $n = 1$, $A = 1$, $B = 0$, $E_1 = 0$, and $R_2 > 0$. Hence, (A, B, E_1) has an ORHP eigenvalue that is uncontrollable and unobservable. In this case, (12.16.4) has the unique solution $P = 0$. Furthermore, since $B = 0$, it follows that $\tilde{A} = A$.

Example 12.16.2. Let $n = 1$, $A = 1$, $B = 0$, $E_1 = 1$, and $R_2 > 0$. Hence, (A, B, E_1) has an ORHP eigenvalue that is uncontrollable and observable. In this case, (12.16.4) has the unique solution $P = -1/2 < 0$. Furthermore, since $B = 0$, it follows that $\tilde{A} = A$.

Example 12.16.3. Let $n = 1$, $A = 0$, $B = 0$, $E_1 = 0$, and $R_2 > 0$. Hence, (A, B, E_1) has an imaginary eigenvalue that is uncontrollable and unobservable. In this case, (12.16.4) has infinitely many solutions $P \in \mathbb{R}$. Hence, (12.16.4) has no maximal solution. Furthermore, since $B = 0$, it follows that, for every solution P , $\tilde{A} = A$.

Example 12.16.4. Let $n = 1$, $A = 0$, $B = 0$, $E_1 = 1$, and $R_2 > 0$. Hence, (A, B, E_1) has an imaginary eigenvalue that is uncontrollable and observable. In this case, (12.16.4) becomes $R_1 = 0$. Thus, (12.16.4) has no solution.

In the remaining examples, (A, B) is controllable.

Example 12.16.5. Let $n = 1$, $A = 1$, $B = 1$, $E_1 = 0$, and $R_2 > 0$. Hence, (A, B, E_1) has an ORHP eigenvalue that is controllable and unobservable. In this case, (12.16.4) has the solutions $P = 0$ and $P = 2R_2 > 0$. The corresponding closed-loop dynamics matrices are $\tilde{A} = 1 > 0$ and $\tilde{A} = -1 < 0$. Hence, the solution $P = 2R_2$ is stabilizing, and the closed-loop eigenvalue 1, which does not depend on R_2 , is the reflection of the open-loop eigenvalue -1 across the imaginary axis.

Example 12.16.6. Let $n = 1$, $A = 1$, $B = 1$, $E_1 = 1$, and $R_2 > 0$. Hence, (A, B, E_1) has an ORHP eigenvalue that is controllable and observable. In this case, (12.16.4) has the solutions $P = R_2 - \sqrt{R_2^2 + R_2} < 0$ and $P = R_2 + \sqrt{R_2^2 + R_2} > 0$. The corresponding closed-loop dynamics matrices are $\tilde{A} = \sqrt{1 + 1/R_2} > 0$ and $\tilde{A} = -\sqrt{1 + 1/R_2} < 0$. Hence, the positive-definite solution $P = R_2 + \sqrt{R_2^2 + R_2}$ is stabilizing.

Example 12.16.7. Let $n = 1$, $A = 0$, $B = 1$, $E_1 = 0$, and $R_2 > 0$. Hence, (A, B, E_1) has an imaginary eigenvalue that is controllable and unobservable. In this case, (12.16.4) has the unique solution $P = 0$, which is not stabilizing.

Example 12.16.8. Let $n = 1$, $A = 0$, $B = 1$, $E_1 = 1$, and $R_2 > 0$. Hence, (A, B, E_1) has an imaginary eigenvalue that is controllable and observable. In this case, (12.16.4) has the solutions $P = -\sqrt{R_2} < 0$ and $P = \sqrt{R_2} > 0$. The corresponding closed-loop dynamics matrices are $\tilde{A} = \sqrt{R_2} > 0$ and $\tilde{A} = -\sqrt{R_2} < 0$. Hence, the positive-definite solution $P = \sqrt{R_2}$ is stabilizing.

Example 12.16.9. Let $n = 2$, $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$, $B = I_2$, $E_1 = 0$, and $R_2 = 1$. Hence, as in Example 12.16.7, both eigenvalues of (A, B, E_1) are imaginary, controllable, and unobservable. Taking the trace of (12.16.4) yields $\text{tr } P^2 = 0$. Thus, the only symmetric matrix P satisfying (12.16.4) is $P = 0$, which implies that $\tilde{A} = A$. Consequently, the open-loop eigenvalues $\pm j$ are unmoved by the feedback gain (12.15.15) even though (A, B) is controllable.

Example 12.16.10. Let $n = 2$, $A = 0$, $B = I_2$, $E_1 = I_2$, and $R_2 = I$. Hence, as in Example 12.16.8, both eigenvalues of (A, B, E_1) are imaginary, controllable, and observable. Furthermore, (12.16.4) becomes $P^2 = I$. Requiring that P be symmetric, it follows that P is a reflector. Hence, $P = I$ is the only positive-semidefinite solution. In fact, P is positive definite and stabilizing since $\tilde{A} = -I$.

Example 12.16.11. Let $A = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$, $B = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, $E_1 = 0$, and $R_2 = 1$ so that (A, B) is controllable, although neither of the states is weighted. In this case, (12.16.4) has four positive-semidefinite solutions, which are given by

$$P_1 = \begin{bmatrix} 18 & -24 \\ -24 & 36 \end{bmatrix}, \quad P_2 = \begin{bmatrix} 2 & 0 \\ 0 & 0 \end{bmatrix}, \quad P_3 = \begin{bmatrix} 0 & 0 \\ 0 & 4 \end{bmatrix}, \quad P_4 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

The corresponding feedback matrices are given by $K_1 = \begin{bmatrix} 6 & -12 \end{bmatrix}$, $K_2 = \begin{bmatrix} -2 & 0 \end{bmatrix}$, $K_3 = \begin{bmatrix} 0 & -4 \end{bmatrix}$, and $K_4 = \begin{bmatrix} 0 & 0 \end{bmatrix}$. Letting $\tilde{A}_i = A - \Sigma P_i$, it follows that $\text{spec}(\tilde{A}_1) = \{-1, -2\}$, $\text{spec}(\tilde{A}_2) = \{-1, 2\}$, $\text{spec}(\tilde{A}_3) = \{1, -2\}$, and $\text{spec}(\tilde{A}_4) = \{1, 2\}$. Thus, P_1 is the only solution that stabilizes the closed-loop system, while the solutions P_2 and P_3 partially stabilize the closed-loop system. Note also that the closed-loop poles that differ from those of the open-loop system are mirror images of the open-loop poles as reflected across the imaginary axis. Finally, note that these solutions satisfy the partial ordering $P_1 \geq P_2 \geq P_4$ and $P_1 \geq P_3 \geq P_4$, and that “larger” solutions are more stabilizing than “smaller” solutions. Moreover, letting $J(K_i) = \text{tr } P_i V$, it can be seen that larger solutions incur a greater closed-loop cost, with the greatest cost incurred by the stabilizing solution P_4 . However, the cost expression $J(K) = \text{tr } P V$ does not follow from (12.15.4) when $A + BK$ is not asymptotically stable.

The following definition concerns solutions of the Riccati equation.

Definition 12.16.12. A matrix $P \in \mathbb{R}^{n \times n}$ is a *solution* of the Riccati equation (12.16.4) if P is symmetric and satisfies (12.16.4). Furthermore, P is the *stabilizing solution* of (12.16.4) if $\tilde{A} = A - \Sigma P$ is asymptotically stable. Finally, a solution $P_{\max}$ of (12.16.4) is the *maximal solution* to (12.16.4) if $P \leq P_{\max}$ for every solution P to (12.16.4).

Since the ordering “ $\leq$ ” is antisymmetric, it follows that (12.16.4) has at most one maximal solution. The uniqueness of the stabilizing solution is shown in the following section.

Next, define the $2n \times 2n$ *Hamiltonian*

$$\mathcal{H} \triangleq \begin{bmatrix} A & \Sigma \\ R_1 & -A^T \end{bmatrix}. \quad (12.16.8)$$

Proposition 12.16.13. The following statements hold:

- i) $\mathcal{H}$ is Hamiltonian.
- ii) $\chi_{\mathcal{H}}$ has a spectral factorization, that is, there exists a monic polynomial $p \in \mathbb{R}[s]$ such that, for all $s \in \mathbb{C}$, $\chi_{\mathcal{H}}(s) = p(s)p(-s)$.
- iii) $\chi_{\mathcal{H}}(j\omega) \geq 0$ for all $\omega \in \mathbb{R}$.
- iv) If either $R_1 = 0$ or $\Sigma = 0$, then $\text{mspec}(\mathcal{H}) = \text{mspec}(A) \cup \text{mspec}(-A)$.
- v) $\chi_{\mathcal{H}}$ is even.
- vi) $\lambda \in \text{spec}(\mathcal{H})$ if and only if $-\lambda \in \text{spec}(\mathcal{H})$.
- vii) If $\lambda \in \text{spec}(\mathcal{H})$, then $\text{amult}_{\mathcal{H}}(\lambda) = \text{amult}_{\mathcal{H}}(-\lambda)$.
- viii) Every imaginary root of $\chi_{\mathcal{H}}$ has even multiplicity.
- ix) Every imaginary eigenvalue of $\mathcal{H}$ has even algebraic multiplicity.

Proof. The result follows from Proposition 4.1.1 and Fact 4.9.23. $\square$

It is helpful to keep in mind that spectral factorizations are not unique. For example, if $\chi_{\mathcal{H}}(s) = (s+1)(s+2)(-s+1)(-s+2)$, then $\chi_{\mathcal{H}}(s) = p(s)p(-s) = \hat{p}(s)\hat{p}(-s)$, where $p(s) = (s+1)(s+2)$ and $\hat{p}(s) = (s+1)(s-2)$. Thus, the spectral factors $p(s)$ and $p(-s)$ can “trade” roots. These roots are the eigenvalues of $\mathcal{H}$.

The following result shows that the Hamiltonian matrix $\mathcal{H}$ is closely linked to the Riccati equation (12.16.4).

Proposition 12.16.14. Let $P \in \mathbb{R}^{n \times n}$ be symmetric. Then, the following statements are equivalent:

i) P is a solution of (12.16.4).

ii) P satisfies

$$\begin{bmatrix} P & I \end{bmatrix} \mathcal{H} \begin{bmatrix} I \\ -P \end{bmatrix} = 0. \quad (12.16.9)$$

iii) P satisfies

$$\mathcal{H} \begin{bmatrix} I \\ -P \end{bmatrix} = \begin{bmatrix} I \\ -P \end{bmatrix} (A - \Sigma P). \quad (12.16.10)$$

iv) P satisfies

$$\mathcal{H} = \begin{bmatrix} I & 0 \\ -P & I \end{bmatrix} \begin{bmatrix} A - \Sigma P & \Sigma \\ 0 & -(A - \Sigma P)^T \end{bmatrix} \begin{bmatrix} I & 0 \\ P & I \end{bmatrix}. \quad (12.16.11)$$

In this case, the following statements hold:

v) $\text{mspec}(\mathcal{H}) = \text{mspec}(A - \Sigma P) \cup \text{mspec}[-(A - \Sigma P)]$.

vi) $\chi_{\mathcal{H}}(s) = (-1)^n \chi_{A - \Sigma P}(s) \chi_{A - \Sigma P}(-s)$.

vii) $\mathcal{R}(\begin{bmatrix} I \\ -P \end{bmatrix})$ is an invariant subspace of $\mathcal{H}$.

Corollary 12.16.15. Assume that (12.16.4) has a stabilizing solution. Then, $\mathcal{H}$ has no imaginary eigenvalues.

For the next two results, P is not necessarily a solution of (12.16.4).

Lemma 12.16.16. Assume that $\lambda \in \text{spec}(A)$ is an observable eigenvalue of (A, R_1) , and let $P \in \mathbb{R}^{n \times n}$ be symmetric. Then, $\lambda \in \text{spec}(A)$ is an observable eigenvalue of $(\tilde{A}, \tilde{R})$.

Proof. Suppose that $\text{rank} \begin{bmatrix} \lambda I - \tilde{A} \\ \tilde{R} \end{bmatrix} < n$. Then, there exists a nonzero vector $v \in \mathbb{C}^n$ such that $\tilde{A}v = \lambda v$ and $\tilde{R}v = 0$. Hence, $v^* R_1 v = -v^* P \Sigma P v \leq 0$, which implies that $R_1 v = 0$ and $P \Sigma P v = 0$. Hence, $\Sigma P v = 0$, and thus $Av = \lambda v$. Therefore, $\text{rank} \begin{bmatrix} \lambda I - A \\ R_1 \end{bmatrix} < n$. $\square$

Lemma 12.16.17. Assume that (A, R_1) is (observable, detectable), and let $P \in \mathbb{R}^{n \times n}$ be symmetric. Then, $(\tilde{A}, \tilde{R})$ is (observable, detectable).

Lemma 12.16.18. Assume that (A, E_1) is observable, and assume that (12.16.4) has a solution P . Then, the following statements hold:

- i) $\nu_-(\tilde{A}) = \nu_+(P)$.
- ii) $\nu_0(\tilde{A}) = \nu_0(P) = 0$.
- iii) $\nu_+(\tilde{A}) = \nu_-(P)$.

Proof. Since (A, R_1) is observable, it follows from Lemma 12.16.17 that $(\tilde{A}, \tilde{R})$ is observable. By writing (12.16.4) as the Lyapunov equation (12.16.6), the result now follows from Fact 12.21.1. $\square$

12.17 The Stabilizing Solution of the Riccati Equation

Proposition 12.17.1. The following statements hold:

- i) (12.16.4) has at most one stabilizing solution.
- ii) If P is the stabilizing solution of (12.16.4), then P is positive semidefinite.
- iii) If P is the stabilizing solution of (12.16.4), then

$$\text{rank } P = \text{rank } \mathcal{O}(\tilde{A}, \tilde{R}). \quad (12.17.1)$$

Proof. To prove i), suppose that (12.16.4) has stabilizing solutions P_1 and P_2 . Then,

$$\begin{aligned} A^T P_1 + P_1 A + R_1 - P_1 \Sigma P_1 &= 0, \\ A^T P_2 + P_2 A + R_1 - P_2 \Sigma P_2 &= 0. \end{aligned}$$

Subtracting these equations and rearranging yields

$$(A - \Sigma P_1)^T (P_1 - P_2) + (P_1 - P_2)(A - \Sigma P_2) = 0.$$

Since $A - \Sigma P_1$ and $A - \Sigma P_2$ are asymptotically stable, it follows from Proposition 11.9.3 and Fact 11.18.33 that $P_1 - P_2 = 0$. Hence, (12.16.4) has at most one stabilizing solution.

Next, to prove ii), suppose that P is a stabilizing solution of (12.16.4). Then, it follows from (12.16.4) that

$$P = \int_0^\infty e^{t(A - \Sigma P)^T} (R_1 + P \Sigma P) e^{t(A - \Sigma P)} dt,$$

which shows that P is positive semidefinite.

Finally, iii) follows from Corollary 12.3.3. $\square$

Theorem 12.17.2. Assume that (12.16.4) has a positive-semidefinite solution P , and assume that (A, E_1) is detectable. Then, P is the stabilizing solution of (12.16.4), and thus P is the only positive-semidefinite solution of (12.16.4). If, in addition, (A, E_1) is observable, then P is positive definite.

Proof. Since (A, R_1) is detectable, it follows from Lemma 12.16.17 that $(\tilde{A}, \tilde{R})$ is detectable. Next, since (12.16.4) has a positive-semidefinite solution P , it follows

from Corollary 12.8.6 that $\tilde{A}$ is asymptotically stable. Hence, P is the stabilizing solution of (12.16.4). The last statement follows from Lemma 12.16.18. $\square$

Corollary 12.17.3. Assume that (A, E_1) is detectable. Then, (12.16.4) has at most one positive-semidefinite solution.

Lemma 12.17.4. Let $\lambda \in \mathbb{C}$, and assume that λ is either an uncontrollable eigenvalue of (A, B) or an unobservable eigenvalue of (A, E_1) . Then, $\lambda \in \text{spec}(\mathcal{H})$.

Proof. Note that

$$\lambda I - \mathcal{H} = \begin{bmatrix} \lambda I - A & -\Sigma \\ -R_1 & \lambda I + A^T \end{bmatrix}.$$

If λ is an uncontrollable eigenvalue of (A, B) , then the first n rows of $\lambda I - \mathcal{H}$ are linearly dependent, and thus $\lambda \in \text{spec}(\mathcal{H})$. On the other hand, if λ is an unobservable eigenvalue of (A, E_1) , then the first n columns of $\lambda I - \mathcal{H}$ are linearly dependent, and thus $\lambda \in \text{spec}(\mathcal{H})$. $\square$

The following result is a consequence of Lemma 12.17.4.

Proposition 12.17.5. Let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that

$$A = S \begin{bmatrix} A_1 & 0 & A_{13} & 0 \\ A_{21} & A_2 & A_{23} & A_{24} \\ 0 & 0 & A_3 & 0 \\ 0 & 0 & A_{43} & A_4 \end{bmatrix} S^{-1}, \quad B = S \begin{bmatrix} B_1 \\ B_2 \\ 0 \\ 0 \end{bmatrix}, \quad (12.17.2)$$

$$E_1 = \begin{bmatrix} E_{11} & 0 & E_{13} & 0 \end{bmatrix} S^{-1}, \quad (12.17.3)$$

where $(\begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix}, \begin{bmatrix} B_1 \\ B_2 \end{bmatrix})$ is controllable and $(\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}, \begin{bmatrix} E_{11} & E_{13} \end{bmatrix})$ is observable. Then,

$$\text{mspec}(A_2) \cup \text{mspec}(-A_2) \subseteq \text{mspec}(\mathcal{H}), \quad (12.17.4)$$

$$\text{mspec}(A_3) \cup \text{mspec}(-A_3) \subseteq \text{mspec}(\mathcal{H}), \quad (12.17.5)$$

$$\text{mspec}(A_4) \cup \text{mspec}(-A_4) \subseteq \text{mspec}(\mathcal{H}). \quad (12.17.6)$$

Next, we present a partial converse of Lemma 12.17.4.

Lemma 12.17.6. Let $\lambda \in \text{spec}(\mathcal{H})$, and assume that $\text{Re } \lambda = 0$. Then, λ is either an uncontrollable eigenvalue of (A, B) or an unobservable eigenvalue of (A, E_1) .

Proof. Suppose that $\lambda = j\omega$ is an eigenvalue of $\mathcal{H}$, where $\omega \in \mathbb{R}$. Then, there exist $x, y \in \mathbb{C}^n$ such that $\begin{bmatrix} x \\ y \end{bmatrix} \neq 0$ and $\mathcal{H} \begin{bmatrix} x \\ y \end{bmatrix} = j\omega \begin{bmatrix} x \\ y \end{bmatrix}$. Consequently,

$$Ax + \Sigma y = j\omega x, \quad R_1 x - A^T y = j\omega y.$$

Rewriting these identities as

$$(A - j\omega I)x = -\Sigma y, \quad (A - j\omega I)^* y = R_1 x$$

yields

$$y^*(A - j\omega I)x = -y^*\Sigma y, \quad x^*(A - j\omega I)^*y = x^*R_1x.$$

Since $x^*(A - j\omega I)^*y$ is real, it follows that $-y^*\Sigma y = x^*R_1x$, and thus $y^*\Sigma y = x^*R_1x = 0$, which implies that $B^T y = 0$ and $E_1x = 0$. Therefore,

$$(A - j\omega I)x = 0, \quad (A - j\omega I)^*y = 0,$$

and hence

$$\begin{bmatrix} A - j\omega I \\ E_1 \end{bmatrix} x = 0, \quad y^* \begin{bmatrix} A - j\omega I & B \end{bmatrix} = 0.$$

Since $\begin{bmatrix} x \\ y \end{bmatrix} \neq 0$, it follows that either $x \neq 0$ or $y \neq 0$, and thus either $\text{rank} \begin{bmatrix} A - j\omega I \\ E_1 \end{bmatrix} < n$ or $\text{rank} \begin{bmatrix} A - j\omega I & B \end{bmatrix} < n$. $\square$

The following result is a restatement of Lemma 12.17.6.

Proposition 12.17.7. Let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that (12.17.2) and (12.17.3) are satisfied, where $(\begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix}, \begin{bmatrix} B_1 \\ B_2 \end{bmatrix})$ is controllable and $(\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}, \begin{bmatrix} E_{11} & E_{13} \end{bmatrix})$ is observable. Then,

$$\begin{aligned} \text{mspec}(\mathcal{H}) \cap j\mathbb{R} &\subseteq \text{mspec}(A_2) \cup \text{mspec}(-A_2) \cup \text{mspec}(A_3) \\ &\cup \text{mspec}(-A_3) \cup \text{mspec}(A_4) \cup \text{mspec}(-A_4). \end{aligned} \quad (12.17.7)$$

Combining Lemma 12.17.4 and Lemma 12.17.6 yields the following result.

Proposition 12.17.8. Let $\lambda \in \mathbb{C}$, assume that $\text{Re } \lambda = 0$, and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that (12.17.2) and (12.17.3) are satisfied, where (A_1, B_1, E_{11}) is controllable and observable, (A_2, B_2) is controllable, and (A_3, E_{13}) is observable. Then, the following statements are equivalent:

- i) λ is either an uncontrollable eigenvalue of (A, B) or an unobservable eigenvalue of (A, E_1) .
- ii) $\lambda \in \text{mspec}(A_2) \cup \text{mspec}(A_3) \cup \text{mspec}(A_4)$.
- iii) λ is an eigenvalue of $\mathcal{H}$.

The next result gives necessary and sufficient conditions under which (12.16.4) has a stabilizing solution. This result also provides a constructive characterization of the stabilizing solution. Result ii) of Proposition 12.10.11 shows that the condition in i) that every imaginary eigenvalue of (A, E_1) is observable is equivalent to the condition that $\left[\begin{array}{c|c} A & B \\ \hline E_1 & E_2 \end{array} \right]$ has no imaginary invariant zeros.

Theorem 12.17.9. The following statements are equivalent:

- i) (A, B) is stabilizable, and every imaginary eigenvalue of (A, E_1) is observable.
- ii) There exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.17.2) and (12.17.3) are satisfied, where $(\begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix}, \begin{bmatrix} B_1 \\ B_2 \end{bmatrix})$ is controllable, $(\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}, \begin{bmatrix} E_{11} & E_{13} \end{bmatrix})$ is observable, $\nu_0(A_2) = 0$, and A_3 and A_4 are asymp-

totically stable.

iii) (12.16.4) has a stabilizing solution.

In this case, let

$$M = \begin{bmatrix} M_1 & M_{12} \\ M_{21} & M_2 \end{bmatrix} \in \mathbb{R}^{2n \times 2n} \quad (12.17.8)$$

be a nonsingular matrix such that $\mathcal{H} = MZM^{-1}$, where

$$Z = \begin{bmatrix} Z_1 & Z_{12} \\ 0 & Z_2 \end{bmatrix} \in \mathbb{R}^{2n \times 2n} \quad (12.17.9)$$

and $Z_1 \in \mathbb{R}^{n \times n}$ is asymptotically stable. Then, M_1 is nonsingular, and

$$P \triangleq -M_{21}M_1^{-1} \quad (12.17.10)$$

is the stabilizing solution of (12.16.4).

Proof. The equivalence of i) and ii) is immediate.

To prove i) $\implies$ iii), first note that Lemma 12.17.6 implies that $\mathcal{H}$ has no imaginary eigenvalues. Hence, since $\mathcal{H}$ is Hamiltonian, it follows that there exists a matrix $M \in \mathbb{R}^{2n \times 2n}$ of the form (12.17.8) such that M is nonsingular and $\mathcal{H} = MZM^{-1}$, where $Z \in \mathbb{R}^{2n \times 2n}$ is of the form (12.17.9) and $Z_1 \in \mathbb{R}^{n \times n}$ is asymptotically stable.

Next, note that $\mathcal{H}M = MZ$ implies that

$$\mathcal{H} \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} = M \begin{bmatrix} Z_1 \\ 0 \end{bmatrix} = \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} Z_1.$$

Therefore,

$$\begin{aligned} \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix}^T J_n \mathcal{H} \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} &= \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix}^T J_n \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} Z_1 \\ &= \begin{bmatrix} M_1^T & M_{21}^T \end{bmatrix} \begin{bmatrix} M_{21} \\ -M_1 \end{bmatrix} Z_1 \\ &= LZ_1, \end{aligned}$$

where $L \triangleq M_1^T M_{21} - M_{21}^T M_1$. Since $J_n \mathcal{H} = (J_n \mathcal{H})^T$, it follows that LZ_1 is symmetric, that is, $LZ_1 = Z_1^T L^T$. Since, in addition, L is skew symmetric, it follows that $0 = Z_1^T L + LZ_1$. Now, since Z_1 is asymptotically stable, it follows that $L = 0$. Hence, $M_1^T M_{21} = M_{21}^T M_1$, which shows that $M_{21}^T M_1$ is symmetric.

To show that M_1 is nonsingular, note that it follows from the identity

$$\begin{bmatrix} I & 0 \end{bmatrix} \mathcal{H} \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} = \begin{bmatrix} I & 0 \end{bmatrix} \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} Z_1$$

that

$$AM_1 + \Sigma M_{21} = M_1 Z_1.$$

Now, let $x \in \mathbb{R}^n$ satisfy $M_1 x = 0$. We thus have

$$\begin{aligned}
x^T M_{21} \Sigma M_{21} x &= x^T M_{21}^T (A M_1 + \Sigma M_{21}) x \\
&= x^T M_{21}^T M_1 Z_1 x \\
&= x^T M_1^T M_{21} Z_1 x \\
&= 0,
\end{aligned}$$

which implies that $B^T M_{21} x = 0$. Hence, $M_1 Z_1 x = (A M_1 + \Sigma M_{21}) x = 0$. Thus, $Z_1 \mathcal{N}(M_1) \subseteq \mathcal{N}(M_1)$.

Now, suppose that M_1 is singular. Since $Z_1 \mathcal{N}(M_1) \subseteq \mathcal{N}(M_1)$, it follows that there exists $\lambda \in \text{spec}(Z_1)$ and $x \in \mathbb{C}^n$ such that $Z_1 x = \lambda x$ and $M_1 x = 0$. Forming

$$\begin{bmatrix} 0 & I \end{bmatrix} \mathcal{H} \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} x = \begin{bmatrix} 0 & I \end{bmatrix} \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} Z_1 x$$

yields $-A^T M_{21} x = M_{21} \lambda x$, and thus $(\lambda I + A^T) M_{21} x = 0$. Since, in addition, as shown above, $B^T M_{21} x = 0$, it follows that $x^* M_{21}^T \begin{bmatrix} -\bar{\lambda} I - A & B \end{bmatrix} = 0$. Since $\lambda \in \text{spec}(Z_1)$, it follows that $\text{Re}(-\bar{\lambda}) > 0$. Furthermore, since, by assumption, (A, B) is stabilizable, it follows that $\text{rank} \begin{bmatrix} \bar{\lambda} I - A & B \end{bmatrix} = n$. Therefore, $M_{21} x = 0$. Combining this fact with $M_1 x = 0$ yields $\begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} x = 0$. Since x is nonzero, it follows that M is singular, which is a contradiction. Consequently, M_1 is nonsingular. Next, define $P \triangleq -M_{21} M_1^{-1}$ and note that, since $M_1^T M_{21}$ is symmetric, it follows that $P = -M_1^{-T} (M_1^T M_{21}) M_1^{-1}$ is also symmetric.

Since $\mathcal{H} \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} = \begin{bmatrix} M_1 \\ M_{21} \end{bmatrix} Z_1$, it follows that

$$\mathcal{H} \begin{bmatrix} I \\ M_{21} M_1^{-1} \end{bmatrix} = \begin{bmatrix} I \\ M_{21} M_1^{-1} \end{bmatrix} M_1 Z_1 M_1^{-1},$$

and thus

$$\mathcal{H} \begin{bmatrix} I \\ -P \end{bmatrix} = \begin{bmatrix} I \\ -P \end{bmatrix} M_1 Z_1 M_1^{-1}.$$

Multiplying on the left by $\begin{bmatrix} P & I \end{bmatrix}$ yields

$$0 = \begin{bmatrix} P & I \end{bmatrix} \mathcal{H} \begin{bmatrix} I \\ -P \end{bmatrix} = A^T P + P A + R_1 - P \Sigma P,$$

which shows that P is a solution of (12.16.4). Similarly, multiplying on the left by $\begin{bmatrix} I & 0 \end{bmatrix}$ yields $A - \Sigma P = M_1 Z_1 M_1^{-1}$. Since Z_1 is asymptotically stable, it follows that $A - \Sigma P$ is also asymptotically stable.

To prove $iii) \implies i)$, note that the existence of a stabilizing solution P implies that (A, B) is stabilizable, and that (12.16.11) implies that $\mathcal{H}$ has no imaginary eigenvalues. $\square$

Corollary 12.17.10. Assume that (A, B) is stabilizable and (A, E_1) is detectable. Then, (12.16.4) has a stabilizing solution.

12.18 The Maximal Solution of the Riccati Equation

In this section we consider the existence of the maximal solution of (12.16.4). Example 12.16.3 shows that the assumptions of Proposition 12.19.1 are not sufficient to guarantee that (12.16.4) has a maximal solution.

Theorem 12.18.1. The following statements are equivalent:

- i) (A, B) is stabilizable.
- ii) (12.16.4) has a solution $P_{\max}$ that is positive semidefinite, maximal, and satisfies $\text{spec}(A - \Sigma P_{\max}) \subset \text{CLHP}$.

Proof. The result $i) \implies ii)$ is given by Theorem 2.1 and Theorem 2.2 of [561]. See also (i) of Theorem 13.11 of [1498]. The converse result follows from Corollary 3 of [1166]. $\square$

Proposition 12.18.2. Assume that (12.16.4) has a maximal solution $P_{\max}$, let P be a solution of (12.16.4), and assume that $\text{spec}(A - \Sigma P_{\max}) \subset \text{CLHP}$ and $\text{spec}(A - \Sigma P) \subset \text{CLHP}$. Then, $P = P_{\max}$.

Proof. It follows from i) of Proposition 12.16.14 that $\text{spec}(A - \Sigma P) = \text{spec}(A - \Sigma P_{\max})$. Since $P_{\max}$ is the maximal solution of (12.16.4), it follows that $P \leq P_{\max}$. Consequently, it follows from the contrapositive form of the second statement of Theorem 8.4.9 that $P = P_{\max}$. $\square$

Proposition 12.18.3. Assume that (12.16.4) has a solution P such that $\text{spec}(A - \Sigma P) \subset \text{CLHP}$. Then, P is stabilizing if and only if $\mathcal{H}$ has no imaginary eigenvalues

It follows from Proposition 12.18.2 that (12.16.4) has at most one positive-semidefinite solution P such that $\text{spec}(A - \Sigma P) \subset \text{CLHP}$. Consequently, (12.16.4) has at most one positive-semidefinite stabilizing solution.

Theorem 12.18.4. The following statements hold:

- i) (12.16.4) has at most one stabilizing solution.
- ii) If P is the stabilizing solution of (12.16.4), then P is positive semidefinite.
- iii) If P is the stabilizing solution of (12.16.4), then P is maximal.

Proof. To prove i), assume that (12.16.4) has stabilizing solutions P_1 and P_2 . Then, (A, B) is stabilizable, and Theorem 12.18.1 implies that (12.16.4) has a maximal solution $P_{\max}$ such that $\text{spec}(A - \Sigma P_{\max}) \subset \text{CLHP}$. Now, Proposition 12.18.2 implies that $P_1 = P_{\max}$ and $P_2 = P_{\max}$. Hence, $P_1 = P_2$.

Alternatively, suppose that (12.16.4) has the stabilizing solutions P_1 and P_2 . Then,

$$A^T P_1 + P_1 A + R_1 - P_1 \Sigma P_1 = 0,$$

$$A^T P_2 + P_2 A + R_1 - P_2 \Sigma P_2 = 0.$$

Subtracting these equations and rearranging yields

$$(A - \Sigma P_1)^T(P_1 - P_2) + (P_1 - P_2)(A - \Sigma P_2) = 0.$$

Since $A - \Sigma P_1$ and $A - \Sigma P_2$ are asymptotically stable, it follows from Proposition 11.9.3 and Fact 11.18.33 that $P_1 - P_2 = 0$. Hence, (12.16.4) has at most one stabilizing solution.

Next, to prove *ii*), suppose that P is a stabilizing solution of (12.16.4). Then, it follows from (12.16.4) that

$$P = \int_0^\infty e^{t(A-\Sigma P)^T} (R_1 + P\Sigma P) e^{t(A-\Sigma P)} dt,$$

which shows that P is positive semidefinite.

To prove *iii*), let P' be a solution of (12.16.4). Then, it follows that

$$(A - \Sigma P')^T(P - P') + (P - P')(A - \Sigma P) + (P - P')\Sigma(P - P') = 0,$$

which implies that $P' \leq P$. Thus, P is also the maximal solution of (12.16.4). $\square$

The following results concerns the monotonicity of solutions of the Riccati equation (12.16.4).

Proposition 12.18.5. Assume that (A, B) is stabilizable, and let $P_{\max}$ denote the maximal solution of (12.16.4). Furthermore, let $\hat{R}_1 \in \mathbb{R}^{n \times n}$ be positive semidefinite, let $\hat{R}_2 \in \mathbb{R}^{m \times m}$ be positive definite, let $\hat{A} \in \mathbb{R}^{n \times n}$, let $\hat{B} \in \mathbb{R}^{n \times m}$, define $\hat{\Sigma} \triangleq \hat{B}\hat{R}_2^{-1}\hat{B}^T$, assume that

$$\begin{bmatrix} \hat{R}_1 & \hat{A}^T \\ \hat{A} & -\hat{\Sigma} \end{bmatrix} \leq \begin{bmatrix} R_1 & A^T \\ A & -\Sigma \end{bmatrix},$$

and let $\hat{P}$ be a solution of

$$\hat{A}^T \hat{P} + \hat{P} \hat{A} + \hat{R}_1 - \hat{P} \hat{\Sigma} \hat{P} = 0. \quad (12.18.1)$$

Then,

$$\hat{P} \leq P_{\max}. \quad (12.18.2)$$

Proof. The result is given by Theorem 1 of [1441]. $\square$

Corollary 12.18.6. Assume that (A, B) is stabilizable, let $\hat{R}_1 \in \mathbb{R}^{n \times n}$ be positive semidefinite, assume that $\hat{R}_1 \leq R_1$, and let $P_{\max}$ and $\hat{P}_{\max}$ denote, respectively, the maximal solutions of (12.16.4) and

$$A^T P + PA + \hat{R}_1 - P\Sigma P = 0. \quad (12.18.3)$$

Then,

$$\hat{P}_{\max} \leq P_{\max}. \quad (12.18.4)$$

Proof. The result follows from Proposition 12.18.5 or Theorem 2.3 of [561]. $\square$

The following result shows that, if $R_1 = 0$, then the closed-loop eigenvalues of the closed-loop dynamics obtained from the maximal solution consist of the CLHP open-loop eigenvalues and reflections of the ORHP open-loop eigenvalues.

Proposition 12.18.7. Assume that (A, B) is stabilizable, assume that $R_1 = 0$, and let $P \in \mathbb{R}^{n \times n}$ be a positive-semidefinite solution of (12.16.4). Then, P is the maximal solution of (12.16.4) if and only if

$$\text{mspec}(A - \Sigma P) = [\text{mspec}(A) \cap \text{CLHP}] \cup [\text{mspec}(-A) \cap \text{OLHP}]. \quad (12.18.5)$$

Proof. Sufficiency follows from Proposition 12.18.2. To prove necessity, note that it follows from the definition (12.16.8) of $\mathcal{H}$ with $R_1 = 0$ and from *iv*) of Proposition 12.16.14 that

$$\text{mspec}(A) \cup \text{mspec}(-A) = \text{mspec}(A - \Sigma P) \cup \text{mspec}[-(A - \Sigma P)].$$

Now, Theorem 12.18.1 implies that $\text{mspec}(A - \Sigma P) \subseteq \text{CLHP}$, which implies that (12.18.5) is satisfied. $\square$

Corollary 12.18.8. Let $R_1 = 0$, and assume that $\text{spec}(A) \subset \text{CLHP}$. Then, $P = 0$ is the only positive-semidefinite solution of (12.16.4).

12.19 Positive-Semidefinite and Positive-Definite Solutions of the Riccati Equation

The following result gives sufficient conditions under which (12.16.4) has a positive-semidefinite solution.

Proposition 12.19.1. Assume that there exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.17.2) and (12.17.3) are satisfied, where $(\begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix}, \begin{bmatrix} B_1 \\ B_2 \end{bmatrix})$ is controllable, $(\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}, \begin{bmatrix} E_{11} & E_{13} \end{bmatrix})$ is observable, and A_3 is asymptotically stable. Then, (12.16.4) has a positive-semidefinite solution.

Proof. First, rewrite (12.17.2) and (12.17.3) as

$$A = S \begin{bmatrix} A_1 & A_{13} & 0 & 0 \\ 0 & A_3 & 0 & 0 \\ A_{21} & A_{23} & A_2 & A_{24} \\ 0 & A_{43} & 0 & A_4 \end{bmatrix} S^{-1}, \quad B = S \begin{bmatrix} B_1 \\ 0 \\ B_2 \\ 0 \end{bmatrix},$$

$$E_1 = \begin{bmatrix} E_{11} & E_{13} & 0 & 0 \end{bmatrix} S^{-1},$$

where $(\begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix}, \begin{bmatrix} B_1 \\ B_2 \end{bmatrix})$ is controllable, $(\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}, \begin{bmatrix} E_{11} & E_{13} \end{bmatrix})$ is observable, and A_3 is asymptotically stable. Since $(\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}, \begin{bmatrix} B_1 \\ 0 \end{bmatrix})$ is stabilizable, it follows from Theorem 12.18.1 that there exists a positive-semidefinite matrix $\hat{P}_1$ that satisfies

$$\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}^T \hat{P}_1 + \hat{P}_1 \begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix} + \begin{bmatrix} E_{11}^T E_{11} & E_{11}^T E_{13} \\ E_{13}^T E_{11} & E_{13}^T E_{13} \end{bmatrix} - \hat{P}_1 \begin{bmatrix} B_1 R_2^{-1} B_1^T & 0 \\ 0 & 0 \end{bmatrix} \hat{P}_1 = 0.$$

Consequently, $P \triangleq S^T \text{diag}(\hat{P}_1, 0, 0) S$ is a positive-semidefinite solution of (12.16.4).

□

Corollary 12.19.2. Assume that (A, B) is stabilizable. Then, (12.16.4) has a positive-semidefinite solution P . If, in addition, (A, E_1) is detectable, then P is the stabilizing solution of (12.16.4), and thus P is the only positive-semidefinite solution of (12.16.4). Finally, if (A, E_1) is observable, then P is positive definite.

Proof. The first statement is given by Theorem 12.18.1. Next, assume that (A, E_1) is detectable. Then, Theorem 12.17.2 implies that P is a stabilizing solution of (12.16.4), which is the only positive-semidefinite solution of (12.16.4). Finally, Theorem 12.17.2 implies that, if (A, E_1) is observable, then P is positive definite. □

The next result gives necessary and sufficient conditions under which (12.16.4) has a positive-definite solution.

Proposition 12.19.3. The following statements are equivalent:

- i) (12.16.4) has a positive-definite solution.
- ii) There exists a nonsingular matrix $S \in \mathbb{R}^{n \times n}$ such that (12.17.2) and (12.17.3) are satisfied, where $(\begin{bmatrix} A_1 & 0 \\ A_{21} & A_2 \end{bmatrix}, \begin{bmatrix} B_1 \\ B_2 \end{bmatrix})$ is controllable, $(\begin{bmatrix} A_1 & A_{13} \\ 0 & A_3 \end{bmatrix}, \begin{bmatrix} E_{11} & E_{13} \end{bmatrix})$ is observable, A_3 is asymptotically stable, $-A_2$ is asymptotically stable, $\text{spec}(A_4) \subset j\mathbb{R}$, and A_4 is semisimple.

In this case, (12.16.4) has exactly one positive-definite solution if and only if A_4 is empty, and infinitely many positive-definite solutions if and only if A_4 is not empty.

Proof. See [1124]. □

Proposition 12.19.4. Assume that (12.16.4) has a stabilizing solution P , and let $S \in \mathbb{R}^{n \times n}$ be a nonsingular matrix such that (12.17.2) and (12.17.3) are satisfied, where (A_1, B_1, E_{11}) is controllable and observable, (A_2, B_2) is controllable, (A_3, E_{13}) is observable, $\nu_0(A_2) = 0$, and A_3 and A_4 are asymptotically stable. Then,

$$\text{def } P = \nu_-(A_2). \quad (12.19.1)$$

Hence, P is positive definite if and only if $\text{spec}(A_2) \subset \text{ORHP}$.

12.20 Facts on Stability, Observability, and Controllability

Fact 12.20.1. Let $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, and $C \in \mathbb{R}^{p \times n}$, and assume that (A, B) is controllable and (A, C) is observable. Then, for all $v \in \mathbb{R}^m$, the step response

$$y(t) = \int_0^t C e^{t\tau} d\tau B v + D v$$

is bounded on $[0, \infty)$ if and only if A is Lyapunov stable and nonsingular.

Fact 12.20.2. Let $A \in \mathbb{R}^{n \times n}$ and $C \in \mathbb{R}^{p \times n}$, assume that (A, C) is detectable, and let $x(t)$ and $y(t)$ satisfy $\dot{x}(t) = Ax(t)$ and $y(t) = Cx(t)$ for $t \in [0, \infty)$. Then, the following statements hold:

- i) y is bounded if and only if x is bounded.
- ii) $\lim_{t \rightarrow \infty} y(t)$ exists if and only if $\lim_{t \rightarrow \infty} x(t)$ exists.
- iii) $y(t) \rightarrow 0$ as $t \rightarrow \infty$ if and only if $x(t) \rightarrow 0$ as $t \rightarrow \infty$.

Fact 12.20.3. Let $x(0) = x_0$, and let $x_f - e^{t_f A} x_0 \in \mathcal{C}(A, B)$. Then, for all $t \in [0, t_f]$, the control $u: [0, t_f] \mapsto \mathbb{R}^m$ defined by

$$u(t) \triangleq B^T e^{(t_f - t)A^T} \left(\int_0^{t_f} e^{\tau A} B B^T e^{\tau A^T} d\tau \right)^+ (x_f - e^{t_f A} x_0)$$

yields $x(t_f) = x_f$.

Fact 12.20.4. Let $x(0) = x_0$, let $x_f \in \mathbb{R}^n$, and assume that (A, B) is controllable. Then, for all $t \in [0, t_f]$, the control $u: [0, t_f] \mapsto \mathbb{R}^m$ defined by

$$u(t) \triangleq B^T e^{(t_f - t)A^T} \left(\int_0^{t_f} e^{\tau A} B B^T e^{\tau A^T} d\tau \right)^{-1} (x_f - e^{t_f A} x_0)$$

yields $x(t_f) = x_f$.

Fact 12.20.5. Let $A \in \mathbb{R}^{n \times n}$, let $B \in \mathbb{R}^{n \times m}$, assume that A is skew symmetric, and assume that (A, B) is controllable. Then, for all $\alpha > 0$, $A - \alpha B B^T$ is asymptotically stable.

Fact 12.20.6. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$. Then, (A, B) is (controllable, stabilizable) if and only if $(A, B B^T)$ is (controllable, stabilizable). Now, assume that B is positive semidefinite. Then, (A, B) is (controllable, stabilizable) if and only if $(A, B^{1/2})$ is (controllable, stabilizable).

Fact 12.20.7. Let $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, and $\hat{B} \in \mathbb{R}^{n \times \hat{m}}$, and assume that (A, B) is (controllable, stabilizable) and $\mathcal{R}(B) \subseteq \mathcal{R}(\hat{B})$. Then, $(A, \hat{B})$ is also (controllable, stabilizable).

Fact 12.20.8. Let $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, and $\hat{B} \in \mathbb{R}^{n \times \hat{m}}$, and assume that (A, B) is (controllable, stabilizable) and $B B^T \leq \hat{B} \hat{B}^T$. Then, $(A, \hat{B})$ is also (controllable, stabilizable). (Proof: Use Lemma 8.6.1 and Fact 12.20.7.)

Fact 12.20.9. Let $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, $\hat{B} \in \mathbb{R}^{n \times \hat{m}}$, and $\hat{C} \in \mathbb{R}^{\hat{m} \times n}$, and assume that (A, B) is (controllable, stabilizable). Then,

$$(A + \hat{B} \hat{C}, [B B^T + \hat{B} \hat{B}^T]^{1/2})$$

is also (controllable, stabilizable). (Proof: See [1455, p. 79].)

Fact 12.20.10. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$. Then, the following statements are equivalent:

- i) (A, B) is controllable.
- ii) There exists $\alpha \in \mathbb{R}$ such that $(A + \alpha I, B)$ is controllable.
- iii) $(A + \alpha I, B)$ is controllable for all $\alpha \in \mathbb{R}$.

Fact 12.20.11. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$. Then, the following statements are equivalent:

- i) (A, B) is stabilizable.
- ii) There exists $\alpha \leq \max\{0, -\text{spabs}(A)\}$ such that $(A + \alpha I, B)$ is stabilizable.
- iii) $(A + \alpha I, B)$ is stabilizable for all $\alpha \leq \max\{0, -\text{spabs}(A)\}$.

Fact 12.20.12. Let $A \in \mathbb{R}^{n \times n}$, assume that A is diagonal, and let $B \in \mathbb{R}^{n \times 1}$. Then, (A, B) is controllable if and only if the diagonal entries of A are distinct and every entry of B is nonzero. (Proof: Note that

$$\begin{aligned} \det \mathcal{K}(A, B) &= \det \begin{bmatrix} b_1 & & 0 \\ & \ddots & \\ 0 & & b_n \end{bmatrix} \begin{bmatrix} 1 & a_1 & \cdots & a_1^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & a_n & \cdots & a_n^{n-1} \end{bmatrix} \\ &= \left(\prod_{i=1}^n b_i \right) \prod_{i < j} (a_i - a_j). \end{aligned}$$

Fact 12.20.13. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times 1}$, and assume that (A, B) is controllable. Then, A is cyclic. (Proof: See Fact 5.14.9.)

Fact 12.20.14. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$, and assume that (A, B) is controllable. Then,

$$\max_{\lambda \in \text{spec}(A)} \text{gmult}_A(\lambda) \leq m.$$

Fact 12.20.15. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$. Then, the following conditions are equivalent:

- i) (A, B) is (controllable, stabilizable) and A is nonsingular.
- ii) (A, AB) is (controllable, stabilizable).

Fact 12.20.16. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$, and assume that (A, B) is controllable. Then, $(A, B^T S^{-T})$ is observable, where $S \in \mathbb{R}^{n \times n}$ is a nonsingular matrix satisfying $A^T = S^{-1}AS$.

Fact 12.20.17. Let (A, B) be controllable, let $t_1 > 0$, and define

$$P = \left(\int_0^{t_1} e^{-tA} B B^T e^{-tA^T} dt \right)^{-1}.$$

Then, $A - BB^T P$ is asymptotically stable. (Proof: P satisfies

$$(A - BB^T P)^T P + P(A - BB^T P) + P(BB^T + e^{t_1 A} B B^T e^{t_1 A^T}) P = 0.$$

Since $(A - BB^T P, BB^T + e^{t_1 A} B B^T e^{t_1 A^T})$ is observable and P is positive definite, Proposition 11.9.5 implies that $A - BB^T P$ is asymptotically stable.) (Remark: This result is due to Lukes and Kleinman. See [1152, pp. 113, 114].)

Fact 12.20.18. Let $A \in \mathbb{R}^{n \times n}$ and $B \in \mathbb{R}^{n \times m}$, assume that A is asymptotically stable, and, for $t \geq 0$, consider the linear system $\dot{x} = Ax + Bu$. Then, if u is bounded, then x is bounded. Furthermore, if $u(t) \rightarrow 0$ as $t \rightarrow \infty$, then $x(t) \rightarrow 0$ as $t \rightarrow \infty$. (Proof: See [1212, p. 330].) (Remark: These results are consequences of *input-to-state stability*.)

Fact 12.20.19. Let $A \in \mathbb{R}^{n \times n}$ and $C \in \mathbb{R}^{l \times n}$, assume that (A, C) is observable, define

$$\mathcal{O}_k(A, C) \triangleq \begin{bmatrix} C \\ CA \\ CA^2 \\ \vdots \\ CA^k \end{bmatrix},$$

and assume that $k \geq n - 1$. Then,

$$A = \begin{bmatrix} 0_{l \times n} \\ \mathcal{O}_k(A, C) \end{bmatrix}^+ \mathcal{O}_{k+1}(A, C).$$

(Remark: This result is due to Palanthandalam-Madapusi.)

12.21 Facts on the Lyapunov Equation and Inertia

Fact 12.21.1. Let $A, P \in \mathbb{F}^{n \times n}$, assume that P is Hermitian, let $C \in \mathbb{F}^{l \times n}$, and assume that $A^* P + PA + C^* C = 0$. Then, the following statements hold:

$$i) |\nu_-(A) - \nu_+(P)| \leq n - \text{rank } \mathcal{O}(A, C).$$

$$ii) |\nu_+(A) - \nu_-(P)| \leq n - \text{rank } \mathcal{O}(A, C).$$

$$iii) \text{ If } \nu_0(A) = 0, \text{ then}$$

$$|\nu_-(A) - \nu_+(P)| + |\nu_+(A) - \nu_-(P)| \leq n - \text{rank } \mathcal{O}(A, C).$$

If, in addition, (A, C) is observable, then the following statements hold:

$$iv) \nu_-(A) = \nu_+(P).$$

$$v) \nu_0(A) = \nu_0(P) = 0.$$

$$vi) \nu_+(A) = \nu_-(P).$$

$$vii) \text{ If } P \text{ is positive definite, then } A \text{ is asymptotically stable.}$$

(Proof: See [64, 312, 930, 1437] and [867, p. 448].) (Remark: $v)$ does not follow

from i – iii .) (Remark: For related results, see [1054] and references given in [930]. See also [289, 372].)

Fact 12.21.2. Let $A, P \in \mathbb{F}^{n \times n}$, assume that P is nonsingular and Hermitian, and assume that $A^*P + PA$ is negative semidefinite. Then, the following statements hold:

- i) $\nu_-(A) \leq \nu_+(P)$.
- ii) $\nu_+(A) \leq \nu_-(P)$.
- iii) If P is positive definite, then $\text{spec}(A) \subset \text{CLHP}$.

(Proof: See [867, p. 447].) (Remark: If P is positive definite, then A is Lyapunov stable, although this result does not follow from i) and ii .)

Fact 12.21.3. Let $A, P \in \mathbb{F}^{n \times n}$, and assume that $\nu_0(A) = 0$, P is Hermitian, and $A^*P + PA$ is negative semidefinite. Then, the following statements hold:

- i) $\nu_-(P) \leq \nu_+(A)$.
- ii) $\nu_+(P) \leq \nu_-(A)$.
- iii) If P is nonsingular, then $\nu_-(P) = \nu_+(A)$ and $\nu_+(P) = \nu_-(A)$.
- iv) If P is positive definite, then A is asymptotically stable.

(Proof: See [867, p. 447].)

Fact 12.21.4. Let $A, P \in \mathbb{F}^{n \times n}$, and assume that $\nu_0(A) = 0$, P is nonsingular and Hermitian, and $A^*P + PA$ is negative semidefinite. Then, the following statements hold:

- i) $\nu_-(A) = \nu_+(P)$.
- ii) $\nu_+(A) = \nu_-(P)$.

(Proof: Combine Fact 12.21.2 and Fact 12.21.3. See [867, p. 448].) (Remark: This result is due to Carlson and Schneider.)

Fact 12.21.5. Let $A, P \in \mathbb{F}^{n \times n}$, assume that P is Hermitian, and assume that $A^*P + PA$ is negative definite. Then, the following statements hold:

- i) $\nu_-(A) = \nu_+(P)$.
- ii) $\nu_0(A) = 0$.
- iii) $\nu_+(A) = \nu_-(P)$.
- iv) P is nonsingular.
- v) If P is positive definite, then A is asymptotically stable.

(Proof: See [447, pp. 441, 442], [867, p. 445], or [1054]. This result follows from Fact 12.21.1 with positive-definite $C = -(A^*P + PA)^{1/2}$.) (Remark: This result is due to Krein, Ostrowski, and Schneider.) (Remark: These conditions are the *classical constraints*. An analogous result holds for the discrete-time Lyapunov equation, where the analogous definition of inertia counts the numbers of eigenvalues inside

the open unit disk, outside the open unit disk, and on the unit circle. See [280, 393].)

Fact 12.21.6. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) $\nu_0(A) = 0$.
- ii) There exists a nonsingular Hermitian matrix $P \in \mathbb{F}^{n \times n}$ such that $A^*P + PA$ is negative definite.
- iii) There exists a Hermitian matrix $P \in \mathbb{F}^{n \times n}$ such that $A^*P + PA$ is negative definite.

In this case, the following statements hold for P given by ii) and iii):

- iv) $\nu_-(A) = \nu_+(P)$.
- v) $\nu_0(A) = \nu_0(P) = 0$.
- vi) $\nu_+(A) = \nu_-(P)$.
- vii) P is nonsingular.
- viii) If P is positive definite, then A is asymptotically stable.

(Proof: For the result i) $\implies$ ii), see [867, p. 445]. The result iii) $\implies$ i) follows from Fact 12.21.5. See [51, 280, 291].)

Fact 12.21.7. Let $A \in \mathbb{F}^{n \times n}$. Then, the following statements are equivalent:

- i) A is Lyapunov stable.
- ii) There exists a positive-definite matrix $P \in \mathbb{F}^{n \times n}$ such that $A^*P + PA$ is negative semidefinite.

Furthermore, the following statements are equivalent:

- iii) A is asymptotically stable.
- iv) There exists a positive-definite matrix $P \in \mathbb{F}^{n \times n}$ such that $A^*P + PA$ is negative definite.
- v) For every positive-definite matrix $R \in \mathbb{F}^{n \times n}$, there exists a positive-definite matrix $P \in \mathbb{F}^{n \times n}$ such that $A^*P + PA$ is negative definite.

(Remark: See Proposition 11.9.5 and Proposition 11.9.6.)

Fact 12.21.8. Let $A, P \in \mathbb{F}^{n \times n}$, and assume P is Hermitian. Then, the following statements hold:

- i) $\nu_+(A^*P + PA) \leq \text{rank } P$.
- ii) $\nu_-(A^*P + PA) \leq \text{rank } P$.

If, in addition, A is asymptotically stable, then the following statement holds:

- iii) $1 \leq \nu_-(A^*P + PA) \leq \text{rank } P$.

(Proof: See [120, 393].)

Fact 12.21.9. Let $A, P \in \mathbb{R}^{n \times n}$, assume that $\nu_0(A) = n$, and assume that P is positive semidefinite. Then, exactly one of the following statements holds:

$$i) \quad A^T P + PA = 0.$$

$$ii) \quad \nu_-(A^T P + PA) \geq 1 \text{ and } \nu_+(A^T P + PA) \geq 1.$$

(Proof: See [1348].)

Fact 12.21.10. Let $R \in \mathbb{F}^{n \times n}$, and assume that R is Hermitian and $\nu_+(R) \geq 1$. Then, there exist an asymptotically stable matrix $A \in \mathbb{F}^{n \times n}$ and a positive-definite matrix $P \in \mathbb{F}^{n \times n}$ such that $A^* P + PA + R = 0$. (Proof: See [120].)

Fact 12.21.11. Let $A \in \mathbb{F}^{n \times n}$, assume that A is cyclic, and let a, b, c, d, e be nonnegative integers such that $a + b = c + d + e = n$, $c \geq 1$, and $e \geq 1$. Then, there exists a nonsingular, Hermitian matrix $P \in \mathbb{F}^{n \times n}$ such that

$$\text{In } P = \begin{bmatrix} a \\ 0 \\ b \end{bmatrix}$$

and

$$\text{In}(A^* P + PA) = \begin{bmatrix} c \\ d \\ e \end{bmatrix}.$$

(Proof: See [1199].) (Remark: See also [1198].)

Fact 12.21.12. Let $P, R \in \mathbb{F}^{n \times n}$, and assume that P is positive and R is Hermitian. Then, the following statements are equivalent:

$$i) \quad \text{tr } RP^{-1} > 0.$$

$$ii) \quad \text{There exists an asymptotically stable matrix } A \in \mathbb{F}^{n \times n} \text{ such that } A^* P + PA + R = 0.$$

(Proof: See [120].)

Fact 12.21.13. Let $A_1 \in \mathbb{R}^{n_1 \times n_1}$, $A_2 \in \mathbb{R}^{n_2 \times n_2}$, $B \in \mathbb{R}^{n_1 \times m}$, and $C \in \mathbb{R}^{m \times n_2}$, assume that $A_1 \oplus A_2$ is nonsingular, and assume that $\text{rank } B = \text{rank } C = m$. Furthermore, let $X \in \mathbb{R}^{n_1 \times n_2}$ be the unique solution of

$$A_1 X + X A_2 + BC = 0.$$

Then,

$$\text{rank } X \leq \min\{\text{rank } \mathcal{K}(A_1, B), \text{rank } \mathcal{O}(A_2, C)\}.$$

Furthermore, equality holds if $m = 1$. (Proof: See [390].) (Remark: Related results are given in [1437, 1443].)

Fact 12.21.14. Let $A_1, A_2 \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^n$, $C \in \mathbb{R}^{1 \times n}$, assume that $A_1 \oplus A_2$ is nonsingular, let $X \in \mathbb{R}^{n \times n}$ satisfy

$$A_1 X + X A_2 + BC = 0,$$

and assume that (A_1, B) is controllable and (A_2, C) is observable. Then, X is nonsingular. (Proof: See Fact 12.21.13 and [1443].)

Fact 12.21.15. Let $A, P, R \in \mathbb{R}^{n \times n}$, and assume that P and R are positive semidefinite, $A^T P + PA + R = 0$, and $\mathcal{N}[\mathcal{O}(A, R)] = \mathcal{N}(A)$. Then, A is semistable. (Proof: See [195].)

Fact 12.21.16. Let $A, V \in \mathbb{R}^{n \times n}$, assume that A is asymptotically stable, assume that V is positive semidefinite, and let $Q \in \mathbb{R}^{n \times n}$ be the unique, positive-definite solution to $AQ + QA^T + V = 0$. Furthermore, let $C \in \mathbb{R}^{l \times n}$, and assume that CVC^T is positive definite. Then, CQC^T is positive definite.

Fact 12.21.17. Let $A, R \in \mathbb{R}^{n \times n}$, assume that A is asymptotically stable, assume that $R \in \mathbb{R}^{n \times n}$ is positive semidefinite, and let $P \in \mathbb{R}^{n \times n}$ satisfy $A^T P + PA + R = 0$. Then, for all $i, j = 1, \dots, n$, there exist $\alpha_{ij} \in \mathbb{R}$ such that

$$P = \sum_{i,j=1}^n \alpha_{ij} A^{(i-1)T} R A^{j-1}.$$

In particular, for all $i, j = 1, \dots, n$, $\alpha_{ij} = \hat{P}_{(i,j)}$, where $\hat{P} \in \mathbb{R}^{n \times n}$ satisfies $\hat{A}^T \hat{P} + \hat{P} \hat{A} + \hat{R} = 0$, where $\hat{A} = C(\chi_A)$ and $\hat{R} = E_{1,1}$. (Proof: See [1204].) (Remark: This identity is *Smith's method*. See [391, 413, 644, 940] for finite-sum solutions of linear matrix equations.)

Fact 12.21.18. Let $\lambda_1, \dots, \lambda_n \in \mathbb{C}$, assume that, for all $i = 1, \dots, n$, $\operatorname{Re} \lambda_i < 0$, define $\Lambda \triangleq \operatorname{diag}(\lambda_1, \dots, \lambda_n)$, let k be a nonnegative integer, and, for all $i, j = 1, \dots, n$, define $P \in \mathbb{C}^{n \times n}$ by

$$P \triangleq \frac{1}{k!} \int_0^\infty t^k e^{\bar{\Lambda} t} e^{\Lambda t} dt.$$

Then, P is positive definite, P satisfies the Lyapunov equation

$$\bar{\Lambda} P + P \Lambda + I = 0,$$

and, for all $i, j = 1, \dots, n$,

$$P_{(i,j)} = \left(\frac{-1}{\bar{\lambda}_i + \lambda_j} \right)^{k+1}.$$

(Proof: For all nonzero $x \in \mathbb{C}^n$, it follows that

$$x^* P x = \int_0^\infty t^k \|e^{\Lambda t} x\|_2^2 dt,$$

is positive. Hence, P is positive definite. Furthermore, note that

$$P_{(i,j)} = \int_0^\infty t^k e^{\bar{\lambda}_i t} e^{\lambda_j t} dt = \frac{(-1)^{k+1} k!}{(\bar{\lambda}_i + \lambda_j)^{k+1}}.$$

(Remark: See [262] and [711, p. 348].) (Remark: See Fact 8.8.16 and Fact 12.21.19.)

Fact 12.21.19. Let $\lambda_1, \dots, \lambda_n \in \mathbb{C}$, assume that, for all $i = 1, \dots, n$, $\operatorname{Re} \lambda_i < 0$, define $\Lambda \triangleq \operatorname{diag}(\lambda_1, \dots, \lambda_n)$, let k be a nonnegative integer, let $R \in \mathbb{C}^{n \times n}$, assume that R is positive semidefinite, and, for all $i, j = 1, \dots, n$, define $P \in \mathbb{C}^{n \times n}$ by

$$P \triangleq \frac{1}{k!} \int_0^\infty t^k e^{\bar{\Lambda} t} R e^{\Lambda t} dt.$$

Then, P is positive semidefinite, P satisfies the Lyapunov equation

$$\bar{\Lambda}P + P\Lambda + R = 0,$$

and, for all $i, j = 1, \dots, n$,

$$P_{(i,j)} = R_{(i,j)} \left(\frac{-1}{\bar{\lambda}_i + \lambda_j} \right)^{k+1}.$$

If, in addition, $I \circ R$ is positive definite, then P is positive definite. (Proof: Use Fact 8.21.12 and Fact 12.21.18.) (Remark: See Fact 8.8.16 and Fact 12.21.18. Note that $P = \hat{P} \circ R$, where $\hat{P}$ is the solution to the Lyapunov equation with $R = I$.)

Fact 12.21.20. Let $A, R \in \mathbb{R}^{n \times n}$, assume that $R \in \mathbb{R}^{n \times n}$ is positive semidefinite, let $q, r \in \mathbb{R}$, where $r > 0$, and assume that there exists a positive-definite matrix $P \in \mathbb{R}^{n \times n}$ satisfying

$$[A - (q + r)I]^T P + P[A - (q + r)I] + \frac{1}{r} A^T P A + R = 0.$$

Then, the spectrum of A is contained in a disk centered at $q + j0$ with radius r . (Remark: The disk is an *eigenvalue inclusion region*. See [141, 614, 1401] for related results concerning elliptical, parabolic, hyperbolic, sector, and vertical strip regions.)

12.22 Facts on Realizations and the H_2 System Norm

Fact 12.22.1. Let $x: [0, \infty) \mapsto \mathbb{R}^n$ and $y: [0, \infty) \mapsto \mathbb{R}^n$, assume that $\int_0^\infty x^T(t)x(t) dt$ and $\int_0^\infty y^T(t)y(t) dt$ exist, and let $\hat{x}: j\mathbb{R} \mapsto \mathbb{C}^n$ and $\hat{y}: j\mathbb{R} \mapsto \mathbb{C}^n$ denote the Fourier transforms of x and y , respectively. Then,

$$\int_0^\infty x^T(t)x(t) dt = \int_{-\infty}^\infty \hat{x}^*(j\omega)\hat{x}(j\omega) d\omega$$

and

$$\int_0^\infty x^T(t)y(t) dt = \int_{-\infty}^\infty \hat{x}^*(j\omega)\hat{y}(j\omega) d\omega.$$

(Remark: These identities are equivalent versions of Parseval's theorem. The second identity follows from the first identity by replacing x with $x + y$.)

Fact 12.22.2. Let $G \in \mathbb{R}^{l \times m}(s)$, where $G \stackrel{\min}{\sim} \left[\frac{A}{C} \middle| \frac{B}{D} \right]$, and assume that, for all $i = 1, \dots, l$ and $j = 1, \dots, m$, $G_{(i,j)} = p_{i,j}/q_{i,j}$, where $p_{i,j}, q_{i,j} \in \mathbb{R}[s]$ are coprime. Then,

$$\text{spec}(A) = \bigcup_{i,j=1}^{l,m} \text{roots}(p_{i,j}).$$

Fact 12.22.3. Let $G \sim \left[\frac{A}{C} \middle| \frac{B}{D} \right]$, let $a, b \in \mathbb{R}$, where $a \neq 0$, and define $H(s) \triangleq G(as + b)$. Then,

$$H \sim \left[\frac{a^{-1}(A - bI)}{a^{-1}C} \middle| \frac{B}{D} \right].$$

Fact 12.22.4. Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$, where A is nonsingular, and define $H(s) \triangleq G(1/s)$. Then,

$$H \sim \left[\begin{array}{c|c} A^{-1} & -A^{-1}B \\ \hline CA^{-1} & D - CA^{-1}B \end{array} \right].$$

Fact 12.22.5. Let $G(s) = C(sI - A)^{-1}B$. Then,

$$G(j\omega) = -CA(\omega^2 I + A^2)^{-1}B - j\omega C(\omega^2 I + A^2)^{-1}B.$$

Fact 12.22.6. Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$ and $H(s) = sG(s)$. Then,

$$H \sim \left[\begin{array}{c|c} A & B \\ \hline CA & CB \end{array} \right].$$

Consequently,

$$sC(sI - A)^{-1}B = CA(sI - A)^{-1}B + CB.$$

Fact 12.22.7. Let $G = \begin{bmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{bmatrix}$, where $G_{ij} \sim \left[\begin{array}{c|c} A_{ij} & B_{ij} \\ \hline C_{ij} & D_{ij} \end{array} \right]$ for all $i, j = 1, 2$. Then,

$$\begin{bmatrix} G_{11} & G_{12} \\ G_{21} & G_{22} \end{bmatrix} \sim \left[\begin{array}{cccc|cc} A_{11} & 0 & 0 & 0 & B_{11} & 0 \\ 0 & A_{12} & 0 & 0 & 0 & B_{12} \\ 0 & 0 & A_{21} & 0 & B_{21} & 0 \\ 0 & 0 & 0 & A_{22} & 0 & B_{22} \\ \hline C_{11} & C_{12} & 0 & 0 & D_{11} & D_{12} \\ 0 & 0 & C_{21} & C_{22} & D_{21} & D_{22} \end{array} \right].$$

Fact 12.22.8. Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$, where $G \in \mathbb{R}^{l \times m}(s)$, and let $M \in \mathbb{R}^{m \times l}$. Then,

$$[I + GM]^{-1} \sim \left[\begin{array}{c|c} A - BMC & B \\ \hline -C & I \end{array} \right]$$

and

$$[I + GM]^{-1}G \sim \left[\begin{array}{c|c} A - BMC & B \\ \hline C & 0 \end{array} \right].$$

Fact 12.22.9. Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$, where $G \in \mathbb{R}^{l \times m}(s)$. If D has a left inverse $D^L \in \mathbb{R}^{m \times l}$, then

$$G^L \sim \left[\begin{array}{c|c} A - BD^L C & BD^L \\ \hline -D^L C & D^L \end{array} \right]$$

satisfies $G^L G = I$. If D has a right inverse $D^R \in \mathbb{R}^{m \times l}$, then

$$G^R \sim \left[\begin{array}{c|c} A - BD^R C & BD^R \\ \hline -D^R C & D^R \end{array} \right]$$

satisfies $G G^R = I$.

Fact 12.22.10. Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & 0 \end{array} \right]$ be a SISO rational transfer function, and let $\lambda \in \mathbb{C}$. Then, there exists a rational function H such that

$$G(s) = \frac{1}{(s + \lambda)^r} H(s)$$

and such that λ is neither a pole nor a zero of H if and only if the Jordan form of A has exactly one block associated with λ , which is of order r .

Fact 12.22.11. Let $G \sim \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$. Then, $G(s)$ is given by the Schur complement

$$G(s) = (A - sI) \left[\begin{array}{cc} A - sI & B \\ C & D \end{array} \right].$$

(Remark: See [151].)

Fact 12.22.12. Let $G \in \mathbb{F}^{n \times m}(s)$, where $G \stackrel{\min}{\sim} \left[\begin{array}{c|c} A & B \\ \hline C & D \end{array} \right]$, and, for all $i = 1, \dots, n$ and $j = 1, \dots, m$, let $G_{(i,j)} = p_{ij}/q_{ij}$, where $p_{ij}, q_{ij} \in \mathbb{F}[s]$ are coprime. Then,

$$\bigcup_{i,j=1}^{n,m} \text{roots}(q_{ij}) = \text{spec}(A).$$

Fact 12.22.13. Let $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, and $C \in \mathbb{R}^{m \times n}$. Then,

$$\det[sI - (A + BC)] = \det[I - C(sI - A)^{-1}B] \det(sI - A).$$

If, in addition, $n = m = 1$, then

$$\det[sI - (A + BC)] = \det(sI - A) - C(sI - A)^A B.$$

(Remark: The last expression is used in [1009] to compute the frequency response of a transfer function.) (Proof: Note that

$$\begin{aligned} \det[I - C(sI - A)^{-1}B] \det(sI - A) &= \det \begin{bmatrix} sI - A & B \\ C & I \end{bmatrix} \\ &= \det \begin{bmatrix} sI - A & B \\ C & I \end{bmatrix} \begin{bmatrix} I & 0 \\ -C & I \end{bmatrix} \\ &= \det \begin{bmatrix} sI - A - BC & B \\ 0 & I \end{bmatrix} \\ &= \det(sI - A - BC). \end{aligned}$$

Fact 12.22.14. Let $A \in \mathbb{R}^{n \times n}$, $B \in \mathbb{R}^{n \times m}$, $C \in \mathbb{R}^{m \times n}$, and $K \in \mathbb{R}^{m \times n}$, and assume that $A + BK$ is nonsingular. Then,

$$\det \begin{bmatrix} A & B \\ C & 0 \end{bmatrix} = (-1)^m \det(A + BK) \det[C(A + BK)^{-1}B].$$

Hence, $\begin{bmatrix} A & B \\ C & 0 \end{bmatrix}$ is nonsingular if and only if $C(A + BK)^{-1}B$ is nonsingular. (Proof: Note that

$$\begin{aligned}
 \det \begin{bmatrix} A & B \\ C & 0 \end{bmatrix} &= \det \begin{bmatrix} A & B \\ C & 0 \end{bmatrix} \begin{bmatrix} I & 0 \\ K & I \end{bmatrix} \\
 &= \det \begin{bmatrix} A+BK & B \\ C & 0 \end{bmatrix} \\
 &= \det(A+BK) \det[-C(A+BK)^{-1}B].
 \end{aligned}$$

Fact 12.22.15. Let $A_1 \in \mathbb{R}^{n \times n}$, $C_1 \in \mathbb{R}^{1 \times n}$, $A_2 \in \mathbb{R}^{m \times m}$, and $B_2 \in \mathbb{R}^{m \times 1}$, let $\lambda \in \mathbb{C}$, assume that λ is an observable eigenvalue of (A_1, C_1) and a controllable eigenvalue of (A_2, B_2) , and define the dynamics matrix $\mathcal{A}$ of the cascaded system by

$$\mathcal{A} \triangleq \begin{bmatrix} A_1 & 0 \\ B_2 C_1 & A_2 \end{bmatrix}.$$

Then,

$$\text{amult}_{\mathcal{A}}(\lambda) = \text{amult}_{A_1}(\lambda) + \text{amult}_{A_2}(\lambda)$$

and

$$\text{gmult}_{\mathcal{A}}(\lambda) = 1.$$

(Remark: The eigenvalue λ is a cyclic eigenvalue of both subsystems as well as the cascaded system. In other words, λ , which occurs in a single Jordan block of each subsystem, occurs in a single Jordan block in the cascaded system. Effectively, the Jordan blocks of the subsystems corresponding to λ are merged.)

Fact 12.22.16. Let $G_1 \in \mathbb{R}^{l_1 \times m}(s)$ and $G_2 \in \mathbb{R}^{l_2 \times m}(s)$ be strictly proper. Then,

$$\left\| \begin{bmatrix} G_1 \\ G_2 \end{bmatrix} \right\|_{H_2}^2 = \|G_1\|_{H_2}^2 + \|G_2\|_{H_2}^2.$$

Fact 12.22.17. Let $G_1, G_2 \in \mathbb{R}^{m \times m}(s)$ be strictly proper. Then,

$$\left\| \begin{bmatrix} G_1 \\ G_2 \end{bmatrix} \right\|_{H_2} = \left\| \begin{bmatrix} G_1 & G_2 \end{bmatrix} \right\|_{H_2}.$$

Fact 12.22.18. Let $G(s) \triangleq \frac{\alpha}{s+\beta}$, where $\beta > 0$. Then,

$$\|G\|_{H_2} = \frac{|\alpha|}{\sqrt{2\beta}}.$$

Fact 12.22.19. Let $G(s) \triangleq \frac{\alpha_1 s + \alpha_0}{s^2 + \beta_1 s + \beta_0}$, where $\beta_0, \beta_1 > 0$. Then,

$$\|G\|_{H_2} = \sqrt{\frac{\alpha_0^2}{2\beta_0\beta_1} + \frac{\alpha_1^2}{2\beta_1}}.$$

Fact 12.22.20. Let $G_1(s) = \frac{\alpha_1}{s+\beta_1}$ and $G_2(s) = \frac{\alpha_2}{s+\beta_2}$, where $\beta_1, \beta_2 > 0$. Then,

$$\|G_1 G_2\|_{H_2} \leq \|G_1\|_{H_2} \|G_2\|_{H_2}$$

if and only if $\beta_1 + \beta_2 \geq 2$. (Remark: The H_2 norm is not submultiplicative.)

12.23 Facts on the Riccati Equation

Fact 12.23.1. Assume that (A, B) is stabilizable, and assume that $\mathcal{H}$ defined by (12.16.8) has an imaginary eigenvalue λ . Then, every Jordan block of $\mathcal{H}$ associated with λ has even order. (Proof: Let P be a solution of (12.16.4), and let $\mathcal{J}$ denote the Jordan form of $A - \Sigma P$. Then, there exists a nonsingular $2n \times 2n$ block-diagonal matrix $\mathcal{S}$ such that $\hat{\mathcal{H}} \triangleq \mathcal{S}^{-1}\mathcal{H}\mathcal{S} = \begin{bmatrix} \mathcal{J} & \hat{\Sigma} \\ 0 & -\mathcal{J}^T \end{bmatrix}$, where $\hat{\Sigma}$ is positive semidefinite. Next, let $\mathcal{J}_\lambda \triangleq \lambda I_r + N_r$ be a Jordan block of $\mathcal{J}$ associated with λ , and consider the submatrix of $\lambda I - \hat{\mathcal{H}}$ consisting of the rows and columns of $\lambda I - \mathcal{J}_\lambda$ and $\lambda I + \mathcal{J}_\lambda^T$. Since (A, B) is stabilizable, it follows that the rank of this submatrix is $2r - 1$. Hence, every Jordan block of $\mathcal{H}$ associated with λ has even order.) (Remark: Canonical forms for symplectic and Hamiltonian matrices are discussed in [873].)

Fact 12.23.2. Let $A, B \in \mathbb{C}^{n \times n}$, assume that A and B are positive definite, let $S \in \mathbb{C}^{n \times n}$, satisfy $A = S^*S$, and define

$$X \triangleq S^{-1}(SBS^*)^{1/2}S^{-*}.$$

Then, X satisfies $XAX = B$. (Proof: See [683, p. 52].)

Fact 12.23.3. Let $A, B \in \mathbb{C}^{n \times n}$, and assume that the $2n \times 2n$ matrix

$$\begin{bmatrix} A & -2I \\ 2B - \frac{1}{2}A^2 & A \end{bmatrix}$$

is simple. Then, there exists a matrix $X \in \mathbb{C}^{n \times n}$ satisfying

$$X^2 + AX + B = 0.$$

(Proof: See [1337].)

Fact 12.23.4. Let $A, B \in \mathbb{F}^{n \times n}$, and assume that A and B are positive semidefinite. Then, the following statements hold:

- i) If A is positive definite, then $X = A\#B$ is the unique positive-definite solution of

$$XA^{-1}X - B = 0.$$

- ii) If A is positive definite, then $X = \frac{1}{2}[-A + A\#(A + 4B)]$ is the unique positive-definite solution of

$$XA^{-1}X + X - B = 0.$$

- iii) If A is positive definite, then $X = \frac{1}{2}[A + A\#(A + 4B)]$ is the unique positive-definite solution of

$$XA^{-1}X - X - B = 0.$$

- iv) If B is positive definite, then $X = A\#B$ is the unique positive-definite solution of

$$XB^{-1}X = A.$$

- v) If A is positive definite, then $X = \frac{1}{2}[A + A\#(A + 4BA^{-1}B)]$ is the unique positive-definite solution of

$$BX^{-1}B - X + A = 0.$$

vi) If A is positive definite, then $X = \frac{1}{2}[-A + A\#(A + 4BA^{-1}B)]$ is the unique positive-definite solution of

$$BX^{-1}B - X - A = 0.$$

vii) If $0 < A \leq B$, then $X = \frac{1}{2}[A + A\#(4B - 3A)]$ is the unique positive-definite solution of

$$XA^{-1}X - X - (B - A) = 0.$$

viii) If $0 < A \leq B$, then $X = \frac{1}{2}[-A + A\#(4B - 3A)]$ is the unique positive-definite solution of

$$XA^{-1}X + X - (B - A) = 0.$$

ix) If $0 < A < B$, $X(0)$ is positive definite, and $X(t)$ satisfies

$$\dot{X} = -XA^{-1}X + X + (B - A),$$

then

$$\lim_{t \rightarrow \infty} X(t) = \frac{1}{2}[A + A\#(4B - 3A)].$$

x) If $0 < A < B$, $X(0)$ is positive definite, and $X(t)$ satisfies

$$\dot{X} = -XA^{-1}X - X + (B - A),$$

then

$$\lim_{t \rightarrow \infty} X(t) = \frac{1}{2}[A + A\#(4B - 3A)].$$

xi) If $0 < A < B$, $X(0)$ and $Y(0)$ are positive definite, $X(t)$ satisfies

$$\dot{X} = -XA^{-1}X + X + (B - A)$$

and $Y(t)$ satisfies

$$\dot{Y} = -YA^{-1}Y - Y + (B - A),$$

then

$$\lim_{t \rightarrow \infty} X(t)\#Y(t) = A\#(B - A).$$

(Proof: See [910].) (Remark: See Fact 8.10.43.) (Remark: The solution X given by vii) is the *golden mean* of A and B . In the scalar case with $A = 1$ and $B = 2$, the solution X of $X^2 - X - 1 = 0$ is the *golden ratio* $\frac{1}{2}(1 + \sqrt{5})$. See Fact 4.11.12.)

Fact 12.23.5. Let $P_0 \in \mathbb{R}^{n \times n}$, assume that P_0 is positive definite, and, for all $t \geq 0$, let $P(t) \in \mathbb{R}^{n \times n}$ satisfy

$$\begin{aligned} \dot{P}(t) &= A^T P(t) + P(t)A + P(t)VP(t), \\ P(0) &= P_0. \end{aligned}$$

Then, for all $t \geq 0$,

$$P(t) = e^{tA^T} \left[P_0^{-1} - \int_0^t e^{\tau A} V e^{\tau A^T} d\tau \right]^{-1} e^{tA}.$$

(Remark: $P(t)$ satisfies a Riccati differential equation.)

Fact 12.23.6. Let $G_c \sim \left[\begin{array}{c|c} A_c & B_c \\ \hline C_c & 0 \end{array} \right]$ denote an n th-order dynamic controller for the standard control problem. If G_c minimizes $\|\tilde{G}\|_2$, then G_c is given by

$$\begin{aligned} A_c &\triangleq A + BC_c - B_cC - B_cDC_c, \\ B_c &\triangleq (QC^T + V_{12})V_2^{-1}, \\ C_c &\triangleq -R_2^{-1}(B^TP + R_{12}^T), \end{aligned}$$

where P and Q are positive-semidefinite solutions to the algebraic Riccati equations

$$\begin{aligned} \hat{A}_R^TP + P\hat{A}_R - PBR_2^{-1}B^TP + \hat{R}_1 &= 0, \\ \hat{A}_EQ + Q\hat{A}_E^T - QC^TV_2^{-1}CQ + \hat{V}_1 &= 0, \end{aligned}$$

where $\hat{A}_R$ and $\hat{R}_1$ are defined by

$$\hat{A}_R \triangleq A - BR_2^{-1}R_{12}^T, \quad \hat{R}_1 \triangleq R_1 - R_{12}R_2^{-1}R_{12}^T,$$

and $\hat{A}_E$ and $\hat{V}_1$ are defined by

$$\hat{A}_E \triangleq A - V_{12}V_2^{-1}C, \quad \hat{V}_1 \triangleq V_1 - V_{12}V_2^{-1}V_{12}^T.$$

Furthermore, the eigenvalues of the closed-loop system are given by

$$\text{mspec} \left(\begin{bmatrix} A & BC_c \\ B_cC & A_c + B_cDC_c \end{bmatrix} \right) = \text{mspec}(A + BC_c) \cup \text{mspec}(A - B_cC).$$

Fact 12.23.7. Let $G_c \sim \left[\begin{array}{c|c} A_c & B_c \\ \hline C_c & 0 \end{array} \right]$ denote an n th-order dynamic controller for the discrete-time standard control problem. If G_c minimizes $\|\tilde{G}\|_2$, then G_c is given by

$$\begin{aligned} A_c &\triangleq A + BC_c - B_cC - B_cDC_c, \\ B_c &\triangleq (AQC^T + V_{12})(V_2 + CQC^T)^{-1}, \\ C_c &\triangleq -(R_2 + B^TPB)^{-1}(R_{12}^T + B^TPA), \\ D_c &\triangleq 0, \end{aligned}$$

and the eigenvalues of the closed-loop system are given by

$$\text{mspec} \left(\begin{bmatrix} A & BC_c \\ B_cC & A_c + B_cDC_c \end{bmatrix} \right) = \text{mspec}(A + BC_c) \cup \text{mspec}(A - B_cC).$$

Now, assume that $D = 0$ and $G_c \sim \left[\begin{array}{c|c} A_c & B_c \\ \hline C_c & D_c \end{array} \right]$. Then,

$$\begin{aligned} A_c &\triangleq A + BC_c - B_cC - BD_cC, \\ B_c &\triangleq (AQC^T + V_{12})(V_2 + CQC^T)^{-1} + BD_c, \\ C_c &\triangleq -(R_2 + B^TPB)^{-1}(R_{12}^T + B^TPA) - D_cC, \\ D_c &\triangleq (R_2 + B^TPB)^{-1}[B^TPAQC^T + R_{12}^TQC^T + B^TPV_{12}](V_2 + CQC^T)^{-1}, \end{aligned}$$

and the eigenvalues of the closed-loop system are given by

$$\text{mspec} \left(\begin{bmatrix} A + BD_cC & BC_c \\ B_cC & A_c \end{bmatrix} \right) = \text{mspec}(A + BC_c) \cup \text{mspec}(A - B_cC).$$

In both cases, P and Q are positive-semidefinite solutions to the discrete-time algebraic Riccati equations

$$\begin{aligned} P &= \hat{A}_R^T P \hat{A}_R - \hat{A}_R^T P B (R_2 + B^T P B)^{-1} B^T P \hat{A}_R + \hat{R}_1, \\ Q &= \hat{A}_E Q \hat{A}_E^T - \hat{A}_E Q C^T (V_2 + C Q C^T)^{-1} C Q \hat{A}_E^T + \hat{V}_1, \end{aligned}$$

where $\hat{A}_R$ and $\hat{R}_1$ are defined by

$$\hat{A}_R \triangleq A - B R_2^{-1} R_{12}^T, \quad \hat{R}_1 \triangleq R_1 - R_{12} R_2^{-1} R_{12}^T,$$

and $\hat{A}_E$ and $\hat{V}_1$ are defined by

$$\hat{A}_E \triangleq A - V_1 V_2^{-1} C, \quad \hat{V}_1 \triangleq V_1 - V_1 V_2^{-1} V_1^T.$$

(Proof: See [618].)

12.24 Notes

Linear system theory is treated in [261, 1150, 1336, 1450]. Time-varying linear systems are considered in [367, 1150], while discrete-time systems are emphasized in [660]. The equivalence of *iv*) and *v*) of Theorem 12.6.18 is the *PBH test*, due to [656]. Spectral factorization results are given in [337]. Stabilization aspects are discussed in [429]. Observable asymptotic stability and controllable asymptotic stability were introduced and used to analyze Lyapunov equations in [1207]. Zeros are treated in [21, 478, 787, 791, 943, 1074, 1154, 1178]. Matrix-based methods for linear system identification are developed in [1363], while stochastic theory is considered in [633].

Solutions of the LQR problem under weak conditions are given in [544]. Solutions of the Riccati equation are considered in [562, 845, 848, 864, 865, 974, 1124, 1434, 1441, 1446]. Proposition 12.16.16 is based on Theorem 3.6 of [1455, p. 79]. A variation of Theorem 12.18.1 is given without proof by Theorem 7.2.1 of [749, p. 125].

There are numerous extensions to the results given in this chapter relating to various generalizations of (12.16.4). These generalizations include the case in which R_1 is indefinite [561, 1438, 1440] as well as the case in which Σ is indefinite [1166]. The latter case is relevant to H_∞ optimal control theory [188]. Additional extensions include the Riccati inequality $A^T P + P A + R_1 - P \Sigma P \geq 0$ [1116, 1165, 1166, 1167], the discrete-time Riccati equation [8, 661, 743, 864, 1116, 1445], and fixed-order control [738].

Bibliography

- [1] A. Abdessemed and E. B. Davies, “Some Commutator Estimates in the Schatten Classes,” *J. London Math. Soc.*, Vol. s2-39, pp. 299–308, 1989. (Cited on page 584.)
- [2] R. Ablamowicz, “Matrix Exponential via Clifford Algebras,” *J. Nonlin. Math. Phys.*, Vol. 5, pp. 294–313, 1998. (Cited on page 673.)
- [3] H. Abou-Kandil, G. Freiling, V. Ionescu, and G. Jank, *Matrix Riccati Equations in Control and Systems Theory*. Basel: Birkhauser, 2003. (Cited on page xvii.)
- [4] S. Abramovich, J. Baric, M. Matic, and J. Pecaric, “On Van De Lune-Alzer’s Inequality,” *J. Math. Ineq.*, Vol. 1, pp. 563–587, 2007. (Cited on page 30.)
- [5] L. Aceto and D. Trigiant, “The Matrices of Pascal and Other Greats,” *Amer. Math. Monthly*, Vol. 108, pp. 232–245, 2001. (Cited on pages 354, 362, 447, and 672.)
- [6] S. Afriat, “Orthogonal and Oblique Projectors and the Characteristics of Pairs of Vector Spaces,” *Proc. Cambridge Phil. Soc.*, Vol. 53, pp. 800–816, 1957. (Cited on page 210.)
- [7] R. Agaev and P. Chebotarev, “On the Spectra of Nonsymmetric Laplacian Matrices,” *Lin. Alg. Appl.*, Vol. 399, pp. 157–168, 2005. (Cited on page 708.)
- [8] C. D. Ahlbrandt and A. C. Peterson, *Discrete Hamiltonian Systems: Difference Equations, Continued Fractions, and Riccati Equations*. Dordrecht: Kluwer, 1996. (Cited on page 805.)
- [9] E. Ahn, S. Kim, and Y. Lim, “An Extended Lie-Trotter Formula and Its Application,” *Lin. Alg. Appl.*, Vol. 427, pp. 190–196, 2007. (Cited on page 683.)
- [10] A. C. Aitken, *Determinants and Matrices*, 9th ed. Edinburgh: Oliver and Boyd, 1956. (Cited on page xix.)
- [11] M. Aivazis, *Group Theory in Physics—Problems and Solutions*. Singapore: World Scientific, 1991. (Cited on page 190.)
- [12] M. Al-Ahmar, “An Identity of Jacobi,” *Amer. Math. Monthly*, Vol. 103, pp. 78–79, 1996. (Cited on page 196.)
- [13] A. A. Albert and B. Muckenhoupt, “On Matrices of Trace Zero,” *Michigan Math. J.*, Vol. 4, pp. 1–3, 1957. (Cited on page 313.)
- [14] A. E. Albert, “Conditions for Positive and Nonnegative Definiteness in Terms of Pseudoinverses,” *SIAM J. Appl. Math.*, Vol. 17, pp. 434–440, 1969. (Cited on page 541.)

- [15] A. E. Albert, *Regression and the Moore-Penrose Pseudoinverse*. New York: Academic Press, 1972. (Cited on pages 380, 382, 385, 390, and 618.)
- [16] R. Aldrovandi, *Special Matrices of Mathematical Physics: Stochastic, Circulant and Bell Matrices*. Singapore: World Scientific, 2001. (Cited on pages xvii, 275, and 357.)
- [17] M. Aleksiejczuk and A. Smoktunowicz, "On Properties of Quadratic Matrices," *Mathematica Pannonica*, Vol. 112, pp. 239–248, 2000. (Cited on page 391.)
- [18] A. Y. Alfakih, "On the Nullspace, the Rangespace and the Characteristic Polynomial of Euclidean Distance Matrices," *Lin. Alg. Appl.*, Vol. 416, pp. 348–354, 2006. (Cited on page 573.)
- [19] M. Alic, P. S. Bullen, J. E. Pecaric, and V. Volenec, "On the Geometric-Arithmetic Mean Inequality for Matrices," *Mathematical Communications*, Vol. 2, pp. 125–128, 1997. (Cited on page 53.)
- [20] M. Alic, B. Mond, J. E. Pecaric, and V. Volenec, "Bounds for the Differences of Matrix Means," *SIAM J. Matrix Anal. Appl.*, Vol. 18, pp. 119–123, 1997. (Cited on page 463.)
- [21] H. Aling and J. M. Schumacher, "A Nine-Fold Decomposition for Linear Systems," *Int. J. Contr.*, Vol. 39, pp. 779–805, 1984. (Cited on page 805.)
- [22] G. Alpargu and G. P. H. Styan, "Some Remarks and a Bibliography on the Kantorovich Inequality," in *Multidimensional Statistical Analysis and Theory of Random Matrices*, A. K. Gupta and V. L. Girko, Eds. Utrecht: VSP, 1996, pp. 1–13. (Cited on page 501.)
- [23] R. C. Alperin, "The Matrix of a Rotation," *College Math. J.*, Vol. 20, p. 230, 1989. (Cited on page 194.)
- [24] C. Alsina and R. Ger, "On Some Inequalities and Stability Results Related to the Exponential Function," *J. Ineq. Appl.*, Vol. 2, pp. 373–380, 1998. (Cited on page 34.)
- [25] C. Alsina and R. B. Nelsen, "On Candido's Identity," *Math. Mag.*, Vol. 80, pp. 226–228, 2007. (Cited on page 32.)
- [26] S. L. Altmann, *Rotations, Quaternions, and Double Groups*. New York: Oxford University Press, 1986. (Cited on pages xvii, 226, and 228.)
- [27] S. L. Altmann, "Hamilton, Rodrigues, and the Quaternion Scandal," *Math. Mag.*, Vol. 62, pp. 291–308, 1989. (Cited on pages 193 and 194.)
- [28] H. Alzer, "A Lower Bound for the Difference between the Arithmetic and Geometric Means," *Nieuw. Arch. Wisk.*, Vol. 8, pp. 195–197, 1990. (Cited on page 52.)
- [29] S. Amghibech, "Problem 11296," *Amer. Math. Monthly*, Vol. 114, p. 452, 2007. (Cited on page 487.)
- [30] B. D. O. Anderson, "Orthogonal Decomposition Defined by a Pair of Skew-Symmetric Forms," *Lin. Alg. Appl.*, Vol. 8, pp. 91–93, 1974. (Cited on pages xvii and 343.)
- [31] B. D. O. Anderson, "Weighted Hankel-Norm Approximation: Calculation of Bounds," *Sys. Contr. Lett.*, Vol. 7, pp. 247–255, 1986. (Cited on page 502.)

- [32] B. D. O. Anderson, E. I. Jury, and M. Mansour, "Schwarz Matrix Properties for Continuous and Discrete Time Systems," *Int. J. Contr.*, Vol. 23, pp. 1–16, 1976. (Cited on page 704.)
- [33] B. D. O. Anderson and J. B. Moore, "A Matrix Kronecker Lemma," *Lin. Alg. Appl.*, Vol. 15, pp. 227–234, 1976. (Cited on page 465.)
- [34] G. Anderson, M. Vamanamurthy, and M. Vuorinen, "Monotonicity Rules in Calculus," *Amer. Math. Monthly*, Vol. 113, pp. 805–816, 2006. (Cited on pages 23, 27, and 29.)
- [35] T. W. Anderson and I. Olkin, "An Extremal Problem for Positive Definite Matrices," *Lin. Multilin. Alg.*, Vol. 6, pp. 257–262, 1978. (Cited on page 459.)
- [36] W. N. Anderson, "Shorted Operators," *SIAM J. Appl. Math.*, Vol. 20, pp. 520–525, 1971. (Cited on pages 529 and 530.)
- [37] W. N. Anderson and R. J. Duffin, "Series and Parallel Addition of Matrices," *J. Math. Anal. Appl.*, Vol. 26, pp. 576–594, 1969. (Cited on page 529.)
- [38] W. N. Anderson, E. J. Harner, and G. E. Trapp, "Eigenvalues of the Difference and Product of Projections," *Lin. Multilin. Alg.*, Vol. 17, pp. 295–299, 1985. (Cited on page 335.)
- [39] W. N. Anderson and M. Schreiber, "On the Infimum of Two Projections," *Acta Sci. Math.*, Vol. 33, pp. 165–168, 1972. (Cited on pages 384, 385, and 460.)
- [40] W. N. Anderson and G. E. Trapp, "Shorted Operators II," *SIAM J. Appl. Math.*, Vol. 28, pp. 60–71, 1975. (Cited on pages 443, 529, 530, and 541.)
- [41] W. N. Anderson and G. E. Trapp, "Inverse Problems for Means of Matrices," *SIAM J. Alg. Disc. Meth.*, Vol. 7, pp. 188–192, 1986. (Cited on page 463.)
- [42] W. N. Anderson and G. E. Trapp, "Symmetric Positive Definite Matrices," *Amer. Math. Monthly*, Vol. 95, pp. 261–262, 1988. (Cited on page 475.)
- [43] T. Ando, "Concavity of Certain Maps on Positive Definite Matrices and Applications to Hadamard Products," *Lin. Alg. Appl.*, Vol. 26, pp. 203–241, 1979. (Cited on pages 439, 536, 537, 540, 542, and 685.)
- [44] T. Ando, "Inequalities for M-Matrices," *Lin. Multilin. Alg.*, Vol. 8, pp. 291–316, 1980. (Cited on page 535.)
- [45] T. Ando, "On the Arithmetic-Geometric-Harmonic-Mean Inequalities for Positive Definite Matrices," *Lin. Alg. Appl.*, Vol. 52/53, pp. 31–37, 1983. (Cited on page 463.)
- [46] T. Ando, "On Some Operator Inequalities," *Math. Ann.*, Vol. 279, pp. 157–159, 1987. (Cited on pages 463 and 464.)
- [47] T. Ando, "Majorizations and Inequalities in Matrix Theory," *Lin. Alg. Appl.*, Vol. 199, pp. 17–67, 1994. (Cited on pages 472, 490, 583, 584, 588, 600, 612, 689, and 694.)
- [48] T. Ando, "Majorization Relations for Hadamard Products," *Lin. Alg. Appl.*, Vol. 223–224, pp. 57–64, 1995. (Cited on page 535.)
- [49] T. Ando, "Matrix Young Inequalities," *Oper. Theory Adv. Appl.*, Vol. 75, pp. 33–38, 1995. (Cited on page 612.)

- [50] T. Ando, "Problem of Infimum in the Positive Cone," in *Analytic and Geometric Inequalities and Applications*, T. M. Rassias and H. M. Srivastava, Eds. Dordrecht: Kluwer, 1999, pp. 1–12. (Cited on pages 460 and 530.)
- [51] T. Ando, "Lowner Inequality of Indefinite Type," *Lin. Alg. Appl.*, Vol. 385, pp. 73–80, 2004. (Cited on pages 457 and 795.)
- [52] T. Ando and R. Bhatia, "Eigenvalue Inequalities Associated with the Cartesian Decomposition," *Lin. Multilin. Alg.*, Vol. 22, pp. 133–147, 1987. (Cited on page 515.)
- [53] T. Ando and F. Hiai, "Log-Majorization and Complementary Golden-Thompson Type Inequalities," *Lin. Alg. Appl.*, Vol. 197/198, pp. 113–131, 1994. (Cited on pages 465, 683, 686, 693, and 694.)
- [54] T. Ando and F. Hiai, "Hölder Type Inequalities for Matrices," *Math. Ineq. Appl.*, Vol. 1, pp. 1–30, 1998. (Cited on pages 442, 460, and 461.)
- [55] T. Ando, F. Hiai, and K. Okubo, "Trace Inequalities for Multiple Products of Two Matrices," *Math. Ineq. Appl.*, Vol. 3, pp. 307–318, 2000. (Cited on pages 477 and 478.)
- [56] T. Ando, R. A. Horn, and C. R. Johnson, "The Singular Values of a Hadamard Product: A Basic Inequality," *Lin. Multilin. Alg.*, Vol. 21, pp. 345–365, 1987. (Cited on page 615.)
- [57] T. Ando, C.-K. Li, and R. Mathias, "Geometric Means," *Lin. Alg. Appl.*, Vol. 385, pp. 305–334, 2004. (Cited on page 463.)
- [58] T. Ando and X. Zhan, "Norm Inequalities Related to Operator Monotone Functions," *Math. Ann.*, Vol. 315, pp. 771–780, 1999. (Cited on pages 514, 590, and 695.)
- [59] T. Andreescu and D. Andrica, *Complex Numbers from A to Z*. Boston: Birkhauser, 2006. (Cited on pages 70, 71, 155, 156, 158, 159, and 160.)
- [60] T. Andreescu and Z. Feng, *103 Trigonometry Problems*. Boston: Birkhauser, 2004. (Cited on page 159.)
- [61] E. Andruchow, G. Corach, and D. Stojanoff, "Geometric Operator Inequalities," *Lin. Alg. Appl.*, Vol. 258, pp. 295–310, 1997. (Cited on pages 576 and 589.)
- [62] E. Angel, *Interactive Computer Graphics*, 3rd ed. Reading: Addison-Wesley, 2002. (Cited on pages xvii and 194.)
- [63] Anonymous, "Like the Carlson's inequality," <http://www.mathlinks.ro/viewtopic.php?t=151210>. (Cited on page 42.)
- [64] A. C. Antoulas and D. C. Sorensen, "Lyapunov, Lanczos, and Inertia," *Lin. Alg. Appl.*, Vol. 326, pp. 137–150, 2001. (Cited on page 793.)
- [65] J. D. Aplevich, *Implicit Linear Systems*. Berlin: Springer, 1991. (Cited on pages xvii and 304.)
- [66] T. Apostol, *Introduction to Analytic Number Theory*. New York: Springer, 1998. (Cited on page 447.)
- [67] T. M. Apostol, "Some Explicit Formulas for the Exponential Matrix," *Amer. Math. Monthly*, Vol. 76, pp. 289–292, 1969. (Cited on page 673.)
- [68] T. M. Apostol, Ed., *Mathematical Analysis*, 2nd ed. Reading: Addison Wesley, 1974. (Cited on pages xxxiv, 13, 62, 67, 632, 634, and 635.)

- [69] H. Araki, "On an Inequality of Lieb and Thirring," *Lett. Math. Phys.*, Vol. 19, pp. 167–170, 1990. (Cited on pages 479 and 583.)
- [70] H. Araki and S. Yamagami, "An Inequality for Hilbert-Schmidt Norm," *Comm. Math. Phys.*, Vol. 81, pp. 89–96, 1981. (Cited on page 584.)
- [71] J. Araujo and J. D. Mitchell, "An Elementary Proof that Every Singular Matrix Is a Product of Idempotent Matrices," *Amer. Math. Monthly*, Vol. 112, pp. 641–645, 2005. (Cited on page 350.)
- [72] A. Arimoto, "A Simple Proof of the Classification of Normal Toeplitz Matrices," *Elec. J. Lin. Alg.*, Vol. 9, pp. 108–111, 2002. (Cited on page 357.)
- [73] T. Arponen, "A Matrix Approach to Polynomials," *Lin. Alg. Appl.*, Vol. 359, pp. 181–196, 2003. (Cited on page 672.)
- [74] V. Arsigny, P. Fillard, X. Pennec, and N. Ayache, "Geometric Means in a Novel Vector Space Structure on Symmetric Positive-Definite Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 29, pp. 328–347, 2007. (Cited on page 686.)
- [75] M. Artin, *Algebra*. Englewood Cliffs: Prentice-Hall, 1991. (Cited on pages 223 and 229.)
- [76] M. Artzrouni, "A Theorem on Products of Matrices," *Lin. Alg. Appl.*, Vol. 49, pp. 153–159, 1983. (Cited on page 715.)
- [77] A. Arvanitoyeorgos, *An Introduction to Lie Groups and the Geometry of Homogeneous Spaces*. Providence: American Mathematical Society, 2003. (Cited on page 229.)
- [78] H. Aslaksen, "SO(2) Invariants of a Set of 2×2 Matrices," *Math. Scand.*, Vol. 65, pp. 59–66, 1989. (Cited on pages 260 and 261.)
- [79] H. Aslaksen, "Laws of Trigonometry on SU(3)," *Trans. Amer. Math. Soc.*, Vol. 317, pp. 127–142, 1990. (Cited on pages 152, 183, and 261.)
- [80] H. Aslaksen, "Quaternionic Determinants," *Mathematical Intelligencer*, Vol. 18, no. 3, pp. 57–65, 1996. (Cited on pages 226 and 229.)
- [81] H. Aslaksen, "Defining Relations of Invariants of Two 3×3 Matrices," *J. Algebra*, Vol. 298, pp. 41–57, 2006. (Cited on page 261.)
- [82] B. A. Asner, "On the Total Nonnegativity of the Hurwitz Matrix," *SIAM J. Appl. Math.*, Vol. 18, pp. 407–414, 1970. (Cited on pages 697 and 702.)
- [83] Y.-H. Au-Yeung, "A note on Some Theorems on Simultaneous Diagonalization of Two Hermitian Matrices," *Proc. Cambridge Phil. Soc.*, Vol. 70, pp. 383–386, 1971. (Cited on pages 504 and 507.)
- [84] Y.-H. Au-Yeung, "Some Inequalities for the Rational Power of a Nonnegative Definite Matrix," *Lin. Alg. Appl.*, Vol. 7, pp. 347–350, 1973. (Cited on page 458.)
- [85] Y.-H. Au-Yeung, "On the Semi-Definiteness of the Real Pencil of Two Hermitian Matrices," *Lin. Alg. Appl.*, Vol. 10, pp. 71–76, 1975. (Cited on page 362.)
- [86] K. M. R. Audenaert, "A Norm Compression Inequality for Block Partitioned Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 413, pp. 155–176, 2006. (Cited on page 596.)

- [87] K. M. R. Audenaert, "On a Norm Compression Inequality for $2 \times N$ Partitioned Block Matrices," *Lin. Alg. Appl.*, Vol. 428, pp. 781–795, 2008. (Cited on page 596.)
- [88] K. M. R. Audenaert, "On the Araki-Lieb-Thirring Inequality," *Int. J. Inform. Sys. Sci.*, Vol. 4, pp. 78–83, 2008. (Cited on pages 479 and 480.)
- [89] J. S. Aujla, "Some Norm Inequalities for Completely Monotone Functions," *SIAM J. Matrix Anal. Appl.*, Vol. 22, pp. 569–573, 2000. (Cited on page 514.)
- [90] J. S. Aujla and J.-C. Bourin, "Eigenvalue Inequalities for Convex and Log-Convex Functions," *Lin. Alg. Appl.*, Vol. 424, pp. 25–35, 2007. (Cited on pages 458, 521, 537, 583, and 592.)
- [91] J. S. Aujla and F. C. Silva, "Weak Majorization Inequalities and Convex Functions," *Lin. Alg. Appl.*, Vol. 369, pp. 217–233, 2003. (Cited on page 514.)
- [92] J. S. Aujla and H. L. Vasudeva, "Inequalities Involving Hadamard Product and Operator Means," *Math. Japonica*, Vol. 42, pp. 265–272, 1995. (Cited on pages 533, 539, and 541.)
- [93] B. Aupetit and J. Zemanek, "A Characterization of Normal Matrices by Their Exponentials," *Lin. Alg. Appl.*, Vol. 132, pp. 119–121, 1990. (Cited on page 695.)
- [94] M. Avriel, *Nonlinear Programming: Analysis and Methods*. Englewood Cliffs: Prentice-Hall, 1976, reprinted by Dover, Mineola, 2003. (Cited on page 505.)
- [95] O. Axelsson, *Iterative Solution Methods*. Cambridge: Cambridge University Press, 1994. (Cited on page xviii.)
- [96] L. E. Azar, "On Some Extensions of Hardy-Hilbert's Inequality and Applications," *J. Ineq. Appl.*, pp. 1–14, 2008, article ID 546829. (Cited on page 63.)
- [97] J. C. Baez, "Symplectic, Quaternionic, Fermionic," <http://math.ucr.edu/home/baez/symplectic.html>. (Cited on pages 222 and 227.)
- [98] J. C. Baez, "The Octonions," *Bull. Amer. Math. Soc.*, Vol. 39, pp. 145–205, 2001. (Cited on page 227.)
- [99] O. Bagdasar, *Inequalities and Applications*. Cluj-Napoca: Babes-Bolyai University, 2006, Bachelor's Degree Thesis, <http://rgmia.vu.edu.au/monographs/index.html>. (Cited on pages 22, 37, 42, and 54.)
- [100] Z. Bai and G. Golub, "Bounds for the Trace of the Inverse and the Determinant of Symmetric Positive Definite Matrices," *Ann. Numer. Math.*, Vol. 4, pp. 29–38, 1997. (Cited on page 476.)
- [101] D. W. Bailey and D. E. Crabtree, "Bounds for Determinants," *Lin. Alg. Appl.*, Vol. 2, pp. 303–309, 1969. (Cited on page 270.)
- [102] H. Bailey and Y. A. Rubinstein, "A Variety of Triangle Inequalities," *College Math. J.*, Vol. 31, pp. 350–355, 2000. (Cited on page 158.)
- [103] A. Baker, *Matrix Groups: An Introduction to Lie Group Theory*. New York: Springer, 2001. (Cited on pages 184, 217, 226, 229, 347, 660, and 682.)
- [104] J. K. Baksalary and O. M. Baksalary, "Nonsingularity of Linear Combinations of Idempotent Matrices," *Lin. Alg. Appl.*, Vol. 388, pp. 25–29, 2004. (Cited on page 203.)

- [105] J. K. Baksalary and O. M. Baksalary, "Particular Formulae for the Moore-Penrose Inverse of a Columnwise Partitioned Matrix," *Lin. Alg. Appl.*, Vol. 421, pp. 16–23, 2007. (Cited on page 391.)
- [106] J. K. Baksalary, O. M. Baksalary, and X. Liu, "Further Relationships between Certain Partial Orders of Matrices and their Squares," *Lin. Alg. Appl.*, Vol. 375, pp. 171–180, 2003. (Cited on page 120.)
- [107] J. K. Baksalary, O. M. Baksalary, and T. Szulc, "Properties of Schur Complements in Partitioned Idempotent Matrices," *Lin. Alg. Appl.*, Vol. 397, pp. 303–318, 2004. (Cited on page 387.)
- [108] J. K. Baksalary, O. M. Baksalary, and G. Trenkler, "A Revisitation of Formulae for the Moore-Penrose Inverse of Modified Matrices," *Lin. Alg. Appl.*, Vol. 372, pp. 207–224, 2003. (Cited on page 378.)
- [109] J. K. Baksalary, K. Nordstrom, and G. P. H. Styan, "Lowner-Ordering Antitonicity of Generalized Inverses of Hermitian Matrices," *Lin. Alg. Appl.*, Vol. 127, pp. 171–182, 1990. (Cited on pages 310, 457, and 527.)
- [110] J. K. Baksalary and F. Pukelsheim, "On the Lowner, Minus, and Star Partial Orderings of Nonnegative Definite Matrices," *Lin. Alg. Appl.*, Vol. 151, pp. 135–141, 1990. (Cited on pages 467, 523, and 526.)
- [111] J. K. Baksalary, F. Pukelsheim, and G. P. H. Styan, "Some Properties of Matrix Partial Orderings," *Lin. Alg. Appl.*, Vol. 119, pp. 57–85, 1989. (Cited on pages 120, 527, and 536.)
- [112] J. K. Baksalary and G. P. H. Styan, "Generalized Inverses of Bordered Matrices," *Lin. Alg. Appl.*, Vol. 354, pp. 41–47, 2002. (Cited on page 391.)
- [113] O. M. Baksalary, A. Mickiewicz, and G. Trenkler, "Rank of a Nonnegative Definite Matrix," *IMAGE*, Vol. 39, pp. 27–28, 2007. (Cited on page 525.)
- [114] O. M. Baksalary and G. Trenkler, "Rank of a Generalized Projector," *IMAGE*, Vol. 39, pp. 25–27, 2007. (Cited on page 178.)
- [115] O. M. Baksalary and G. Trenkler, "Characterizations of EP, Normal, and Hermitian Matrices," *Lin. Multilin. Alg.*, Vol. 56, pp. 299–304, 2008. (Cited on pages 180, 181, 315, 373, 396, 397, and 398.)
- [116] K. Ball, E. Carlen, and E. Lieb, "Sharp Uniform Convexity and Smoothness Inequalities for Trace Norms," *Invent. Math.*, Vol. 115, pp. 463–482, 1994. (Cited on pages 570 and 586.)
- [117] C. S. Ballantine, "Products of Positive Semidefinite Matrices," *Pac. J. Math.*, Vol. 23, pp. 427–433, 1967. (Cited on pages 350 and 542.)
- [118] C. S. Ballantine, "Products of Positive Definite Matrices II," *Pac. J. Math.*, Vol. 24, pp. 7–17, 1968. (Cited on page 542.)
- [119] C. S. Ballantine, "Products of Positive Definite Matrices III," *J. Algebra*, Vol. 10, pp. 174–182, 1968. (Cited on page 542.)
- [120] C. S. Ballantine, "A Note on the Matrix Equation $H = AP + PA^*$," *Lin. Alg. Appl.*, Vol. 2, pp. 37–47, 1969. (Cited on pages 314, 795, and 796.)
- [121] C. S. Ballantine, "Products of Positive Definite Matrices IV," *Lin. Alg. Appl.*, Vol. 3, pp. 79–114, 1970. (Cited on page 542.)

- [122] C. S. Ballantine, "Products of EP Matrices," *Lin. Alg. Appl.*, Vol. 12, pp. 257–267, 1975. (Cited on page 178.)
- [123] C. S. Ballantine, "Some Involutory Similarities," *Lin. Multilin. Alg.*, Vol. 3, pp. 19–23, 1975. (Cited on page 350.)
- [124] C. S. Ballantine, "Products of Involutory Matrices I," *Lin. Multilin. Alg.*, Vol. 5, pp. 53–62, 1977. (Cited on page 351.)
- [125] C. S. Ballantine, "Products of Idempotent Matrices," *Lin. Alg. Appl.*, Vol. 19, pp. 81–86, 1978. (Cited on page 350.)
- [126] C. S. Ballantine and C. R. Johnson, "Accretive Matrix Products," *Lin. Multilin. Alg.*, Vol. 3, pp. 169–185, 1975. (Cited on page 701.)
- [127] S. Banerjee, "Revisiting Spherical Trigonometry with Orthogonal Projectors," *College Math. J.*, Vol. 35, pp. 375–381, 2004. (Cited on page 161.)
- [128] J. Bang-Jensen and G. Gutin, *Digraphs: Theory, Algorithms and Applications*. New York: Springer, 2001. (Cited on page xvii.)
- [129] W. Bani-Domi and F. Kittaneh, "Norm Equalities and Inequalities for Operator Matrices," *Lin. Alg. Appl.*, Vol. 429, pp. 57–67, 2008. (Cited on pages 594 and 595.)
- [130] R. B. Bapat, "Two Inequalities for the Perron Root," *Lin. Alg. Appl.*, Vol. 85, pp. 241–248, 1987. (Cited on pages 66 and 280.)
- [131] R. B. Bapat, "On Generalized Inverses of Banded Matrices," *Elec. J. Lin. Alg.*, Vol. 16, pp. 284–290, 2007. (Cited on page 218.)
- [132] R. B. Bapat and M. K. Kwong, "A Generalization of $A \circ A^{-1} \geq I$," *Lin. Alg. Appl.*, Vol. 93, pp. 107–112, 1987. (Cited on page 533.)
- [133] R. B. Bapat and T. E. S. Raghavan, *Nonnegative Matrices and Applications*. Cambridge: Cambridge University Press, 1997. (Cited on page 275.)
- [134] R. B. Bapat and B. Zheng, "Generalized Inverses of Bordered Matrices," *Elec. J. Lin. Alg.*, Vol. 10, pp. 16–30, 2003. (Cited on pages 124 and 391.)
- [135] I. Y. Bar-Itzhack, D. Hershkowitz, and L. Rodman, "Pointing in Real Euclidean Space," *J. Guid. Contr. Dyn.*, Vol. 20, pp. 916–922, 1997. (Cited on pages 191 and 674.)
- [136] E. J. Barbeau, *Polynomials*. New York: Springer, 1989. (Cited on pages 31, 39, 42, 47, 158, 708, and 710.)
- [137] E. R. Barnes and A. J. Hoffman, "On Bounds for Eigenvalues of Real Symmetric Matrices," *Lin. Alg. Appl.*, Vol. 40, pp. 217–223, 1981. (Cited on page 271.)
- [138] S. Barnett, "A Note on the Bezoutian Matrix," *SIAM J. Appl. Math.*, Vol. 22, pp. 84–86, 1972. (Cited on page 257.)
- [139] S. Barnett, "Congenial Matrices," *Lin. Alg. Appl.*, Vol. 41, pp. 277–298, 1981. (Cited on pages 354, 361, and 362.)
- [140] S. Barnett, "Inversion of Partitioned Matrices with Patterned Blocks," *Int. J. Sys. Sci.*, Vol. 14, pp. 235–237, 1983. (Cited on page 215.)
- [141] S. Barnett, *Polynomials and Linear Control Systems*. New York: Marcel Dekker, 1983. (Cited on pages 362 and 798.)

- [142] S. Barnett, *Matrices in Control Theory*, revised ed. Malabar: Krieger, 1984. (Cited on page xvii.)
- [143] S. Barnett, "Leverrier's Algorithm: A New Proof and Extensions," *SIAM J. Matrix Anal. Appl.*, Vol. 10, pp. 551–556, 1989. (Cited on page 281.)
- [144] S. Barnett, *Matrices: Methods and Applications*. Oxford: Clarendon, 1990. (Cited on pages 146, 353, 362, 377, 393, 454, and 618.)
- [145] S. Barnett and P. Lancaster, "Some Properties of the Bezoutian for Polynomial Matrices," *Lin. Multilin. Alg.*, Vol. 9, pp. 99–110, 1980. (Cited on page 257.)
- [146] S. Barnett and C. Storey, *Matrix Methods in Stability Theory*. New York: Barnes and Noble, 1970. (Cited on pages xvii, 135, 361, 450, and 702.)
- [147] W. Barrett, "A Theorem on Inverses of Tridiagonal Matrices," *Lin. Alg. Appl.*, Vol. 27, pp. 211–217, 1979. (Cited on page 220.)
- [148] W. Barrett, "Hermitian and Positive Definite Matrices," in *Handbook of Linear Algebra*, L. Hogben, Ed. Boca Raton: Chapman & Hall/CRC, 2007, pp. 8–1–8–12. (Cited on page 538.)
- [149] W. Barrett, C. R. Johnson, and P. Tarazaga, "The Real Positive Definite Completion Problem for a Simple Cycle," *Lin. Alg. Appl.*, Vol. 192, pp. 3–31, 1993. (Cited on page 449.)
- [150] J. Barria and P. R. Halmos, "Vector Bases for Two Commuting Matrices," *Lin. Multilin. Alg.*, Vol. 27, pp. 147–157, 1990. (Cited on page 319.)
- [151] H. Bart, I. Gohberg, M. A. Kaashoek, and A. C. M. Ran, "Schur Complements and State Space Realizations," *Lin. Alg. Appl.*, Vol. 399, pp. 203–224, 2005. (Cited on page 800.)
- [152] H. Baruh, *Analytical Dynamics*. Boston: McGraw-Hill, 1999. (Cited on pages 193, 226, and 676.)
- [153] A. Barvinok, *A Course in Convexity*. Providence: American Mathematical Society, 2002. (Cited on pages 43, 47, 110, 111, 115, 498, 511, and 634.)
- [154] S. Barza, J. Pecaric, and L.-E. Persson, "Carlson Type Inequalities," *J. Ineq. Appl.*, Vol. 2, pp. 121–135, 1998. (Cited on page 58.)
- [155] F. L. Bauer, J. Stoer, and C. Witzgall, "Absolute and Monotonic Norms," *Numer. Math.*, Vol. 3, pp. 257–264, 1961. (Cited on page 619.)
- [156] D. S. Bayard, "An Optimization Result with Application to Optimal Spacecraft Reaction Wheel Orientation Design," in *Proc. Amer. Contr. Conf.*, Arlington, VA, June 2001, pp. 1473–1478. (Cited on page 334.)
- [157] M. S. Bazaraa, H. D. Sherali, and C. M. Shetty, *Nonlinear Programming*, 2nd ed. New York: Wiley, 1993. (Cited on pages 279, 624, and 626.)
- [158] N. Bebiano and J. da Providencia, "On the Determinant of Certain Strictly Dissipative Matrices," *Lin. Alg. Appl.*, Vol. 83, pp. 117–128, 1986. (Cited on page 486.)
- [159] N. Bebiano, J. da Providencia, and R. Lemos, "Matrix Inequalities in Statistical Mechanics," *Lin. Alg. Appl.*, Vol. 376, pp. 265–273, 2004. (Cited on pages xvii, 686, 687, and 688.)

- [160] N. Bebiano, R. Lemos, and J. da Providencia, “Inequalities for Quantum Relative Entropy,” *Lin. Alg. Appl.*, Vol. 401, pp. 159–172, 2005. (Cited on pages xvii and 686.)
- [161] N. Bebiano and M. E. Miranda, “On a Recent Determinantal Inequality,” *Lin. Alg. Appl.*, Vol. 201, pp. 99–102, 1994. (Cited on page 335.)
- [162] E. F. Beckenbach and R. Bellman, *Inequalities*. Berlin: Springer, 1965. (Cited on pages 76 and 544.)
- [163] R. I. Becker, “Necessary and Sufficient Conditions for the Simultaneous Diagonalizability of Two Quadratic Forms,” *Lin. Alg. Appl.*, Vol. 30, pp. 129–139, 1980. (Cited on pages 362 and 504.)
- [164] W. Beckner, “Inequalities in Fourier Analysis,” *Ann. Math.*, Vol. 102, p. 159182, 1975. (Cited on page 586.)
- [165] L. W. Beineke and R. J. Wilson, Eds., *Topics in Algebraic Graph Theory*. Cambridge: Cambridge University Press, 2005. (Cited on page xvii.)
- [166] T. N. Bekjan, “On Joint Convexity of Trace Functions,” *Lin. Alg. Appl.*, Vol. 390, pp. 321–327, 2004. (Cited on page 500.)
- [167] P. A. Bekker, “The Positive Semidefiniteness of Partitioned Matrices,” *Lin. Alg. Appl.*, Vol. 111, pp. 261–278, 1988. (Cited on pages 483, 484, and 541.)
- [168] J. G. Belinfante, B. Kolman, and H. A. Smith, “An Introduction to Lie Groups and Lie Algebras with Applications,” *SIAM Rev.*, Vol. 8, pp. 11–46, 1966. (Cited on page 722.)
- [169] G. R. Belitskii and Y. I. Lyubich, *Matrix Norms and Their Applications*. Basel: Birkhauser, 1988. (Cited on page 619.)
- [170] R. Bellman, *Introduction to Matrix Analysis*, 2nd ed. New York: McGraw-Hill, 1960, reprinted by SIAM, Philadelphia, 1995. (Cited on pages 146, 280, 454, and 682.)
- [171] R. Bellman, “Some Inequalities for the Square Root of a Positive Definite Matrix,” *Lin. Alg. Appl.*, Vol. 1, pp. 321–324, 1968. (Cited on page 460.)
- [172] A. Ben-Israel, “A Note on Partitioned Matrices and Equations,” *SIAM Rev.*, Vol. 11, pp. 247–250, 1969. (Cited on page 391.)
- [173] A. Ben-Israel, “The Moore of the Moore-Penrose Inverse,” *Elect. J. Lin. Alg.*, Vol. 9, pp. 150–157, 2002. (Cited on page 398.)
- [174] A. Ben-Israel and T. N. E. Greville, *Generalized Inverses: Theory and Applications*, 2nd ed. New York: Springer, 2003. (Cited on pages 372, 376, 391, 394, 395, 398, and 507.)
- [175] B. Ben Taher and M. Rachidi, “Some Explicit Formulas for the Polynomial Decomposition of the Matrix Exponential and Applications,” *Lin. Alg. Appl.*, Vol. 350, pp. 171–184, 2002. (Cited on page 673.)
- [176] A. Ben-Tal and A. Nemirovski, *Lectures on Modern Convex Optimization*. Philadelphia: SIAM, 2001. (Cited on pages 110 and 164.)
- [177] A. T. Benjamin and J. J. Quinn, *Proofs That Really Count: The Art of Combinatorial Proof*. Washington, DC: Mathematical Association of America, 2003. (Cited on pages 11, 17, and 278.)

- [178] L. Berg, "Three Results in Connection with Inverse Matrices," *Lin. Alg. Appl.*, Vol. 84, pp. 63–77, 1986. (Cited on page 129.)
- [179] C. Berge, *The Theory of Graphs*. London: Methuen, 1962, reprinted by Dover, Mineola, 2001. (Cited on page xvii.)
- [180] L. D. Berkovitz, *Convexity and Optimization in $\mathbb{R}^n$* . New York: Wiley, 2002. (Cited on pages 164 and 635.)
- [181] A. Berman, M. Neumann, and R. J. Stern, *Nonnegative Matrices in Dynamic Systems*. New York: Wiley, 1989. (Cited on pages 275, 706, and 707.)
- [182] A. Berman and R. J. Plemmons, *Nonnegative Matrices in the Mathematical Sciences*. New York: Academic Press, 1979, reprinted by SIAM, Philadelphia, 1994. (Cited on pages 176, 230, 275, 277, 706, and 707.)
- [183] A. Berman, R. S. Varga, and R. C. Ward, "ALPS: Matrices with Nonpositive Off-Diagonal Entries," *Lin. Alg. Appl.*, Vol. 21, pp. 233–244, 1978. (Cited on page 706.)
- [184] D. S. Bernstein, "Inequalities for the Trace of Matrix Exponentials," *SIAM J. Matrix Anal. Appl.*, Vol. 9, pp. 156–158, 1988. (Cited on pages 689 and 692.)
- [185] D. S. Bernstein, "Some Open Problems in Matrix Theory Arising in Linear Systems and Control," *Lin. Alg. Appl.*, Vol. 162–164, pp. 409–432, 1992. (Cited on page 701.)
- [186] D. S. Bernstein and S. P. Bhat, "Lyapunov Stability, Semistability, and Asymptotic Stability of Matrix Second-Order Systems," *ASME Trans. J. Vibr. Acoustics*, Vol. 117, pp. 145–153, 1995. (Cited on pages 338, 705, and 722.)
- [187] D. S. Bernstein and S. P. Bhat, "Nonnegativity, Reducibility, and Semistability of Mass Action Kinetics," in *Proc. Conf. Dec. Contr.*, Phoenix, AZ, December 1999, pp. 2206–2211. (Cited on page 707.)
- [188] D. S. Bernstein and W. M. Haddad, "LQG Control with an H_∞ Performance Bound: A Riccati Equation Approach," *IEEE Trans. Autom. Contr.*, Vol. 34, pp. 293–305, 1989. (Cited on page 805.)
- [189] D. S. Bernstein, W. M. Haddad, D. C. Hyland, and F. Tyan, "Maximum Entropy-Type Lyapunov Functions for Robust Stability and Performance Analysis," *Sys. Contr. Lett.*, Vol. 21, pp. 73–87, 1993. (Cited on page 343.)
- [190] D. S. Bernstein and D. C. Hyland, "Compartmental Modeling and Second-Moment Analysis of State Space Systems," *SIAM J. Matrix Anal. Appl.*, Vol. 14, pp. 880–901, 1993. (Cited on pages 230, 706, and 708.)
- [191] D. S. Bernstein and W. So, "Some Explicit Formulas for the Matrix Exponential," *IEEE Trans. Autom. Contr.*, Vol. 38, pp. 1228–1232, 1993. (Cited on page 673.)
- [192] D. S. Bernstein and C. F. Van Loan, "Rational Matrix Functions and Rank-1 Updates," *SIAM J. Matrix Anal. Appl.*, Vol. 22, pp. 145–154, 2000. (Cited on page 127.)
- [193] K. V. Bhagwat and R. Subramanian, "Inequalities between Means of Positive Operators," *Math. Proc. Camb. Phil. Soc.*, Vol. 83, pp. 393–401, 1978. (Cited on pages 433, 459, and 683.)
- [194] S. P. Bhat and D. S. Bernstein, "Average-Preserving Symmetries and Energy Equipartition in Linear Hamiltonian Systems," preprint. (Cited on page 227.)

- [195] S. P. Bhat and D. S. Bernstein, "Nontangency-Based Lyapunov Tests for Convergence and Stability in Systems Having a Continuum of Equilibria," *SIAM J. Contr. Optim.*, Vol. 42, pp. 1745–1775, 2003. (Cited on pages 722 and 797.)
- [196] R. Bhatia, *Perturbation Bounds for Matrix Eigenvalues*. Essex: Longman Scientific and Technical, 1987. (Cited on page 600.)
- [197] R. Bhatia, *Matrix Analysis*. New York: Springer, 1997. (Cited on pages 163, 326, 327, 433, 436, 439, 441, 442, 443, 479, 480, 509, 510, 513, 520, 521, 541, 542, 561, 578, 580, 583, 585, 588, 589, 590, 600, 601, 605, 686, 693, and 695.)
- [198] R. Bhatia, "Linear Algebra to Quantum Cohomology: The Story of Alfred Horn's Inequalities," *Amer. Math. Monthly*, Vol. 108, pp. 289–318, 2001. (Cited on pages 511, 513, and 542.)
- [199] R. Bhatia, "Infinitely Divisible Matrices," *Amer. Math. Monthly*, Vol. 113, pp. 221–235, 2006. (Cited on pages 221, 446, 447, and 531.)
- [200] R. Bhatia, *Perturbation Bounds for Matrix Eigenvalues*. Philadelphia: SIAM, 2007. (Cited on pages 272, 518, 575, 600, and 601.)
- [201] R. Bhatia, *Positive Definite Matrices*. Princeton: Princeton University Press, 2007. (Cited on pages 441, 442, 445, 446, 447, 449, 457, 463, 686, and 688.)
- [202] R. Bhatia and C. Davis, "More Matrix Forms of the Arithmetic-Geometric Mean Inequality," *SIAM J. Matrix Anal. Appl.*, Vol. 14, pp. 132–136, 1993. (Cited on page 589.)
- [203] R. Bhatia and C. Davis, "A Cauchy-Schwarz Inequality for Operators with Applications," *Lin. Alg. Appl.*, Vol. 223/224, pp. 119–129, 1995. (Cited on pages 582 and 589.)
- [204] R. Bhatia and D. Drissi, "Generalized Lyapunov Equations and Positive Definite Functions," *SIAM J. Matrix Anal. Appl.*, Vol. 27, pp. 103–114, 2005. (Cited on page 448.)
- [205] R. Bhatia and L. Elsner, "Higher Order Logarithmic Derivatives of Matrices in the Spectral Norm," *SIAM J. Matrix Anal. Appl.*, Vol. 25, pp. 662–668, 2003. (Cited on page 691.)
- [206] R. Bhatia and J. Holbrook, "Frechet Derivatives of the Power Function," *Indiana University Math. J.*, Vol. 49, pp. 1155–1173, 2000. (Cited on pages 24 and 515.)
- [207] R. Bhatia and J. Holbrook, "Riemannian Geometry and Matrix Geometric Means," *Lin. Alg. Appl.*, Vol. 413, pp. 594–618, 2006. (Cited on page 688.)
- [208] R. Bhatia and F. Kittaneh, "Norm Inequalities for Partitioned Operators and an Application," *Math. Ann.*, Vol. 287, pp. 719–726, 1990. (Cited on page 594.)
- [209] R. Bhatia and F. Kittaneh, "On the Singular Values of a Product of Operators," *SIAM J. Matrix Anal. Appl.*, Vol. 11, pp. 272–277, 1990. (Cited on pages 458, 589, and 613.)
- [210] R. Bhatia and F. Kittaneh, "Norm Inequalities for Positive Operators," *Lett. Math. Phys.*, Vol. 43, pp. 225–231, 1998. (Cited on page 590.)
- [211] R. Bhatia and F. Kittaneh, "Cartesian Decompositions and Schatten Norms," *Lin. Alg. Appl.*, Vol. 318, pp. 109–116, 2000. (Cited on pages 586 and 587.)

- [212] R. Bhatia and F. Kittaneh, "Notes on Matrix Arithmetic-Geometric Mean Inequalities," *Lin. Alg. Appl.*, Vol. 308, pp. 203–211, 2000. (Cited on pages 582 and 583.)
- [213] R. Bhatia and F. Kittaneh, "Clarkson Inequalities with Several Operators," *Bull. London Math. Soc.*, Vol. 36, pp. 820–832, 2004. (Cited on page 586.)
- [214] R. Bhatia and F. Kittaneh, "Commutators, Pinchings, and Spectral Variation," *Oper. Matrices*, Vol. 2, pp. 143–151, 2008. (Cited on page 585.)
- [215] R. Bhatia and F. Kittaneh, "The Matrix Arithmetic-Geometric Mean Inequality," *Lin. Alg. Appl.*, Vol. 428, pp. 2177–2191, 2008. (Cited on pages 515, 580, and 590.)
- [216] R. Bhatia and K. R. Parthasarathy, "Positive Definite Functions and Operator Inequalities," *Bull. London Math. Soc.*, Vol. 32, pp. 214–228, 2000. (Cited on pages 446, 589, 590, and 695.)
- [217] R. Bhatia and P. Rosenthal, "How and Why to Solve the Operator Equation $AX - XB = Y$," *Bull. London Math. Soc.*, Vol. 29, pp. 1–21, 1997. (Cited on pages 320 and 335.)
- [218] R. Bhatia and P. Semrl, "Orthogonality of Matrices and Some Distance Problems," *Lin. Alg. Appl.*, Vol. 287, pp. 77–85, 1999. (Cited on page 570.)
- [219] R. Bhatia and X. Zhan, "Norm Inequalities for Operators with Positive Real Part," *J. Operator Theory*, Vol. 50, pp. 67–76, 2003. (Cited on pages 486 and 587.)
- [220] R. Bhattacharya and K. Mukherjea, "On Unitary Similarity of Matrices," *Lin. Alg. Appl.*, Vol. 126, pp. 95–105, 1989. (Cited on page 318.)
- [221] S. P. Bhattacharyya, H. Chapellat, and L. Keel, *Robust Control: The Parametric Approach*. Englewood Cliffs: Prentice-Hall, 1995. (Cited on page 697.)
- [222] M. R. Bicknell, "The Lambda Number of a Matrix: The Sum of Its n^2 Cofactors," *Amer. Math. Monthly*, Vol. 72, pp. 260–264, 1965. (Cited on page 142.)
- [223] N. Biggs, *Algebraic Graph Theory*, 2nd ed. Cambridge: Cambridge University Press, 2000. (Cited on pages xvii and 337.)
- [224] P. Binding, B. Najman, and Q. Ye, "A Variational Principle for Eigenvalues of Pencils of Hermitian Matrices," *Integ. Equ. Oper. Theory*, Vol. 35, pp. 398–422, 1999. (Cited on page 362.)
- [225] K. Binmore, *Fun and Games: A Text on Game Theory*. Lexington: D. C. Heath and Co., 1992. (Cited on page xvii.)
- [226] A. Bjorck, *Numerical Methods for Least Squares Problems*. Philadelphia: SIAM, 1996. (Cited on page 618.)
- [227] S. Blanes and F. Casas, "On the Convergence and Optimization of the Baker-Campbell-Hausdorff Formula," *Lin. Alg. Appl.*, Vol. 378, pp. 135–158, 2004. (Cited on page 683.)
- [228] S. Blanes, F. Casas, J. A. Oteo, and J. Ros, "Magnus and Fer Expansions for Matrix Differential Equations: The Convergence Problem," *J. Phys. A: Math. Gen.*, Vol. 31, pp. 259–268, 1998. (Cited on page 678.)
- [229] E. D. Bloch, *Proofs and Fundamentals: A First Course in Abstract Mathematics*. Boston: Birkhauser, 2000. (Cited on pages 10 and 76.)

- [230] V. Blondel and J. N. Tsitsiklis, "When Is a Pair of Matrices Mortal?" *Inform. Proc. Lett.*, Vol. 63, pp. 283–286, 1997. (Cited on page 715.)
- [231] V. Blondel and J. N. Tsitsiklis, "The Boundedness of All Products of a Pair of Matrices Is Undecidable," *Sys. Contr. Lett.*, Vol. 41, pp. 135–140, 2000. (Cited on page 715.)
- [232] L. M. Blumenthal, *Theory and Applications of Distance Geometry*. Oxford: Oxford University Press, 1953. (Cited on page 161.)
- [233] W. Boehm, "An Operator Limit," *SIAM Rev.*, Vol. 36, p. 659, 1994. (Cited on page 714.)
- [234] A. W. Bojanczyk and A. Lutoborski, "Computation of the Euler Angles of a Symmetric 3×3 Matrix," *SIAM J. Matrix Anal. Appl.*, Vol. 12, pp. 41–48, 1991. (Cited on page 265.)
- [235] B. Bollobas, *Modern Graph Theory*. New York: Springer, 1998. (Cited on page xvii.)
- [236] J. V. Bondar, "Comments on and Complements to *Inequalities: Theory of Majorization and Its Applications*," *Lin. Alg. Appl.*, Vol. 199, pp. 115–129, 1994. (Cited on pages 45 and 60.)
- [237] A. Borck, *Numerical Methods for Least Squares Problems*. Philadelphia: SIAM, 1996. (Cited on page 398.)
- [238] J. Borwein, D. Bailey, and R. Girgensohn, *Experimentation in Mathematics: Computational Paths to Discovery*. Natick: A K Peters, 2004. (Cited on pages 17, 18, 20, 21, 56, 161, and 275.)
- [239] J. M. Borwein and A. S. Lewis, *Convex Analysis and Nonlinear Optimization*. New York: Springer, 2000. (Cited on pages 110, 111, 164, 279, 333, 334, 441, 460, 633, and 635.)
- [240] A. J. Bosch, "The Factorization of a Square Matrix into Two Symmetric Matrices," *Amer. Math. Monthly*, Vol. 93, pp. 462–464, 1986. (Cited on pages 349 and 361.)
- [241] A. J. Bosch, "Note on the Factorization of a Square Matrix into Two Hermitian or Symmetric Matrices," *SIAM Rev.*, Vol. 29, pp. 463–468, 1987. (Cited on pages 349 and 361.)
- [242] A. Bottcher and D. Wenzel, "How Big Can the Commutator of Two Matrices Be and How Big Is it Typically?" *Lin. Alg. Appl.*, Vol. 403, pp. 216–228, 2005. (Cited on page 584.)
- [243] A. Bottcher and D. Wenzel, "The Frobenius Norm and the Commutator," *Lin. Alg. Appl.*, Vol. 429, pp. 1864–1885, 2008. (Cited on page 584.)
- [244] O. Bottema, Z. Djordjovic, R. R. Janic, D. S. Mitrinovic, and P. M. Vasic, *Inequalities: Theory of Majorization and Its Applications*. Groningen: Wolters-Noordhoff, 1969. (Cited on page 158.)
- [245] T. L. Boullion and P. L. Odell, *Generalized Inverse Matrices*. New York: Wiley, 1971. (Cited on page 398.)
- [246] J.-C. Bourin, "Some Inequalities for Norms on Matrices and Operators," *Lin. Alg. Appl.*, Vol. 292, pp. 139–154, 1999. (Cited on pages 480, 481, 576, and 608.)

- [247] J.-C. Bourin, "Convexity or Concavity Inequalities for Hermitian Operators," *Math. Ineq. Appl.*, Vol. 7, pp. 607–620, 2004. (Cited on page 482.)
- [248] J.-C. Bourin, "Hermitian Operators and Convex Functions," *J. Ineq. Pure Appl. Math.*, Vol. 6, no. 5, pp. 1–6, 2005, Article 139. (Cited on pages 482 and 483.)
- [249] J.-C. Bourin, "Reverse Inequality to Araki's Inequality Comparison of $A^p Z^p Z^p$ and $(AZA)^p$," *Math. Ineq. Appl.*, Vol. 232, pp. 373–378, 2005. (Cited on page 465.)
- [250] J.-C. Bourin, "Matrix Versions of Some Classical Inequalities," *Lin. Alg. Appl.*, Vol. 407, pp. 890–907, 2006. (Cited on pages 483 and 599.)
- [251] J.-C. Bourin, "Reverse Rearrangement Inequalities via Matrix Technics," *J. Ineq. Pure Appl. Math.*, Vol. 7, no. 2, pp. 1–6, 2006, Article 43. (Cited on pages 61 and 458.)
- [252] J.-C. Bourin and Y. Seo, "Reverse Inequality to Golden-Thompson Type Inequalities: Comparison of e^{A+B} and $e^A e^B$," *Lin. Alg. Appl.*, Vol. 426, pp. 312–316, 2007. (Cited on page 687.)
- [253] K. Bourque and S. Ligh, "Matrices Associated with Classes of Arithmetical Functions," *J. Number Theory*, Vol. 45, pp. 367–376, 1993. (Cited on page 447.)
- [254] S. Boyd, "Entropy and Random Feedback," in *Open Problems in Mathematical Systems and Control Theory*, V. D. Blondel, E. D. Sontag, M. Vidyasagar, and J. C. Willems, Eds. New York: Springer, 1998, pp. 71–74. (Cited on page 578.)
- [255] S. Boyd and L. Vandenberghe, *Convex Optimization*. Cambridge: Cambridge University Press, 2004. (Cited on pages xvii and 164.)
- [256] J. L. Brenner, "A Bound for a Determinant with Dominant Main Diagonal," *Proc. Amer. Math. Soc.*, Vol. 5, pp. 631–634, 1954. (Cited on page 270.)
- [257] J. L. Brenner, "Expanded Matrices from Matrices with Complex Elements," *SIAM Rev.*, Vol. 3, pp. 165–166, 1961. (Cited on pages 154 and 229.)
- [258] J. L. Brenner and J. S. Lim, "The Matrix Equations $A = XYZ$ and $B = ZYX$ and Related Ones," *Canad. Math. Bull.*, Vol. 17, pp. 179–183, 1974. (Cited on page 346.)
- [259] J. W. Brewer, "Kronecker Products and Matrix Calculus in System Theory," *IEEE Trans. Circ. Sys.*, Vol. CAS-25, pp. 772–781, 1978, correction: CAS-26, p. 360, 1979. (Cited on page 416.)
- [260] L. Brickman, "On the Field of Values of a Matrix," *Proc. Amer. Math. Soc.*, Vol. 12, pp. 61–66, 1961. (Cited on page 498.)
- [261] R. Brockett, *Finite Dimensional Linear Systems*. New York: Wiley, 1970. (Cited on page 805.)
- [262] R. W. Brockett, "Using Feedback to Improve System Identification," in *Control of Uncertain Systems*, B. A. Francis, M. C. Smith, and J. C. Willems, Eds. Berlin: Springer, 2006, pp. 45–65. (Cited on pages 511 and 797.)
- [263] G. Brown, "Convexity and Minkowski's Inequality," *Amer. Math. Monthly*, Vol. 112, pp. 740–742, 2005. (Cited on page 66.)
- [264] E. T. Browne, *Introduction to the Theory of Determinants and Matrices*. Chapel Hill: University of North Carolina Press, 1958. (Cited on page 541.)

- [265] R. Bru, J. J. Climent, and M. Neumann, "On the Index of Block Upper Triangular Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 16, pp. 436–447, 1995. (Cited on page 343.)
- [266] R. A. Brualdi, *Combinatorial Matrix Classes*. Cambridge: Cambridge University Press, 2006. (Cited on page xvii.)
- [267] R. A. Brualdi and J. Q. Massey, "Some Applications of Elementary Linear Algebra in Combinatorics," *College Math. J.*, Vol. 24, pp. 10–19, 1993. (Cited on page 131.)
- [268] R. A. Brualdi and S. Mellendorf, "Regions in the Complex Plane Containing the Eigenvalues of a Matrix," *Amer. Math. Monthly*, Vol. 101, pp. 975–985, 1994. (Cited on pages 269 and 704.)
- [269] R. A. Brualdi and H. J. Ryser, *Combinatorial Matrix Theory*. Cambridge: Cambridge University Press, 1991. (Cited on pages xvii, 126, 131, and 506.)
- [270] R. A. Brualdi and H. Schneider, "Determinantal Identities: Gauss, Schur, Cauchy, Sylvester, Kronecker, Jacobi, Binet, Laplace, Muir, and Cayley," *Lin. Alg. Appl.*, Vol. 52/53, pp. 769–791, 1983. (Cited on page 129.)
- [271] D. Buckholtz, "Inverting the Difference of Hilbert Space Projections," *Amer. Math. Monthly*, Vol. 104, pp. 60–61, 1997. (Cited on page 210.)
- [272] D. Buckholtz, "Hilbert Space Idempotents and Involutions," *Proc. Amer. Math. Soc.*, Vol. 128, pp. 1415–1418, 1999. (Cited on pages 336 and 337.)
- [273] P. S. Bullen, *A Dictionary of Inequalities*. Essex: Longman, 1998. (Cited on pages 29, 35, 42, 55, 57, 64, 76, 546, and 571.)
- [274] P. S. Bullen, *Handbook of Means and Their Inequalities*. Dordrecht: Kluwer, 2003. (Cited on pages 23, 25, 27, 29, and 76.)
- [275] P. S. Bullen, D. S. Mitrinovic, and P. M. Vasic, *Means and Their Inequalities*. Dordrecht: Reidel, 1988. (Cited on pages 38, 52, 76, and 567.)
- [276] A. Bultheel and M. Van Barel, *Linear Algebra, Rational Approximation and Orthogonal Polynomials*. Amsterdam: Elsevier, 1997. (Cited on page 281.)
- [277] F. Burns, D. Carlson, E. V. Haynsworth, and T. L. Markham, "Generalized Inverse Formulas Using the Schur-Complement," *SIAM J. Appl. Math.*, Vol. 26, pp. 254–259, 1974. (Cited on page 391.)
- [278] P. J. Bushell and G. B. Trustrum, "Trace Inequalities for Positive Definite Matrix Power Products," *Lin. Alg. Appl.*, Vol. 132, pp. 173–178, 1990. (Cited on page 519.)
- [279] N. D. Cahill, J. R. D'Errico, D. A. Narayan, and J. Y. Narayan, "Fibonacci Determinants," *College Math. J.*, Vol. 33, pp. 221–225, 2002. (Cited on page 278.)
- [280] B. E. Cain, "Inertia Theory," *Lin. Alg. Appl.*, Vol. 30, pp. 211–240, 1980. (Cited on pages 309 and 795.)
- [281] D. Callan, "When Is "Rank" Additive?" *College Math. J.*, Vol. 29, pp. 145–147, 1998. (Cited on page 118.)
- [282] S. L. Campbell, *Singular Systems*. London: Pitman, 1980. (Cited on page 398.)
- [283] S. L. Campbell and C. D. Meyer, *Generalized Inverses of Linear Transformations*. London: Pitman, 1979, reprinted by Dover, Mineola, 1991. (Cited on pages 377, 384, 391, 395, 398, and 722.)

- [284] S. L. Campbell and N. J. Rose, "Singular Perturbation of Autonomous Linear Systems," *SIAM J. Math. Anal.*, Vol. 10, pp. 542–551, 1979. (Cited on page 699.)
- [285] L. Cao and H. M. Schwartz, "A Decomposition Method for Positive Semidefinite Matrices and its Application to Recursive Parameter Estimation," *SIAM J. Matrix Anal. Appl.*, Vol. 22, pp. 1095–1111, 2001. (Cited on pages 418 and 457.)
- [286] E. A. Carlen and E. H. Lieb, "A Minkowski Type Trace Inequality and Strong Subadditivity of Quantum Entropy," *Amer. Math. Soc. Transl.*, Vol. 189, pp. 59–62, 1999. (Cited on page 441.)
- [287] D. Carlson, "Controllability, Inertia, and Stability for Tridiagonal Matrices," *Lin. Alg. Appl.*, Vol. 56, pp. 207–220, 1984. (Cited on page 702.)
- [288] D. Carlson, "What Are Schur Complements Anyway?" *Lin. Alg. Appl.*, Vol. 74, pp. 257–275, 1986. (Cited on page 542.)
- [289] D. Carlson, "On the Controllability of Matrix Pairs (A, K) with K Positive Semidefinite, II," *SIAM J. Matrix Anal. Appl.*, Vol. 15, pp. 129–133, 1994. (Cited on page 794.)
- [290] D. Carlson, E. V. Haynsworth, and T. L. Markham, "A Generalization of the Schur Complement by Means of the Moore-Penrose Inverse," *SIAM J. Appl. Math.*, Vol. 26, pp. 169–175, 1974. (Cited on pages 387, 442, and 542.)
- [291] D. Carlson and R. Hill, "Generalized Controllability and Inertia Theory," *Lin. Alg. Appl.*, Vol. 15, pp. 177–187, 1976. (Cited on page 795.)
- [292] D. Carlson, C. R. Johnson, D. C. Lay, and A. D. Porter, Eds., *Linear Algebra Gems: Assets for Undergraduate Mathematics*. Washington, DC: Mathematical Association of America, 2002. (Cited on page 164.)
- [293] D. Carlson, C. R. Johnson, D. C. Lay, A. D. Porter, A. E. Watkins, and W. Watkins, Eds., *Resources for Teaching Linear Algebra*. Washington, DC: Mathematical Association of America, 1997. (Cited on page 164.)
- [294] P. Cartier, "Mathemagics, A Tribute to L. Euler and R. Feynman," in *Noise, Oscillators and Algebraic Randomness*, M. Planat, Ed. New York: Springer, 2000, pp. 6–67. (Cited on page 722.)
- [295] D. I. Cartwright and M. J. Field, "A Refinement of the Arithmetic Mean-Geometric Mean Inequality," *Proc. Amer. Math. Soc.*, Vol. 71, pp. 36–38, 1978. (Cited on pages 52 and 722.)
- [296] N. Castro-Gonzalez and E. Dopazo, "Representations of the Drazin Inverse for a Class of Block Matrices," *Lin. Alg. Appl.*, Vol. 400, pp. 253–269, 2005. (Cited on page 391.)
- [297] H. Caswell, *Matrix Population Models: Construction, Analysis, and Interpretation*, 2nd ed. Sunderland: Sinauer Associates, 2000. (Cited on page xvii.)
- [298] F. S. Cater, "Products of Central Collineations," *Lin. Alg. Appl.*, Vol. 19, pp. 251–274, 1978. (Cited on page 348.)
- [299] Y. Chabrillac and J.-P. Crouzeix, "Definiteness and Semidefiniteness of Quadratic Forms Revisited," *Lin. Alg. Appl.*, Vol. 63, pp. 283–292, 1984. (Cited on page 504.)
- [300] N. N. Chan and M. K. Kwong, "Hermitian Matrix Inequalities and a Conjecture," *Amer. Math. Monthly*, Vol. 92, pp. 533–541, 1985. (Cited on page 458.)

- [301] H. Chapellat, M. Mansour, and S. P. Bhattacharyya, "Elementary Proofs of Some Classical Stability Criteria," *IEEE Trans. Educ.*, Vol. 33, pp. 232–239, 1990. (Cited on pages 696 and 697.)
- [302] G. Chartrand, *Graphs and Digraphs*, 4th ed. Boca Raton: Chapman & Hall, 2004. (Cited on page xvii.)
- [303] G. Chartrand and L. Lesniak, *Graphs and Digraphs*, 4th ed. Boca Raton: Chapman & Hall/CRC, 2004. (Cited on page xvii.)
- [304] F. Chatelin, *Eigenvalues of Matrices*. New York: Wiley, 1993. (Cited on page xviii.)
- [305] J.-J. Chattot, *Computational Aerodynamics and Fluid Dynamics*. Berlin: Springer, 2002. (Cited on page xvii.)
- [306] N. A. Chaturvedi, N. H. McClamroch, and D. S. Bernstein, "Asymptotic Smooth Stabilization of the Inverted 3D Pendulum," *IEEE Trans. Autom. Contr.*, preprint. (Cited on page 189.)
- [307] J.-P. Chehab and M. Raydan, "Geometrical Properties of the Frobenius Condition Number for Positive Definite Matrices," *Lin. Alg. Appl.*, Vol. 429, pp. 2089–2097, 2008. (Cited on page 476.)
- [308] V.-S. Chellaboina and W. M. Haddad, "Is the Frobenius Matrix Norm Induced?" *IEEE Trans. Autom. Contr.*, Vol. 40, pp. 2137–2139, 1995. (Cited on page 579.)
- [309] V.-S. Chellaboina and W. M. Haddad, "Solution to 'Some Matrix Integral Identities'," *SIAM Rev.*, Vol. 39, pp. 763–765, 1997. (Cited on page 767.)
- [310] V.-S. Chellaboina, W. M. Haddad, D. S. Bernstein, and D. A. Wilson, "Induced Convolution Operator Norms of Linear Dynamical Systems," *Math. Contr. Sig. Sys.*, Vol. 13, pp. 216–239, 2000. (Cited on page 619.)
- [311] B. M. Chen, Z. Lin, and Y. Shamash, *Linear Systems Theory: A Structural Decomposition Approach*. Boston: Birkhauser, 2004. (Cited on pages xvii, 114, 325, 333, and 362.)
- [312] C. T. Chen, "A Generalization of the Inertia Theorem," *SIAM J. Appl. Math.*, Vol. 25, pp. 158–161, 1973. (Cited on page 793.)
- [313] C.-T. Chen, *Linear System Theory and Design*. New York: Holt, Rinehart, Winston, 1984. (Cited on pages xvii, 704, and 709.)
- [314] H. Chen, "A Unified Elementary Approach to Some Classical Inequalities," *Int. J. Math. Educ. Sci. Tech.*, Vol. 31, pp. 289–292, 2000. (Cited on pages 52 and 61.)
- [315] J. Chen, "On Bounds of Matrix Eigenvalues," *Math. Ineq. Appl.*, Vol. 10, pp. 723–726, 2007. (Cited on page 598.)
- [316] J.-Q. Chen, J. Ping, and F. Wang, *Group Representation for Physicists*, 2nd ed. Singapore: World Scientific, 2002. (Cited on page 224.)
- [317] L. Chen and C. S. Wong, "Inequalities for Singular Values and Traces," *Lin. Alg. Appl.*, Vol. 171, pp. 109–120, 1992. (Cited on pages 478 and 613.)
- [318] S. Chen, "Inequalities for M-Matrices and Inverse M-Matrices," *Lin. Alg. Appl.*, Vol. 426, pp. 610–618, 2007. (Cited on page 535.)

- [319] C.-M. Cheng, "Cases of Equality for a Singular Value Inequality for the Hadamard Product," *Lin. Alg. Appl.*, Vol. 177, pp. 209–231, 1992. (Cited on page 615.)
- [320] C.-M. Cheng, R. A. Horn, and C.-K. Li, "Inequalities and Equalities for the Cartesian Decomposition of Complex Matrices," *Lin. Alg. Appl.*, Vol. 341, pp. 219–237, 2002. (Cited on pages 443, 486, and 515.)
- [321] H.-W. Cheng and S. S.-T. Yau, "More Explicit Formulas for the Matrix Exponential," *Lin. Alg. Appl.*, Vol. 262, pp. 131–163, 1997. (Cited on page 673.)
- [322] S. Cheng and Y. Tian, "Moore-Penrose Inverses of Products and Differences of Orthogonal Projectors," *Acta Scientiarum Math.*, Vol. 69, pp. 533–542, 2003. (Cited on pages 380, 383, and 388.)
- [323] S. Cheng and Y. Tian, "Two Sets of New Characterizations for Normal and EP Matrices," *Lin. Alg. Appl.*, Vol. 375, pp. 181–195, 2003. (Cited on pages 178, 180, 372, 373, 394, and 395.)
- [324] M.-T. Chien and M. Neumann, "Positive Definiteness of Tridiagonal Matrices via the Numerical Range," *Elec. J. Lin. Alg.*, Vol. 3, pp. 93–102, 1998. (Cited on page 219.)
- [325] M.-D. Choi, "Tricks or Treats with the Hilbert Matrix," *Amer. Math. Monthly*, Vol. 90, pp. 301–312, 1983. (Cited on page 215.)
- [326] M. D. Choi, T. Y. Lam, and B. Reznick, "Sums of Squares of Real Polynomials," in *K Theory and Algebraic Geometry*, B. Jacob and A. Rosenberg, Eds. Providence: American Mathematical Society, 1995, pp. 103–126. (Cited on page 42.)
- [327] M.-D. Choi and P. Y. Wu, "Convex Combinations of Projections," *Lin. Alg. Appl.*, Vol. 136, pp. 25–42, 1990. (Cited on pages 208 and 360.)
- [328] J. Chollet, "Some Inequalities for Principal Submatrices," *Amer. Math. Monthly*, Vol. 104, pp. 609–617, 1997. (Cited on page 474.)
- [329] A. Choudhry, "Extraction of n th Roots of 2×2 Matrices," *Lin. Alg. Appl.*, Vol. 387, pp. 183–192, 2004. (Cited on page 348.)
- [330] M. T. Chu, R. E. Funderlic, and G. H. Golub, "A Rank-One Reduction Formula and Its Application to Matrix Factorizations," *SIAM Rev.*, Vol. 37, pp. 512–530, 1995. (Cited on pages 378 and 388.)
- [331] M. T. Chu, R. E. Funderlic, and G. H. Golub, "Rank Modifications of Semidefinite Matrices Associated with a Secant Update Formula," *SIAM J. Matrix Anal. Appl.*, Vol. 20, pp. 428–436, 1998. (Cited on page 457.)
- [332] X.-G. Chu, C.-E. Zhang, and F. Qi, "Two New Algebraic Inequalities with $2n$ Variables," *J. Ineq. Pure Appl. Math.*, Vol. 8, no. 4, pp. 1–6, 2007, Article 102. (Cited on page 17.)
- [333] J. Chuai and Y. Tian, "Rank Equalities and Inequalities for Kronecker Products of Matrices with Applications," *Appl. Math. Comp.*, Vol. 150, pp. 129–137, 2004. (Cited on pages 407 and 410.)
- [334] N. L. C. Chui and J. M. Maciejowski, "Realization of Stable Models with Subspace Methods," *Automatica*, Vol. 32, pp. 1587–1595, 1996. (Cited on page 714.)
- [335] F. R. K. Chung, *Spectral Graph Theory*. Providence: American Mathematical Society, 2000. (Cited on page xvii.)

- [336] A. Cizmesija and J. Pecaric, "Classical Hardy's and Carleman's Inequalities and Mixed Means," in *Survey on Classical Inequalities*, T. M. Rassias, Ed. Dordrecht: Kluwer, 2000, pp. 27–65. (Cited on page 58.)
- [337] D. J. Clements, B. D. O. Anderson, A. J. Laub, and J. B. Matson, "Spectral Factorization with Imaginary-Axis Zeros," *Lin. Alg. Appl.*, Vol. 250, pp. 225–252, 1997. (Cited on page 805.)
- [338] R. E. Cline, "Representations for the Generalized Inverse of a Partitioned Matrix," *SIAM J. Appl. Math.*, Vol. 12, pp. 588–600, 1964. (Cited on page 390.)
- [339] R. E. Cline and R. E. Funderlic, "The Rank of a Difference of Matrices and Associated Generalized Inverses," *Lin. Alg. Appl.*, Vol. 24, pp. 185–215, 1979. (Cited on pages 119, 121, and 383.)
- [340] M. J. Cloud and B. C. Drachman, *Inequalities with Applications to Engineering*. New York: Springer, 1998. (Cited on page 76.)
- [341] E. S. Coakley, F. M. Dopico, and C. R. Johnson, "Matrices for Orthogonal Groups Admitting Only Determinant One," *Lin. Alg. Appl.*, Vol. 428, pp. 796–813, 2008. (Cited on page 225.)
- [342] J. E. Cohen, "Spectral Inequalities for Matrix Exponentials," *Lin. Alg. Appl.*, Vol. 111, pp. 25–28, 1988. (Cited on page 690.)
- [343] J. E. Cohen, S. Friedland, T. Kato, and F. P. Kelly, "Eigenvalue Inequalities for Products of Matrix Exponentials," *Lin. Alg. Appl.*, Vol. 45, pp. 55–95, 1982. (Cited on page 692.)
- [344] D. K. Cohoon, "Sufficient Conditions for the Zero Matrix," *Amer. Math. Monthly*, Vol. 96, pp. 448–449, 1989. (Cited on page 127.)
- [345] D. Constales, "A Closed Formula for the Moore-Penrose Generalized Inverse of a Complex Matrix of Given Rank," *Acta Math. Hung.*, Vol. 80, pp. 83–88, 1998. (Cited on page 371.)
- [346] J. C. Conway and D. A. Smith, *On Quaternions and Octonions: Their Geometry, Arithmetic, and Symmetry*. Natick: A. K. Peters, 2003. (Cited on pages 47, 48, 223, 226, and 227.)
- [347] G. Corach, H. Porta, and L. Recht, "An Operator Inequality," *Lin. Alg. Appl.*, Vol. 142, pp. 153–158, 1990. (Cited on page 589.)
- [348] M. J. Corless and A. E. Frazho, *Linear Systems and Control: An Operator Perspective*. New York: Marcel Dekker, 2003. (Cited on page xvii.)
- [349] E. B. Corrachano and G. Sobczyk, Eds., *Geometric Algebra with Applications in Science and Engineering*. Boston: Birkhauser, 2001. (Cited on pages 188 and 227.)
- [350] P. J. Costa and S. Rabinowitz, "Matrix Differentiation Identities," *SIAM Rev.*, Vol. 36, pp. 657–659, 1994. (Cited on page 641.)
- [351] R. W. Cottle, "Quartic Barriers," *Comp. Optim. Appl.*, Vol. 12, pp. 81–105, 1999. (Cited on pages 497, 507, and 696.)
- [352] T. M. Cover and J. A. Thomas, "Determinant Inequalities via Information Theory," *SIAM J. Matrix Anal. Appl.*, Vol. 9, pp. 384–392, 1988. (Cited on pages 442, 487, and 506.)

- [353] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, 2nd ed. New York: Wiley, 2006. (Cited on pages xvii, 442, 487, and 492.)
- [354] D. E. Crabtree, "The Characteristic Vector of the Adjoint Matrix," *Amer. Math. Monthly*, Vol. 75, pp. 1127–1128, 1968. (Cited on page 342.)
- [355] C. R. Crawford and Y. S. Moon, "Finding a Positive Definite Linear Combination of Two Hermitian Matrices," *Lin. Alg. Appl.*, Vol. 51, pp. 37–48, 1983. (Cited on page 504.)
- [356] T. Crilly, "Cayley's Anticipation of a Generalised Cayley-Hamilton Theorem," *Historia Mathematica*, Vol. 5, pp. 211–219, 1978. (Cited on page 261.)
- [357] M. D. Crossley, *Essential Topology*. New York: Springer, 2005. (Cited on page 192.)
- [358] C. G. Cullen, "A Note on Normal Matrices," *Amer. Math. Monthly*, Vol. 72, pp. 643–644, 1965. (Cited on page 343.)
- [359] C. G. Cullen, *Matrices and Linear Transformations*, 2nd ed. Reading: Addison-Wesley, 1972, reprinted by Dover, Mineola, 1990. (Cited on pages 629 and 640.)
- [360] W. J. Culver, "On the Existence and Uniqueness of the Real Logarithm of a Matrix," *Proc. Amer. Math. Soc.*, Vol. 17, pp. 1146–1151, 1966. (Cited on page 722.)
- [361] R. F. Curtain and H. J. Zwart, *An Introduction to Infinite-Dimensional Linear Systems Theory*. New York: Springer, 1995. (Cited on page 691.)
- [362] M. L. Curtis, *Matrix Groups*, 2nd ed. New York: Springer, 1984. (Cited on pages 193, 226, and 229.)
- [363] D. Cvetkovic, P. Rowlinson, and S. Simic, *Eigenspaces of Graphs*. Cambridge: Cambridge University Press, 1997. (Cited on page xvii.)
- [364] P. J. Daboul and R. Delbourgo, "Matrix Representations of Octonions and Generalizations," *J. Math. Phys.*, Vol. 40, pp. 4134–4150, 1999. (Cited on page 227.)
- [365] G. Dahlquist, "On Matrix Majorants and Minorants, with Application to Differential Equations," *Lin. Alg. Appl.*, Vol. 52/53, pp. 199–216, 1983. (Cited on page 706.)
- [366] R. D'Andrea, "Extension of Parrott's Theorem to Nondefinite Scalings," *IEEE Trans. Autom. Contr.*, Vol. 45, pp. 937–940, 2000. (Cited on page 610.)
- [367] H. D'Angelo, *Linear Time-Varying Systems: Analysis and Synthesis*. Boston: Allyn and Bacon, 1970. (Cited on page 805.)
- [368] J. P. D'Angelo, *Inequalities from Complex Analysis*. Washington, DC: The Mathematical Association of America, 2002. (Cited on pages 70 and 565.)
- [369] F. M. Dannan, "Matrix and Operator Inequalities," *J. Ineq. Pure. Appl. Math.*, Vol. 2, no. 3/34, pp. 1–7, 2001. (Cited on pages 480 and 488.)
- [370] R. Datko and V. Seshadri, "A Characterization and a Canonical Decomposition of Hurwitzian Matrices," *Amer. Math. Monthly*, Vol. 77, pp. 732–733, 1970. (Cited on page 704.)
- [371] B. N. Datta, *Numerical Linear Algebra and Applications*. Pacific Grove: Brooks/Cole, 1995. (Cited on page xvii.)

- [372] B. N. Datta, "Stability and Inertia," *Lin. Alg. Appl.*, Vol. 302–303, pp. 563–600, 1999. (Cited on page 794.)
- [373] B. N. Datta, *Numerical Methods for Linear Control Systems*. San Diego, CA: Elsevier Academic Press, 2003. (Cited on page xvii.)
- [374] J. Dattorro, *Convex Optimization and Euclidean Distance Geometry*. Palo Alto: Meboo Publishing, 2005. (Cited on pages xvii, 118, 126, 323, and 642.)
- [375] I. Daubechies and J. C. Lagarias, "Sets of Matrices all Infinite Products of Which Converge," *Lin. Alg. Appl.*, Vol. 162, pp. 227–263, 1992. (Cited on page 715.)
- [376] P. J. Davis, *The Mathematics of Matrices*, 2nd ed. Boston: Ginn, 1965. (Cited on page xix.)
- [377] P. J. Davis, *Circulant Matrices*, 2nd ed. New York: Chelsea, 1994. (Cited on page 357.)
- [378] R. J. H. Dawlings, "Products of Idempotents in the Semigroup of Singular Endomorphisms of a Finite-Dimensional Vector Space," *Proc. Royal Soc. Edinburgh*, Vol. 91A, pp. 123–133, 1981. (Cited on page 350.)
- [379] J. Day, W. So, and R. C. Thompson, "Some Properties of the Campbell-Baker-Hausdorff Formula," *Lin. Multilin. Alg.*, Vol. 29, pp. 207–224, 1991. (Cited on page 655.)
- [380] J. Day, W. So, and R. C. Thompson, "The Spectrum of a Hermitian Matrix Sum," *Lin. Alg. Appl.*, Vol. 280, pp. 289–332, 1998. (Cited on page 513.)
- [381] P. W. Day, "Rearrangement Inequalities," *Canad. J. Math.*, Vol. 24, pp. 930–943, 1972. (Cited on page 61.)
- [382] C. de Boor, "An Empty Exercise," *SIGNUM*, Vol. 25, pp. 2–6, 1990. (Cited on page 164.)
- [383] P. P. N. de Groen, "A Counterexample on Vector Norms and the Subordinate Matrix Norms," *Amer. Math. Monthly*, Vol. 97, pp. 406–407, 1990. (Cited on page 579.)
- [384] F. R. de Hoog, R. P. Speed, and E. R. Williams, "On a Matrix Identity Associated with Generalized Least Squares," *Lin. Alg. Appl.*, Vol. 127, pp. 449–456, 1990. (Cited on page 378.)
- [385] W. de Launey and J. Seberry, "The Strong Kronecker Product," *J. Combinatorial Thy., Series A*, Vol. 66, pp. 192–213, 1994. (Cited on page 416.)
- [386] J. de Pillis, "Transformations on Partitioned Matrices," *Duke Math. J.*, Vol. 36, pp. 511–515, 1969. (Cited on pages 484 and 494.)
- [387] J. de Pillis, "Inequalities for Partitioned Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 4, pp. 79–94, 1971. (Cited on page 494.)
- [388] J. E. De Pillis, "Linear Operators and their Partitioned Matrices," *Notices Amer. Math. Soc.*, Vol. 14, p. 636, 1967. (Cited on page 494.)
- [389] L. G. de Pillis, "Determinants and Polynomial Root Structure," *Int. J. Math. Educ. Sci. Technol.*, Vol. 36, pp. 469–481, 2005. (Cited on page 255.)

- [390] E. de Souza and S. P. Bhattacharyya, "Controllability, Observability and the Solution of $AX - XB = C$," *Lin. Alg. Appl.*, Vol. 39, pp. 167–188, 1981. (Cited on page 796.)
- [391] E. de Souza and S. P. Bhattacharyya, "Controllability, Observability, and the Solution of $AX - XB = -C$," *Lin. Alg. Appl.*, Vol. 39, pp. 167–188, 1981. (Cited on page 797.)
- [392] P. N. De Souza and J.-N. Silva, *Berkeley Problems in Mathematics*, 3rd ed. New York: Springer, 2004. (Cited on pages 113, 118, 150, 151, and 457.)
- [393] L. M. DeAlba and C. R. Johnson, "Possible Inertia Combinations in the Stein and Lyapunov Equations," *Lin. Alg. Appl.*, Vol. 222, pp. 227–240, 1995. (Cited on pages 428 and 795.)
- [394] H. P. Decell, "An Application of the Cayley-Hamilton Theorem to Generalized Matrix Inversion," *SIAM Rev.*, Vol. 7, pp. 526–528, 1965. (Cited on page 374.)
- [395] N. Del Buono, L. Lopez, and T. Politi, "Computation of Functions of Hamiltonian and Skew-Symmetric Matrices," *Math. Computers Simulation*, preprint. (Cited on page 217.)
- [396] J. W. Demmel, *Applied Numerical Linear Algebra*. Philadelphia: SIAM, 1997. (Cited on page xviii.)
- [397] E. D. Denman and A. N. Beavers, "The Matrix Sign Function and Computations in Systems," *Appl. Math. Computation*, Vol. 2, pp. 63–94, 1976. (Cited on page 349.)
- [398] C. R. DePrima and C. R. Johnson, "The Range of $A^{-1}A^*$ in $GL(n, C)$," *Lin. Alg. Appl.*, Vol. 9, pp. 209–222, 1974. (Cited on page 346.)
- [399] C. A. Desoer and H. Haneda, "The Measure of a Matrix as a Tool to Analyze Computer Algorithms for Circuit Analysis," *IEEE Trans. Circ. Thy.*, Vol. 19, pp. 480–486, 1972. (Cited on page 691.)
- [400] E. Deutsch, "Matricial Norms," *Numer. Math.*, Vol. 16, pp. 73–84, 1970. (Cited on page 593.)
- [401] E. Deutsch, "Matricial Norms and the Zeros of Polynomials," *Lin. Alg. Appl.*, Vol. 3, pp. 483–489, 1970. (Cited on pages 709 and 710.)
- [402] E. Deutsch and M. Mlynarski, "Matricial Logarithmic Derivatives," *Lin. Alg. Appl.*, Vol. 19, pp. 17–31, 1978. (Cited on page 691.)
- [403] P. J. Dhrymes, *Mathematics for Econometrics*, 3rd ed. New York: Springer, 2000. (Cited on page xvii.)
- [404] L. Dieci, "Real Hamiltonian Logarithm of a Symplectic Matrix," *Lin. Alg. Appl.*, Vol. 281, pp. 227–246, 1998. (Cited on pages 660 and 685.)
- [405] R. Diestel, *Graph Theory*, 3rd ed. Berlin: Springer, 2006. (Cited on page xvii.)
- [406] L. L. Dines, "On the Mapping of Quadratic Forms," *Bull. Amer. Math. Soc.*, Vol. 47, pp. 494–498, 1941. (Cited on page 498.)
- [407] J. Ding, "Perturbation of Systems in Linear Algebraic Equations," *Lin. Multilin. Alg.*, Vol. 47, pp. 119–127, 2000. (Cited on page 592.)
- [408] J. Ding, "Lower and Upper Bounds for the Perturbation of General Linear Algebraic Equations," *Appl. Math. Lett.*, Vol. 14, pp. 49–52, 2001. (Cited on page 592.)

- [409] J. Ding and W. C. Pye, "On the Spectrum and Pseudoinverse of a Special Bordered Matrix," *Lin. Alg. Appl.*, Vol. 331, pp. 11–20, 2001. (Cited on page 263.)
- [410] A. Dittmer, "Cross Product Identities in Arbitrary Dimension," *Amer. Math. Monthly*, Vol. 101, pp. 887–891, 1994. (Cited on page 188.)
- [411] G. M. Dixon, *Division Algebras: Octonions, Quaternions, Complex Numbers and the Algebraic Design of Physics*. Dordrecht: Kluwer, 1994. (Cited on page 227.)
- [412] J. D. Dixon, "How Good Is Hadamard's Inequality for Determinants?" *Canad. Math. Bull.*, Vol. 27, pp. 260–264, 1984. (Cited on page 491.)
- [413] T. E. Djaferis and S. K. Mitter, "Algebraic Methods for the Study of Some Linear Matrix Equations," *Lin. Alg. Appl.*, Vol. 44, pp. 125–142, 1982. (Cited on page 797.)
- [414] D. Z. Djokovic, "Product of Two Involutions," *Arch. Math.*, Vol. 18, pp. 582–584, 1967. (Cited on page 350.)
- [415] D. Z. Djokovic, "On Some Representations of Matrices," *Lin. Multilin. Alg.*, Vol. 4, pp. 33–40, 1976. (Cited on pages 150 and 316.)
- [416] D. Z. Djokovic and O. P. Lossers, "A Determinant Inequality," *Amer. Math. Monthly*, Vol. 83, pp. 483–484, 1976. (Cited on page 153.)
- [417] D. Z. Djokovic and J. Malzan, "Products of Reflections in the Unitary Group," *Proc. Amer. Math. Soc.*, Vol. 73, pp. 157–160, 1979. (Cited on page 348.)
- [418] D. Z. Dokovic, "On the Product of Two Alternating Matrices," *Amer. Math. Monthly*, Vol. 98, pp. 935–936, 1991. (Cited on page 343.)
- [419] D. Z. Dokovic, "Unitary Similarity of Projectors," *Aequationes Mathematicae*, Vol. 42, pp. 220–224, 1991. (Cited on pages 314, 315, and 318.)
- [420] D. Z. Dokovic, F. Szechtman, and K. Zhao, "An Algorithm that Carries a Square Matrix into Its Transpose by an Involutory Congruence Transformation," *Elec. J. Lin. Alg.*, Vol. 10, pp. 320–340, 2003. (Cited on page 312.)
- [421] V. Dolotin and A. Morozov, *Introduction to Non-Linear Algebra*. Singapore: World Scientific, 2007. (Cited on page 416.)
- [422] W. F. Donoghue, *Monotone Matrix Functions and Analytic Continuation*. New York: Springer, 1974. (Cited on pages 446, 540, and 541.)
- [423] F. M. Dopico and C. R. Johnson, "Complementary Bases in Symplectic Matrices and a Proof That Their Determinant Is One," *Lin. Alg. Appl.*, Vol. 419, pp. 772–778, 2006. (Cited on page 217.)
- [424] F. M. Dopico, C. R. Johnson, and J. M. Molera, "Multiple LU Factorizations of a Singular Matrix," *Lin. Alg. Appl.*, Vol. 419, pp. 24–36, 2006. (Cited on page 346.)
- [425] C. Doran and A. Lasenby, *Geometric Algebra for Physicists*. Cambridge: Cambridge University Press, 2005. (Cited on pages 188 and 227.)
- [426] L. Dorst, D. Fontijne, and S. Mann, *Geometric Algebra for Computer Science: An Object-Oriented Approach to Geometry*. Amsterdam: Elsevier, 2007. (Cited on pages 227 and 565.)

- [427] R. G. Douglas, "On Majorization, Factorization, and Range Inclusion of Operators on Hilbert Space," *Proc. Amer. Math. Soc.*, Vol. 17, pp. 413–415, 1966. (Cited on page 431.)
- [428] P. G. Doyle and J. L. Snell, *Random Walks and Electric Networks*. Washington, DC: Mathematical Association of America, 1984. (Cited on page xvii.)
- [429] V. Dragan and A. Halanay, *Stabilization of Linear Systems*. Boston: Birkhauser, 1999. (Cited on page 805.)
- [430] S. S. Dragomir, "A Survey on Cauchy-Bunyakovsky-Schwarz Type Discrete Inequalities," *J. Ineq. Pure Appl. Math.*, Vol. 4, no. 3, pp. 1–142, 2003, Article 3. (Cited on pages 36, 57, 61, 63, 64, 65, 66, and 70.)
- [431] S. S. Dragomir, *Discrete Inequalities of the Cauchy-Bunyakovsky-Schwarz Type*. Hauppauge: Nova Science Publishers, 2004. (Cited on pages 36, 57, 61, 63, 64, 65, 66, and 70.)
- [432] S. S. Dragomir, "Some Reverses of the Generalised Triangle Inequality in Complex Inner Product Spaces," *Lin. Alg. Appl.*, Vol. 402, pp. 245–254, 2005. (Cited on page 70.)
- [433] S. S. Dragomir and C. J. Goh, "A Counterpart of Jensen's Discrete Inequality for Differentiable Convex Mappings and Applications in Information Theory," *Math. Comput. Modelling*, Vol. 24, pp. 1–11, 1996. (Cited on page 59.)
- [434] S. S. Dragomir and C. J. Goh, "Some Bounds on Entropy Measures in Information Theory," *Appl. Math. Lett.*, Vol. 10, pp. 23–28, 1997. (Cited on page 59.)
- [435] S. S. Dragomir, C. E. Pearce, and J. Pecaric, "Some New Inequalities for the Logarithmic Map, with Applications to Entropy and Mutual Information," *Kyungpook Math. J.*, Vol. 41, pp. 115–125, 2001. (Cited on pages 59 and 64.)
- [436] M. P. Drazin, "A Note on Skew-Symmetric Matrices," *Math. Gaz.*, Vol. 36, pp. 253–255, 1952. (Cited on page 259.)
- [437] D. Drissi, "Sharp Inequalities for Some Operator Means," *SIAM J. Matrix Anal. Appl.*, Vol. 28, pp. 822–828, 2006. (Cited on page 38.)
- [438] D. Drivaliaris, S. Karanasios, and D. Pappas, "Factorizations of EP Operators," *Lin. Alg. Appl.*, Vol. 429, pp. 1555–1567, 2008. (Cited on page 372.)
- [439] R. Drnovsek, H. Radjavi, and P. Rosenthal, "A Characterization of Commutators of Idempotents," *Lin. Alg. Appl.*, Vol. 347, pp. 91–99, 2002. (Cited on page 200.)
- [440] S. W. Drury and G. P. H. Styan, "Normal Matrix and a Commutator," *IMAGE*, Vol. 31, p. 24, 2003. (Cited on page 185.)
- [441] H. Du, C. Deng, and Q. Li, "On the Infimum Problem of Hilbert Space Effects," *Science in China: Series A Math.*, Vol. 49, pp. 545–556, 2006. (Cited on page 460.)
- [442] J. J. Duistermaat and J. A. C. Kolk, Eds., *Multidimensional Real Analysis I: Differentiation*. Cambridge: Cambridge University Press, 2004. (Cited on pages 634, 638, and 682.)
- [443] I. Duleba, "On a Computationally Simple Form of the Generalized Campbell-Baker-Hausdorff-Dynkin Formula," *Sys. Contr. Lett.*, Vol. 34, pp. 191–202, 1998. (Cited on page 678.)

- [444] G. E. Dullerud and F. Paganini, *A Course in Robust Control Theory: A Convex Approach*, 2nd ed. New York: Springer, 1999. (Cited on page xvii.)
- [445] D. S. Dummit and R. M. Foote, *Abstract Algebra*, 3rd ed. New York: Wiley, 2004. (Cited on pages 222, 223, 357, and 361.)
- [446] C. F. Dunkl and K. S. Williams, "A Simple Inequality," *Amer. Math. Monthly*, Vol. 71, pp. 53–54, 1964. (Cited on page 565.)
- [447] H. Dym, *Linear Algebra in Action*. Providence: American Mathematical Society, 2006. (Cited on pages 56, 129, 131, 134, 160, 280, 309, 310, 311, 335, 336, 391, 431, 458, 507, 510, 525, 590, 591, 610, 634, 636, 638, 698, and 794.)
- [448] A. Edelman and G. Strang, "Pascal Matrices," *Amer. Math. Monthly*, Vol. 111, pp. 189–197, 2004. (Cited on page 447.)
- [449] O. Egecioglu, "Parallelogram-Law-Type Identities," *Lin. Alg. Appl.*, Vol. 225, pp. 1–12, 1995. (Cited on page 70.)
- [450] H. G. Eggleston, *Convexity*. Cambridge: Cambridge University Press, 1958. (Cited on page 164.)
- [451] L. Elsner, D. Hershkowitz, and H. Schneider, "Bounds on Norms of Compound Matrices and on Products of Eigenvalues," *Bull. London Math. Soc.*, Vol. 32, pp. 15–24, 2000. (Cited on pages 17 and 413.)
- [452] L. Elsner and K. D. Ikramov, "Normal Matrices: An Update," *Lin. Alg. Appl.*, Vol. 285, pp. 291–303, 1998. (Cited on pages 180, 650, and 695.)
- [453] L. Elsner, C. R. Johnson, and J. A. D. Da Silva, "The Perron Root of a Weighted Geometric Mean of Nonnegative Matrices," *Lin. Multilin. Alg.*, Vol. 24, pp. 1–13, 1988. (Cited on page 415.)
- [454] L. Elsner and M. H. C. Paardekooper, "On Measures of Nonnormality of Matrices," *Lin. Alg. Appl.*, Vol. 92, pp. 107–124, 1987. (Cited on page 180.)
- [455] L. Elsner and P. Rozsa, "On Eigenvectors and Adjoints of Modified Matrices," *Lin. Multilin. Alg.*, Vol. 10, pp. 235–247, 1981. (Cited on page 142.)
- [456] L. Elsner and T. Szulc, "Convex Sets of Schur Stable and Stable Matrices," *Lin. Multilin. Alg.*, Vol. 48, pp. 1–19, 2000. (Cited on page 269.)
- [457] A. Engel, *Problem-Solving Strategies*. New York: Springer, 1998. (Cited on pages 31, 32, 34, 39, 40, 41, 42, 43, 44, 45, 46, 49, 52, 58, 60, and 158.)
- [458] K. Engo, "On the BCH Formula in $\mathfrak{so}(3)$," *Numer. Math. BIT*, Vol. 41, pp. 629–632, 2001. (Cited on page 683.)
- [459] K. Erdmann and M. J. Wildon, *Introduction to Lie Algebras*. New York: Springer, 2006. (Cited on page 229.)
- [460] J. A. Erdos, "On Products of Idempotent Matrices," *Glasgow Math. J.*, Vol. 8, pp. 118–122, 1967. (Cited on page 350.)
- [461] R. Eriksson, "On the Measure of Solid Angles," *Math. Mag.*, Vol. 63, pp. 184–187, 1990. (Cited on page 161.)
- [462] J.-C. Evard and F. Uhlig, "On the Matrix Equation $f(X) = A$," *Lin. Alg. Appl.*, Vol. 162–164, pp. 447–519, 1992. (Cited on page 722.)

- [463] S. Fallat and M. J. Tsatsomeros, "On the Cayley Transform of Positivity Classes of Matrices," *Elec. J. Lin. Alg.*, Vol. 9, pp. 190–196, 2002. (Cited on page 453.)
- [464] K. Fan, "Generalized Cayley Transforms and Strictly Dissipative Matrices," *Lin. Alg. Appl.*, Vol. 5, pp. 155–172, 1972. (Cited on page 453.)
- [465] K. Fan, "On Real Matrices with Positive Definite Symmetric Component," *Lin. Multilin. Alg.*, Vol. 1, pp. 1–4, 1973. (Cited on pages 453 and 487.)
- [466] K. Fan, "On Strictly Dissipative Matrices," *Lin. Alg. Appl.*, Vol. 9, pp. 223–241, 1974. (Cited on pages 453 and 485.)
- [467] M. Fang, "Bounds on Eigenvalues of the Hadamard Product and the Fan Product of Matrices," *Lin. Alg. Appl.*, Vol. 425, pp. 7–15, 2007. (Cited on page 415.)
- [468] Y. Fang, K. A. Loparo, and X. Feng, "Inequalities for the Trace of Matrix Product," *IEEE Trans. Autom. Contr.*, Vol. 39, pp. 2489–2490, 1994. (Cited on page 334.)
- [469] R. W. Farebrother, "A Class of Square Roots of Involutory Matrices," *IMAGE*, Vol. 28, pp. 26–28, 2002. (Cited on page 398.)
- [470] R. W. Farebrother, J. Gross, and S.-O. Troschke, "Matrix Representation of Quaternions," *Lin. Alg. Appl.*, Vol. 362, pp. 251–255, 2003. (Cited on page 229.)
- [471] R. W. Farebrother and I. Wrobel, "Regular and Reflected Rotation Matrices," *IMAGE*, Vol. 29, pp. 24–25, 2002. (Cited on page 348.)
- [472] D. R. Farenick, M. Krupnik, N. Krupnik, and W. Y. Lee, "Normal Toeplitz Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 17, pp. 1037–1043, 1996. (Cited on page 357.)
- [473] A. Fassler and E. Stiefel, *Group Theoretical Methods and Their Applications*. Boston: Birkhauser, 1992. (Cited on page 229.)
- [474] A. E. Fekete, *Real Linear Algebra*. New York: Marcel Dekker, 1985. (Cited on pages 188 and 200.)
- [475] B. Q. Feng, "Equivalence Constants for Certain Matrix Norms," *Lin. Alg. Appl.*, Vol. 374, pp. 247–253, 2003. (Cited on page 583.)
- [476] B. Q. Feng and A. Tonge, "Equivalence Constants for Certain Matrix Norms, II," *Lin. Alg. Appl.*, Vol. 420, pp. 388–399, 2007. (Cited on pages 574 and 583.)
- [477] R. Fenn, *Geometry*. New York: Springer, 2001. (Cited on pages 30, 161, 188, 193, 226, 227, and 278.)
- [478] P. G. Ferreira and S. P. Bhattacharyya, "On Blocking Zeros," *IEEE Trans. Autom. Contr.*, Vol. AC-22, pp. 258–259, 1977. (Cited on page 805.)
- [479] J. H. Ferziger and M. Peric, *Computational Methods for Fluid Dynamics*, 3rd ed. Berlin: Springer, 2002. (Cited on page xvii.)
- [480] M. Fiedler, "A Note on the Hadamard Product of Matrices," *Lin. Alg. Appl.*, Vol. 49, pp. 233–235, 1983. (Cited on pages 532 and 535.)
- [481] M. Fiedler, *Special Matrices and Their Applications in Numerical Mathematics*. Dordrecht: Martinus Nijhoff, 1986. (Cited on pages xvii, 255, 257, 275, 329, 355, 390, 412, and 702.)

- [482] M. Fiedler, "Some Applications of Matrices and Graphs in Euclidean Geometry," in *Handbook of Linear Algebra*, L. Hogben, Ed. Boca Raton: Chapman & Hall/CRC, 2007, pp. 66–1–66–15. (Cited on pages 160 and 532.)
- [483] M. Fiedler and T. L. Markham, "A Characterization of the Moore-Penrose Inverse," *Lin. Alg. Appl.*, Vol. 179, pp. 129–133, 1993. (Cited on pages 149 and 376.)
- [484] M. Fiedler and T. L. Markham, "An Observation on the Hadamard Product of Hermitian Matrices," *Lin. Alg. Appl.*, Vol. 215, pp. 179–182, 1995. (Cited on page 534.)
- [485] M. Fiedler and T. L. Markham, "Some Results on the Bergstrom and Minkowski Inequalities," *Lin. Alg. Appl.*, Vol. 232, pp. 199–211, 1996. (Cited on page 468.)
- [486] M. Fiedler and V. Ptak, "A New Positive Definite Geometric Mean of Two Positive Definite Matrices," *Lin. Alg. Appl.*, Vol. 251, pp. 1–20, 1997. (Cited on page 463.)
- [487] J. A. Fill and D. E. Fishkind, "The Moore-Penrose Generalized Inverse for Sums of Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 21, pp. 629–635, 1999. (Cited on page 378.)
- [488] P. A. Fillmore, "On Similarity and the Diagonal of a Matrix," *Amer. Math. Monthly*, Vol. 76, pp. 167–169, 1969. (Cited on page 313.)
- [489] P. A. Fillmore, "On Sums of Projections," *J. Funct. Anal.*, Vol. 4, pp. 146–152, 1969. (Cited on page 360.)
- [490] P. A. Fillmore and J. P. Williams, "On Operator Ranges," *Adv. Math.*, Vol. 7, pp. 254–281, 1971. (Cited on page 431.)
- [491] C. H. Fitzgerald and R. A. Horn, "On Fractional Hadamard Powers of Positive Definite Matrices," *J. Math. Anal. Appl.*, Vol. 61, pp. 633–642, 1977. (Cited on page 531.)
- [492] H. Flanders, "Methods of Proof in Linear Algebra," *Amer. Math. Monthly*, Vol. 63, pp. 1–15, 1956. (Cited on page 214.)
- [493] H. Flanders, "On the Norm and Spectral Radius," *Lin. Multilin. Alg.*, Vol. 2, pp. 239–240, 1974. (Cited on page 603.)
- [494] H. Flanders, "An Extremal Problem in the Space of Positive Definite Matrices," *Lin. Multilin. Alg.*, Vol. 3, pp. 33–39, 1975. (Cited on page 459.)
- [495] W. Fleming, *Functions of Several Variables*, 2nd ed. New York: Springer, 1987. (Cited on page 494.)
- [496] T. M. Flett, *Differential Analysis*. Cambridge: Cambridge University Press, 1980. (Cited on page 642.)
- [497] S. Foldes, *Fundamental Structures of Algebra and Discrete Mathematics*. New York: Wiley, 1994. (Cited on page 222.)
- [498] J. Foley, A. van Dam, S. Feiner, and J. Hughes, *Computer Graphics Principles and Practice*, 2nd ed. Reading: Addison-Wesley, 1990. (Cited on pages xvii and 194.)
- [499] E. Formanek, "Polynomial Identities and the Cayley-Hamilton Theorem," *Mathematical Intelligencer*, Vol. 11, pp. 37–39, 1989. (Cited on pages 149, 214, and 260.)
- [500] E. Formanek, *The Polynomial Identities and Invariants of $n \times n$ Matrices*. Providence: American Mathematical Society, 1991. (Cited on pages 149, 214, and 260.)

- [501] L. R. Foulds, *Graph Theory Applications*. New York: Springer, 1992. (Cited on page xvii.)
- [502] B. A. Francis, *A Course in H_∞ Control Theory*. New York: Springer, 1987. (Cited on page xvii.)
- [503] J. Franklin, *Matrix Theory*. Englewood Cliffs: Prentice-Hall, 1968. (Cited on page xix.)
- [504] M. Frazier, *An Introduction to Wavelets through Linear Algebra*. New York: Springer, 1999. (Cited on page xvii.)
- [505] S. Friedland, "A Note on a Determinantal Inequality," *Lin. Alg. Appl.*, Vol. 141, pp. 221–222, 1990. (Cited on page 601.)
- [506] S. Friedland, D. Hershkowitz, and S. M. Rump, "Positive Entries of Stable Matrices," *Elec. J. Lin. Alg.*, Vol. 12, pp. 17–24, 2005. (Cited on page 708.)
- [507] S. Friedland and A. Torokhti, "Generalized Rank-Constrained Matrix Approximation," *SIAM J. Matrix Anal. Appl.*, Vol. 29, pp. 656–659, 2007. (Cited on pages 614 and 618.)
- [508] P. A. Fuhrmann, *A Polynomial Approach to Linear Algebra*. New York: Springer, 1996. (Cited on pages 197, 255, 257, 258, 281, 309, and 473.)
- [509] J. I. Fujii and M. Fujii, "Kolmogorov's Complexity for Positive Definite Matrices," *Lin. Alg. Appl.*, Vol. 341, pp. 171–180, 2002. (Cited on page 502.)
- [510] J. I. Fujii, M. Fujii, T. Furuta, and R. Nakamoto, "Norm Inequalities Equivalent to Heinz Inequality," *Proc. Amer. Math. Soc.*, Vol. 118, pp. 827–830, 1993. (Cited on page 589.)
- [511] J.-I. Fujii, S. Izumino, and Y. Seo, "Determinant for Positive Operators and Specht's Theorem," *Scientiae Mathematicae*, Vol. 1, pp. 307–310, 1998. (Cited on page 53.)
- [512] M. Fujii, E. Kamei, and R. Nakamoto, "On a Question of Furuta on Chaotic Order," *Lin. Alg. Appl.*, Vol. 341, pp. 119–129, 2002. (Cited on pages 461, 466, 522, and 523.)
- [513] M. Fujii, E. Kamei, and R. Nakamoto, "On a Question of Furuta on Chaotic Order, II," *Math. J. Okayama University*, Vol. 45, pp. 123–131, 2003. (Cited on page 466.)
- [514] M. Fujii and F. Kubo, "Buzano's Inequality and Bounds for Roots of Algebraic Equations," *Proc. Amer. Math. Soc.*, Vol. 117, pp. 359–361, 1993. (Cited on pages 67, 565, and 711.)
- [515] M. Fujii, S. H. Lee, Y. Seo, and D. Jung, "Reverse Inequalities on Chaotically Geometric Mean via Specht Ratio," *Math. Ineq. Appl.*, Vol. 6, pp. 509–519, 2003. (Cited on page 34.)
- [516] M. Fujii, S. H. Lee, Y. Seo, and D. Jung, "Reverse Inequalities on Chaotically Geometric Mean via Specht Ratio, II," *J. Ineq. Pure Appl. Math.*, Vol. 4/40, pp. 1–8, 2003. (Cited on page 53.)
- [517] M. Fujii and R. Nakamoto, "Rota's Theorem and Heinz Inequalities," *Lin. Alg. Appl.*, Vol. 214, pp. 271–275, 1995. (Cited on page 589.)

- [518] M. Fujii, Y. Seo, and M. Tominaga, “Golden-Thompson Type Inequalities Related to a Geometric Mean via Specht’s Ratio,” *Math. Ineq. Appl.*, Vol. 5, pp. 573–582, 2002. (Cited on pages 685 and 686.)
- [519] A. T. Fuller, “Conditions for a Matrix to Have Only Characteristic Roots with Negative Real Parts,” *J. Math. Anal. Appl.*, Vol. 23, pp. 71–98, 1968. (Cited on pages 406, 413, 416, and 697.)
- [520] W. Fulton and J. Harris, *Representation Theory*. New York: Springer, 2004. (Cited on page 223.)
- [521] S. Furuichi, “Matrix Trace Inequalities on the Tsallis Entropies,” *J. Ineq. Pure Appl. Math.*, Vol. 9, no. 1, pp. 1–7, 2008, Article 1. (Cited on page 479.)
- [522] T. Furuta, “ $A \geq B \geq 0$ Assures $(B^r A^p B^r)^{1/q} \geq B^{(p+2r)/q}$ for $r \geq 0$, $p \geq 0$, $q \geq 1$ with $(1+2r)q \geq p+2r$,” *Proc. Amer. Math. Soc.*, Vol. 101, pp. 85–88, 1987. (Cited on page 433.)
- [523] T. Furuta, “Norm Inequalities Equivalent to Loewner-Heinz Theorem,” *Rev. Math. Phys.*, Vol. 1, pp. 135–137, 1989. (Cited on pages 520 and 582.)
- [524] T. Furuta, “Two Operator Functions with Monotone Property,” *Proc. Amer. Math. Soc.*, Vol. 111, pp. 511–516, 1991. (Cited on page 461.)
- [525] T. Furuta, “A Note on the Arithmetic-Geometric-Mean Inequality for Every Unitarily Invariant Matrix Norm,” *Lin. Alg. Appl.*, Vol. 208/209, pp. 223–228, 1994. (Cited on page 589.)
- [526] T. Furuta, “Extension of the Furuta Inequality and Ando-Hiai Log-Majorization,” *Lin. Alg. Appl.*, Vol. 219, pp. 139–155, 1995. (Cited on page 461.)
- [527] T. Furuta, “Generalizations of Kosaki Trace Inequalities and Related Trace Inequalities on Chaotic Order,” *Lin. Alg. Appl.*, Vol. 235, pp. 153–161, 1996. (Cited on page 481.)
- [528] T. Furuta, “Operator Inequalities Associated with Hölder-McCarthy and Kantorovich Inequalities,” *J. Ineq. Appl.*, Vol. 2, pp. 137–148, 1998. (Cited on page 465.)
- [529] T. Furuta, “Simple Proof of the Concavity of Operator Entropy $f(A) = -A \log A$,” *Math. Ineq. Appl.*, Vol. 3, pp. 305–306, 2000. (Cited on page 441.)
- [530] T. Furuta, *Invitation to Linear Operators: From Matrices to Bounded Linear Operators on a Hilbert Space*. London: Taylor and Francis, 2001. (Cited on pages 24, 209, 433, 434, 436, 455, 461, 464, 465, 502, 511, 522, and 568.)
- [531] T. Furuta, “Spectral Order $A \succ B$ if and only if $A^{2p-r} \geq \left(A^{-r/2} B^p A^{-r/2}\right)^{(2p-r)/(p-r)}$ for all $p > r \geq 0$ and Its Application,” *Math. Ineq. Appl.*, Vol. 4, pp. 619–624, 2001. (Cited on pages 466 and 523.)
- [532] T. Furuta, “The Hölder-McCarthy and the Young Inequalities Are Equivalent for Hilbert Space Operators,” *Amer. Math. Monthly*, Vol. 108, pp. 68–69, 2001. (Cited on page 502.)
- [533] T. Furuta, “Convergence of Logarithmic Trace Inequalities via Generalized Lie-Trotter Formulae,” *Lin. Alg. Appl.*, Vol. 396, pp. 353–372, 2005. (Cited on pages 683 and 686.)

- [534] T. Furuta, “Concrete Examples of Operator Monotone Functions Obtained by an Elementary Method without Appealing to Löwner Integral Representation,” *Lin. Alg. Appl.*, Vol. 429, pp. 972–980, 2008. (Cited on page 467.)
- [535] F. Gaines, “A Note on Matrices with Zero Trace,” *Amer. Math. Monthly*, Vol. 73, pp. 630–631, 1966. (Cited on pages 184, 185, and 313.)
- [536] A. Galantai, *Projectors and Projection Methods*. Dordrecht: Kluwer, 2004. (Cited on pages 204, 230, 315, 316, 329, 335, 336, 571, 572, and 636.)
- [537] A. Galantai, “Subspaces, Angles and Pairs of Orthogonal Projections,” *Lin. Multilin. Alg.*, Vol. 56, pp. 227–260, 2008. (Cited on pages 112, 209, 210, 315, 316, 329, 335, 336, 337, and 380.)
- [538] J. Gallier and D. Xu, “Computing Exponentials of Skew-Symmetric Matrices and Logarithms of Orthogonal Matrices,” *Int. J. Robotics Automation*, Vol. 17, pp. 1–11, 2002. (Cited on page 193.)
- [539] L. Gangsong and Z. Guobiao, “Inverse Forms of Hadamard Inequality,” *SIAM J. Matrix Anal. Appl.*, Vol. 23, pp. 990–997, 2002. (Cited on page 511.)
- [540] F. R. Gantmacher, *The Theory of Matrices*. New York: Chelsea, 1959, Vol. I. (Cited on pages xix and 541.)
- [541] F. R. Gantmacher, *The Theory of Matrices*. New York: Chelsea, 1959, Vol. II. (Cited on pages xix, 304, 318, and 722.)
- [542] D. J. H. Garling, *Inequalities: A Journey into Linear Analysis*. Cambridge: Cambridge University Press, 2007. (Cited on pages 33, 53, 54, 58, 63, 75, 162, 163, 569, and 574.)
- [543] J. Garloff and D. G. Wagner, “Hadamard Products of Stable Polynomials Are Stable,” *J. Math. Anal. Appl.*, Vol. 202, pp. 797–809, 1996. (Cited on page 697.)
- [544] T. Geerts, “A Necessary and Sufficient Condition for Solvability of the Linear-Quadratic Control Problem without Stability,” *Sys. Contr. Lett.*, Vol. 11, pp. 47–51, 1988. (Cited on page 805.)
- [545] I. M. Gelfand, M. M. Kapranov, and A. V. Zelevinsky, *Discriminants, Resultants, and Multidimensional Determinants*. Boston: Birkhauser, 1994. (Cited on page 416.)
- [546] M. G. Genton, “Classes of Kernels for Machine Learning: A Statistics Perspective,” *J. Machine Learning Res.*, Vol. 2, pp. 299–312, 2001. (Cited on page 445.)
- [547] A. George and K. D. Ikramov, “Common Invariant Subspaces of Two Matrices,” *Lin. Alg. Appl.*, Vol. 287, pp. 171–179, 1999. (Cited on pages 342 and 358.)
- [548] P. Gerdes, *Adventures in the World of Matrices*. Hauppauge: Nova Publishers, 2007. (Cited on page 357.)
- [549] A. Gerrard and J. M. Burch, *Introduction to Matrix Methods in Optics*. New York: Wiley, 1975. (Cited on page xvii.)
- [550] A. Gheondea, S. Gudder, and P. Jonas, “On the Infimum of Quantum Effects,” *J. Math. Phys.*, Vol. 46, pp. 1–11, 2005, paper 062102. (Cited on page 460.)
- [551] F. A. A. Ghouraba and M. A. Seoud, “Set Matrices,” *Int. J. Math. Educ. Sci. Technol.*, Vol. 30, pp. 651–659, 1999. (Cited on page 230.)

- [552] M. I. Gil, "On Inequalities for Eigenvalues of Matrices," *Lin. Alg. Appl.*, Vol. 184, pp. 201–206, 1993. (Cited on page 325.)
- [553] R. Gilmore, *Lie Groups, Lie Algebras, and Some of Their Applications*. New York: Wiley, 1974, reprinted by Dover, Mineola, 2005. (Cited on page 229.)
- [554] R. Gilmore, *Lie Groups, Physics, and Geometry: An Introduction for Physicists, Engineers, and Chemists*. Cambridge: Cambridge University Press, 2008. (Cited on page 229.)
- [555] P. R. Girard, *Quaternions, Clifford Algebras and Relativistic Physics*. Boston: Birkhauser, 2007. (Cited on pages 188, 193, and 225.)
- [556] M. L. Glasser, "Exponentials of Certain Hilbert Space Operators," *SIAM Rev.*, Vol. 34, pp. 498–500, 1992. (Cited on page 683.)
- [557] C. Godsil and G. Royle, *Algebraic Graph Theory*. New York: Springer, 2001. (Cited on page xvii.)
- [558] S. K. Godunov, *Modern Aspects of Linear Algebra*. Providence: American Mathematical Society, 1998. (Cited on pages xix, 211, and 699.)
- [559] I. Gohberg, P. Lancaster, and L. Rodman, *Matrix Polynomials*. New York: Academic Press, 1982. (Cited on pages 234, 281, and 361.)
- [560] I. Gohberg, P. Lancaster, and L. Rodman, *Invariant Subspaces of Matrices with Applications*. New York: Wiley, 1986. (Cited on pages 336, 361, and 636.)
- [561] I. Gohberg, P. Lancaster, and L. Rodman, "On Hermitian Solutions of the Symmetric Algebraic Riccati Equation," *SIAM J. Contr. Optim.*, Vol. 24, pp. 1323–1334, 1986. (Cited on pages 787, 788, and 805.)
- [562] I. Gohberg, P. Lancaster, and L. Rodman, *Indefinite Linear Algebra and Applications*. Boston: Birkhauser, 2005. (Cited on pages 361 and 805.)
- [563] M. A. Golberg, "The Derivative of a Determinant," *Amer. Math. Monthly*, Vol. 79, pp. 1124–1126, 1972. (Cited on pages 640 and 678.)
- [564] M. Goldberg, "Mixed Multiplicativity and l_p Norms for Matrices," *Lin. Alg. Appl.*, Vol. 73, pp. 123–131, 1986. (Cited on page 583.)
- [565] M. Goldberg, "Equivalence Constants for l_p Norms of Matrices," *Lin. Multilin. Alg.*, Vol. 21, pp. 173–179, 1987. (Cited on page 583.)
- [566] M. Goldberg, "Multiplicativity Factors and Mixed Multiplicativity," *Lin. Alg. Appl.*, Vol. 97, pp. 45–56, 1987. (Cited on page 583.)
- [567] M. Goldberg and G. Zwas, "On Matrices Having Equal Spectral Radius and Spectral Norm," *Lin. Alg. Appl.*, Vol. 8, pp. 427–434, 1974. (Cited on page 603.)
- [568] H. Goller, "Shorted Operators and Rank Decomposition Matrices," *Lin. Alg. Appl.*, Vol. 81, pp. 207–236, 1986. (Cited on page 530.)
- [569] G. H. Golub and C. F. Van Loan, *Matrix Computations*, 3rd ed. Baltimore: Johns Hopkins University Press, 1996. (Cited on pages xviii, 614, and 646.)
- [570] N. C. Gonzalez, J. J. Koliha, and Y. Wei, "Integral Representation of the Drazin Inverse," *Elect. J. Lin. Alg.*, Vol. 9, pp. 129–131, 2002. (Cited on page 680.)

- [571] F. M. Goodman, *Algebra: Abstract and Concrete*, 2nd ed. Englewood Cliffs: Prentice-Hall, 2003. (Cited on page 223.)
- [572] L. E. Goodman and W. H. Warner, *Statics*. Belmont: Wadsworth Publishing Company, 1964, reprinted by Dover, Mineola, 2001. (Cited on page 188.)
- [573] N. Gordon and D. Salmond, "Bayesian Pattern Matching Technique for Target Acquisition," *J. Guid. Contr. Dyn.*, Vol. 22, pp. 68–77, 1999. (Cited on page 139.)
- [574] A. Goroncy and T. Rychlik, "How Deviant Can You Be? The Complete Solution," *Math. Ineq. Appl.*, Vol. 9, pp. 633–647, 2006. (Cited on page 51.)
- [575] W. Govaerts and J. D. Pryce, "A Singular Value Inequality for Block Matrices," *Lin. Alg. Appl.*, Vol. 125, pp. 141–145, 1989. (Cited on page 610.)
- [576] W. Govaerts and B. Sijnave, "Matrix Manifolds and the Jordan Structure of the Bialternate Matrix Product," *Lin. Alg. Appl.*, Vol. 292, pp. 245–266, 1999. (Cited on pages 413 and 416.)
- [577] R. Gow, "The Equivalence of an Invertible Matrix to Its Transpose," *Lin. Alg. Appl.*, Vol. 8, pp. 329–336, 1980. (Cited on pages 312 and 351.)
- [578] R. Gow and T. J. Laffey, "Pairs of Alternating Forms and Products of Two Skew-Symmetric Matrices," *Lin. Alg. Appl.*, Vol. 63, pp. 119–132, 1984. (Cited on pages 343 and 351.)
- [579] A. Graham, *Kronecker Products and Matrix Calculus with Applications*. Chichester: Ellis Horwood, 1981. (Cited on page 416.)
- [580] F. A. Graybill, *Matrices with Applications in Statistics*, 2nd ed. Belmont: Wadsworth, 1983. (Cited on page xvii.)
- [581] J. Grcar, "Linear Algebra People," <http://seesar.lbl.gov/ccse/people/grcar/index.html>. (Cited on page 164.)
- [582] J. F. Grcar, "A Matrix Lower Bound," Lawrence Berkeley National Laboratory, Report LBNL-50635, 2002. (Cited on pages 559 and 560.)
- [583] W. L. Green and T. D. Morley, "Operator Means and Matrix Functions," *Lin. Alg. Appl.*, Vol. 137/138, pp. 453–465, 1990. (Cited on pages 463 and 529.)
- [584] D. H. Greene and D. E. Knuth, *Mathematics for the Analysis of Algorithms*, 3rd ed. Boston: Birkhauser, 1990. (Cited on page 17.)
- [585] W. H. Greub, *Multilinear Algebra*. New York: Springer, 1967. (Cited on page 416.)
- [586] W. H. Greub, *Linear Algebra*. New York: Springer, 1981. (Cited on page xix.)
- [587] T. N. E. Greville, "Notes on the Generalized Inverse of a Matrix Product," *SIAM Rev.*, Vol. 8, pp. 518–521, 1966. (Cited on page 380.)
- [588] T. N. E. Greville, "Solutions of the Matrix Equation $XAX = X$ and Relations between Oblique and Orthogonal projectors," *SIAM J. Appl. Math.*, Vol. 26, pp. 828–832, 1974. (Cited on pages 210 and 381.)
- [589] R. Grone, C. R. Johnson, E. M. Sa, and H. Wolkowicz, "Normal Matrices," *Lin. Alg. Appl.*, Vol. 87, pp. 213–225, 1987. (Cited on pages 180, 190, and 341.)
- [590] J. Gross, "A Note on a Partial Ordering in the Set of Hermitian Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 18, pp. 887–892, 1997. (Cited on pages 523 and 526.)

- [591] J. Gross, "Some Remarks on Partial Orderings of Hermitian Matrices," *Lin. Multilin. Alg.*, Vol. 42, pp. 53–60, 1997. (Cited on pages 523 and 526.)
- [592] J. Gross, "More on Concavity of a Matrix Function," *SIAM J. Matrix Anal. Appl.*, Vol. 19, pp. 365–368, 1998. (Cited on page 530.)
- [593] J. Gross, "On Oblique Projection, Rank Additivity and the Moore-Penrose Inverse of the Sum of Two Matrices," *Lin. Multilin. Alg.*, Vol. 46, pp. 265–275, 1999. (Cited on page 381.)
- [594] J. Gross, "On the Product of Orthogonal Projectors," *Lin. Alg. Appl.*, Vol. 289, pp. 141–150, 1999. (Cited on pages 209 and 381.)
- [595] J. Gross, "The Moore-Penrose Inverse of a Partitioned Nonnegative Definite Matrix," *Lin. Alg. Appl.*, Vol. 321, pp. 113–121, 2000. (Cited on page 391.)
- [596] J. Gross and G. Trenkler, "On the Product of Oblique Projectors," *Lin. Multilin. Alg.*, Vol. 44, pp. 247–259, 1998. (Cited on page 204.)
- [597] J. Gross and G. Trenkler, "Nonsingularity of the Difference of Two Oblique Projectors," *SIAM J. Matrix Anal. Appl.*, Vol. 21, pp. 390–395, 1999. (Cited on pages 202, 203, and 204.)
- [598] J. Gross and G. Trenkler, "Product and Sum of Projections," *Amer. Math. Monthly*, Vol. 111, pp. 261–262, 2004. (Cited on page 335.)
- [599] J. Gross, G. Trenkler, and S.-O. Troschke, "The Vector Cross Product in $\mathbb{C}$," *Int. J. Math. Educ. Sci. Tech.*, Vol. 30, pp. 549–555, 1999. (Cited on page 188.)
- [600] J. Gross, G. Trenkler, and S.-O. Troschke, "Quaternions: Further Contributions to a Matrix Oriented Approach," *Lin. Alg. Appl.*, Vol. 326, pp. 205–213, 2001. (Cited on page 229.)
- [601] J. Gross and S.-O. Troschke, "Some Remarks on Partial Orderings of Nonnegative Definite Matrices," *Lin. Alg. Appl.*, Vol. 264, pp. 457–461, 1997. (Cited on pages 524 and 526.)
- [602] J. Gross and J. Yellen, *Graph Theory and Its Applications*, 2nd ed. Boca Raton: Chapman & Hall/CRC, 2005. (Cited on page xvii.)
- [603] L. C. Grove and C. T. Benson, *Finite Reflection Groups*, 2nd ed. New York: Springer, 1996. (Cited on page 223.)
- [604] K. Guan and H. Zhu, "A Generalized Heronian Mean and Its Inequalities," *University Beogradu Publ. Elektrotehn. Fak.*, Vol. 17, pp. 1–16, 2006, <http://pefm.math.etf.bg.ac.yu/to%20appear/rad528.pdf>. (Cited on page 38.)
- [605] S. Gull, A. Lasenby, and C. Doran, "Imaginary Numbers Are not Real: The Geometric Algebra of Spacetime," *Found. Phys.*, Vol. 23, pp. 1175–1201, 1993. (Cited on pages 188, 227, and 565.)
- [606] A. K. Gupta and D. K. Nagar, *Matrix Variate Distributions*. Boca Raton: CRC, 1999. (Cited on page xvii.)
- [607] K. Gurlebeck and W. Sprossig, *Quaternionic and Clifford Calculus for Physicists and Engineers*. New York: Chichester, 1997. (Cited on pages 227, 228, and 675.)
- [608] L. Gurvits, "Stability of Discrete Linear Inclusion," *Lin. Alg. Appl.*, Vol. 231, pp. 47–85, 1995. (Cited on page 715.)

- [609] K. E. Gustafson, "Matrix Trigonometry," *Lin. Alg. Appl.*, Vol. 217, pp. 117–140, 1995. (Cited on page 577.)
- [610] K. E. Gustafson and D. K. M. Rao, *Numerical Range*. New York: Springer, 1997. (Cited on page 497.)
- [611] W. H. Gustafson, P. R. Halmos, and H. Radjavi, "Products of Involutions," *Lin. Alg. Appl.*, Vol. 13, pp. 157–162, 1976. (Cited on page 351.)
- [612] P. W. Gwanyama, "The HM-GM-QM-QM Inequalities," *College Math. J.*, Vol. 35, pp. 47–50, 2004. (Cited on page 52.)
- [613] W. M. Haddad and D. S. Bernstein, "Robust Stabilization with Positive Real Uncertainty: Beyond the Small Gain Theorem," *Sys. Contr. Lett.*, Vol. 17, pp. 191–208, 1991. (Cited on page 491.)
- [614] W. M. Haddad and D. S. Bernstein, "Controller Design with Regional Pole Constraints," *IEEE Trans. Autom. Contr.*, Vol. 37, pp. 54–69, 1992. (Cited on page 798.)
- [615] W. M. Haddad and V. Chellaboina, "Stability and Dissipativity Theory for Non-negative Dynamical Systems: A Unified Analysis Framework for Biological and Physiological Systems," *Nonlinear Anal. Real World Appl.*, Vol. 6, pp. 35–65, 2005. (Cited on page 707.)
- [616] W. M. Haddad and V. Chellaboina, *Nonlinear Dynamical Systems and Control*. Princeton: Princeton University Press, 2008. (Cited on page xvii.)
- [617] W. M. Haddad, V. Chellaboina, and S. G. Nersesov, *Thermodynamics: A Dynamical Systems Approach*. Princeton: Princeton University Press, 2005. (Cited on pages 230 and 708.)
- [618] W. M. Haddad, V. Kapila, and E. G. Collins, "Optimality Conditions for Reduced-Order Modeling, Estimation, and Control for Discrete-Time Linear Periodic Plants," *J. Math. Sys. Est. Contr.*, Vol. 6, pp. 437–460, 1996. (Cited on page 805.)
- [619] W. W. Hager, "Updating the Inverse of a Matrix," *SIAM Rev.*, Vol. 31, pp. 221–239, 1989. (Cited on page 164.)
- [620] W. Hahn, *Stability of Motion*. Berlin: Springer, 1967. (Cited on page 722.)
- [621] E. Hairer, C. Lubich, and G. Wanner, *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*. Berlin: Springer, 2002. (Cited on page 678.)
- [622] M. Hajja, "A Method for Establishing Certain Trigonometric Inequalities," *J. Ineq. Pure Appl. Math.*, Vol. 8, no. 1, pp. 1–11, 2007, Article 29. (Cited on page 158.)
- [623] A. Hall, "Conditions for a Matrix Kronecker Lemma," *Lin. Alg. Appl.*, Vol. 76, pp. 271–277, 1986. (Cited on page 465.)
- [624] B. C. Hall, *Lie Groups, Lie Algebras, and Representations: An Elementary Introduction*. New York: Springer, 2003. (Cited on pages 217, 654, 655, 657, 658, 659, 660, 680, 684, and 722.)
- [625] G. T. Halliwell and P. R. Mercer, "A Refinement of an Inequality from Information Theory," *J. Ineq. Pure Appl. Math.*, Vol. 5, no. 1, pp. 1–3, 2004, Article 3. (Cited on pages 27 and 67.)

- [626] P. R. Halmos, *Finite-Dimensional Vector Spaces*. Princeton: Van Nostrand, 1958, reprinted by Springer, New York, 1974. (Cited on pages 313, 340, and 376.)
- [627] P. R. Halmos, *A Hilbert Space Problem Book*, 2nd ed. New York: Springer, 1982. (Cited on pages 318, 343, 384, and 385.)
- [628] P. R. Halmos, “Bad Products of Good Matrices,” *Lin. Multilin. Alg.*, Vol. 29, pp. 1–20, 1991. (Cited on pages 349 and 350.)
- [629] P. R. Halmos, *Problems for Mathematicians Young and Old*. Washington, DC: Mathematical Association of America, 1991. (Cited on pages 113, 279, 451, and 681.)
- [630] P. R. Halmos, *Linear Algebra Problem Book*. Washington, DC: Mathematical Association of America, 1995. (Cited on pages 90, 202, and 343.)
- [631] M. Hamermesh, *Group Theory and its Application to Physical Problems*. Reading: Addison-Wesley, 1962, reprinted by Dover, Mineola, 1989. (Cited on page 224.)
- [632] J. H. Han and M. D. Hirschhorn, “Another Look at an Amazing Identity of Ramanujan,” *Math. Mag.*, Vol. 79, pp. 302–304, 2006. (Cited on page 128.)
- [633] E. J. Hannan and M. Deistler, *The Statistical Theory of Linear Systems*. New York: Wiley, 1988. (Cited on page 805.)
- [634] F. Hansen, “Extrema for Concave Operator Mappings,” *Math. Japonica*, Vol. 40, pp. 331–338, 1994. (Cited on page 483.)
- [635] F. Hansen, “Operator Inequalities Associated with Jensen’s Inequality,” in *Survey on Classical Inequalities*, T. M. Rassias, Ed. Dordrecht: Kluwer, 2000, pp. 67–98. (Cited on page 483.)
- [636] A. J. Hanson, *Visualizing Quaternions*. Amsterdam: Elsevier, 2006. (Cited on pages 227 and 228.)
- [637] G. Hardy, J. E. Littlewood, and G. Polya, *Inequalities*. Cambridge: Cambridge University Press, 1988. (Cited on page 76.)
- [638] L. A. Harris, “Factorizations of Operator Matrices,” *Lin. Alg. Appl.*, Vol. 225, pp. 37–41, 1995. (Cited on pages 473 and 474.)
- [639] L. A. Harris, “The Inverse of a Block Matrix,” *Amer. Math. Monthly*, Vol. 102, pp. 656–657, 1995. (Cited on page 148.)
- [640] W. A. Harris, J. P. Fillmore, and D. R. Smith, “Matrix Exponentials—Another Approach,” *SIAM Rev.*, Vol. 43, pp. 694–706, 2001. (Cited on page 673.)
- [641] G. W. Hart, *Multidimensional Analysis: Algebras and Systems for Science and Engineering*. New York: Springer, 1995. (Cited on pages xvii and 230.)
- [642] D. J. Hartfiel, *Nonhomogeneous Matrix Products*. Singapore: World Scientific, 2002. (Cited on pages xvii and 715.)
- [643] R. Hartwig, X. Li, and Y. Wei, “Representations for the Drazin Inverse of a 2×2 Block Matrix,” *SIAM J. Matrix Anal. Appl.*, Vol. 27, pp. 757–771, 2006. (Cited on page 391.)
- [644] R. E. Hartwig, “Resultants and the Solutions of $AX - XB = -C$,” *SIAM J. Appl. Math.*, Vol. 23, pp. 104–117, 1972. (Cited on page 797.)

- [645] R. E. Hartwig, "Block Generalized Inverses," *Arch. Rat. Mech. Anal.*, Vol. 61, pp. 197–251, 1976. (Cited on page 391.)
- [646] R. E. Hartwig, "A Note on the Partial Ordering of Positive Semi-Definite Matrices," *Lin. Multilin. Alg.*, Vol. 6, pp. 223–226, 1978. (Cited on pages 524, 526, and 527.)
- [647] R. E. Hartwig, "How to Partially Order Regular Elements," *Math. Japonica*, Vol. 25, pp. 1–13, 1980. (Cited on page 119.)
- [648] R. E. Hartwig and I. J. Katz, "On Products of EP Matrices," *Lin. Alg. Appl.*, Vol. 252, pp. 339–345, 1997. (Cited on page 382.)
- [649] R. E. Hartwig and M. S. Putcha, "When Is a Matrix a Difference of Two Idempotents?" *Lin. Multilin. Alg.*, Vol. 26, pp. 267–277, 1990. (Cited on pages 322 and 337.)
- [650] R. E. Hartwig and M. S. Putcha, "When Is a Matrix a Sum of Idempotents?" *Lin. Multilin. Alg.*, Vol. 26, pp. 279–286, 1990. (Cited on page 361.)
- [651] R. E. Hartwig and K. Spindelbock, "Matrices for which A^* and A^+ Commute," *Lin. Multilin. Alg.*, Vol. 14, pp. 241–256, 1984. (Cited on pages 315, 372, 373, and 396.)
- [652] R. E. Hartwig and G. P. H. Styan, "On Some Characterizations of the "Star" Partial Ordering for Matrices and Rank Subtractivity," *Lin. Alg. Appl.*, Vol. 82, pp. 145–161, 1989. (Cited on pages 119, 120, 385, and 526.)
- [653] R. E. Hartwig, G. Wang, and Y. Wei, "Some Additive Results on Drazin Inverse," *Lin. Alg. Appl.*, Vol. 322, pp. 207–217, 2001. (Cited on page 394.)
- [654] D. A. Harville, *Matrix Algebra from a Statistician's Perspective*. New York: Springer, 1997. (Cited on pages xvii, 199, 202, 371, 377, 378, 381, 383, 486, 506, and 642.)
- [655] J. Hauke and A. Markiewicz, "On Partial Orderings on the Set of Rectangular Matrices," *Lin. Alg. Appl.*, Vol. 219, pp. 187–193, 1995. (Cited on page 524.)
- [656] M. L. J. Hautus, "Controllability and Observability Conditions of Linear Autonomous Systems," *Proc. Koninklijke Akademik Van Wetenschappen*, Vol. 72, pp. 443–448, 1969. (Cited on page 805.)
- [657] T. Haynes, "Stable Matrices, the Cayley Transform, and Convergent Matrices," *Int. J. Math. Math. Sci.*, Vol. 14, pp. 77–81, 1991. (Cited on page 713.)
- [658] E. V. Haynsworth, "Applications of an Inequality for the Schur Complement," *Proc. Amer. Math. Soc.*, Vol. 24, pp. 512–516, 1970. (Cited on pages 442, 474, and 542.)
- [659] E. Hecht, *Optics*, 4th ed. Reading: Addison Wesley, 2002. (Cited on page xvii.)
- [660] C. Heij, A. Ran, and F. van Schagen, Eds., *Introduction to Mathematical Systems Theory: Linear Systems, Identification and Control*. Basel: Birkhauser, 2007. (Cited on page 805.)
- [661] C. Heij, A. Ran, and F. van Schagen, *Introduction to Mathematical Systems Theory: Linear Systems, Identification and Control*. Basel: Birkhauser, 2007. (Cited on page 805.)

- [662] G. Heinig, "Matrix Representations of Bezoutians," *Lin. Alg. Appl.*, Vol. 223/224, pp. 337–354, 1995. (Cited on page 257.)
- [663] U. Helmke and P. A. Fuhrmann, "Bezoutians," *Lin. Alg. Appl.*, Vol. 122–124, pp. 1039–1097, 1989. (Cited on page 257.)
- [664] B. W. Helton, "Logarithms of Matrices," *Proc. Amer. Math. Soc.*, Vol. 19, pp. 733–738, 1968. (Cited on page 722.)
- [665] H. V. Henderson, F. Pukelsheim, and S. R. Searle, "On the History of the Kronecker Product," *Lin. Multilin. Alg.*, Vol. 14, pp. 113–120, 1983. (Cited on page 416.)
- [666] H. V. Henderson and S. R. Searle, "On Deriving the Inverse of a Sum of Matrices," *SIAM Rev.*, Vol. 23, pp. 53–60, 1981. (Cited on pages 145 and 164.)
- [667] H. V. Henderson and S. R. Searle, "The Vec-Permutation Matrix, The Vec Operator and Kronecker Products: A Review," *Lin. Multilin. Alg.*, Vol. 9, pp. 271–288, 1981. (Cited on page 416.)
- [668] J. Herman, R. Kucera, and J. Simsa, *Equations and Inequalities*. New York: Springer, 2000. (Cited on pages 17, 18, 23, 24, 25, 26, 32, 36, 37, 39, 40, 41, 42, 43, 44, 46, 47, 49, 52, and 60.)
- [669] D. Hershkwitz, "Positive Semidefinite Pattern Decompositions," *SIAM J. Matrix Anal. Appl.*, Vol. 11, pp. 612–619, 1990. (Cited on page 475.)
- [670] D. Hestenes, *Space-Time Algebra*. New York: Gordon and Breach, 1966. (Cited on page 227.)
- [671] D. Hestenes, *New Foundations for Classical Mechanics*. Dordrecht: Kluwer, 1986. (Cited on pages 188 and 227.)
- [672] D. Hestenes and G. Sobczyk, *Clifford Algebra to Geometric Calculus: A Unified Language for Mathematics and Physics*. Dordrecht: D. Riedel, 1984. (Cited on pages 188 and 227.)
- [673] F. Hiai and H. Kosaki, Eds., *Means of Hilbert Space Operators*. Berlin: Springer, 2003. (Cited on pages 29, 35, and 447.)
- [674] F. Hiai and D. Petz, "The Golden-Thompson Trace Inequality Is Complemented," *Lin. Alg. Appl.*, Vol. 181, pp. 153–185, 1993. (Cited on pages 461 and 686.)
- [675] F. Hiai and X. Zhan, "Submultiplicativity vs Subadditivity for Unitarily Invariant Norms," *Lin. Alg. Appl.*, Vol. 377, pp. 155–164, 2004. (Cited on page 580.)
- [676] D. J. Higham and N. J. Higham, *Matlab Guide*, 2nd ed. Philadelphia: SIAM, 2005. (Cited on page 164.)
- [677] N. J. Higham, "Newton's Method for the Matrix Square Root," *Math. Computation*, Vol. 46, pp. 537–549, 1986. (Cited on page 349.)
- [678] N. J. Higham, "Computing Real Square Roots of a Real Matrix," *Lin. Alg. Appl.*, Vol. 88/89, pp. 405–430, 1987. (Cited on page 348.)
- [679] N. J. Higham, "Matrix Nearness Problems and Applications," in *Applications of Matrix Theory*, M. J. C. Gover and S. Barnett, Eds. Oxford: Oxford University Press, 1989, pp. 1–27. (Cited on pages 608, 700, and 701.)
- [680] N. J. Higham, "Estimating the Matrix p -Norm," *Numer. Math.*, Vol. 62, pp. 539–555, 1992. (Cited on pages 571 and 575.)

- [681] N. J. Higham, *Accuracy and Stability of Numerical Algorithms*, 2nd ed. Philadelphia: SIAM, 2002. (Cited on pages xviii, 215, 221, 327, 328, 331, 571, 573, 575, 576, 577, 603, 608, 609, 615, and 619.)
- [682] N. J. Higham, "Cayley, Sylvester, and Early Matrix Theory," *Lin. Alg. Appl.*, Vol. 428, pp. 39–43, 2008. (Cited on page 261.)
- [683] N. J. Higham, *Functions of Matrices: Theory and Computation*. Philadelphia: SIAM, 2008. (Cited on pages xviii, 72, 74, 348, 359, 360, 584, 629, 637, 657, 658, 677, 681, 685, 692, 693, 694, 722, and 802.)
- [684] G. N. Hile and P. Lounesto, "Matrix Representations of Clifford Algebras," *Lin. Alg. Appl.*, Vol. 128, pp. 51–63, 1990. (Cited on page 227.)
- [685] R. D. Hill, R. G. Bates, and S. R. Waters, "On Centrohermitian Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 11, pp. 128–133, 1990. (Cited on page 221.)
- [686] R. D. Hill and E. E. Underwood, "On the Matrix Adjoint (Adjugate)," *SIAM J. Alg. Disc. Meth.*, Vol. 6, pp. 731–737, 1985. (Cited on page 142.)
- [687] C.-J. Hillar, "Advances on the Bessis-Moussa-Villani Trace Conjecture," *Lin. Alg. Appl.*, Vol. 426, pp. 130–142, 2007. (Cited on page 482.)
- [688] L. O. Hilliard, "The Case of Equality in Hopf's Inequality," *SIAM J. Alg. Disc. Meth.*, Vol. 8, pp. 691–709, 1987. (Cited on page 280.)
- [689] D. Hinrichsen, E. Plischke, and F. Wirth, "Robustness of Transient Behavior," in *Unsolved Problems in Mathematical Systems and Control Theory*, V. D. Blondel and A. Megretski, Eds. Princeton: Princeton University Press, 2004. (Cited on page 699.)
- [690] D. Hinrichsen and A. J. Pritchard, *Mathematical Systems Theory I: Modelling, State Space Analysis, Stability and Robustness*. Berlin: Springer, 2005. (Cited on pages 318, 326, 327, 554, 555, 617, 638, 639, 691, 695, and 709.)
- [691] M. W. Hirsch and S. Smale, *Differential Equations, Dynamical Systems, and Linear Algebra*. San Diego: Academic Press, 1974. (Cited on pages xvii and 311.)
- [692] M. W. Hirsch, S. Smale, and R. L. Devaney, *Differential Equations, Dynamical Systems and an Introduction to Chaos*, 2nd ed. New York: Elsevier, 2003. (Cited on page xvii.)
- [693] O. Hirzallah, "Inequalities for Sums and Products of Operators," *Lin. Alg. Appl.*, Vol. 407, pp. 32–42, 2005. (Cited on page 613.)
- [694] O. Hirzallah and F. Kittaneh, "Matrix Young Inequalities for the Hilbert-Schmidt Norm," *Lin. Alg. Appl.*, Vol. 308, pp. 77–84, 2000. (Cited on page 612.)
- [695] O. Hirzallah and F. Kittaneh, "Commutator Inequalities for Hilbert-Schmidt Norm," *J. Math. Anal. Appl.*, Vol. 268, pp. 67–73, 2002. (Cited on pages 38 and 70.)
- [696] O. Hirzallah and F. Kittaneh, "Non-Commutative Clarkson Inequalities for Unitarily Invariant Norms," *Pac. J. Math.*, Vol. 202, pp. 363–369, 2002. (Cited on page 586.)
- [697] O. Hirzallah and F. Kittaneh, "Norm Inequalities for Weighted Power Means of Operators," *Lin. Alg. Appl.*, Vol. 341, pp. 181–193, 2002. (Cited on page 502.)

- [698] O. Hirzallah and F. Kittaneh, "Inequalities for Sums and Direct Sums of Hilbert Space Operators," *Lin. Alg. Appl.*, Vol. 424, pp. 71–82, 2007. (Cited on pages 612 and 613.)
- [699] A. Hmamed, "A Matrix Inequality," *Int. J. Contr.*, Vol. 49, pp. 363–365, 1989. (Cited on page 541.)
- [700] J. B. Hoagg, J. Chandrasekar, and D. S. Bernstein, "On the Zeros, Initial Under-shoot, and Relative Degree of Lumped-Parameter Structures," *ASME J. Dyn. Sys. Meas. Contr.*, Vol. 129, pp. 493–502, 2007. (Cited on page 219.)
- [701] K. Hoffman and R. Kunze, *Linear Algebra*, 2nd ed. Englewood Cliffs: Prentice-Hall, 1971. (Cited on page xix.)
- [702] L. Hogben, Ed., *Handbook of Linear Algebra*. Boca Raton: Chapman & Hall/CRC, 2007. (Cited on page xvii.)
- [703] R. R. Holmes and T. Y. Tam, "Group Representations," in *Handbook of Linear Algebra*, L. Hogben, Ed. Boca Raton: Chapman & Hall/CRC, 2007, pp. 68–1–68–11. (Cited on page 224.)
- [704] O. Holtz, "On Convergence of Infinite Matrix Products," *Elec. J. Lin. Alg.*, Vol. 7, pp. 178–181, 2000. (Cited on page 715.)
- [705] O. Holtz, "Hermite-Biehler, Routh-Hurwitz, and Total Positivity," *Lin. Alg. Appl.*, Vol. 372, pp. 105–110, 2003. (Cited on page 697.)
- [706] Y. Hong and R. A. Horn, "The Jordan Canonical Form of a Product of a Hermitian and a Positive Semidefinite Matrix," *Lin. Alg. Appl.*, Vol. 147, pp. 373–386, 1991. (Cited on page 350.)
- [707] Y. P. Hong and C.-T. Pan, "A Lower Bound for the Smallest Singular Value," *Lin. Alg. Appl.*, Vol. 172, pp. 27–32, 1992. (Cited on page 606.)
- [708] A. Horn, "Doubly Stochastic Matrices and the Diagonal of a Rotation Matrix," *Amer. J. Math.*, Vol. 76, pp. 620–630, 1954. (Cited on page 511.)
- [709] R. A. Horn and C. R. Johnson, *Matrix Analysis*. Cambridge: Cambridge University Press, 1985. (Cited on pages 139, 163, 214, 254, 271, 273, 275, 280, 281, 293, 314, 320, 342, 345, 346, 358, 360, 412, 428, 431, 446, 474, 492, 505, 507, 510, 511, 512, 532, 537, 538, 541, 546, 549, 550, 553, 572, 573, 574, 578, 580, 601, and 609.)
- [710] R. A. Horn and C. R. Johnson, "Hadamard and Conventional Submultiplicativity for Unitarily Invariant Norms on Matrices," *Lin. Multilin. Alg.*, Vol. 20, pp. 91–106, 1987. (Cited on page 578.)
- [711] R. A. Horn and C. R. Johnson, *Topics in Matrix Analysis*. Cambridge: Cambridge University Press, 1991. (Cited on pages 163, 197, 276, 326, 327, 341, 342, 348, 405, 407, 416, 431, 443, 470, 497, 505, 509, 561, 562, 578, 592, 603, 605, 606, 612, 615, 641, 642, 654, 657, 678, 689, 692, 693, 703, 707, and 797.)
- [712] R. A. Horn and R. Mathias, "An Analog of the Cauchy-Schwarz Inequality for Hadamard Products and Unitarily Invariant Norms," *SIAM J. Matrix Anal. Appl.*, Vol. 11, pp. 481–498, 1990. (Cited on pages 591, 603, and 617.)
- [713] R. A. Horn and R. Mathias, "Cauchy-Schwarz Inequalities Associated with Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 142, pp. 63–82, 1990. (Cited on pages 539, 576, and 581.)

- [714] R. A. Horn and R. Mathias, "Block-Matrix Generalizations of Schur's Basic Theorems on Hadamard Products," *Lin. Alg. Appl.*, Vol. 172, pp. 337–346, 1992. (Cited on page 416.)
- [715] R. A. Horn and I. Olkin, "When Does $A^*A = B^*B$ and Why Does One Want to Know?" *Amer. Math. Monthly*, Vol. 103, pp. 470–482, 1996. (Cited on page 319.)
- [716] R. A. Horn and G. G. Piepmeyer, "Two Applications of the Theory of Primary Matrix Functions," *Lin. Alg. Appl.*, Vol. 361, pp. 99–106, 2003. (Cited on page 642.)
- [717] R. A. Horn and F. Zhang, "Basic Properties of the Schur Complement," in *The Schur Complement and Its Applications*, F. Zhang, Ed. New York: Springer, 2004, pp. 17–46. (Cited on pages 333, 393, 461, 490, and 493.)
- [718] B. G. Horne, "Lower Bounds for the Spectral Radius of a Matrix," *Lin. Alg. Appl.*, Vol. 263, pp. 261–273, 1997. (Cited on page 604.)
- [719] H.-C. Hou and H.-K. Du, "Norm Inequalities of Positive Operator Matrices," *Integ. Eqns. Operator Theory*, Vol. 22, pp. 281–294, 1995. (Cited on pages 469, 521, and 610.)
- [720] S.-H. Hou, "A Simple Proof of the Leverrier-Faddeev Characteristic Polynomial Algorithm," *SIAM Rev.*, Vol. 40, pp. 706–709, 1998. (Cited on page 281.)
- [721] A. S. Householder, *The Theory of Matrices in Numerical Analysis*. New York: Blaisdell, 1964, reprinted by Dover, New York, 1975. (Cited on pages xviii, 378, 611, and 634.)
- [722] A. S. Householder, "Bezoutiants, Elimination and Localization," *SIAM Rev.*, Vol. 12, pp. 73–78, 1970. (Cited on page 257.)
- [723] A. S. Householder and J. A. Carpenter, "The Singular Values of Involutory and of Idempotent Matrices," *Numer. Math.*, Vol. 5, pp. 234–237, 1963. (Cited on page 329.)
- [724] R. Howe, "Very Basic Lie Theory," *Amer. Math. Monthly*, Vol. 90, pp. 600–623, 1983. (Cited on pages 229 and 722.)
- [725] J. M. Howie, *Complex Analysis*. New York: Springer, 2003. (Cited on page 76.)
- [726] R. A. Howland, *Intermediate Dynamics: A Linear Algebraic Approach*. New York: Springer, 2006. (Cited on page xvii.)
- [727] P.-F. Hsieh and Y. Sibuya, *Basic Theory of Ordinary Differential Equations*. New York: Springer, 1999. (Cited on pages xvii and 311.)
- [728] G.-D. Hu and G.-H. Hu, "A Relation between the Weighted Logarithmic Norm of a Matrix and the Lyapunov Equation," *Numer. Math. BIT*, Vol. 40, pp. 606–610, 2000. (Cited on page 700.)
- [729] S. Hu-yun, "Estimation of the Eigenvalues of AB for $A > 0$ and $B > 0$," *Lin. Alg. Appl.*, Vol. 73, pp. 147–150, 1986. (Cited on pages 127 and 519.)
- [730] R. Huang, "Some Inequalities for Hadamard Products of Matrices," *Lin. Multilin. Alg.*, Vol. 56, pp. 543–554, 2008. (Cited on pages 603, 616, and 617.)
- [731] R. Huang, "Some Inequalities for the Hadamard Product and the Fan Product of Matrices," *Lin. Alg. Appl.*, Vol. 428, pp. 1551–1559, 2008. (Cited on pages 415 and 416.)

- [732] T.-Z. Huang and L. Wang, "Improving Bounds for Eigenvalues of Complex Matrices Using Traces," *Lin. Alg. Appl.*, Vol. 426, pp. 841–854, 2007. (Cited on pages 51 and 598.)
- [733] P. C. Hughes, *Spacecraft Attitude Dynamics*. New York: Wiley, 1986. (Cited on pages xvii and 676.)
- [734] M. Huhtanen, "Real Linear Kronecker Product Operations," *Lin. Alg. Appl.*, Vol. 418, pp. 347–361, 2006. (Cited on page 70.)
- [735] S. Humphries and C. Krattenthaler, "Trace Identities from Identities for Determinants," *Lin. Alg. Appl.*, Vol. 411, pp. 328–342, 2005. (Cited on page 132.)
- [736] C. H. Hung and T. L. Markham, "The Moore-Penrose Inverse of a Partitioned Matrix," *Lin. Alg. Appl.*, Vol. 11, pp. 73–86, 1975. (Cited on page 391.)
- [737] J. J. Hunter, "Generalized Inverses and Their Application to Applied Probability Problems," *Lin. Alg. Appl.*, Vol. 45, pp. 157–198, 1982. (Cited on page 398.)
- [738] D. C. Hyland and D. S. Bernstein, "The Optimal Projection Equations for Fixed-Order Dynamic Compensation," *IEEE Trans. Autom. Contr.*, Vol. AC-29, pp. 1034–1037, 1984. (Cited on page 805.)
- [739] D. C. Hyland and E. G. Collins, "Block Kronecker Products and Block Norm Matrices in Large-Scale Systems Analysis," *SIAM J. Matrix Anal. Appl.*, Vol. 10, pp. 18–29, 1989. (Cited on page 416.)
- [740] N. H. Ibragimov, *Elementary Lie Group Analysis and Ordinary Differential Equations*. Chichester: Wiley, 1999. (Cited on page 722.)
- [741] Y. Ikebe and T. Inagaki, "An Elementary Approach to the Functional Calculus for Matrices," *Amer. Math. Monthly*, Vol. 93, pp. 390–392, 1986. (Cited on page 642.)
- [742] K. D. Ikramov, "A Simple Proof of the Generalized Schur Inequality," *Lin. Alg. Appl.*, Vol. 199, pp. 143–149, 1994. (Cited on pages 597 and 598.)
- [743] V. Ionescu, C. Oar, and M. Weiss, *Generalized Riccati Theory and Robust Control*. Chichester: Wiley, 1999. (Cited on pages xvii and 805.)
- [744] I. Ipsen and C. Meyer, "The Angle between Complementary Subspaces," *Amer. Math. Monthly*, Vol. 102, pp. 904–911, 1995. (Cited on pages 112, 210, 329, 336, and 381.)
- [745] A. Iserles, "Solving Linear Ordinary Differential Equations by Exponentials of Iterated Commutators," *Numer. Math.*, Vol. 45, pp. 183–199, 1984. (Cited on page 678.)
- [746] A. Iserles, H. Z. Munthe-Kaas, S. P. Norsett, and A. Zanna, "Lie-Group Methods," *Acta Numerica*, Vol. 9, pp. 215–365, 2000. (Cited on pages 188, 674, 678, and 682.)
- [747] Y. Ito, S. Hattori, and H. Maeda, "On the Decomposition of a Matrix into the Sum of Stable Matrices," *Lin. Alg. Appl.*, Vol. 297, pp. 177–182, 1999. (Cited on page 704.)
- [748] S. Izumino, H. Mori, and Y. Seo, "On Ozeki's Inequality," *J. Ineq. Appl.*, Vol. 2, pp. 235–253, 1998. (Cited on pages 65 and 501.)
- [749] D. H. Jacobson, D. H. Martin, M. Pachter, and T. Geveci, *Extensions of Linear Quadratic Control Theory*. Berlin: Springer, 1980. (Cited on page 805.)

- [750] A. Jeffrey, *Handbook of Mathematical Formulas and Integrals*. San Diego: Academic Press, 1995. (Cited on pages 74, 75, and 76.)
- [751] C. Jeffries, V. Klee, and P. van der Driessche, "Quality Stability of Linear Systems," *Lin. Alg. Appl.*, Vol. 87, pp. 1–48, 1987. (Cited on page 708.)
- [752] A. Jennings and J. J. McKeown, *Matrix Computation*, 2nd ed. New York: Wiley, 1992. (Cited on page xviii.)
- [753] G. A. Jennings, *Modern Geometry with Applications*. New York: Springer, 1997. (Cited on page 161.)
- [754] S. T. Jensen and G. P. H. Styan, "Some Comments and a Bibliography on the Laguerre-Samuelson Inequality with Extensions and Applications in Statistics and Matrix Theory," in *Analytic and Geometric Inequalities and Applications*, T. M. Rassias and H. M. Srivastava, Eds. Dordrecht: Kluwer, 1999, pp. 151–181. (Cited on pages 51 and 454.)
- [755] J. Ji, "Explicit Expressions of the Generalized Inverses and Condensed Cramer Rules," *Lin. Alg. Appl.*, Vol. 404, pp. 183–192, 2005. (Cited on page 129.)
- [756] G. Jia and J. Cao, "A New Upper Bound of the Logarithmic Mean," *J. Ineq. Pure Appl. Math.*, Vol. 4, no. 4, pp. 1–4, 2003, Article 80. (Cited on pages 27 and 38.)
- [757] Y. Jiang, W. W. Hager, and J. Li, "The Geometric Mean Decomposition," *Lin. Alg. Appl.*, Vol. 396, pp. 373–384, 2005. (Cited on page 316.)
- [758] D. Jocić and F. Kittaneh, "Some Perturbation Inequalities for Self-Adjoint Operators," *J. Operator Theory*, Vol. 31, pp. 3–10, 1994. (Cited on page 588.)
- [759] C. R. Johnson, "An Inequality for Matrices Whose Symmetric Part Is Positive Definite," *Lin. Alg. Appl.*, Vol. 6, pp. 13–18, 1973. (Cited on page 487.)
- [760] C. R. Johnson, "Inequalities for a Complex Matrix whose Real Part Is Positive Definite," *Trans. Amer. Math. Soc.*, Vol. 212, pp. 149–154, 1975. (Cited on page 485.)
- [761] C. R. Johnson, "The Inertia of a Product of Two Hermitian Matrices," *J. Math. Anal. Appl.*, Vol. 57, pp. 85–90, 1977. (Cited on page 309.)
- [762] C. R. Johnson, "Closure Properties of Certain Positivity Classes of Matrices under Various Algebraic Operations," *Lin. Alg. Appl.*, Vol. 97, pp. 243–247, 1987. (Cited on page 697.)
- [763] C. R. Johnson, "A Gersgorin-Type Lower Bound for the Smallest Singular Value," *Lin. Alg. Appl.*, Vol. 112, pp. 1–7, 1989. (Cited on page 606.)
- [764] C. R. Johnson, "A Gersgorin-Type Lower Bound for the Smallest Singular Value," *Lin. Alg. Appl.*, Vol. 112, pp. 1–7, 1989. (Cited on page 604.)
- [765] C. R. Johnson and L. Elsner, "The Relationship between Hadamard and Conventional Multiplication for Positive Definite Matrices," *Lin. Alg. Appl.*, Vol. 92, pp. 231–240, 1987. (Cited on page 534.)
- [766] C. R. Johnson, M. Neumann, and M. J. Tsatsomeros, "Conditions for the Positivity of Determinants," *Lin. Multilin. Alg.*, Vol. 40, pp. 241–248, 1996. (Cited on page 219.)
- [767] C. R. Johnson and M. Newman, "A Surprising Determinantal Inequality for Real Matrices," *Math. Ann.*, Vol. 247, pp. 179–186, 1980. (Cited on page 132.)

- [768] C. R. Johnson and P. Nylén, “Monotonicity Properties of Norms,” *Lin. Alg. Appl.*, Vol. 148, pp. 43–58, 1991. (Cited on page 619.)
- [769] C. R. Johnson, K. Okubo, and R. Beams, “Uniqueness of Matrix Square Roots,” *Lin. Alg. Appl.*, Vol. 323, pp. 51–60, 2001. (Cited on page 348.)
- [770] C. R. Johnson and L. Rodman, “Convex Sets of Hermitian Matrices with Constant Inertia,” *SIAM J. Alg. Disc. Meth.*, Vol. 6, pp. 351–359, 1985. (Cited on pages 308 and 310.)
- [771] C. R. Johnson and R. Schreiner, “The Relationship between AB and BA ,” *Amer. Math. Monthly*, Vol. 103, pp. 578–582, 1996. (Cited on page 346.)
- [772] C. R. Johnson and H. M. Shapiro, “Mathematical Aspects of the Relative Gain Array $A \circ A^{-T}$,” *SIAM J. Alg. Disc. Meth.*, Vol. 7, pp. 627–644, 1986. (Cited on pages 415 and 532.)
- [773] C. R. Johnson and R. L. Smith, “Linear Interpolation Problems for Matrix Classes and a Transformational Characterization of M-Matrices,” *Lin. Alg. Appl.*, Vol. 330, pp. 43–48, 2001. (Cited on pages 185, 191, and 674.)
- [774] C. R. Johnson and T. Szulc, “Further Lower Bounds for the Smallest Singular Value,” *Lin. Alg. Appl.*, Vol. 272, pp. 169–179, 1998. (Cited on page 604.)
- [775] C. R. Johnson and F. Zhang, “An Operator Inequality and Matrix Normality,” *Lin. Alg. Appl.*, Vol. 240, pp. 105–110, 1996. (Cited on page 502.)
- [776] M. Jolly, “On the Calculus of Complex Matrices,” *Int. J. Contr.*, Vol. 61, pp. 749–755, 1995. (Cited on page 642.)
- [777] T. F. Jordan, *Quantum Mechanics in Simple Matrix Form*. New York: Wiley, 1986, reprinted by Dover, Mineola, 2005. (Cited on page 228.)
- [778] E. A. Jorswieck and H. Boche, “Majorization and Matrix Monotone Functions in Wireless Communications,” *Found. Trends Comm. Inform. Theory*, Vol. 3, pp. 553–701, 2006. (Cited on page xvii.)
- [779] E. A. Jorswieck and H. Boche, “Performance Analysis of MIMO Systems in Spatially Correlated Fading Using Matrix-Monotone Functions,” *IEICE Trans. Fund.*, Vol. E89-A, pp. 1454–1472, 2006. (Cited on pages xvii and 162.)
- [780] A. Joseph, A. Melnikov, and R. Rentschler, Eds., *Studies in Memory of Issai Schur*. Boston: Birkhauser, 2002. (Cited on page 542.)
- [781] D. Joyner, *Adventures in Group Theory*. Baltimore: Johns Hopkins University Press, 2002. (Cited on page 230.)
- [782] E. I. Jury, *Inners and Stability of Dynamic Systems*, 2nd ed. Malabar: Krieger, 1982. (Cited on pages 413, 416, and 709.)
- [783] W. J. Kaczor and M. T. Nowak, *Problems in Mathematical Analysis II*. American Mathematical Society, 2001. (Cited on pages 24, 27, 29, 34, and 66.)
- [784] R. V. Kadison, “Order Properties of Bounded Self-Adjoint Operators,” *Proc. Amer. Math. Soc.*, Vol. 2, pp. 505–510, 1951. (Cited on page 459.)
- [785] A. Kagan and P. J. Smith, “A Stronger Version of Matrix Convexity as Applied to Functions of Hermitian Matrices,” *J. Ineq. Appl.*, Vol. 3, pp. 143–152, 1999. (Cited on page 483.)

- [786] J. B. Kagstrom, "Bounds and Perturbation Bounds for the Matrix Exponential," *Numer. Math. BIT*, Vol. 17, pp. 39–57, 1977. (Cited on page 699.)
- [787] T. Kailath, *Linear Systems*. Englewood Cliffs: Prentice-Hall, 1980. (Cited on pages 237, 281, 304, 353, and 805.)
- [788] D. Kalman and J. E. White, "Polynomial Equations and Circulant Matrices," *Amer. Math. Monthly*, Vol. 108, pp. 821–840, 2001. (Cited on page 357.)
- [789] T. R. Kane, P. W. Likins, and D. A. Levinson, *Spacecraft Dynamics*. New York: McGraw-Hill, 1983. (Cited on page xvii.)
- [790] I. Kaplansky, *Linear Algebra and Geometry: A Second Course*. New York: Chelsea, 1974, reprinted by Dover, Mineola, 2003. (Cited on pages xix and 318.)
- [791] N. Karcanias, "Matrix Pencil Approach to Geometric System Theory," *Proc. IEE*, Vol. 126, pp. 585–590, 1979. (Cited on pages 362 and 805.)
- [792] S. Karlin and F. Ost, "Some Monotonicity Properties of Schur Powers of Matrices and Related Inequalities," *Lin. Alg. Appl.*, Vol. 68, pp. 47–65, 1985. (Cited on page 415.)
- [793] E. Kaszkurewicz and A. Bhaya, *Matrix Diagonal Stability in Systems and Computation*. Boston: Birkhauser, 2000. (Cited on page 722.)
- [794] M. Kato, K.-S. Saito, and T. Tamura, "Sharp Triangle Inequality and Its Reverse in Banach Space," *Math. Ineq. Appl.*, Vol. 10, pp. 451–460, 2007. (Cited on page 568.)
- [795] T. Kato, "Spectral Order and a Matrix Limit Theorem," *Lin. Multilin. Alg.*, Vol. 8, pp. 15–19, 1979. (Cited on page 460.)
- [796] T. Kato, *Perturbation Theory for Linear Operators*. Berlin: Springer, 1980. (Cited on pages 600 and 619.)
- [797] H. Katsuura, "Generalizations of the Arithmetic-Geometric Mean Inequality and a Three Dimensional Puzzle," *College Math. J.*, Vol. 34, pp. 280–282, 2003. (Cited on pages 46, 53, and 54.)
- [798] M. Kauderer, *Symplectic Matrices: First Order Systems and Special Relativity*. Singapore: World Scientific, 1994. (Cited on page xvii.)
- [799] J. Y. Kazakia, "Orthogonal Transformation of a Trace Free Symmetric Matrix into One with Zero Diagonal Elements," *Int. J. Eng. Sci.*, Vol. 26, pp. 903–906, 1988. (Cited on page 313.)
- [800] N. D. Kazarinoff, *Analytic Inequalities*. New York: Holt, Rinehart, and Winston, 1961, reprinted by Dover, Mineola, 2003. (Cited on pages 36 and 546.)
- [801] M. G. Kendall, *A Course in the Geometry of n Dimensions*. London: Griffin, 1961, reprinted by Dover, Mineola, 2004. (Cited on page 184.)
- [802] C. Kenney and A. J. Laub, "Controllability and Stability Radii for Companion Form Systems," *Math. Contr. Sig. Sys.*, Vol. 1, pp. 239–256, 1988. (Cited on pages 327 and 711.)
- [803] C. Kenney and A. J. Laub, "Rational Iteration Methods for the Matrix Sign Function," *SIAM J. Matrix Anal. Appl.*, Vol. 12, pp. 273–291, 1991. (Cited on page 637.)
- [804] H. Kestelman, "Eigenvectors of a Cross-Diagonal Matrix," *Amer. Math. Monthly*, Vol. 93, p. 566, 1986. (Cited on page 340.)

- [805] N. Keyfitz, *Introduction to the Mathematics of Population*. Reading: Addison-Wesley, 1968. (Cited on pages xvii and 275.)
- [806] W. Khalil and E. Dombre, *Modeling, Identification, and Control of Robots*. New York: Taylor and Francis, 2002. (Cited on page xvii.)
- [807] C. G. Khatri, "A Note on Idempotent Matrices," *Lin. Alg. Appl.*, Vol. 70, pp. 185–195, 1985. (Cited on page 374.)
- [808] C. G. Khatri and S. K. Mitra, "Hermitian and Nonnegative Definite Solutions of Linear Matrix Equations," *SIAM J. Appl. Math.*, Vol. 31, pp. 579–585, 1976. (Cited on page 384.)
- [809] S. Kim and Y. Lim, "A Converse Inequality of Higher Order Weighted Arithmetic and Geometric Means of Positive Definite Operators," *Lin. Alg. Appl.*, Vol. 426, pp. 490–496, 2007. (Cited on pages 53 and 463.)
- [810] C. King, "Inequalities for Trace Norms of 2×2 Block Matrices," *Comm. Math. Phys.*, Vol. 242, pp. 531–545, 2003. (Cited on page 595.)
- [811] C. King and M. Nathanson, "New Trace Norm Inequalities for 2×2 Blocks of Diagonal Matrices," *Lin. Alg. Appl.*, Vol. 389, pp. 77–93, 2004. (Cited on page 586.)
- [812] F. Kittaneh, "Inequalities for the Schatten p -Norm III," *Commun. Math. Phys.*, Vol. 104, pp. 307–310, 1986. (Cited on page 584.)
- [813] F. Kittaneh, "Inequalities for the Schatten p -Norm. IV," *Commun. Math. Phys.*, Vol. 106, pp. 581–585, 1986. (Cited on page 583.)
- [814] F. Kittaneh, "On Zero-Trace Matrices," *Lin. Alg. Appl.*, Vol. 151, pp. 119–124, 1991. (Cited on page 313.)
- [815] F. Kittaneh, "A Note on the Arithmetic-Geometric-Mean Inequality for Matrices," *Lin. Alg. Appl.*, Vol. 171, pp. 1–8, 1992. (Cited on page 589.)
- [816] F. Kittaneh, "Norm Inequalities for Fractional Powers of Positive Operators," *Lett. Math. Phys.*, Vol. 27, pp. 279–285, 1993. (Cited on pages 515 and 613.)
- [817] F. Kittaneh, "Singular Values of Companion Matrices and Bounds on Zeros of Polynomials," *SIAM J. Matrix Anal. Appl.*, Vol. 16, pp. 333–340, 1995. (Cited on pages 327 and 711.)
- [818] F. Kittaneh, "Norm Inequalities for Certain Operator Sums," *J. Funct. Anal.*, Vol. 143, pp. 337–348, 1997. (Cited on pages 516, 517, and 589.)
- [819] F. Kittaneh, "Some Norm Inequalities for Operators," *Canad. Math. Bull.*, Vol. 42, pp. 87–96, 1999. (Cited on page 589.)
- [820] F. Kittaneh, "Commutator Inequalities Associated with the Polar Decomposition," *Proc. Amer. Math. Soc.*, Vol. 130, pp. 1279–1283, 2001. (Cited on pages 581, 584, 589, and 603.)
- [821] F. Kittaneh, "Norm Inequalities for Sums of Positive Operators," *J. Operator Theory*, Vol. 48, pp. 95–103, 2002. (Cited on pages 517, 589, 610, and 611.)
- [822] F. Kittaneh, "Bounds for the Zeros of Polynomials from Matrix Inequalities," *Arch. Math.*, Vol. 81, pp. 601–608, 2003. (Cited on pages 331 and 711.)

- [823] F. Kittaneh, "A Numerical Radius Inequality and an Estimate for the Numerical Radius of the Frobenius Companion Matrix," *Studia Mathematica*, Vol. 158, pp. 11–17, 2003. (Cited on pages 578 and 712.)
- [824] F. Kittaneh, "Norm Inequalities for Sums and Differences of Positive Operators," *Lin. Alg. Appl.*, Vol. 383, pp. 85–91, 2004. (Cited on pages 516, 517, and 603.)
- [825] F. Kittaneh, "Norm Inequalities for Sums of Positive Operators II," *Positivity*, Vol. 10, pp. 251–260, 2006. (Cited on pages 589 and 592.)
- [826] F. Kittaneh, "Bounds for the Zeros of Polynomials from Matrix Inequalities-II," *Lin. Multilin. Alg.*, Vol. 55, pp. 147–158, 2007. (Cited on page 711.)
- [827] F. Kittaneh and H. Kosaki, "Inequalities for the Schatten p -Norm. V," *Publications of RIMS Kyoto University*, Vol. 23, pp. 433–443, 1987. (Cited on pages 584 and 585.)
- [828] A.-L. Klaus and C.-K. Li, "Isometries for the Vector (p, q) Norm and the Induced (p, q) Norm," *Lin. Multilin. Alg.*, Vol. 38, pp. 315–332, 1995. (Cited on page 583.)
- [829] J. A. Knox and H. J. Brothers, "Novel Series-Based Approximations to e ," *College Math. J.*, Vol. 30, pp. 269–275, 1999. (Cited on pages 25 and 26.)
- [830] C. T. Koch and J. C. H. Spence, "A Useful Expansion of the Exponential of the Sum of Two Non-Commuting Matrices, One of Which is Diagonal," *J. Phys. A: Math. Gen.*, Vol. 36, pp. 803–816, 2003. (Cited on page 678.)
- [831] D. Koks, *Explorations in Mathematical Physics*. New York: Springer, 2006. (Cited on page 227.)
- [832] J. J. Koliha, "A Simple Proof of the Product Theorem for EP Matrices," *Lin. Alg. Appl.*, Vol. 294, pp. 213–215, 1999. (Cited on page 382.)
- [833] J. J. Koliha and V. Rakocevic, "Invertibility of the Sum of Idempotents," *Lin. Multilin. Alg.*, Vol. 50, pp. 285–292, 2002. (Cited on page 203.)
- [834] J. J. Koliha and V. Rakocevic, "Invertibility of the Difference of Idempotents," *Lin. Multilin. Alg.*, Vol. 51, pp. 97–110, 2003. (Cited on page 203.)
- [835] J. J. Koliha and V. Rakocevic, "The Nullity and Rank of Linear Combinations of Idempotent Matrices," *Lin. Alg. Appl.*, Vol. 418, pp. 11–14, 2006. (Cited on page 202.)
- [836] J. J. Koliha, V. Rakocevic, and I. Straskraba, "The Difference and Sum of Projectors," *Lin. Alg. Appl.*, Vol. 388, pp. 279–288, 2004. (Cited on pages 202, 203, and 205.)
- [837] N. Komaroff, "Bounds on Eigenvalues of Matrix Products with an Application to the Algebraic Riccati Equation," *IEEE Trans. Autom. Contr.*, Vol. 35, pp. 348–350, 1990. (Cited on page 333.)
- [838] N. Komaroff, "Rearrangement and Matrix Product Inequalities," *Lin. Alg. Appl.*, Vol. 140, pp. 155–161, 1990. (Cited on pages 61 and 518.)
- [839] V. Komornik, "Another Short Proof of Descartes's Rule of Signs," *Amer. Math. Monthly*, Vol. 113, pp. 829–830, 2006. (Cited on page 696.)
- [840] R. H. Koning, H. Neudecker, and T. Wansbeek, "Block Kronecker Products and the vecb Operator," *Lin. Alg. Appl.*, Vol. 149, pp. 165–184, 1991. (Cited on page 416.)

- [841] T. Koshy, *Fibonacci and Lucas Numbers with Applications*. New York: Wiley, 2001. (Cited on pages xvii, 218, and 278.)
- [842] J. Kovac-Striko and K. Veselic, "Trace Minimization and Definiteness of Symmetric Pencils," *Lin. Alg. Appl.*, Vol. 216, pp. 139–158, 1995. (Cited on page 362.)
- [843] O. Krafft, "An Arithmetic-Harmonic-Mean Inequality for Nonnegative Definite Matrices," *Lin. Alg. Appl.*, Vol. 268, pp. 243–246, 1998. (Cited on page 529.)
- [844] C. Krattenthaler, "Advanced Determinant Calculus: A Complement," *Lin. Alg. Appl.*, Vol. 411, pp. 68–166, 2005. (Cited on page xvii.)
- [845] W. Kratz, *Quadratic Functionals in Variational Analysis and Control Theory*. New York: Wiley, 1995. (Cited on page 805.)
- [846] E. Kreindler and A. Jameson, "Conditions for Nonnegativeness of Partitioned Matrices," *IEEE Trans. Autom. Contr.*, Vol. AC-17, pp. 147–148, 1972. (Cited on page 541.)
- [847] R. Kress, H. L. de Vries, and R. Wegmann, "On Nonnormal Matrices," *Lin. Alg. Appl.*, Vol. 8, pp. 109–120, 1974. (Cited on page 597.)
- [848] V. Kucera, "On Nonnegative Definite Solutions to Matrix Quadratic Equations," *Automatica*, Vol. 8, pp. 413–423, 1972. (Cited on page 805.)
- [849] A. Kufner, L. Maligranda, and L.-E. Persson, "The Prehistory of the Hardy Inequality," *Amer. Math. Monthly*, Vol. 113, pp. 715–732, 2006. (Cited on pages 58, 59, and 63.)
- [850] J. B. Kuipers, *Quaternions and Rotation Sequences: A Primer with Applications to Orbits, Aerospace, and Virtual Reality*. Princeton: Princeton University Press, 1999. (Cited on pages xvii, 226, and 228.)
- [851] S. Kurepa, "A Note on Logarithms of Normal Operators," *Proc. Amer. Math. Soc.*, Vol. 13, pp. 307–311, 1962. (Cited on page 689.)
- [852] K. Kwakernaak and R. Sivan, *Linear Optimal Control Systems*. New York: Wiley, 1972. (Cited on page xvii.)
- [853] M. Kwapisz, "The Power of a Matrix," *SIAM Rev.*, Vol. 40, pp. 703–705, 1998. (Cited on page 712.)
- [854] M. K. Kwong, "Some Results on Matrix Monotone Functions," *Lin. Alg. Appl.*, Vol. 118, pp. 129–153, 1989. (Cited on pages 221 and 460.)
- [855] I. I. Kyrchei, "Analogues of the Adjoint Matrix for Generalized Inverses and Corresponding Cramer Rules," *Lin. Multilin. Alg.*, Vol. 56, pp. 453–469, 2008. (Cited on page 129.)
- [856] K. R. Laberteaux, "Hermitian Matrices," *Amer. Math. Monthly*, Vol. 104, p. 277, 1997. (Cited on page 181.)
- [857] T. J. Laffey, "Products of Skew-Symmetric Matrices," *Lin. Alg. Appl.*, Vol. 68, pp. 249–251, 1985. (Cited on page 351.)
- [858] T. J. Laffey, "Products of Matrices," in *Generators and Relations in Groups and Geometries*, A. Barlotti, E. W. Ellers, P. Plaumann, and K. Strambach, Eds. Dordrecht: Kluwer, 1991, pp. 95–123. (Cited on page 351.)

- [859] T. J. Laffey and S. Lazarus, "Two-Generated Commutative Matrix Subalgebras," *Lin. Alg. Appl.*, Vol. 147, pp. 249–273, 1991. (Cited on page 319.)
- [860] T. J. Laffey and E. Meehan, "A Refinement of an Inequality of Johnson, Loewy and London on Nonnegative Matrices and Some Applications," *Elec. J. Lin. Alg.*, Vol. 3, pp. 119–128, 1998. (Cited on page 281.)
- [861] J. C. Lagarias and Y. Wang, "The Finiteness Conjecture for the Generalized Spectral Radius of a Set of Matrices," *Lin. Alg. Appl.*, Vol. 214, pp. 17–42, 1995. (Cited on page 715.)
- [862] S. Lakshminarayanan, S. L. Shah, and K. Nandakumar, "Cramer's Rule for Non-Square Matrices," *Amer. Math. Monthly*, Vol. 106, p. 865, 1999. (Cited on page 129.)
- [863] P. Lancaster, *Lambda-Matrices and Vibrating Systems*. Oxford: Pergamon, 1966, reprinted by Dover, Mineola, 2002. (Cited on page xvii.)
- [864] P. Lancaster and L. Rodman, "Solutions of the Continuous and Discrete Time Algebraic Riccati Equations: A Review," in *The Riccati Equation*, S. Bittanti, J. C. Willems, and A. Laub, Eds. New York: Springer, 1991, pp. 11–51. (Cited on page 805.)
- [865] P. Lancaster and L. Rodman, *Algebraic Riccati Equations*. Oxford: Clarendon, 1995. (Cited on pages xvii and 805.)
- [866] P. Lancaster and L. Rodman, "Canonical Forms for Hermitian Matrix Pairs under Strict Equivalence and Congruence," *SIAM Rev.*, Vol. 47, pp. 407–443, 2005. (Cited on pages 304 and 504.)
- [867] P. Lancaster and M. Tismenetsky, *The Theory of Matrices*, 2nd ed. Orlando: Academic Press, 1985. (Cited on pages 320, 340, 406, 560, 579, 619, 793, 794, and 795.)
- [868] L. Larson, *Problem Solving Through Problems*. New York: Springer, 1983. (Cited on pages 24, 29, 33, 35, 38, 40, 41, 42, 43, 44, 49, 155, and 158.)
- [869] L. Larsson, L. Maligranda, J. Pecaric, and L.-E. Persson, *Multiplicative Inequalities of Carlson Type and Interpolation*. Singapore: World Scientific, 2006. (Cited on page 58.)
- [870] J. Lasenby, A. N. Lasenby, and C. J. L. Doran, "A Unified Mathematical Language for Physics and Engineering in the 21st Century," *Phil. Trans. Math. Phys. Eng. Sci.*, Vol. 358, pp. 21–39, 2000. (Cited on pages 188, 227, and 565.)
- [871] J. B. Lasserre, "A Trace Inequality for Matrix Product," *IEEE Trans. Autom. Contr.*, Vol. 40, pp. 1500–1501, 1995. (Cited on page 333.)
- [872] A. J. Laub, *Matrix Analysis for Scientists and Engineers*. Philadelphia: SIAM, 2004. (Cited on pages xix, 113, 304, 306, and 338.)
- [873] A. J. Laub and K. Meyer, "Canonical Forms for Symplectic and Hamiltonian Matrices," *Celestial Mechanics*, Vol. 9, pp. 213–238, 1974. (Cited on page 802.)
- [874] C. Laurie, B. Mathes, and H. Radjavi, "Sums of Idempotents," *Lin. Alg. Appl.*, Vol. 208/209, pp. 175–197, 1994. (Cited on page 361.)
- [875] J. L. Lavoie, "A Determinantal Inequality Involving the Moore-Penrose Inverse," *Lin. Alg. Appl.*, Vol. 31, pp. 77–80, 1980. (Cited on page 392.)

- [876] C. L. Lawson, *Solving Least Squares Problems*. Englewood Cliffs: Prentice-Hall, 1974, reprinted by SIAM, Philadelphia, 1995. (Cited on page 398.)
- [877] J. D. Lawson and Y. Lim, "The Geometric Mean, Matrices, Metrics, and More," *Amer. Math. Monthly*, Vol. 108, pp. 797–812, 2001. (Cited on pages 431 and 463.)
- [878] P. D. Lax, *Linear Algebra*. New York: Wiley, 1997. (Cited on pages 160, 259, 457, and 600.)
- [879] S. R. Lay, *Convex Sets and Their Applications*. New York: Wiley, 1982. (Cited on pages 93, 164, and 635.)
- [880] B. Leclerc, "On Identities Satisfied by Minors of a Matrix," *Adv. Math.*, Vol. 100, pp. 101–132, 1993. (Cited on page 129.)
- [881] K. J. LeCouteur, "Representation of the Function $\text{Tr}(\exp(A-\lambda B))$ as a Laplace Transform with Positive Weight and Some Matrix Inequalities," *J. Phys. A: Math. Gen.*, Vol. 13, pp. 3147–3159, 1980. (Cited on pages 519 and 682.)
- [882] A. Lee, "On S-Symmetric, S-Skewsymmetric, and S-Orthogonal Matrices," *Periodica Math. Hungar.*, Vol. 7, pp. 71–76, 1976. (Cited on page 313.)
- [883] A. Lee, "Centrohermitian and Skew-Centrohermitian Matrices," *Lin. Alg. Appl.*, Vol. 29, pp. 205–210, 1980. (Cited on pages 230 and 376.)
- [884] J. M. Lee and D. A. Weinberg, "A Note on Canonical Forms for Matrix Congruence," *Lin. Alg. Appl.*, Vol. 249, pp. 207–215, 1996. (Cited on page 361.)
- [885] S. H. Lehnigk, *Stability Theorems for Linear Motions*. Englewood Cliffs: Prentice-Hall, 1966. (Cited on page 722.)
- [886] T.-G. Lei, C.-W. Woo, and F. Zhang, "Matrix Inequalities by Means of Embedding," *Elect. J. Lin. Alg.*, Vol. 11, pp. 66–77, 2004. (Cited on pages 453, 483, 493, 501, 505, 508, and 510.)
- [887] F. S. Leite, "Bounds on the Order of Generation of $\text{so}(n, r)$ by One-Parameter Subgroups," *Rocky Mountain J. Math.*, Vol. 21, pp. 879–911, 1183–1188, 1991. (Cited on pages 348 and 675.)
- [888] E. Leonard, "The Matrix Exponential," *SIAM Rev.*, Vol. 38, pp. 507–512, 1996. (Cited on page 646.)
- [889] G. Letac, "A Matrix and Its Matrix of Reciprocals Both Positive Semi-Definite," *Amer. Math. Monthly*, Vol. 82, pp. 80–81, 1975. (Cited on page 531.)
- [890] J. S. Lew, "The Cayley Hamilton Theorem in n Dimensions," *Z. Angew. Math. Phys.*, Vol. 17, pp. 650–653, 1966. (Cited on pages 260 and 261.)
- [891] A. S. Lewis, "Convex Analysis on the Hermitian Matrices," *SIAM J. Optim.*, Vol. 6, pp. 164–177, 1996. (Cited on pages 333 and 334.)
- [892] D. C. Lewis, "A Qualitative Analysis of S-Systems: Hopf Bifurcations," in *Canonical Nonlinear Modeling*, E. O. Voit, Ed. New York: Van Nostrand Reinhold, 1991, pp. 304–344. (Cited on page 414.)
- [893] C.-K. Li and R.-C. Li, "A Note on Eigenvalues of Perturbed Hermitian Matrices," *Lin. Alg. Appl.*, Vol. 395, pp. 183–190, 2005. (Cited on page 272.)
- [894] C.-K. Li and R. Mathias, "The Determinant of the Sum of Two Matrices," *Bull. Austral. Math. Soc.*, Vol. 52, pp. 425–429, 1995. (Cited on pages 611 and 612.)

- [895] C.-K. Li and R. Mathias, "The Lidskii-Mirsky-Wielandt Theorem—Additive and Multiplicative Versions," *Numer. Math.*, Vol. 81, pp. 377–413, 1999. (Cited on page 600.)
- [896] C.-K. Li and R. Mathias, "Extremal Characterizations of the Schur Complement and Resulting Inequalities," *SIAM Rev.*, Vol. 42, pp. 233–246, 2000. (Cited on pages 436, 442, 533, and 542.)
- [897] C.-K. Li and R. Mathias, "Inequalities on Singular Values of Block Triangular Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 24, pp. 126–131, 2002. (Cited on page 328.)
- [898] C.-K. Li and S. Nataraj, "Some Matrix Techniques in Game Theory," *Math. Ineq. Appl.*, Vol. 3, pp. 133–141, 2000. (Cited on page xvii.)
- [899] C.-K. Li and E. Poon, "Additive Decomposition of Real Matrices," *Lin. Multilin. Alg.*, Vol. 50, pp. 321–326, 2002. (Cited on page 360.)
- [900] C.-K. Li and L. Rodman, "Some Extremal Problems for Positive Definite Matrices and Operators," *Lin. Alg. Appl.*, Vol. 140, pp. 139–154, 1990. (Cited on pages 459 and 460.)
- [901] C.-K. Li and H. Schneider, "Orthogonality of Matrices," *Lin. Alg. Appl.*, Vol. 347, pp. 115–122, 2002. (Cited on page 570.)
- [902] J.-L. Li and Y.-L. Li, "On the Strengthened Jordan's Inequality," *J. Ineq. Appl.*, pp. 1–8, 2007, article ID 74328. (Cited on page 29.)
- [903] Q. Li, "Commutators of Orthogonal Projections," *Nihonkai. Math. J.*, Vol. 15, pp. 93–99, 2004. (Cited on page 581.)
- [904] X. Li and Y. Wei, "A Note on the Representations for the Drazin Inverse of 2×2 Block Matrices," *Lin. Alg. Appl.*, Vol. 423, pp. 332–338, 2007. (Cited on page 391.)
- [905] E. H. Lieb, "Convex Trace Functions and the Wigner-Yanase-Dyson Conjecture," *Adv. Math.*, Vol. 11, pp. 267–288, 1973. (Cited on pages 441, 500, 542, and 688.)
- [906] E. H. Lieb and M. Loss, *Analysis*. Providence: American Mathematical Society, 2001. (Cited on page 570.)
- [907] E. H. Lieb and M. B. Ruskai, "Some Operator Inequalities of the Schwarz Type," *Adv. Math.*, Vol. 12, pp. 269–273, 1974. (Cited on pages 474 and 542.)
- [908] E. H. Lieb and R. Seiringer, "Equivalent Forms of the Bessis-Moussa-Villani Conjecture," *J. Stat. Phys.*, Vol. 115, pp. 185–190, 2004. (Cited on page 482.)
- [909] E. H. Lieb and W. E. Thirring, "Inequalities for the Moments of the Eigenvalues of the Schrodinger Hamiltonian and Their Relation to Sobolev Inequalities," in *Studies in Mathematical Physics*, E. H. Lieb, B. Simon, and A. Wightman, Eds. Princeton: Princeton University Press, 1976, pp. 269–303. (Cited on page 519.)
- [910] Y. Lim, "The Matrix Golden Mean and its Applications to Riccati Matrix Equations," *SIAM J. Matrix Anal. Appl.*, Vol. 29, pp. 54–66, 2006. (Cited on page 803.)
- [911] C.-S. Lin, "On Halmos's Sharpening of Reid's Inequality," *Math. Reports Acad. Sci. Canada*, Vol. 20, pp. 62–64, 1998. (Cited on page 503.)
- [912] C.-S. Lin, "Inequalities of Reid Type and Furuta," *Proc. Amer. Math. Soc.*, Vol. 129, pp. 855–859, 2000. (Cited on page 503.)

- [913] C.-S. Lin, "Heinz-Kato-Furuta Inequalities with Bounds and Equality Conditions," *Math. Ineq. Appl.*, Vol. 5, pp. 735–743, 2002. (Cited on page 565.)
- [914] C.-S. Lin, "On Operator Order and Chaotic Operator Order for Two Operators," *Lin. Alg. Appl.*, Vol. 425, pp. 1–6, 2007. (Cited on pages 466 and 522.)
- [915] C.-S. Lin, "On Operator Order and Chaotic Operator Order for Two Operators," *Lin. Alg. Appl.*, Vol. 425, pp. 1–6, 2007. (Cited on page 522.)
- [916] T.-P. Lin, "The Power Mean and the Logarithmic Mean," *Amer. Math. Monthly*, Vol. 81, pp. 879–883, 1974. (Cited on pages 36 and 37.)
- [917] W.-W. Lin, "The Computation of the Kronecker Canonical Form of an Arbitrary Symmetric Pencil," *Lin. Alg. Appl.*, Vol. 103, pp. 41–71, 1988. (Cited on page 362.)
- [918] H. Linden, "Numerical Radii of Some Companion Matrices and Bounds for the Zeros of Polynomials," in *Analytic and Geometric Inequalities and Applications*, T. M. Rassias and H. M. Srivastava, Eds. Dordrecht: Kluwer, 1999, pp. 205–229. (Cited on pages 710 and 711.)
- [919] L. Lipsky, *Queueing Theory: A Linear Algebraic Approach*, 2nd ed. New York: Springer, 2008. (Cited on page xvii.)
- [920] B. Liu and H.-J. Lai, *Matrices in Combinatorics and Graph Theory*. New York: Springer, 2000. (Cited on page xvii.)
- [921] H. Liu and L. Zhu, "New Strengthened Carleman's Inequality and Hardy's Inequality," *J. Ineq. Appl.*, pp. 1–7, 2007, article ID 84104. (Cited on page 26.)
- [922] J. Liu and J. Wang, "Some Inequalities for Schur Complements," *Lin. Alg. Appl.*, Vol. 293, pp. 233–241, 1999. (Cited on pages 442 and 542.)
- [923] J. Liu and Q. Xie, "Inequalities Involving Khatri-Rao Products of Positive Semidefinite Hermitian Matrices," *Int. J. Inform. Sys. Sci.*, Vol. 4, pp. 30–40–135, 2008. (Cited on page 416.)
- [924] R.-W. Liu and R. J. Leake, "Exhaustive Equivalence Classes of Optimal Systems with Separable Controls," *SIAM Rev.*, Vol. 4, pp. 678–685, 1966. (Cited on page 186.)
- [925] S. Liu, "Matrix Results on the Khatri-Rao and Tracy-Singh Products," *Lin. Alg. Appl.*, Vol. 289, pp. 267–277, 1999. (Cited on pages 416, 533, 536, and 539.)
- [926] S. Liu, "Several Inequalities Involving Khatri-Rao Products of Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 354, pp. 175–186, 2002. (Cited on page 416.)
- [927] S. Liu and H. Neudecker, "Several Matrix Kantorovich-Type Inequalities," *Math. Anal. Appl.*, Vol. 197, pp. 23–26, 1996. (Cited on pages 57 and 501.)
- [928] S. Liu and G. Trenkler, "Hadamard, Khatri-Rao, Kronecker and Other Matrix Products," *Int. J. Inform. Sys. Sci.*, Vol. 4, pp. 160–177, 2008. (Cited on page 416.)
- [929] E. Liz, "A Note on the Matrix Exponential," *SIAM Rev.*, Vol. 40, pp. 700–702, 1998. (Cited on page 646.)
- [930] R. Loewy, "An Inertia Theorem for Lyapunov's Equation and the Dimension of a Controllability Space," *Lin. Alg. Appl.*, Vol. 260, pp. 1–7, 1997. (Cited on pages 793 and 794.)

- [931] D. O. Logofet, *Matrices and Graphs: Stability Problems in Mathematical Ecology*. Boca Raton: CRC Press, 1993. (Cited on page xvii.)
- [932] R. Lopez-Valcarce and S. Dasgupta, "Some Properties of the Matrix Exponential," *IEEE Trans. Circ. Sys. Analog. Dig. Sig. Proc.*, Vol. 48, pp. 213–215, 2001. (Cited on page 680.)
- [933] M. Loss and M. B. Ruskai, Eds., *Inequalities: Selecta of Elliott H. Lieb*. New York: Springer, 2002. (Cited on pages 500 and 688.)
- [934] P. Lounesto, *Clifford Algebras and Spinors*, 2nd ed. Cambridge: Cambridge University Press, 2001. (Cited on pages 188 and 227.)
- [935] D. G. Luenberger, *Optimization by Vector Space Methods*. New York: Wiley, 1969. (Cited on page xvii.)
- [936] D. G. Luenberger, *Introduction to Linear and Nonlinear Programming*, 2nd ed. Reading: Addison-Wesley, 1984. (Cited on page 505.)
- [937] M. Lundquist and W. Barrett, "Rank Inequalities for Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 248, pp. 91–100, 1996. (Cited on page 124.)
- [938] M. Lundquist and W. Barrett, "Rank Inequalities for Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 248, pp. 91–100, 1996. (Cited on pages 444 and 492.)
- [939] H. Lutkepohl, *Handbook of Matrices*. Chichester: Wiley, 1996. (Cited on page xix.)
- [940] E.-C. Ma, "A Finite Series Solution of the Matrix Equation $AX - XB = C$," *SIAM J. Appl. Math.*, Vol. 14, pp. 490–495, 1966. (Cited on page 797.)
- [941] Y. Ma, S. Soatto, J. Kosecka, and S. S. Sastry, *An Invitation to 3-D Vision*. New York: Springer, 2004. (Cited on page xvii.)
- [942] C. C. MacDuffee, *The Theory of Matrices*. New York: Chelsea, 1956. (Cited on pages 406, 410, 413, and 416.)
- [943] A. G. J. Macfarlane and N. Karcnias, "Poles and Zeros of Linear Multivariable Systems: A Survey of the Algebraic, Geometric, and Complex-Variable Theory," *Int. J. Contr.*, Vol. 24, pp. 33–74, 1976. (Cited on page 805.)
- [944] D. S. Mackey, N. Mackey, and F. Tisseur, "Structured Tools for Structured Matrices," *Elec. J. Lin. Alg.*, Vol. 10, pp. 106–145, 2003. (Cited on page 230.)
- [945] J. H. Maddocks, "Restricted Quadratic Forms, Inertia Theorems, and the Schur Complement," *Lin. Alg. Appl.*, Vol. 108, pp. 1–36, 1988. (Cited on pages 311 and 504.)
- [946] J. R. Magnus, "A Representation Theorem for $(\text{tr } A^p)^{1/p}$," *Lin. Alg. Appl.*, Vol. 95, pp. 127–134, 1987. (Cited on pages 476, 477, and 481.)
- [947] J. R. Magnus, *Linear Structures*. London: Griffin, 1988. (Cited on pages xvii, 390, and 416.)
- [948] J. R. Magnus and H. Neudecker, *Matrix Differential Calculus with Applications in Statistics and Econometrics*. Chichester: Wiley, 1988. (Cited on pages xvii, 370, 384, 388, 391, 398, 416, 476, 477, 503, and 642.)
- [949] W. Magnus, "On the Exponential Solution of Differential Equations for a Linear Operator," *Commun. Pure Appl. Math.*, Vol. VII, pp. 649–673, 1954. (Cited on page 678.)

- [950] K. N. Majindar, "On Simultaneous Hermitian Congruence Transformations of Matrices," *Amer. Math. Monthly*, Vol. 70, pp. 842–844, 1963. (Cited on page 508.)
- [951] L. Maligranda, "Simple Norm Inequalities," *Amer. Math. Monthly*, Vol. 113, pp. 256–260, 2006. (Cited on pages 567, 568, and 569.)
- [952] A. N. Malyshev and M. Sadkane, "On the Stability of Large Matrices," *J. Comp. Appl. Math.*, Vol. 102, pp. 303–313, 1999. (Cited on page 699.)
- [953] O. Mangasarian, *Nonlinear Programming*. New York: McGraw-Hill, 1969, reprinted by Krieger, Malabar, 1982. (Cited on page xvii.)
- [954] S. M. Manjegani, "Hölder and Young Inequalities for the Trace of Operators," *Positivity*, Vol. 11, pp. 239–250, 2007. (Cited on page 477.)
- [955] H. B. Mann, "Quadratic Forms with Linear Constraints," *Amer. Math. Monthly*, Vol. 50, pp. 430–433, 1943. (Cited on page 505.)
- [956] L. E. Mansfield, *Linear Algebra with Geometric Application*. New York: Marcel-Dekker, 1976. (Cited on page xix.)
- [957] M. Marcus, "An Eigenvalue Inequality for the Product of Normal Matrices," *Amer. Math. Monthly*, Vol. 63, pp. 173–174, 1956. (Cited on page 333.)
- [958] M. Marcus, *Finite Dimensional Multilinear Algebra, Part I*. Marcel Dekker, 1973. (Cited on pages 412 and 416.)
- [959] M. Marcus, *Finite Dimensional Multilinear Algebra, Part II*. Marcel Dekker, 1975. (Cited on page 416.)
- [960] M. Marcus, "Two Determinant Condensation Formulas," *Lin. Multilin. Alg.*, Vol. 22, pp. 95–102, 1987. (Cited on page 136.)
- [961] M. Marcus and S. M. Katz, "Matrices of Schur Functions," *Duke Math. J.*, Vol. 36, pp. 343–352, 1969. (Cited on page 494.)
- [962] M. Marcus and N. A. Khan, "A Note on the Hadamard Product," *Canad. Math. J.*, Vol. 2, pp. 81–83, 1959. (Cited on pages 405, 416, and 536.)
- [963] M. Marcus and H. Minc, *A Survey of Matrix Theory and Matrix Inequalities*. Boston: Prindle, Weber, and Schmidt, 1964, reprinted by Dover, New York, 1992. (Cited on pages xix, 55, 76, 229, 325, 544, and 598.)
- [964] M. Marcus and W. Watkins, "Partitioned Hermitian Matrices," *Duke Math. J.*, Vol. 38, pp. 237–249, 1971. (Cited on pages 450, 484, 494, and 513.)
- [965] M. Margaliot and G. Langholz, "The Routh-Hurwitz Array and Realization of Characteristic Polynomials," *IEEE Trans. Autom. Contr.*, Vol. 45, pp. 2424–2427, 2000. (Cited on page 704.)
- [966] T. L. Markham, "An Application of Theorems of Schur and Albert," *Proc. Amer. Math. Soc.*, Vol. 59, pp. 205–210, 1976. (Cited on page 541.)
- [967] T. L. Markham, "Oppenheim's Inequality for Positive Definite Matrices," *Amer. Math. Monthly*, Vol. 93, pp. 642–644, 1986. (Cited on page 535.)
- [968] G. Marsaglia and G. P. H. Styan, "Equalities and Inequalities for Ranks of Matrices," *Lin. Multilin. Alg.*, Vol. 2, pp. 269–292, 1974. (Cited on pages 121, 164, 387, and 389.)

- [969] J. E. Marsden and R. S. Ratiu, *Introduction to Mechanics and Symmetry: A Basic Exposition of Classical Mechanical Systems*, 2nd ed. New York: Springer, 1994. (Cited on page 193.)
- [970] J. E. Marsden and T. S. Ratiu, *Introduction to Mechanics and Symmetry*. New York: Springer, 1994. (Cited on page xvii.)
- [971] A. W. Marshall and I. Olkin, *Inequalities: Theory of Majorization and Its Applications*. New York: Academic Press, 1979. (Cited on pages 44, 45, 60, 61, 76, 158, 162, 163, 164, 326, 334, 412, 442, 443, 510, 541, 542, 618, and 619.)
- [972] A. W. Marshall and I. Olkin, "Matrix Versions of the Cauchy and Kantorovich Inequalities," *Aequationes Math.*, Vol. 40, pp. 89–93, 1990. (Cited on page 455.)
- [973] A. W. Marshall, I. Olkin, and B. Arnold, *Inequalities: Theory of Majorization and Its Applications*, 2nd ed. New York: Springer, 2009. (Cited on page 164.)
- [974] K. Martensson, "On the Matrix Riccati Equation," *Inform. Sci.*, Vol. 3, pp. 17–49, 1971. (Cited on page 805.)
- [975] A. M. Mathai, *Jacobians of Matrix Transformations and Functions of Matrix Argument*. Singapore: World Scientific, 1997. (Cited on page 642.)
- [976] R. Mathias, "The Spectral Norm of a Nonnegative Matrix," *Lin. Alg. Appl.*, Vol. 131, pp. 269–284, 1990. (Cited on page 615.)
- [977] R. Mathias, "Evaluating the Frechet Derivative of the Matrix Exponential," *Numer. Math.*, Vol. 63, pp. 213–226, 1992. (Cited on page 682.)
- [978] R. Mathias, "Matrices with Positive Definite Hermitian Part: Inequalities and Linear Systems," *SIAM J. Matrix Anal. Appl.*, Vol. 13, pp. 640–654, 1992. (Cited on pages 440 and 512.)
- [979] R. Mathias, "An Arithmetic-Geometric-Harmonic Mean Inequality Involving Hadamard Products," *Lin. Alg. Appl.*, Vol. 184, pp. 71–78, 1993. (Cited on page 448.)
- [980] R. Mathias, "A Chain Rule for Matrix Functions and Applications," *SIAM J. Matrix Anal. Appl.*, Vol. 17, pp. 610–620, 1996. (Cited on page 642.)
- [981] R. Mathias, "Singular Values and Singular Value Inequalities," in *Handbook of Linear Algebra*, L. Hogben, Ed. Boca Raton: Chapman & Hall/CRC, 2007, pp. 17–1–17–16. (Cited on page 332.)
- [982] M. Matic, C. E. M. Pearce, and J. Pecaric, "Shannon's and Related Inequalities in Information Theory," in *Survey on Classical Inequalities*, T. M. Rassias, Ed. Dordrecht: Kluwer, 2000, pp. 127–164. (Cited on pages 59 and 67.)
- [983] T. Matsuda, "An Inductive Proof of a Mixed Arithmetic-Geometric Mean Inequality," *Amer. Math. Monthly*, Vol. 102, pp. 634–637, 1995. (Cited on page 58.)
- [984] V. Mazorchuk and S. Rabanovich, "Multicommutators and Multianticommutators of Orthogonal Projectors," *Lin. Multilin. Alg.*, preprint. (Cited on page 581.)
- [985] J. E. McCarthy, "Pick's Theorem—What's the Big Deal?" *Amer. Math. Monthly*, Vol. 110, pp. 36–45, 2003. (Cited on page 450.)
- [986] J. M. McCarthy, *Geometric Design of Linkages*. New York: Springer, 2000. (Cited on page 722.)

- [987] J. P. McCloskey, "Characterizations of r -Potent Matrices," *Math. Proc. Camb. Phil. Soc.*, Vol. 96, pp. 213–222, 1984. (Cited on page 213.)
- [988] A. R. Meenakshi and C. Rajian, "On a Product of Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 295, pp. 3–6, 1999. (Cited on page 530.)
- [989] C. L. Mehta, "Some Inequalities Involving Traces of Operators," *J. Math. Phys.*, Vol. 9, pp. 693–697, 1968. (Cited on page 500.)
- [990] Y. A. Melnikov, *Influence Functions and Matrices*. New York: Marcel Dekker, 1998. (Cited on page xvii.)
- [991] P. R. Mercer, "The Dunkl-Williams Inequality in an Inner Product Space," *Math. Ineq. Appl.*, Vol. 10, pp. 447–450, 2007. (Cited on pages 567 and 568.)
- [992] J. K. Merikoski, H. Sarria, and P. Tarazaga, "Bounds for Singular Values Using Traces," *Lin. Alg. Appl.*, Vol. 210, pp. 227–254, 1994. (Cited on page 604.)
- [993] R. Merris, "Laplacian Matrices of Graphs: A Survey," *Lin. Alg. Appl.*, Vol. 198, pp. 143–176, 1994. (Cited on pages 277 and 506.)
- [994] R. Merris, *Multilinear Algebra*. Amsterdam: Gordon and Breach, 1997. (Cited on pages 416 and 536.)
- [995] R. Merris and S. Pierce, "Monotonicity of Positive Semidefinite Hermitian Matrices," *Proc. Amer. Math. Soc.*, Vol. 31, pp. 437–440, 1972. (Cited on pages 461 and 469.)
- [996] C. D. Meyer, "The Moore-Penrose Inverse of a Bordered Matrix," *Lin. Alg. Appl.*, Vol. 5, pp. 375–382, 1972. (Cited on page 391.)
- [997] C. D. Meyer, "Generalized Inverses and Ranks of Block Matrices," *SIAM J. Appl. Math.*, Vol. 25, pp. 597–602, 1973. (Cited on pages 357 and 391.)
- [998] C. D. Meyer, *Matrix Analysis and Applied Linear Algebra*. Philadelphia: SIAM, 2000. (Cited on pages 160, 175, 176, 177, 212, 243, 279, 375, and 714.)
- [999] C. D. Meyer and N. J. Rose, "The Index and the Drazin Inverse of Block Triangular Matrices," *SIAM J. Appl. Math.*, Vol. 33, pp. 1–7, 1977. (Cited on pages 343, 391, and 395.)
- [1000] J.-M. Miao, "General Expressions for the Moore-Penrose Inverse of a 2×2 Block Matrix," *Lin. Alg. Appl.*, Vol. 151, pp. 1–15, 1991. (Cited on page 391.)
- [1001] L. Mihalyffy, "An Alternative Representation of the Generalized Inverse of Partitioned Matrices," *Lin. Alg. Appl.*, Vol. 4, pp. 95–100, 1971. (Cited on page 391.)
- [1002] K. S. Miller, *Some Eclectic Matrix Theory*. Malabar: Krieger, 1987. (Cited on pages 254, 410, and 506.)
- [1003] G. A. Milliken and F. Akdeniz, "A Theorem on the Difference of the Generalized Inverses of Two Nonnegative Matrices," *Commun. Statist. Theory Methods*, Vol. 6, pp. 73–79, 1977. (Cited on pages 526 and 527.)
- [1004] G. V. Milovanovic, D. S. Mitrinovic, and T. M. Rassias, *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*. Singapore: World Scientific, 1994. (Cited on pages 709 and 712.)

- [1005] G. V. Milovanovic and T. M. Rassias, "Inequalities for Polynomial Zeros," in *Survey on Classical Inequalities*, T. M. Rassias, Ed. Dordrecht: Kluwer, 2000, pp. 165–202. (Cited on pages 638, 709, 710, 711, and 712.)
- [1006] N. Minamide, "An Extension of the Matrix Inversion Lemma," *SIAM J. Alg. Disc. Meth.*, Vol. 6, pp. 371–377, 1985. (Cited on page 378.)
- [1007] H. Miranda and R. C. Thompson, "A Trace Inequality with a Subtracted Term," *Lin. Alg. Appl.*, Vol. 185, pp. 165–172, 1993. (Cited on page 541.)
- [1008] L. Mirsky, *An Introduction to Linear Algebra*. Oxford: Clarendon, 1972, reprinted by Dover, Mineola, 1990. (Cited on pages xix, 190, and 198.)
- [1009] P. Misra and R. V. Patel, "A Determinant Identity and its Application in Evaluating Frequency Response Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 9, pp. 248–255, 1988. (Cited on page 800.)
- [1010] D. S. Mitrinovic, J. E. Pecaric, and A. M. Fink, *Classical and New Inequalities in Analysis*. Dordrecht: Kluwer, 1993. (Cited on pages 23, 70, 76, 271, 565, 568, and 569.)
- [1011] D. S. Mitrinovic, J. E. Pecaric, and V. Volenec, *Recent Advances in Geometric Inequalities*. Dordrecht: Kluwer, 1989. (Cited on page 158.)
- [1012] B. Mityagin, "An Inequality in Linear Algebra," *SIAM Rev.*, Vol. 33, pp. 125–127, 1991. (Cited on pages 458 and 541.)
- [1013] M. Moakher, "Means and Averaging in the Group of Rotations," *SIAM J. Matrix Anal. Appl.*, Vol. 24, pp. 1–16, 2002. (Cited on pages 196, 360, 674, 688, and 692.)
- [1014] M. Moakher, "A Differential Geometric Approach to the Geometric Mean of Symmetric Positive-Definite Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 26, pp. 735–747, 2005. (Cited on pages 463, 640, 688, and 689.)
- [1015] C. Moler and C. F. Van Loan, "Nineteen Dubious Ways to Compute the Exponential of a Matrix, Twenty-Five Years Later," *SIAM Rev.*, Vol. 45, pp. 3–49, 2000. (Cited on pages 692 and 722.)
- [1016] B. Mond, "Generalized Inverse Extensions of Matrix Inequalities," *Lin. Alg. Appl.*, Vol. 2, pp. 393–399, 1969. (Cited on page 502.)
- [1017] B. Mond and J. E. Pecaric, "Reverse Forms of a Convex Matrix Inequality," *Lin. Alg. Appl.*, Vol. 220, pp. 359–364, 1995. (Cited on page 457.)
- [1018] B. Mond and J. E. Pecaric, "Inequalities for the Hadamard Product of Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 19, pp. 66–70, 1998. (Cited on pages 537 and 538.)
- [1019] B. Mond and J. E. Pecaric, "On Inequalities Involving the Hadamard Product of Matrices," *Elec. J. Lin. Alg.*, Vol. 6, pp. 56–61, 2000. (Cited on pages 537 and 539.)
- [1020] V. V. Monov, "On the Spectrum of Convex Sets of Matrices," *IEEE Trans. Autom. Contr.*, Vol. 44, pp. 1009–1012, 1992. (Cited on page 269.)
- [1021] T. Moreland and S. Gudder, "Infima of Hilbert Space Effects," *Lin. Alg. Appl.*, Vol. 286, pp. 1–17, 1999. (Cited on page 460.)
- [1022] T. Mori, "Comments on 'A Matrix Inequality Associated with Bounds on Solutions of Algebraic Riccati and Lyapunov Equation'," *IEEE Trans. Autom. Contr.*, Vol. 33, p. 1088, 1988. (Cited on page 541.)

- [1023] T. Muir, *The Theory of Determinants in the Historical Order of Development*. New York: Dover, 1966. (Cited on page 164.)
- [1024] W. W. Muir, "Inequalities Concerning the Inverses of Positive Definite Matrices," *Proc. Edinburgh Math. Soc.*, Vol. 19, pp. 109–113, 1974–75. (Cited on pages 441, 442, and 542.)
- [1025] I. S. Murphy, "A Note on the Product of Complementary Principal Minors of a Positive Definite Matrix," *Lin. Alg. Appl.*, Vol. 44, pp. 169–172, 1982. (Cited on page 492.)
- [1026] R. M. Murray, Z. Li, and S. S. Sastry, *A Mathematical Introduction to Robotic Manipulation*. Boca Raton: CRC, 1994. (Cited on pages xvii and 722.)
- [1027] I. Najfeld and T. F. Havel, "Derivatives of the Matrix Exponential and Their Computation," *Adv. Appl. Math.*, Vol. 16, pp. 321–375, 1995. (Cited on page 682.)
- [1028] R. Nakamoto, "A Norm Inequality for Hermitian Operators," *Amer. Math. Monthly*, Vol. 110, pp. 238–239, 2003. (Cited on page 695.)
- [1029] Y. Nakamura, "Any Hermitian Matrix is a Linear Combination of Four Projections," *Lin. Alg. Appl.*, Vol. 61, pp. 133–139, 1984. (Cited on page 361.)
- [1030] A. W. Naylor and G. R. Sell, *Linear Operator Theory in Engineering and Science*. New York: Springer, 1986. (Cited on pages 70, 76, 565, 568, 623, 624, and 642.)
- [1031] T. Needham, *Visual Complex Analysis*. Oxford: Oxford University Press, 1997. (Cited on page 76.)
- [1032] C. N. Nett and W. M. Haddad, "A System-Theoretic Appropriate Realization of the Empty Matrix Concept," *IEEE Trans. Autom. Contr.*, Vol. 38, pp. 771–775, 1993. (Cited on page 164.)
- [1033] M. G. Neubauer, "An Inequality for Positive Definite Matrices with Applications to Combinatorial Matrices," *Lin. Alg. Appl.*, Vol. 267, pp. 163–174, 1997. (Cited on page 491.)
- [1034] M. F. Neuts, *Matrix-Geometric Solutions in Stochastic Models*. Baltimore: Johns Hopkins University Press, 1981, reprinted by Dover, Mineola, 1994. (Cited on page xvii.)
- [1035] R. W. Newcomb, "On the Simultaneous Diagonalization of Two Semi-Definite Matrices," *Quart. Appl. Math.*, Vol. 19, pp. 144–146, 1961. (Cited on page 541.)
- [1036] M. Newman, "Lyapunov Revisited," *Lin. Alg. Appl.*, Vol. 52/53, pp. 525–528, 1983. (Cited on page 309.)
- [1037] M. Newman, W. So, and R. C. Thompson, "Convergence Domains for the Campbell-Baker-Hausdorff Formula," *Lin. Multilin. Alg.*, Vol. 24, pp. 301–310, 1989. (Cited on page 655.)
- [1038] D. W. Nicholson, "Eigenvalue Bounds for of $AB + BA$ for A, B Positive Definite Matrices," *Lin. Alg. Appl.*, Vol. 24, pp. 173–183, 1979. (Cited on page 520.)
- [1039] C. Niculescu and L.-E. Persson, *Convex Functions and Their Applications: A Contemporary Approach*. New York: Springer, 2006. (Cited on pages 21, 22, 35, 37, 42, 54, 55, 56, 62, 132, 483, 565, 638, and 639.)

- [1040] C. P. Niculescu, "Convexity According to the Geometric Mean," *Math. Ineq. Appl.*, Vol. 3, pp. 155–167, 2000. (Cited on pages 39, 54, 158, and 712.)
- [1041] C. P. Niculescu and F. Popovici, "A Refinement of Popoviciu's Inequality," *Bull. Math. Soc. Sci. Math. Roumanie*, Vol. 49(97), pp. 285–290, 2006. (Cited on page 22.)
- [1042] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information*. Cambridge: Cambridge University Press, 2000. (Cited on page xvii.)
- [1043] M. Niezgoda, "Laguerre-Samuelson Type Inequalities," *Lin. Alg. Appl.*, Vol. 422, pp. 574–581, 2007. (Cited on pages 51 and 454.)
- [1044] K. Nishio, "The Structure of a Real Linear Combination of Two Projections," *Lin. Alg. Appl.*, Vol. 66, pp. 169–176, 1985. (Cited on page 200.)
- [1045] B. Noble and J. W. Daniel, *Applied Linear Algebra*, 3rd ed. Englewood Cliffs: Prentice-Hall, 1988. (Cited on pages xix and 164.)
- [1046] K. Nomakuchi, "On the Characterization of Generalized Inverses by Bordered Matrices," *Lin. Alg. Appl.*, Vol. 33, pp. 1–8, 1980. (Cited on page 391.)
- [1047] K. Nordstrom, "Some Further Aspects of the Lowner-Ordering Antitonicity of the Moore-Penrose Inverse," *Commun. Statist. Theory Meth.*, Vol. 18, pp. 4471–4489, 1989. (Cited on pages 457 and 527.)
- [1048] J. Nunemacher, "Which Real Matrices Have Real Logarithms?" *Math. Mag.*, Vol. 62, pp. 132–135, 1989. (Cited on page 722.)
- [1049] H. Ogawa, "An Operator Pseudo-Inversion Lemma," *SIAM J. Appl. Math.*, Vol. 48, pp. 1527–1531, 1988. (Cited on page 378.)
- [1050] I. Olkin, "An Inequality for a Sum of Forms," *Lin. Alg. Appl.*, Vol. 52–53, pp. 529–532, 1983. (Cited on page 456.)
- [1051] J. M. Ortega, *Matrix Theory, A Second Course*. New York: Plenum, 1987. (Cited on page xix.)
- [1052] B. Ortner and A. R. Krauter, "Lower Bounds for the Determinant and the Trace of a Class of Hermitian Matrices," *Lin. Alg. Appl.*, Vol. 236, pp. 147–180, 1996. (Cited on page 476.)
- [1053] S. L. Osburn and D. S. Bernstein, "An Exact Treatment of the Achievable Closed-Loop H_2 Performance of Sampled-Data Controllers: From Continuous-Time to Open-Loop," *Automatica*, Vol. 31, pp. 617–620, 1995. (Cited on page 681.)
- [1054] A. Ostrowski and H. Schneider, "Some Theorems on the Inertia of General Matrices," *J. Math. Anal. Appl.*, Vol. 4, pp. 72–84, 1962. (Cited on page 794.)
- [1055] A. M. Ostrowski, "On Some Metrical Properties of Operator Matrices and Matrices Partitioned into Blocks," *J. Math. Anal. Appl.*, Vol. 2, pp. 161–209, 1961. (Cited on page 593.)
- [1056] J. A. Oteo, "The Baker-Campbell-Hausdorff Formula and Nested Commutator Identities," *J. Math. Phys.*, Vol. 32, pp. 419–424, 1991. (Cited on page 678.)
- [1057] D. V. Ouellette, "Schur Complements and Statistics," *Lin. Alg. Appl.*, Vol. 36, pp. 187–295, 1981. (Cited on pages 468, 487, and 542.)

- [1058] D. A. Overdijk, "Skew-Symmetric Matrices in Classical Mechanics," Eindhoven University, Memorandum COSOR 89-23, 1989. (Cited on page 188.)
- [1059] C. C. Paige and M. Saunders, "Towards a Generalized Singular Value Decomposition," *SIAM J. Numer. Anal.*, Vol. 18, pp. 398–405, 1981. (Cited on page 316.)
- [1060] C. C. Paige, G. P. H. Styan, B. Y. Wang, and F. Zhang, "Hua's Matrix Equality and Schur Complements," *Int. J. Inform. Sys. Sci.*, Vol. 4, pp. 124–135, 2008. (Cited on pages 310, 472, 473, 490, and 503.)
- [1061] C. C. Paige and M. Wei, "History and Generality of the CS Decomposition," *Lin. Alg. Appl.*, Vol. 208/209, pp. 303–326, 1994. (Cited on page 316.)
- [1062] H. Palanthandalam-Madapusi, D. S. Bernstein, and R. Venugopal, "Dimensional Analysis of Matrices: State-Space Models and Dimensionless Units," *IEEE Contr. Sys. Mag.*, Vol. 27, no. December, pp. 100–109, 2007. (Cited on pages 230 and 679.)
- [1063] H. Palanthandalam-Madapusi, T. Van Pelt, and D. S. Bernstein, "Existence Conditions for Quadratic Programming with a Sign-Indefinite Quadratic Equality Constraint," preprint. (Cited on page 498.)
- [1064] H. Palanthandalam-Madapusi, T. Van Pelt, and D. S. Bernstein, "Parameter Consistency and Quadratically Constrained Errors-in-Variables Least-Squares Identification," preprint. (Cited on page 498.)
- [1065] F. C. Paliogiannis, "On Commuting Operator Exponentials," *Proc. Amer. Math. Soc.*, Vol. 131, pp. 3777–3781, 2003. (Cited on page 681.)
- [1066] B. P. Palka, *An Introduction to Complex Function Theory*. New York: Springer, 1991. (Cited on page 76.)
- [1067] C. V. Pao, "Logarithmic Derivatives of a Square Matrix," *Lin. Alg. Appl.*, Vol. 6, pp. 159–164, 1973. (Cited on page 691.)
- [1068] J. G. Papastavridis, *Tensor Calculus and Analytical Dynamics*. Boca Raton: CRC, 1998. (Cited on page xvii.)
- [1069] J. G. Papastavridis, *Analytical Mechanics*. Oxford: Oxford University Press, 2002. (Cited on pages xvii and 451.)
- [1070] F. C. Park, "Computational Aspects of the Product-of-Exponentials Formula for Robot Kinematics," *IEEE Trans. Autom. Contr.*, Vol. 39, pp. 643–647, 1994. (Cited on page 722.)
- [1071] P. Park, "On the Trace Bound of a Matrix Product," *IEEE Trans. Autom. Contr.*, Vol. 41, pp. 1799–1802, 1996. (Cited on page 334.)
- [1072] D. F. Parker, *Fields, Flow and Waves: An Introduction to Continuum Models*. London: Springer, 2003. (Cited on page 362.)
- [1073] P. C. Parks, "A New Proof of the Routh-Hurwitz Stability Criterion Using the Second Method of Liapunov," *Proc. Camb. Phil. Soc.*, Vol. 58, pp. 694–702, 1962. (Cited on page 704.)
- [1074] R. V. Patel, "On Blocking Zeros in Linear Multivariable Systems," *IEEE Trans. Autom. Contr.*, Vol. AC-31, pp. 239–241, 1986. (Cited on page 805.)
- [1075] R. V. Patel and M. Toda, "Trace Inequalities Involving Hermitian Matrices," *Lin. Alg. Appl.*, Vol. 23, pp. 13–20, 1979. (Cited on pages 484 and 692.)

- [1076] C. Pearcy, "A Complete Set of Unitary Invariants for Operators Generating Finite W^* -Algebras of Type I," *Pac. J. Math.*, Vol. 12, pp. 1405–1414, 1962. (Cited on page 318.)
- [1077] M. C. Pease, *Methods of Matrix Algebra*. New York: Academic Press, 1965. (Cited on page 229.)
- [1078] J. Pecaric and R. Rajic, "The Dunkl-Williams Inequality with n Elements in Normed Linear Spaces," *Math. Ineq. Appl.*, Vol. 10, pp. 461–470, 2007. (Cited on page 568.)
- [1079] J. E. Pecaric, S. Puntanen, and G. P. H. Styan, "Some Further Matrix Extensions of the Cauchy-Schwarz and Kantorovich Inequalities, with Some Statistical Applications," *Lin. Alg. Appl.*, Vol. 237/238, pp. 455–476, 1996. (Cited on pages 501 and 502.)
- [1080] J. E. Pecaric, S. Puntanen, and G. P. H. Styan, "Some Further Matrix Extensions of the Cauchy-Schwarz and Kantorovich Inequalities, with Some Statistical Applications," *Lin. Alg. Appl.*, Vol. 237/238, pp. 455–476, 1996. (Cited on page 616.)
- [1081] S. Perlis, *Theory of Matrices*. Reading: Addison-Wesley, 1952, reprinted by Dover, New York, 1991. (Cited on pages 164, 233, 234, 237, 281, and 361.)
- [1082] T. Peter, "Maximizing the Area of a Quadrilateral," *College Math. J.*, Vol. 34, pp. 315–316, 2003. (Cited on page 159.)
- [1083] I. R. Petersen and C. V. Hollot, "A Riccati Equation Approach to the Stabilization of Uncertain Systems," *Automatica*, Vol. 22, pp. 397–411, 1986. (Cited on page 504.)
- [1084] D. Petz and R. Temesi, "Means of Positive Numbers and Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 27, pp. 712–720, 2005. (Cited on pages 463 and 467.)
- [1085] L. A. Pipes, "Applications of Laplace Transforms of Matrix Functions," *J. Franklin Inst.*, Vol. 285, pp. 436–451, 1968. (Cited on pages 672 and 673.)
- [1086] A. O. Pittenger and M. H. Rubin, "Convexity and Separation Problem of Quantum Mechanical Density Matrices," *Lin. Alg. Appl.*, Vol. 346, pp. 47–71, 2002. (Cited on page xvii.)
- [1087] I. Polik and T. Terlaky, "A Survey of the S-Lemma," *SIAM Rev.*, Vol. 49, pp. 371–418, 2007. (Cited on page 498.)
- [1088] T. Politi, "A Formula for the Exponential of a Real Skew-Symmetric Matrix of Order 4," *Numer. Math. BIT*, Vol. 41, pp. 842–845, 2001. (Cited on pages 673 and 675.)
- [1089] D. S. G. Pollock, "Tensor Products and Matrix Differential Calculus," *Lin. Alg. Appl.*, Vol. 67, pp. 169–193, 1985. (Cited on page 642.)
- [1090] B. T. Polyak, "Convexity of Quadratic Transformations and Its Use in Control and Optimization," *J. Optim. Theory Appl.*, Vol. 99, pp. 553–583, 1998. (Cited on pages 497, 498, and 499.)
- [1091] B. Poonen, "A Unique $(2k+1)$ -th Root of a Matrix," *Amer. Math. Monthly*, Vol. 98, p. 763, 1991. (Cited on page 349.)
- [1092] B. Poonen, "Positive Deformations of the Cauchy Matrix," *Amer. Math. Monthly*, Vol. 102, pp. 842–843, 1995. (Cited on page 447.)

- [1093] D. Popa and I. Rasa, "Inequalities Involving the Inner Product," *J. Ineq. Pure Appl. Math.*, Vol. 8, no. 3, pp. 1–4, 2007, Article 86. (Cited on page 565.)
- [1094] V. M. Popov, *Hyperstability of Control Systems*. Berlin: Springer, 1973. (Cited on pages xvii and 722.)
- [1095] G. J. Porter, "Linear Algebra and Affine Planar Transformations," *College Math. J.*, Vol. 24, pp. 47–51, 1993. (Cited on page 194.)
- [1096] B. H. Pourciau, "Modern Multiplier Rules," *Amer. Math. Monthly*, Vol. 87, pp. 433–452, 1980. (Cited on page 635.)
- [1097] C. R. Pranesachar, "Ratio Circum-to-In-Radius," *Amer. Math. Monthly*, Vol. 114, p. 648, 2007. (Cited on page 158.)
- [1098] V. V. Prasolov, *Problems and Theorems in Linear Algebra*. Providence: American Mathematical Society, 1994. (Cited on pages xix, 143, 150, 184, 185, 197, 208, 214, 227, 255, 257, 259, 271, 275, 313, 320, 328, 334, 335, 340, 341, 343, 345, 349, 350, 358, 393, 412, 442, 474, 486, 487, 488, 493, 507, 508, 533, 534, 574, 588, 599, 601, 608, 640, 641, and 660.)
- [1099] U. Prells, M. I. Friswell, and S. D. Garvey, "Use of Geometric Algebra: Compound Matrices and the Determinant of the Sum of Two Matrices," *Proc. Royal Soc. Lond. A*, Vol. 459, pp. 273–285, 2003. (Cited on pages 412 and 413.)
- [1100] J. S. Przemieniecki, *Theory of Matrix Structural Analysis*. New York: McGraw-Hill, 1968. (Cited on page xvii.)
- [1101] P. J. Psarrakos, "On the m th Roots of a Complex Matrix," *Elec. J. Lin. Alg.*, Vol. 9, pp. 32–41, 2002. (Cited on page 348.)
- [1102] N. J. Pullman, *Matrix Theory and Its Applications: Selected Topics*. New York: Marcel Dekker, 1976. (Cited on page 722.)
- [1103] S. Puntanen and G. P. H. Styan, "Historical Introduction: Issai Schur and the Early Development of the Schur Complement," in *The Schur Complement and Its Applications*, F. Zhang, Ed. New York: Springer, 2004, pp. 1–16. (Cited on page 386.)
- [1104] F. Qi, "Inequalities between the Sum of Squares and the Exponential of Sum of a Nonnegative Sequence," *J. Ineq. Pure Appl. Math.*, Vol. 8, no. 3, pp. 1–5, 2007, Article 78. (Cited on page 60.)
- [1105] F. Qi and L. Debnath, "Inequalities for Power-Exponential Functions," *J. Ineq. Pure Appl. Math.*, Vol. 1, no. 2/15, pp. 1–5, 2000. (Cited on page 38.)
- [1106] C. Qian and J. Li, "Global Finite-Time Stabilization by Output Feedback for Planar Systems without Observation Linearization," *IEEE Trans. Autom. Contr.*, Vol. 50, pp. 885–890, 2005. (Cited on page 38.)
- [1107] R. X. Qian and C. L. DeMarco, "An Approach to Robust Stability of Matrix Polytopes through Copositive Homogeneous Polynomials," *IEEE Trans. Autom. Contr.*, Vol. 37, pp. 848–852, 1992. (Cited on page 269.)
- [1108] L. Qiu, B. Bernhardsson, A. Rantzer, E. J. Davison, P. M. Young, and J. C. Doyle, "A Formula for Computation of the Real Stability Radius," *Automatica*, Vol. 31, pp. 879–890, 1995. (Cited on page 701.)

- [1109] L. Qiu and X. Zhan, "On the Span of Hadamard Products of Vectors," *Lin. Alg. Appl.*, Vol. 422, pp. 304–307, 2007. (Cited on page 413.)
- [1110] J. F. Queiro and A. Kovacec, "A Bound for the Determinant of the Sum of Two Normal Matrices," *Lin. Multilin. Alg.*, Vol. 33, pp. 171–173, 1993. (Cited on page 335.)
- [1111] V. Rabanovich, "Every Matrix Is a Linear Combination of Three Idempotents," *Lin. Alg. Appl.*, Vol. 390, pp. 137–143, 2004. (Cited on page 361.)
- [1112] H. Radjavi, "Decomposition of Matrices into Simple Involutions," *Lin. Alg. Appl.*, Vol. 12, pp. 247–255, 1975. (Cited on page 348.)
- [1113] H. Radjavi and P. Rosenthal, *Simultaneous Triangularization*. New York: Springer, 2000. (Cited on page 358.)
- [1114] H. Radjavi and J. P. Williams, "Products of Self-Adjoint Operators," *Michigan Math. J.*, Vol. 16, pp. 177–185, 1969. (Cited on pages 319, 348, and 376.)
- [1115] V. Rakocevic, "On the Norm of Idempotent Operators in a Hilbert Space," *Amer. Math. Monthly*, Vol. 107, pp. 748–750, 2000. (Cited on pages 210 and 336.)
- [1116] A. C. M. Ran and R. Vreugdenhil, "Existence and Comparison Theorems for Algebraic Riccati Equations for Continuous- and Discrete-Time Systems," *Lin. Alg. Appl.*, Vol. 99, pp. 63–83, 1988. (Cited on page 805.)
- [1117] A. Rantzer, "On the Kalman-Yakubovich-Popov Lemma," *Sys. Contr. Lett.*, Vol. 28, pp. 7–10, 1996. (Cited on page 320.)
- [1118] C. R. Rao and S. K. Mitra, *Generalized Inverse of Matrices and Its Applications*. New York: Wiley, 1971. (Cited on pages 398, 529, and 541.)
- [1119] C. R. Rao and M. B. Rao, *Matrix Algebra and Its Applications to Statistics and Econometrics*. Singapore: World Scientific, 1998. (Cited on pages xvii, 278, and 416.)
- [1120] J. V. Rao, "Some More Representations for the Generalized Inverse of a Partitioned Matrix," *SIAM J. Appl. Math.*, Vol. 24, pp. 272–276, 1973. (Cited on page 391.)
- [1121] U. A. Rauhala, "Array Algebra Expansion of Matrix and Tensor Calculus: Part 1," *SIAM J. Matrix Anal. Appl.*, Vol. 24, pp. 490–508, 2003. (Cited on page 416.)
- [1122] P. A. Regalia and S. K. Mitra, "Kronecker Products, Unitary Matrices and Signal Processing Applications," *SIAM Rev.*, Vol. 31, pp. 586–613, 1989. (Cited on page 416.)
- [1123] M. W. Reinsch, "A Simple Expression for the Terms in the Baker-Campbell-Hausdorff Series," *J. Math. Physics*, Vol. 41, pp. 2434–2442, 2000. (Cited on page 683.)
- [1124] T. J. Richardson and R. H. Kwong, "On Positive Definite Solutions to the Algebraic Riccati Equation," *Sys. Contr. Lett.*, Vol. 7, pp. 99–104, 1986. (Cited on pages 790 and 805.)
- [1125] A. N. Richmond, "Expansions for the Exponential of a Sum of Matrices," in *Applications of Matrix Theory*, M. J. C. Gover and S. Barnett, Eds. Oxford: Oxford University Press, 1989, pp. 283–289. (Cited on page 684.)
- [1126] K. S. Riedel, "A Sherman-Morrison-Woodbury Identity for Rank Augmenting Matrices with Application to Centering," *SIAM J. Matrix Anal. Appl.*, Vol. 13, pp. 659–662, 1992. (Cited on page 378.)

- [1127] J. R. Ringrose, *Compact Non-Self-Adjoint Operators*. New York: Van Nostrand Reinhold, 1971. (Cited on page 619.)
- [1128] R. S. Rivlin, "Further Remarks on the Stress Deformation Relations for Isotropic Materials," *J. Rational Mech. Anal.*, Vol. 4, pp. 681–702, 1955. (Cited on pages 260 and 261.)
- [1129] J. W. Robbin, *Matrix Algebra Using MINimal MATlab*. Wellesley: A. K. Peters, 1995. (Cited on pages 76, 115, 164, 211, 229, 281, 283, 319, 346, and 640.)
- [1130] J. W. Robbin and D. A. Salamon, "The Exponential Vandermonde Matrix," *Lin. Alg. Appl.*, Vol. 317, pp. 225–226, 2000. (Cited on page 414.)
- [1131] D. W. Robinson, "Nullities of Submatrices of the Moore-Penrose Inverse," *Lin. Alg. Appl.*, Vol. 94, pp. 127–132, 1987. (Cited on page 218.)
- [1132] P. Robinson and A. J. Wathen, "Variational Bounds on the Entries of the Inverse of a Matrix," *IMA J. Numerical Anal.*, Vol. 12, pp. 463–486, 1992. (Cited on page 476.)
- [1133] R. T. Rockafellar, *Convex Analysis*. Princeton: Princeton University Press, 1990. (Cited on pages 164, 624, 632, 635, and 642.)
- [1134] R. T. Rockafellar and R. J. B. Wets, *Variational Analysis*. Berlin: Springer, 1998. (Cited on pages 632 and 642.)
- [1135] L. Rodman, "Products of Symmetric and Skew Symmetric Matrices," *Lin. Multilin. Alg.*, Vol. 43, pp. 19–34, 1997. (Cited on page 351.)
- [1136] G. S. Rogers, *Matrix Derivatives*. New York: Marcel Dekker, 1980. (Cited on page 642.)
- [1137] C. A. Rohde, "Generalized Inverses of Partitioned Matrices," *SIAM J. Appl. Math.*, Vol. 13, pp. 1033–1035, 1965. (Cited on page 391.)
- [1138] J. Rohn, "Computing the Norm $\|A\|_{\infty,1}$ Is NP-Hard," *Lin. Multilin. Alg.*, Vol. 47, pp. 195–204, 2000. (Cited on page 575.)
- [1139] O. Rojo, "Further Bounds for the Smallest Singular Value and the Spectral Condition Number," *Computers Math. Appl.*, Vol. 38, pp. 215–228, 1999. (Cited on page 604.)
- [1140] O. Rojo, "Inequalities Involving the Mean and the Standard Deviation of Nonnegative Real Numbers," *J. Ineq. Appl.*, Vol. 2006, pp. 1–15, 2006, article ID 43465. (Cited on page 51.)
- [1141] J. Rooij, "On Ky Fan's Inequality and its Additive Analogues," *Math. Ineq. Appl.*, Vol. 6, pp. 595–604, 2003. (Cited on page 55.)
- [1142] D. J. Rose, "Matrix Identities of the Fast Fourier Transform," *Lin. Alg. Appl.*, Vol. 29, pp. 423–443, 1980. (Cited on page 357.)
- [1143] K. H. Rosen, Ed., *Handbook of Discrete and Combinatorial Mathematics*. Boca Raton: CRC, 2000. (Cited on pages xvii and xix.)
- [1144] H. H. Rosenbrock, *State-Space and Multivariable Theory*. New York: Wiley, 1970. (Cited on pages 763 and 764.)
- [1145] M. Rosenfeld, "A Sufficient Condition for Nilpotence," *Amer. Math. Monthly*, Vol. 103, pp. 907–909, 1996. (Cited on pages xvii and 215.)

- [1146] A. Rosoiu, "Triangle Radii Inequality," *Amer. Math. Monthly*, Vol. 114, p. 640, 2007. (Cited on page 158.)
- [1147] W. Rossmann, *Lie Groups: An Introduction Through Linear Groups*. Oxford: Oxford University Press, 2002. (Cited on page 229.)
- [1148] U. G. Rothblum, "Nonnegative Matrices and Stochastic Matrices," in *Handbook of Linear Algebra*, L. Hogben, Ed. Boca Raton: Chapman & Hall/CRC, 2007, pp. 9–1–9–25. (Cited on pages 275, 280, and 398.)
- [1149] J. J. Rotman, *An Introduction to the Theory of Groups*, 4th ed. New York: Springer, 1999. (Cited on pages 223 and 357.)
- [1150] W. J. Rugh, *Linear System Theory*, 2nd ed. Upper Saddle River: Prentice Hall, 1996. (Cited on pages 252, 253, 678, 743, 756, and 805.)
- [1151] A. M. Russell and C. J. F. Upton, "A Class of Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 93, pp. 121–126, 1987. (Cited on page 449.)
- [1152] D. L. Russell, *Mathematics of Finite-Dimensional Control Systems*. New York: Marcel Dekker, 1979. (Cited on page 793.)
- [1153] A. Saberi, P. Sannuti, and B. M. Chen, *H₂ Optimal Control*. Englewood Cliffs: Prentice-Hall, 1995. (Cited on page xvii.)
- [1154] M. K. Sain and C. B. Schrader, "The Role of Zeros in the Performance of Multi-input, Multioutput Feedback Systems," *IEEE Trans. Educ.*, Vol. 33, pp. 244–257, 1990. (Cited on page 805.)
- [1155] J. Sandor, "On Certain Inequalities for Means, II," *J. Math. Anal. Appl.*, Vol. 199, pp. 629–635, 1996. (Cited on page 37.)
- [1156] J. Sandor, "Inequalities for Generalized Convex Functions with Applications I," *Studia University Babes-Bolyai, Mathematica*, Vol. XLIV, pp. 93–107, 1999. (Cited on page 22.)
- [1157] J. Sandor, "On Certain Limits Related to the Number e ," *Libertas Mathematica*, Vol. XX, pp. 155–159, 2000. (Cited on page 26.)
- [1158] J. Sandor, "Inequalities for Generalized Convex Functions with Applications II," *Studia University Babes-Bolyai, Mathematica*, Vol. XLVI, pp. 79–91, 2001. (Cited on page 22.)
- [1159] J. Sandor, "On an Inequality of Ky Fan," *Int. J. Math. Educ. Sci. Tech.*, Vol. 32, pp. 133–160, 2001. (Cited on page 55.)
- [1160] J. Sandor and L. Debnath, "On Certain Inequalities Involving the Constant e and Their Applications," *J. Math. Anal. Appl.*, Vol. 249, pp. 569–582, 2000. (Cited on pages 25, 26, and 66.)
- [1161] R. A. Satnoianu, "General Power Inequalities between the Sides and the Circumscribed and Inscribed Radii Related to the Fundamental Triangle Inequality," *Math. Ineq. Appl.*, Vol. 5, pp. 745–751, 2002. (Cited on page 158.)
- [1162] D. H. Sattinger and O. L. Weaver, *Lie Groups and Algebras with Applications to Physics, Geometry, and Mechanics*. New York: Springer, 1986. (Cited on pages 150, 172, 655, 660, and 722.)

- [1163] A. H. Sayed, *Fundamentals of Adaptive Filtering*. New York: Wiley, 2003. (Cited on page xvii.)
- [1164] H. Schaub, P. Tsiotras, and J. L. Junkins, "Principal Rotation Representations of Proper $N \times N$ Orthogonal Matrices," *Int. J. Eng. Sci.*, Vol. 33, pp. 2277–2295, 1995. (Cited on page 674.)
- [1165] C. W. Scherer, "The Solution Set of the Algebraic Riccati Equation and the Algebraic Riccati Inequality," *Lin. Alg. Appl.*, Vol. 153, pp. 99–122, 1991. (Cited on page 805.)
- [1166] C. W. Scherer, "The Algebraic Riccati Equation and Inequality for Systems with Uncontrollable Modes on the Imaginary Axis," *SIAM J. Matrix Anal. Appl.*, Vol. 16, pp. 1308–1327, 1995. (Cited on pages 787 and 805.)
- [1167] C. W. Scherer, "The General Nonstrict Algebraic Riccati Inequality," *Lin. Alg. Appl.*, Vol. 219, pp. 1–33, 1995. (Cited on page 805.)
- [1168] P. Scherk, "On the Decomposition of Orthogonalities into Symmetries," *Proc. Amer. Math. Soc.*, Vol. 1, pp. 481–491, 1950. (Cited on page 347.)
- [1169] C. Schmoeger, "Remarks on Commuting Exponentials in Banach Algebras," *Proc. Amer. Math. Soc.*, Vol. 127, pp. 1337–1338, 1999. (Cited on page 681.)
- [1170] C. Schmoeger, "Remarks on Commuting Exponentials in Banach Algebras, II," *Proc. Amer. Math. Soc.*, Vol. 128, pp. 3405–3409, 2000. (Cited on page 681.)
- [1171] C. Schmoeger, "On Normal Operator Exponentials," *Proc. Amer. Math. Soc.*, Vol. 130, pp. 697–702, 2001. (Cited on page 681.)
- [1172] C. Schmoeger, "On Logarithms of Linear Operators on Hilbert Spaces," *Demonstratio Math.*, Vol. XXXV, pp. 375–384, 2002. (Cited on pages 650 and 689.)
- [1173] C. Schmoeger, "On the Operator Equation $e^A = e^B$," *Lin. Alg. Appl.*, Vol. 359, pp. 169–179, 2003. (Cited on page 682.)
- [1174] H. Schneider, "Olga Taussky-Todd's Influence on Matrix Theory and Matrix Theorists," *Lin. Multilin. Alg.*, Vol. 5, pp. 197–224, 1977. (Cited on page 269.)
- [1175] B. Scholkopf and A. J. Smola, *Learning with Kernels: Support Vector Machines, Regularization, Optimization, and Beyond*. Cambridge: MIT Press, 2001. (Cited on page 445.)
- [1176] D. Scholz and M. Weyrauch, "A Note on the Zassenhaus Product Formula," *J. Math. Phys.*, Vol. 47, pp. 033 505/1–7, 2006. (Cited on page 685.)
- [1177] J. R. Schott, *Matrix Analysis for Statistics*. New York: Wiley, 2005. (Cited on pages xvii, 379, 443, 513, and 518.)
- [1178] C. B. Schrader and M. K. Sain, "Research on System Zeros: A Survey," *Int. J. Contr.*, Vol. 50, pp. 1407–1433, 1989. (Cited on page 805.)
- [1179] B. S. W. Schroder, *Ordered Sets: An Introduction*. Boston: Birkhauser, 2003. (Cited on page 76.)
- [1180] A. J. Schwenk, "Tight Bounds on the Spectral Radius of Asymmetric Nonnegative Matrices," *Lin. Alg. Appl.*, Vol. 75, pp. 257–265, 1986. (Cited on page 415.)
- [1181] S. R. Searle, *Matrix Algebra Useful for Statistics*. New York: Wiley, 1982. (Cited on page xvii.)

- [1182] P. Sebastian, “On the Derivatives of Matrix Powers,” *SIAM J. Matrix Anal. Appl.*, Vol. 17, pp. 640–648, 1996. (Cited on page 642.)
- [1183] P. Sebastiani, “On the Derivatives of Matrix Powers,” *SIAM J. Matrix Anal. Appl.*, Vol. 17, pp. 640–648, 1996. (Cited on page 640.)
- [1184] G. A. F. Seber, *A Matrix Handbook for Statisticians*. New York: Wiley, 2008. (Cited on pages 67, 111, 118, 139, 154, 156, 184, 206, 208, 213, 217, 220, 230, 269, 271, 374, 375, 379, 393, 487, 500, 526, 535, 572, 607, and 706.)
- [1185] J. M. Selig, *Geometric Fundamentals of Robotics*, 2nd ed. New York: Springer, 2005. (Cited on pages xvii, 193, 227, and 229.)
- [1186] D. Serre, *Matrices: Theory and Applications*. New York: Springer, 2002. (Cited on pages 172, 217, 260, and 390.)
- [1187] J.-P. Serre and L. L. Scott, *Linear Representations of Finite Groups*. New York: Springer, 1996. (Cited on page 223.)
- [1188] C. Shafroth, “A Generalization of the Formula for Computing the Inverse of a Matrix,” *Amer. Math. Monthly*, Vol. 88, pp. 614–616, 1981. (Cited on pages 139 and 196.)
- [1189] W. M. Shah and A. Liman, “On the Zeros of a Class of Polynomials,” *Math. Ineq. Appl.*, Vol. 10, pp. 733–744, 2007. (Cited on page 709.)
- [1190] H. Shapiro, “A Survey of Canonical Forms and Invariants for Unitary Similarity,” *Lin. Alg. Appl.*, Vol. 147, pp. 101–167, 1991. (Cited on page 318.)
- [1191] H. Shapiro, “Notes from Math 223: Olga Taussky Todd’s Matrix Theory Course, 1976–1977,” *Mathematical Intelligencer*, Vol. 19, no. 1, pp. 21–27, 1997. (Cited on page 351.)
- [1192] R. Shaw and F. I. Yeadon, “On $(a \times b) \times c$,” *Amer. Math. Monthly*, Vol. 96, pp. 623–629, 1989. (Cited on page 188.)
- [1193] S.-Q. Shen and T.-Z. Huang, “Several Inequalities for the Largest Singular Value and the Spectral Radius of Matrices,” *Math. Ineq. Appl.*, Vol. 10, pp. 713–722, 2007. (Cited on pages 275, 416, and 617.)
- [1194] G. E. Shilov, *Linear Algebra*. Englewood Cliffs: Prentice-Hall, 1971, reprinted by Dover, New York, 1977. (Cited on page xix.)
- [1195] M. D. Shuster, “A Survey of Attitude Representations,” *J. Astron. Sci.*, Vol. 41, pp. 439–517, 1993. (Cited on pages 226 and 674.)
- [1196] M. D. Shuster, “Attitude Analysis in Flatland: The Plane Truth,” *J. Astronautical Sci.*, Vol. 52, pp. 195–209, 2004. (Cited on page 193.)
- [1197] D. D. Siljak, *Large-Scale Dynamic Systems: Stability and Structure*. New York: North-Holland, 1978. (Cited on pages xvii and 706.)
- [1198] F. C. Silva and R. Simoes, “On the Lyapunov and Stein Equations,” *Lin. Alg. Appl.*, Vol. 420, pp. 329–338, 2007. (Cited on page 796.)
- [1199] F. C. Silva and R. Simoes, “On the Lyapunov and Stein Equations, II,” *Lin. Alg. Appl.*, Vol. 426, pp. 305–311, 2007. (Cited on page 796.)
- [1200] S. F. Singer, *Symmetry in Mechanics: A Gentle, Modern Introduction*. Boston: Birkhauser, 2001. (Cited on page xvii.)

- [1201] R. E. Skelton, T. Iwasaki, and K. Grigoriadis, *A Unified Algebraic Approach to Linear Control Design*. London: Taylor and Francis, 1998. (Cited on pages xvii and 722.)
- [1202] D. M. Smiley and M. F. Smiley, "The Polygonal Inequalities," *Amer. Math. Monthly*, Vol. 71, pp. 755–760, 1964. (Cited on page 565.)
- [1203] O. K. Smith, "Eigenvalues of a Symmetric 3×3 Matrix," *Comm. ACM*, Vol. 4, p. 168, 1961. (Cited on page 265.)
- [1204] R. A. Smith, "Matrix Calculations for Lyapunov Quadratic Forms," *J. Diff. Eqns.*, Vol. 2, pp. 208–217, 1966. (Cited on page 797.)
- [1205] A. Smoktunowicz, "Block Matrices and Symmetric Perturbations," *Lin. Alg. Appl.*, Vol. 429, pp. 2628–2635, 2008. (Cited on pages 186 and 593.)
- [1206] R. Snieder, *A Guided Tour of Mathematical Methods for the Physical Sciences*, 2nd ed. Cambridge: Cambridge University Press, 2004. (Cited on page 370.)
- [1207] J. Snyders and M. Zakai, "On Nonnegative Solutions of the Equation $AD + DA' = C$," *SIAM J. Appl. Math.*, Vol. 18, pp. 704–714, 1970. (Cited on pages 722, 746, 747, and 805.)
- [1208] W. So, "Equality Cases in Matrix Exponential Inequalities," *SIAM J. Matrix Anal. Appl.*, Vol. 13, pp. 1154–1158, 1992. (Cited on pages 180, 476, 650, 681, 682, 689, and 693.)
- [1209] W. So, "The High Road to an Exponential Formula," *Lin. Alg. Appl.*, Vol. 379, pp. 69–75, 2004. (Cited on page 688.)
- [1210] W. So and R. C. Thompson, "Product of Exponentials of Hermitian and Complex Symmetric Matrices," *Lin. Multilin. Alg.*, Vol. 29, pp. 225–233, 1991. (Cited on page 688.)
- [1211] W. So and R. C. Thompson, "Singular Values of Matrix Exponentials," *Lin. Multilin. Alg.*, Vol. 47, pp. 249–258, 2000. (Cited on pages 509 and 690.)
- [1212] E. D. Sontag, *Mathematical Control Theory: Deterministic Finite-Dimensional Systems*, 2nd ed. New York: Springer, 1998. (Cited on pages xvii and 793.)
- [1213] E. D. Sontag, "Passivity Gains and the 'Secant Condition' for Stability," *Sys. Contr. Lett.*, Vol. 55, pp. 177–183, 2006. (Cited on page 704.)
- [1214] A. R. Sourour, "A Factorization Theorem for Matrices," *Lin. Multilin. Alg.*, Vol. 19, pp. 141–147, 1986. (Cited on page 351.)
- [1215] A. R. Sourour, "Nilpotent Factorization of Matrices," *Lin. Multilin. Alg.*, Vol. 31, pp. 303–308, 1992. (Cited on page 350.)
- [1216] E. Spiegel, "Sums of Projections," *Lin. Alg. Appl.*, Vol. 187, pp. 239–249, 1993. (Cited on page 361.)
- [1217] M. Spivak, *A Comprehensive Introduction to Differential Geometry*, 3rd ed. Houston: Publish or Perish, 1999. (Cited on pages 227 and 624.)
- [1218] R. P. Stanley, *Enumerative Combinatorics, Volume 1*, 2nd ed. Cambridge: Cambridge University Press, 2000. (Cited on page 11.)
- [1219] W.-H. Steeb, *Matrix Calculus and Kronecker Product with Applications and C++ Programs*. Singapore: World Scientific, 2001. (Cited on page 416.)

- [1220] W.-H. Steeb and F. Wilhelm, "Exponential Functions of Kronecker Products and Trace Calculation," *Lin. Multilin. Alg.*, Vol. 9, pp. 345–346, 1981. (Cited on page 689.)
- [1221] J. M. Steele, *The Cauchy-Schwarz Master Class*. Washington, DC: Mathematical Association of America, 2004. (Cited on page 76.)
- [1222] R. F. Stengel, *Flight Dynamics*. Princeton: Princeton University Press, 2004. (Cited on page xvii.)
- [1223] C. Stepniak, "Ordering of Nonnegative Definite Matrices with Application to Comparison of Linear Models," *Lin. Alg. Appl.*, Vol. 70, pp. 67–71, 1985. (Cited on page 526.)
- [1224] H. J. Stetter, *Numerical Polynomial Algebra*. Philadelphia: SIAM, 2004. (Cited on page xviii.)
- [1225] G. W. Stewart, *Introduction to Matrix Computations*. New York: Academic Press, 1973. (Cited on page xviii.)
- [1226] G. W. Stewart, "On the Perturbation of Pseudo-Inverses, Projections and Linear Least Squares Problems," *SIAM Rev.*, Vol. 19, pp. 634–662, 1977. (Cited on page 618.)
- [1227] G. W. Stewart, *Matrix Algorithms Volume I: Basic Decompositions*. Philadelphia: SIAM, 1998. (Cited on page xviii.)
- [1228] G. W. Stewart, "On the Adjugate Matrix," *Lin. Alg. Appl.*, Vol. 283, pp. 151–164, 1998. (Cited on pages xxxvi and 164.)
- [1229] G. W. Stewart, *Matrix Algorithms Volume II: Eigensystems*. Philadelphia: SIAM, 2001. (Cited on page xviii.)
- [1230] G. W. Stewart and J. Sun, *Matrix Perturbation Theory*. Boston: Academic Press, 1990. (Cited on pages 304, 306, 316, 336, 358, 504, 541, 570, 578, 579, 600, 614, 615, 619, 635, and 636.)
- [1231] E. U. Stickel, "Fast Computation of Matrix Exponential and Logarithm," *Analysis*, Vol. 5, pp. 163–173, 1985. (Cited on page 705.)
- [1232] E. U. Stickel, "An Algorithm for Fast High Precision Computation of Matrix Exponential and Logarithm," *Analysis*, Vol. 10, pp. 85–95, 1990. (Cited on pages 705 and 722.)
- [1233] L. Stiller, "Multilinear Algebra and Chess Endgames," in *Games of No Chance*, R. Nowakowski, Ed. Berkeley: Mathematical Sciences Research Institute, 1996, pp. 151–192. (Cited on page xvii.)
- [1234] J. Stoer, "On the Characterization of Least Upper Bound Norms in Matrix Space," *Numer. Math.*, Vol. 6, pp. 302–314, 1964. (Cited on page 579.)
- [1235] J. Stoer and C. Witzgall, *Convexity and Optimization in Finite Dimensions I*. Berlin: Springer, 1970. (Cited on pages 164 and 635.)
- [1236] K. B. Stolarsky, "Generalizations of the Logarithmic Mean," *J. Math. Anal. Appl.*, Vol. 48, pp. 87–92, 1975. (Cited on page 37.)
- [1237] M. G. Stone, "A Mnemonic for Areas of Polygons," *Amer. Math. Monthly*, Vol. 93, pp. 479–480, 1986. (Cited on page 160.)

- [1238] G. Strang, *Linear Algebra and Its Applications*, 3rd ed. San Diego: Harcourt, Brace, Jovanovich, 1988. (Cited on pages xvii and xix.)
- [1239] G. Strang, "The Fundamental Theorem of Linear Algebra," *Amer. Math. Monthly*, Vol. 100, pp. 848–855, 1993. (Cited on page 164.)
- [1240] G. Strang, "Every Unit Matrix is a LULU," *Lin. Alg. Appl.*, Vol. 265, pp. 165–172, 1997. (Cited on page 347.)
- [1241] G. Strang and K. Borre, *Linear Algebra, Geodesy, and GPS*. Wellesley-Cambridge Press, 1997. (Cited on page xvii.)
- [1242] G. Strang and T. Nguyen, "The Interplay of Ranks of Submatrices," *SIAM Rev.*, Vol. 46, pp. 637–646, 2004. (Cited on pages 125 and 218.)
- [1243] S. Strelitz, "On the Routh-Hurwitz Problem," *Amer. Math. Monthly*, Vol. 84, pp. 542–544, 1977. (Cited on page 698.)
- [1244] R. S. Strichartz, "The Campbell-Baker-Hausdorff-Dynkin Formula and Solutions of Differential Equations," *J. Funct. Anal.*, Vol. 72, pp. 320–345, 1987. (Cited on page 678.)
- [1245] T. Strom, "On Logarithmic Norms," *SIAM J. Numer. Anal.*, Vol. 12, pp. 741–753, 1975. (Cited on page 691.)
- [1246] J. Stuelpnagel, "On the Parametrization of the Three-Dimensional Rotation Group," *SIAM Rev.*, Vol. 6, pp. 422–430, 1964. (Cited on page 674.)
- [1247] G. P. H. Styan, "On Lavoie's Determinantal Inequality," *Lin. Alg. Appl.*, Vol. 37, pp. 77–80, 1981. (Cited on page 392.)
- [1248] R. P. Sullivan, "Products of Nilpotent Matrices," *Lin. Multilin. Alg.*, Vol. 56, pp. 311–317, 2008. (Cited on page 350.)
- [1249] K. N. Swamy, "On Sylvester's Criterion for Positive-Semidefinite Matrices," *IEEE Trans. Autom. Contr.*, Vol. AC-18, p. 306, 1973. (Cited on page 541.)
- [1250] P. Szekeres, *A Course in Modern Mathematical Physics: Groups, Hilbert Space and Differential Geometry*. Cambridge: Cambridge University Press, 2004. (Cited on page 227.)
- [1251] G. Szep, "Simultaneous Triangularization of Projector Matrices," *Acta Math. Hung.*, Vol. 48, pp. 285–288, 1986. (Cited on page 358.)
- [1252] T. Szirtes, *Applied Dimensional Analysis and Modeling*. New York: McGraw-Hill, 1998. (Cited on page xvii.)
- [1253] H. Takagi, T. Miura, T. Kanzo, and S.-E. Takahasi, "A Reconsideration of Hua's Inequality," *J. Ineq. Appl.*, Vol. 2005, pp. 15–23, 2005. (Cited on pages 51 and 566.)
- [1254] Y. Takahashi, S.-E. Takahasi, and S. Wada, "A General Hlawka Inequality and Its Reverse Inequality," *Math. Ineq. Appl.*, preprint. (Cited on page 566.)
- [1255] Y. Tao, "More Results on Singular Value Inequalities of Matrices," *Lin. Alg. Appl.*, Vol. 416, pp. 724–729, 2006. (Cited on page 515.)
- [1256] K. Tapp, *Matrix Groups for Undergraduates*. Providence: American Mathematical Society, 2005. (Cited on pages 193, 222, 223, 226, 227, and 638.)

- [1257] O. Taussky, "Positive-Definite Matrices and Their Role in the Study of the Characteristic Roots of General Matrices," *Adv. Math.*, Vol. 2, pp. 175–186, 1968. (Cited on page 361.)
- [1258] O. Taussky, "The Role of Symmetric Matrices in the Study of General Matrices," *Lin. Alg. Appl.*, Vol. 5, pp. 147–154, 1972. (Cited on page 361.)
- [1259] O. Taussky, "How I Became a Torchbearer for Matrix Theory," *Amer. Math. Monthly*, Vol. 95, pp. 801–812, 1988. (Cited on page 164.)
- [1260] O. Taussky and J. Todd, "Another Look at a Matrix of Mark Kac," *Lin. Alg. Appl.*, Vol. 150, pp. 341–360, 1991. (Cited on page 330.)
- [1261] O. Taussky and H. Zassenhaus, "On the Similarity Transformation between a Matrix and Its Transpose," *Pac. J. Math.*, Vol. 9, pp. 893–896, 1959. (Cited on page 361.)
- [1262] W. Tempelman, "The Linear Algebra of Cross Product Operations," *J. Astron. Sci.*, Vol. 36, pp. 447–461, 1988. (Cited on page 188.)
- [1263] G. ten Have, "Structure of the n th Roots of a Matrix," *Lin. Alg. Appl.*, Vol. 187, pp. 59–66, 1993. (Cited on page 348.)
- [1264] R. E. Terrell, "Solution to 'Exponentials of Certain Hilbert Space Operators'," *SIAM Rev.*, Vol. 34, pp. 498–500, 1992. (Cited on page 683.)
- [1265] R. E. Terrell, "Matrix Exponentials," *SIAM Rev.*, Vol. 38, pp. 313–314, 1996. (Cited on page 684.)
- [1266] R. C. Thompson, "On Matrix Commutators," *J. Washington Acad. Sci.*, Vol. 48, pp. 306–307, 1958. (Cited on page 185.)
- [1267] R. C. Thompson, "A Determinantal Inequality for Positive Definite Matrices," *Canad. Math. Bull.*, Vol. 4, pp. 57–62, 1961. (Cited on page 494.)
- [1268] R. C. Thompson, "Some Matrix Factorization Theorems," *Pac. J. Math.*, Vol. 33, pp. 763–810, 1970. (Cited on page 351.)
- [1269] R. C. Thompson, "Dissipative Matrices and Related Results," *Lin. Alg. Appl.*, Vol. 11, pp. 155–169, 1975. (Cited on page 485.)
- [1270] R. C. Thompson, "A Matrix Inequality," *Comment. Math. Univ. Carolinae*, Vol. 17, pp. 393–397, 1976. (Cited on page 490.)
- [1271] R. C. Thompson, "Matrix Type Metric Inequalities," *Lin. Multilin. Alg.*, Vol. 5, pp. 303–319, 1978. (Cited on pages 68, 70, and 583.)
- [1272] R. C. Thompson, "Proof of a Conjectured Exponential Formula," *Lin. Multilin. Alg.*, Vol. 19, pp. 187–197, 1986. (Cited on page 688.)
- [1273] R. C. Thompson, "Special Cases of a Matrix Exponential Formula," *Lin. Alg. Appl.*, Vol. 107, pp. 283–292, 1988. (Cited on page 688.)
- [1274] R. C. Thompson, "Convergence Proof for Goldberg's Exponential Series," *Lin. Alg. Appl.*, Vol. 121, pp. 3–7, 1989. (Cited on page 678.)
- [1275] R. C. Thompson, "Pencils of Complex and Real Symmetric and Skew Matrices," *Lin. Alg. Appl.*, Vol. 147, pp. 323–371, 1991. (Cited on page 361.)

- [1276] R. C. Thompson, "High, Low, and Quantitative Roads in Linear Algebra," *Lin. Alg. Appl.*, Vol. 162–164, pp. 23–64, 1992. (Cited on page 164.)
- [1277] Y. Tian, "EP Matrices Revisited," preprint. (Cited on pages 178 and 312.)
- [1278] Y. Tian, "The Moore-Penrose Inverses of $m \times n$ Block Matrices and Their Applications," *Lin. Alg. Appl.*, Vol. 283, pp. 35–60, 1998. (Cited on pages 386 and 391.)
- [1279] Y. Tian, "Matrix Representations of Octonions and Their Applications," *Adv. Appl. Clifford Algebras*, Vol. 10, pp. 61–90, 2000. (Cited on page 227.)
- [1280] Y. Tian, "Commutativity of EP Matrices," *IMAGE*, Vol. 27, pp. 25–27, 2001. (Cited on page 382.)
- [1281] Y. Tian, "How to Characterize Equalities for the Moore-Penrose Inverse of a Matrix," *Kyungpook Math. J.*, Vol. 41, pp. 1–15, 2001. (Cited on pages 152, 202, 372, and 375.)
- [1282] Y. Tian, "Some Equalities for Generalized Inverses of Matrix Sums and Block Circulant Matrices," *Archivum Mathematicum (BRNO)*, Vol. 37, pp. 301–306, 2001. (Cited on pages 148, 153, 386, and 393.)
- [1283] Y. Tian, "Some Inequalities for Sums of Matrices," *Scientiae Mathematicae Japonicae*, Vol. 54, pp. 355–361, 2001. (Cited on page 465.)
- [1284] Y. Tian, "How to Express a Parallel Sum of k Matrices," *J. Math. Anal. Appl.*, Vol. 266, pp. 333–341, 2002. (Cited on page 529.)
- [1285] Y. Tian, "The Minimal Rank of the Matrix Expression $A - BX - YC$," *Missouri J. Math. Sci.*, Vol. 14, pp. 40–48, 2002. (Cited on page 387.)
- [1286] Y. Tian, "Upper and Lower Bounds for Ranks of Matrix Expressions Using Generalized Inverses," *Lin. Alg. Appl.*, Vol. 355, pp. 187–214, 2002. (Cited on page 388.)
- [1287] Y. Tian, "A Range Equality for Idempotent Matrix," *IMAGE*, Vol. 30, pp. 26–27, 2003. (Cited on page 199.)
- [1288] Y. Tian, "A Range Equality for the Difference of Orthogonal Projectors," *IMAGE*, Vol. 30, p. 36, 2003. (Cited on page 380.)
- [1289] Y. Tian, "On Mixed-Type Reverse Order Laws for the Moore-Penrose Inverse of a Matrix Product," *Int. J. Math. Math. Sci.*, Vol. 58, pp. 3103–3116, 2004. (Cited on pages 380 and 391.)
- [1290] Y. Tian, "Rank Equalities for Block Matrices and Their Moore-Penrose Inverses," *Houston J. Math.*, Vol. 30, pp. 483–510, 2004. (Cited on page 386.)
- [1291] Y. Tian, "Using Rank Formulas to Characterize Equalities for Moore-Penrose Inverses of Matrix Products," *Appl. Math. Comp.*, Vol. 147, pp. 581–600, 2004. (Cited on page 380.)
- [1292] Y. Tian, "A Range Equality for the Commutator with Two Involutory Matrices," *IMAGE*, Vol. 35, pp. 32–33, 2005. (Cited on page 212.)
- [1293] Y. Tian, "A Range Equality for the Kronecker Product of Matrices," *IMAGE*, Vol. 34, pp. 26–27, 2005. (Cited on page 407.)
- [1294] Y. Tian, "Similarity of Two Block Matrices," *IMAGE*, Vol. 34, pp. 27–32, 2005. (Cited on page 321.)

- [1295] Y. Tian, "The Schur Complement in an Orthogonal Projector," *IMAGE*, Vol. 35, pp. 34–36, 2005. (Cited on page 387.)
- [1296] Y. Tian, "Two Characterizations of an EP Matrix," *IMAGE*, Vol. 34, p. 32, 2005. (Cited on page 372.)
- [1297] Y. Tian, "Inequalities Involving Rank and Kronecker Product," *Amer. Math. Monthly*, Vol. 113, p. 851, 2006. (Cited on page 407.)
- [1298] Y. Tian, "On the Inverse of a Product of Orthogonal Projectors," *Amer. Math. Monthly*, Vol. 113, pp. 467–468, 2006. (Cited on page 381.)
- [1299] Y. Tian, "Two Commutativity Equalities for the Regularized Tikhonov Inverse," *IMAGE*, Vol. 39, p. 31, 2007. (Cited on page 455.)
- [1300] Y. Tian, "Two Equalities for the Moore-Penrose Inverse of a Row Block Matrix," *IMAGE*, Vol. 39, p. 31, 2007. (Cited on page 391.)
- [1301] Y. Tian and S. Cheng, "Some Identities for Moore-Penrose Inverses of Matrix Products," *Lin. Multilin. Alg.*, Vol. 52, pp. 405–420, 2004. (Cited on page 379.)
- [1302] Y. Tian and S. Cheng, "The Moore-Penrose Inverse for Sums of Matrices under Rank Additivity Conditions," *Lin. Multilin. Alg.*, Vol. 53, pp. 45–65, 2005. (Cited on pages 383, 386, and 531.)
- [1303] Y. Tian and Y. Liu, "Extremal Ranks of Some Symmetric Matrix Expressions with Applications," *SIAM J. Matrix Anal. Appl.*, Vol. 28, pp. 890–905, 2006. (Cited on page 387.)
- [1304] Y. Tian and G. P. H. Styan, "Universal Factorization Inequalities for Quaternion Matrices and their Applications," *Math. J. Okayama University*, Vol. 41, pp. 45–62, 1999. (Cited on page 229.)
- [1305] Y. Tian and G. P. H. Styan, "How to Establish Universal Block-Matrix Factorizations," *Elec. J. Lin. Alg.*, Vol. 8, pp. 115–127, 2001. (Cited on pages 135 and 229.)
- [1306] Y. Tian and G. P. H. Styan, "Rank Equalities for Idempotent and Involutory Matrices," *Lin. Alg. Appl.*, Vol. 335, pp. 101–117, 2001. (Cited on pages 202, 203, 210, and 371.)
- [1307] Y. Tian and G. P. H. Styan, "A New Rank Formula for Idempotent Matrices with Applications," *Comment. Math. University Carolinae*, Vol. 43, pp. 379–384, 2002. (Cited on pages 124 and 201.)
- [1308] Y. Tian and G. P. H. Styan, "When Does $\text{rank}(ABC) = \text{rank}(AB) + \text{rank}(BC) - \text{rank}(B)$ Hold?" *Int. J. Math. Educ. Sci. Tech.*, Vol. 33, pp. 127–137, 2002. (Cited on pages 118, 124, 203, 208, 388, and 389.)
- [1309] Y. Tian and G. P. H. Styan, "Rank Equalities for Idempotent Matrices with Applications," *J. Comp. Appl. Math.*, Vol. 191, pp. 77–97, 2006. (Cited on pages 202, 203, and 210.)
- [1310] Y. Tian and Y. Takane, "Schur Complements and Banachiewicz-Schur Forms," *Elec. J. Lin. Alg.*, Vol. 13, pp. 405–418, 2005. (Cited on page 391.)
- [1311] T. Toffoli and J. Quick, "Three-Dimensional Rotations by Three Shears," *Graph. Models Image Process.*, Vol. 59, pp. 89–96, 1997. (Cited on page 347.)

- [1312] M. Tominaga, "A Generalized Reverse Inequality of the Cordes Inequality," *Math. Ineq. Appl.*, Vol. 11, pp. 221–227, 2008. (Cited on page 520.)
- [1313] A. Tonge, "Equivalence Constants for Matrix Norms: A Problem of Goldberg," *Lin. Alg. Appl.*, Vol. 306, pp. 1–13, 2000. (Cited on page 583.)
- [1314] G. E. Trapp, "Hermitian Semidefinite Matrix Means and Related Matrix Inequalities—An Introduction," *Lin. Multilin. Alg.*, Vol. 16, pp. 113–123, 1984. (Cited on page 463.)
- [1315] L. N. Trefethen and D. Bau, *Numerical Linear Algebra*. Philadelphia: SIAM, 1997. (Cited on page xviii.)
- [1316] L. N. Trefethen and M. Embree, *Spectra and Pseudospectra*. Princeton: Princeton University Press, 2005. (Cited on page 691.)
- [1317] D. Trenkler, G. Trenkler, C.-K. Li, and H. J. Werner, "Square Roots and Additivity," *IMAGE*, Vol. 29, p. 30, 2002. (Cited on page 458.)
- [1318] G. Trenkler, "A Trace Inequality," *Amer. Math. Monthly*, Vol. 102, pp. 362–363, 1995. (Cited on page 477.)
- [1319] G. Trenkler, "The Moore-Penrose Inverse and the Vector Product," *Int. J. Math. Educ. Sci. Tech.*, Vol. 33, pp. 431–436, 2002. (Cited on pages 188 and 389.)
- [1320] G. Trenkler, "A Matrix Related to an Idempotent Matrix," *IMAGE*, Vol. 31, pp. 39–40, 2003. (Cited on page 375.)
- [1321] G. Trenkler, "On the Product of Orthogonal Projectors," *IMAGE*, Vol. 31, p. 43, 2003. (Cited on pages 209 and 381.)
- [1322] G. Trenkler, "An Extension of Lagrange's Identity to Matrices," *Int. J. Math. Educ. Sci. Technol.*, Vol. 35, pp. 245–249, 2004. (Cited on page 61.)
- [1323] G. Trenkler, "A Property for the Sum of a Matrix A and its Moore-Penrose Inverse A^+ ," *IMAGE*, Vol. 35, pp. 38–40, 2005. (Cited on page 372.)
- [1324] G. Trenkler, "Factorization of a Projector," *IMAGE*, Vol. 34, pp. 33–35, 2005. (Cited on page 350.)
- [1325] G. Trenkler, "On the Product of Orthogonal Projectors," *IMAGE*, Vol. 35, pp. 42–43, 2005. (Cited on page 458.)
- [1326] G. Trenkler, "Projectors and Similarity," *IMAGE*, Vol. 34, pp. 35–36, 2005. (Cited on pages 375 and 391.)
- [1327] G. Trenkler, "Property of the Cross Product," *IMAGE*, Vol. 34, pp. 36–37, 2005. (Cited on page 188.)
- [1328] G. Trenkler, "A Range Equality for Idempotent Hermitian Matrices," *IMAGE*, Vol. 36, pp. 34–35, 2006. (Cited on page 209.)
- [1329] G. Trenkler, "Necessary and Sufficient Conditions for $A + A^*$ to be a Nonnegative Definite Matrix," *IMAGE*, Vol. 37, pp. 28–30, 2006. (Cited on page 525.)
- [1330] G. Trenkler, "Another Property for the Sum of a Matrix A and its Moore-Penrose Inverse A^+ ," *IMAGE*, Vol. 39, pp. 23–25, 2007. (Cited on page 372.)
- [1331] G. Trenkler, "Characterization of EP-ness," *IMAGE*, Vol. 39, pp. 30–31, 2007. (Cited on page 372.)

- [1332] G. Trenkler and S. Puntanen, "A Multivariate Version of Samuelson's Inequality," *Lin. Alg. Appl.*, Vol. 410, pp. 143–149, 2005. (Cited on pages 51 and 454.)
- [1333] G. Trenkler and D. Trenkler, "The Sherman-Morrison Formula and Eigenvalues of a Special Bordered Matrix," *Acta Math. University Comenianae*, Vol. 74, pp. 255–258, 2005. (Cited on page 263.)
- [1334] G. Trenkler and D. Trenkler, "The Vector Cross Product and 4×4 Skew-Symmetric Matrices," in *Recent Advances in Linear Models and Related Areas*, Shalabh and C. Heumann, Eds. Berlin: Springer, 2008, pp. 95–104. (Cited on pages 188, 264, and 676.)
- [1335] G. Trenkler and H. J. Werner, "Partial Isometry and Idempotent Matrices," *IM-AGE*, Vol. 29, pp. 30–32, 2002. (Cited on page 207.)
- [1336] H. L. Trentelman, A. A. Stoorvogel, and M. L. J. Hautus, *Control Theory for Linear Systems*. New York: Springer, 2001. (Cited on pages xvii and 805.)
- [1337] P. Treuenfels, "The Matrix Equation $X^2 - 2AX + B = 0$," *Amer. Math. Monthly*, Vol. 66, pp. 145–146, 1959. (Cited on page 802.)
- [1338] B. Tromborg and S. Waldenstrom, "Bounds on the Diagonal Elements of a Unitary Matrix," *Lin. Alg. Appl.*, Vol. 20, pp. 189–195, 1978. (Cited on page 196.)
- [1339] M. Tsatsomeros, J. S. Maybee, D. D. Olesky, and P. Van Den Driessche, "Nullspaces of Matrices and Their Compounds," *Lin. Multilin. Alg.*, Vol. 34, pp. 291–300, 1993. (Cited on page 338.)
- [1340] N.-K. Tsing and F. Uhlig, "Inertia, Numerical Range, and Zeros of Quadratic Forms for Matrix Pencils," *SIAM J. Matrix Anal. Appl.*, Vol. 12, pp. 146–159, 1991. (Cited on pages 362 and 504.)
- [1341] P. Tsiotras, "Further Passivity Results for the Attitude Control Problem," *IEEE Trans. Autom. Contr.*, Vol. 43, pp. 1597–1600, 1998. (Cited on page 188.)
- [1342] P. Tsiotras, J. L. Junkins, and H. Schaub, "Higher-Order Cayley Transforms with Applications to Attitude Representations," *J. Guid. Contr. Dyn.*, Vol. 20, pp. 528–534, 1997. (Cited on page 191.)
- [1343] S. H. Tung, "On Lower and Upper Bounds of the Difference between the Arithmetic and the Geometric Mean," *Math. Comput.*, Vol. 29, pp. 834–836, 1975. (Cited on page 52.)
- [1344] D. A. Turkington, *Matrix Calculus and Zero-One Matrices*. Cambridge: Cambridge University Press, 2002. (Cited on page 416.)
- [1345] R. Turkmen and D. Bozkurt, "A Note for Bounds of Norms of Hadamard Product of Matrices," *Math. Ineq. Appl.*, Vol. 9, pp. 211–217, 2006. (Cited on page 615.)
- [1346] H. W. Turnbull, *The Theory of Determinants, Matrices and Invariants*. London: Blackie, 1950. (Cited on page 164.)
- [1347] G. M. Tuynman, "The Derivation of the Exponential Map of Matrices," *Amer. Math. Monthly*, Vol. 102, pp. 818–820, 1995. (Cited on page 682.)
- [1348] F. Tyan and D. S. Bernstein, "Global Stabilization of Systems Containing a Double Integrator Using a Saturated Linear Controller," *Int. J. Robust Nonlin. Contr.*, Vol. 9, pp. 1143–1156, 1999. (Cited on page 796.)

- [1349] M. Uchiyama, "Norms and Determinants of Products of Logarithmic Functions of Positive Semi-Definite Operators," *Math. Ineq. Appl.*, Vol. 1, pp. 279–284, 1998. (Cited on pages 522 and 694.)
- [1350] M. Uchiyama, "Some Exponential Operator Inequalities," *Math. Ineq. Appl.*, Vol. 2, pp. 469–471, 1999. (Cited on page 688.)
- [1351] F. E. Udwardia and R. E. Kalaba, *Analytical Dynamics: A New Approach*. Cambridge: Cambridge University Press, 1996. (Cited on page xvii.)
- [1352] F. Uhlig, "A Recurring Theorem About Pairs of Quadratic Forms and Extensions: A Survey," *Lin. Alg. Appl.*, Vol. 25, pp. 219–237, 1979. (Cited on pages 362, 504, 507, and 508.)
- [1353] F. Uhlig, "On the Matrix Equation $AX = B$ with Applications to the Generators of a Controllability Matrix," *Lin. Alg. Appl.*, Vol. 85, pp. 203–209, 1987. (Cited on page 117.)
- [1354] F. Uhlig, "Constructive Ways for Generating (Generalized) Real Orthogonal Matrices as Products of (Generalized) Symmetries," *Lin. Alg. Appl.*, Vol. 332–334, pp. 459–467, 2001. (Cited on page 347.)
- [1355] F. A. Valentine, *Convex Sets*. New York: McGraw-Hill, 1964. (Cited on page 164.)
- [1356] M. Van Barel, V. Ptak, and Z. Vavrin, "Bezout and Hankel Matrices Associated with Row Reduced Matrix Polynomials, Barnett-Type Formulas," *Lin. Alg. Appl.*, Vol. 332–334, pp. 583–606, 2001. (Cited on page 257.)
- [1357] R. van der Merwe, *Sigma-Point Kalman Filters for Probabilistic Inference in Dynamic State-Space Models*. Portland: Oregon Health and Science University, 2004, Ph.D. Dissertation. (Cited on page 676.)
- [1358] P. Van Dooren, "The Computation of Kronecker's Canonical Form of a Singular Pencil," *Lin. Alg. Appl.*, Vol. 27, pp. 103–140, 1979. (Cited on page 362.)
- [1359] C. F. Van Loan, "Computing Integrals Involving the Matrix Exponential," *IEEE Trans. Autom. Contr.*, Vol. AC-23, pp. 395–404, 1978. (Cited on page 681.)
- [1360] C. F. Van Loan, "How Near Is a Stable Matrix to an Unstable Matrix," *Contemporary Math.*, Vol. 47, pp. 465–478, 1985. (Cited on page 701.)
- [1361] C. F. Van Loan, *Computational Frameworks for the Fast Fourier Transform*. Philadelphia: SIAM, 1992. (Cited on pages xvii and 357.)
- [1362] C. F. Van Loan, "The Ubiquitous Kronecker Product," *J. Comp. Appl. Math.*, Vol. 123, pp. 85–100, 2000. (Cited on page 416.)
- [1363] P. Van Overschee and B. De Moor, *Subspace Identification for Linear Systems: Theory, Implementation, Applications*. Dordrecht: Kluwer, 1996. (Cited on page 805.)
- [1364] R. Vandebril and M. Van Barel, "A Note on the Nullity Theorem," *J. Comp. Appl. Math.*, Vol. 189, pp. 179–190, 2006. (Cited on page 125.)
- [1365] R. Vandebril, M. Van Barel, and N. Mastronardi, *Matrix Computations and Semiseparable Matrices*. Baltimore: Johns Hopkins University Press, 2008. (Cited on pages 125 and 129.)

- [1366] V. S. Varadarajan, *Lie Groups, Lie Algebras, and Their Representations*. New York: Springer, 1984. (Cited on pages 230, 655, 682, and 722.)
- [1367] J. M. Varah, "A Lower Bound for the Smallest Singular Value of a Matrix," *Lin. Alg. Appl.*, Vol. 11, pp. 3–5, 1975. (Cited on page 606.)
- [1368] A. I. G. Vardulakis, *Linear Multivariable Control: Algebraic Analysis and Synthesis Methods*. Chichester: Wiley, 1991. (Cited on pages xvii and 281.)
- [1369] R. S. Varga, *Matrix Iterative Analysis*. Englewood Cliffs: Prentice-Hall, 1962. (Cited on pages xviii and 275.)
- [1370] R. S. Varga, *Gersgorin and His Circles*. New York: Springer, 2004. (Cited on page 269.)
- [1371] P. R. Vein, "A Short Survey of Some Recent Applications of Determinants," *Lin. Alg. Appl.*, Vol. 42, pp. 287–297, 1982. (Cited on page 164.)
- [1372] R. Vein and P. Dale, *Determinants and Their Applications in Mathematical Physics*. New York: Springer, 1999. (Cited on pages xvii and 164.)
- [1373] D. Veljan, "The Sine Theorem and Inequalities for Volumes of Simplices and Determinants," *Lin. Alg. Appl.*, Vol. 219, pp. 79–91, 1995. (Cited on page 160.)
- [1374] D. Veljan and S. Wu, "Parametrized Klamkin's Inequality and Improved Euler's Inequality," *Math. Ineq. Appl.*, preprint. (Cited on pages 42 and 158.)
- [1375] J. Vermeer, "Orthogonal Similarity of a Real Matrix and Its Transpose," *Lin. Alg. Appl.*, Vol. 428, pp. 382–392, 2008. (Cited on page 345.)
- [1376] K. Veselic, "On Real Eigenvalues of Real Tridiagonal Matrices," *Lin. Alg. Appl.*, Vol. 27, pp. 167–171, 1979. (Cited on page 330.)
- [1377] K. Veselic, "Estimating the Operator Exponential," *Lin. Alg. Appl.*, Vol. 280, pp. 241–244, 1998. (Cited on page 701.)
- [1378] K. Veselic, "Bounds for Exponentially Stable Semigroups," *Lin. Alg. Appl.*, Vol. 358, pp. 309–333, 2003. (Cited on page 701.)
- [1379] W. J. Vetter, "Matrix Calculus Operations and Taylor Expansions," *SIAM Rev.*, Vol. 15, pp. 352–369, 1973. (Cited on page 416.)
- [1380] M. Vidyasagar, "On Matrix Measures and Convex Liapunov Functions," *J. Math. Anal. Appl.*, Vol. 62, pp. 90–103, 1978. (Cited on page 691.)
- [1381] G. Visick, "A Weak Majorization Involving the Matrices $A \circ B$ and AB ," *Lin. Alg. Appl.*, Vol. 223/224, pp. 731–744, 1995. (Cited on page 535.)
- [1382] G. Visick, "Majorizations of Hadamard Products of Matrix Powers," *Lin. Alg. Appl.*, Vol. 269, pp. 233–240, 1998. (Cited on pages 535 and 540.)
- [1383] G. Visick, "A Quantitative Version of the Observation that the Hadamard Product Is a Principal Submatrix of the Kronecker Product," *Lin. Alg. Appl.*, Vol. 304, pp. 45–68, 2000. (Cited on pages 532, 537, and 539.)
- [1384] G. Visick, "Another Inequality for Hadamard Products," *IMAGE*, Vol. 29, pp. 32–33, 2002. (Cited on page 532.)
- [1385] S.-W. Vong and X.-Q. Jin, "Proof of Bottcher and Wenzel's Conjecture," *Oper. Matrices*, Vol. 2, pp. 435–442, 2008. (Cited on page 584.)

- [1386] S. Wada, "On Some Refinement of the Cauchy-Schwarz Inequality," *Lin. Alg. Appl.*, Vol. 420, pp. 433–440, 2007. (Cited on pages 60, 63, and 64.)
- [1387] G. G. Walter and M. Contreras, *Compartmental Modeling with Networks*. Boston: Birkhauser, 1999. (Cited on pages xvii and 708.)
- [1388] B. Wang and F. Zhang, "Some Inequalities for the Eigenvalues of the Product of Positive Semidefinite Hermitian Matrices," *Lin. Alg. Appl.*, Vol. 160, pp. 113–118, 1992. (Cited on pages 518, 519, and 614.)
- [1389] B.-Y. Wang and M.-P. Gong, "Some Eigenvalue Inequalities for Positive Semidefinite Matrix Power Products," *Lin. Alg. Appl.*, Vol. 184, pp. 249–260, 1993. (Cited on page 519.)
- [1390] B.-Y. Wang, B.-Y. Xi, and F. Zhang, "Some Inequalities for Sum and Product of Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 293, pp. 39–49, 1999. (Cited on pages 528 and 614.)
- [1391] B.-Y. Wang and F. Zhang, "A Trace Inequality for Unitary Matrices," *Amer. Math. Monthly*, Vol. 101, pp. 453–455, 1994. (Cited on page 198.)
- [1392] B.-Y. Wang and F. Zhang, "Trace and Eigenvalue Inequalities for Ordinary and Hadamard Products of Positive Semidefinite Hermitian Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 16, pp. 1173–1183, 1995. (Cited on pages 519 and 534.)
- [1393] B.-Y. Wang and F. Zhang, "Schur Complements and Matrix Inequalities of Hadamard Products," *Lin. Multilin. Alg.*, Vol. 43, pp. 315–326, 1997. (Cited on page 541.)
- [1394] C.-L. Wang, "On Development of Inverses of the Cauchy and Hölder Inequalities," *SIAM Rev.*, Vol. 21, pp. 550–557, 1979. (Cited on pages 57 and 65.)
- [1395] D. Wang, "The Polar Decomposition and a Matrix Inequality," *Amer. Math. Monthly*, Vol. 96, pp. 517–519, 1989. (Cited on page 433.)
- [1396] G. Wang, Y. Wei, and S. Qiao, *Generalized Inverses: Theory and Computations*. Beijing/New York: Science Press, 2004. (Cited on pages 129, 202, 378, and 398.)
- [1397] J.-H. Wang, "The Length Problem for a Sum of Idempotents," *Lin. Alg. Appl.*, Vol. 215, pp. 135–159, 1995. (Cited on page 361.)
- [1398] L.-C. Wang and C.-L. Li, "On Some New Mean Value Inequalities," *J. Ineq. Pure Appl. Math.*, Vol. 8, no. 3, pp. 1–8, 2007, Article 87. (Cited on page 58.)
- [1399] Q.-G. Wang, "Necessary and Sufficient Conditions for Stability of a Matrix Polytope with Normal Vertex Matrices," *Automatica*, Vol. 27, pp. 887–888, 1991. (Cited on page 271.)
- [1400] X. Wang, "A Simple Proof of Descartes's Rule of Signs," *Amer. Math. Monthly*, Vol. 111, pp. 525–526, 2004. (Cited on page 696.)
- [1401] Y. W. Wang and D. S. Bernstein, "Controller Design with Regional Pole Constraints: Hyperbolic and Horizontal Strip Regions," *AIAA J. Guid. Contr. Dyn.*, Vol. 16, pp. 784–787, 1993. (Cited on page 798.)
- [1402] Y. W. Wang and D. S. Bernstein, " L_2 Controller Synthesis with L_∞ -Bounded Closed-Loop Impulse Response," *Int. J. Contr.*, Vol. 60, pp. 1295–1306, 1994. (Cited on page 689.)

- [1403] A. J. B. Ward and F. Gerrish, "A Constructive Proof by Elementary Transformations of Roth's Removal Theorems in the Theory of Matrix Equations," *Int. J. Math. Educ. Sci. Tech.*, Vol. 31, pp. 425–429, 2000. (Cited on page 320.)
- [1404] J. Warga, *Optimal Control of Differential and Functional Equations*. New York: Academic Press, 1972. (Cited on pages 625 and 626.)
- [1405] W. E. Waterhouse, "A Determinant Identity with Matrix Entries," *Amer. Math. Monthly*, Vol. 97, pp. 249–250, 1990. (Cited on pages 136 and 137.)
- [1406] W. Watkins, "Convex Matrix Functions," *Proc. Amer. Math. Soc.*, Vol. 44, pp. 31–34, 1974. (Cited on pages 488 and 536.)
- [1407] W. Watkins, "Generating Functions," *Coll. Math. J.*, Vol. 18, pp. 195–211, 1987. (Cited on page 278.)
- [1408] W. Watkins, "A Determinantal Inequality for Correlation Matrices," *Lin. Alg. Appl.*, Vol. 104, pp. 59–63, 1988. (Cited on page 533.)
- [1409] G. S. Watson, G. Alpargu, and G. P. H. Styan, "Some Comments on Six Inequalities Associated with the Inefficiency of Ordinary Least Squares with One Regressor," *Lin. Alg. Appl.*, Vol. 264, pp. 13–54, 1997. (Cited on pages 57 and 66.)
- [1410] J. R. Weaver, "Centrosymmetric (Cross-Symmetric) Matrices, Their Basic Properties, Eigenvalues, and Eigenvectors," *Amer. Math. Monthly*, Vol. 92, pp. 711–717, 1985. (Cited on pages 230 and 314.)
- [1411] J. H. Webb, "An Inequality in Algebra and Geometry," *College Math. J.*, Vol. 36, p. 164, 2005. (Cited on page 45.)
- [1412] R. Webster, *Convexity*. Oxford: Oxford University Press, 1994. (Cited on page 164.)
- [1413] E. Wegert and L. N. Trefethen, "From the Buffon Needle Problem to the Kreiss Matrix Theorem," *Amer. Math. Monthly*, Vol. 101, pp. 132–139, 1994. (Cited on page 715.)
- [1414] J. Wei and E. Norman, "Lie Algebraic Solution of Linear Differential Equations," *J. Math. Phys.*, Vol. 4, pp. 575–581, 1963. (Cited on page 678.)
- [1415] J. Wei and E. Norman, "On Global Representations of the Solutions of Linear Differential Equations as a Product of Exponentials," *Proc. Amer. Math. Soc.*, Vol. 15, pp. 327–334, 1964. (Cited on page 678.)
- [1416] M. Wei, "Reverse Order Laws for Generalized Inverses of Multiple Matrix Products," *Lin. Alg. Appl.*, Vol. 293, pp. 273–288, 1999. (Cited on page 380.)
- [1417] Y. Wei, "A Characterization and Representation of the Drazin Inverse," *SIAM J. Matrix Anal. Appl.*, Vol. 17, pp. 744–747, 1996. (Cited on page 393.)
- [1418] Y. Wei, "Expressions for the Drazin Inverse of a 2×2 Block Matrix," *Lin. Multilin. Alg.*, Vol. 45, pp. 131–146, 1998. (Cited on page 391.)
- [1419] G. H. Weiss and A. A. Maradudin, "The Baker-Hausdorff Formula and a Problem in Crystal Physics," *J. Math. Phys.*, Vol. 3, pp. 771–777, 1962. (Cited on page 678.)
- [1420] E. M. E. Wermuth, "Two Remarks on Matrix Exponentials," *Lin. Alg. Appl.*, Vol. 117, pp. 127–132, 1989. (Cited on pages 681 and 694.)

- [1421] E. M. E. Wermuth, "A Remark on Commuting Operator Exponentials," *Proc. Amer. Math. Soc.*, Vol. 125, pp. 1685–1688, 1997. (Cited on page 681.)
- [1422] H.-J. Werner, "On the Matrix Monotonicity of Generalized Inversion," *Lin. Alg. Appl.*, Vol. 27, pp. 141–145, 1979. (Cited on page 527.)
- [1423] H. J. Werner, "On the Product of Orthogonal Projectors," *IMAGE*, Vol. 32, pp. 30–37, 2004. (Cited on pages 209, 375, and 380.)
- [1424] H. J. Werner, "Solution 34-4.2 to 'A Range Equality for the Commutator with Two Involutory Matrices'," *IMAGE*, Vol. 35, pp. 32–33, 2005. (Cited on page 200.)
- [1425] J. R. Wertz and W. J. Larson, Eds., *Space Mission Analysis and Design*. Dordrecht: Kluwer, 1999. (Cited on page 161.)
- [1426] P. Wesseling, *Principles of Computational Fluid Dynamics*. Berlin: Springer, 2001. (Cited on page xvii.)
- [1427] J. R. Westlake, *A Handbook of Numerical Matrix Inversion and Solution of Linear Equations*. New York: Wiley, 1968. (Cited on page xviii.)
- [1428] N. A. Wiegmann, "Normal Products of Matrices," *Duke Math. J.*, Vol. 15, pp. 633–638, 1948. (Cited on page 343.)
- [1429] Z. Wiener, "An Interesting Matrix Exponent Formula," *Lin. Alg. Appl.*, Vol. 257, pp. 307–310, 1997. (Cited on page 714.)
- [1430] E. P. Wigner and M. M. Yanase, "On the Positive Semidefinite Nature of a Certain Matrix Expression," *Canad. J. Math.*, Vol. 16, pp. 397–406, 1964. (Cited on page 457.)
- [1431] R. M. Wilcox, "Exponential Operators and Parameter Differentiation in Quantum Physics," *J. Math. Phys.*, Vol. 8, pp. 962–982, 1967. (Cited on pages xviii and 684.)
- [1432] J. B. Wilker, J. S. Sumner, A. A. Jagers, M. Vowe, and J. Anglesio, "Inequalities Involving Trigonometric Functions," *Amer. Math. Monthly*, Vol. 98, pp. 264–266, 1991. (Cited on page 29.)
- [1433] J. H. Wilkinson, *The Algebraic Eigenvalue Problem*. London: Oxford University Press, 1965. (Cited on page xviii.)
- [1434] J. C. Willems, "Least Squares Stationary Optimal Control and the Algebraic Riccati Equation," *IEEE Trans. Autom. Contr.*, Vol. AC-16, pp. 621–634, 1971. (Cited on page 805.)
- [1435] D. A. Wilson, "Convolution and Hankel Operator Norms for Linear Systems," *IEEE Trans. Autom. Contr.*, Vol. AC-34, pp. 94–97, 1989. (Cited on page 619.)
- [1436] P. M. H. Wilson, *Curved Spaces: From Classical Geometries to Elementary Differential Geometry*. Cambridge: Cambridge University Press, 2008. (Cited on pages 161 and 193.)
- [1437] H. K. Wimmer, "Inertia Theorems for Matrices, Controllability, and Linear Vibrations," *Lin. Alg. Appl.*, Vol. 8, pp. 337–343, 1974. (Cited on pages 793 and 796.)
- [1438] H. K. Wimmer, "The Algebraic Riccati Equation without Complete Controllability," *SIAM J. Alg. Disc. Math.*, Vol. 3, pp. 1–12, 1982. (Cited on page 805.)
- [1439] H. K. Wimmer, "On Ostrowski's Generalization of Sylvester's Law of Inertia," *Lin. Alg. Appl.*, Vol. 52/53, pp. 739–741, 1983. (Cited on page 310.)

- [1440] H. K. Wimmer, "The Algebraic Riccati Equation: Conditions for the Existence and Uniqueness of Solutions," *Lin. Alg. Appl.*, Vol. 58, pp. 441–452, 1984. (Cited on page 805.)
- [1441] H. K. Wimmer, "Monotonicity of Maximal Solutions of Algebraic Riccati Equations," *Sys. Contr. Lett.*, Vol. 5, pp. 317–319, 1985. (Cited on pages 788 and 805.)
- [1442] H. K. Wimmer, "Extremal Problems for Hölder Norms of Matrices and Realizations of Linear Systems," *SIAM J. Matrix Anal. Appl.*, Vol. 9, pp. 314–322, 1988. (Cited on pages 459 and 585.)
- [1443] H. K. Wimmer, "Linear Matrix Equations, Controllability and Observability, and the Rank of Solutions," *SIAM J. Matrix Anal. Appl.*, Vol. 9, pp. 570–578, 1988. (Cited on page 796.)
- [1444] H. K. Wimmer, "On the History of the Bezoutian and the Resultant Matrix," *Lin. Alg. Appl.*, Vol. 128, pp. 27–34, 1990. (Cited on page 257.)
- [1445] H. K. Wimmer, "Normal Forms of Symplectic Pencils and the Discrete-Time Algebraic Riccati Equation," *Lin. Alg. Appl.*, Vol. 147, pp. 411–440, 1991. (Cited on page 805.)
- [1446] H. K. Wimmer, "Lattice Properties of Sets of Semidefinite Solutions of Continuous-Time Algebraic Riccati Equations," *Automatica*, Vol. 31, pp. 173–182, 1995. (Cited on page 805.)
- [1447] A. Witkowski, "A New Proof of the Monotonicity of Power Means," *J. Ineq. Pure Appl. Math.*, Vol. 5, no. 1, pp. 1–2, 2004, Article 6. (Cited on pages 23 and 27.)
- [1448] H. Wolkowicz and G. P. H. Styan, "Bounds for Eigenvalues Using Traces," *Lin. Alg. Appl.*, Vol. 29, pp. 471–506, 1980. (Cited on pages 51, 322, and 332.)
- [1449] H. Wolkowicz and G. P. H. Styan, "More Bounds for Eigenvalues Using Traces," *Lin. Alg. Appl.*, Vol. 31, pp. 1–17, 1980. (Cited on page 332.)
- [1450] W. A. Wolovich, *Linear Multivariable Systems*. New York: Springer, 1974. (Cited on page 805.)
- [1451] M. J. Wonenburger, "A Decomposition of Orthogonal Transformations," *Canad. Math. Bull.*, Vol. 7, pp. 379–383, 1964. (Cited on page 350.)
- [1452] M. J. Wonenburger, "Transformations Which are Products of Two Involutions," *J. Math. Mech.*, Vol. 16, pp. 327–338, 1966. (Cited on page 350.)
- [1453] C. S. Wong, "Characterizations of Products of Symmetric Matrices," *Lin. Alg. Appl.*, Vol. 42, pp. 243–251, 1982. (Cited on page 350.)
- [1454] E. Wong, Ed., *Stochastic Processes in Information and Dynamical Systems*. New York: McGraw-Hill, 1971. (Cited on page 447.)
- [1455] W. M. Wonham, *Linear Multivariable Control: A Geometric Approach*, 2nd ed. New York: Springer, 1979. (Cited on pages xvii, 262, 646, 791, and 805.)
- [1456] C.-F. Wu, "On Some Ordering Properties of the Generalized Inverses of Nonnegative Definite Matrices," *Lin. Alg. Appl.*, Vol. 32, pp. 49–60, 1980. (Cited on page 527.)
- [1457] P. Y. Wu, "Products of Nilpotent Matrices," *Lin. Alg. Appl.*, Vol. 96, pp. 227–232, 1987. (Cited on page 350.)

- [1458] P. Y. Wu, "Products of Positive Semidefinite Matrices," *Lin. Alg. Appl.*, Vol. 111, pp. 53–61, 1988. (Cited on pages 350 and 542.)
- [1459] P. Y. Wu, "The Operator Factorization Problems," *Lin. Alg. Appl.*, Vol. 117, pp. 35–63, 1989. (Cited on pages 349, 350, 351, and 701.)
- [1460] P. Y. Wu, "Sums of Idempotent Matrices," *Lin. Alg. Appl.*, Vol. 142, pp. 43–54, 1990. (Cited on pages 360 and 361.)
- [1461] S. Wu, "Note on a Conjecture of R. A. Satnoianu," *Math. Ineq. Appl.*, preprint. (Cited on page 59.)
- [1462] S. Wu, "A Further Generalization of Aczel's Inequality and Popoviciu's Inequality," *Math. Ineq. Appl.*, Vol. 10, pp. 565–573, 2007. (Cited on page 64.)
- [1463] S. Wu, "A Sharpened Version of the Fundamental Triangle Inequality," *Math. Ineq. Appl.*, Vol. 11, pp. 477–482, 2008. (Cited on page 158.)
- [1464] Y.-D. Wu, Z.-H. Zhang, and V. Loksha, "Sharpening on Mircea's Inequality," *J. Ineq. Pure Appl. Math.*, Vol. 8, no. 4, pp. 1–6, 2007, Article 116. (Cited on page 158.)
- [1465] Z.-G. Xiao and Z.-H. Zhang, "The Inequalities $G \leq L \leq I \leq A$ in n Variables," *J. Ineq. Pure Appl. Math.*, Vol. 4, no. 2/39, pp. 1–6, 2003. (Cited on page 54.)
- [1466] C. Xu, "Bellman's Inequality," *Lin. Alg. Appl.*, Vol. 229, pp. 9–14, 1995. (Cited on page 480.)
- [1467] C. Xu, Z. Xu, and F. Zhang, "Revisiting Hua-Marcus-Bellman-Ando Inequalities on Contractive Matrices," *Lin. Alg. Appl.*, preprint. (Cited on pages 70, 145, 472, 490, and 565.)
- [1468] H. Xu, "Two Results About the Matrix Exponential," *Lin. Alg. Appl.*, Vol. 262, pp. 99–109, 1997. (Cited on pages 650 and 701.)
- [1469] A. Yakub, "Symmetric Inequality," *Amer. Math. Monthly*, Vol. 114, p. 649, 2007. (Cited on page 45.)
- [1470] T. Yamazaki, "An Extension of Specht's Theorem via Kantorovich Inequality and Related Results," *Math. Ineq. Appl.*, Vol. 3, pp. 89–96, 2000. (Cited on page 53.)
- [1471] T. Yamazaki, "Further Characterizations of Chaotic Order via Specht's Ratio," *Math. Ineq. Appl.*, Vol. 3, pp. 259–268, 2000. (Cited on page 522.)
- [1472] B. Yang, "On a New Inequality Similar to Hardy-Hilbert's Inequality," *Math. Ineq. Appl.*, Vol. 6, pp. 37–44, 2003. (Cited on page 63.)
- [1473] B. Yang and T. M. Rassias, "On the Way of Weight Coefficient and Research for the Hilbert-Type Inequalities," *Math. Ineq. Appl.*, Vol. 6, pp. 625–658, 2003. (Cited on page 63.)
- [1474] X. Yang, "Necessary Conditions of Hurwitz Polynomials," *Lin. Alg. Appl.*, Vol. 359, pp. 21–27, 2003. (Cited on page 697.)
- [1475] X. Yang, "Some Necessary Conditions for Hurwitz Stability," *Automatica*, Vol. 40, pp. 527–529, 2004. (Cited on page 697.)
- [1476] Z. P. Yang and X. X. Feng, "A Note on the Trace Inequality for Products of Hermitian Matrix Power," *J. Ineq. Pure Appl. Math.*, Vol. 3, no. 5/78, pp. 1–12, 2002. (Cited on pages 476, 478, and 480.)

- [1477] S. F. Yau and Y. Bresler, "A Generalization of Bergstrom's Inequality and Some Applications," *Lin. Alg. Appl.*, Vol. 161, pp. 135–151, 1992. (Cited on pages 535 and 538.)
- [1478] D. M. Young, *Iterative Solution of Large Linear Systems*. New York: Academic Press, 1971, reprinted by Dover, Mineola, 2003. (Cited on page xviii.)
- [1479] N. J. Young, "Matrices which Maximise any Analytic Function," *Acta Math. Hung.*, Vol. 34, pp. 239–243, 1979. (Cited on page 607.)
- [1480] A. Zanna and H. Z. Munthe-Kaas, "Generalized Polar Decompositions for the Approximation of the Matrix Exponential," *SIAM J. Matrix Anal. Appl.*, Vol. 23, pp. 840–862, 2002. (Cited on page 677.)
- [1481] X. Zhan, "Inequalities for the Singular Values of Hadamard Products," *SIAM J. Matrix Anal. Appl.*, Vol. 18, pp. 1093–1095, 1997. (Cited on page 615.)
- [1482] X. Zhan, "Inequalities for Unitarily Invariant Norms," *SIAM J. Matrix Anal. Appl.*, Vol. 20, pp. 466–470, 1998. (Cited on page 448.)
- [1483] X. Zhan, "Singular Values of Differences of Positive Semidefinite Matrices," *SIAM J. Matrix Anal. Appl.*, Vol. 22, pp. 819–823, 2000. (Cited on pages 515 and 590.)
- [1484] X. Zhan, "Span of the Orthogonal Orbit of Real Matrices," *Lin. Multilin. Alg.*, Vol. 49, pp. 337–346, 2001. (Cited on page 360.)
- [1485] X. Zhan, *Matrix Inequalities*. New York: Springer, 2002. (Cited on pages 164, 327, 434, 442, 447, 448, 509, 510, 511, 521, 529, 530, 531, 535, 537, 539, 540, 541, 549, 582, 612, and 685.)
- [1486] X. Zhan, "On Some Matrix Inequalities," *Lin. Alg. Appl.*, Vol. 376, pp. 299–303, 2004. (Cited on page 516.)
- [1487] F. Zhang, *Linear Algebra: Challenging Problems for Students*. Baltimore: Johns Hopkins University Press, 1996. (Cited on pages 131, 184, and 185.)
- [1488] F. Zhang, "Quaternions and Matrices of Quaternions," *Lin. Alg. Appl.*, Vol. 251, pp. 21–57, 1997. (Cited on page 229.)
- [1489] F. Zhang, "A Compound Matrix with Positive Determinant," *Amer. Math. Monthly*, Vol. 105, p. 958, 1998. (Cited on page 153.)
- [1490] F. Zhang, *Matrix Theory: Basic Results and Techniques*. New York: Springer, 1999. (Cited on pages 126, 127, 219, 312, 342, 357, 456, 465, 471, 476, 478, 489, 541, and 565.)
- [1491] F. Zhang, "Schur Complements and Matrix Inequalities in the Lowner Ordering," *Lin. Alg. Appl.*, Vol. 321, pp. 399–410, 2000. (Cited on page 490.)
- [1492] F. Zhang, "Matrix Inequalities by Means of Block Matrices," *Math. Ineq. Appl.*, Vol. 4, pp. 481–490, 2001. (Cited on pages 452, 469, 493, 508, 525, 533, 534, and 539.)
- [1493] F. Zhang, "Inequalities Involving Square Roots," *IMAGE*, Vol. 29, pp. 33–34, 2002. (Cited on page 458.)
- [1494] F. Zhang, "Block Matrix Techniques," in *The Schur Complement and Its Applications*, F. Zhang, Ed. New York: Springer, 2004, pp. 83–110. (Cited on pages 453, 483, 493, 501, 505, 508, and 510.)

- [1495] F. Zhang, "A Matrix Identity on the Schur Complement," *Lin. Multilin. Alg.*, Vol. 52, pp. 367–373, 2004. (Cited on page 393.)
- [1496] L. Zhang, "A Characterization of the Drazin Inverse," *Lin. Alg. Appl.*, Vol. 335, pp. 183–188, 2001. (Cited on page 393.)
- [1497] Q.-C. Zhong, "J-Spectral Factorization of Regular Para-Hermitian Transfer Matrices," *Automatica*, Vol. 41, pp. 1289–1293, 2005. (Cited on page 206.)
- [1498] K. Zhou, *Robust and Optimal Control*. Upper Saddle River: Prentice-Hall, 1996. (Cited on pages xvii, 281, 607, 610, and 787.)
- [1499] L. Zhu, "A New Simple Proof of Wilker's Inequality," *Math. Ineq. Appl.*, Vol. 8, pp. 749–750, 2005. (Cited on page 29.)
- [1500] L. Zhu, "On Wilker-Type Inequalities," *Math. Ineq. Appl.*, Vol. 10, pp. 727–731, 2005. (Cited on page 29.)
- [1501] G. Zielke, "Some Remarks on Matrix Norms, Condition Numbers, and Error Estimates for Linear Equations," *Lin. Alg. Appl.*, Vol. 110, pp. 29–41, 1988. (Cited on pages 573 and 592.)
- [1502] S. Zlobec, "An Explicit Form of the Moore-Penrose Inverse of an Arbitrary Complex Matrix," *SIAM Rev.*, Vol. 12, pp. 132–134, 1970. (Cited on page 366.)
- [1503] D. Zwillinger, *Standard Mathematical Tables and Formulae*, 31st ed. Boca Raton: Chapman & Hall/CRC, 2003. (Cited on pages 75, 158, 161, 255, and 357.)

Author Index

- Abdessemed, A. 584
Ablamowicz, R. 673
Abou-Kandil, H. xvii
Abramovich, S. 30
Aceto, L. 354, 362, 447, 672
Afriat, S. 210
Agaev, R. 708
Ahlbrandt, C. D. 805
Ahn, E. 683
Aitken, A. C. xix
Aivazis, M. 190
Akdeniz, F. 526, 527
Al-Ahmar, M. 196
Albert, A. A. 313
Albert, A. E. 380, 382, 385, 390, 541, 618
Aldrovandi, R. xvii, 275, 357
Aleksiejczuk, M. 391
Alfakih, A. Y. 573
Alic, M. 53, 463
Aling, H. 805
Alpargu, G. 57, 66, 501
Alperin, R. C. 194
Alsina, C. 32, 34
Altmann, S. L. xvii, 193, 194, 226, 228
Alzer, H. 52
Amghibech, S. 487
Anderson, B. D. O. xvii, 343, 465, 502, 704, 805
Anderson, G. 23, 27, 29
Anderson, T. W. 459
Anderson, W. N. 335, 384, 385, 443, 460, 463, 475, 529, 530, 541
Ando, T. 439, 442, 457, 460, 461, 463–465, 472, 477, 478, 490, 514, 515, 530, 535–537, 540, 542, 583, 584, 588, 590, 600, 612, 615, 683, 685, 686, 689, 693–695, 795
Andreescu, T. 70, 71, 155, 156, 158–160
Andrica, D. 70, 71, 155, 156, 158–160
Andruchow, E. 576, 589
Angel, E. xvii, 194
Anglesio, J. 29
Anonymous 42
Antoulas, A. C. 793
Aplevich, J. D. xvii, 304
Apostol, T. 447
Apostol, T. M. 673
Araki, H. 479, 583, 584
Araujo, J. 350
Arimoto, A. 357
Arnold, B. 164
Arponen, T. 672
Arsigny, V. 686
Artin, M. 223, 229
Artzrouni, M. 715
Arvanitoyeorgos, A. 229
Aslaksen, H. 152, 183, 226, 229, 260, 261
Asner, B. A. 697, 702
Au-Yeung, Y.-H. 362, 458, 504, 507
Audenaert, K. M. R. 479, 480, 596
Aujla, J. S. 458, 514, 521, 533, 537, 539, 541, 583, 592
Aupetit, B. 695
Avriel, M. 505
Axelsson, O. xviii
Ayache, N. 686
Azar, L. E. 63
Baez, J. C. 222, 227
Bagdasar, O. 22, 37, 42, 54
Bai, Z. 476
Bailey, D. 17, 18, 20, 21, 56, 161, 275
Bailey, D. W. 270
Bailey, H. 158
Baker, A. 184, 217, 226, 229, 347, 660, 682
Baksalary, J. K. 120, 203, 310, 378, 387, 391, 457, 467, 523, 526, 527, 536
Baksalary, O. M. 120, 178, 180, 181, 203, 315, 373, 378, 387, 391, 396–398, 525
Ball, K. 570, 586
Ballantine, C. S. 178, 314, 350, 351, 542, 701, 795, 796
Banerjee, S. 161
Bang-Jensen, J. xvii
Bani-Domi, W. 594, 595
Bapat, R. B. 66, 124, 218, 275, 280, 391, 533
Bar-Itzhack, I. Y. 191, 674
Barbeau, E. J. 31, 39, 42, 47, 158, 708, 710
Baric, J. 30
Barnes, E. R. 271
Barnett, S. xvii, 135, 146, 215, 257, 281, 353, 354, 361, 362, 377, 393, 450, 454, 618, 702, 798
Barrett, W. 124, 220, 444, 449, 492, 538
Barria, J. 319
Bart, H. 800
Baruh, H. 193, 226, 676
Barvinok, A. 43, 47, 110, 111, 115, 498, 511, 634
Barza, S. 58
Bates, R. G. 221
Bau, D. xviii
Bauer, F. L. 619

- Bayard, D. S. 334
 Bazaraa, M. S. 279, 624, 626
 Beams, R. 348
 Beavers, A. N. 349
 Bebiano, N. xvii, 335, 486, 686–688
 Beckenbach, E. F. 76, 544
 Becker, R. I. 362, 504
 Beckner, W. 586
 Bekjan, T. N. 500
 Bekker, P. A. 483, 484, 541
 Belinfante, J. G. 722
 Belitskii, G. R. 619
 Bellman, R. 76, 146, 280, 454, 460, 544, 682
 Ben-Israel, A. 372, 376, 391, 394, 395, 398, 507
 Ben Taher, B. 673
 Ben-Tal, A. 110, 164
 Benjamin, A. T. 11, 17, 278
 Benson, C. T. 223
 Berg, L. 129
 Berge, C. xvii
 Berkovitz, L. D. 164, 635
 Berman, A. 176, 230, 275, 277, 706, 707
 Bernhardsson, B. 701
 Bernstein, D. S. 127, 189, 219, 227, 230, 338, 343, 491, 498, 619, 673, 679, 681, 689, 692, 701, 705–708, 722, 796–798, 805
 Bhagwat, K. V. 433, 459, 683
 Bhat, S. P. 227, 338, 705, 707, 722, 797
 Bhatia, R. 24, 163, 221, 272, 320, 326, 327, 335, 433, 436, 439, 441–443, 445–449, 457, 458, 463, 479, 480, 486, 509–511, 513, 515, 518, 520, 521, 531, 541, 542, 561, 570, 575, 578, 580, 582, 583, 585–590, 594, 600, 601, 605, 613, 686, 688, 691, 693, 695
 Bhattacharya, R. 318
 Bhattacharyya, S. P. 696, 697, 796, 805
 Bhattacharyya, S. P. 797
 Bhaya, A. 722
 Bicknell, M. R. 142
 Biggs, N. xvii, 337
 Binding, P. 362
 Binmore, K. xvii
 Bjorck, A. 618
 Blanes, S. 678, 683
 Bloch, E. D. 10, 76
 Blondel, V. 715
 Blumenthal, L. M. 161
 Boche, H. xvii, 162
 Boehm, W. 714
 Bojanczyk, A. W. 265
 Bollobas, B. xvii
 Bondar, J. V. 45, 60
 Borck, A. 398
 Borre, K. xvii
 Borwein, J. 17, 18, 20, 21, 56, 161, 275
 Borwein, J. M. 110, 111, 164, 279, 333, 334, 441, 460, 633, 635
 Bosch, A. J. 349, 361
 Bottcher, A. 584
 Bottema, O. 158
 Boullion, T. L. 398
 Bourin, J.-C. 61, 458, 465, 480–483, 521, 537, 576, 583, 592, 599, 608, 687
 Bourque, K. 447
 Boyd, S. xvii, 164, 578
 Bozkurt, D. 615
 Brenner, J. L. 154, 229, 270, 346
 Bresler, Y. 535, 538
 Brewer, J. W. 416
 Brickman, L. 498
 Brockett, R. 805
 Brockett, R. W. 511, 797
 Brothers, H. J. 25, 26
 Brown, G. 66
 Browne, E. T. 541
 Bru, R. 343
 Brualdi, R. A. xvii, 126, 129, 131, 269, 506, 704
 Buckholtz, D. 210, 336, 337
 Bullen, P. S. 23, 25, 27, 29, 35, 38, 42, 52, 53, 55, 57, 64, 76, 546, 567, 571
 Bultheel, A. 281
 Burch, J. M. xvii
 Burns, F. 391
 Bushell, P. J. 519
 Cahill, N. D. 278
 Cain, B. E. 309, 795
 Callan, D. 118
 Campbell, S. L. 377, 384, 391, 395, 398, 699, 722
 Cao, J. 27, 38
 Cao, L. 418, 457
 Carlen, E. 570, 586
 Carlen, E. A. 441
 Carlson, D. 387, 391, 442, 542, 702, 794, 795
 Carpenter, J. A. 329
 Cartier, P. 722
 Cartwright, D. I. 52, 722
 Casas, F. 678, 683
 Castro-Gonzalez, N. 391
 Caswell, H. xvii
 Cater, F. S. 348
 Chabrilac, Y. 504
 Chan, N. N. 458
 Chandrasekar, J. 219
 Chapellat, H. 696, 697
 Chartrand, G. xvii
 Chatelin, F. xviii
 Chattot, J.-J. xvii
 Chaturvedi, N. A. 189
 Chebotarev, P. 708
 Chehab, J.-P. 476
 Chellaboina, V. xvii, 230, 707, 708
 Chellaboina, V.-S. 579, 619, 767
 Chen, B. M. xvii, 114, 325, 333, 362
 Chen, C. T. 793
 Chen, H. 52, 61
 Chen, J. 598
 Chen, J.-Q. 224
 Chen, L. 478, 613
 Chen, S. 535
 Cheng, C.-M. 443, 486, 515, 615
 Cheng, H.-W. 673
 Cheng, S. 178, 180, 372, 373, 379, 380, 383, 386, 388, 394, 395, 531
 Chien, M.-T. 219
 Choi, M.-D. 208, 215, 360
 Chollet, J. 474
 Choudhry, A. 348
 Chu, M. T. 378, 388, 457
 Chu, X.-G. 17
 Chuai, J. 407, 410
 Chuang, I. L. xvii
 Chui, N. L. C. 714

- Chung, F. R. K. xvii
 Cizmesija, A. 58
 Clements, D. J. 805
 Climent, J. J. 343
 Cline, R. E. 119, 121, 383, 390
 Cloud, M. J. 76
 Coakley, E. S. 225
 Cohen, J. E. 690, 692
 Cohoon, D. K. 127
 Collins, E. G. 416, 805
 Constales, D. 371
 Contreras, M. xvii, 708
 Conway, J. C. 47, 48, 223, 226, 227
 Corach, G. 576, 589
 Corless, M. J. xvii
 Costa, P. J. 641
 Cottle, R. W. 497, 507, 696
 Cover, T. M. xvii, 442, 487, 492, 506
 Crabtree, D. E. 270, 342
 Crawford, C. R. 504
 Crilly, T. 261
 Crossley, M. D. 192
 Crouzeix, J.-P. 504
 Cullen, C. G. 343, 629, 640
 Culver, W. J. 722
 Curtain, R. F. 691
 Curtis, M. L. 193, 226, 229
 Cvetkovic, D. xvii

 da Providencia, J. xvii, 486, 686–688
 Da Silva, J. A. D. 415
 Daboul, P. J. 227
 Dahlquist, G. 706
 Dale, P. xvii, 164
 D'Andrea, R. 610
 D'Angelo, H. 805
 D'Angelo, J. P. 70, 565
 Daniel, J. W. xix, 164
 Dannan, F. M. 480, 488
 Dasgupta, S. 680
 Datko, R. 704
 Datta, B. N. xvii, 794
 Dattorro, J. xvii, 118, 126, 323, 642
 Daubechies, I. 715
 Davies, E. B. 584
 Davis, C. 582, 589
 Davis, P. J. xix, 357
 Davison, E. J. 701
 Dawlings, R. J. H. 350
 Day, J. 513, 655
 Day, P. W. 61
 de Boor, C. 164
 de Groen, P. P. N. 579
 de Hoog, F. R. 378
 de Launey, W. 416
 De Moor, B. 805
 de Pillis, J. 484, 494
 De Pillis, J. E. 494
 de Pillis, L. G. 255
 de Souza, E. 796, 797
 De Souza, P. N. 113, 118, 150, 151, 457
 de Vries, H. L. 597
 DeAlba, L. M. 428, 795
 Debnath, L. 25, 26, 38, 66
 Decell, H. P. 374
 Deistler, M. 805
 Del Buono, N. 217
 Delbourgo, R. 227
 DeMarco, C. L. 269
 Demmel, J. W. xviii
 Deng, C. 460
 Denman, E. D. 349
 DePrima, C. R. 346
 D'Errico, J. R. 278
 Desoer, C. A. 691
 Deutsch, E. 593, 691, 709, 710
 Devaney, R. L. xvii
 Dhrymes, P. J. xvii
 Dieci, L. 660, 685
 Diestel, R. xvii
 Dines, L. L. 498
 Ding, J. 263, 592
 Dittmer, A. 188
 Dixon, G. M. 227
 Dixon, J. D. 491
 Djafaris, T. E. 797
 Djokovic, D. Z. 150, 153, 316, 348, 350
 Djordjovic, Z. 158
 Dokovic, D. Z. 312, 314, 315, 318, 343
 Dolotin, V. 416
 Dombre, E. xvii
 Donoghue, W. F. 446, 540, 541
 Dopazo, E. 391
 Dopico, F. M. 217, 225, 346
 Doran, C. 188, 227, 565
 Doran, C. J. L. 188, 227, 565
 Dorst, L. 227, 565
 Douglas, R. G. 431
 Doyle, J. C. 701
 Doyle, P. G. xvii
 Drachman, B. C. 76
 Dragan, V. 805
 Dragomir, S. S. 36, 57, 59, 61, 63–66, 70
 Drazin, M. P. 259
 Drissi, D. 38, 448
 Drivaliaris, D. 372
 Drnovsek, R. 200
 Drury, S. W. 185
 Du, H. 460
 Du, H.-K. 469, 521, 610
 Duffin, R. J. 529
 Duleba, I. 678
 Dullerud, G. E. xvii
 Dummit, D. S. 222, 223, 357, 361
 Dunkl, C. F. 565
 Dym, H. 56, 129, 131, 134, 160, 280, 309–311, 335, 336, 391, 431, 458, 507, 510, 525, 590, 591, 610, 634, 636, 638, 698, 794
 Edelman, A. 447
 Egecioglu, O. 70
 Eggleston, H. G. 164
 Elsner, L. 17, 142, 180, 269, 413, 415, 534, 650, 691, 695
 Embree, M. 691
 Engel, A. 31, 32, 34, 39–46, 49, 52, 58, 60, 158
 Engo, K. 683
 Erdmann, K. 229
 Erdos, J. A. 350
 Eriksson, R. 161
 Evard, J.-C. 722

 Fallat, S. 453
 Fan, K. 453, 485, 487
 Fang, M. 415
 Fang, Y. 334
 Farebrother, R. W. 229, 348, 398
 Farenick, D. R. 357
 Fassler, A. 229
 Feiner, S. xvii, 194
 Fekete, A. E. 188, 200
 Feng, B. Q. 574, 583
 Feng, X. 334
 Feng, X. X. 476, 478, 480

- Feng, Z. 159
 Fenn, R. 30, 161, 188, 193,
 226, 227, 278
 Ferreira, P. G. 805
 Ferziger, J. H. xvii
 Fiedler, M. xvii, 149, 160,
 255, 257, 275, 329, 355,
 376, 390, 412, 463, 468,
 532, 534, 535, 702
 Field, M. J. 52, 722
 Fill, J. A. 378
 Fillard, P. 686
 Fillmore, J. P. 673
 Fillmore, P. A. 313, 360,
 431
 Fink, A. M. 23, 70, 76, 271,
 565, 568, 569
 Fishkind, D. E. 378
 Fitzgerald, C. H. 531
 Flanders, H. 214, 459, 603
 Fleming, W. 494
 Flett, T. M. 642
 Foldes, S. 222
 Foley, J. xvii, 194
 Fontijne, D. 227, 565
 Foote, R. M. 222, 223, 357,
 361
 Formanek, E. 149, 214, 260
 Foulds, L. R. xvii
 Francis, B. A. xvii
 Franklin, J. xix
 Frazho, A. E. xvii
 Frazier, M. xvii
 Freiling, G. xvii
 Friedland, S. 601, 614, 618,
 692, 708
 Friswell, M. I. 412, 413
 Fuhrmann, P. A. 197, 255,
 257, 258, 281, 309, 473
 Fujii, J. I. 502, 589
 Fujii, M. 34, 53, 67, 461,
 466, 502, 522, 523, 565,
 589, 685, 686, 711
 Fuller, A. T. 406, 413, 416,
 697
 Fulton, W. 223
 Funderlic, R. E. 119, 121,
 378, 383, 388, 457
 Furuichi, S. 479
 Furuta, T. 24, 209, 433,
 434, 436, 441, 455, 461,
 464–467, 481, 502, 511,
 520, 522, 523, 568, 582,
 589, 683, 686
 Gaines, F. 184, 185, 313
 Galantai, A. 112, 204, 209,
 210, 230, 315, 316, 329,
 335–337, 380, 571, 572,
 636
 Gallier, J. 193
 Gangsong, L. 511
 Gantmacher, F. R. xix,
 304, 318, 541, 722
 Garling, D. J. H. 33, 53,
 54, 58, 63, 75, 162, 163,
 569, 574
 Garloff, J. 697
 Garvey, S. D. 412, 413
 Geerts, T. 805
 Gelfand, I. M. 416
 Genton, M. G. 445
 George, A. 342, 358
 Ger, R. 34
 Gerdes, P. 357
 Gerrard, A. xvii
 Gerrish, F. 320
 Geveci, T. 805
 Gheondea, A. 460
 Ghouraba, F. A. A. 230
 Gil, M. I. 325
 Gilmore, R. 229
 Girard, P. R. 188, 193, 225
 Girgensohn, R. 17, 18, 20,
 21, 56, 161, 275
 Glasser, M. L. 683
 Godsil, C. xvii
 Godunov, S. K. xix, 211,
 699
 Goh, C. J. 59
 Gohberg, I. 234, 281, 336,
 361, 636, 787, 788, 800,
 805
 Golberg, M. A. 640, 678
 Goldberg, M. 583, 603
 Goller, H. 530
 Golub, G. H. xviii, 378,
 388, 457, 614, 646
 Golub, G. H. 476
 Gong, M.-P. 519
 Gonzalez, N. C. 680
 Goodman, F. M. 223
 Goodman, L. E. 188
 Gordon, N. 139
 Goroncy, A. 51
 Govaerts, W. 413, 416, 610
 Gow, R. 312, 343, 351
 Graham, A. 416
 Graybill, F. A. xvii
 Grcar, J. 164
 Grcar, J. F. 559, 560
 Green, W. L. 463, 529
 Greene, D. H. 17
 Greub, W. H. xix, 416
 Greville, T. N. E. 210, 372,
 376, 380, 381, 391, 394,
 395, 398, 507
 Grigoriadis, K. xvii, 722
 Grone, R. 180, 190, 341
 Gross, J. xvii, 188,
 202–204, 209, 229, 335,
 381, 391, 523, 524, 526,
 530
 Grove, L. C. 223
 Guan, K. 38
 Gudder, S. 460
 Gull, S. 188, 227, 565
 Guobiao, Z. 511
 Gupta, A. K. xvii
 Gurlebeck, K. 227, 228, 675
 Gurvits, L. 715
 Gustafson, K. E. 497, 577
 Gustafson, W. H. 351
 Gutin, G. xvii
 Gwanyama, P. W. 52
 Haddad, W. M. xvii, 164,
 230, 343, 491, 579, 619,
 707, 708, 767, 798, 805
 Hager, W. W. 164, 316
 Hahn, W. 722
 Hairer, E. 678
 Hajja, M. 158
 Halanay, A. 805
 Hall, A. 465
 Hall, B. C. 217, 654, 655,
 657–660, 680, 684, 722
 Halliwell, G. T. 27, 67
 Halmos, P. R. 90, 113, 202,
 279, 313, 318, 319, 340,
 343, 349–351, 376, 384,
 385, 451, 681
 Hamermesh, M. 224
 Han, J. H. 128
 Haneda, H. 691
 Hannan, E. J. 805
 Hansen, F. 483
 Hanson, A. J. 227, 228
 Hardy, G. 76
 Harner, E. J. 335
 Harris, J. 223
 Harris, L. A. 148, 473, 474
 Harris, W. A. 673

- Hart, G. W. xvii, 230
 Hartfiel, D. J. xvii, 715
 Hartwig, R. 391
 Hartwig, R. E. 119, 120, 315, 322, 337, 361, 372, 373, 382, 385, 391, 394, 396, 524, 526, 527, 797
 Harville, D. A. xvii, 199, 202, 371, 377, 378, 381, 383, 486, 506, 642
 Hattori, S. 704
 Hauke, J. 524
 Hautus, M. L. J. xvii, 805
 Havel, T. F. 682
 Haynes, T. 713
 Haynsworth, E. V. 387, 391, 442, 474, 542
 Hecht, E. xvii
 Heij, C. 805
 Heinig, G. 257
 Helmke, U. 257
 Helton, B. W. 722
 Henderson, H. V. 145, 164, 416
 Herman, J. 17, 18, 23–26, 32, 36, 37, 39–44, 46, 47, 49, 52, 60
 Hershkowitz, D. 17, 191, 413, 475, 674, 708
 Hestenes, D. 188, 227
 Hiai, F. 442, 460, 461, 465, 477, 478, 580, 683, 686, 693, 694
 Higham, D. J. 164
 Higham, N. J. xviii, 72, 74, 164, 215, 221, 261, 327, 328, 331, 348, 349, 359, 360, 571, 573, 575–577, 584, 603, 608, 609, 615, 619, 629, 637, 657, 658, 677, 681, 685, 692–694, 700, 701, 722, 802
 Hile, G. N. 227
 Hill, R. 795
 Hill, R. D. 142, 221
 Hillar, C.-J. 482
 Hilliard, L. O. 280
 Hinrichsen, D. 318, 326, 327, 554, 555, 617, 638, 639, 691, 695, 699, 709
 Hirsch, M. W. xvii, 311
 Hirschhorn, M. D. 128
 Hirzallah, O. 38, 70, 502, 586, 612, 613
 Hmamed, A. 541
 Hoagg, J. B. 219
 Hoffman, A. J. 271
 Hoffman, K. xix
 Holbrook, J. 24, 515, 688
 Hollot, C. V. 504
 Holmes, R. R. 224
 Holtz, O. 697, 715
 Hong, Y. 350
 Hong, Y. P. 606
 Horn, A. 511
 Horn, R. A. 139, 163, 197, 214, 254, 271, 273, 275, 276, 280, 281, 293, 314, 319, 320, 326, 327, 333, 341, 342, 345, 346, 348, 350, 358, 360, 393, 405, 407, 412, 416, 428, 431, 443, 446, 461, 470, 474, 486, 490, 492, 493, 497, 505, 507, 509–512, 515, 531, 532, 537–539, 541, 546, 549, 550, 553, 561, 562, 572–574, 576, 578, 580, 581, 591, 592, 601, 603, 605, 606, 609, 612, 615, 617, 641, 642, 654, 657, 678, 689, 692, 693, 703, 707, 797
 Horne, B. G. 604
 Hou, H.-C. 469, 521, 610
 Hou, S.-H. 281
 Householder, A. S. xviii, 257, 329, 378, 611, 634
 Howe, R. 229, 722
 Howie, J. M. 76
 Howland, R. A. xvii
 Hsieh, P.-F. xvii, 311
 Hu, G.-D. 700
 Hu, G.-H. 700
 Hu-yun, S. 127, 519
 Huang, R. 415, 416, 603, 616, 617
 Huang, T.-Z. 51, 275, 416, 598, 617
 Hughes, J. xvii, 194
 Hughes, P. C. xvii, 676
 Huhtanen, M. 70
 Humphries, S. 132
 Hung, C. H. 391
 Hunter, J. J. 398
 Hyland, D. C. 230, 343, 416, 706, 708, 805
 Ibragimov, N. H. 722
 Ikebe, Y. 642
 Ikramov, K. D. 180, 342, 358, 597, 598, 650, 695
 Inagaki, T. 642
 Ionescu, V. xvii, 805
 Ipsen, I. 112, 210, 329, 336, 381
 Iserles, A. 188, 674, 678, 682
 Ito, Y. 704
 Iwasaki, T. xvii, 722
 Izumino, S. 53, 65, 501
 Jacobson, D. H. 805
 Jagers, A. A. 29
 Jameson, A. 541
 Janic, R. R. 158
 Jank, G. xvii
 Jeffrey, A. 74–76
 Jeffries, C. 708
 Jennings, A. xviii
 Jennings, G. A. 161
 Jensen, S. T. 51, 454
 Ji, J. 129
 Jia, G. 27, 38
 Jiang, Y. 316
 Jin, X.-Q. 584
 Jodic, D. 588
 Johnson, C. R. 132, 139, 163, 180, 185, 190, 191, 197, 214, 217, 219, 225, 254, 271, 273, 275, 276, 280, 281, 293, 308–310, 314, 320, 326, 327, 341, 342, 345, 346, 348, 358, 360, 405, 407, 412, 415, 416, 428, 431, 443, 446, 449, 470, 474, 485, 487, 492, 497, 502, 505, 507, 509–512, 532, 534, 537, 538, 541, 546, 549, 550, 553, 561, 562, 572–574, 578, 580, 592, 601, 603–606, 609, 612, 615, 619, 641, 642, 654, 657, 674, 678, 689, 692, 693, 697, 701, 703, 707, 795, 797
 Jolly, M. 642
 Jonas, P. 460
 Jordan, T. F. 228
 Jorswieck, E. A. xvii, 162
 Joyner, D. 230

- Jung, D. 34, 53
 Junkins, J. L. 191, 674
 Jury, E. I. 413, 416, 704, 709
 Kaashoek, M. A. 800
 Kaczor, W. J. 24, 27, 29, 34, 66
 Kadison, R. V. 459
 Kagan, A. 483
 Kagstrom, J. B. 699
 Kailath, T. 237, 281, 304, 353, 805
 Kalaba, R. E. xvii
 Kalman, D. 357
 Kamei, E. 461, 466, 522, 523
 Kane, T. R. xvii
 Kanzo, T. 51, 566
 Kapila, V. 805
 Kaplansky, I. xix, 318
 Kapranov, M. M. 416
 Karanasios, S. 372
 Karcianas, N. 362, 805
 Karlin, S. 415
 Kaszkurewicz, E. 722
 Kato, M. 568
 Kato, T. 460, 600, 619, 692
 Katsuura, H. 46, 53, 54
 Katz, I. J. 382
 Katz, S. M. 494
 Kauderer, M. xvii
 Kazakia, J. Y. 313
 Kazarinoff, N. D. 36, 546
 Keel, L. 697
 Kelly, F. P. 692
 Kendall, M. G. 184
 Kenney, C. 327, 637, 711
 Kestelman, H. 340
 Keyfitz, N. xvii, 275
 Khalil, W. xvii
 Khan, N. A. 405, 416, 536
 Khatri, C. G. 374, 384
 Kim, S. 53, 463, 683
 King, C. 586, 595
 Kittaneh, F. 38, 70, 313, 327, 331, 458, 502, 515–517, 578, 580–590, 592, 594, 595, 603, 610–613, 711, 712
 Klaus, A.-L. 583
 Klee, V. 708
 Knox, J. A. 25, 26
 Knuth, D. E. 17
 Koch, C. T. 678
 Koks, D. 227
 Koliha, J. J. 202, 203, 205, 382, 680
 Kolman, B. 722
 Komaroff, N. 61, 333, 518
 Komornik, V. 696
 Koning, R. H. 416
 Kosaki, H. 584, 585
 Kosecka, J. xvii
 Koshy, T. xvii, 218, 278
 Kovac-Striko, J. 362
 Kovacec, A. 335
 Krafft, O. 529
 Krattenthaler, C. xvii, 132
 Kratz, W. 805
 Krauter, A. R. 476
 Kreindler, E. 541
 Kress, R. 597
 Krupnik, M. 357
 Krupnik, N. 357
 Kubo, F. 67, 565, 711
 Kucera, R. 17, 18, 23–26, 32, 36, 37, 39–44, 46, 47, 49, 52, 60
 Kucera, V. 805
 Kufner, A. 58, 59, 63
 Kuipers, J. B. xvii, 226, 228
 Kunze, R. xix
 Kurepa, S. 689
 Kwakernaak, K. xvii
 Kwapisz, M. 712
 Kwong, M. K. 221, 458, 460, 533
 Kwong, R. H. 790, 805
 Kyrchei, I. I. 129
 Laberteaux, K. R. 181
 Laffey, T. J. 281, 319, 343, 351
 Lagarias, J. C. 715
 Lai, H.-J. xvii
 Lakshminarayanan, S. 129
 Lam, T. Y. 42
 Lancaster, P. xvii, 234, 257, 281, 304, 320, 336, 340, 361, 406, 504, 560, 579, 619, 636, 787, 788, 793–795, 805
 Langholz, G. 704
 Larson, L. 24, 29, 33, 35, 38, 40–44, 49, 155, 158
 Larsson, L. 58
 Lasenby, A. 188, 227, 565
 Lasenby, A. N. 188, 227, 565
 Lasenby, J. 188, 227, 565
 Lasserre, J. B. 333
 Laub, A. J. xix, 113, 304, 306, 327, 338, 637, 711, 802, 805
 Laurie, C. 361
 Lavoie, J. L. 392
 Lawson, C. L. 398
 Lawson, J. D. 431, 463
 Lax, P. D. 160, 259, 457, 600
 Lay, S. R. 93, 164, 635
 Lazarus, S. 319
 Leake, R. J. 186
 Leclerc, B. 129
 LeCouteur, K. J. 519, 682
 Lee, A. 230, 313, 376
 Lee, J. M. 361
 Lee, S. H. 34, 53
 Lee, W. Y. 357
 Lehnigk, S. H. 722
 Lei, T.-G. 453, 483, 493, 501, 505, 508, 510
 Leite, F. S. 348, 675
 Lemos, R. xvii, 686–688
 Leonard, E. 646
 Lesniak, L. xvii
 Letac, G. 531
 Levinson, D. A. xvii
 Lew, J. S. 260, 261
 Lewis, A. S. 110, 111, 164, 279, 333, 334, 441, 460, 633, 635
 Lewis, D. C. 414
 Li, C.-K. xvii, 272, 328, 360, 436, 442, 443, 458–460, 463, 486, 515, 533, 542, 570, 583, 600, 611, 612
 Li, C.-L. 58
 Li, J. 38, 316
 Li, J.-L. 29
 Li, Q. 460, 581
 Li, R.-C. 272
 Li, X. 391
 Li, Y.-L. 29
 Li, Z. xvii, 722
 Lieb, E. 570, 586
 Lieb, E. H. 441, 474, 482, 500, 519, 542, 570, 688
 Ligh, S. 447

- Likins, P. W. xvii
 Lim, J. S. 346
 Lim, Y. 53, 431, 463, 683, 803
 Liman, A. 709
 Lin, C.-S. 466, 503, 522, 565
 Lin, T.-P. 36, 37
 Lin, W.-W. 362
 Lin, Z. xvii, 114, 325, 333, 362
 Linden, H. 710, 711
 Lipsky, L. xvii
 Littlewood, J. E. 76
 Liu, B. xvii
 Liu, H. 26
 Liu, J. 416, 442, 542
 Liu, R.-W. 186
 Liu, S. 57, 416, 501, 533, 536, 539
 Liu, X. 120
 Liu, Y. 387
 Liz, E. 646
 Loewy, R. 793, 794
 Logofet, D. O. xvii
 Loksha, V. 158
 Loparo, K. A. 334
 Lopez, L. 217
 Lopez-Valcarce, R. 680
 Loss, M. 570
 Lossers, O. P. 153
 Lounesto, P. 188, 227
 Lubich, C. 678
 Luenberger, D. G. xvii, 505
 Lundquist, M. 124, 444, 492
 Lutkepohl, H. xix
 Lutoborski, A. 265
 Lyubich, Y. I. 619
 Ma, E.-C. 797
 Ma, Y. xvii
 MacDuffee, C. C. 406, 410, 413, 416
 Macfarlane, A. G. J. 805
 Maciejowski, J. M. 714
 Mackey, D. S. 230
 Mackey, N. 230
 Maddocks, J. H. 311, 504
 Maeda, H. 704
 Magnus, J. R. xvii, 370, 384, 388, 390, 391, 398, 416, 476, 477, 481, 503, 642
 Magnus, W. 678
 Majindar, K. N. 508
 Maligranda, L. 58, 59, 63, 567–569
 Malyshev, A. N. 699
 Malzan, J. 348
 Mangasarian, O. xvii
 Manjegani, S. M. 477
 Mann, H. B. 505
 Mann, S. 227, 565
 Mansfield, L. E. xix
 Mansour, M. 696, 697, 704
 Maradudin, A. A. 678
 Marcus, M. xix, 55, 76, 136, 229, 325, 333, 405, 412, 416, 450, 484, 494, 513, 536, 544, 598
 Margaliot, M. 704
 Markham, T. L. 149, 376, 387, 391, 442, 468, 534, 535, 541, 542
 Markiewicz, A. 524
 Marsaglia, G. 121, 164, 387, 389
 Marsden, J. E. xvii, 193
 Marshall, A. W. 44, 45, 60, 61, 76, 158, 162–164, 326, 334, 412, 442, 443, 455, 510, 541, 542, 618, 619
 Martensson, K. 805
 Martin, D. H. 805
 Massey, J. Q. 131
 Mastronardi, N. 125, 129
 Mathai, A. M. 642
 Mathes, B. 361
 Mathias, R. 328, 332, 416, 436, 440, 442, 448, 463, 512, 533, 539, 542, 576, 581, 591, 600, 603, 611, 612, 615, 617, 642, 682
 Matic, M. 30, 59, 67
 Matson, J. B. 805
 Matsuda, T. 58
 Maybee, J. S. 338
 Mazorchuk, V. 581
 McCarthy, J. E. 450
 McCarthy, J. M. 722
 McClamroch, N. H. 189
 McCloskey, J. P. 213
 McKeown, J. J. xviii
 Meehan, E. 281
 Meenakshi, A. R. 530
 Mehta, C. L. 500
 Mellendorf, S. 269, 704
 Melnikov, Y. A. xvii
 Mercer, P. R. 27, 67, 567, 568
 Merikoski, J. K. 604
 Merris, R. 277, 416, 461, 469, 506, 536
 Meyer, C. 112, 210, 329, 336, 381
 Meyer, C. D. 160, 175–177, 212, 243, 279, 343, 357, 375, 377, 384, 391, 395, 398, 714, 722
 Meyer, K. 802
 Miao, J.-M. 391
 Mickiewicz, A. 525
 Mihalyffy, L. 391
 Miller, K. S. 254, 410, 506
 Milliken, G. A. 526, 527
 Milovanovic, G. V. 638, 709–712
 Minamide, N. 378
 Minc, H. xix, 55, 76, 229, 325, 544, 598
 Miranda, H. 541
 Miranda, M. E. 335
 Mirsky, L. xix, 190, 198
 Misra, P. 800
 Mitchell, J. D. 350
 Mitra, S. K. 384, 398, 416, 529, 541
 Mitrinovic, D. S. 23, 38, 52, 70, 76, 158, 271, 565, 567–569, 709, 712
 Mitter, S. K. 797
 Mityagin, B. 458, 541
 Miura, T. 51, 566
 Mlynarski, M. 691
 Moakher, M. 196, 360, 463, 640, 674, 688, 689, 692
 Moler, C. 692, 722
 Molera, J. M. 346
 Mond, B. 457, 463, 502, 537–539
 Monov, V. V. 269
 Moon, Y. S. 504
 Moore, J. B. 465
 Moreland, T. 460
 Mori, H. 65, 501
 Mori, T. 541
 Morley, T. D. 463, 529
 Morozov, A. 416
 Muckenhoupt, B. 313
 Muir, T. 164
 Muir, W. W. 441, 442, 542

- Mukherjea, K. 318
 Munthe-Kaas, H. Z. 188, 674, 677, 678, 682
 Murphy, I. S. 492
 Murray, R. M. xvii, 722

 Nagar, D. K. xvii
 Najfeld, I. 682
 Najman, B. 362
 Nakamoto, R. 461, 466, 522, 523, 589, 695
 Nakamura, Y. 361
 Nandakumar, K. 129
 Narayan, D. A. 278
 Narayan, J. Y. 278
 Nataraj, S. xvii
 Nathanson, M. 586
 Naylor, A. W. 70, 76, 565, 568, 623, 624, 642
 Needham, T. 76
 Nelsen, R. B. 32
 Nemirovski, A. 110, 164
 Nersesov, S. G. 230, 708
 Nett, C. N. 164
 Neubauer, M. G. 491
 Neudecker, H. xvii, 57, 370, 384, 388, 391, 398, 416, 476, 477, 501, 503, 642
 Neumann, M. 219, 275, 343, 706, 707
 Neuts, M. F. xvii
 Newcomb, R. W. 541
 Newman, M. 132, 309, 655
 Nguyen, T. 125, 218
 Nicholson, D. W. 520
 Niculescu, C. 21, 22, 35, 37, 42, 54–56, 62, 132, 483, 565, 638, 639
 Niculescu, C. P. 22, 39, 54, 158, 712
 Nielsen, M. A. xvii
 Niezgoda, M. 51, 454
 Nishio, K. 200
 Noble, B. xix, 164
 Nomakuchi, K. 391
 Nordstrom, K. 310, 457, 527
 Norman, E. 678
 Norsett, S. P. 188, 674, 678, 682
 Nowak, M. T. 24, 27, 29, 34, 66
 Nunemacher, J. 722
 Nylen, P. 619

 Oar, C. xvii, 805
 Odell, P. L. 398
 Ogawa, H. 378
 Okubo, K. 348, 477, 478
 Olesky, D. D. 338
 Olkin, I. 44, 45, 60, 61, 76, 158, 162–164, 319, 326, 334, 412, 442, 443, 455, 456, 459, 510, 541, 542, 618, 619
 Ortega, J. M. xix
 Ortner, B. 476
 Osburn, S. L. 681
 Ost, F. 415
 Ostrowski, A. 794
 Ostrowski, A. M. 593
 Oteo, J. A. 678
 Ouellette, D. V. 468, 487, 542
 Overdijk, D. A. 188

 Paardekooper, M. H. C. 180
 Pachter, M. 805
 Paganini, F. xvii
 Paige, C. C. 310, 316, 472, 473, 490, 503
 Palanthandalam-Madapusi, H. 230, 498, 679
 Paliogiannis, F. C. 681
 Palka, B. P. 76
 Pan, C.-T. 606
 Pao, C. V. 691
 Papastavridis, J. G. xvii, 451
 Pappas, D. 372
 Park, F. C. 722
 Park, P. 334
 Parker, D. F. 362
 Parks, P. C. 704
 Parthasarathy, K. R. 446, 589, 590, 695
 Patel, R. V. 484, 692, 800, 805
 Pearce, C. E. 59, 64
 Pearce, C. E. M. 59, 67
 Percy, C. 318
 Pease, M. C. 229
 Pecaric, J. 30, 58, 59, 64, 67, 568
 Pecaric, J. E. 23, 53, 70, 76, 158, 271, 457, 463, 501, 502, 537–539, 565, 568, 569, 616

 Pennec, X. 686
 Peric, M. xvii
 Perlis, S. 164, 233, 234, 237, 281, 361
 Persson, L.-E. 21, 22, 35, 37, 42, 54–56, 58, 59, 62, 63, 132, 483, 565, 638, 639
 Peter, T. 159
 Petersen, I. R. 504
 Peterson, A. C. 805
 Petz, D. 461, 463, 467, 686
 Piepmeyer, G. G. 642
 Pierce, S. 461, 469
 Ping, J. 224
 Pipes, L. A. 672, 673
 Pittenger, A. O. xvii
 Plemmons, R. J. 176, 230, 275, 277, 706, 707
 Plischke, E. 699
 Polik, I. 498
 Politi, T. 217, 673, 675
 Pollock, D. S. G. 642
 Polya, G. 76
 Polyak, B. T. 497–499
 Poon, E. 360
 Poonen, B. 349, 447
 Popa, D. 565
 Popov, V. M. xvii, 722
 Popovici, F. 22
 Porta, H. 589
 Porter, G. J. 194
 Pourciau, B. H. 635
 Pranesachar, C. R. 158
 Prasolov, V. V. xix, 143, 150, 184, 185, 197, 208, 214, 227, 255, 257, 259, 271, 275, 313, 320, 328, 334, 335, 340, 341, 343, 345, 349, 350, 358, 393, 412, 442, 474, 486–488, 493, 507, 508, 533, 534, 574, 588, 599, 601, 608, 640, 641, 660
 Prells, U. 412, 413
 Pritchard, A. J. 318, 326, 327, 554, 555, 617, 638, 639, 691, 695, 709
 Pryce, J. D. 610
 Przemieniecki, J. S. xvii
 Psarrakos, P. J. 348
 Ptak, V. 257, 463
 Pukelsheim, F. 120, 416, 467, 523, 526, 527, 536

- Pullman, N. J. 722
 Puntanen, S. 51, 386, 454, 501, 502, 616
 Putcha, M. S. 322, 337, 361
 Pye, W. C. 263

 Qi, F. 17, 38, 60
 Qian, C. 38
 Qian, R. X. 269
 Qiao, S. 129, 202, 378, 398
 Qiu, L. 413, 701
 Queiro, J. F. 335
 Quick, J. 347
 Quinn, J. J. 11, 17, 278

 Rabanovich, S. 581
 Rabanovich, V. 361
 Rabinowitz, S. 641
 Rachidi, M. 673
 Radjavi, H. 200, 319, 348, 351, 358, 361, 376
 Raghavan, T. E. S. 275
 Rajian, C. 530
 Rajic, R. 568
 Rakocovic, V. 202, 203, 205, 210, 336
 Ran, A. 805
 Ran, A. C. M. 800, 805
 Rantzer, A. 320, 701
 Rao, C. R. xvii, 278, 398, 416, 529, 541
 Rao, D. K. M. 497
 Rao, J. V. 391
 Rao, M. B. xvii, 278, 416
 Rasa, I. 565
 Rassias, T. M. 63, 638, 709–712
 Ratiu, R. S. 193
 Ratiu, T. S. xvii
 Rauhala, U. A. 416
 Raydan, M. 476
 Recht, L. 589
 Regalia, P. A. 416
 Reinsch, M. W. 683
 Reznick, B. 42
 Richardson, T. J. 790, 805
 Richmond, A. N. 684
 Riedel, K. S. 378
 Ringrose, J. R. 619
 Rivlin, R. S. 260, 261
 Robbin, J. W. 76, 115, 164, 211, 229, 281, 283, 319, 346, 414, 640
 Robinson, D. W. 218
 Robinson, P. 476
 Rockafellar, R. T. 164, 624, 632, 635, 642
 Rodman, L. xvii, 191, 234, 281, 304, 308, 310, 336, 351, 361, 459, 460, 504, 636, 674, 787, 788, 805
 Rogers, G. S. 642
 Rohde, C. A. 391
 Rohn, J. 575
 Rojo, O. 51, 604
 Rooin, J. 55
 Ros, J. 678
 Rose, D. J. 357
 Rose, N. J. 343, 391, 395, 699
 Rosenbrock, H. H. 763, 764
 Rosenfeld, M. xvii, 215
 Rosenthal, P. 200, 320, 335, 358
 Rosoiu, A. 158
 Rossmann, W. 229
 Rothblum, U. G. 275, 280, 398
 Rotman, J. J. 223, 357
 Rowlinson, P. xvii
 Royle, G. xvii
 Rozsa, P. 142
 Rubin, M. H. xvii
 Rubinstein, Y. A. 158
 Rugh, W. J. 252, 253, 678, 743, 756, 805
 Rump, S. M. 708
 Ruskai, M. B. 474, 542
 Russell, A. M. 449
 Russell, D. L. 793
 Rychlik, T. 51
 Ryser, H. J. xvii, 126, 131, 506

 Sa, E. M. 180, 190, 341
 Saberi, A. xvii
 Sadkane, M. 699
 Sain, M. K. 805
 Saito, K.-S. 568
 Salamon, D. A. 414
 Salmond, D. 139
 Sandor, J. 22, 25, 26, 37, 55, 66
 Sannuti, P. xvii
 Sarria, H. 604
 Sastry, S. S. xvii, 722
 Satnoianu, R. A. 158
 Sattinger, D. H. 150, 172, 655, 660, 722
 Saunders, M. 316
 Sayed, A. H. xvii
 Schaub, H. 191, 674
 Scherer, C. W. 787, 805
 Scherk, P. 347
 Schmoeger, C. 650, 681, 682, 689
 Schneider, H. 17, 129, 269, 413, 570, 794
 Scholkopf, B. 445
 Scholz, D. 685
 Schott, J. R. xvii, 379, 443, 513, 518
 Schrader, C. B. 805
 Schreiber, M. 384, 385, 460
 Schreiner, R. 346
 Schroder, B. S. W. 76
 Schumacher, J. M. 805
 Schwartz, H. M. 418, 457
 Schwenk, A. J. 415
 Scott, L. L. 223
 Searle, S. R. xvii, 145, 164, 416
 Sebastian, P. 642
 Sebastiani, P. 640
 Seber, G. A. F. 67, 111, 118, 139, 154, 156, 184, 206, 208, 213, 217, 220, 230, 269, 271, 374, 375, 379, 393, 487, 500, 526, 535, 572, 607, 706
 Seberry, J. 416
 Seiringer, R. 482
 Selig, J. M. xvii, 193, 227, 229
 Sell, G. R. 70, 76, 565, 568, 623, 624, 642
 Semrl, P. 570
 Seo, Y. 34, 53, 65, 501, 685–687
 Seoud, M. A. 230
 Serre, D. 172, 217, 260, 390
 Serre, J.-P. 223
 Seshadri, V. 704
 Shafroth, C. 139, 196
 Shah, S. L. 129
 Shah, W. M. 709
 Shamash, Y. xvii, 114, 325, 333, 362
 Shapiro, H. 318, 351
 Shapiro, H. M. 415, 532
 Shaw, R. 188

- Shen, S.-Q. 275, 416, 617
 Sherali, H. D. 279, 624, 626
 Shetty, C. M. 279, 624, 626
 Shilov, G. E. xix
 Shuster, M. D. 193, 226, 674
 Sibuya, Y. xvii, 311
 Sijnave, B. 413, 416
 Siljak, D. D. xvii, 706
 Silva, F. C. 514, 796
 Silva, J.-N. 113, 118, 150, 151, 457
 Simic, S. xvii
 Simoes, R. 796
 Simsa, J. 17, 18, 23–26, 32, 36, 37, 39–44, 46, 47, 49, 52, 60
 Singer, S. F. xvii
 Sivan, R. xvii
 Skelton, R. E. xvii, 722
 Smale, S. xvii, 311
 Smiley, D. M. 565
 Smiley, M. F. 565
 Smith, D. A. 47, 48, 223, 226, 227
 Smith, D. R. 673
 Smith, H. A. 722
 Smith, O. K. 265
 Smith, P. J. 483
 Smith, R. A. 797
 Smith, R. L. 185, 191, 674
 Smoktunowicz, A. 186, 391, 593
 Smola, A. J. 445
 Snell, J. L. xvii
 Snieder, R. 370
 Snijders, J. 722, 746, 747, 805
 So, W. 180, 476, 509, 513, 650, 655, 673, 681, 682, 688–690, 693
 Soatto, S. xvii
 Sobczyk, G. 188, 227
 Sontag, E. D. xvii, 704, 793
 Sorensen, D. C. 793
 Sourour, A. R. 350, 351
 Speed, R. P. 378
 Spence, J. C. H. 678
 Spiegel, E. 361
 Spindelbock, K. 315, 372, 373, 396
 Spivak, M. 227, 624
 Sprossig, W. 227, 228, 675
 Stanley, R. P. 11
 Steeb, W.-H. 416, 689
 Steele, J. M. 76
 Stengel, R. F. xvii
 Stepniak, C. 526
 Stern, R. J. 275, 706, 707
 Stetter, H. J. xviii
 Stewart, G. W. xviii, xxxvi, 164, 304, 306, 316, 336, 358, 504, 541, 570, 578, 579, 600, 614, 615, 618, 619, 635, 636
 Stickel, E. U. 705, 722
 Stiefel, E. 229
 Stillier, L. xvii
 Stoer, J. 164, 579, 619, 635
 Stojanoff, D. 576, 589
 Stolarsky, K. B. 37
 Stone, M. G. 160
 Stoorvogel, A. A. xvii, 805
 Storey, C. xvii, 135, 361, 450, 702
 Strang, G. xvii, xix, 125, 164, 218, 347, 447
 Straskraba, I. 202, 203, 205
 Strelitz, S. 698
 Strichartz, R. S. 678
 Strom, T. 691
 Stuelpnagel, J. 674
 Styán, G. P. H. 51, 57, 66, 118–121, 124, 135, 164, 185, 201–203, 208, 210, 229, 310, 322, 332, 371, 385–389, 391, 392, 454, 457, 472, 473, 490, 501–503, 526, 527, 536, 616
 Subramanian, R. 433, 459, 683
 Sullivan, R. P. 350
 Sumner, J. S. 29
 Sun, J. 304, 306, 316, 336, 358, 504, 541, 570, 578, 579, 600, 614, 615, 619, 635, 636
 Swamy, K. N. 541
 Szechtman, F. 312
 Szekeres, P. 227
 Szep, G. 358
 Szirtes, T. xvii
 Szulc, T. 269, 387, 604
 Takagi, H. 51, 566
 Takahashi, Y. 566
 Takahasi, S.-E. 51, 566
 Takane, Y. 391
 Tam, T. Y. 224
 Tamura, T. 568
 Tao, Y. 515
 Tapp, K. 193, 222, 223, 226, 227, 638
 Tarazaga, P. 449, 604
 Taussky, O. 164, 330, 361
 Temesi, R. 463, 467
 Tempelman, W. 188
 ten Have, G. 348
 Terlaky, T. 498
 Terrell, R. E. 683, 684
 Thirring, W. E. 519
 Thomas, J. A. xvii, 442, 487, 492, 506
 Thompson, R. C. 68, 70, 164, 185, 351, 361, 485, 490, 494, 509, 513, 541, 583, 655, 678, 688, 690
 Tian, Y. 118, 124, 135, 148, 152, 153, 178, 180, 199, 201–203, 208, 210, 212, 227, 229, 312, 321, 371–373, 375, 379–383, 386–389, 391, 393–395, 407, 410, 455, 465, 529, 531
 Tismenetsky, M. 320, 340, 406, 560, 579, 619, 793–795
 Tisseur, F. 230
 Toda, M. 484, 692
 Todd, J. 330
 Toffoli, T. 347
 Tominaga, M. 520, 685, 686
 Tonge, A. 574, 583
 Torokhti, A. 614, 618
 Trapp, G. E. 335, 443, 463, 475, 529, 530, 541
 Trefethen, L. N. xviii, 691, 715
 Trenkler, D. 188, 263, 264, 458, 676
 Trenkler, G. 51, 61, 178, 180, 181, 188, 202–204, 207, 209, 229, 263, 264, 315, 335, 350, 372, 373, 375, 378, 381, 389, 391, 396–398, 416, 454, 458, 477, 525, 676
 Trentelman, H. L. xvii, 805
 Treuenfels, P. 802

- Trigiante, D. 354, 362, 447, 672
 Tromborg, B. 196
 Troschke, S.-O. 188, 229, 524, 526
 Trustrum, G. B. 519
 Tsatsomeros, M. 338
 Tsatsomeros, M. J. 219, 453
 Tsing, N.-K. 362, 504
 Tsiotras, P. 188, 191, 674
 Tsitsiklis, J. N. 715
 Tung, S. H. 52
 Turkington, D. A. 416
 Turkmen, R. 615
 Turnbull, H. W. 164
 Tuynman, G. M. 682
 Tyan, F. 343, 796

 Uchiyama, M. 522, 688, 694
 Udawadia, F. E. xvii
 Uhlig, F. 117, 347, 362, 504, 507, 508, 722
 Underwood, E. E. 142
 Upton, C. J. F. 449

 Valentine, F. A. 164
 Vamanamurthy, M. 23, 27, 29
 Van Barel, M. 125, 129, 257, 281
 van Dam, A. xvii, 194
 Van Den Driessche, P. 338
 van der Driessche, P. 708
 van der Merwe, R. 676
 Van Dooren, P. 362
 Van Loan, C. F. xvii, xviii, 127, 357, 416, 614, 646, 681, 692, 701, 722
 Van Overschee, P. 805
 Van Pelt, T. 498
 van Schagen, F. 805
 Vandebril, R. 125, 129
 Vandenberghe, L. xvii, 164
 Varadarajan, V. S. 230, 655, 682, 722
 Varah, J. M. 606
 Vardulakis, A. I. G. xvii, 281
 Varga, R. S. xviii, 269, 275, 706
 Vasic, P. M. 38, 52, 76, 158, 567
 Vasudeva, H. L. 533, 539, 541
 Vavrin, Z. 257
 Vein, P. R. 164
 Vein, R. xvii, 164
 Veljan, D. 42, 158, 160
 Venugopal, R. 230, 679
 Vermeer, J. 345
 Veselic, K. 330, 362, 701
 Vetter, W. J. 416
 Vidyasagar, M. 691
 Visick, G. 532, 535, 537, 539, 540
 Volenec, V. 53, 158, 463
 Vong, S.-W. 584
 Vowe, M. 29
 Vreugdenhil, R. 805
 Vuorinen, M. 23, 27, 29

 Wada, S. 60, 63, 64, 566
 Wagner, D. G. 697
 Waldenstrom, S. 196
 Walter, G. G. xvii, 708
 Wang, B. 518, 519, 614
 Wang, B.-Y. 198, 519, 528, 534, 541, 614
 Wang, C.-L. 57, 65
 Wang, D. 433
 Wang, F. 224
 Wang, G. 129, 202, 378, 394, 398
 Wang, J. 442, 542
 Wang, J.-H. 361
 Wang, L. 51, 598
 Wang, L.-C. 58
 Wang, Q.-G. 271
 Wang, X. 696
 Wang, Y. 715
 Wang, Y. W. 689, 798
 Wanner, G. 678
 Wansbeek, T. 416
 Ward, A. J. B. 320
 Ward, R. C. 706
 Warga, J. 625, 626
 Warner, W. H. 188
 Waterhouse, W. E. 136, 137
 Waters, S. R. 221
 Wathen, A. J. 476
 Watkins, W. 278, 450, 484, 488, 494, 513, 533, 536
 Watson, G. S. 57, 66
 Weaver, J. R. 230, 314
 Weaver, O. L. 150, 172, 655, 660, 722
 Webb, J. H. 45
 Webster, R. 164
 Wegert, E. 715
 Wegmann, R. 597
 Wei, J. 678
 Wei, M. 316, 380
 Wei, Y. 129, 202, 378, 391, 393, 394, 398, 680
 Weinberg, D. A. 361
 Weiss, G. H. 678
 Weiss, M. xvii, 805
 Wenzel, D. 584
 Wermuth, E. M. E. 681, 694
 Werner, H.-J. 527
 Wesseling, P. xvii
 Westlake, J. R. xviii
 Wets, R. J. B. 632, 642
 Weyrauch, M. 685
 White, J. E. 357
 Wiegmann, N. A. 343
 Wiener, Z. 714
 Wigner, E. P. 457
 Wilcox, R. M. xviii, 684
 Wildon, M. J. 229
 Wilhelm, F. 689
 Wilker, J. B. 29
 Wilkinson, J. H. xviii
 Willems, J. C. 805
 Williams, E. R. 378
 Williams, J. P. 319, 348, 376, 431
 Williams, K. S. 565
 Wilson, D. A. 619
 Wilson, P. M. H. 161, 193
 Wimmer, H. K. 257, 310, 459, 585, 788, 793, 796, 805
 Wirth, F. 699
 Witkowski, A. 23, 27
 Witzgall, C. 164, 619, 635
 Wolkowicz, H. 51, 180, 190, 322, 332, 341
 Wolovich, W. A. 805
 Wonenburger, M. J. 350
 Wong, C. S. 350, 478, 613
 Wonham, W. M. xvii, 262, 646, 791, 805
 Woo, C.-W. 453, 483, 493, 501, 505, 508, 510
 Wrobel, I. 348
 Wu, C.-F. 527

- Wu, P. Y. 208, 349–351,
 360, 361, 542, 701
 Wu, S. 42, 59, 64, 158
 Wu, Y.-D. 158

 Xi, B.-Y. 528, 614
 Xiao, Z.-G. 54
 Xie, Q. 416
 Xu, C. 70, 145, 472, 480,
 490, 565
 Xu, D. 193
 Xu, H. 650, 701
 Xu, Z. 70, 145, 472, 490,
 565

 Yakub, A. 45
 Yamagami, S. 584
 Yamazaki, T. 53, 522
 Yanase, M. M. 457
 Yang, B. 63
 Yang, X. 697
 Yang, Z. P. 476, 478, 480
 Yau, S. F. 535, 538
 Yau, S. S.-T. 673

 Ye, Q. 362
 Yeadon, F. I. 188
 Yellen, J. xvii
 Young, D. M. xviii
 Young, N. J. 607
 Young, P. M. 701

 Zakai, M. 722, 746, 747,
 805
 Zanna, A. 188, 674, 677,
 678, 682
 Zassenhaus, H. 361
 Zelevinsky, A. V. 416
 Zemanek, J. 695
 Zhan, X. 164, 327, 360,
 413, 434, 442, 447, 448,
 486, 509–511, 514–516,
 521, 529–531, 535, 537,
 539–541, 549, 580, 582,
 587, 590, 612, 615, 685,
 695
 Zhang, C.-E. 17
 Zhang, F. 70, 126, 127,
 131, 145, 153, 184, 185,
 198, 219, 229, 310, 312,
 333, 342, 357, 393, 452,
 453, 456, 458, 461, 465,
 469, 471–473, 476, 478,
 483, 489, 490, 493,
 501–503, 505, 508, 510,
 518, 519, 525, 528, 533,
 534, 539, 541, 565, 614
 Zhang, L. 393
 Zhang, Z.-H. 54, 158
 Zhao, K. 312
 Zheng, B. 124, 391
 Zhong, Q.-C. 206
 Zhou, K. xvii, 281, 607,
 610, 787
 Zhu, H. 38
 Zhu, L. 26, 29
 Zielke, G. 573, 592
 Zlobec, S. 366
 Zwart, H. J. 691
 Zwas, G. 603
 Zwillinger, D. 75, 158, 161,
 255, 357

Index

Symbols

- $0_{n \times m}$
 $n \times m$ zero matrix
definition, 83
- $1_{n \times m}$
 $n \times m$ ones matrix
definition, 84
- 2×2 matrices
commutator
Fact 2.18.1, 149
- 2×2 matrix
discrete-time
asymptotically
stable matrix
Fact 11.21.1, 712
eigenvalue inequality
Fact 8.17.1, 508
singular value
Fact 5.11.31, 328
- 2×2
positive-semidefinite
matrix
square root
Fact 8.9.6, 451
- 2×2 trace
Fact 2.12.9, 126
- 3×3 matrix identity
trace
Fact 4.9.5, 261
- 3×3 symmetric matrix
eigenvalue
Fact 4.10.1, 265
- $A \oplus B$
Kronecker sum
definition, 403
- $A \# B$
geometric mean
definition, 461
- $A \#_{\alpha} B$
generalized
geometric mean
definition, 464
- A^{-1}
inverse matrix
definition, 101
- $A \stackrel{\text{GL}}{\leq} B$
generalized Löwner
partial ordering
definition, 524
- $A \stackrel{\text{rs}}{\leq} B$
rank subtractivity
partial ordering
definition, 119
- $A \stackrel{*}{\leq} B$
star partial ordering
definition, 120
- $A \stackrel{i}{\leftarrow} b$
column replacement
definition, 80
- $A \circ B$
Schur product
definition, 404
- $A \otimes B$
Kronecker product
definition, 400
- $A : B$
parallel sum
definition, 528
- $A^{\hat{*}}$
reverse complex
conjugate transpose
definition, 88
- $A^{\circ \alpha}$
Schur power
definition, 404
- A^{+}
generalized inverse
definition, 363
- $A^{1/2}$
positive-semidefinite
matrix square root
definition, 431
- $A^{\#}$
group generalized
inverse
definition, 369
- A^A
adjugate
definition, 105
- A^D
Drazin generalized
inverse
definition, 367
- A^L
left inverse
definition, 98
- A^R
right inverse
definition, 98

A^T **transpose**

definition, 86

 $A^{\hat{T}}$ **reverse transpose**

definition, 88

 $A_{[i;j]}$ **submatrix**

definition, 105

 $A_{\perp}$ **complementary****idempotent matrix**

definition, 176

complementary**projector**

definition, 175

 $B(p, q)$ **Bezout matrix**

definition, 255

 $C(p)$ **companion matrix**

definition, 283

 C^* **complex conjugate****transpose**

definition, 87

 $D|\mathcal{A}$ **Schur complement**

definition, 367

 $E_{i,j,n \times m}$ $n \times m$ **matrix with a single unit entry**

definition, 84

 $E_{i,j}$ **matrix with a single unit entry**

definition, 84

 $H(g)$ **Hankel matrix**

definition, 257

 I_n **identity matrix**

definition, 83

 J

$$\begin{bmatrix} 0 & I \\ -I & 0 \end{bmatrix}$$

definition, 169

 J_{2n}

$$\begin{bmatrix} 0 & I_n \\ -I_n & 0 \end{bmatrix}$$

definition, 169

 $K(x)$ **cross-product matrix**

definition, 82

 N **standard nilpotent matrix**

definition, 166

 N_n $n \times n$ **standard nilpotent matrix**

definition, 166

 $P_{A,B}$ **pencil**

definition, 304

 $P_{n,m}$ **Kronecker****permutation matrix**

definition, 402

 $V(\lambda_1, \dots, \lambda_n)$ **Vandermonde matrix**

definition, 354

 $[A, B]$ **commutator**

definition, 82

 $\mathbb{B}_{\varepsilon}(x)$ **open ball**

definition, 621

 $\mathbb{C}^{n \times m}$ $n \times m$ **complex matrices**

definition, 79

 $\mathbb{F}$ **real or complex numbers**

definition, 78

 $\mathbb{F}(s)$ **rational functions**

definition, 249

 $\mathbb{F}[s]$ **polynomials with coefficients in $\mathbb{F}$**

definition, 231

 $\mathbb{F}^{n \times m}$ $n \times m$ **real or complex matrices**

definition, 79

 $\mathbb{F}^{n \times m}[s]$ **polynomial matrices with coefficients in $\mathbb{F}^{n \times m}$**

definition, 234

 $\mathbb{F}^{n \times m}(s)$ $n \times m$ **rational transfer functions**

definition, 249

 $\mathbb{F}_{\text{prop}}^{n \times m}(s)$ $n \times m$ **proper rational transfer functions**

definition, 249

 $\mathbb{F}_{\text{prop}}(s)$ **proper rational functions**

definition, 249

 $\mathbb{R}$ **complex numbers**

definition, 78

real numbers

definition, 78

 $\mathbb{R}^{n \times m}$ $n \times m$ **real matrices**

definition, 79

 $\mathbb{S}_{\varepsilon}(x)$ **sphere**

definition, 621

 $\mathbb{H}^n$ $n \times n$ **Hermitian matrices**

definition, 417

 $\mathbb{N}^n$

- $n \times n$ **positive-semidefinite matrices**
definition, 417
- $\mathbf{P}^n$
 $n \times n$ **positive-definite matrices**
definition, 417
- Im x**
imaginary part
definition, 77
- In A**
inertia
definition, 245
- Re x**
real part
definition, 77
- $\mathcal{C}(A, B)$
controllable subspace
definition, 737
- $\mathcal{H}$
Hamiltonian
definition, 780
- $\mathcal{H}(G)$
Markov block-Hankel matrix
definition, 754
- $\mathcal{H}_l(q)$
hypercompanion matrix
definition, 288
- $\mathcal{H}_{i,j,k}(G)$
Markov block-Hankel matrix
definition, 754
- $\mathcal{J}_l(q)$
real Jordan matrix
definition, 289
- $\mathcal{K}(A, B)$
controllability matrix
definition, 737
- $\mathcal{L}\{x(t)\}$
Laplace transform
definition, 646
- $\mathcal{N}(A)$
null space
definition, 94
- $\mathcal{O}(A, C)$
observability matrix
definition, 728
- $\mathcal{R}(A)$
range
definition, 93
- $\mathcal{S}^\perp$
orthogonal complement
definition, 91
- $\mathcal{S}_s(A)$
asymptotically stable subspace
definition, 665
- $\mathcal{S}_u(A)$
unstable subspace
definition, 665
- $\mathcal{U}(A, C)$
unobservable subspace
definition, 728
- $\mathcal{X}^\sim$
complement
definition, 2
- $\mathcal{Y} \setminus \mathcal{X}$
relative complement
definition, 2
- $\|A\|_p$
Hölder norm
definition, 547
- $\|A\|_F$
Frobenius norm
definition, 547
- $\|A\|_{\text{col}}$
column norm
definition, 556
- $\|A\|_{\text{row}}$
row norm
definition, 556
- $\|A\|_{\sigma p}$
Schatten norm
definition, 548
- $\|A\|_{q,p}$
Hölder-induced norm
definition, 554
- $\|x\|_p$
Hölder norm
definition, 544
- $\|y\|_D$
dual norm
definition, 570
- ad_A
adjoint operator
definition, 82
- $\text{aff } \mathcal{S}$
affine hull
definition, 90
- $\text{bd } \mathcal{S}$
boundary
definition, 622
- $\text{bd}_g, \mathcal{S}$
relative boundary
definition, 622
- χ_A
characteristic equation
definition, 240
- $\chi_{A,B}$
characteristic polynomial
definition, 305
- $\text{cl } \mathcal{S}$
closure
definition, 621
- $\text{cl}_g, \mathcal{S}$
relative closure
definition, 622
- $\text{co } \mathcal{S}$
convex hull

- definition, 89
- coco $\mathcal{S}$**
 - convex conical hull**
 - definition, 89
- col _{i} (A)**
 - column**
 - definition, 79
- cone $\mathcal{S}$**
 - conical hull**
 - definition, 89
- dcone $\mathcal{S}$**
 - dual cone**
 - definition, 91
- def A**
 - defect**
 - definition, 96
- deg p**
 - degree**
 - definition, 231
- det A**
 - determinant**
 - definition, 103
- diag($A_1, \dots, A_k$)**
 - block-diagonal matrix**
 - definition, 167
- diag($a_1, \dots, a_n$)**
 - diagonal matrix**
 - definition, 167
- dim $\mathcal{S}$**
 - dimension of a set**
 - definition, 90
- $\ell(A)$**
 - lower bound**
 - definition, 558
- $\ell_{q,p}(A)$**
 - Hölder-induced lower bound**
 - definition, 559
- $\hat{I}_n$**
 - reverse identity matrix**
 - definition, 84
- ind A**
 - index of a matrix**
 - definition, 176
- ind _{A} (λ)**
 - index of an eigenvalue**
 - definition, 295
- int $\mathcal{S}$**
 - interior**
 - definition, 621
- int _{$\mathcal{S}'$} $\mathcal{S}$**
 - relative interior**
 - definition, 621
- $\lambda_1(A)$**
 - maximum eigenvalue**
 - definition, 240
 - minimum eigenvalue**
 - definition, 240
- $\lambda_i(A)$**
 - eigenvalue**
 - definition, 240
- log(A)**
 - matrix logarithm**
 - definition, 654
- mroots(p)**
 - multiset of roots**
 - definition, 232
- mspec(A)**
 - multispectrum**
 - definition, 240
- μ_A**
 - minimal polynomial**
 - definition, 247
- $\nu_-(A), \nu_0(A)$**
 - inertia**
 - definition, 245
- $\overline{C}$**
 - complex conjugate**
 - definition, 87
- π**
 - prime numbers**
 - Fact 1.7.8, 19
- polar $\mathcal{S}$**
- dual cone**
 - definition, 91
- rank A**
 - rank**
 - definition, 95
- rank G**
 - normal rank for a rational transfer function**
 - definition, 249
- rank P**
 - normal rank for a polynomial matrix**
 - definition, 235
- reldeg G**
 - relative degree**
 - definition, 249
- revdiag($a_1, \dots, a_n$)**
 - reverse diagonal matrix**
 - definition, 167
- $d_{\max}(A)$**
 - maximum diagonal entry**
 - definition, 80
- $d_{\min}(A)$**
 - minimum diagonal entry**
 - definition, 80
- $d_i(A)$**
 - diagonal entry**
 - definition, 80
- roots(p)**
 - set of roots**
 - definition, 232
- row _{i} (A)**
 - row**
 - definition, 79
- sig A**
 - signature**
 - definition, 245
- $\sigma_{\max}(A)$**

maximum singular value
definition, 301

$\sigma_{\min}(A)$
minimum singular value
definition, 301

$\sigma_i(A)$
singular value
definition, 301

sign x
sign
definition, 89

sign α
sign
definition, xxi

spabs(A)
spectral abscissa
definition, 245

spec(A)
spectrum
definition, 240

sprad(A)
spectral abscissa
definition, 245

tr A
trace
definition, 86

vccone($\mathcal{D}, x_0$)
variational cone
definition, 625

vec A
column-stacking operator
definition, 399

$|x|$
absolute value
definition, 88

e^A
matrix exponential
definition, 643

e_i

i **th column of the identity matrix**
definition, 84

$e_{i,n}$
 i **th column of the $n \times n$ identity matrix**
definition, 84

$f^{(k)}(x_0)$
 k **th Fréchet derivative**
definition, 627

$f'(x_0)$
Fréchet derivative
definition, 626

k th Fréchet derivative
definition, 627

n -tuple
definition, 3

$x \gg 0$
positive vector
definition, 79

$x \geq 0$
nonnegative vector
definition, 79

SO(3)
logarithm
Fact 11.15.10, 692

SO(n)
eigenvalue
Fact 5.11.2, 321

amult $_A(\lambda)$
algebraic multiplicity
definition, 240

circ($a_0, \dots, a_{n-1}$)
circulant matrix
definition, 355

exp(A)
matrix exponential
definition, 643

glb(S)
greatest lower bound
definition, 7

gmult $_A$
geometric multiplicity
definition, 245

inf(S)
infimum
definition, 7

lub(S)
least upper bound
definition, 7

mult $_p(\lambda)$
multiplicity
definition, 232

sh(A, B)
shorted operator
definition, 530

sup(S)
supremum
definition, 7

D $_+f(x_0; \xi)$
one-sided directional differential
definition, 625

(1)-inverse
definition, 364
determinant
Fact 6.5.28, 393
left inverse
Proposition 6.1.3, 364
right inverse
Proposition 6.1.2, 364

(1,2)-inverse
definition, 364

A

Abel
quintic polynomial
Fact 3.21.7, 223

Abelian group
definition
Definition 3.3.3, 172
equivalence relation
Proposition 3.4.2, 173

absolute norm

monotone norm
Proposition 9.1.2, 543

absolute sum norm

definition, 545

absolute value

Frobenius norm
Fact 9.13.11, 603

Hölder-induced
norm

Fact 9.8.26, 576

inequality

Fact 1.11.24, 45

Fact 1.11.25, 45

irreducible matrix

Fact 3.20.4, 218

matrix, 88

maximum singular
value

Fact 9.13.10, 603

reducible matrix

Fact 3.20.4, 218

scalar inequality

Fact 1.11.1, 39

Fact 1.11.12, 43

Fact 1.12.3, 46

Schatten norm

Fact 9.13.11, 603

spectral radius

Fact 4.11.16, 279

vector, 88**absolute-value function**

Niculescu's
inequality
Fact 1.10.19, 33

absolute-value matrix

positive-semidefinite
matrix
Fact 8.9.1, 450

**absolutely convergent
sequence**

convergent sequence
Proposition 10.2.7, 623
Proposition 10.2.9, 623

**absolutely convergent
series****definition**

Definition 10.2.6, 623

Definition 10.2.8, 623

Aczel's inequality**norm inequality**

Fact 9.7.4, 563

quadratic inequality

Fact 1.16.19, 64

additive compound

asymptotically stable
polynomial

Fact 11.17.12, 697

additive decomposition

diagonalizable
matrix

Fact 5.9.3, 311

Hermitian matrix

Fact 3.7.29, 183

nilpotent matrix

Fact 5.9.3, 311

orthogonal matrix

Fact 5.19.2, 360

Fact 5.19.3, 360

unitary matrix

Fact 5.19.1, 360

adjacency matrix**definition**

Definition 3.2.1, 170

graph of a matrix

Proposition 3.2.5, 171

inbound Laplacian
matrix

Theorem 3.2.2, 170

Laplacian matrix

Theorem 3.2.2, 170

Theorem 3.2.3, 171

Fact 4.11.11, 277

outbound Laplacian
matrix

Theorem 3.2.2, 170

symmetric graph

Fact 4.11.1, 272

adjacent

Definition 1.4.2, 8

adjoint norm**definition**

Fact 9.8.8, 572

dual norm

Fact 9.8.8, 572

**Hölder-induced
norm**

Fact 9.8.10, 572

adjoint operator**commutator**

Fact 2.18.5, 149

Fact 2.18.6, 150

adjugate

basic properties, 106

characteristic**polynomial**

Fact 4.9.8, 261

cross product

Fact 6.5.16, 389

defect

Fact 2.16.7, 143

definition, 105**derivative**

Fact 10.11.19, 640

Fact 10.11.21, 641

determinant

Fact 2.14.27, 139

Fact 2.16.3, 141

Fact 2.16.5, 142

Fact 2.16.6, 142

diagonalizable**matrix**

Fact 5.14.5, 339

eigenvalue

Fact 4.10.7, 267

eigenvector

Fact 5.14.26, 342

elementary matrix

Fact 2.16.1, 141

factor

Fact 2.16.9, 143

Frobenius norm

Fact 9.8.15, 573

generalized inverse

Fact 6.3.6, 370

Fact 6.3.7, 371

Fact 6.5.16, 389

Hermitian matrix

Fact 3.7.10, 179

iterated

Fact 2.16.5, 142
matrix powers
 Fact 4.9.8, 261
matrix product
 Fact 2.16.10, 143
nilpotent matrix
 Fact 6.3.6, 370
null space
 Fact 2.16.7, 143
outer-product
perturbation
 Fact 2.16.3, 141
partitioned matrix
 Fact 2.14.27, 139
range
 Fact 2.16.7, 143
rank
 Fact 2.16.7, 143
 Fact 2.16.8, 143
scalar factor
 Fact 2.16.5, 142
singular value
 Fact 5.11.36, 328
skew-Hermitian matrix
 Fact 3.7.10, 179
 Fact 3.7.11, 179
skew-symmetric matrix
 Fact 4.9.20, 263
spectrum
 Fact 4.10.7, 267
trace
 Fact 4.9.8, 261
transpose
 Fact 2.16.5, 142
affine closed half space
 closed half space
 Fact 2.9.6, 111
 definition, 91
affine function
 definition, 81
affine hull
 closure
 Fact 10.8.11, 633
 constructive
 characterization
 Theorem 2.3.5, 91

convex hull
 Fact 2.9.3, 110
convex set
 Theorem 10.3.2, 624
 Fact 10.8.8, 632
 definition, 90
 linear mapping
 Fact 2.10.4, 115
affine hyperplane
 affine subspace
 Fact 2.9.6, 111
 definition, 91
 determinant
 Fact 2.20.3, 154
affine mapping
 Hermitian matrix
 Fact 3.7.14, 181
 normal matrix
 Fact 3.7.14, 181
affine open half space
 definition, 91
 open half space
 Fact 2.9.6, 111
affine subspace
 affine hull of image
 Fact 2.9.26, 113
 affine hyperplane
 Fact 2.9.6, 111
 definition, 89
 image under linear
 mapping
 Fact 2.9.26, 113
 left inverse
 Fact 2.9.26, 113
 span
 Fact 2.9.7, 111
 Fact 2.20.4, 154
 Fact 10.8.12, 633
 subspace
 Fact 2.9.8, 111
Afriat
 spectrum of a
 product of
 projectors
 Fact 5.12.15, 335
Akers

maximum singular
 value of a product
 of elementary
 projectors
 Fact 9.14.1, 607
algebraic multiplicity
 block-triangular
 matrix
 Proposition 5.5.13, 298
 definition
 Definition 4.4.4, 240
 geometric
 multiplicity
 Proposition 5.5.3, 295
 index of an
 eigenvalue
 Proposition 5.5.6, 296
orthogonal matrix
 Fact 5.11.2, 321
outer-product matrix
 Fact 5.14.3, 338
almost nonnegative matrix
 asymptotically stable
 matrix
 Fact 11.19.5, 707
 definition, 230
 Definition 3.1.4, 168
 group-invertible
 matrix
 Fact 11.19.4, 706
irreducible matrix
 Fact 11.19.2, 706
Lyapunov-stable matrix
 Fact 11.19.4, 706
matrix exponential
 Fact 11.19.1, 706
 Fact 11.19.2, 706
N-matrix
 Fact 11.19.3, 706
 Fact 11.19.5, 707
nonnegative matrix
 Fact 11.19.1, 706
positive matrix
 Fact 11.19.2, 706
alternating group

- group
 - Fact 3.21.7, 223
- Alzer's inequality**
 - sum of integers
 - Fact 1.9.31, 30
- Amemiya's inequality**
 - Schur product
 - Fact 8.21.39, 539
- Anderson**
 - rank of a tripotent matrix
 - Fact 2.10.23, 118
- Ando**
 - convex function
 - Proposition 8.6.17, 542
 - inertia of congruent, normal matrices
 - Fact 5.10.17, 319
- angle**
 - definition, 85
- angular velocity vector**
 - quaternions
 - Fact 11.11.15, 675
- antieigenvalue**
 - definition
 - Fact 9.8.37, 577
- antisymmetric graph**
 - Laplacian
 - Fact 4.11.1, 272
- antisymmetric relation**
 - definition
 - Definition 1.3.8, 7
 - one-sided cone
 - induced by
 - Proposition 2.3.6, 93
 - positive-semidefinite matrix
 - Proposition 8.1.1, 417
- aperiodic graph**
 - Definition 1.4.3, 9
 - nonnegative matrix
 - Fact 4.11.5, 273
- Araki**
 - positive-semidefinite matrix inequality
 - Fact 8.12.21, 480
- Araki-Lieb-Thirring inequality**
 - positive-semidefinite matrix inequality
 - Fact 8.12.20, 479
- arc**
 - definition, 8
- area**
 - parallelogram
 - Fact 2.20.17, 160
 - Fact 9.7.5, 565
 - polygon
 - Fact 2.20.14, 159
 - triangle
 - Fact 2.20.7, 155
 - Fact 2.20.8, 156
 - Fact 2.20.10, 156
- arithmetic mean**
 - Carleman's inequality
 - Fact 1.15.40, 58
 - geometric mean
 - Fact 1.10.36, 37
 - Fact 1.15.21, 53
 - Fact 1.15.23, 53
 - Fact 1.15.24, 54
 - Fact 1.15.25, 54
 - Fact 1.15.26, 54
 - Fact 1.15.27, 54
 - identric mean
 - Fact 1.10.36, 37
 - logarithmic mean
 - Fact 1.15.26, 54
 - mixed arithmetic-geometric mean inequality
 - Fact 1.15.39, 58
 - Muirhead's theorem
 - Fact 1.15.25, 54
 - positive-definite matrix
 - Fact 8.10.34, 460
 - scalar inequality
 - Fact 1.11.6, 39
- Fact 1.11.7, 39
- Fact 1.11.8, 40
- Fact 1.11.9, 41
- Fact 1.11.10, 41
- arithmetic-mean inequality**
 - harmonic mean
 - Fact 1.15.16, 52
 - Fact 1.15.17, 52
- arithmetic-mean–geometric-mean inequality**
 - alternative form
 - Fact 1.15.33, 56
 - difference
 - Fact 1.15.29, 55
 - harmonic mean
 - Fact 1.15.15, 52
 - Jensen's inequality
 - Fact 1.8.4, 21
 - main form
 - Fact 1.15.14, 51
 - Fact 1.15.28, 54
- Popoviciu**
 - Fact 1.15.29, 55
- positive-definite matrix**
 - Fact 8.13.8, 486
- quartic identity**
 - Fact 1.12.5, 47
- Rado**
 - Fact 1.15.29, 55
- ratio**
 - Fact 1.15.29, 55
- reverse inequality**
 - Fact 1.15.18, 52
 - Fact 1.15.19, 52
- sextic identity**
 - Fact 1.13.1, 47
- variation**
 - Fact 1.10.13, 32
- weighted arithmetic-mean–geometric-mean inequality**
 - Fact 1.15.32, 56

- arithmetic-mean–harmonic-mean inequality**
 - scalar inequality
 - Fact 1.15.37, 57
- associative identities**
 - definition, 82
- associativity**
 - composition
 - Proposition 1.2.1, 3
- asymptotic stability**
 - eigenvalue
 - Proposition 11.8.2, 662
 - input-to-state stability
 - Fact 12.20.18, 793
 - linear dynamical system
 - Proposition 11.8.2, 662
 - Lyapunov equation**
 - Corollary 11.9.1, 666
 - matrix exponential**
 - Proposition 11.8.2, 662
 - nonlinear system**
 - Theorem 11.7.2, 661
- asymptotically stable equilibrium**
 - definition
 - Definition 11.7.1, 660
- asymptotically stable matrix**
 - 2×2 matrix
 - Fact 11.18.35, 705
 - almost nonnegative matrix
 - Fact 11.19.5, 707
 - asymptotically stable polynomial
 - Proposition 11.8.4, 663
 - Cayley transform**
 - Fact 11.21.8, 713
 - compartmental matrix**
 - Fact 11.19.6, 707
 - controllability**
 - Fact 12.20.5, 791
 - controllability Gramian**
 - Proposition 12.7.9, 747
 - Corollary 12.7.10, 747
 - controllable pair**
 - Proposition 12.7.9, 747
 - Corollary 12.7.10, 747
 - controllably asymptotically stable**
 - Proposition 12.8.3, 747
 - Proposition 12.8.5, 748
 - cyclic matrix**
 - Fact 11.18.25, 702
 - definition**
 - Definition 11.8.1, 662
 - detectability**
 - Proposition 12.5.5, 735
 - Corollary 12.5.6, 735
 - diagonalizable over $\mathbb{R}$**
 - Fact 11.17.10, 697
 - discrete-time asymptotically stable matrix**
 - Fact 11.21.8, 713
 - dissipative matrix**
 - Fact 11.18.21, 701
 - Fact 11.18.37, 705
 - factorization**
 - Fact 11.18.22, 701
 - integral**
 - Lemma 11.9.2, 667
 - inverse matrix**
 - Fact 11.18.15, 700
 - Kronecker sum**
 - Fact 11.18.32, 704
 - Fact 11.18.33, 704
 - Fact 11.18.34, 705
 - linear matrix equation**
 - Proposition 11.9.3, 667
 - logarithmic derivative**
 - Fact 11.18.11, 699
 - Lyapunov equation**
 - Proposition 11.9.5, 668
 - Corollary 11.9.4, 668
 - Corollary 11.9.7, 669
 - Corollary 12.4.4, 734
 - Corollary 12.5.6, 735
 - Corollary 12.7.4, 746
 - Corollary 12.8.6, 749
 - Fact 12.21.7, 795
 - Fact 12.21.17, 797
 - matrix exponential**
 - Lemma 11.9.2, 667
 - Fact 11.18.8, 699
 - Fact 11.18.9, 699
 - Fact 11.18.10, 699
 - Fact 11.18.15, 700
 - Fact 11.18.18, 701
 - Fact 11.18.19, 701
 - Fact 11.21.7, 713
 - minimal realization**
 - Definition 12.9.17, 757
 - negative-definite matrix**
 - Fact 11.18.30, 704
 - nonsingular N-matrix**
 - Fact 11.19.5, 707
 - normal matrix**
 - Fact 11.18.37, 705
 - observability Gramian**
 - Corollary 12.4.10, 734
 - observable pair**
 - Proposition 12.4.9, 734
 - Corollary 12.4.10, 734
 - observably asymptotically stable**
 - Proposition 11.9.5, 735
 - Proposition 12.5.5, 735
 - perturbation**
 - Fact 11.18.16, 700
 - positive-definite matrix**
 - Proposition 11.9.5, 668
 - Proposition 12.4.9, 734
 - Corollary 11.9.7, 669
 - Fact 11.18.21, 701
 - secant condition**
 - Fact 11.18.29, 704
 - sign of entries**
 - Fact 11.19.5, 708
 - sign stability**
 - Fact 11.19.5, 708

- similar matrices**
 - Fact 11.18.4, 698
 - skew-Hermitian matrix**
 - Fact 11.18.30, 704
 - spectrum**
 - Fact 11.18.13, 700
 - square root**
 - Fact 11.18.36, 705
 - stability radius**
 - Fact 11.18.17, 700
 - stabilizability**
 - Proposition 11.9.5, 735
 - Proposition 12.8.3, 747
 - Proposition 12.8.5, 748
 - Corollary 12.8.6, 749
 - subdeterminant**
 - Fact 11.19.1, 707
 - trace**
 - Fact 11.18.31, 704
 - tridiagonal matrix**
 - Fact 11.18.24, 702
 - Fact 11.18.25, 702
 - Fact 11.18.26, 702
 - Fact 11.18.27, 703
 - Fact 11.18.28, 703
 - asymptotically stable polynomial**
 - additive compound**
 - Fact 11.17.12, 697
 - asymptotically stable matrix**
 - Proposition 11.8.4, 663
 - definition**
 - Definition 11.8.3, 663
 - even polynomial**
 - Fact 11.17.6, 696
 - Hermite-Biehler theorem**
 - Fact 11.17.6, 696
 - interlacing theorem**
 - Fact 11.17.6, 696
 - Kharitonov's theorem**
 - Fact 11.17.13, 698
 - Kronecker sum**
 - Fact 11.17.11, 697
 - odd polynomial**
 - Fact 11.17.6, 696
 - polynomial coefficients**
 - Fact 11.17.2, 696
 - Fact 11.17.3, 696
 - Fact 11.17.7, 697
 - Fact 11.17.8, 697
 - Fact 11.17.10, 697
 - Fact 11.17.11, 697
 - Fact 11.17.12, 697
 - reciprocal argument**
 - Fact 11.17.4, 696
 - Schur product of polynomials**
 - Fact 11.17.9, 697
 - subdeterminant**
 - Fact 11.18.23, 702
 - asymptotically stable subspace**
 - definition**, 665
 - asymptotically stable transfer function**
 - minimal realization**
 - Proposition 12.9.18, 757
 - SISO entries**
 - Proposition 12.9.19, 757
 - average**
 - positive-semidefinite matrix**
 - Fact 5.19.5, 360
 - averaged limit**
 - integral**
 - Fact 10.11.6, 638
- B**
- Baker-Campbell-Hausdorff series**
 - matrix exponential**
 - Proposition 11.4.7, 655
 - Baker-Campbell-Hausdorff-Dynkin expansion**
 - time-varying dynamics**
 - Fact 11.13.4, 678
 - balanced realization**
 - definition**
 - Definition 12.9.20, 757
 - minimal realization**
 - Proposition 12.9.21, 757
 - balancing transformation**
 - existence**
 - Corollary 8.3.3, 423
 - Bandila's inequality triangle**
 - Fact 2.20.11, 156
 - Barnett**
 - asymptotic stability of a tridiagonal matrix**
 - Fact 11.18.24, 702
 - Barnett factorization**
 - Bezout matrix**
 - Fact 4.8.6, 255
 - barycentric coordinates**
 - conjugate parameters**
 - Fact 1.16.11, 62
 - definition**, 89
 - basis**
 - definition**, 90
 - Beckner's two-point inequality**
 - powers**
 - Fact 1.10.15, 33
 - Fact 9.9.35, 586
 - Bellman**
 - quadratic form inequality**
 - Fact 8.15.7, 501
 - Ben-Israel**
 - generalized inverse**
 - Fact 6.3.35, 376

Bencze

arithmetic-mean–
geometric-mean–
logarithmic-mean
inequality

Fact 1.15.26, 54

Bendixson's theorem

eigenvalue bound

Fact 5.11.21, 325

Fact 9.11.8, 598

Berezin

trace of a convex
function

Fact 8.12.33, 482

Bergstrom

positive-definite
matrix determinant

Fact 8.13.15, 488

Bergstrom's inequality

quadratic form

Fact 8.11.3, 468

Fact 8.15.18, 503

Bernoulli matrix

Vandermonde matrix

Fact 5.16.3, 354

Bernoulli's inequality

scalar inequality

Fact 1.9.1, 22

Fact 1.9.2, 23

Bernstein matrix

Vandermonde matrix

Fact 5.16.3, 354

Bernstein's inequality

matrix exponential

Fact 11.15.4, 689

Berwald

polynomial root
bounds

Fact 11.20.11, 711

Bessel's inequality

norm inequality

Fact 9.7.4, 563

Bessis-Moussa-Villani

trace conjecture

derivative of a

matrix exponential

Fact 8.12.31, 482

power of a positive-
semidefinite

matrix

Fact 8.12.30, 482

Bezout equation

coprime polynomials

Fact 4.8.5, 255

Bezout identity

right coprime

polynomial

matrices

Theorem 4.7.14, 252

Bezout matrix

coprime polynomials

Fact 4.8.6, 255

Fact 4.8.7, 257

Fact 4.8.8, 257

definition

Fact 4.8.6, 255

distinct roots

Fact 4.8.9, 258

factorization

Fact 5.15.24, 349

polynomial roots

Fact 4.8.9, 258

Bhatia

Schatten norm

inequality

Fact 9.9.45, 588

unitarily invariant

norm inequality

Fact 9.9.44, 588

bialternate product

compound matrix

Fact 7.5.17, 411

Kronecker product,

416

bidiagonal matrix

singular value

Fact 5.11.47, 332

biequivalent matrices

congruent matrices

Proposition 3.4.5, 174

definition

Definition 3.4.3, 174

Kronecker product

Fact 7.4.11, 405

rank

Proposition 5.1.3, 283

similar matrices

Proposition 3.4.5, 174

Smith form

Theorem 5.1.1, 283

Corollary 5.1.2, 283

unitarily similar

matrices

Proposition 3.4.5, 174

bijjective function

definition, 76

bilinear function

definition, 627

Binet-Cauchy formula

determinant

Fact 2.13.4, 129

Binet-Cauchy theorem

compound of a

matrix product

Fact 7.5.17, 411

binomial identity

sum

Fact 1.7.1, 14

Fact 1.7.2, 17

binomial series

infinite series

Fact 1.18.8, 73

bivector

parallelogram

Fact 9.7.5, 565

block

definition, 80

block decomposition

Hamiltonian

Proposition 12.17.5,
783

minimal realization

Proposition 12.9.10,
753

block-circulant matrix

circulant matrix

Fact 3.18.3, 215

Drazin generalized

inverse

Fact 6.6.1, 393

generalized inverse

Fact 6.5.2, 386

inverse matrix

Fact 2.17.6, 148

block-diagonal matrix

companion matrix

Proposition 5.2.8, 286

Lemma 5.2.2, 285

definition

Definition 3.1.3, 167

geometric

multiplicity

Proposition 5.5.13, 298

Hermitian matrix

Fact 3.7.8, 179

least common

multiple

Lemma 5.2.7, 286

matrix exponential

Proposition 11.2.8, 649

maximum singular

value

Fact 5.11.33, 328

minimal polynomial

Lemma 5.2.7, 286

normal matrix

Fact 3.7.8, 179

shifted-unitary

matrix

Fact 3.11.25, 196

similar matrices

Theorem 5.3.2, 288

Theorem 5.3.3, 289

singular value

Fact 8.18.9, 515

Fact 8.18.10, 515

Fact 9.14.21, 612

Fact 9.14.25, 613

skew-Hermitian

matrix

Fact 3.7.8, 179

unitary matrix

Fact 3.11.25, 196

block-Hankel matrix

definition

Definition 3.1.3, 167

Hankel matrix

Fact 3.18.3, 215

Markov

block-Hankel

matrix

definition, 754

block-Kronecker

product

Kronecker product,
416

block-Toeplitz matrix

definition

Definition 3.1.3, 167

Toeplitz matrix

Fact 3.18.3, 215

block-triangular matrix

algebraic multiplicity

Proposition 5.5.13, 298

controllable

dynamics

Theorem 12.6.8, 739

controllable subspace

Proposition 12.6.9, 739

Proposition 12.6.10,

740

controllably

asymptotically

stable

Proposition 12.7.3, 743

detectability

Proposition 12.5.4, 735

determinant

Fact 2.14.8, 134

index of a matrix

Fact 5.14.32, 343

Fact 6.6.13, 395

inverse matrix

Fact 2.17.1, 146

maximum singular

value

Fact 5.11.32, 328

minimal polynomial

Fact 4.10.12, 268

observable dynamics

Theorem 12.3.8, 730

observably

asymptotically

stable

Proposition 12.4.3, 732

spectrum

Proposition 5.5.13, 298

stabilizability

Proposition 12.8.4, 747

unobservable

subspace

Proposition 12.3.9, 730

Proposition 12.3.10,

730

blocking zero

definition

Definition 4.7.10, 251

rational transfer

function

Definition 4.7.4, 249

Smith-McMillan

form

Proposition 4.7.11, 251

Blundon

triangle inequality

Fact 2.20.11, 156

blunt cone

definition, 89

Bonami's inequality

powers

Fact 1.10.16, 33

Fact 9.7.20, 569

Borchers

trace norm of a

matrix difference

Fact 9.9.24, 584

Borobia

asymptotically stable

polynomial

Fact 11.17.8, 697

both

definition, 1

boundary

definition, 622

interior

Fact 10.8.7, 632

union
Fact 10.9.2, 634

boundary relative to a set
definition, 622

bounded set
continuous function
Theorem 10.3.10, 625
Corollary 10.3.11, 625
definition, 622
image under linear mapping
Fact 9.8.1, 571
open ball
Fact 10.8.2, 632

Bourbaki
polynomial root bound
Fact 11.20.4, 709

Bourin
spectral radius of a product
Fact 8.18.25, 520

Brahmagupta's formula
quadrilateral
Fact 2.20.13, 159

Brauer
spectrum bounds
Fact 4.10.21, 271

Brouwer fixed-point theorem
image of a continuous function
Corollary 10.3.11, 625

Brown
trace of a convex function
Fact 8.12.33, 482

Browne's theorem
eigenvalue bound
Fact 5.11.21, 325
Fact 5.11.22, 325
Fact 9.11.7, 598

Brownian motion

positive-semidefinite matrix
Fact 8.8.4, 446

Buzano's inequality
Cauchy-Schwarz inequality
Fact 1.17.2, 67
norm inequality
Fact 9.7.4, 563

C

Callan
determinant of a partitioned matrix
Fact 2.14.15, 136

Callebaut
monotonicity
Fact 1.16.1, 60

Callebaut's inequality
refined
Cauchy-Schwarz inequality
Fact 1.16.16, 63

canonical form
definition, 4

canonical mapping
definition, 4

Cantor intersection theorem
intersection of closed sets
Fact 10.9.11, 635

Cardano's trigonometric solution
cubic polynomial
Fact 4.10.1, 265
eigenvalue
Fact 4.10.1, 265

cardinality
definition, 2
inclusion-exclusion principle
Fact 1.5.5, 11

union
Fact 1.5.5, 11

Carleman's inequality
arithmetic mean
Fact 1.15.40, 58

Carlson
inertia of a Hermitian matrix
Fact 12.21.4, 794

Carlson inequality
sum of powers
Fact 1.15.41, 58

Carmichael
polynomial root bound
Fact 11.20.10, 711

Cartesian decomposition
determinant
Fact 8.13.4, 485
Fact 8.13.11, 486
eigenvalue
Fact 5.11.21, 325
Hermitian matrix
Fact 3.7.27, 182
Fact 3.7.28, 183
Fact 3.7.29, 183
positive-semidefinite matrix
Fact 9.9.40, 587

Schatten norm
Fact 9.9.37, 586
Fact 9.9.38, 587
Fact 9.9.39, 587
Fact 9.9.40, 587

singular value
Fact 8.18.7, 514

skew-Hermitian matrix
Fact 3.7.27, 182
Fact 3.7.28, 183
Fact 3.7.29, 183

spectrum
Fact 5.11.21, 325

Cartesian product
definition, 3

cascade**interconnection**

definition, 770

transfer functionProposition 12.13.2,
770**cascaded systems****geometric****multiplicity**

Fact 12.22.15, 801

Cauchy**polynomial root****bounds**

Fact 11.20.11, 711

**Cauchy interlacing
theorem****Hermitian matrix****eigenvalue**

Lemma 8.4.4, 425

Cauchy matrix**determinant**

Fact 3.20.14, 220

Fact 3.20.15, 221

positive-definite**matrix**

Fact 8.8.16, 449

Fact 12.21.18, 797

positive-semidefinite**matrix**

Fact 8.8.7, 447

Fact 8.8.9, 448

Fact 12.21.19, 797

Cauchy's estimate**polynomial root****bound**

Fact 11.20.6, 709

Cauchy-Schwarz**inequality****Buzano's inequality**

Fact 1.17.2, 67

Callebaut's**inequality**

Fact 1.16.16, 63

De Bruijn's**inequality**

Fact 1.16.20, 64

determinant

Fact 8.13.22, 489

Frobenius norm

Corollary 9.3.9, 553

inner product bound

Corollary 9.1.7, 546

McLaughlin's**inequality**

Fact 1.16.17, 64

Milne's inequality

Fact 1.16.15, 63

Ozeki's inequality

Fact 1.16.23, 65

Polya-Szego**inequality**

Fact 1.16.21, 64

**positive-semidefinite
matrix**

Fact 8.11.14, 470

Fact 8.11.15, 470

Fact 8.15.8, 501

vector case

Fact 1.16.9, 62

Cayley transform**asymptotically stable
matrix**

Fact 11.21.8, 713

cross product

Fact 3.11.8, 190

cross-product matrix

Fact 3.10.1, 186

definition

Fact 3.11.29, 197

discrete-time**asymptotically****stable matrix**

Fact 11.21.8, 713

Hamiltonian matrix

Fact 3.19.12, 217

Hermitian matrix

Fact 3.11.29, 197

orthogonal matrix

Fact 3.11.8, 190

Fact 3.11.28, 196

Fact 3.11.30, 197

Fact 3.11.31, 198

positive-definite**matrix**

Fact 8.9.30, 453

**skew-Hermitian
matrix**

Fact 3.11.28, 196

**skew-symmetric
matrix**

Fact 3.11.8, 190

Fact 3.11.28, 196

Fact 3.11.30, 197

Fact 3.11.31, 198

symplectic matrix

Fact 3.19.12, 217

unitary matrix

Fact 3.11.28, 196

**Cayley-Hamilton
theorem****characteristic****polynomial**

Theorem 4.4.7, 243

generalized version

Fact 4.9.7, 261

center subgroup**commutator**

Fact 2.18.10, 150

centralizer**commutator**

Fact 2.18.9, 150

Fact 7.5.2, 409

commuting matrices

Fact 5.14.22, 341

Fact 5.14.24, 342

centrohermitian matrix**complex conjugate****transpose**

Fact 3.20.16, 221

definition

Definition 3.1.2, 166

generalized inverse

Fact 6.3.31, 376

matrix product

Fact 3.20.17, 221

**centrosymmetric
matrix****definition**

Definition 3.1.2, 166

matrix product

Fact 3.20.17, 221

matrix transpose

Fact 3.20.16, 221

Cesaro summable

discrete-time

Lyapunov-stable matrix

Fact 11.21.11, 714

chaotic order

matrix logarithm

Fact 8.19.1, 522

Fact 8.19.2, 523

positive-semidefinite order

Fact 8.19.2, 523

characteristic equation

definition, 240

characteristic polynomial

2×2 matrix

Fact 4.9.1, 260

3×3 matrix

Fact 4.9.2, 260

adjugate

Fact 4.9.8, 261

Cayley-Hamilton theorem

Theorem 4.4.7, 243

companion matrix

Proposition 5.2.1, 284

Corollary 5.2.4, 286

Corollary 5.2.5, 286

cross-product matrix

Fact 4.9.19, 263

Fact 4.9.20, 263

cyclic matrix

Proposition 5.5.15, 299

definition

Definition 4.4.1, 239

degree

Proposition 4.4.3, 240

derivative

Lemma 4.4.8, 244

eigenvalue

Proposition 4.4.6, 242

generalized inverse

Fact 6.3.20, 374

Hamiltonian matrix

Fact 4.9.21, 264

Fact 4.9.23, 264

identities

Proposition 4.4.5, 241

inverse matrix

Fact 4.9.9, 261

Leverrier's algorithm

Proposition 4.4.9, 244

matrix product

Proposition 4.4.10, 244

Corollary 4.4.11, 245

minimal polynomial

Fact 4.9.24, 265

monic

Proposition 4.4.3, 240

outer-product

matrix

Fact 4.9.16, 262

Fact 4.9.18, 263

output feedback

Fact 12.22.13, 800

partitioned matrix

Fact 4.9.14, 262

Fact 4.9.15, 262

Fact 4.9.17, 263

Fact 4.9.18, 263

Fact 4.9.22, 264

Fact 4.9.23, 264

similar matrices

Fact 4.9.10, 262

similarity invariant

Proposition 4.4.2, 240

Proposition 4.6.2, 248

skew-Hermitian

matrix

Fact 4.9.13, 262

skew-symmetric

matrix

Fact 4.9.12, 262

Fact 4.9.19, 263

Fact 4.9.20, 263

Fact 5.14.34, 343

sum of derivatives

Fact 4.9.11, 262

upper

block-triangular

matrix

Fact 4.10.11, 267

Chebyshev's inequality rearrangement

Fact 1.16.3, 60

Chen form

tridiagonal matrix

Fact 11.18.27, 703

child

Definition 1.4.2, 8

Cholesky

decomposition

existence

Fact 8.9.37, 454

circle

complex numbers

Fact 2.20.12, 158

circulant matrix

block-circulant matrix

Fact 3.18.3, 215

companion matrix

Fact 5.16.7, 355

Fourier matrix

Fact 5.16.7, 355

group

Fact 3.21.7, 224

Fact 3.21.8, 224

permutation matrix

Fact 5.16.8, 357

primary circulant

Fact 5.16.7, 355

spectrum

Fact 5.16.7, 355

Clarkson inequalities

complex numbers

Fact 1.18.2, 69

Schatten norm

Fact 9.9.34, 586

CLHP

closed left half plane

definition, 77

Clifford algebra

real matrix

representation

Fact 3.22.1, 225

Cline

factorization

expression for the

- group generalized inverse
 - Fact 6.6.12, 395
- generalized inverse of a matrix product
 - Fact 6.4.10, 379
- closed half space**
 - affine closed half space
 - Fact 2.9.6, 111
 - definition, 91
- closed relative to a set**
 - continuous function
 - Theorem 10.3.4, 624
 - definition
 - Definition 10.1.4, 622
- closed set**
 - complement
 - Fact 10.8.4, 632
 - continuous function
 - Theorem 10.3.10, 625
 - Corollary 10.3.5, 624
 - Corollary 10.3.11, 625
 - definition
 - Definition 10.1.3, 621
 - image under linear mapping
 - Fact 10.9.8, 635
 - intersection
 - Fact 10.9.10, 635
 - Fact 10.9.11, 635
 - polar
 - Fact 2.9.4, 110
 - positive-semidefinite matrix
 - Fact 10.8.18, 633
 - subspace
 - Fact 10.8.21, 633
 - union
 - Fact 10.9.10, 635
- closed-loop spectrum**
 - detectability
 - Lemma 12.16.17, 781
 - Hamiltonian
 - Proposition 12.16.14, 781
- maximal solution of the Riccati equation
 - Proposition 12.18.2, 787
- observability
 - Lemma 12.16.17, 781
- observable eigenvalue
 - Lemma 12.16.16, 781
- Riccati equation**
 - Proposition 12.16.14, 781
 - Proposition 12.18.2, 787
 - Proposition 12.18.3, 787
 - Proposition 12.18.7, 789
- closure**
 - affine hull
 - Fact 10.8.11, 633
 - complement
 - Fact 10.8.6, 632
 - convex hull
 - Fact 10.8.13, 633
 - convex set
 - Fact 10.8.8, 632
 - Fact 10.8.19, 633
 - definition
 - Definition 10.1.3, 621
 - smallest closed set
 - Fact 10.8.3, 632
 - subset
 - Fact 10.9.1, 634
 - union
 - Fact 10.9.2, 634
- closure point**
 - definition
 - Definition 10.1.3, 621
- closure point relative to a set**
 - definition
 - Definition 10.1.4, 622
- closure relative to a set**
 - definition
 - Definition 10.1.4, 622
- codomain**
 - definition, 3
- cofactor**
 - definition, 105
 - determinant expansion
 - Proposition 2.7.5, 105
- cogredient**
 - diagonalization**
 - commuting matrices
 - Fact 8.16.1, 507
 - definition, 422
 - diagonalizable matrix
 - Fact 8.16.2, 507
 - Fact 8.16.3, 507
 - positive-definite matrix
 - Theorem 8.3.1, 423
 - Fact 8.16.5, 507
 - positive-semidefinite matrix
 - Theorem 8.3.4, 423
 - unitary matrix
 - Fact 8.16.1, 507
- cogredient**
 - diagonalization of positive-definite matrices**
 - Weierstrass
 - Fact 8.16.2, 507
- cogredient transformation**
 - Hermitian matrix
 - Fact 8.16.4, 507
 - Fact 8.16.6, 507
 - simultaneous diagonalization
 - Fact 8.16.4, 507
 - Fact 8.16.6, 507
 - simultaneous triangularization
 - Fact 5.17.9, 358
- Cohn**
 - polynomial root bounds
 - Fact 11.20.11, 711

colinear**determinant**

Fact 2.20.1, 154

Fact 2.20.5, 155

Fact 2.20.9, 156

colleague form

definition, 362

column

definition, 79

column norm

definition, 556

Hölder-induced**norm**

Fact 9.8.21, 575

Fact 9.8.23, 575

Kronecker product

Fact 9.9.61, 591

partitioned matrix

Fact 9.8.11, 572

row norm

Fact 9.8.10, 572

spectral radius

Corollary 9.4.10, 556

column vector

definition, 78

column-stackingoperator, *see* **vec****common divisor**

definition, 233

common eigenvector**commuting matrices**

Fact 5.14.28, 342

norm equality

Fact 9.9.33, 585

simultaneous**triangularization**

Fact 5.17.1, 358

subspace

Fact 5.14.27, 342

common multiple

definition, 234

commutant**commutator**

Fact 2.18.9, 150

Fact 7.5.2, 409

commutator **2×2 matrices**

Fact 2.18.1, 149

adjoint operator

Fact 2.18.5, 149

Fact 2.18.6, 150

center subgroup

Fact 2.18.10, 150

centralizer

Fact 2.18.9, 150

Fact 7.5.2, 409

convergent sequence

Fact 11.14.9, 683

definition, 82**derivative of a****matrix**

Fact 11.14.11, 683

determinant

Fact 2.18.7, 150

dimension

Fact 2.18.9, 150

Fact 2.18.10, 150

Fact 2.18.11, 150

Fact 7.5.2, 409

factorization

Fact 5.15.33, 351

Frobenius norm

Fact 9.9.26, 584

Fact 9.9.27, 584

Hermitian matrix

Fact 3.8.1, 184

Fact 3.8.3, 185

Fact 9.9.30, 585

idempotent matrix

Fact 3.12.16, 200

Fact 3.12.17, 200

Fact 3.12.30, 204

Fact 3.12.31, 204

Fact 3.12.32, 205

Fact 3.15.4, 200

identities

Fact 2.12.19, 127

Fact 2.18.4, 149

infinite product

Fact 11.14.18, 685

involutory matrix

Fact 3.15.4, 212

lower triangular**matrix**

Fact 3.17.11, 214

matrix exponential

Fact 11.14.9, 683

Fact 11.14.11, 683

Fact 11.14.12, 683

Fact 11.14.13, 684

Fact 11.14.14, 684

Fact 11.14.15, 684

Fact 11.14.16, 684

Fact 11.14.17, 684

Fact 11.14.18, 685

maximum eigenvalue

Fact 9.9.30, 585

Fact 9.9.31, 585

maximum singular**value**

Fact 9.9.29, 584

Fact 9.14.9, 609

nilpotent matrix

Fact 3.12.16, 200

Fact 3.17.11, 214

Fact 3.17.12, 214

Fact 3.17.13, 214

normal matrix

Fact 3.8.6, 185

Fact 3.8.7, 185

Fact 9.9.31, 585

power

Fact 2.18.2, 149

powers

Fact 2.18.3, 149

projector

Fact 3.13.23, 210

Fact 9.9.9, 581

rank

Fact 3.12.31, 204

Fact 3.13.23, 210

Fact 5.17.5, 358

Fact 6.3.9, 371

Schatten norm

Fact 9.9.27, 584

series

Fact 11.14.17, 684

simultaneous**triangularization**

Fact 5.17.5, 358

Fact 5.17.6, 358

skew-Hermitian**matrix**

- Fact 3.8.1, 184
- Fact 3.8.4, 185
- skew-symmetric matrix**
 - Fact 3.8.5, 185
- spectrum**
 - Fact 5.12.14, 335
- spread**
 - Fact 9.9.30, 585
 - Fact 9.9.31, 585
- submultiplicative norm**
 - Fact 9.9.8, 580
- subspace**
 - Fact 2.18.9, 150
 - Fact 2.18.10, 150
 - Fact 2.18.12, 151
- sum**
 - Fact 2.18.12, 151
- trace**
 - Fact 2.18.1, 149
 - Fact 2.18.2, 149
 - Fact 5.9.18, 313
- triangularization**
 - Fact 5.17.5, 358
- unitarily invariant norm**
 - Fact 9.9.29, 584
 - Fact 9.9.30, 585
 - Fact 9.9.31, 585
- upper triangular matrix**
 - Fact 3.17.11, 214
- zero diagonal**
 - Fact 3.8.2, 184
- zero trace**
 - Fact 2.18.11, 150
- commutator realization**
 - Shoda's theorem**
 - Fact 5.9.18, 313
- commuting matrices**
 - centralizer**
 - Fact 5.14.22, 341
 - Fact 5.14.24, 342
 - cogredient diagonalization**
 - Fact 8.16.1, 507
 - common eigenvector**
 - Fact 5.14.28, 342
 - cyclic matrix**
 - Fact 5.14.22, 341
 - diagonalizable matrix**
 - Fact 5.17.8, 358
 - dimension**
 - Fact 5.10.15, 319
 - Fact 5.10.16, 319
 - Drazin generalized inverse**
 - Fact 6.6.4, 394
 - Fact 6.6.5, 394
 - eigenvector**
 - Fact 5.14.25, 342
 - generalized Cayley-Hamilton theorem**
 - Fact 4.9.7, 261
 - Hermitian matrix**
 - Fact 5.14.29, 342
 - idempotent matrix**
 - Fact 3.16.5, 213
 - Kronecker sum**
 - Fact 7.5.4, 409
 - matrix exponential**
 - Proposition 11.1.5, 645
 - Corollary 11.1.6, 645
 - Fact 11.14.2, 681
 - Fact 11.14.5, 682
 - nilpotent matrix**
 - Fact 3.17.9, 214
 - Fact 3.17.10, 214
 - normal matrix**
 - Fact 3.7.28, 183
 - Fact 3.7.29, 183
 - Fact 5.14.29, 342
 - Fact 5.17.7, 358
 - Fact 11.14.5, 682
 - polynomial representation**
 - Fact 5.14.22, 341
 - Fact 5.14.23, 342
 - Fact 5.14.24, 342
 - positive-definite matrix**
 - Fact 8.9.40, 455
 - positive-semidefinite matrix**
 - Fact 8.19.5, 467, 523
 - projector**
 - Fact 6.4.33, 383
 - Fact 8.10.23, 458
 - Fact 8.10.25, 458
 - range-Hermitian matrix**
 - Fact 6.4.26, 382
 - Fact 6.4.27, 382
 - rank subtractivity**
 - partial ordering**
 - Fact 8.19.5, 523
 - simple matrix**
 - Fact 5.14.23, 342
 - simultaneous diagonalization**
 - Fact 8.16.1, 507
 - simultaneous triangularization**
 - Fact 5.17.4, 358
 - spectral radius**
 - Fact 5.12.11, 334
 - spectrum**
 - Fact 5.12.14, 335
 - square root**
 - Fact 5.18.1, 359
 - Fact 8.10.25, 458
 - star partial ordering**
 - Fact 2.10.36, 120
 - time-varying dynamics**
 - Fact 11.13.4, 678
 - triangularization**
 - Fact 5.17.4, 358
 - compact set**
 - continuous function**
 - Theorem 10.3.8, 624
 - convergent subsequence**
 - Theorem 10.2.5, 623
 - convex hull**
 - Fact 10.8.15, 633
 - definition, 622**
 - existence of minimizer**
 - Corollary 10.3.9, 624
 - companion form matrix**

- discrete-time semistable matrix
 - Fact 11.21.18, 715
- companion matrix**
 - block-diagonal matrix
 - Proposition 5.2.8, 286
 - Lemma 5.2.2, 285
 - bottom, right, top, left
 - Fact 5.16.1, 352
 - characteristic polynomial
 - Proposition 5.2.1, 284
 - Corollary 5.2.4, 286
 - Corollary 5.2.5, 286
 - circulant matrix
 - Fact 5.16.7, 355
 - cyclic matrix
 - Fact 5.16.5, 354
 - definition, 283
 - elementary divisor
 - Theorem 5.2.9, 287
 - example
 - Example 5.3.6, 290
 - Example 5.3.7, 291
 - hypercompanion matrix
 - Corollary 5.3.4, 289
 - Lemma 5.3.1, 288
 - inverse matrix
 - Fact 5.16.2, 353
 - minimal polynomial
 - Proposition 5.2.1, 284
 - Corollary 5.2.4, 286
 - Corollary 5.2.5, 286
 - nonnegative matrix
 - Fact 4.11.13, 279
 - oscillator
 - Fact 5.14.35, 344
 - similar matrices
 - Fact 5.16.5, 354
 - singular value
 - Fact 5.11.30, 327
 - Vandermonde matrix
 - Fact 5.16.4, 354
- compartmental matrix**
- asymptotically stable matrix
 - Fact 11.19.6, 707
- Lyapunov-stable matrix
 - Fact 11.19.6, 707
- semistable matrix
 - Fact 11.19.6, 707
- compatible norm**
 - induced norm
 - Proposition 9.4.3, 553
- compatible norms**
 - definition, 549
 - Hölder norm
 - Proposition 9.3.5, 550
 - Schatten norm
 - Proposition 9.3.6, 551
 - Corollary 9.3.7, 552
 - Corollary 9.3.8, 552
 - submultiplicative norm
 - Proposition 9.3.1, 550
 - trace norm
 - Corollary 9.3.8, 552
- complement**
 - closed set
 - Fact 10.8.4, 632
 - closure
 - Fact 10.8.6, 632
 - definition, 2
 - interior
 - Fact 10.8.6, 632
 - open set
 - Fact 10.8.4, 632
 - relatively closed set
 - Fact 10.8.5, 632
 - relatively open set
 - Fact 10.8.5, 632
- complement of a graph**
 - Definition 1.4.1, 8
- complement of a relation**
 - definition
 - Definition 1.3.4, 5
- complementary relation**
 - relation
 - Proposition 1.3.5, 6
- complementary submatrix**
 - defect
 - Fact 2.11.20, 125
- complementary subspaces**
 - complex conjugate transpose
 - Fact 3.12.1, 198
 - definition, 90
 - group-invertible matrix
 - Corollary 3.5.8, 176
 - idempotent matrix
 - Proposition 3.5.3, 176
 - Proposition 3.5.4, 176
 - Fact 3.12.1, 198
 - Fact 3.12.33, 205
 - index of a matrix
 - Proposition 3.5.7, 176
 - partitioned matrix
 - Fact 3.12.33, 205
 - projector
 - Fact 3.13.24, 210
 - simultaneous
 - Fact 2.9.23, 113
 - stable subspace
 - Proposition 11.8.8, 665
 - sum of dimensions
 - Corollary 2.3.2, 90
 - unstable subspace
 - Proposition 11.8.8, 665
- completely solid set**
 - convex set
 - Fact 10.8.9, 632
 - definition, 622
 - open ball
 - Fact 10.8.1, 632
 - positive-semidefinite matrix
 - Fact 10.8.18, 633
 - solid set
 - Fact 10.8.9, 632
- complex conjugate determinant**

- Fact 2.19.8, 153
- Fact 2.19.9, 153
- matrix exponential**
 - Proposition 11.2.8, 649
- partitioned matrix**
 - Fact 2.19.9, 153
- similar matrices**
 - Fact 5.9.31, 316
- complex conjugate of a matrix**
 - definition, 87
- complex conjugate of a vector**
 - definition, 85
- complex conjugate transpose**
- complementary subspaces**
 - Fact 3.12.1, 198
- definition, 87**
- determinant**
 - Fact 9.11.1, 596
- diagonalizable matrix**
 - Fact 5.14.5, 339
- factorization**
 - Fact 5.15.23, 349
- generalized inverse**
 - Fact 6.3.9, 371
 - Fact 6.3.10, 371
 - Fact 6.3.13, 372
 - Fact 6.3.16, 373
 - Fact 6.3.17, 373
 - Fact 6.3.18, 373
 - Fact 6.3.22, 374
 - Fact 6.3.27, 375
 - Fact 6.3.28, 375
 - Fact 6.4.7, 379
 - Fact 6.6.16, 396
 - Fact 6.6.17, 397
 - Fact 6.6.18, 397
- group generalized inverse**
 - Fact 6.6.10, 394
- Hermitian matrix**
 - Fact 3.7.13, 180
 - Fact 5.9.8, 312
 - Fact 6.6.18, 397
- idempotent matrix**
 - Fact 5.9.21, 314
- identity**
 - Fact 2.10.33, 119
 - Fact 2.10.34, 120
- Kronecker product**
 - Proposition 7.1.3, 400
- left inverse**
 - Fact 2.15.1, 140
 - Fact 2.15.2, 140
- matrix exponential**
 - Proposition 11.2.8, 649
 - Fact 11.15.4, 689
 - Fact 11.15.6, 690
- maximum singular value**
 - Fact 8.17.3, 508
 - Fact 8.18.11, 515
 - Fact 8.21.10, 533
- nonsingular matrix**
 - Fact 2.16.30, 146
- norm**
 - Fact 9.8.8, 572
- normal matrix**
 - Fact 5.14.30, 343
 - Fact 6.3.16, 373
 - Fact 6.3.17, 373
 - Fact 6.6.10, 394
 - Fact 6.6.17, 397
- partitioned matrix**
 - Fact 6.5.3, 386
- positive-definite matrix**
 - Fact 8.9.39, 455
- projector**
 - Fact 3.13.1, 206
- range**
 - Fact 6.5.3, 386
 - Fact 8.7.2, 443
- range-Hermitian matrix**
 - Fact 3.6.4, 178
 - Fact 6.3.10, 371
 - Fact 6.6.16, 396
- Schur product**
 - Fact 8.21.9, 533
- similarity transformation**
 - Fact 5.9.8, 312
- Fact 5.15.4, 345
- singular value**
 - Fact 5.11.34, 328
- subspace**
 - Fact 2.9.28, 114
- trace**
 - Fact 8.12.4, 476
 - Fact 8.12.5, 476
 - Fact 9.13.16, 604
- unitarily invariant norm**
 - Fact 9.8.30, 576
- unitarily left-equivalent matrices**
 - Fact 5.10.18, 319
 - Fact 5.10.19, 319
- unitarily right-equivalent matrices**
 - Fact 5.10.18, 319
- unitarily similar matrices**
 - Fact 5.9.20, 314
 - Fact 5.9.21, 314
- complex conjugate transpose of a vector**
 - definition, 85
- complex inequality Petrovich**
 - Fact 1.18.2, 69
- complex matrix**
 - block 2×2 representation**
 - Fact 2.19.3, 151
- complex conjugate**
 - Fact 2.19.4, 152
- determinant**
 - Fact 2.19.3, 151
 - Fact 2.19.10, 153
- partitioned matrix**
 - Fact 2.19.4, 152
 - Fact 2.19.5, 152
 - Fact 2.19.6, 152
 - Fact 2.19.7, 153
 - Fact 3.11.27, 196
- positive-definite matrix**

- Fact 3.7.9, 179
- positive-semidefinite matrix**
 - Fact 3.7.9, 179
- rank**
 - Fact 2.19.3, 151
- complex numbers**
 - 2×2 representation
 - Fact 2.19.1, 151
- circle**
 - Fact 2.20.12, 158
- Clarkson inequalities**
 - Fact 1.18.2, 69
- Dunkl-Williams inequality**
 - Fact 1.18.5, 71
- equilateral triangle**
 - Fact 2.20.6, 155
- exponential function**
 - Fact 1.18.6, 71
- identities**
 - Fact 1.18.1, 68
 - Fact 1.18.2, 69
- identity**
 - Fact 1.18.4, 71
- inequalities**
 - Fact 1.18.1, 68
 - Fact 1.18.2, 69
 - Fact 1.18.5, 71
- inequality**
 - Fact 1.12.4, 47
- infinite series**
 - Fact 1.18.8, 73
- logarithm function**
 - Fact 1.18.7, 72
- Maligranda inequality**
 - Fact 1.18.5, 71
- Massera-Schaffer inequality**
 - Fact 1.18.5, 71
- parallelogram law**
 - Fact 1.18.2, 69
- polarization identity**
 - Fact 1.18.2, 69
- quadratic formula**
 - Fact 1.18.3, 70
- trigonometric function**
 - Fact 1.19.3, 76
- complex symmetric Jordan form**
- similarity transformation**
 - Fact 5.15.2, 345
 - Fact 5.15.3, 345
- complex-symmetric matrix**
 - T-congruence**
 - Fact 5.9.22, 314
 - T-congruent diagonalization**
 - Fact 5.9.22, 314
 - unitary matrix**
 - Fact 5.9.22, 314
- component**
 - definition, 78
- composition**
 - associativity
 - Proposition 1.2.1, 3
 - definition, 3
- composition of functions**
 - one-to-one function**
 - Fact 1.5.16, 13
 - onto function**
 - Fact 1.5.16, 13
- compound matrix**
 - matrix product**
 - Fact 7.5.17, 411
- compound of a matrix product**
 - Binet-Cauchy theorem**
 - Fact 7.5.17, 411
- comrade form**
 - definition, 362
- concave function**
 - definition
 - Definition 8.6.14, 436
 - function composition
 - Lemma 8.6.16, 436
- nonincreasing function**
- condition number**
 - linear system solution**
 - Fact 9.9.64, 592
 - Fact 9.9.65, 592
 - Fact 9.9.66, 592
- cone**
 - blunt**
 - definition, 89
 - cone of image**
 - Fact 2.9.26, 113
 - constructive characterization**
 - Theorem 2.3.5, 91
 - definition, 89**
 - dictionary ordering**
 - Fact 2.9.31, 115
 - image under linear mapping**
 - Fact 2.9.26, 113
 - intersection**
 - Fact 2.9.9, 111
 - left inverse**
 - Fact 2.9.26, 113
 - lexicographic ordering**
 - Fact 2.9.31, 115
 - one-sided**
 - definition, 89
 - pointed**
 - definition, 89
 - quadratic form**
 - Fact 8.14.11, 498
 - Fact 8.14.13, 498
 - Fact 8.14.14, 498
 - sum**
 - Fact 2.9.9, 111
 - variational**
 - definition, 625
- confederate form**
 - definition, 362
- congenial matrix**
 - definition, 362
- congruence**
 - equivalence relation**
 - Fact 5.10.3, 317

- generalized inverse
 - Fact 8.20.5, 525
- congruence transformation**
 - normal matrix
 - Fact 5.10.17, 319
- congruent matrices**
 - biequivalent matrices
 - Proposition 3.4.5, 174
 - definition
 - Definition 3.4.4, 174
 - Hermitian matrix
 - Proposition 3.4.5, 174
 - Corollary 5.4.7, 294
 - inertia
 - Corollary 5.4.7, 294
 - Fact 5.8.22, 311
 - Kronecker product
 - Fact 7.4.12, 406
 - matrix classes
 - Proposition 3.4.5, 174
 - positive-definite matrix
 - Proposition 3.4.5, 174
 - Corollary 8.1.3, 419
 - positive-semidefinite matrix
 - Proposition 3.4.5, 174
 - Corollary 8.1.3, 419
 - range-Hermitian matrix
 - Proposition 3.4.5, 174
 - Fact 5.9.6, 312
 - skew-Hermitian matrix
 - Proposition 3.4.5, 174
 - skew-symmetric matrix
 - Fact 3.7.34, 184
 - Fact 5.9.16, 313
 - Sylvester's law of inertia, 294
 - symmetric matrix
 - Fact 5.9.16, 313
 - unit imaginary matrix
 - Fact 3.7.34, 184
- conical hull**
- definition, 89
- conjugate parameters**
 - barycentric coordinates
 - Fact 1.16.11, 62
- connected graph**
 - Definition 1.4.3, 9
 - irreducible matrix
 - Fact 4.11.2, 273
 - walk
 - Fact 4.11.4, 273
- constant polynomial**
 - definition, 231
- contained**
 - definition, 2
- continuity**
 - spectrum
 - Fact 10.11.8, 638
 - Fact 10.11.9, 639
- continuity of roots**
 - coefficients
 - polynomial
 - Fact 10.11.2, 638
- continuous function**
 - bounded set
 - Theorem 10.3.10, 625
 - Corollary 10.3.11, 625
 - closed relative to a set
 - Theorem 10.3.4, 624
 - closed set
 - Theorem 10.3.10, 625
 - Corollary 10.3.5, 624
 - Corollary 10.3.11, 625
 - compact set
 - Theorem 10.3.8, 624
 - convex function
 - Theorem 10.3.2, 624
 - Fact 10.11.12, 639
 - convex set
 - Theorem 10.3.10, 625
 - Corollary 10.3.11, 625
 - definition
 - Definition 10.3.1, 623
 - differentiable function
 - Proposition 10.4.4, 626
 - existence of minimizer
 - Corollary 10.3.9, 624
 - fixed-point theorem
 - Theorem 10.3.10, 625
 - Corollary 10.3.11, 625
 - linear function
 - Corollary 10.3.3, 624
 - maximization
 - Fact 10.11.4, 638
 - open relative to a set
 - Theorem 10.3.4, 624
 - open set
 - Corollary 10.3.5, 624
 - open set image
 - Theorem 10.3.7, 624
 - pathwise-connected set
 - Fact 10.11.5, 638
- continuous-time control problem**
 - LQG controller
 - Fact 12.23.6, 804
- continuously differentiable function**
 - definition, 627
- contractive matrix**
 - complex conjugate transpose
 - Fact 3.20.12, 220
 - definition
 - Definition 3.1.2, 166
 - partitioned matrix
 - Fact 8.11.24, 473
 - positive-definite matrix
 - Fact 8.11.13, 470
- contradiction**
 - definition, 1
- contragredient diagonalization**
 - definition, 422
 - positive-definite matrix
 - Theorem 8.3.2, 423

Corollary 8.3.3, 423
positive-semidefinite matrix
 Theorem 8.3.5, 424
 Corollary 8.3.7, 424

contrapositive
 definition, 1

controllability
 asymptotically stable matrix
 Fact 12.20.5, 791
 cyclic matrix
 Fact 12.20.13, 792
 diagonal matrix
 Fact 12.20.12, 792
 final state
 Fact 12.20.4, 791
 geometric multiplicity
 Fact 12.20.14, 792
Gramian
 Fact 12.20.17, 792
 input matrix
 Fact 12.20.15, 792
 positive-semidefinite matrix
 Fact 12.20.6, 791
 positive-semidefinite ordering
 Fact 12.20.8, 791
 range
 Fact 12.20.7, 791
 shift
 Fact 12.20.10, 792
 shifted dynamics
 Fact 12.20.9, 791
 skew-symmetric matrix
 Fact 12.20.5, 791
 stabilization
 Fact 12.20.17, 792
Sylvester's equation
 Fact 12.21.14, 796
 transpose
 Fact 12.20.16, 792
controllability Gramian
 asymptotically stable matrix

Proposition 12.7.9, 747
 Corollary 12.7.10, 747
controllably asymptotically stable
 Proposition 12.7.3, 743
 Proposition 12.7.4, 745
 Proposition 12.7.5, 746
 Proposition 12.7.6, 746
 Proposition 12.7.7, 746
frequency domain
 Corollary 12.11.5, 767
 H_2 norm
 Corollary 12.11.4, 767
 Corollary 12.11.5, 767
 L_2 norm
 Theorem 12.11.1, 765
controllability matrix
 controllable pair
 Theorem 12.6.18, 742
definition, 737
 rank
 Corollary 12.6.3, 737
Sylvester's equation
 Fact 12.21.13, 796
controllability pencil
 definition
 Definition 12.6.12, 740
Smith form
 Proposition 12.6.15, 741
Smith zeros
 Proposition 12.6.16, 741
uncontrollable eigenvalue
 Proposition 12.6.13, 740
uncontrollable spectrum
 Proposition 12.6.16, 741
controllable canonical form
 definition, 750
 equivalent realizations
 Corollary 12.9.9, 752

realization
 Proposition 12.9.3, 750
controllable dynamics
 block-triangular matrix
 Theorem 12.6.8, 739
orthogonal matrix
 Theorem 12.6.8, 739
controllable eigenvalue
 controllable subspace
 Proposition 12.6.17, 742
controllable pair
 asymptotically stable matrix
 Proposition 12.7.9, 747
 Corollary 12.7.10, 747
controllability matrix
 Theorem 12.6.18, 742
cyclic matrix
 Fact 5.14.9, 340
eigenvalue placement
 Proposition 12.6.19, 743
equivalent realizations
 Proposition 12.9.8, 752
Markov
block-Hankel matrix
 Proposition 12.9.11, 754
minimal realization
 Proposition 12.9.10, 753
 Corollary 12.9.15, 756
positive-definite matrix
 Theorem 12.6.18, 742
rank
 Fact 5.14.10, 340
controllable subspace
 block-triangular matrix
 Proposition 12.6.9, 739

- Proposition 12.6.10, 740
- controllable eigenvalue**
 - Proposition 12.6.17, 742
- definition**
 - Definition 12.6.1, 735
- equivalent expressions**
 - Lemma 12.6.2, 736
- final state**
 - Fact 12.20.3, 791
- full-state feedback**
 - Proposition 12.6.5, 737
- identity shift**
 - Lemma 12.6.7, 738
- invariant subspace**
 - Corollary 12.6.4, 737
- nonsingular matrix**
 - Proposition 12.6.10, 740
- orthogonal matrix**
 - Proposition 12.6.9, 739
- projector**
 - Lemma 12.6.6, 738
- controllably asymptotically stable**
- asymptotically stable matrix**
 - Proposition 12.8.3, 747
 - Proposition 12.8.5, 748
- block-triangular matrix**
 - Proposition 12.7.3, 743
- controllability Gramian**
 - Proposition 12.7.3, 743
 - Proposition 12.7.4, 745
 - Proposition 12.7.5, 746
 - Proposition 12.7.6, 746
 - Proposition 12.7.7, 746
- definition**
 - Definition 12.7.1, 743
- full-state feedback**
 - Proposition 12.7.2, 743
- Lyapunov equation**
 - Proposition 12.7.3, 743
- orthogonal matrix**
 - Proposition 12.7.3, 743
- rank**
 - Proposition 12.7.4, 745
 - Proposition 12.7.5, 746
- stabilizability**
 - Proposition 12.8.3, 747
 - Proposition 12.8.5, 748
- convergent sequence**
- absolutely convergent sequence**
 - Proposition 10.2.7, 623
 - Proposition 10.2.9, 623
- closure point**
 - Proposition 10.2.4, 623
- commutator**
 - Fact 11.14.9, 683
- discrete-time semistable matrix**
 - Fact 11.21.14, 714
- generalized inverse**
 - Fact 6.3.35, 376
 - Fact 6.3.36, 377
- Hermitian matrix**
 - Fact 11.14.7, 683
 - Fact 11.14.8, 683
- inverse matrix**
 - Fact 2.16.29, 146
 - Fact 4.10.5, 266
- matrix exponential**
 - Proposition 11.1.3, 644
 - Fact 11.14.7, 683
 - Fact 11.14.8, 683
 - Fact 11.14.9, 683
 - Fact 11.21.14, 714
- matrix sign function**
 - Fact 5.15.21, 348
- spectral radius**
 - Fact 4.10.5, 266
 - Fact 9.8.4, 572
- square root**
 - Fact 5.15.21, 348
 - Fact 8.9.32, 454
- unitary matrix**
 - Fact 8.9.33, 454
- vectors**
 - Fact 10.11.1, 638
- convergent sequence of matrices**
- definition**
 - Definition 10.2.3, 622
- convergent sequence of scalars**
- definition**
 - Definition 10.2.2, 622
- convergent sequence of vectors**
- definition**
 - Definition 10.2.3, 622
- convergent series**
- definition**
 - Definition 10.2.6, 623
 - Definition 10.2.8, 623
- matrix exponential**
 - Proposition 11.1.2, 644
- convergent subsequence**
- compact set**
 - Theorem 10.2.5, 623
- converse**
- definition, 1**
- convex combination**
- definition, 89**
- determinant**
 - Fact 8.13.16, 488
- norm inequality**
 - Fact 9.7.15, 568
- positive-semidefinite matrix**
 - Fact 5.19.6, 360
 - Fact 8.13.16, 488
- convex cone**
- definition, 89**
- induced by transitive relation**
 - Proposition 2.3.6, 93
- inner product**
 - Fact 10.9.13, 635
- intersection**
 - Fact 2.9.9, 111
- polar**
 - Fact 2.9.4, 110

- positive-semidefinite matrix**, 417
- quadratic form**
 - Fact 8.14.11, 498
 - Fact 8.14.13, 498
 - Fact 8.14.14, 498
- separation theorem**
 - Fact 10.9.13, 635
- sum**
 - Fact 2.9.9, 111
- union**
 - Fact 2.9.10, 111
- convex conical hull**
 - constructive characterization**
 - Theorem 2.3.5, 91
 - convex hull**
 - Fact 2.9.3, 110
 - definition**, 89
 - dual cone**
 - Fact 2.9.3, 110
- convex function**
 - constant function**
 - Fact 1.8.3, 21
 - continuous function**
 - Theorem 10.3.2, 624
 - Fact 10.11.12, 639
 - convex set**
 - Fact 10.11.11, 639
 - Fact 10.11.12, 639
 - Fact 10.11.13, 639
 - definition**
 - Definition 1.2.3, 5
 - Definition 8.6.14, 436
 - derivative**
 - Fact 10.11.14, 639
 - determinant**
 - Proposition 8.6.17, 437
 - Fact 2.13.17, 132
 - directional differential**
 - Fact 10.11.14, 639
 - eigenvalue**
 - Corollary 8.6.19, 442
 - Fact 8.18.5, 513
 - function composition**
 - Lemma 8.6.16, 436
 - Hermite-Hadamard inequality**
 - Fact 1.8.6, 22
 - Hermitian matrix**
 - Fact 8.12.32, 482
 - Fact 8.12.33, 482
 - increasing function**
 - Theorem 8.6.15, 436
 - Jensen**
 - Fact 10.11.7, 638
 - Jensen's inequality**
 - Fact 1.8.4, 21
 - Fact 1.15.35, 57
 - Kronecker product**
 - Proposition 8.6.17, 437
 - log majorization**
 - Fact 2.21.12, 163
 - logarithm**
 - Fact 11.16.14, 695
 - Fact 11.16.15, 695
 - logarithm of determinant**
 - Proposition 8.6.17, 437
 - logarithm of trace**
 - Proposition 8.6.17, 437
 - matrix exponential**
 - Fact 8.14.18, 500
 - Fact 11.16.14, 695
 - Fact 11.16.15, 695
 - matrix functions**
 - Proposition 8.6.17, 437
 - matrix logarithm**
 - Proposition 8.6.17, 437
 - midpoint convex**
 - Fact 10.11.7, 638
 - minimizer**
 - Fact 8.14.15, 499
 - Niculescu's inequality**
 - Fact 1.8.5, 22
 - nondecreasing function**
 - Lemma 8.6.16, 436
 - one-sided directional differential**
 - Proposition 10.4.1, 626
 - Popoviciu's inequality**
 - Fact 1.8.6, 22
 - positive-definite matrix**
 - Fact 8.14.17, 499
 - positive-semidefinite matrix**
 - Fact 8.14.15, 499
 - Fact 8.20.20, 530
 - reverse inequality**
 - Fact 8.10.9, 457
 - scalar inequality**
 - Fact 1.8.1, 21
 - Schur complement**
 - Proposition 8.6.17, 437
 - Lemma 8.6.16, 436
 - singular value**
 - Fact 11.16.14, 695
 - Fact 11.16.15, 695
 - strong log majorization**
 - Fact 2.21.9, 163
 - strong majorization**
 - Fact 2.21.8, 163
 - Fact 2.21.11, 163
 - subdifferential**
 - Fact 10.11.14, 639
 - trace**
 - Proposition 8.6.17, 437
 - Fact 8.14.17, 499
 - transformation**
 - Fact 1.8.2, 21
 - weak majorization**
 - Fact 2.21.8, 163
 - Fact 2.21.9, 163
 - Fact 2.21.10, 163
 - Fact 2.21.11, 163
 - Fact 8.18.5, 513
 - convex hull**
 - affine hull**
 - Fact 2.9.3, 110
 - closure**
 - Fact 10.8.13, 633
 - compact set**
 - Fact 10.8.15, 633
 - constructive characterization**
 - Theorem 2.3.5, 91
 - definition**, 89
 - Hermitian matrix diagonal**

- Fact 8.17.8, 510
- open set**
 - Fact 10.8.14, 633
- simplex**
 - Fact 2.20.4, 154
- solid set**
 - Fact 10.8.10, 632
- spectrum**
 - Fact 8.14.7, 496
 - Fact 8.14.8, 497
- strong majorization**
 - Fact 2.21.7, 163
- convex polyhedron**
 - volume**
 - Fact 2.20.20, 160
- convex set**
 - affine hull**
 - Theorem 10.3.2, 624
 - Fact 10.8.8, 632
 - closure**
 - Fact 10.8.8, 632
 - Fact 10.8.19, 633
 - completely solid set**
 - Fact 10.8.9, 632
 - continuous function**
 - Theorem 10.3.10, 625
 - Corollary 10.3.11, 625
 - convexity of image**
 - Fact 2.9.26, 113
 - definition, 89**
 - distance from a point**
 - Fact 10.9.15, 636
 - Fact 10.9.16, 636
 - extreme point**
 - Fact 10.8.23, 634
 - image under linear mapping**
 - Fact 2.9.26, 113
 - interior**
 - Fact 10.8.8, 632
 - Fact 10.8.19, 633
 - intersection**
 - Fact 2.9.9, 111
 - Fact 10.9.6, 634
 - left inverse**
 - Fact 2.9.26, 113
 - norm**
 - Fact 9.7.23, 570
 - open ball**
 - Fact 10.8.1, 632
 - positive-semidefinite matrix**
 - Fact 8.14.2, 494
 - Fact 8.14.3, 495
 - Fact 8.14.4, 495
 - Fact 8.14.5, 495
 - Fact 8.14.6, 496
 - quadratic form**
 - Fact 8.14.2, 494
 - Fact 8.14.3, 495
 - Fact 8.14.4, 495
 - Fact 8.14.5, 495
 - Fact 8.14.6, 496
 - Fact 8.14.9, 497
 - Fact 8.14.11, 498
 - Fact 8.14.12, 498
 - Fact 8.14.13, 498
 - Fact 8.14.14, 498
 - set cancellation**
 - Fact 10.9.7, 635
 - solid set**
 - Fact 10.8.9, 632
 - sublevel set**
 - Fact 8.14.1, 494
 - sum**
 - Fact 2.9.9, 111
 - sum of sets**
 - Fact 2.9.1, 110
 - Fact 2.9.2, 110
 - Fact 10.9.4, 634
 - Fact 10.9.5, 634
 - Fact 10.9.7, 635
 - union**
 - Fact 10.9.7, 634
 - convex sets**
 - proper separation theorem**
 - Fact 10.9.14, 635
 - coplanar**
 - determinant**
 - Fact 2.20.2, 154
 - copositive matrix**
 - nonnegative matrix**
 - Fact 8.15.37, 507
 - positive-semidefinite matrix**
 - Fact 8.15.37, 507
 - coprime**
 - polynomial**
 - Fact 4.8.3, 254
 - Fact 4.8.4, 254
 - coprime polynomials**
 - Bezout matrix**
 - Fact 4.8.6, 255
 - Fact 4.8.7, 257
 - Fact 4.8.8, 257
 - definition, 233**
 - resultant**
 - Fact 4.8.4, 254
 - Smith-McMillan form**
 - Fact 4.8.15, 259
 - Sylvester matrix**
 - Fact 4.8.4, 254
 - coprime right polynomial fraction description**
 - Smith-McMillan form**
 - Proposition 4.7.16, 253
 - unimodular matrix**
 - Proposition 4.7.15, 253
 - Copson inequality**
 - sum of powers**
 - Fact 1.15.43, 59
 - Cordes inequality**
 - maximum singular value**
 - Fact 8.18.26, 520
 - corollary**
 - definition, 1**
 - cosine law**
 - vector identity**
 - Fact 9.7.4, 563
 - cosine rule**
 - triangle**
 - Fact 2.20.11, 156
 - Crabtree**

- Schur complement of a partitioned matrix
 - Fact 6.5.29, 393
- Crabtree-Haynsworth quotient formula**
 - Schur complement of a partitioned matrix
 - Fact 6.5.29, 393
- Cramer's rule**
 - linear system solution
 - Fact 2.13.6, 129
- creation matrix**
 - upper triangular matrix
 - Fact 11.11.4, 672
- CRHP**
 - closed right half plane
 - definition, 77
- Crimmins**
 - product of projectors
 - Fact 6.3.32, 376
- cross product**
 - adjugate
 - Fact 6.5.16, 389
 - Cayley transform
 - Fact 3.11.8, 190
 - identities
 - Fact 3.10.1, 186
 - matrix exponential
 - Fact 11.11.7, 673
 - Fact 11.11.8, 674
 - Fact 11.11.9, 674
 - orthogonal matrix
 - Fact 3.10.2, 189
 - Fact 3.10.3, 189
 - Fact 3.11.8, 190
 - outer-product matrix
 - Fact 3.11.8, 190
 - parallelogram
 - Fact 9.7.5, 565
- cross-product matrix**
- Cayley transform
 - Fact 3.10.1, 186
- characteristic polynomial
 - Fact 4.9.19, 263
 - Fact 4.9.20, 263
- identities
 - Fact 3.10.1, 186
- matrix exponential
 - Fact 11.11.6, 673
 - Fact 11.11.12, 674
 - Fact 11.11.13, 675
 - Fact 11.11.16, 676
 - Fact 11.11.17, 676
- orthogonal matrix
 - Fact 11.11.12, 674
 - Fact 11.11.13, 675
- spectrum
 - Fact 4.9.19, 263
- CS decomposition**
 - unitary matrix
 - Fact 5.9.29, 316
- cube root**
 - identity
 - Fact 2.12.23, 128
- cubes**
 - identity
 - Fact 2.12.24, 128
- cubic**
 - scalar inequality
 - Fact 1.11.14, 43
 - Fact 1.11.15, 43
 - Fact 1.11.16, 43
- cubic polynomial**
 - Cardano's trigonometric solution
 - Fact 4.10.1, 265
- CUD**
 - closed unit disk
 - definition, 670
- cycle**
 - definition
 - Definition 1.4.3, 9
 - graph
 - Fact 1.6.4, 13
- symmetric graph
 - Fact 1.6.5, 14
- cyclic eigenvalue**
 - definition
 - Definition 5.5.4, 296
- eigenvector
 - Fact 5.14.2, 338
- semisimple eigenvalue
 - Proposition 5.5.5, 296
- simple eigenvalue
 - Proposition 5.5.5, 296
- cyclic group**
 - group
 - Fact 3.21.7, 223
- cyclic inequality**
 - scalar inequality
 - Fact 1.11.11, 42
- cyclic matrix**
 - asymptotically stable matrix
 - Fact 11.18.25, 702
 - companion matrix
 - Fact 5.16.5, 354
 - characteristic polynomial
 - Proposition 5.5.15, 299
 - commuting matrices
 - Fact 5.14.22, 341
 - controllability
 - Fact 12.20.13, 792
 - controllable pair
 - Fact 5.14.9, 340
 - definition
 - Definition 5.5.4, 296
 - determinant
 - Fact 5.14.9, 340
 - identity perturbation
 - Fact 5.14.16, 341
 - linear independence
 - Fact 5.14.9, 340
 - matrix power
 - Fact 5.14.9, 340
 - minimal polynomial
 - Proposition 5.5.15, 299
 - nonsingular matrix
 - Fact 5.14.9, 340

rank

Fact 5.11.1, 321

semisimple matrix

Fact 5.14.11, 340

similar matrices

Fact 5.16.5, 354

simple matrix

Fact 5.14.11, 340

tridiagonal matrix

Fact 11.18.25, 702

D**damped natural****frequency**

definition, 654

Fact 5.14.35, 344

damping

definition, 654

damping matrix

partitioned matrix

Fact 5.12.21, 337

damping ratio

definition, 654

Fact 5.14.35, 344

Davenport

orthogonal matrices

and matrix

exponentials

Fact 11.11.13, 675

De Bruijn's inequality

refined

Cauchy-Schwarz

inequality

Fact 1.16.20, 64

De Morgan's laws

logical equivalents

Fact 1.5.1, 10

Decell

generalized inverse

Fact 6.4.31, 382

decreasing function

definition

Definition 8.6.12, 434

defect

adjugate

Fact 2.16.7, 143

definition, 96

geometric

multiplicity

Proposition 4.5.2, 246

group-invertible

matrix

Fact 3.6.1, 177

Hermitian matrix

Fact 5.8.7, 308

Fact 8.9.7, 451

identity

Fact 2.10.20, 117

identity involving

defect

Corollary 2.5.5, 97

identity with powers

Proposition 2.5.8, 97

identity with

transpose

Corollary 2.5.3, 96

Kronecker sum

Fact 7.5.2, 409

partitioned matrix

Fact 2.11.3, 121

Fact 2.11.8, 122

Fact 2.11.11, 123

product

Proposition 2.6.3, 99

product of matrices

Fact 2.10.14, 116

semisimple

eigenvalue

Proposition 5.5.8, 296

submatrix

Fact 2.11.20, 125

Sylvester's law of

nullity

Fact 2.10.15, 117

defective eigenvalue

definition

Definition 5.5.4, 296

defective matrix

definition

Definition 5.5.4, 296

identity perturbation

Fact 5.14.16, 341

nilpotent matrix

Fact 5.14.18, 341

outer-product

matrix

Fact 5.14.3, 338

deflating subspace

pencil

Fact 5.13.1, 338

degree

graph

Definition 1.4.3, 9

degree matrix

definition

Definition 3.2.1, 170

symmetric graph

Fact 4.11.1, 272

degree of a polynomial

definition, 231

degree of a polynomial

matrix

definition, 234

derivative

adjugate

Fact 10.11.19, 640

Fact 10.11.21, 641

convex function

Fact 10.11.14, 639

determinant

Proposition 10.7.3, 631

Fact 10.11.19, 640

Fact 10.11.21, 641

Fact 10.11.22, 641

Fact 10.11.23, 641

inverse matrix

Proposition 10.7.2, 630

Fact 10.11.18, 640

Fact 10.11.19, 641

logarithm of

determinant

Proposition 10.7.3, 631

matrix

definition, 628

matrix exponential

Fact 8.12.31, 482

Fact 11.14.3, 682

- Fact 11.14.4, 682
- Fact 11.14.10, 683
- Fact 11.15.2, 689
- matrix power**
 - Proposition 10.7.2, 630
- maximum singular value**
 - Fact 11.15.2, 689
- realization**
 - Fact 12.22.6, 799
- squared matrix**
 - Fact 10.11.17, 640
- trace**
 - Proposition 10.7.4, 631
 - Fact 11.14.3, 682
- transfer function**
 - Fact 12.22.6, 799
- derivative of a matrix**
- commutator**
 - Fact 11.14.11, 683
- matrix exponential**
 - Fact 11.14.11, 683
- matrix product**
 - Fact 11.13.8, 679
- derivative of a matrix exponential**
 - Bessis-Moussa-Villani trace conjecture
 - Fact 8.12.31, 482
- derivative of an integral**
- Leibniz's rule**
 - Fact 10.11.10, 639
- derogatory eigenvalue**
- definition**
 - Definition 5.5.4, 296
- derogatory matrix**
- definition**
 - Definition 5.5.4, 296
- identity perturbation**
 - Fact 5.14.16, 341
- Descartes rule of signs**
- polynomial**
 - Fact 11.17.1, 695
- detectability**
- asymptotically stable**
- matrix**
 - Proposition 12.5.5, 735
 - Corollary 12.5.6, 735
- block-triangular matrix**
 - Proposition 12.5.4, 735
- closed-loop spectrum**
 - Lemma 12.16.17, 781
- definition**
 - Definition 12.5.1, 734
- Lyapunov equation**
 - Corollary 12.5.6, 735
- observably**
- asymptotically stable**
 - Proposition 12.5.5, 735
- orthogonal matrix**
 - Proposition 12.5.4, 735
- output convergence**
 - Fact 12.20.2, 791
- output injection**
 - Proposition 12.5.2, 734
- Riccati equation**
 - Corollary 12.17.3, 783
 - Corollary 12.19.2, 790
- state convergence**
 - Fact 12.20.2, 791
- determinant**
- (1)-inverse**
 - Fact 6.5.28, 393
- adjugate**
 - Fact 2.14.27, 139
 - Fact 2.16.3, 141
 - Fact 2.16.5, 142
- affine hyperplane**
 - Fact 2.20.3, 154
- basic properties**
 - Proposition 2.7.2, 103
- Binet-Cauchy**
- formula**
 - Fact 2.13.4, 129
- block-triangular matrix**
 - Fact 2.14.8, 134
- Cartesian**
- decomposition**
 - Fact 8.13.4, 485
 - Fact 8.13.11, 486
- Cauchy matrix**
 - Fact 3.20.14, 220
 - Fact 3.20.15, 221
- Cauchy-Schwarz inequality**
 - Fact 8.13.22, 489
- cofactor expansion**
 - Proposition 2.7.5, 105
- colinear**
 - Fact 2.20.1, 154
 - Fact 2.20.5, 155
 - Fact 2.20.9, 156
- column interchange**
 - Proposition 2.7.2, 103
- commutator**
 - Fact 2.18.7, 150
- complex conjugate**
 - Fact 2.19.8, 153
 - Fact 2.19.9, 153
- complex conjugate transpose**
 - Proposition 2.7.1, 103
 - Fact 9.11.1, 596
- complex matrix**
 - Fact 2.19.3, 151
 - Fact 2.19.10, 153
- convex combination**
 - Fact 8.13.16, 488
- convex function**
 - Proposition 8.6.17, 437
 - Fact 2.13.17, 132
- coplanar**
 - Fact 2.20.2, 154
- cyclic matrix**
 - Fact 5.14.9, 340
- definition, 103**
- derivative**
 - Proposition 10.7.3, 631
 - Fact 10.11.19, 640
 - Fact 10.11.21, 641
 - Fact 10.11.22, 641
 - Fact 10.11.23, 641
- dissipative matrix**
 - Fact 8.13.2, 485
 - Fact 8.13.11, 486, 487
 - Fact 8.13.31, 491
- eigenvalue**
 - Fact 5.11.28, 326
 - Fact 5.11.29, 327

- Fact 8.13.1, 485
- elementary matrix**
 - Fact 2.16.1, 141
- factorization**
 - Fact 5.15.7, 346
 - Fact 5.15.34, 351
- Fibonacci numbers**
 - Fact 4.11.12, 277
- Frobenius norm**
 - Fact 9.8.39, 578
- full-state feedback**
 - Fact 12.22.14, 800
- generalized inverse**
 - Fact 6.5.26, 392
 - Fact 6.5.27, 392
 - Fact 6.5.28, 393
- geometric mean**
 - Fact 8.10.43, 461
- group**
 - Proposition 3.3.6, 172
- Hadamard's inequality**
 - Fact 8.13.33, 491
 - Fact 8.13.34, 491
- Hankel matrix**
 - Fact 3.18.4, 215
- Hermitian matrix**
 - Corollary 8.4.10, 427
 - Fact 3.7.21, 182
 - Fact 8.13.7, 486
- Hua's inequalities**
 - Fact 8.13.25, 489
- identity**
 - Fact 2.13.10, 130
 - Fact 2.13.11, 130
 - Fact 2.13.12, 130
 - Fact 2.13.13, 131
- induced norm**
 - Fact 9.12.11, 601
- inequality**
 - Fact 8.13.24, 489
 - Fact 8.13.25, 489
 - Fact 8.13.26, 490
 - Fact 8.13.27, 490
 - Fact 8.13.28, 490
 - Fact 8.13.30, 490
 - Fact 8.21.19, 534
- integral**
 - Fact 11.13.15, 680
- invariant zero**
 - Fact 12.22.14, 800
- inverse**
 - Fact 2.13.5, 129
- inverse function theorem**
 - Theorem 10.4.5, 627
- involutory matrix**
 - Fact 3.15.1, 212
 - Fact 5.15.32, 351
- Kronecker product**
 - Proposition 7.1.11, 402
- Kronecker sum**
 - Fact 7.5.11, 410
- linear combination**
 - Fact 8.13.18, 488
- lower**
 - block-triangular matrix**
 - Proposition 2.7.1, 103
 - reverse-triangular matrix**
 - Fact 2.13.8, 130
- matrix exponential**
 - Proposition 11.4.6, 655
 - Corollary 11.2.4, 648
 - Corollary 11.2.5, 648
 - Fact 11.13.15, 680
 - Fact 11.15.5, 689
- matrix logarithm**
 - Fact 8.13.8, 486
 - Fact 8.18.30, 521
 - Fact 9.8.39, 578
 - Fact 11.14.24, 686
- maximum singular value**
 - Fact 9.14.17, 611
 - Fact 9.14.18, 611
- minimum singular value**
 - Fact 9.14.18, 611
- nilpotent matrix**
 - Fact 3.17.9, 214
- nonsingular matrix**
 - Corollary 2.7.4, 104
 - Lemma 2.8.6, 108
- normal matrix**
 - Fact 5.12.12, 335
- ones matrix**
 - Fact 2.13.2, 129
- ones matrix perturbation**
 - Fact 2.16.6, 142
- orthogonal matrix**
 - Fact 3.11.21, 196
 - Fact 3.11.22, 196
- Ostrowski-Taussky inequality**
 - Fact 8.13.2, 485
- outer-product perturbation**
 - Fact 2.16.3, 141
- output feedback**
 - Fact 12.22.13, 800
- partitioned matrix**
 - Corollary 2.8.5, 107
 - Lemma 8.2.6, 421
 - Fact 2.14.2, 133
 - Fact 2.14.3, 133
 - Fact 2.14.4, 133
 - Fact 2.14.5, 134
 - Fact 2.14.6, 134
 - Fact 2.14.7, 134
 - Fact 2.14.9, 134
 - Fact 2.14.10, 135
 - Fact 2.14.11, 135
 - Fact 2.14.13, 135
 - Fact 2.14.14, 136
 - Fact 2.14.15, 136
 - Fact 2.14.16, 136
 - Fact 2.14.17, 136
 - Fact 2.14.18, 137
 - Fact 2.14.19, 137
 - Fact 2.14.20, 137
 - Fact 2.14.21, 137
 - Fact 2.14.22, 138
 - Fact 2.14.23, 138
 - Fact 2.14.24, 138
 - Fact 2.14.25, 138
 - Fact 2.14.26, 139
 - Fact 2.14.28, 139
 - Fact 2.17.5, 147
 - Fact 2.19.3, 151
 - Fact 2.19.9, 153
 - Fact 5.12.21, 337

- Fact 6.5.28, 393
- Fact 8.13.35, 492
- Fact 8.13.36, 492
- Fact 8.13.38, 492
- Fact 8.13.39, 493
- Fact 8.13.40, 493
- Fact 8.13.41, 493
- Fact 8.13.42, 493
- partitioned positive-semidefinite matrix**
 - Proposition 8.2.3, 420
- permutation matrix**
 - Fact 2.13.9, 130
- positive-definite matrix**
 - Proposition 8.4.14, 429
 - Fact 8.12.1, 475
 - Fact 8.13.6, 486
 - Fact 8.13.7, 486
 - Fact 8.13.8, 486
 - Fact 8.13.9, 486
 - Fact 8.13.10, 487
 - Fact 8.13.12, 487
 - Fact 8.13.13, 487
 - Fact 8.13.14, 487
 - Fact 8.13.15, 488
 - Fact 8.13.17, 488
 - Fact 8.13.19, 488
 - Fact 8.13.21, 488
 - Fact 8.13.23, 489
- positive-semidefinite matrix**
 - Corollary 8.4.15, 429
 - Fact 8.13.16, 488
 - Fact 8.13.18, 488
 - Fact 8.13.20, 488
 - Fact 8.13.21, 488
 - Fact 8.13.24, 489
 - Fact 8.13.29, 490
 - Fact 8.13.35, 492
 - Fact 8.13.36, 492
 - Fact 8.13.38, 492
 - Fact 8.13.39, 493
 - Fact 8.13.40, 493
 - Fact 8.13.41, 493
 - Fact 8.17.11, 511
 - Fact 8.18.30, 521
 - Fact 8.21.8, 533
- Fact 8.21.19, 534
- Fact 8.21.20, 535
- Fact 9.8.39, 578
- product**
 - Proposition 2.7.3, 104
- rank-deficient matrix**
 - Fact 2.13.3, 129
- reverse identity matrix**
 - Fact 2.13.1, 128
- row interchange**
 - Proposition 2.7.2, 103
- Schur complement**
 - Proposition 8.2.3, 420
- semidissipative matrix**
 - Fact 8.13.3, 485
 - Fact 8.13.4, 485
 - Fact 8.13.11, 486, 487
- singular value**
 - Fact 5.11.28, 326
 - Fact 5.11.29, 327
 - Fact 8.13.1, 485
 - Fact 9.13.23, 606
- singular values**
 - Fact 5.12.13, 335
- skew-Hermitian matrix**
 - Fact 3.7.11, 179
 - Fact 3.7.16, 181
 - Fact 8.13.6, 486
- skew-symmetric matrix**
 - Fact 3.7.15, 181
 - Fact 3.7.33, 184
 - Fact 4.8.14, 259
 - Fact 4.9.20, 263
 - Fact 4.10.2, 266
- strongly increasing function**
 - Proposition 8.6.13, 435
- subdeterminant**
 - Fact 2.13.4, 129
 - Fact 2.14.12, 135
- subdeterminant expansion**
 - Corollary 2.7.6, 106
- submatrix**
 - Fact 2.14.1, 132
- sum of Kronecker product**
 - Fact 7.5.12, 410
 - Fact 7.5.13, 410
- sum of matrices**
 - Fact 5.12.12, 335
 - Fact 9.14.18, 611
- sum of orthogonal matrices**
 - Fact 3.11.22, 196
- Sylvester's identity**
 - Fact 2.14.1, 132
- symplectic matrix**
 - Fact 3.19.10, 217
 - Fact 3.19.11, 217
- time-varying dynamics**
 - Fact 11.13.4, 678
- Toeplitz matrix**
 - Fact 2.13.13, 131
 - Fact 3.20.7, 219
- trace**
 - Proposition 8.4.14, 429
 - Corollary 11.2.4, 648
 - Corollary 11.2.5, 648
 - Fact 2.13.16, 132
 - Fact 8.12.1, 475
 - Fact 8.13.20, 488
 - Fact 11.14.20, 685
- transpose**
 - Proposition 2.7.1, 103
- tridiagonal matrix**
 - Fact 3.20.6, 218
 - Fact 3.20.7, 219
 - Fact 3.20.8, 219
 - Fact 3.20.9, 219
 - Fact 3.20.11, 220
- unimodular matrix**
 - Proposition 4.3.7, 238
- unitary matrix**
 - Fact 3.11.15, 194
 - Fact 3.11.20, 196
 - Fact 3.11.23, 196
 - Fact 3.11.24, 196
- upper bound**
 - Fact 2.13.14, 131
 - Fact 2.13.15, 131
 - Fact 8.13.32, 491
 - Fact 8.13.33, 491

- Fact 8.13.34, 491
- Vandermonde matrix**
 - Fact 5.16.3, 354
- determinant identities**
 - Magnus
 - Fact 2.13.16, 132
- determinant inequality**
 - Hua's inequalities
 - Fact 8.11.21, 472
- determinant lower bound**
 - nonsingular matrix
 - Fact 4.10.18, 269
- determinant of a partitioned matrix**
 - Hadamard's inequality
 - Fact 6.5.26, 392
- determinant of an outer-product perturbation**
 - Sherman-Morrison-Woodbury formula
 - Fact 2.16.3, 141
- determinantal compression**
 - partitioned matrix
 - Fact 8.13.42, 493
- diagonal**
 - eigenvalue
 - Fact 8.12.3, 476
 - positive-semidefinite matrix
 - Fact 8.12.3, 476
 - zero
 - Fact 5.9.18, 313
- diagonal dominance**
 - rank
 - Fact 4.10.23, 271
- diagonal dominance theorem**
 - nonsingular matrix
 - Fact 4.10.17, 269
 - Fact 4.10.18, 269

- diagonal entries**
 - definition, 80
 - Hermitian matrix
 - Fact 8.17.13, 512
 - similar matrices
 - Fact 5.9.13, 313
 - unitarily similar matrices
 - Fact 5.9.17, 313
 - Fact 5.9.19, 313
 - unitary matrix
 - Fact 3.11.19, 195
 - Fact 8.17.10, 511
- diagonal entries of a unitary matrix**
 - Schur-Horn theorem
 - Fact 3.11.19, 195
 - Fact 8.17.10, 511
- diagonal entry**
 - eigenvalue
 - Fact 8.17.8, 510
 - Hermitian matrix
 - Corollary 8.4.7, 427
 - Fact 8.17.8, 510
 - Fact 8.17.9, 510
 - positive-semidefinite matrix
 - Fact 8.10.16, 457
 - strong majorization
 - Fact 8.17.8, 510
- diagonal matrix**
 - controllability
 - Fact 12.20.12, 792
 - definition
 - Definition 3.1.3, 167
 - Hermitian matrix
 - Corollary 5.4.5, 294
 - Kronecker product
 - Fact 7.4.3, 405
 - matrix exponential
 - Fact 11.13.16, 680
 - orthogonally similar matrices
 - Fact 5.9.15, 313
 - unitary matrix
 - Theorem 5.6.4, 302
- diagonalizable matrix**

- $S - N$ decomposition**
 - Fact 5.9.3, 311
- additive**
 - decomposition
 - Fact 5.9.3, 311
- adjugate**
 - Fact 5.14.5, 339
- cogredient**
 - diagonalization
 - Fact 8.16.2, 507
 - Fact 8.16.3, 507
- commuting matrices**
 - Fact 5.17.8, 358
- complex conjugate**
 - transpose
 - Fact 5.14.5, 339
- eigenvector**
 - Fact 5.14.6, 339
- example**
 - Example 5.5.18, 299
- factorization**
 - Fact 5.15.27, 350
- involutory matrix**
 - Fact 5.14.20, 341
- Jordan-Chevalley decomposition**
 - Fact 5.9.3, 311
- simultaneous**
 - diagonalization
 - Fact 8.16.2, 507
 - Fact 8.16.3, 507
- transpose**
 - Fact 5.14.5, 339
- diagonalizable over $\mathbb{C}$**
 - definition
 - Definition 5.5.4, 296
- diagonalizable over $\mathbb{F}$**
 - identity perturbation
 - Fact 5.14.16, 341
- diagonalizable over $\mathbb{R}$**
 - asymptotically stable matrix
 - Fact 11.17.10, 697
 - definition
 - Definition 5.5.4, 296
 - factorization
 - Proposition 5.5.12, 297
 - Corollary 5.5.22, 301

- similar matrices
 - Proposition 5.5.12, 297
 - Corollary 5.5.22, 301
- diagonally dominant matrix**
 - nonsingular matrix
 - Fact 4.10.17, 269
- diagonally located block**
 - definition, 80
- Diaz-Goldman-Metcalf inequality**
 - Hölder's inequality
 - Fact 1.16.22, 65
- dictionary ordering**
 - cone
 - Fact 2.9.31, 115
 - total ordering
 - Fact 1.5.8, 12
- difference**
 - Frobenius norm
 - Fact 9.9.25, 584
 - generalized inverse
 - Fact 6.4.37, 384
 - idempotent matrix
 - Fact 5.12.19, 337
 - maximum singular value
 - Fact 8.18.8, 515
 - Fact 9.9.32, 585
 - projector
 - Fact 3.13.24, 210
 - Fact 5.12.17, 335
 - Fact 6.4.20, 381
 - Schatten norm
 - Fact 9.9.23, 584
 - singular value
 - Fact 8.18.9, 515
 - Fact 8.18.10, 515
 - trace norm
 - Fact 9.9.24, 584
- difference equation**
 - golden ratio
 - Fact 4.11.12, 277
 - nonnegative matrix
 - Fact 4.11.12, 277
- difference of idempotent matrices**
 - Makelainen
 - Fact 5.12.19, 337
 - Styan
 - Fact 5.12.19, 337
- difference of matrices**
 - idempotent matrix
 - Fact 3.12.25, 202
 - Fact 3.12.27, 203
 - Fact 3.12.28, 203
 - Fact 3.12.30, 204
 - Fact 3.12.32, 205
- differentiable function**
 - continuous function
 - Proposition 10.4.4, 626
 - definition
 - Definition 10.4.3, 626
- dihedral group**
 - group
 - Fact 3.21.7, 223
 - Klein four-group
 - Fact 3.21.7, 223
- dimension**
 - commuting matrices
 - Fact 5.10.15, 319
 - Fact 5.10.16, 319
 - product of matrices
 - Fact 2.10.14, 116
 - rank inequality
 - Fact 2.10.4, 115
 - solid set
 - Fact 10.8.16, 633
 - subspace
 - Fact 2.10.4, 115
 - subspace dimension theorem
 - Theorem 2.3.1, 90
 - subspace intersection
 - Fact 2.9.20, 112
 - Fact 2.9.21, 113
 - Fact 2.9.22, 113
 - variational cone
 - Fact 10.8.20, 633
 - zero trace
 - Fact 2.18.11, 150
- dimension of a subspace**
 - definition, 90
- dimension of an affine subspace**
 - definition, 90
- dimension of an arbitrary set**
 - definition, 90
- dimension theorem**
 - rank and defect
 - Corollary 2.5.5, 97
- directed cut**
 - graph
 - Fact 4.11.2, 273
- direction cosines**
 - Euler parameters
 - Fact 3.11.10, 192
 - orthogonal matrix
 - Fact 3.11.10, 192
- directional differential**
 - convex function
 - Fact 10.11.14, 639
- discrete Fourier analysis**
 - circulant matrix
 - Fact 5.16.7, 355
- discrete-time**
 - asymptotic stability
 - eigenvalue
 - Proposition 11.10.2, 670
 - linear dynamical system
 - Proposition 11.10.2, 670
 - matrix exponential
 - Proposition 11.10.2, 670
- discrete-time asymptotically stable matrix**
 - 2×2 matrix
 - Fact 11.21.1, 712

- asymptotically stable matrix
 - Fact 11.21.8, 713
- Cayley transform**
 - Fact 11.21.8, 713
- definition**
 - Definition 11.10.1, 670
- discrete-time asymptotically stable polynomial**
 - Proposition 11.10.4, 671
- dissipative matrix**
 - Fact 11.21.4, 712
- Kronecker product**
 - Fact 11.21.5, 713
 - Fact 11.21.6, 713
- Lyapunov equation**
 - Proposition 11.10.5, 671
- matrix exponential**
 - Fact 11.21.7, 713
- matrix limit**
 - Fact 11.21.12, 714
- matrix power**
 - Fact 11.21.2, 712
- normal matrix**
 - Fact 11.21.4, 712
- partitioned matrix**
 - Fact 11.21.9, 713
- positive-definite matrix**
 - Proposition 11.10.5, 671
 - Fact 11.21.9, 713
 - Fact 11.21.15, 714
- similar matrices**
 - Fact 11.18.4, 698
- discrete-time asymptotically stable polynomial**
 - definition**
 - Definition 11.10.3, 671
- discrete-time asymptotically stable matrix**
 - Proposition 11.10.4, 671
- polynomial coefficients**
 - Fact 11.20.1, 708
 - Fact 11.20.2, 709
 - Fact 11.20.3, 709
- discrete-time control problem**
 - LQG controller**
 - Fact 12.23.7, 804
- discrete-time dynamics matrix power**
 - Fact 11.21.3, 712
- discrete-time Lyapunov equation**
 - discrete-time asymptotically stable matrix**
 - Fact 11.21.15, 714
 - discrete-time Lyapunov-stable matrix**
 - Proposition 11.10.6, 671
 - Stein equation**
 - Fact 11.21.15, 714
- discrete-time Lyapunov stability eigenvalue**
 - Proposition 11.10.2, 670
- linear dynamical system**
 - Proposition 11.10.2, 670
- matrix exponential**
 - Proposition 11.10.2, 670
- discrete-time Lyapunov-stable matrix**
 - definition**
 - Definition 11.10.1, 670
 - discrete-time Lyapunov equation**
 - Proposition 11.10.6, 671
- discrete-time Lyapunov-stable polynomial**
 - definition**
 - Fact 11.21.8, 713
- discrete-time Lyapunov-stable polynomial**
 - Proposition 11.10.4, 671
- group generalized inverse**
 - Fact 11.21.11, 714
- Kreiss matrix theorem**
 - Fact 11.21.17, 715
- Kronecker product**
 - Fact 11.21.5, 713
 - Fact 11.21.6, 713
- logarithm**
 - Fact 11.14.19, 685
- matrix exponential**
 - Fact 11.21.7, 713
- matrix limit**
 - Fact 11.21.11, 714
- matrix power**
 - Fact 11.21.2, 712
 - Fact 11.21.10, 714
- maximum singular value**
 - Fact 11.21.17, 715
- normal matrix**
 - Fact 11.21.4, 712
- positive-definite matrix**
 - Proposition 11.10.6, 671
- positive-semidefinite matrix**
 - Fact 11.21.15, 714
- semicontractive matrix**
 - Fact 11.21.4, 712
- semidissipative matrix**
 - Fact 11.21.4, 712
- similar matrices**
 - Fact 11.18.4, 698
- unitary matrix**
 - Fact 11.21.13, 714
- discrete-time Lyapunov-stable polynomial**
 - definition**
 - Fact 11.21.8, 713

- Definition 11.10.3, 671
- discrete-time**
 - Lyapunov-stable matrix**
 - Proposition 11.10.4, 671
- discrete-time semistability**
 - eigenvalue**
 - Proposition 11.10.2, 670
 - linear dynamical system**
 - Proposition 11.10.2, 670
 - matrix exponential**
 - Proposition 11.10.2, 670
- discrete-time semistable matrix**
 - companion form matrix**
 - Fact 11.21.18, 715
 - convergent sequence**
 - Fact 11.21.14, 714
 - definition**
 - Definition 11.10.1, 670
 - discrete-time semistable polynomial**
 - Proposition 11.10.4, 671
 - idempotent matrix**
 - Fact 11.21.10, 714
 - Kronecker product**
 - Fact 11.21.5, 713
 - Fact 11.21.6, 713
 - limit**
 - Fact 11.21.10, 714
 - matrix exponential**
 - Fact 11.21.7, 713
 - Fact 11.21.14, 714
 - similar matrices**
 - Fact 11.18.4, 698
- discrete-time semistable polynomial**
 - definition**
- Definition 11.10.3, 671
- discrete-time**
 - semistable matrix**
 - Proposition 11.10.4, 671
- discrete-time time-varying system**
 - state convergence**
 - Fact 11.21.16, 715
- discriminant**
 - compound matrix**
 - Fact 7.5.17, 411
- disjoint**
 - definition, 3**
- dissipative matrix**
 - asymptotically stable matrix**
 - Fact 11.18.21, 701
 - Fact 11.18.37, 705
 - definition**
 - Definition 3.1.1, 165
 - determinant**
 - Fact 8.13.2, 485
 - Fact 8.13.11, 486, 487
 - Fact 8.13.31, 491
 - discrete-time asymptotically stable matrix**
 - Fact 11.21.4, 712
 - Frobenius norm**
 - Fact 11.15.3, 689
 - inertia**
 - Fact 5.8.12, 309
 - Kronecker sum**
 - Fact 7.5.8, 409
 - matrix exponential**
 - Fact 11.15.3, 689
 - maximum singular value**
 - Fact 8.17.12, 511
 - nonsingular matrix**
 - Fact 3.20.13, 220
 - normal matrix**
 - Fact 11.18.37, 705
 - positive-definite matrix**
 - Fact 8.17.12, 511
- Fact 11.18.21, 701
- range-Hermitian matrix**
 - Fact 5.14.31, 343
- semidissipative matrix**
 - Fact 8.13.31, 491
- spectrum**
 - Fact 8.13.31, 491
- strictly dissipative matrix**
 - Fact 8.9.31, 453
- unitary matrix**
 - Fact 8.9.31, 453
- distance from a point**
 - set**
 - Fact 10.9.15, 636
 - Fact 10.9.16, 636
- distance to singularity**
 - nonsingular matrix**
 - Fact 9.14.7, 608
- distinct eigenvalues**
 - eigenvector**
 - Proposition 4.5.4, 246
- distinct roots**
 - Bezout matrix**
 - Fact 4.8.9, 258
- distributive identities**
 - definition, 82**
- divides**
 - definition, 233**
- division of polynomial matrices**
 - quotient and remainder**
 - Lemma 4.2.1, 234
- Dixmier**
 - projectors and unitarily similar matrices**
 - Fact 5.10.12, 319
- Djokovic**
 - maximum singular value of a product**

- of elementary projectors
 - Fact 9.14.1, 607
- rank of a Kronecker product
- Fact 8.21.16, 534
- Schur product of positive-definite matrices
- Fact 8.21.13, 533
- Djokovic inequality**
- Euclidean norm
- Fact 9.7.7, 565
- domain**
- definition, 3
- Dormido**
- asymptotically stable polynomial
- Fact 11.17.8, 697
- double cover**
- orthogonal matrix parameterization
- Fact 3.11.10, 192
- spin group
- Fact 3.11.10, 192
- doublet**
- definition
- Fact 2.10.24, 118
- outer-product matrix
- Fact 2.10.24, 118
- Fact 2.12.6, 126
- spectrum
- Fact 5.11.13, 323
- doubly stochastic matrix**
- strong majorization
- Fact 2.21.7, 163
- Douglas-Fillmore-Williams lemma**
- factorization
- Theorem 8.6.2, 431
- Dragomir's inequality**
- harmonic mean
- Fact 1.16.24, 65
- Dragomir-Yang inequalities**
- Euclidean norm
- Fact 9.7.8, 566
- Fact 9.7.9, 566
- Drazin**
- real eigenvalues
- Fact 5.14.13, 340
- Drazin generalized inverse**
- block-circulant matrix
- Fact 6.6.1, 393
- commuting matrices
- Fact 6.6.4, 394
- Fact 6.6.5, 394
- definition, 367
- idempotent matrix
- Proposition 6.2.2, 368
- integral
- Fact 11.13.12, 679
- Fact 11.13.14, 680
- Kronecker product**
- Fact 7.4.31, 408
- matrix exponential**
- Fact 11.13.12, 679
- Fact 11.13.14, 680
- matrix limit**
- Fact 6.6.11, 395
- matrix product**
- Fact 6.6.3, 393
- Fact 6.6.4, 394
- matrix sum**
- Fact 6.6.5, 394
- null space**
- Proposition 6.2.2, 368
- partitioned matrix**
- Fact 6.6.1, 393
- Fact 6.6.2, 393
- positive-semidefinite matrix**
- Fact 8.20.2, 525
- range**
- Proposition 6.2.2, 368
- sum**
- Fact 6.6.1, 393
- tripotent matrix**
- Proposition 6.2.2, 368
- uniqueness
- Theorem 6.2.1, 367
- dual cone**
- convex conical hull
- Fact 2.9.3, 110
- definition, 91
- intersection
- Fact 2.9.5, 111
- sum of sets
- Fact 2.9.5, 111
- dual norm**
- adjoint norm
- Fact 9.8.8, 572
- definition
- Fact 9.7.22, 570
- induced norm
- Fact 9.7.22, 570
- quadratic form
- Fact 9.8.34, 577
- Dunkl-Williams inequality**
- complex numbers
- Fact 1.18.5, 71
- norm**
- Fact 9.7.10, 566
- Fact 9.7.13, 567
- dynamic compensator**
- LQG controller**
- Fact 12.23.6, 804
- Fact 12.23.7, 804
- E**
- Eckart-Young theorem**
- fixed-rank approximation
- Fact 9.14.28, 614
- eigensolution**
- eigenvector
- Fact 11.13.6, 679
- Fact 11.13.7, 679
- eigenvalue**
- $SO(n)$
- Fact 5.11.2, 321
- adjugate

Fact 4.10.7, 267
asymptotic spectrum
 Fact 4.10.28, 272
asymptotic stability
 Proposition 11.8.2, 662
bound
 Fact 4.10.22, 271
 Fact 5.11.22, 325
 Fact 5.11.23, 325
 Fact 9.11.7, 598
bounds
 Fact 4.10.16, 269
 Fact 4.10.20, 270
Cardano's trigonometric solution
 Fact 4.10.1, 265
Cartesian decomposition
 Fact 5.11.21, 325
convex function
 Corollary 8.6.19, 442
 Fact 8.18.5, 513
definition, 240
determinant
 Fact 5.11.28, 326, 327
 Fact 8.13.1, 485
diagonal entry
 Fact 8.12.3, 476
 Fact 8.17.8, 510
discrete-time asymptotic stability
 Proposition 11.10.2, 670
discrete-time Lyapunov stability
 Proposition 11.10.2, 670
discrete-time semistability
 Proposition 11.10.2, 670
Frobenius norm
 Fact 9.11.3, 597
 Fact 9.11.5, 598
generalized eigenvector
 Fact 5.14.8, 339

generalized Schur inequality
 Fact 9.11.6, 598
Hermitian matrix
 Theorem 8.4.5, 426
 Theorem 8.4.9, 427
 Theorem 8.4.11, 428
 Corollary 8.4.2, 425
 Corollary 8.4.6, 426
 Corollary 8.4.7, 427
 Corollary 8.6.19, 442
 Lemma 8.4.3, 425
 Lemma 8.4.4, 425
 Fact 8.10.4, 456
 Fact 8.15.20, 503
 Fact 8.15.31, 505
 Fact 8.17.8, 510
 Fact 8.17.9, 510
 Fact 8.17.13, 512
 Fact 8.17.15, 512
 Fact 8.17.16, 512
 Fact 8.18.4, 513
 Fact 8.18.17, 517
 Fact 8.21.28, 536
Hermitian part
 Fact 5.11.24, 325
Hölder matrix norm
 Fact 9.11.6, 598
Kronecker product
 Proposition 7.1.10, 401
 Fact 7.4.13, 406
 Fact 7.4.15, 406
 Fact 7.4.21, 406
 Fact 7.4.28, 407
 Fact 7.4.32, 408
Kronecker sum
 Proposition 7.2.3, 403
 Fact 7.5.5, 409
 Fact 7.5.7, 409
 Fact 7.5.16, 411
Lyapunov stability
 Proposition 11.8.2, 662
matrix logarithm
 Theorem 11.5.1, 656
matrix sum
 Fact 5.12.2, 333
 Fact 5.12.3, 333
normal matrix
 Fact 5.14.15, 341

orthogonal matrix
 Fact 5.11.2, 321
partitioned matrix
 Proposition 5.6.6, 303
 Fact 5.12.20, 337
 Fact 5.12.21, 337
 Fact 5.12.22, 338
positive-definite matrix
 Fact 8.10.24, 458
 Fact 8.15.20, 503
 Fact 8.15.29, 505
 Fact 8.15.30, 505
 Fact 8.18.29, 521
 Fact 8.21.21, 535
positive-semidefinite matrix
 Fact 8.12.3, 476
 Fact 8.15.11, 501
 Fact 8.18.6, 514
 Fact 8.18.19, 518
 Fact 8.18.20, 518
 Fact 8.18.22, 519
 Fact 8.18.23, 519
 Fact 8.18.24, 520
 Fact 8.18.27, 521
 Fact 8.20.17, 528
 Fact 8.21.18, 534
 Fact 8.21.20, 535
quadratic form
 Lemma 8.4.3, 425
 Fact 8.15.20, 503
root locus
 Fact 4.10.28, 272
Schatten norm
 Fact 9.11.6, 598
Schur product
 Fact 8.21.18, 534
Schur's inequality
 Fact 8.17.5, 509
 Fact 9.11.3, 597
semistability
 Proposition 11.8.2, 662
singular value
 Fact 8.17.5, 509
 Fact 8.17.6, 509
 Fact 9.13.22, 606
skew-Hermitian matrix

- Fact 5.11.6, 321
- skew-symmetric matrix**
 - Fact 4.10.2, 266
- spectral abscissa**
 - Fact 5.11.24, 325
- strong majorization**
 - Corollary 8.6.19, 442
 - Fact 8.17.8, 510
 - Fact 8.18.4, 513
 - Fact 8.18.29, 521
- subscript convention, 240**
- symmetric matrix**
 - Fact 4.10.1, 265
- trace**
 - Proposition 8.4.13, 428
 - Fact 5.11.11, 322
 - Fact 8.17.5, 509
 - Fact 8.18.18, 518
- weak log majorization**
 - Fact 8.18.27, 521
- weak majorization**
 - Fact 8.17.5, 509
 - Fact 8.18.5, 513
 - Fact 8.18.6, 514
 - Fact 8.18.27, 521
- eigenvalue bound**
 - Bendixson's theorem**
 - Fact 5.11.21, 325
 - Fact 9.11.8, 598
 - Browne's theorem**
 - Fact 5.11.21, 325
 - Frobenius norm**
 - Fact 9.12.3, 599
 - Henrici**
 - Fact 9.11.3, 597
 - Hermitian matrix**
 - Fact 9.12.3, 599
 - Hirsch's theorem**
 - Fact 5.11.21, 325
 - Hirsch's theorems**
 - Fact 9.11.8, 598
 - Hölder norm**
 - Fact 9.11.8, 598
 - trace**
 - Fact 5.11.45, 331
- eigenvalue bounds**
 - ovals of Cassini**
 - Fact 4.10.21, 271
- eigenvalue characterization**
 - minimum principle**
 - Fact 8.17.15, 512
- eigenvalue inclusion region**
 - Lyapunov equation**
 - Fact 12.21.20, 798
- eigenvalue inequality**
 - 2×2 matrix**
 - Fact 8.17.1, 508
 - Hermitian matrix**
 - Lemma 8.4.1, 424
 - Fact 8.18.3, 513
 - Poincaré separation theorem**
 - Fact 8.17.16, 512
- eigenvalue of Hermitian part**
 - maximum singular value**
 - Fact 5.11.25, 326
 - minimum singular value**
 - Fact 5.11.25, 326
 - singular value**
 - Fact 5.11.27, 326
 - Fact 8.17.4, 509
 - weak majorization**
 - Fact 5.11.27, 326
- eigenvalue perturbation**
 - Frobenius norm**
 - Fact 9.12.4, 599
 - Fact 9.12.9, 601
 - Fact 9.12.10, 601
 - Hermitian matrix**
 - Fact 4.10.27, 272
 - maximum singular value**
 - Fact 9.12.4, 599
 - Fact 9.12.8, 601
 - normal matrix**
 - Fact 9.12.8, 601
- partitioned matrix**
 - Fact 4.10.27, 272
- unitarily invariant norm**
 - Fact 9.12.4, 599
- eigenvalue placement**
 - controllable pair**
 - Proposition 12.6.19, 743
 - observable pair**
 - Proposition 12.3.19, 732
- eigenvector**
 - adjugate**
 - Fact 5.14.26, 342
 - commuting matrices**
 - Fact 5.14.25, 342
 - cyclic eigenvalue**
 - Fact 5.14.2, 338
 - definition, 245**
 - diagonalizable matrix**
 - Fact 5.14.6, 339
 - distinct eigenvalues**
 - Proposition 4.5.4, 246
 - eigensolution**
 - Fact 11.13.6, 679
 - Fact 11.13.7, 679
 - generalized eigensolution**
 - Fact 11.13.7, 679
 - Kronecker product**
 - Proposition 7.1.10, 401
 - Fact 7.4.21, 406
 - Fact 7.4.32, 408
 - Kronecker sum**
 - Proposition 7.2.3, 403
 - Fact 7.5.16, 411
 - M-matrix**
 - Fact 4.11.10, 276
 - normal matrix**
 - Proposition 4.5.4, 246
 - Lemma 4.5.3, 246
 - similarity transformation**
 - Fact 5.14.6, 339
 - Fact 5.14.7, 339

upper triangular
matrix
Fact 5.17.1, 358

either
definition, 1

element
definition, 2

elementary divisor
companion matrix
Theorem 5.2.9, 287
definition, 287
factorization
Fact 5.15.37, 351
hypercompanion
matrix
Lemma 5.3.1, 288

elementary matrix
definition
Definition 3.1.2, 166
inverse matrix
Fact 3.7.20, 182
nonsingular matrix
Fact 5.15.12, 347
properties and
matrix types
Fact 3.7.19, 181
semisimple matrix
Fact 5.14.17, 341
spectrum
Proposition 5.5.21, 300
unitarily similar
matrices
Proposition 5.6.3, 302

elementary
multicompanion
form
definition, 287

elementary polynomial
matrix
definition, 236

elementary projector
definition
Definition 3.1.1, 165

elementary reflector
Fact 3.13.7, 207
Fact 3.14.3, 211

hyperplane
Fact 3.13.8, 207
maximum singular
value
Fact 9.14.1, 607
reflector
Fact 5.15.13, 347
spectrum
Proposition 5.5.21, 300
trace
Fact 5.8.11, 309
unitarily similar
matrices
Proposition 5.6.3, 302

elementary reflector
definition
Definition 3.1.1, 165
elementary projector
Fact 3.13.7, 207
Fact 3.14.3, 211
hyperplane
Fact 3.14.5, 211
null space
Fact 3.13.7, 207
orthogonal matrix
Fact 5.15.15, 347
range
Fact 3.13.7, 207
rank
Fact 3.13.7, 207
reflection theorem
Fact 3.14.4, 211
reflector
Fact 5.15.14, 347
spectrum
Proposition 5.5.21, 300
trace
Fact 5.8.11, 309
unitarily similar
matrices
Proposition 5.6.3, 302

elementary symmetric
function
Schur concave
function
Fact 1.15.20, 53

elementary symmetric
mean

Newton's inequality
Fact 1.15.11, 50

elementary symmetric
polynomial
inequality
Fact 1.15.11, 50
Newton's identities
Fact 4.8.2, 254

ellipsoid
positive-definite
matrix
Fact 3.7.35, 184
volume
Fact 3.7.35, 184

Embry
commuting matrices
Fact 5.12.14, 335

empty matrix
definition, 83

empty set
definition, 2

Enestrom-Kakeya
theorem
polynomial root
locations
Fact 11.20.3, 709

entropy
logarithm
Fact 1.15.45, 59
Fact 1.15.46, 59
Fact 1.15.47, 59
Fact 1.16.30, 67

Schur concave
function
Fact 2.21.6, 162
strong majorization
Fact 2.21.6, 162

entry
definition, 79

EP matrix, see
range-Hermitian
matrix
definition, 229

equi-induced norm

- definition**
 - Definition 9.4.1, 553
- normalized norm**
 - Theorem 9.4.2, 553
- spectral radius**
 - Corollary 9.4.5, 554
- submultiplicative norm**
 - Corollary 9.4.4, 554
 - Fact 9.8.45, 579
- equi-induced self-adjoint norm**
 - maximum singular value
 - Fact 9.13.5, 602
- equi-induced unitarily invariant norm**
 - maximum singular value
 - Fact 9.13.4, 602
- equilateral triangle**
 - complex numbers
 - Fact 2.20.6, 155
- equilibrium**
 - definition, 660
- equivalence**
 - equivalence relation
 - Fact 5.10.3, 317
- equivalence class**
 - equivalent matrices
 - Fact 5.10.4, 317
 - induced by
 - equivalence relation
 - Theorem 1.3.6, 6
 - similar matrices
 - Fact 5.10.4, 317
 - unitarily similar matrices
 - Fact 5.10.4, 317
- equivalence class induced by**
 - definition, 6
- equivalence hull**
 - definition
 - Definition 1.3.4, 5
- relation**
 - Proposition 1.3.5, 6
- equivalence relation**
 - Abelian group**
 - Proposition 3.4.2, 173
 - congruence**
 - Fact 5.10.3, 317
 - definition**
 - Definition 1.3.2, 5
 - equivalence**
 - Fact 5.10.3, 317
 - equivalence class**
 - Theorem 1.3.6, 6
 - group**
 - Proposition 3.4.1, 173
 - Proposition 3.4.2, 173
 - intersection**
 - Proposition 1.3.3, 5
 - left equivalence**
 - Fact 5.10.3, 317
 - partition**
 - Theorem 1.3.7, 7
 - right equivalence**
 - Fact 5.10.3, 317
 - similarity**
 - Fact 5.10.3, 317
 - unitary**
 - biequivalence**
 - Fact 5.10.3, 317
 - unitary left equivalence**
 - Fact 5.10.3, 317
 - unitary right equivalence**
 - Fact 5.10.3, 317
 - unitary similarity**
 - Fact 5.10.3, 317
- equivalent matrices**
 - equivalence class
 - Fact 5.10.4, 317
- equivalent norms**
 - equivalence
 - Theorem 9.1.8, 546
 - norms**
 - Fact 9.8.12, 573
- equivalent realizations**
- controllable**
 - canonical form**
 - Corollary 12.9.9, 752
 - controllable pair**
 - Proposition 12.9.8, 752
 - invariant zero**
 - Proposition 12.10.10, 764
 - observable canonical form**
 - Corollary 12.9.9, 752
 - observable pair**
 - Proposition 12.9.8, 752
 - similar matrices**
 - Definition 12.9.6, 751
- ergodic theorem**
 - unitary matrix limit**
 - Fact 6.3.34, 376
- Euclidean distance matrix**
 - negative-semidefinite matrix**
 - Fact 9.8.14, 573
- Schoenberg**
 - Fact 9.8.14, 573
- Euclidean norm**
 - Cauchy-Schwarz inequality**
 - Corollary 9.1.7, 546
 - definition, 545**
 - Djokovic inequality**
 - Fact 9.7.7, 565
 - Dragomir-Yang inequalities**
 - Fact 9.7.8, 566
 - Fact 9.7.9, 566
 - generalized Hlawka inequality**
 - Fact 9.7.7, 565
 - inequality**
 - Fact 9.7.4, 563
 - Fact 9.7.6, 565
 - Fact 9.7.7, 565
 - Fact 9.7.8, 566
 - Fact 9.7.9, 566
 - Fact 9.7.18, 569
 - Kronecker product**
 - Fact 9.7.27, 570

- outer-product matrix**
Fact 9.7.27, 570
- projector**
Fact 9.8.2, 571
Fact 9.8.3, 571
Fact 10.9.17, 636
- reverse triangle inequality**
Fact 9.7.6, 565
- Euler constant**
logarithm
Fact 1.7.5, 18
- Euler parameters**
direction cosines
Fact 3.11.10, 192
orthogonal matrix
Fact 3.11.10, 192
Fact 3.11.11, 193
Rodrigues's formulas
Fact 3.11.11, 193
- Euler product formula**
prime numbers
Fact 1.7.8, 19
zeta function
Fact 1.7.8, 19
- Euler totient function**
positive-semidefinite matrix
Fact 8.8.5, 447
- Euler's inequality**
triangle
Fact 2.20.11, 156
- Euler's polyhedron formula**
face
Fact 1.6.7, 14
- even polynomial**
asymptotically stable polynomial
Fact 11.17.6, 696
definition, 232
- Everitt**
determinant of a partitioned
- positive-semidefinite matrix**
Fact 8.13.38, 492
- exactly proper rational function**
definition
Definition 4.7.1, 249
- exactly proper rational transfer function**
definition
Definition 4.7.2, 249
- existence of transformation**
Hermitian matrix
Fact 3.9.2, 185
orthogonal matrix
Fact 3.9.5, 186
outer-product matrix
Fact 3.9.1, 185
skew-Hermitian matrix
Fact 3.9.4, 186
- existential statement**
definition, 2
logical equivalents
Fact 1.5.4, 11
- exogenous input**
definition, 772
- exponent**
scalar inequality
Fact 1.9.1, 22
- exponential, see matrix exponential**
inequality
Fact 1.15.48, 60
matrix logarithm
Fact 11.14.26, 686
positive-definite matrix
Fact 11.14.26, 686
- exponential function**
complex numbers
Fact 1.18.6, 71
- convex function**
Fact 1.10.26, 34
inequality
Fact 1.10.27, 34
limit
Fact 1.9.18, 26
scalar inequalities
Fact 1.10.28, 35
scalar inequality
Fact 1.9.14, 25
Fact 1.9.15, 25
Fact 1.9.16, 25
Fact 1.9.17, 26
- exponential inequality**
scalar case
Fact 1.9.13, 24
- extended infinite interval**
definition, xxxv
- extreme point**
convex set
Fact 10.8.23, 634
Krein-Milman theorem
Fact 10.8.23, 634
- F**
- face**
Euler's polyhedron formula
Fact 1.6.7, 14
- fact**
definition, 1
- factorial**
bounds
Fact 1.9.20, 26
inequality
Fact 1.9.31, 30
Stirling's formula
Fact 1.9.19, 26
- factorization**
asymptotically stable matrix
Fact 11.18.22, 701
Bezout matrix

Fact 5.15.24, 349
commutator
 Fact 5.15.33, 351
complex conjugate transpose
 Fact 5.15.23, 349
determinant
 Fact 5.15.7, 346
 Fact 5.15.34, 351
diagonalizable matrix
 Fact 5.15.27, 350
diagonalizable over $\mathbb{R}$
 Proposition 5.5.12, 297
 Corollary 5.5.22, 301
Douglas-Fillmore-Williams lemma
 Theorem 8.6.2, 431
elementary divisor
 Fact 5.15.37, 351
full rank
 Fact 5.15.40, 351
generalized inverse
 Fact 6.5.25, 392
group generalized inverse
 Fact 6.6.12, 395
Hermitian matrix
 Fact 5.15.17, 348
 Fact 5.15.25, 349
 Fact 5.15.26, 349
 Fact 5.15.41, 351
 Fact 8.16.1, 507
idempotent matrix
 Fact 5.15.28, 350
 Fact 5.15.30, 350
involutory matrix
 Fact 5.15.18, 348
 Fact 5.15.31, 350
 Fact 5.15.32, 351
Jordan form
 Fact 5.15.5, 346
lower triangular matrix
 Fact 5.15.10, 346
LULU decomposition
 Fact 5.15.11, 346

nilpotent matrix
 Fact 5.15.29, 350
nonsingular matrix
 Fact 5.15.12, 347
 Fact 5.15.36, 351
orthogonal matrix
 Fact 5.15.15, 347
 Fact 5.15.16, 347
 Fact 5.15.31, 350
 Fact 5.15.35, 351
partitioned matrix, 420
 Proposition 2.8.3, 107
 Proposition 2.8.4, 107
 Fact 2.14.9, 134
 Fact 2.16.2, 141
 Fact 2.17.3, 147
 Fact 2.17.4, 147
 Fact 2.17.5, 147
 Fact 6.5.25, 392
 Fact 8.11.25, 473
 Fact 8.11.26, 473
positive-definite matrix
 Fact 5.15.26, 349
 Fact 5.18.4, 359
 Fact 5.18.5, 359
 Fact 5.18.6, 359
 Fact 5.18.8, 360
positive-semidefinite matrix
 Fact 5.15.22, 349
 Fact 5.15.26, 349
 Fact 5.18.2, 359
 Fact 5.18.3, 359
 Fact 5.18.7, 359
 Fact 8.9.36, 454
 Fact 8.9.37, 454
projector
 Fact 5.15.13, 347
 Fact 5.15.17, 348
 Fact 6.3.32, 376
range
 Theorem 8.6.2, 431
reflector
 Fact 5.15.14, 347
reverse-symmetric matrix
 Fact 5.9.12, 313

rotation-dilation
 Fact 2.19.2, 151
shear
 Fact 5.15.11, 346
similar matrices
 Fact 5.15.6, 346
skew-symmetric matrix
 Fact 5.15.37, 351
 Fact 5.15.38, 351
symmetric matrix
 Corollary 5.3.9, 292
 Fact 5.15.24, 349
ULU decomposition
 Fact 5.15.11, 346
unitary matrix
 Fact 5.15.8, 346
 Fact 5.18.6, 359
upper triangular matrix
 Fact 5.15.8, 346
 Fact 5.15.10, 346

Fan

convex function
 Proposition 8.6.17, 542
trace of a Hermitian matrix product
 Fact 5.12.4, 333
trace of a product of orthogonal matrices
 Fact 5.12.10, 334

Fan constant

definition
 Fact 8.10.48, 465

Fan dominance theorem

singular value
 Fact 9.14.19, 611

Farkas theorem

linear system solution
 Fact 4.11.14, 279

fast Fourier transform

circulant matrix
 Fact 5.16.7, 355

feedback**interconnection****realization**

Proposition 12.13.4,
772

Proposition 12.14.1,
774

Fact 12.22.8, 799

transfer function

Fact 12.22.8, 799

feedback signal

definition, 772

Fejer's theorem

positive-semidefinite
matrix

Fact 8.21.35, 538

Fer expansion

time-varying
dynamics

Fact 11.13.4, 678

Fibonacci numbers**determinant**

Fact 4.11.12, 277

generating function

Fact 4.11.12, 277

nonnegative matrix

Fact 4.11.12, 277

field of values

spectrum of convex
hull

Fact 8.14.7, 496

Fact 8.14.8, 497

final state**controllability**

Fact 12.20.4, 791

controllable subspace

Fact 12.20.3, 791

finite group**group**

Fact 3.21.7, 223

representation

Fact 3.21.9, 224

finite interval

definition, xxxv

finite-sum solution**Lyapunov equation**

Fact 12.21.17, 797

Finsler's lemma**positive-definite****linear combination**

Fact 8.15.24, 504

Fact 8.15.25, 504

Fischer's inequality**positive-semidefinite****matrix determinant**

Fact 8.13.35, 492

Fact 8.13.36, 492

positive-semidefinite**matrix determinant****reverse inequality**

Fact 8.13.41, 493

fixed-point theorem**continuous function**

Theorem 10.3.10, 625

Corollary 10.3.11, 625

fixed-rank**approximation****Eckart-Young****theorem**

Fact 9.14.28, 614

Frobenius norm

Fact 9.14.28, 614

Fact 9.15.4, 618

least squares

Fact 9.14.28, 614

Fact 9.15.4, 618

Schmidt-Mirsky**theorem**

Fact 9.14.28, 614

singular value

Fact 9.14.28, 614

Fact 9.15.4, 618

unitarily invariant**norm**

Fact 9.14.28, 614

forced response

definition, 725

forest**symmetric graph**

Fact 1.6.5, 14

Fourier matrix**circulant matrix**

Fact 5.16.7, 355

Vandermonde matrix

Fact 5.16.7, 355

Fourier transform**Parseval's theorem**

Fact 12.22.1, 798

Frame**finite sequence for****inverse matrix**

Fact 2.16.28, 146

Franck

maximum singular
value lower bound
on distance to
singularity

Fact 9.14.6, 608

Fréchet derivative**definition**

Definition 10.4.3, 626

uniqueness

Proposition 10.4.2, 626

free response

definition, 725

frequency domain**controllability****Gramian**

Corollary 12.11.5, 767

frequency response**imaginary part**

Fact 12.22.5, 799

real part

Fact 12.22.5, 799

transfer function

Fact 12.22.5, 799

Friedland

matrix exponential
and singular value

Fact 11.16.15, 695

Frobenius

similar to transpose

Corollary 5.3.8, 291

singular value

Corollary 9.6.7, 562

symmetric matrix
factorization
Fact 5.15.24, 349

Frobenius canonical form, see multicompanion form
definition, 362

Frobenius inequality
rank of partitioned matrix
Fact 2.11.14, 123
Fact 6.5.15, 389

Frobenius matrix
definition, 362

Frobenius norm
absolute value
Fact 9.13.11, 603
adjugate
Fact 9.8.15, 573
Cauchy-Schwarz inequality
Corollary 9.3.9, 553
commutator
Fact 9.9.26, 584
Fact 9.9.27, 584
definition, 547
determinant
Fact 9.8.39, 578
dissipative matrix
Fact 11.15.3, 689
eigenvalue
Fact 9.11.3, 597
Fact 9.11.5, 598
eigenvalue bound
Fact 9.12.3, 599
eigenvalue perturbation
Fact 9.12.4, 599
Fact 9.12.9, 601
Fact 9.12.10, 601
fixed-rank
approximation
Fact 9.14.28, 614
Fact 9.15.4, 618
Hermitian matrix
Fact 9.9.41, 588

inequality
Fact 9.9.25, 584
Kronecker product
Fact 9.14.37, 617
matrix difference
Fact 9.9.25, 584
matrix exponential
Fact 11.14.32, 688
Fact 11.15.3, 689
maximum singular value bound
Fact 9.13.13, 604
normal matrix
Fact 9.12.9, 601
outer-product matrix
Fact 9.7.26, 570
polar decomposition
Fact 9.9.42, 588
positive-semidefinite matrix
Fact 9.8.39, 578
Fact 9.9.12, 581
Fact 9.9.15, 582
Fact 9.9.27, 584
rank
Fact 9.11.4, 598
Fact 9.14.28, 614
Fact 9.15.4, 618
Schatten norm, 549
Fact 9.8.20, 575
Schur product
Fact 9.14.34, 616
Schur's inequality
Fact 9.11.3, 597
spectral radius
Fact 9.13.12, 603
trace
Fact 9.11.3, 597
Fact 9.11.4, 598
Fact 9.11.5, 598
Fact 9.12.2, 599
trace norm
Fact 9.9.11, 581
triangle inequality
Fact 9.9.13, 582
unitarily invariant norm
Fact 9.14.34, 616

unitary matrix
Fact 9.9.42, 588

Fujii-Kubo
polynomial root bound
Fact 11.20.9, 710

Fujiwara's bound
polynomial
Fact 11.20.8, 710

full column rank
definition, 95
equivalent properties
Theorem 2.6.1, 98
nonsingular equivalence
Corollary 2.6.6, 101

full rank
definition, 96

full row rank
definition, 95
equivalent properties
Theorem 2.6.1, 98
nonsingular equivalence
Corollary 2.6.6, 101

full-rank factorization
generalized inverse
Fact 6.4.9, 379
idempotent matrix
Fact 3.12.23, 202

full-state feedback
controllable subspace
Proposition 12.6.5, 737
controllably asymptotically stable
Proposition 12.7.2, 743
determinant
Fact 12.22.14, 800
invariant zero
Proposition 12.10.10, 764
Fact 12.22.14, 800
stabilizability
Proposition 12.8.2, 747

uncontrollable
eigenvalue
Proposition 12.6.14,
740

unobservable
eigenvalue
Proposition 12.3.14,
731

unobservable
subspace
Proposition 12.3.5, 729

function

definition, 3

graph

Fact 1.6.1, 13

Fact 1.6.2, 13

Fact 1.6.3, 13

intersection

Fact 1.5.11, 12

Fact 1.5.12, 12

relation

Proposition 1.3.1, 5

union

Fact 1.5.11, 12

Fact 1.5.12, 12

function composition

matrix

multiplication

Theorem 2.1.3, 81

fundamental theorem of algebra

definition, 232

fundamental triangle inequality

Ramus

Fact 2.20.11, 156

triangle

Fact 2.20.11, 156

Wu

Fact 2.20.11, 156

Furuta inequality

positive-definite

matrix

Fact 8.10.50, 465

positive-semidefinite

matrix inequality

Proposition 8.6.7, 433

spectral order
Fact 8.19.4, 523

G

Galois

quintic polynomial

Fact 3.21.7, 223

gamma

logarithm

Fact 1.7.5, 18

gap topology

minimal principal

angle

Fact 10.9.18, 636

subspace

Fact 10.9.18, 636

Gastinel

distance to

singularity of a

nonsingular matrix

Fact 9.14.7, 608

generalized algebraic multiplicity

definition, 305

generalized

Cayley-Hamilton

theorem

commuting matrices

Fact 4.9.7, 261

generalized eigensolution

eigenvector

Fact 11.13.7, 679

generalized eigenvalue

definition, 304

pencil

Proposition 5.7.3, 305

Proposition 5.7.4, 306

regular pencil

Proposition 5.7.3, 305

Proposition 5.7.4, 306

singular pencil

Proposition 5.7.3, 305

generalized eigenvector

eigenvalue

Fact 5.14.8, 339

generalized geometric mean

positive-definite

matrix

Fact 8.10.45, 464

generalized geometric multiplicity

definition, 305

generalized Hölder inequality

vector

Fact 9.7.34, 571

generalized inverse (1,3) inverse

Fact 6.3.14, 372

(1,4) inverse

Fact 6.3.14, 372

adjugate

Fact 6.3.6, 370

Fact 6.3.7, 371

Fact 6.5.16, 389

basic properties

Proposition 6.1.6, 365

block-circulant

matrix

Fact 6.5.2, 386

centrohermitian

matrix

Fact 6.3.31, 376

characteristic

polynomial

Fact 6.3.20, 374

characterization

Fact 6.4.1, 377

complex conjugate

transpose

Fact 6.3.9, 371

Fact 6.3.10, 371

Fact 6.3.13, 372

Fact 6.3.16, 373

Fact 6.3.17, 373

Fact 6.3.18, 373

Fact 6.3.22, 374

- Fact 6.3.27, 375
- Fact 6.3.28, 375
- Fact 6.4.7, 379
- Fact 6.6.16, 396
- Fact 6.6.17, 397
- Fact 6.6.18, 397
- congruence**
 - Fact 8.20.5, 525
- convergent sequence**
 - Fact 6.3.35, 376
 - Fact 6.3.36, 377
- definition, 363**
- determinant**
 - Fact 6.5.26, 392
 - Fact 6.5.27, 392
 - Fact 6.5.28, 393
- difference**
 - Fact 6.4.33, 383
- factorization**
 - Fact 6.5.25, 392
- full-rank**
 - factorization**
 - Fact 6.4.9, 379
- group generalized**
 - inverse**
 - Fact 6.6.7, 394
- Hermitian matrix**
 - Fact 6.3.21, 374
 - Fact 6.4.3, 378
 - Fact 8.20.12, 527
- idempotent matrix**
 - Fact 5.12.18, 336
 - Fact 6.3.22, 374
 - Fact 6.3.23, 374
 - Fact 6.3.24, 374
 - Fact 6.3.25, 375
 - Fact 6.3.26, 375
 - Fact 6.3.27, 375
 - Fact 6.4.18, 381
 - Fact 6.4.19, 381
 - Fact 6.4.20, 381
 - Fact 6.4.22, 381
 - Fact 6.4.25, 381
- identity**
 - Fact 6.3.33, 376
- inertia**
 - Fact 6.3.21, 374
 - Fact 8.20.12, 527
- integral**
 - Fact 11.13.10, 679
- Jordan canonical form**
 - Fact 6.6.9, 394
- Kronecker product**
 - Fact 7.4.30, 408
- least squares**
 - Fact 9.15.1, 618
 - Fact 9.15.2, 618
 - Fact 9.15.3, 618
- left inverse**
 - Corollary 6.1.4, 364
 - Fact 6.4.39, 384
 - Fact 6.4.40, 384
- left-inner matrix**
 - Fact 6.3.8, 371
- left-invertible matrix**
 - Proposition 6.1.5, 364
- linear matrix equation**
 - Fact 6.4.38, 384
- linear system**
 - Proposition 6.1.7, 366
- matrix difference**
 - Fact 6.4.37, 384
- matrix exponential**
 - Fact 11.13.10, 679
- matrix inversion lemma**
 - Fact 6.4.4, 378
- matrix limit**
 - Fact 6.3.19, 374
- matrix product**
 - Fact 6.4.5, 378
 - Fact 6.4.6, 378
 - Fact 6.4.8, 379
 - Fact 6.4.9, 379
 - Fact 6.4.10, 379
 - Fact 6.4.11, 379
 - Fact 6.4.12, 379
 - Fact 6.4.13, 380
 - Fact 6.4.14, 380
 - Fact 6.4.16, 380
 - Fact 6.4.17, 380
 - Fact 6.4.21, 381
 - Fact 6.4.22, 381
 - Fact 6.4.23, 381
 - Fact 6.4.30, 382
 - Fact 6.4.31, 382
- matrix sum**
 - Fact 6.4.34, 383
 - Fact 6.4.35, 383
 - Fact 6.4.36, 383
- maximum singular value**
 - Fact 9.14.8, 608
 - Fact 9.14.30, 615
- Newton-Raphson algorithm**
 - Fact 6.3.35, 376
- normal matrix**
 - Proposition 6.1.6, 365
 - Fact 6.3.16, 373
 - Fact 6.3.17, 373
- null space**
 - Proposition 6.1.6, 365
 - Fact 6.3.24, 374
- observability matrix**
 - Fact 12.20.19, 793
- outer-product matrix**
 - Fact 6.3.2, 370
- outer-product perturbation**
 - Fact 6.4.2, 377
- partial isometry**
 - Fact 6.3.28, 375
- partitioned matrix**
 - Fact 6.3.30, 376
 - Fact 6.5.1, 385
 - Fact 6.5.2, 386
 - Fact 6.5.3, 386
 - Fact 6.5.4, 386
 - Fact 6.5.13, 388
 - Fact 6.5.17, 390
 - Fact 6.5.18, 390
 - Fact 6.5.19, 390
 - Fact 6.5.20, 391
 - Fact 6.5.21, 391
 - Fact 6.5.22, 391
 - Fact 6.5.23, 391
 - Fact 6.5.24, 391
 - Fact 8.20.22, 530
- positive-definite matrix**
 - Proposition 6.1.6, 365
 - Fact 6.4.7, 379

positive-semidefinite matrix

Proposition 6.1.6, 365
 Fact 6.4.4, 378
 Fact 8.20.1, 525
 Fact 8.20.2, 525
 Fact 8.20.3, 525
 Fact 8.20.4, 525
 Fact 8.20.6, 526
 Fact 8.20.7, 526
 Fact 8.20.8, 526
 Fact 8.20.9, 526
 Fact 8.20.10, 526
 Fact 8.20.11, 527
 Fact 8.20.13, 527
 Fact 8.20.15, 527
 Fact 8.20.16, 527
 Fact 8.20.17, 528
 Fact 8.20.18, 528
 Fact 8.20.19, 530
 Fact 8.20.20, 530
 Fact 8.20.22, 530
 Fact 8.20.23, 531

projector

Fact 6.3.3, 370
 Fact 6.3.4, 370
 Fact 6.3.5, 370
 Fact 6.3.26, 375
 Fact 6.3.27, 375
 Fact 6.3.32, 376
 Fact 6.4.15, 380
 Fact 6.4.16, 380
 Fact 6.4.17, 380
 Fact 6.4.18, 381
 Fact 6.4.19, 381
 Fact 6.4.21, 381
 Fact 6.4.23, 381
 Fact 6.4.24, 381
 Fact 6.4.25, 381
 Fact 6.4.33, 383
 Fact 6.4.41, 384
 Fact 6.4.46, 385
 Fact 6.5.10, 388

range

Proposition 6.1.6, 365
 Fact 6.3.24, 374
 Fact 6.4.42, 384
 Fact 6.4.43, 385
 Fact 6.5.3, 386

range-Hermitian matrix

Proposition 6.1.6, 365
 Fact 6.3.10, 371
 Fact 6.3.11, 372
 Fact 6.3.12, 372
 Fact 6.3.16, 373
 Fact 6.3.17, 373
 Fact 6.4.26, 382
 Fact 6.4.27, 382
 Fact 6.4.28, 382
 Fact 6.4.29, 382

rank

Fact 6.3.9, 371
 Fact 6.3.22, 374
 Fact 6.3.36, 377
 Fact 6.4.2, 377
 Fact 6.4.44, 385
 Fact 6.5.6, 386
 Fact 6.5.8, 387
 Fact 6.5.9, 387
 Fact 6.5.12, 388
 Fact 6.5.13, 388
 Fact 6.5.14, 388

rank subtractivity partial ordering

Fact 6.5.30, 393

right inverse

Corollary 6.1.4, 364

right-inner matrix

Fact 6.3.8, 371

right-invertible matrix

Proposition 6.1.5, 364

sequence

Fact 6.3.36, 377

singular value

Fact 6.3.29, 376

singular value decomposition

Fact 6.3.15, 373

square root

Fact 8.20.4, 525

star partial ordering

Fact 8.19.8, 524

star-dagger matrix

Fact 6.3.13, 372

sum

Fact 6.5.1, 385

Fact 6.5.2, 386

trace

Fact 6.3.22, 374

uniqueness

Theorem 6.1.1, 363

unitary matrix

Fact 6.3.34, 376

Urquhart

Fact 6.3.14, 372

generalized Löwner partial ordering**definition**

Fact 8.19.10, 524

generalized multispectrum**definition, 304****generalized projector****range-Hermitian matrix**

Fact 3.6.4, 178

generalized Schur inequality**eigenvalues**

Fact 9.11.6, 598

generalized spectrum**definition, 304****generating function****Fibonacci numbers**

Fact 4.11.12, 277

geometric mean**arithmetic mean**

Fact 1.15.21, 53

Fact 1.15.23, 53

Fact 1.15.24, 54

Fact 1.15.25, 54

Fact 1.15.26, 54

Fact 1.15.27, 54

determinant

Fact 8.10.43, 461

matrix exponential

Fact 8.10.44, 464

matrix logarithm

Fact 11.14.39, 689

Muirhead's theorem

Fact 1.15.25, 54

- nondecreasing function
 - Fact 8.10.43, 461
 - Fact 8.10.44, 464
- positive-definite matrix
 - Fact 8.10.43, 461
 - Fact 8.10.46, 464
 - Fact 8.21.51, 541
- positive-semidefinite matrix
 - Fact 8.10.43, 461
- Riccati equation
 - Fact 12.23.4, 802
- scalar inequality
 - Fact 1.11.6, 39
- Schur product
 - Fact 8.21.51, 541
- geometric multiplicity**
 - algebraic multiplicity
 - Proposition 5.5.3, 295
- block-diagonal matrix
 - Proposition 5.5.13, 298
- cascaded systems
 - Fact 12.22.15, 801
- controllability
 - Fact 12.20.14, 792
- defect
 - Proposition 4.5.2, 246
- definition
 - Definition 4.5.1, 245
- partitioned matrix
 - Proposition 5.5.14, 298
- rank
 - Proposition 4.5.2, 246
- similar matrices
 - Proposition 5.5.10, 297
- geometric-mean decomposition**
 - unitary matrix
 - Fact 5.9.30, 316
- Gershgorin circle theorem**
 - eigenvalue bounds
 - Fact 4.10.16, 269
 - Fact 4.10.20, 270
- Gerstenhaber**
 - dimension of the algebra generated by two commuting matrices
 - Fact 5.10.21, 319
- Gibson**
 - diagonal entries of similar matrices
 - Fact 5.9.13, 313
- Givens rotation**
 - orthogonal matrix
 - Fact 5.15.16, 347
- global asymptotic stability**
 - nonlinear system
 - Theorem 11.7.2, 661
- globally asymptotically stable equilibrium**
 - definition
 - Definition 11.7.1, 660
- Gohberg-Semencul formulas**
 - Bezout matrix
 - Fact 4.8.6, 255
- golden mean**
 - positive-definite solution of a Riccati equation
 - Fact 12.23.4, 802
 - Riccati equation
 - Fact 12.23.4, 802
- golden ratio**
 - difference equation
 - Fact 4.11.12, 277
 - Riccati equation
 - Fact 12.23.4, 802
- Golden-Thompson inequality**
 - matrix exponential
 - Fact 11.14.28, 687
 - Fact 11.16.4, 692
- Gordan's theorem**
 - positive vector
 - Fact 4.11.15, 279
- gradient**
 - definition, 627
- Gram matrix**
 - positive-semidefinite matrix
 - Fact 8.9.36, 454
- Gram-Schmidt orthonormalization**
 - upper triangular matrix factorization
 - Fact 5.15.9, 346
- Gramian**
 - controllability
 - Fact 12.20.17, 792
 - stabilization
 - Fact 12.20.17, 792
- Graph**
 - definition, 3
- graph**
 - antisymmetric graph
 - Fact 4.11.1, 272
 - cycle
 - Fact 1.6.4, 13
 - definition, 8
 - directed cut
 - Fact 4.11.2, 273
 - function
 - Fact 1.6.1, 13
 - Fact 1.6.2, 13
 - Fact 1.6.3, 13
 - Hamiltonian cycle
 - Fact 1.6.6, 14
 - irreducible matrix
 - Fact 4.11.2, 273
 - Laplacian matrix
 - Fact 8.15.36, 506
 - spanning path
 - Fact 1.6.6, 14
 - symmetric graph
 - Fact 4.11.1, 272
 - tournament
 - Fact 1.6.6, 14
 - walk
 - Fact 4.11.3, 273
- graph of a matrix**
 - adjacency matrix

- Proposition 3.2.5, 171
- definition**
 - Definition 3.2.4, 171
- greatest common divisor**
 - definition, 233
- greatest lower bound**
 - projector
 - Fact 6.4.41, 384
- greatest upper bound for a partial ordering**
 - definition
 - Definition 1.3.9, 7
- Gregory's series**
 - infinite series
 - Fact 1.18.8, 73
- Greville**
 - generalized inverse of a matrix product
 - Fact 6.4.10, 379
 - Fact 6.4.12, 379
 - generalized inverse of a partitioned matrix
 - Fact 6.5.17, 390
- group**
 - alternating group
 - Fact 3.21.7, 223
 - circulant matrix
 - Fact 3.21.7, 224
 - Fact 3.21.8, 224
 - classical
 - Proposition 3.3.6, 172
 - cyclic group
 - Fact 3.21.7, 223
 - definition**
 - Definition 3.3.3, 172
 - dihedral group
 - Fact 3.21.7, 223
 - equivalence relation
 - Proposition 3.4.1, 173
 - Proposition 3.4.2, 173
 - finite group
 - Fact 3.21.7, 223
 - Fact 3.21.9, 224
 - icosahedral group
 - Fact 3.21.7, 223
 - isomorphism
 - Proposition 3.3.5, 172
 - Lie group**
 - Definition 11.6.1, 658
 - Proposition 11.6.2, 658
 - matrix exponential**
 - Proposition 11.6.7, 659
 - octahedral group**
 - Fact 3.21.7, 223
 - orthogonal matrix**
 - Fact 3.21.11, 225
 - pathwise connected**
 - Proposition 11.6.8, 660
 - permutation group**
 - Fact 3.21.7, 223
 - real numbers**
 - Fact 3.21.1, 221
 - symmetry group**
 - Fact 3.21.7, 223
 - tetrahedral group**
 - Fact 3.21.7, 223
 - transpose**
 - Fact 3.21.10, 225
 - unipotent matrix**
 - Fact 3.21.5, 222
 - Fact 11.22.1, 715
 - unit sphere**
 - Fact 3.21.2, 221
 - upper triangular matrix**
 - Fact 3.21.5, 222
 - Fact 11.22.1, 715
- group generalized inverse**
 - complex conjugate transpose
 - Fact 6.6.10, 394
 - definition, 369**
 - discrete-time Lyapunov-stable matrix
 - Fact 11.21.11, 714
 - factorization**
 - Fact 6.6.12, 395
 - generalized inverse
 - Fact 6.6.7, 394
 - idempotent matrix**
 - Proposition 6.2.3, 369
- integral**
 - Fact 11.13.13, 680
 - Fact 11.13.14, 680
- irreducible matrix**
 - Fact 6.6.20, 398
- Kronecker product**
 - Fact 7.4.31, 408
- limit**
 - Fact 6.6.14, 395
- matrix exponential**
 - Fact 11.13.13, 680
 - Fact 11.13.14, 680
 - Fact 11.18.5, 698
 - Fact 11.18.6, 698
- normal matrix**
 - Fact 6.6.10, 394
- null space**
 - Proposition 6.2.3, 369
- positive-semidefinite matrix**
 - Fact 8.20.1, 525
- range**
 - Proposition 6.2.3, 369
- range-Hermitian matrix**
 - Fact 6.6.8, 394
- singular value decomposition**
 - Fact 6.6.15, 395
- trace**
 - Fact 6.6.6, 394
- group-invertible matrix**
 - almost nonnegative matrix
 - Fact 11.19.4, 706
 - complementary subspaces**
 - Corollary 3.5.8, 176
 - definition**
 - Definition 3.1.1, 165
 - equivalent characterizations**
 - Fact 3.6.1, 177
 - Hermitian matrix**
 - Fact 6.6.18, 397
 - idempotent matrix**
 - Proposition 3.1.6, 169
 - Proposition 3.5.9, 176
 - Proposition 6.2.3, 369

Fact 5.11.8, 322
index of a matrix
 Proposition 3.5.6, 176
 Corollary 5.5.9, 297
 Fact 5.14.4, 339
inertia
 Fact 5.8.5, 308
Jordan canonical form
 Fact 6.6.9, 394
Kronecker product
 Fact 7.4.16, 406
 Fact 7.4.31, 408
Lyapunov-stable matrix
 Fact 11.18.2, 698
matrix exponential
 Fact 11.18.14, 700
matrix power
 Fact 3.6.2, 177
 Fact 6.6.19, 398
N-matrix
 Fact 11.19.4, 706
normal matrix
 Fact 6.6.17, 397
outer-product matrix
 Fact 5.14.3, 338
positive-definite matrix
 Fact 8.10.12, 457
positive-semidefinite matrix
 Fact 8.10.12, 457
projector
 Fact 3.13.21, 209
range
 Fact 5.14.4, 339
range-Hermitian matrix
 Proposition 3.1.6, 169
 Fact 6.6.16, 396
rank
 Fact 5.8.5, 308
 Fact 5.14.4, 339
semistable matrix
 Fact 11.18.3, 698
similar matrices
 Proposition 3.4.5, 174

Fact 5.9.5, 312
spectrum
 Proposition 5.5.21, 300
square root
 Fact 5.15.20, 348
stable subspace
 Proposition 11.8.8, 665
tripotent matrix
 Proposition 3.1.6, 169
unitarily similar matrices
 Proposition 3.4.5, 174

groups

complex representation
 Fact 3.21.8, 224
representation
 Fact 3.21.8, 224

H**H₂ norm**

controllability Gramian
 Corollary 12.11.4, 767
 Corollary 12.11.5, 767
definition
 Definition 12.11.2, 766
observability Gramian
 Corollary 12.11.4, 767
Parseval's theorem
 Theorem 12.11.3, 766
partitioned transfer function
 Fact 12.22.16, 801
 Fact 12.22.17, 801
quadratic performance measure
 Proposition 12.15.1, 776
submultiplicative norm
 Fact 12.22.20, 801
sum of transfer functions

Proposition 12.11.6, 767

transfer function

Fact 12.22.16, 801
 Fact 12.22.17, 801
 Fact 12.22.18, 801
 Fact 12.22.19, 801

Hadamard product, see Schur product**Hadamard's inequality**

determinant
 Fact 8.13.33, 491
 Fact 8.13.34, 491
determinant bound
 Fact 9.11.1, 596
determinant of a partitioned matrix
 Fact 6.5.26, 392
positive-semidefinite matrix determinant
 Fact 8.17.11, 511

Hadamard-Fischer inequality

positive-semidefinite matrix
 Fact 8.13.36, 492

Hahn-Banach theorem

inner product inequality
 Fact 10.9.12, 635

half-vectorization operator

Kronecker product, 416

Hamiltonian

block decomposition
 Proposition 12.17.5, 783
closed-loop spectrum
 Proposition 12.16.14, 781
definition, 780
Jordan form
 Fact 12.23.1, 802
Riccati equation
 Theorem 12.17.9, 784

- Proposition 12.16.14, 781
- Corollary 12.16.15, 781
- spectral factorization**
 - Proposition 12.16.13, 780
- spectrum**
 - Theorem 12.17.9, 784
 - Proposition 12.16.13, 780
 - Proposition 12.17.5, 783
 - Proposition 12.17.7, 784
 - Proposition 12.17.8, 784
 - Lemma 12.17.4, 783
 - Lemma 12.17.6, 783
- stabilizability**
 - Fact 12.23.1, 802
- stabilizing solution**
 - Corollary 12.16.15, 781
- uncontrollable**
 - eigenvalue**
 - Proposition 12.17.7, 784
 - Proposition 12.17.8, 784
 - Lemma 12.17.4, 783
 - Lemma 12.17.6, 783
- unobservable**
 - eigenvalue**
 - Proposition 12.17.7, 784
 - Proposition 12.17.8, 784
 - Lemma 12.17.4, 783
 - Lemma 12.17.6, 783
- Hamiltonian cycle**
 - definition**
 - Definition 1.4.3, 9
 - graph**
 - Fact 1.6.6, 14
 - tournament**
 - Fact 1.6.6, 14
- Hamiltonian graph**
 - definition**
 - Definition 1.4.3, 9
- Hamiltonian matrix**
 - Cayley transform**
 - Fact 3.19.12, 217
 - characteristic polynomial**
 - Fact 4.9.21, 264
 - Fact 4.9.23, 264
 - definition**
 - Definition 3.1.5, 169
 - identity**
 - Fact 3.19.1, 216
 - inverse matrix**
 - Fact 3.19.5, 216
 - matrix exponential**
 - Proposition 11.6.7, 659
 - matrix logarithm**
 - Fact 11.14.19, 685
 - matrix sum**
 - Fact 3.19.5, 216
 - orthogonal matrix**
 - Fact 3.19.13, 217
 - orthosymplectic matrix**
 - Fact 3.19.13, 217
 - partitioned matrix**
 - Proposition 3.1.7, 169
 - Fact 3.19.6, 216
 - Fact 3.19.8, 217
 - Fact 4.9.22, 264
 - Fact 5.12.21, 337
 - skew reflector**
 - Fact 3.19.3, 216
 - skew-involutory matrix**
 - Fact 3.19.2, 216
 - Fact 3.19.3, 216
 - skew-symmetric matrix**
 - Fact 3.7.34, 184
 - Fact 3.19.3, 216
 - Fact 3.19.8, 217
 - spectrum**
 - Proposition 5.5.21, 300
 - symplectic matrix**
 - Fact 3.19.2, 216
 - Fact 3.19.12, 217
 - Fact 3.19.13, 217
 - symplectic similarity**
 - Fact 3.19.4, 216
- trace**
 - Fact 3.19.7, 216
- unit imaginary matrix**
 - Fact 3.19.3, 216
- Hamiltonian path**
 - definition**
 - Definition 1.4.3, 9
- Hankel matrix**
 - block-Hankel matrix**
 - Fact 3.18.3, 215
 - definition**
 - Definition 3.1.3, 167
- Hilbert matrix**
 - Fact 3.18.4, 215
- Markov**
 - block-Hankel matrix**
 - definition, 754
 - rational function**
 - Fact 4.8.8, 257
 - symmetric matrix**
 - Fact 3.18.2, 215
 - Toeplitz matrix**
 - Fact 3.18.1, 215
- Hanner inequality**
 - Hölder norm**
 - Fact 9.7.21, 569
 - Schatten norm**
 - Fact 9.9.36, 586
- Hansen**
 - trace of a convex function**
 - Fact 8.12.33, 482
- Hardy**
 - Hölder-induced norm**
 - Fact 9.8.17, 574
- Hardy inequality**
 - sum of powers**
 - Fact 1.15.42, 58
- Hardy-Hilbert inequality**
 - sum of powers**
 - Fact 1.16.13, 63
 - Fact 1.16.14, 63

Hardy-Littlewood rearrangement inequality

- sum of products
 - Fact 1.16.4, 60
- sum of products inequality
 - Fact 1.16.5, 60

Hardy-Littlewood-Polya theorem

- doubly stochastic matrix
 - Fact 2.21.7, 163

harmonic mean

- arithmetic-mean inequality
 - Fact 1.15.16, 52
 - Fact 1.15.17, 52
- arithmetic-mean–geometric-mean inequality
 - Fact 1.15.15, 52

Dragomir's inequality

- Fact 1.16.24, 65

harmonic steady-state response

- linear system
 - Theorem 12.12.1, 768

Hartwig

- rank of an idempotent matrix
 - Fact 3.12.27, 203

Haynsworth

- positive-semidefinite matrix
 - Fact 5.14.13, 340
- Schur complement of a partitioned matrix
 - Fact 6.5.29, 393

Haynsworth inertia additivity formula

- Schur complement
 - Fact 6.5.5, 386

Heinz inequality

- unitarily invariant norm
 - Fact 9.9.49, 589

Heinz mean

- scalar inequality
 - Fact 1.10.38, 38

Heisenberg group

- unipotent matrix
 - Fact 3.21.5, 222
 - Fact 11.22.1, 715
- upper triangular matrix
 - Fact 3.21.5, 222
 - Fact 11.22.1, 715

Henrici

- eigenvalue bound
 - Fact 9.11.3, 597

Hermite-Biehler theorem

- asymptotically stable polynomial
 - Fact 11.17.6, 696

Hermite-Hadamard inequality

- convex function
 - Fact 1.8.6, 22

Hermitian matrix

- additive decomposition
 - Fact 3.7.29, 183
- adjugate
 - Fact 3.7.10, 179
- affine mapping
 - Fact 3.7.14, 181
- block-diagonal matrix
 - Fact 3.7.8, 179

Cartesian

- decomposition
 - Fact 3.7.27, 182
 - Fact 3.7.28, 183
 - Fact 3.7.29, 183

cogredient

- transformation
 - Fact 8.16.4, 507
 - Fact 8.16.6, 507

commutator

- Fact 3.8.1, 184
- Fact 3.8.3, 185
- Fact 9.9.30, 585

commuting matrices

- Fact 5.14.29, 342

complex conjugate transpose

- Fact 3.7.13, 180
- Fact 5.9.8, 312
- Fact 6.6.18, 397

congruent matrices

- Proposition 3.4.5, 174
- Corollary 5.4.7, 294

convergent sequence

- Fact 11.14.7, 683
- Fact 11.14.8, 683

convex function

- Fact 8.12.32, 482
- Fact 8.12.33, 482

convex hull

- Fact 8.17.8, 510

defect

- Fact 5.8.7, 308
- Fact 8.9.7, 451

definition

- Definition 3.1.1, 165

determinant

- Corollary 8.4.10, 427
- Fact 3.7.21, 182
- Fact 8.13.7, 486

diagonal

- Fact 8.17.8, 510

diagonal entries

- Fact 8.17.13, 512

diagonal entry

- Corollary 8.4.7, 427
- Fact 8.17.8, 510
- Fact 8.17.9, 510

diagonal matrix

- Corollary 5.4.5, 294

eigenvalue

- Theorem 8.4.5, 426
- Theorem 8.4.9, 427
- Theorem 8.4.11, 428
- Corollary 8.4.2, 425
- Corollary 8.4.6, 426
- Corollary 8.4.7, 427
- Corollary 8.6.19, 442

- Lemma 8.4.3, 425
- Lemma 8.4.4, 425
- Fact 8.10.4, 456
- Fact 8.15.20, 503
- Fact 8.15.31, 505
- Fact 8.17.8, 510
- Fact 8.17.9, 510
- Fact 8.17.15, 512
- Fact 8.17.16, 512
- Fact 8.18.4, 513
- Fact 8.18.17, 517
- Fact 8.21.28, 536
- eigenvalue bound**
 - Fact 9.12.3, 599
- eigenvalue inequality**
 - Lemma 8.4.1, 424
 - Fact 8.18.3, 513
- eigenvalue perturbation**
 - Fact 4.10.27, 272
- eigenvalues**
 - Fact 8.17.13, 512
- existence of transformation**
 - Fact 3.9.2, 185
- factorization**
 - Fact 5.15.17, 348
 - Fact 5.15.25, 349
 - Fact 5.15.26, 349
 - Fact 5.15.41, 351
 - Fact 8.16.1, 507
- Frobenius norm**
 - Fact 9.9.41, 588
- generalized inverse**
 - Fact 6.3.21, 374
 - Fact 6.4.3, 378
 - Fact 8.20.12, 527
- group-invertible matrix**
 - Fact 6.6.18, 397
- inequality**
 - Fact 8.9.13, 452
 - Fact 8.9.15, 452
 - Fact 8.9.20, 452
 - Fact 8.13.26, 490
 - Fact 8.13.30, 490
- inertia**
 - Theorem 8.4.11, 428
 - Proposition 5.4.6, 294
- Fact 5.8.6, 308
- Fact 5.8.8, 308
- Fact 5.8.12, 309
- Fact 5.8.13, 309
- Fact 5.8.14, 309
- Fact 5.8.15, 309
- Fact 5.8.16, 310
- Fact 5.8.17, 310
- Fact 5.8.18, 310
- Fact 5.8.19, 310
- Fact 5.12.1, 333
- Fact 6.3.21, 374
- Fact 8.10.15, 457
- Fact 8.20.12, 527
- Fact 8.20.14, 527
- Fact 12.21.1, 793
- Fact 12.21.2, 794
- Fact 12.21.3, 794
- Fact 12.21.4, 794
- Fact 12.21.5, 794
- Fact 12.21.6, 795
- Fact 12.21.7, 795
- Fact 12.21.8, 795
- Fact 12.21.10, 796
- Fact 12.21.11, 796
- Fact 12.21.12, 796
- Kronecker product**
 - Fact 7.4.16, 406
 - Fact 8.21.28, 536
- Kronecker sum**
 - Fact 7.5.8, 409
- limit**
 - Fact 8.10.1, 456
- linear combination**
 - Fact 8.15.24, 504
 - Fact 8.15.25, 504
 - Fact 8.15.26, 504
- linear combination of projectors**
 - Fact 5.19.10, 361
- matrix exponential**
 - Proposition 11.2.8, 649
 - Proposition 11.2.9, 650
 - Proposition 11.4.5, 654
 - Corollary 11.2.6, 648
 - Fact 11.14.7, 683
 - Fact 11.14.8, 683
 - Fact 11.14.21, 685
 - Fact 11.14.28, 687
- Fact 11.14.29, 687
- Fact 11.14.31, 688
- Fact 11.14.32, 688
- Fact 11.14.34, 688
- Fact 11.15.1, 689
- Fact 11.16.4, 692
- Fact 11.16.5, 694
- Fact 11.16.13, 695
- Fact 11.16.17, 695
- maximum eigenvalue**
 - Lemma 8.4.3, 425
 - Fact 5.11.5, 321
 - Fact 8.10.3, 456
- maximum singular value**
 - Fact 5.11.5, 321
 - Fact 9.9.41, 588
- minimum eigenvalue**
 - Lemma 8.4.3, 425
 - Fact 8.10.3, 456
- normal matrix**
 - Proposition 3.1.6, 169
- outer-product matrix**
 - Fact 3.7.18, 181
 - Fact 3.9.2, 185
- partitioned matrix**
 - Fact 3.7.27, 182
 - Fact 4.10.27, 272
 - Fact 5.8.19, 310
 - Fact 5.12.1, 333
 - Fact 6.5.5, 386
- positive-definite matrix**
 - Fact 5.15.41, 351
 - Fact 8.10.13, 457
 - Fact 8.13.7, 486
- positive-semidefinite matrix**
 - Fact 5.15.41, 351
 - Fact 8.9.11, 452
 - Fact 8.10.13, 457
- product**
 - Example 5.5.19, 300
- projector**
 - Fact 3.13.2, 206
 - Fact 3.13.13, 208
 - Fact 3.13.20, 209
 - Fact 5.15.17, 348

properties of $<$ and $\leq$

Proposition 8.1.2, 418

quadratic form

Fact 3.7.6, 178

Fact 3.7.7, 179

Fact 8.15.24, 504

Fact 8.15.25, 504

Fact 8.15.26, 504

Fact 8.15.31, 505

quadratic matrix equation

Fact 5.11.4, 321

range

Lemma 8.6.1, 431

rank

Fact 3.7.22, 182

Fact 3.7.30, 183

Fact 5.8.6, 308

Fact 5.8.7, 308

Fact 8.9.7, 451

Rayleigh quotient

Lemma 8.4.3, 425

reflector

Fact 3.14.2, 211

Schatten norm

Fact 9.9.27, 584

Fact 9.9.39, 587

Schur decomposition

Corollary 5.4.5, 294

Schur product

Fact 8.21.28, 536

Fact 8.21.32, 537

signature

Fact 5.8.6, 308

Fact 5.8.7, 308

Fact 8.10.17, 457

similar matrices

Proposition 5.5.12, 297

simultaneous

diagonalization

Fact 8.16.1, 507

Fact 8.16.4, 507

Fact 8.16.6, 507

skew-Hermitian matrix

Fact 3.7.9, 179

Fact 3.7.28, 183

skew-symmetric matrix

Fact 3.7.9, 179

spectral abscissa

Fact 5.11.5, 321

spectral radius

Fact 5.11.5, 321

spectral variation

Fact 9.12.5, 600

Fact 9.12.7, 601

spectrum

Proposition 5.5.21, 300

Lemma 8.4.8, 427

spread

Fact 8.15.31, 505

strong majorization

Fact 8.17.8, 510

submatrix

Theorem 8.4.5, 426

Corollary 8.4.6, 426

Lemma 8.4.4, 425

Fact 5.8.8, 308

symmetric matrix

Fact 3.7.9, 179

trace

Proposition 8.4.13, 428

Corollary 8.4.10, 427

Lemma 8.4.12, 428

Fact 3.7.13, 180

Fact 3.7.22, 182

Fact 8.12.38, 483

trace of a product

Fact 8.12.6, 476

Fact 8.12.7, 477

Fact 8.12.8, 477

Fact 8.12.16, 478

trace of product

Fact 5.12.4, 333

Fact 5.12.5, 333

Fact 8.18.18, 518

tripotent matrix

Fact 3.16.3, 213

unitarily invariant norm

Fact 9.9.5, 580

Fact 9.9.41, 588

Fact 9.9.43, 588

Fact 11.16.13, 695

unitarily similar matrices

Proposition 3.4.5, 174

Proposition 5.6.3, 302

Corollary 5.4.5, 294

unitary matrix

Fact 3.11.29, 197

Fact 8.16.1, 507

Fact 11.14.34, 688

Hermitian matrix eigenvalue

Cauchy interlacing theorem

Lemma 8.4.4, 425

inclusion principle

Theorem 8.4.5, 426

Hermitian matrix eigenvalues

monotonicity theorem

Theorem 8.4.9, 427

Fact 8.10.4, 456

Weyl's inequality

Theorem 8.4.9, 427

Fact 8.10.4, 456

Hermitian matrix inertia identity

Styan

Fact 8.10.15, 457

Hermitian part

eigenvalue

Fact 5.11.24, 325

Hermitian perturbation

Lidskii-Mirsky-

Wielandt

theorem

Fact 9.12.4, 599

Heron mean

logarithmic mean

Fact 1.10.37, 37

Heron's formula

triangle

Fact 2.20.11, 156

Hessenberg matrix

lower or upper

- Definition 3.1.3, 167
- Hessian**
 - definition, 627
- hidden convexity**
 - quadratic form
 - Fact 8.14.11, 498
- Hilbert matrix**
 - Hankel matrix
 - Fact 3.18.4, 215
 - positive-definite matrix
 - Fact 3.18.4, 215
- Hille-Yosida theorem**
 - matrix exponential bound
 - Fact 11.15.8, 691
- Hirsch's theorem**
 - eigenvalue bound
 - Fact 5.11.21, 325
 - Fact 9.11.8, 598
- Hlawka's equality**
 - norm identity
 - Fact 9.7.4, 563
- Hlawka's inequality**
 - Euclidean norm
 - Fact 9.7.7, 565
 - norm inequality
 - Fact 9.7.4, 563
- Hoffman**
 - eigenvalue perturbation
 - Fact 9.12.9, 601
- Hoffman-Wielandt theorem**
 - eigenvalue perturbation
 - Fact 9.12.9, 601
- Hölder norm**
 - compatible norms
 - Proposition 9.3.5, 550
 - complex conjugate
 - Fact 9.7.33, 571
 - definition, 544
 - eigenvalue
 - Fact 9.11.6, 598
- eigenvalue bound**
 - Fact 9.11.8, 598
- Hanner inequality**
 - Fact 9.7.21, 569
- Hölder-induced norm**
 - Proposition 9.4.11, 556
 - Fact 9.7.28, 571
 - Fact 9.8.12, 573
 - Fact 9.8.17, 574
 - Fact 9.8.18, 574
 - Fact 9.8.19, 575
 - Fact 9.8.29, 576
- inequality**
 - Proposition 9.1.5, 545
 - Proposition 9.1.6, 545
 - Fact 9.7.18, 569
 - Fact 9.7.19, 569
 - Fact 9.7.21, 569
 - Fact 9.7.29, 571
- Kronecker product**
 - Fact 9.9.61, 591
- matrix**
 - definition, 547
- Minkowski's inequality**
 - Lemma 9.1.3, 544
- monotonicity**
 - Proposition 9.1.5, 545
- power-sum inequality**
 - Fact 1.15.34, 57
- Schatten norm**
 - Proposition 9.2.5, 549
 - Fact 9.11.6, 598
- submultiplicative norm**
 - Fact 9.9.20, 583
- vector**
 - Fact 9.7.34, 571
- vector norm**
 - Proposition 9.1.4, 544
- Hölder's inequality**
 - Diaz-Goldman-Metcalf inequality
 - Fact 1.16.22, 65
- positive-semidefinite matrix**
 - Fact 8.12.12, 477
- positive-semidefinite matrix trace**
 - Fact 8.12.11, 477
- reversal**
 - Fact 1.16.22, 65
- scalar case**
 - Fact 1.16.11, 62
 - Fact 1.16.12, 62
- vector inequality**
 - Proposition 9.1.6, 545
- Hölder-induced lower bound**
 - definition, 559
- Hölder-induced norm**
 - absolute value
 - Fact 9.8.26, 576
 - adjoint norm
 - Fact 9.8.10, 572
 - column norm
 - Fact 9.8.21, 575
 - Fact 9.8.23, 575
 - Fact 9.8.25, 576
 - complex conjugate
 - Fact 9.8.27, 576
 - complex conjugate transpose
 - Fact 9.8.28, 576
 - definition, 554
 - field
 - Proposition 9.4.7, 554
 - formulas
 - Proposition 9.4.9, 555
 - Hardy**
 - Fact 9.8.17, 574
 - Hölder norm**
 - Proposition 9.4.11, 556
 - Fact 9.7.28, 571
 - Fact 9.8.12, 573
 - Fact 9.8.17, 574
 - Fact 9.8.18, 574
 - Fact 9.8.19, 575
 - Fact 9.8.29, 576
 - inequality**
 - Fact 9.8.21, 575
 - Fact 9.8.22, 575

- Littlewood**
 - Fact 9.8.17, 574
 - Fact 9.8.18, 574
 - maximum singular value**
 - Fact 9.8.21, 575
 - monotonicity**
 - Proposition 9.4.6, 554
 - Orlicz**
 - Fact 9.8.18, 574
 - partitioned matrix**
 - Fact 9.8.11, 572
 - quadratic form**
 - Fact 9.8.35, 577
 - Fact 9.8.36, 577
 - row norm**
 - Fact 9.8.21, 575
 - Fact 9.8.23, 575
 - Fact 9.8.25, 576
 - Hölder-McCarthy inequality**
 - quadratic form
 - Fact 8.15.14, 502
 - Hopf's theorem**
 - eigenvalues of a positive matrix
 - Fact 4.11.20, 280
 - Horn**
 - diagonal entries of a unitary matrix
 - Fact 8.17.10, 511
 - Householder matrix, see elementary reflector**
 - definition, 229
 - Householder reflector, see elementary reflector**
 - definition, 229
 - Hsu**
 - orthogonally similar matrices
 - Fact 5.9.15, 313
 - Hua's inequalities**
 - determinant inequality
 - Fact 8.11.21, 472
 - positive-semidefinite matrix
 - Fact 8.11.21, 472
 - Hua's inequality**
 - scalar inequality
 - Fact 1.15.13, 51
 - Hua's matrix equality**
 - positive-semidefinite matrix
 - Fact 8.11.21, 472
 - Hurwitz matrix, see asymptotically stable matrix**
 - Hurwitz polynomial, see asymptotically stable polynomial**
 - asymptotically stable polynomial
 - Fact 11.18.23, 702
 - Huygens**
 - polynomial bound
 - Fact 11.20.13, 712
 - Huygens's inequality**
 - trigonometric inequality
 - Fact 1.9.29, 28
 - hyperbolic identities**
 - Fact 1.19.2, 75
 - hyperbolic inequality**
 - scalar
 - Fact 1.9.29, 28
 - Fact 1.10.29, 35
 - hypercompanion form**
 - existence
 - Theorem 5.3.2, 288
 - Theorem 5.3.3, 289
 - hypercompanion matrix**
 - companion matrix
 - Corollary 5.3.4, 289
 - Lemma 5.3.1, 288
 - definition, 288**
 - elementary divisor**
 - Lemma 5.3.1, 288
 - example**
 - Example 5.3.6, 290
 - Example 5.3.7, 291
 - real Jordan form**
 - Fact 5.10.1, 316
 - similarity transformation**
 - Fact 5.10.1, 316
 - hyperellipsoid**
 - volume
 - Fact 3.7.35, 184
 - hyperplane**
 - definition, 91
 - elementary projector
 - Fact 3.13.8, 207
 - elementary reflector
 - Fact 3.14.5, 211
- I**
- icosahedral group**
 - group
 - Fact 3.21.7, 223
- idempotent matrix**
 - commutator
 - Fact 3.12.16, 200
 - Fact 3.12.17, 200
 - Fact 3.12.30, 204
 - Fact 3.12.31, 204
 - Fact 3.12.32, 205
 - Fact 3.15.4, 200
 - commuting matrices
 - Fact 3.16.5, 213
 - complementary idempotent matrix
 - Fact 3.12.12, 199
 - complementary subspaces
 - Proposition 3.5.3, 176
 - Proposition 3.5.4, 176
 - Fact 3.12.1, 198
 - Fact 3.12.33, 205
 - complex conjugate
 - Fact 3.12.7, 199

complex conjugate
transpose

Fact 3.12.7, 199

Fact 5.9.21, 314

definition

Definition 3.1.1, 165

difference

Fact 3.12.25, 202

Fact 3.12.30, 204

Fact 5.12.19, 337

difference of
matrices

Fact 3.12.27, 203

Fact 3.12.28, 203

Fact 3.12.32, 205

discrete-time
semistable matrix

Fact 11.21.10, 714

Drazin generalized
inverse

Proposition 6.2.2, 368

factorization

Fact 5.15.28, 350

Fact 5.15.30, 350

full-rank
factorization

Fact 3.12.23, 202

generalized inverse

Fact 5.12.18, 336

Fact 6.3.22, 374

Fact 6.3.23, 374

Fact 6.3.24, 374

Fact 6.3.25, 375

Fact 6.3.26, 375

Fact 6.3.27, 375

Fact 6.4.18, 381

Fact 6.4.19, 381

Fact 6.4.20, 381

Fact 6.4.22, 381

Fact 6.4.25, 381

group generalized
inverse

Proposition 6.2.3, 369

group-invertible
matrix

Proposition 3.1.6, 169

Proposition 3.5.9, 176

Proposition 6.2.3, 369

Fact 5.11.8, 322

identities

Fact 3.12.18, 200

identity perturbation

Fact 3.12.13, 199

inertia

Fact 5.8.1, 307

involutory matrix

Fact 3.15.2, 212

Kronecker product

Fact 7.4.16, 406

left inverse

Fact 3.12.10, 199

linear combination

Fact 3.12.26, 203

Fact 3.12.28, 203

Fact 5.19.9, 361

matrix exponential

Fact 11.11.1, 671

Fact 11.16.12, 695

matrix product

Fact 3.12.21, 201

Fact 3.12.23, 202

matrix sum

Fact 3.12.26, 203

Fact 5.19.7, 361

Fact 5.19.8, 361

Fact 5.19.9, 361

maximum singular
value

Fact 5.11.38, 328

Fact 5.11.39, 329

Fact 5.12.18, 336

nilpotent matrix

Fact 3.12.16, 200

nonsingular matrix

Fact 3.12.11, 199

Fact 3.12.26, 203

Fact 3.12.28, 203

Fact 3.12.32, 205

norm

Fact 11.16.12, 695

normal matrix

Fact 3.13.3, 206

null space

Fact 3.12.3, 199

Fact 3.15.4, 200

Fact 6.3.24, 374

onto a subspace
along another
subspace

definition, 176

outer-product
matrix

Fact 3.7.18, 181

Fact 3.12.6, 199

partitioned matrix

Fact 3.12.14, 200

Fact 3.12.20, 201

Fact 3.12.33, 205

Fact 5.10.22, 320

positive-definite
matrix

Fact 5.15.30, 350

positive-semidefinite
matrix

Fact 5.15.30, 350

power

Fact 3.12.3, 198

product

Fact 3.12.29, 203

projector

Fact 3.13.3, 206

Fact 3.13.13, 208

Fact 3.13.20, 209

Fact 3.13.24, 210

Fact 5.10.13, 319

Fact 5.12.18, 336

Fact 6.3.26, 375

Fact 6.4.18, 381

Fact 6.4.19, 381

Fact 6.4.20, 381

Fact 6.4.25, 381

quadratic form

Fact 3.13.11, 208

range

Fact 3.12.3, 199

Fact 3.12.4, 199

Fact 3.15.4, 200

Fact 6.3.24, 374

range-Hermitian
matrix

Fact 3.13.3, 206

Fact 6.3.27, 375

rank

Fact 3.12.6, 199

Fact 3.12.9, 199

Fact 3.12.19, 201
 Fact 3.12.20, 201
 Fact 3.12.22, 201
 Fact 3.12.24, 202
 Fact 3.12.25, 202
 Fact 3.12.27, 203
 Fact 3.12.31, 204
 Fact 5.8.1, 307
 Fact 5.11.7, 322
right inverse
 Fact 3.12.10, 199
semisimple matrix
 Fact 5.14.21, 341
similar matrices
 Proposition 3.4.5, 174
 Proposition 5.6.3, 302
 Corollary 5.5.22, 301
 Fact 5.10.9, 318
 Fact 5.10.13, 319
 Fact 5.10.14, 319
 Fact 5.10.22, 320
singular value
 Fact 5.11.38, 328
skew-Hermitian matrix
 Fact 3.12.8, 199
skew-idempotent matrix
 Fact 3.12.5, 199
spectrum
 Proposition 5.5.21, 300
 Fact 5.11.7, 322
stable subspace
 Proposition 11.8.8, 665
submultiplicative norm
 Fact 9.8.6, 572
sum
 Fact 3.12.22, 201
trace
 Fact 5.8.1, 307
 Fact 5.11.7, 322
transpose
 Fact 3.12.7, 199
tripotent matrix
 Fact 3.16.1, 212
 Fact 3.16.5, 213
unitarily similar matrices

Proposition 3.4.5, 174
 Fact 5.9.21, 314
 Fact 5.9.26, 315
 Fact 5.9.27, 315
 Fact 5.10.10, 318
unstable subspace
 Proposition 11.8.8, 665
idempotent matrix onto a subspace along another subspace
 definition, 176
identity
 cube root
 Fact 2.12.23, 128
identity function
 definition, 3
identity matrix
 definition, 83
 symplectic matrix
 Fact 3.19.3, 216
identity perturbation
 cyclic matrix
 Fact 5.14.16, 341
 defective matrix
 Fact 5.14.16, 341
 derogatory matrix
 Fact 5.14.16, 341
 diagonalizable over $\mathbb{F}$
 Fact 5.14.16, 341
 inverse matrix
 Fact 4.8.12, 259
 semisimple matrix
 Fact 5.14.16, 341
 simple matrix
 Fact 5.14.16, 341
 spectrum
 Fact 4.10.13, 268
 Fact 4.10.14, 269
identity shift
 controllable subspace
 Lemma 12.6.7, 738
 unobservable subspace
 Lemma 12.3.7, 730
identity theorem

matrix function
 evaluation
 Theorem 10.5.3, 629
identric mean
 arithmetic mean
 Fact 1.10.36, 37
 logarithmic mean
 Fact 1.10.36, 37
image
 definition, 3
imaginary part
 frequency response
 Fact 12.22.5, 799
 transfer function
 Fact 12.22.5, 799
imaginary vector
 definition, 85
implication
 definition, 1
improper rational function
 definition
 Definition 4.7.1, 249
improper rational transfer function
 definition
 Definition 4.7.2, 249
impulse function
 definition, 724
impulse response
 definition, 725
impulse response function
 definition, 725
inbound Laplacian matrix
 adjacency matrix
 Theorem 3.2.2, 170
 definition
 Definition 3.2.1, 170
incidence matrix
 definition
 Definition 3.2.1, 170

- Laplacian matrix
 - Theorem 3.2.2, 170
 - Theorem 3.2.3, 171
- inclusion principle**
 - Hermitian matrix eigenvalue
 - Theorem 8.4.5, 426
- inclusion-exclusion principle**
 - cardinality
 - Fact 1.5.5, 11
- increasing function**
 - convex function
 - Theorem 8.6.15, 436
 - definition
 - Definition 8.6.12, 434
 - log majorization
 - Fact 2.21.12, 163
 - logarithm
 - Proposition 8.6.13, 435
 - matrix functions
 - Proposition 8.6.13, 435
 - positive-definite matrix
 - Fact 8.10.53, 466
 - Schur complement
 - Proposition 8.6.13, 435
 - weak majorization
 - Fact 2.21.10, 163
- increasing sequence**
 - positive-semidefinite matrix
 - Proposition 8.6.3, 432
- indecomposable matrix, see irreducible matrix**
 - definition, 229
- indegree**
 - graph
 - Definition 1.4.3, 9
- indegree matrix**
 - definition
 - Definition 3.2.1, 170
- index of a matrix**
 - block-triangular matrix
 - Fact 5.14.32, 343
 - Fact 6.6.13, 395
 - complementary subspaces
 - Proposition 3.5.7, 176
 - definition
 - Definition 3.5.5, 176
 - group-invertible matrix
 - Proposition 3.5.6, 176
 - Corollary 5.5.9, 297
 - Fact 5.14.4, 339
 - Kronecker product
 - Fact 7.4.26, 407
 - outer-product matrix
 - Fact 5.14.3, 338
 - partitioned matrix
 - Fact 5.14.32, 343
 - Fact 6.6.13, 395
 - range
 - Fact 5.14.4, 339
 - rank
 - Proposition 5.5.2, 295
- index of an eigenvalue**
 - algebraic multiplicity
 - Proposition 5.5.6, 296
 - definition
 - Definition 5.5.1, 295
 - Jordan block
 - Proposition 5.5.3, 295
 - minimal polynomial
 - Proposition 5.5.15, 299
 - rank
 - Proposition 5.5.2, 295
 - semisimple eigenvalue
 - Proposition 5.5.8, 296
- induced lower bound**
 - definition
 - Definition 9.5.1, 558
 - Proposition 9.5.2, 558
 - lower bound
 - Fact 9.8.43, 579
 - maximum singular value
 - Corollary 9.5.5, 560
 - minimum singular value
 - Corollary 9.5.5, 560
 - properties
 - Proposition 9.5.2, 558
 - Proposition 9.5.3, 559**
 - singular value
 - Proposition 9.5.4, 560
 - supermultiplicativity
 - Proposition 9.5.6, 560
- induced norm**
 - compatible norm
 - Proposition 9.4.3, 553
 - definition
 - Definition 9.4.1, 553
 - determinant
 - Fact 9.12.11, 601
 - dual norm
 - Fact 9.7.22, 570
 - field
 - Example 9.4.8, 554
 - maximum singular value
 - Fact 9.8.24, 575
 - norm
 - Theorem 9.4.2, 553
 - quadratic form
 - Fact 9.8.34, 577
 - spectral radius
 - Corollary 9.4.5, 554
 - Corollary 9.4.10, 556
- induced norms**
 - symmetry property
 - Fact 9.8.16, 574
- inequality**
 - elementary
 - symmetric function
 - Fact 1.15.20, 53
 - sum of products
 - Fact 1.15.20, 53
- inertia**
 - congruent matrices
 - Corollary 5.4.7, 294
 - Fact 5.8.22, 311
 - definition, 245

dissipative matrix

Fact 5.8.12, 309

generalized inverse

Fact 6.3.21, 374

Fact 8.20.12, 527

group-invertible matrix

Fact 5.8.5, 308

Hermitian matrix

Theorem 8.4.11, 428

Proposition 5.4.6, 294

Fact 5.8.6, 308

Fact 5.8.8, 308

Fact 5.8.12, 309

Fact 5.8.13, 309

Fact 5.8.14, 309

Fact 5.8.15, 309

Fact 5.8.16, 310

Fact 5.8.17, 310

Fact 5.8.18, 310

Fact 5.8.19, 310

Fact 5.12.1, 333

Fact 6.3.21, 374

Fact 8.10.15, 457

Fact 8.20.12, 527

Fact 8.20.14, 527

Fact 12.21.1, 793

Fact 12.21.2, 794

Fact 12.21.3, 794

Fact 12.21.4, 794

Fact 12.21.5, 794

Fact 12.21.6, 795

Fact 12.21.7, 795

Fact 12.21.8, 795

Fact 12.21.10, 796

Fact 12.21.11, 796

Fact 12.21.12, 796

idempotent matrix

Fact 5.8.1, 307

inequalities

Fact 5.8.16, 310

involutory matrix

Fact 5.8.2, 307

Lyapunov equation

Fact 12.21.1, 793

Fact 12.21.2, 794

Fact 12.21.3, 794

Fact 12.21.4, 794

Fact 12.21.5, 794

Fact 12.21.6, 795

Fact 12.21.7, 795

Fact 12.21.8, 795

Fact 12.21.9, 796

Fact 12.21.10, 796

Fact 12.21.11, 796

Fact 12.21.12, 796

nilpotent matrix

Fact 5.8.4, 307

normal matrix

Fact 5.10.17, 319

partitioned matrix

Fact 5.8.19, 310

Fact 5.8.20, 310

Fact 5.8.21, 311

Fact 5.12.1, 333

Fact 6.5.5, 386

positive-definite matrix

Fact 5.8.10, 308

positive-semidefinite matrix

Fact 5.8.9, 308

Fact 5.8.10, 308

Fact 12.21.9, 796

rank

Fact 5.8.5, 308

Fact 5.8.18, 310

Riccati equation

Lemma 12.16.18, 781

Schur complement

Fact 6.5.5, 386

skew-Hermitian matrix

Fact 5.8.4, 307

skew-involutory matrix

Fact 5.8.4, 307

submatrix

Fact 5.8.8, 308

tripotent matrix

Fact 5.8.3, 307

inertia matrix**positive-definite matrix**

Fact 8.9.5, 451

rigid body

Fact 8.9.5, 451

infinite finite interval

definition, xxxv

infinite matrix product

convergence

Fact 11.21.16, 715

infinite product

commutator

Fact 11.14.18, 685

convergence

Fact 11.21.16, 715

identity

Fact 1.7.10, 20

Fact 1.7.11, 20

matrix exponential

Fact 11.14.18, 685

infinite series

binomial series

Fact 1.18.8, 73

complex numbers

Fact 1.18.8, 73

Gregory's series

Fact 1.18.8, 73

identity

Fact 1.7.6, 18

Fact 1.7.7, 19

Fact 1.7.9, 19

Mercator's series

Fact 1.18.8, 73

spectral radius

Fact 10.11.24, 641

infinity norm

definition, 545

Kronecker product

Fact 9.9.61, 591

submultiplicative

norm

Fact 9.9.1, 580

Fact 9.9.2, 580

injective function

definition, 76

inner product

convex cone

Fact 10.9.13, 635

inequality

Fact 2.12.1, 126

open ball

Fact 9.7.24, 570
separation theorem
 Fact 10.9.13, 635
 Fact 10.9.14, 635
subspace
 Fact 10.9.12, 635
inner product of complex matrices
 definition, 87
inner product of complex vectors
 definition, 85
inner product of real matrices
 definition, 86
inner product of real vectors
 definition, 85
inner-product minimization
 positive-definite matrix
 Fact 8.15.12, 502
input matrix
 controllability
 Fact 12.20.15, 792
 stabilizability
 Fact 12.20.15, 792
input-to-state stability
 asymptotic stability
 Fact 12.20.18, 793
integers
 identity
 Fact 1.10.1, 30
 Fact 1.10.2, 30
integral
 asymptotically stable matrix
 Lemma 11.9.2, 667
 averaged limit
 Fact 10.11.6, 638
determinant
 Fact 11.13.15, 680
Drazin generalized inverse

Fact 11.13.12, 679
 Fact 11.13.14, 680
generalized inverse
 Fact 11.13.10, 679
group generalized inverse
 Fact 11.13.13, 680
 Fact 11.13.14, 680
inverse matrix
 Fact 11.13.11, 679
matrix
 definition, 628
matrix exponential
 Proposition 11.1.4, 645
 Lemma 11.9.2, 667
 Fact 11.13.10, 679
 Fact 11.13.11, 679
 Fact 11.13.12, 679
 Fact 11.13.13, 680
 Fact 11.13.14, 680
 Fact 11.13.15, 680
 Fact 11.14.1, 681
 Fact 11.18.5, 698
 Fact 11.18.6, 698
positive-definite matrix
 Fact 8.15.32, 505
 Fact 8.15.33, 506
 Fact 8.15.34, 506
 Fact 8.15.35, 506
positive-semidefinite matrix
 Proposition 8.6.10, 433
quadratic form
 Fact 8.15.34, 506
 Fact 8.15.35, 506
integral representation
Kronecker sum
 Fact 11.18.34, 705

interior
 boundary
 Fact 10.8.7, 632
complement
 Fact 10.8.6, 632
convex set
 Fact 10.8.8, 632
 Fact 10.8.19, 633
definition

Definition 10.1.1, 621
intersection
 Fact 10.9.2, 634
largest open set
 Fact 10.8.3, 632
simplex
 Fact 2.20.4, 154
subset
 Fact 10.9.1, 634
union
 Fact 10.9.2, 634
 Fact 10.9.3, 634
interior point
 definition
 Definition 10.1.1, 621
interior point relative to a set
 definition
 Definition 10.1.2, 621
interior relative to a set
 definition
 Definition 10.1.2, 621
interlacing
 singular value
 Fact 9.14.10, 609
interlacing theorem
 asymptotically stable polynomial
 Fact 11.17.6, 696
interpolation
 polynomial
 Fact 4.8.11, 259
intersection
 closed set
 Fact 10.9.10, 635
 Fact 10.9.11, 635
convex set
 Fact 10.9.7, 634
definition, 2
dual cone
 Fact 2.9.5, 111
equivalence relation
 Proposition 1.3.3, 5
interior
 Fact 10.9.2, 634
open set

- Fact 10.9.9, 635
- reflexive relation**
 - Proposition 1.3.3, 5
- span**
 - Fact 2.9.12, 111
- symmetric relation**
 - Proposition 1.3.3, 5
- transitive relation**
 - Proposition 1.3.3, 5
- intersection of closed sets**
 - Cantor intersection theorem**
 - Fact 10.9.11, 635
- intersection of ranges**
 - projector**
 - Fact 6.4.41, 384
- intersection of subspaces**
 - subspace dimension theorem**
 - Theorem 2.3.1, 90
- interval**
 - definition, xxxv**
- invariance of domain**
 - open set image**
 - Theorem 10.3.7, 624
- invariant subspace**
 - controllable subspace**
 - Corollary 12.6.4, 737
 - definition, 94**
 - lower triangular matrix**
 - Fact 5.9.2, 311
 - matrix representation**
 - Fact 2.9.25, 113
 - stable subspace**
 - Proposition 11.8.8, 665
 - unobservable subspace**
 - Corollary 12.3.4, 729
 - unstable subspace**
 - Proposition 11.8.8, 665
 - upper triangular matrix**
 - Fact 5.9.2, 311
- invariant zero definition**
 - Definition 12.10.1, 757
- determinant**
 - Fact 12.22.14, 800
- equivalent realizations**
 - Proposition 12.10.10, 764
- full actuation**
 - Definition 12.10.2, 758
- full observation**
 - Definition 12.10.2, 758
- full-state feedback**
 - Proposition 12.10.10, 764
 - Fact 12.22.14, 800
- observable pair**
 - Corollary 12.10.12, 765
- pencil**
 - Corollary 12.10.4, 759
 - Corollary 12.10.5, 760
 - Corollary 12.10.6, 761
- regular pencil**
 - Corollary 12.10.4, 759
 - Corollary 12.10.5, 760
 - Corollary 12.10.6, 761
- transmission zero**
 - Theorem 12.10.8, 762
 - Theorem 12.10.9, 762
- uncontrollable spectrum**
 - Theorem 12.10.9, 762
- uncontrollable-unobservable spectrum**
 - Theorem 12.10.9, 762
- unobservable eigenvalue**
 - Proposition 12.10.11, 764
- unobservable spectrum**
 - Theorem 12.10.9, 762
- inverse determinant**
 - Fact 2.13.5, 129
- left-invertible matrix**
 - Proposition 2.6.5, 101
- polynomial matrix**
 - definition, 235**
- positive-definite matrix**
 - Fact 8.11.10, 469
- rank**
 - Fact 2.11.21, 125
 - Fact 2.11.22, 125
- right-invertible matrix**
 - Proposition 2.6.5, 101
- subdeterminant**
 - Fact 2.13.5, 129
- inverse function**
 - definition, 4**
 - uniqueness**
 - Theorem 1.2.2, 4
- inverse function theorem**
 - determinant**
 - Theorem 10.4.5, 627
 - existence of local inverse**
 - Theorem 10.4.5, 627
- inverse image**
 - definition, 4**
 - subspace intersection**
 - Fact 2.9.30, 114
 - subspace sum**
 - Fact 2.9.30, 114
- inverse matrix**
 - 2×2
 - Fact 2.16.12, 143
 - 2×2 **block triangular**
 - Lemma 2.8.2, 107
 - 3×3
 - Fact 2.16.12, 143
 - asymptotically stable matrix**
 - Fact 11.18.15, 700
 - block-circulant matrix**
 - Fact 2.17.6, 148
 - block-triangular matrix**

- Fact 2.17.1, 146
- characteristic polynomial**
 - Fact 4.9.9, 261
- companion matrix**
 - Fact 5.16.2, 353
- convergent sequence**
 - Fact 2.16.29, 146
 - Fact 4.10.5, 266
- definition, 101**
- derivative**
 - Proposition 10.7.2, 630
 - Fact 10.11.19, 641
- elementary matrix**
 - Fact 2.16.1, 141
 - Fact 3.7.20, 182
- finite sequence**
 - Fact 2.16.28, 146
- Hamiltonian matrix**
 - Fact 3.19.5, 216
- Hankel matrix**
 - Fact 3.18.4, 215
- identity**
 - Fact 2.16.13, 143
 - Fact 2.16.14, 144
 - Fact 2.16.15, 144
 - Fact 2.16.16, 144
 - Fact 2.16.17, 144
 - Fact 2.16.18, 144
 - Fact 2.16.19, 144
 - Fact 2.16.20, 144
 - Fact 2.16.21, 145
 - Fact 2.16.22, 145
 - Fact 2.16.23, 145
 - Fact 2.16.24, 145
 - Fact 2.16.25, 145
 - Fact 2.16.26, 145
 - Fact 2.16.27, 146
- identity perturbation**
 - Fact 4.8.12, 259
- integral**
 - Fact 11.13.11, 679
- Kronecker product**
 - Proposition 7.1.7, 401
- lower bound**
 - Fact 8.9.17, 452
- matrix exponential**
 - Proposition 11.2.8, 649
 - Fact 11.13.11, 679
- matrix inversion lemma**
 - Corollary 2.8.8, 108
- matrix sum**
 - Corollary 2.8.10, 110
- maximum singular value**
 - Fact 9.14.8, 608
- Newton-Raphson algorithm**
 - Fact 2.16.29, 146
- normalized submultiplicative norm**
 - Fact 9.8.44, 579
 - Fact 9.9.56, 590
 - Fact 9.9.57, 590
 - Fact 9.9.58, 591
 - Fact 9.9.59, 591
- outer-product perturbation**
 - Fact 2.16.3, 141
- partitioned matrix**
 - Fact 2.16.4, 142
 - Fact 2.17.2, 146
 - Fact 2.17.3, 147
 - Fact 2.17.4, 147
 - Fact 2.17.5, 147
 - Fact 2.17.6, 148
 - Fact 2.17.8, 148
 - Fact 5.12.21, 337
- perturbation**
 - Fact 9.9.60, 591
- polynomial representation**
 - Fact 4.8.13, 259
- positive-definite matrix**
 - Proposition 8.6.6, 432
 - Lemma 8.6.5, 432
 - Fact 8.9.17, 452
 - Fact 8.9.41, 455
- positive-semidefinite matrix**
 - Fact 8.10.37, 461
- product**
 - Proposition 2.6.9, 102
- rank**
 - Fact 2.17.10, 149
- Fact 6.5.11, 388
- series**
 - Proposition 9.4.13, 557
- similar matrices**
 - Fact 5.15.31, 350
- similarity transformation**
 - Fact 5.15.4, 345
- spectral radius**
 - Proposition 9.4.13, 557
- spectrum**
 - Fact 5.11.14, 324
- sum**
 - Fact 2.17.6, 148
- tridiagonal matrix**
 - Fact 3.20.9, 219
 - Fact 3.20.10, 219
 - Fact 3.20.11, 220
- upper block-triangular matrix**
 - Fact 2.17.7, 148
 - Fact 2.17.9, 148
- inverse operation composition**
 - Fact 1.5.10, 12
- iterated**
 - Fact 1.5.9, 12
- invertible function definition, 4**
- involutory matrix commutator**
 - Fact 3.15.4, 212
- definition**
 - Definition 3.1.1, 165
- determinant**
 - Fact 3.15.1, 212
 - Fact 5.15.32, 351
- diagonalizable matrix**
 - Fact 5.14.20, 341
- factorization**
 - Fact 5.15.18, 348
 - Fact 5.15.31, 350
 - Fact 5.15.32, 351
- idempotent matrix**
 - Fact 3.15.2, 212
- identity**

Fact 3.15.3, 212
inertia
 Fact 5.8.2, 307
Kronecker product
 Fact 7.4.16, 406
matrix exponential
 Fact 11.11.1, 671
normal matrix
 Fact 5.9.9, 312
 Fact 5.9.10, 312
null space
 Fact 3.15.4, 212
partitioned matrix
 Fact 3.15.5, 212
range
 Fact 3.15.4, 212
reflector
 Fact 3.14.2, 211
semisimple matrix
 Fact 5.14.19, 341
signature
 Fact 5.8.2, 307
similar matrices
 Proposition 3.4.5, 174
 Corollary 5.5.22, 301
 Fact 5.15.31, 350
spectrum
 Proposition 5.5.21, 300
symmetric matrix
 Fact 5.15.36, 351
trace
 Fact 5.8.2, 307
transpose
 Fact 5.9.7, 312
tripotent matrix
 Fact 3.16.2, 212
unitarily similar matrices
 Proposition 3.4.5, 174
irreducible matrix
 absolute value
 Fact 3.20.4, 218
 almost nonnegative
 matrix
 Fact 11.19.2, 706
 connected graph
 Fact 4.11.2, 273
definition
 Definition 3.1.1, 165

graph
 Fact 4.11.2, 273
group generalized inverse
 Fact 6.6.20, 398
M-matrix
 Fact 4.11.10, 276
permutation matrix
 Fact 3.20.3, 217
positive matrix
 Fact 4.11.5, 273
primary circulant
 Fact 3.20.3, 217
spectral radius
 convexity
 Fact 4.11.18, 280
 spectral radius
 monotonicity
 Fact 4.11.18, 280

irreducible polynomial
 definition, 233

isomorphic groups
 symplectic group
 and unitary group
 Fact 3.21.3, 222

isomorphism
 definition
 Definition 3.3.4, 172
group
 Proposition 3.3.5, 172

J

Jacobi identity
 commutator
 Fact 2.18.4, 149

Jacobi's identity
 determinant
 Fact 2.14.28, 139
 matrix differential
 equation
 Fact 11.13.4, 678

Jacobian
 definition, 627

Jacobson

nilpotent
 commutator
 Fact 3.17.12, 214

Jensen
 convex function
 Fact 10.11.7, 638

Jensen's inequality
 arithmetic-mean–
 geometric-mean
 inequality
 Fact 1.8.4, 21
 convex function
 Fact 1.8.4, 21
 Fact 1.15.35, 57

JLL inequality
 trace of a matrix
 power
 Fact 4.11.22, 281

Jordan block
 index of an
 eigenvalue
 Proposition 5.5.3, 295

Jordan canonical form
 generalized inverse
 Fact 6.6.9, 394
 group-invertible
 matrix
 Fact 6.6.9, 394

Jordan form
 existence
 Theorem 5.3.3, 289

factorization
 Fact 5.15.5, 346

Hamiltonian
 Fact 12.23.1, 802

minimal polynomial
 Proposition 5.5.15, 299

normal matrix
 Fact 5.10.6, 317

real Jordan form
 Fact 5.10.2, 317

Schur decomposition
 Fact 5.10.6, 317

square root
 Fact 5.15.19, 348

transfer function

Fact 12.22.10, 800

Jordan matrix

example

Example 5.3.6, 290

Example 5.3.7, 291

Jordan structure

logarithm

Corollary 11.4.4, 654

matrix exponential

Corollary 11.4.4, 654

Jordan's inequality

trigonometric

inequality

Fact 1.9.29, 28

Jordan-Chevalley

decomposition

diagonalizable

matrix

Fact 5.9.3, 311

nilpotent matrix

Fact 5.9.3, 311

Joyal

polynomial root

bound

Fact 11.20.7, 710

Jury test

discrete-time

asymptotically

stable polynomial

Fact 11.20.1, 708

K

Kalman decomposition

minimal realization

Proposition 12.9.10,
753

Kantorovich inequality

positive-semidefinite

matrix

Fact 8.15.9, 501

quadratic form

Fact 8.15.9, 501

scalar case

Fact 1.15.36, 57

Kato

maximum singular

value of a matrix

difference

Fact 9.9.32, 585

kernel function

positive-semidefinite

matrix

Fact 8.8.1, 444

Fact 8.8.2, 445

Kharitonov's theorem

asymptotically stable

polynomial

Fact 11.17.13, 698

Khatri-Rao product

Kronecker product,

416

Kittaneh

Schatten norm

inequality

Fact 9.9.45, 588

Klamkin's inequality

triangle

Fact 2.20.11, 156

Klein four-group

dihedral group

Fact 3.21.7, 223

Klein's inequality

trace of a matrix

logarithm

Fact 11.14.25, 686

Kleinman

stabilization and

Gramian

Fact 12.20.17, 792

Kojima's bound

polynomial

Fact 11.20.8, 710

Kosaki

Schatten norm

inequality

Fact 9.9.45, 588

trace norm of a

matrix difference

Fact 9.9.24, 584

trace of a convex

function

Fact 8.12.33, 482

unitarily invariant

norm inequality

Fact 9.9.44, 588

Krein

inertia of a

Hermitian matrix

Fact 12.21.5, 794

Krein-Milman theorem

extreme points of a

convex set

Fact 10.8.23, 634

Kreiss matrix theorem

maximum singular

value

Fact 11.21.17, 715

Kristof

least squares and

unitary

biequivalence

Fact 9.15.6, 619

Kronecker canonical

form

pencil

Theorem 5.7.1, 304

regular pencil

Proposition 5.7.2, 305

Kronecker permutation

matrix

definition, 402

Kronecker product

Fact 7.4.29, 407

orthogonal matrix

Fact 7.4.29, 407

trace

Fact 7.4.29, 407

transpose

Proposition 7.1.13, 402

vec

Fact 7.4.29, 407

Kronecker product

biequivalent matrices

Fact 7.4.11, 405

- column norm**
 - Fact 9.9.61, 591
- complex conjugate transpose**
 - Proposition 7.1.3, 400
- congruent matrices**
 - Fact 7.4.12, 406
- convex function**
 - Proposition 8.6.17, 437
- definition**
 - Definition 7.1.2, 400
- determinant**
 - Proposition 7.1.11, 402
 - Fact 7.5.12, 410
 - Fact 7.5.13, 410
- diagonal matrix**
 - Fact 7.4.3, 405
- discrete-time**
 - asymptotically stable matrix**
 - Fact 11.21.5, 713
 - Fact 11.21.6, 713
- discrete-time Lyapunov-stable matrix**
 - Fact 11.21.5, 713
 - Fact 11.21.6, 713
- discrete-time semistable matrix**
 - Fact 11.21.5, 713
 - Fact 11.21.6, 713
- Drazin generalized inverse**
 - Fact 7.4.31, 408
- eigenvalue**
 - Proposition 7.1.10, 401
 - Fact 7.4.13, 406
 - Fact 7.4.15, 406
 - Fact 7.4.21, 406
 - Fact 7.4.28, 407
 - Fact 7.4.32, 408
- eigenvector**
 - Proposition 7.1.10, 401
 - Fact 7.4.21, 406
 - Fact 7.4.32, 408
- Euclidean norm**
 - Fact 9.7.27, 570
- Frobenius norm**
 - Fact 9.14.37, 617
- generalized inverse**
 - Fact 7.4.30, 408
- group generalized inverse**
 - Fact 7.4.31, 408
- group-invertible matrix**
 - Fact 7.4.16, 406
 - Fact 7.4.31, 408
- Hermitian matrix**
 - Fact 7.4.16, 406
 - Fact 8.21.28, 536
- Hölder norm**
 - Fact 9.9.61, 591
- idempotent matrix**
 - Fact 7.4.16, 406
- index of a matrix**
 - Fact 7.4.26, 407
- infinity norm**
 - Fact 9.9.61, 591
- inverse matrix**
 - Proposition 7.1.7, 401
- involutory matrix**
 - Fact 7.4.16, 406
- Kronecker permutation matrix**
 - Fact 7.4.29, 407
- Kronecker sum**
 - Fact 11.14.37, 688
- left-equivalent matrices**
 - Fact 7.4.11, 405
- lower triangular matrix**
 - Fact 7.4.3, 405
- matrix exponential**
 - Proposition 11.1.7, 645
 - Fact 11.14.37, 688
 - Fact 11.14.38, 688
- matrix multiplication**
 - Proposition 7.1.6, 400
- matrix power**
 - Fact 7.4.4, 405
 - Fact 7.4.10, 405
 - Fact 7.4.21, 406
- matrix sum**
 - Proposition 7.1.4, 400
- maximum singular value**
 - Fact 9.14.37, 617
- nilpotent matrix**
 - Fact 7.4.16, 406
- normal matrix**
 - Fact 7.4.16, 406
- orthogonal matrix**
 - Fact 7.4.16, 406
- outer-product matrix**
 - Proposition 7.1.8, 401
- partitioned matrix**
 - Fact 7.4.18, 406
 - Fact 7.4.19, 406
 - Fact 7.4.24, 407
- positive-definite matrix**
 - Fact 7.4.16, 406
- positive-semidefinite matrix**
 - Fact 7.4.16, 406
 - Fact 8.21.16, 534
 - Fact 8.21.22, 536
 - Fact 8.21.23, 536
 - Fact 8.21.24, 536
 - Fact 8.21.26, 536
 - Fact 8.21.27, 536
 - Fact 8.21.29, 536
- projector**
 - Fact 7.4.16, 406
- range**
 - Fact 7.4.22, 407
- range-Hermitian matrix**
 - Fact 7.4.16, 406
- rank**
 - Fact 7.4.23, 407
 - Fact 7.4.24, 407
 - Fact 7.4.25, 407
 - Fact 8.21.16, 534
- reflector**
 - Fact 7.4.16, 406
- right-equivalent matrices**
 - Fact 7.4.11, 405
- row norm**
 - Fact 9.9.61, 591
- Schatten norm**

- Fact 9.14.37, 617
 - Schur product**
 - Proposition 7.3.1, 404
 - semisimple matrix**
 - Fact 7.4.16, 406
 - similar matrices**
 - Fact 7.4.12, 406
 - singular matrix**
 - Fact 7.4.27, 407
 - skew-Hermitian matrix**
 - Fact 7.4.17, 406
 - spectral radius**
 - Fact 7.4.14, 406
 - square root**
 - Fact 8.21.29, 536
 - Fact 8.21.30, 537
 - submatrix**
 - Proposition 7.3.1, 404
 - trace**
 - Proposition 7.1.12, 402
 - Fact 11.14.38, 688
 - transpose**
 - Proposition 7.1.3, 400
 - triple product**
 - Proposition 7.1.5, 400
 - Fact 7.4.7, 405
 - tripotent matrix**
 - Fact 7.4.16, 406
 - unitarily similar matrices**
 - Fact 7.4.12, 406
 - unitary matrix**
 - Fact 7.4.16, 406
 - upper triangular matrix**
 - Fact 7.4.3, 405
 - vec**
 - Fact 7.4.5, 405
 - Fact 7.4.6, 405
 - Fact 7.4.8, 405
 - vector**
 - Fact 7.4.1, 405
 - Fact 7.4.2, 405
 - Fact 7.4.20, 406
 - Kronecker sum**
 - associativity**
 - Proposition 7.2.2, 403
 - asymptotically stable matrix**
 - Fact 11.18.32, 704
 - Fact 11.18.33, 704
 - Fact 11.18.34, 705
 - asymptotically stable polynomial**
 - Fact 11.17.11, 697
 - commuting matrices**
 - Fact 7.5.4, 409
 - defect**
 - Fact 7.5.2, 409
 - definition**
 - Definition 7.2.1, 403
 - determinant**
 - Fact 7.5.11, 410
 - dissipative matrix**
 - Fact 7.5.8, 409
 - eigenvalue**
 - Proposition 7.2.3, 403
 - Fact 7.5.5, 409
 - Fact 7.5.7, 409
 - Fact 7.5.16, 411
 - eigenvector**
 - Proposition 7.2.3, 403
 - Fact 7.5.16, 411
 - Hermitian matrix**
 - Fact 7.5.8, 409
 - integral representation**
 - Fact 11.18.34, 705
 - Kronecker product**
 - Fact 11.14.37, 688
 - linear matrix equation**
 - Proposition 11.9.3, 667
 - linear system**
 - Fact 7.5.15, 411
 - Lyapunov equation**
 - Corollary 11.9.4, 668
 - Lyapunov-stable matrix**
 - Fact 11.18.32, 704
 - Fact 11.18.33, 704
 - matrix exponential**
 - Proposition 11.1.7, 645
 - Fact 11.14.36, 688
 - Fact 11.14.37, 688
 - matrix power**
 - Fact 7.5.1, 409
 - nilpotent matrix**
 - Fact 7.5.3, 409
 - Fact 7.5.8, 409
 - normal matrix**
 - Fact 7.5.8, 409
 - positive matrix**
 - Fact 7.5.8, 409
 - positive-semidefinite matrix**
 - Fact 7.5.8, 409
 - range-Hermitian matrix**
 - Fact 7.5.8, 409
 - rank**
 - Fact 7.5.2, 409
 - Fact 7.5.9, 409
 - Fact 7.5.10, 410
 - semidissipative matrix**
 - Fact 7.5.8, 409
 - semistable matrix**
 - Fact 11.18.32, 704
 - Fact 11.18.33, 704
 - similar matrices**
 - Fact 7.5.9, 409
 - skew-Hermitian matrix**
 - Fact 7.5.8, 409
 - spectral abscissa**
 - Fact 7.5.6, 409
 - trace**
 - Fact 11.14.36, 688
- L**
- L₂ norm**
 - controllability Gramian**
 - Theorem 12.11.1, 765
 - definition, 765**
 - observability Gramian**
 - Theorem 12.11.1, 765
 - Löwner-Heinz inequality**
 - positive-semidefinite matrix inequality**

Corollary 8.6.11, 434

Labelle

polynomial root

bound

Fact 11.20.7, 710

Laffey

simultaneous

triangularization

Fact 5.17.5, 358

Lagrange identity

product identity

Fact 1.16.8, 61

Lagrange interpolation

formula

polynomial

interpolation

Fact 4.8.11, 259

Lagrange-Hermite

interpolation

polynomial

matrix function

Theorem 10.5.2, 629

Laguerre-Samuelson

inequality

mean

Fact 1.15.12, 51

Fact 8.9.35, 454

Lancaster's formulas

quadratic form

integral

Fact 8.15.34, 506

Laplace transform

matrix exponential

Proposition 11.2.2, 647

resolvent

Proposition 11.2.2, 647

Laplacian

symmetric graph

Fact 4.11.1, 272

Laplacian matrix

adjacency matrix

Theorem 3.2.2, 170

Theorem 3.2.3, 171

Fact 4.11.11, 277

definition

Definition 3.2.1, 170

incidence matrix

Theorem 3.2.2, 170

Theorem 3.2.3, 171

quadratic form

Fact 8.15.36, 506

spectrum

Fact 11.19.7, 708

symmetric graph

Fact 8.15.36, 506

lattice

definition

Definition 1.3.9, 7

positive-semidefinite

matrix

Fact 8.10.32, 459

Fact 8.10.33, 459

leading principal

submatrix

definition, 80

leaf

Definition 1.4.2, 8

least common multiple

block-diagonal

matrix

Lemma 5.2.7, 286

definition, 234

least lower bound for a

partial ordering

definition

Definition 1.3.9, 7

least squares

fixed-rank

approximation

Fact 9.14.28, 614

Fact 9.15.4, 618

generalized inverse

Fact 9.15.1, 618

Fact 9.15.2, 618

Fact 9.15.3, 618

singular value

decomposition

Fact 9.14.28, 614

Fact 9.15.4, 618

Fact 9.15.5, 618

Fact 9.15.6, 619

least squares and

unitary

biequivalence

Kristof

Fact 9.15.6, 619

least upper bound

projector

Fact 6.4.41, 385

left divides

definition, 234

left equivalence

equivalence relation

Fact 5.10.3, 317

left inverse

(1)-inverse

Proposition 6.1.3, 364

affine subspace

Fact 2.9.26, 113

complex conjugate

transpose

Fact 2.15.1, 140

Fact 2.15.2, 140

cone

Fact 2.9.26, 113

convex set

Fact 2.9.26, 113

definition, 4

generalized inverse

Corollary 6.1.4, 364

Fact 6.4.39, 384

Fact 6.4.40, 384

idempotent matrix

Fact 3.12.10, 199

left-inner matrix

Fact 3.11.5, 190

matrix product

Fact 2.15.5, 141

positive-definite

matrix

Fact 3.7.25, 182

representation

Fact 2.15.3, 140

subspace

Fact 2.9.26, 113

uniqueness

Theorem 1.2.2, 4

left-equivalent matrices

definition

Definition 3.4.3, 174

group-invertible

matrix

Fact 3.6.1, 177

Kronecker product

Fact 7.4.11, 405

null space

Proposition 5.1.3, 283

positive-semidefinite matrix

Fact 5.10.19, 319

left-inner matrix

definition

Definition 3.1.2, 166

generalized inverse

Fact 6.3.8, 371

left inverse

Fact 3.11.5, 190

left-invertible function

definition, 4

left-invertible matrix

definition, 98

equivalent properties

Theorem 2.6.1, 98

generalized inverse

Proposition 6.1.5, 364

inverse

Proposition 2.6.5, 101

matrix product

Fact 2.10.3, 115

nonsingular

equivalence

Corollary 2.6.6, 101

unique left inverse

Proposition 2.6.2, 99

Lehmer matrix

positive-semidefinite matrix

Fact 8.8.5, 447

Lehmer mean

power inequality

Fact 1.10.35, 36

Leibniz's rule

derivative of an integral

Fact 10.11.10, 639

lemma

definition, 1

Leslie matrix

definition, 362

Leverrier's algorithm

characteristic

polynomial

Proposition 4.4.9, 244

lexicographic ordering

cone

Fact 2.9.31, 115

total ordering

Fact 1.5.8, 12

Lidskii-Mirsky-Wielandt theorem

Hermitian

perturbation

Fact 9.12.4, 599

Lidskii-Wielandt

inequalities

eigenvalue inequality for Hermitian matrices

Fact 8.18.3, 513

Lie algebra

classical examples

Proposition 3.3.2, 171

definition

Definition 3.3.1, 171

Lie group

Proposition 11.6.4, 658

Proposition 11.6.5, 659

Proposition 11.6.6, 659

matrix exponential

Proposition 11.6.7, 659

strictly upper

triangular matrix

Fact 3.21.4, 222

Fact 11.22.1, 715

upper triangular

matrix

Fact 3.21.4, 222

Fact 11.22.1, 715

Lie algebra of a Lie group

matrix exponential

Proposition 11.6.3, 658

Lie group

definition

Definition 11.6.1, 658

group

Proposition 11.6.2, 658

Lie algebra

Proposition 11.6.4, 658

Proposition 11.6.5, 659

Proposition 11.6.6, 659

Lie-Trotter formula

matrix exponential

Fact 11.14.7, 683

Lie-Trotter product formula

matrix exponential

Corollary 11.4.8, 656

Fact 11.16.2, 692

Fact 11.16.3, 692

Lieb concavity

theorem, 542

Lieb-Thirring inequality

positive-semidefinite

matrix

Fact 8.12.22, 480

Fact 8.18.20, 518

limit

discrete-time

semistable matrix

Fact 11.21.10, 714

Drazin generalized

inverse

Fact 6.6.11, 395

Hermitian matrix

Fact 8.10.1, 456

matrix exponential

Fact 11.18.5, 698

Fact 11.18.6, 698

Fact 11.18.7, 699

matrix logarithm

Proposition 8.6.4, 432

positive-definite

matrix

Fact 8.10.47, 465
positive-semidefinite matrix
 Proposition 8.6.3, 432
 Fact 8.10.47, 465
projector
 Fact 6.4.41, 384
 Fact 6.4.46, 385
semistable matrix
 Fact 11.18.7, 699

Linden

polynomial root bound
 Fact 11.20.9, 710

linear combination

determinant
 Fact 8.13.18, 488

Hermitian matrix

Fact 8.15.24, 504
 Fact 8.15.25, 504
 Fact 8.15.26, 504

idempotent matrix

Fact 5.19.9, 361

positive-semidefinite matrix

Fact 8.13.18, 488

linear combination of projectors

Hermitian matrix
 Fact 5.19.10, 361

linear combination of two vectors

definition, 79

linear constraint

quadratic form
 Fact 8.14.10, 497

linear dependence

absolute value
 Fact 9.7.1, 563
triangle inequality
 Fact 9.7.3, 563

linear dependence of two matrices

definition, 80

linear dependence of two vectors

definition, 79

linear dependence of vectors

definition, 90

linear dynamical system

asymptotically stable
 Proposition 11.8.2, 662

discrete-time asymptotically stable
 Proposition 11.10.2, 670

discrete-time Lyapunov stable
 Proposition 11.10.2, 670

discrete-time semistable
 Proposition 11.10.2, 670

discrete-time semistable
 Proposition 11.10.2, 670
Lyapunov stable
 Proposition 11.8.2, 662
semistable
 Proposition 11.8.2, 662

linear function

continuous function
 Corollary 10.3.3, 624
definition, 81

linear independence

cyclic matrix
 Fact 5.14.9, 340
definition, 90
outer-product matrix
 Fact 2.12.8, 126

linear matrix equation

asymptotically stable matrix
 Proposition 11.9.3, 667
existence of solutions
 Fact 5.10.20, 320
 Fact 5.10.21, 320
generalized inverse
 Fact 6.4.38, 384

Kronecker sum

Proposition 11.9.3, 667

matrix exponential

Proposition 11.9.3, 667

rank

Fact 2.10.16, 117

skew-symmetric matrix

Fact 3.7.3, 178

solution

Fact 6.4.38, 384

Sylvester's equation

Proposition 7.2.4, 403
 Proposition 11.9.3, 667
 Fact 5.10.20, 320
 Fact 5.10.21, 320
 Fact 6.5.7, 387

symmetric matrix

Fact 3.7.3, 178

linear system

generalized inverse
 Proposition 6.1.7, 366
harmonic steady-state response
 Theorem 12.12.1, 768

Kronecker sum

Fact 7.5.15, 411

right inverse

Fact 6.3.1, 369

solutions

Proposition 6.1.7, 366
 Fact 2.10.6, 116

linear system solution

Cramer's rule
 Fact 2.13.6, 129

nonnegative vector

Fact 4.11.14, 279

norm

Fact 9.9.64, 592
 Fact 9.9.65, 592
 Fact 9.9.66, 592

rank

Theorem 2.6.4, 100
 Corollary 2.6.7, 101

right-invertible matrix

Fact 2.13.7, 129

linear-quadratic control problem

definition, 775

Riccati equation

Theorem 12.15.2, 776

solution

Theorem 12.15.2, 776

linearly independent rational functions

definition, 250

Littlewood

Hölder-induced norm

Fact 9.8.17, 574

Fact 9.8.18, 574

Ljance

minimal principal

angle and subspaces

Fact 5.11.39, 329

log majorization

convex function

Fact 2.21.12, 163

increasing function

Fact 2.21.12, 163

positive-semidefinite matrix

Fact 8.11.9, 469

logarithm, see matrix logarithm

SO(3)

Fact 11.15.10, 692

convex function

Fact 11.16.14, 695

Fact 11.16.15, 695

determinant

Fact 8.13.8, 486

determinant and convex function

Proposition 8.6.17, 437

entropy

Fact 1.15.45, 59

Fact 1.15.46, 59

Fact 1.15.47, 59

Fact 1.16.30, 67

Euler constant

Fact 1.7.5, 18

gamma

Fact 1.7.5, 18

increasing function

Proposition 8.6.13, 435

inequality

Fact 1.15.45, 59

Fact 1.15.46, 59

Fact 1.15.47, 59

Jordan structure

Corollary 11.4.4, 654

orthogonal matrix

Fact 11.15.10, 692

rotation matrix

Fact 11.15.10, 692

scalar inequalities

Fact 1.9.21, 26

Fact 1.9.22, 26

Fact 1.9.23, 27

Fact 1.9.24, 27

Fact 1.9.25, 27

Fact 1.10.24, 34

Fact 1.10.25, 34

Fact 1.10.40, 38

Shannon's inequality

Fact 1.16.30, 67

trace and convex function

Proposition 8.6.17, 437

logarithm function

complex numbers

Fact 1.18.7, 72

principal branch

Fact 1.18.7, 72

scalar inequalities

Fact 1.9.26, 27

Fact 1.9.27, 27

Fact 1.9.28, 27

logarithmic derivative

asymptotically stable matrix

Fact 11.18.11, 699

Lyapunov equation

Fact 11.18.11, 699

properties

Fact 11.15.7, 690

logarithmic mean

arithmetic mean

Fact 1.15.26, 54

Heron mean

Fact 1.10.37, 37

identric mean

Fact 1.10.36, 37

Polya's inequality

Fact 1.10.36, 37

logical equivalents

De Morgan's laws

Fact 1.5.1, 10

existential statement

Fact 1.5.4, 11

implication

Fact 1.5.1, 10

Fact 1.5.2, 10

Fact 1.5.3, 11

universal statement

Fact 1.5.4, 11

loop

Definition 1.4.2, 8

lower block-triangular matrix

definition

Definition 3.1.3, 167

determinant

Proposition 2.7.1, 103

lower bound

induced lower bound

Fact 9.8.43, 579

minimum singular value

Fact 9.13.15, 604

Fact 9.13.21, 606

lower bound for a partial ordering

definition

Definition 1.3.9, 7

lower Hessenberg matrix

definition

Definition 3.1.3, 167

lower

reverse-triangular matrix

definition

Fact 2.13.8, 130

determinant

Fact 2.13.8, 130

lower triangular matrix**commutator**

Fact 3.17.11, 214

definition

Definition 3.1.3, 167

factorization

Fact 5.15.10, 346

invariant subspace

Fact 5.9.2, 311

Kronecker product

Fact 7.4.3, 405

matrix exponential

Fact 11.13.1, 677

Fact 11.13.16, 680

matrix power

Fact 3.18.7, 216

matrix product

Fact 3.20.18, 221

nilpotent matrix

Fact 3.17.11, 214

similar matrices

Fact 5.9.2, 311

Toeplitz matrix

Fact 3.18.7, 216

Fact 11.13.1, 677

LQG controller**continuous-time****control problem**

Fact 12.23.6, 804

discrete-time control**problem**

Fact 12.23.7, 804

dynamic**compensator**

Fact 12.23.6, 804

Fact 12.23.7, 804

LU decomposition**existence**

Fact 5.15.10, 346

Lucas numbers**nonnegative matrix**

Fact 4.11.12, 277

Lukes**stabilization and
Gramian**

Fact 12.20.17, 792

LULU decomposition**factorization**

Fact 5.15.11, 346

Lyapunov equation**asymptotic stability**

Corollary 11.9.1, 666

**asymptotically stable
matrix**

Proposition 11.9.5, 668

Corollary 11.9.4, 668

Corollary 11.9.7, 669

Corollary 12.4.4, 734

Corollary 12.5.6, 735

Corollary 12.7.4, 746

Corollary 12.8.6, 749

Fact 12.21.7, 795

Fact 12.21.17, 797

controllably**asymptotically****stable**

Proposition 12.7.3, 743

detectability

Corollary 12.5.6, 735

discrete-time**asymptotically****stable matrix**

Proposition 11.10.5,

671

eigenvalue inclusion**region**

Fact 12.21.20, 798

finite-sum solution

Fact 12.21.17, 797

inertia

Fact 12.21.1, 793

Fact 12.21.2, 794

Fact 12.21.3, 794

Fact 12.21.4, 794

Fact 12.21.5, 794

Fact 12.21.6, 795

Fact 12.21.7, 795

Fact 12.21.8, 795

Fact 12.21.9, 796

Fact 12.21.10, 796

Fact 12.21.11, 796

Fact 12.21.12, 796

Kronecker sum

Corollary 11.9.4, 668

logarithmic**derivative**

Fact 11.18.11, 699

Lyapunov stability

Corollary 11.9.1, 666

Lyapunov-stable**matrix**

Proposition 11.9.6, 669

Corollary 11.9.7, 669

matrix exponential

Corollary 11.9.4, 668

Fact 11.18.18, 701

Fact 11.18.19, 701

null space

Fact 12.21.15, 797

observability matrix

Fact 12.21.15, 797

observably**asymptotically****stable**

Proposition 12.4.3, 732

positive-definite**matrix**

Fact 12.21.16, 797

Fact 12.21.18, 797

positive-semidefinite**matrix**

Fact 12.21.15, 797

Fact 12.21.19, 797

Schur power

Fact 8.8.16, 449

semistability

Corollary 11.9.1, 666

semistable matrix

Fact 12.21.15, 797

skew-Hermitian**matrix**

Fact 11.18.12, 700

stabilizability

Corollary 12.8.6, 749

Lyapunov stability**eigenvalue**

Proposition 11.8.2, 662

linear dynamical**system**

Proposition 11.8.2, 662

Lyapunov equation

Corollary 11.9.1, 666

matrix exponential

Proposition 11.8.2, 662

nonlinear system

Theorem 11.7.2, 661

Lyapunov's direct method

stability theory

Theorem 11.7.2, 661

Lyapunov-stable equilibrium

definition

Definition 11.7.1, 660

Lyapunov-stable matrix

almost nonnegative matrix

Fact 11.19.4, 706

compartmental matrix

Fact 11.19.6, 707

definition

Definition 11.8.1, 662

group-invertible matrix

Fact 11.18.2, 698

Kronecker sum

Fact 11.18.32, 704

Fact 11.18.33, 704

Lyapunov equation

Proposition 11.9.6, 669

Corollary 11.9.7, 669

Lyapunov-stable polynomial

Proposition 11.8.4, 663

matrix exponential

Fact 11.18.6, 698

Fact 11.21.7, 713

minimal realization

Definition 12.9.17, 757

N-matrix

Fact 11.19.4, 706

normal matrix

Fact 11.18.37, 705

positive-definite matrix

Proposition 11.9.6, 669

Corollary 11.9.7, 669

semidissipative matrix

Fact 11.18.37, 705

semistable matrix

Fact 11.18.1, 698

similar matrices

Fact 11.18.4, 698

step response

Fact 12.20.1, 790

Lyapunov-stable polynomial

definition

Definition 11.8.3, 663

Lyapunov-stable matrix

Proposition 11.8.4, 663

subdeterminant

Fact 11.18.23, 702

Lyapunov-stable transfer function

minimal realization

Proposition 12.9.18, 757

SISO entries

Proposition 12.9.19, 757

M

M-matrix

definition

Fact 4.11.6, 275

determinant

Fact 4.11.8, 276

eigenvector

Fact 4.11.10, 276

inverse

Fact 4.11.8, 276

irreducible matrix

Fact 4.11.10, 276

nonnegative matrix

Fact 4.11.6, 275

rank

Fact 8.7.7, 444

Schur product

Fact 7.6.15, 415

submatrix

Fact 4.11.7, 276

Z-matrix

Fact 4.11.6, 275

Fact 4.11.8, 276

Magnus

determinant

identities

Fact 2.13.16, 132

Magnus expansion

time-varying

dynamics

Fact 11.13.4, 678

Makelainen

difference of

idempotent

matrices

Fact 5.12.19, 337

Maligranda inequality

complex numbers

Fact 1.18.5, 71

norm

Fact 9.7.10, 566

Fact 9.7.13, 567

Mann

positivity of a quadratic form on a subspace

Fact 8.15.27, 504

Marcus

quadratic form

inequality

Fact 8.15.19, 503

similar matrices and nonzero diagonal entries

Fact 5.9.14, 313

Markov block-Hankel matrix

controllable pair

Proposition 12.9.11, 754

definition, 754

minimal realization

Proposition 12.9.12, 755

observable pair

Proposition 12.9.11, 754

rational transfer function

- Proposition 12.9.11, 754
- Proposition 12.9.12, 755
- Proposition 12.9.13, 755
- Markov parameter**
 - definition, 727
 - rational transfer function
 - Proposition 12.9.7, 751
- Martins's inequality**
 - sum of integers
 - Fact 1.9.31, 30
- Mason**
 - polynomial root bound
 - Fact 11.20.10, 711
- mass**
 - definition, 654
- mass matrix**
 - partitioned matrix
 - Fact 5.12.21, 337
- mass-spring system**
 - spectrum
 - Fact 5.12.21, 337
 - stability
 - Fact 11.18.38, 705
- Massera-Schaffer inequality**
 - complex numbers
 - Fact 1.18.5, 71
 - norm
 - Fact 9.7.10, 566
 - Fact 9.7.13, 567
- matricial norm**
 - partitioned matrix
 - Fact 9.10.1, 593
- matrix**
 - definition, 79
- matrix cosine**
 - matrix exponential
 - Fact 11.12.1, 677
 - matrix sine
 - Fact 11.12.1, 677
- matrix derivative**
 - definition, 630
- matrix differential equation**
 - Jacobi's identity
 - Fact 11.13.4, 678
 - matrix exponential
 - Fact 11.13.3, 677
 - Riccati differential equation
 - Fact 12.23.5, 803
 - time-varying dynamics
 - Fact 11.13.4, 678
 - Fact 11.13.5, 678
- matrix exponential**
 - 2×2 matrix
 - Proposition 11.3.2, 651
 - Corollary 11.3.3, 652
 - Lemma 11.3.1, 651
 - Example 11.3.4, 652
 - Example 11.3.5, 652
 - 3×3 matrix
 - Fact 11.11.5, 673
 - 3×3 orthogonal matrix
 - Fact 11.11.10, 674
 - Fact 11.11.11, 674
 - 3×3 skew-symmetric matrix
 - Fact 11.11.6, 673
 - Fact 11.11.10, 674
 - Fact 11.11.11, 674
 - 4×4 skew-symmetric matrix
 - Fact 11.11.14, 675
 - Fact 11.11.15, 675
 - Fact 11.11.16, 676
 - Fact 11.11.17, 676
 - $SO(n)$
 - Fact 11.11.3, 672
 - almost nonnegative matrix
 - Fact 11.19.1, 706
 - Fact 11.19.2, 706
 - asymptotic stability
 - Proposition 11.8.2, 662
- asymptotically stable matrix**
 - Lemma 11.9.2, 667
 - Fact 11.18.8, 699
 - Fact 11.18.9, 699
 - Fact 11.18.10, 699
 - Fact 11.18.15, 700
 - Fact 11.18.18, 701
 - Fact 11.18.19, 701
 - Fact 11.21.7, 713
- block-diagonal matrix**
 - Proposition 11.2.8, 649
- commutator**
 - Fact 11.14.9, 683
 - Fact 11.14.11, 683
 - Fact 11.14.12, 683
 - Fact 11.14.13, 684
 - Fact 11.14.14, 684
 - Fact 11.14.15, 684
 - Fact 11.14.16, 684
 - Fact 11.14.17, 684
 - Fact 11.14.18, 685
- commuting matrices**
 - Proposition 11.1.5, 645
 - Corollary 11.1.6, 645
 - Fact 11.14.2, 681
 - Fact 11.14.5, 682
- complex conjugate**
 - Proposition 11.2.8, 649
- complex conjugate transpose**
 - Proposition 11.2.8, 649
 - Fact 11.15.4, 689
 - Fact 11.15.6, 690
- convergence in time**
 - Proposition 11.8.7, 665
- convergent sequence**
 - Proposition 11.1.3, 644
 - Fact 11.14.7, 683
 - Fact 11.14.8, 683
 - Fact 11.14.9, 683
 - Fact 11.21.14, 714
- convergent series**
 - Proposition 11.1.2, 644
- convex function**
 - Fact 8.14.18, 500
 - Fact 11.16.14, 695
 - Fact 11.16.15, 695

- cross product**
 - Fact 11.11.7, 673
 - Fact 11.11.8, 674
 - Fact 11.11.9, 674
- cross-product matrix**
 - Fact 11.11.6, 673
 - Fact 11.11.12, 674
 - Fact 11.11.13, 675
 - Fact 11.11.16, 676
 - Fact 11.11.17, 676
- definition**
 - Definition 11.1.1, 643
- derivative**
 - Fact 8.12.31, 482
 - Fact 11.14.3, 682
 - Fact 11.14.4, 682
 - Fact 11.14.10, 683
 - Fact 11.15.2, 689
- derivative of a matrix**
 - Fact 11.14.11, 683
- determinant**
 - Proposition 11.4.6, 655
 - Corollary 11.2.4, 648
 - Corollary 11.2.5, 648
 - Fact 11.13.15, 680
 - Fact 11.15.5, 689
- diagonal matrix**
 - Fact 11.13.16, 680
- discrete-time**
 - asymptotic stability**
 - Proposition 11.10.2, 670
- discrete-time**
 - asymptotically stable matrix**
 - Fact 11.21.7, 713
- discrete-time Lyapunov stability**
 - Proposition 11.10.2, 670
- discrete-time Lyapunov-stable matrix**
 - Fact 11.21.7, 713
- discrete-time semistability**
 - Proposition 11.10.2, 670
- discrete-time semistable matrix**
 - Fact 11.21.7, 713
 - Fact 11.21.14, 714
- dissipative matrix**
 - Fact 11.15.3, 689
- Drazin generalized inverse**
 - Fact 11.13.12, 679
 - Fact 11.13.14, 680
- eigenstructure**
 - Proposition 11.2.7, 648
- Frobenius norm**
 - Fact 11.14.32, 688
 - Fact 11.15.3, 689
- generalized inverse**
 - Fact 11.13.10, 679
- geometric mean**
 - Fact 8.10.44, 464
- Golden-Thompson inequality**
 - Fact 11.14.28, 687
 - Fact 11.16.4, 692
- group**
 - Proposition 11.6.7, 659
- group generalized inverse**
 - Fact 11.13.13, 680
 - Fact 11.13.14, 680
 - Fact 11.18.5, 698
 - Fact 11.18.6, 698
- group-invertible matrix**
 - Fact 11.18.14, 700
- Hamiltonian matrix**
 - Proposition 11.6.7, 659
- Hermitian matrix**
 - Proposition 11.2.8, 649
 - Proposition 11.2.9, 650
 - Proposition 11.4.5, 654
 - Corollary 11.2.6, 648
 - Fact 11.14.7, 683
 - Fact 11.14.8, 683
 - Fact 11.14.21, 685
 - Fact 11.14.28, 687
 - Fact 11.14.29, 687
 - Fact 11.14.31, 688
 - Fact 11.14.32, 688
 - Fact 11.14.34, 688
- Fact 11.15.1, 689**
- Fact 11.16.4, 692**
- Fact 11.16.5, 694**
- Fact 11.16.13, 695**
- Fact 11.16.17, 695**
- idempotent matrix**
 - Fact 11.11.1, 671
 - Fact 11.16.12, 695
- infinite product**
 - Fact 11.14.18, 685
- integral**
 - Proposition 11.1.4, 645
 - Lemma 11.9.2, 667
 - Fact 11.13.10, 679
 - Fact 11.13.11, 679
 - Fact 11.13.12, 679
 - Fact 11.13.13, 680
 - Fact 11.13.14, 680
 - Fact 11.13.15, 680
 - Fact 11.14.1, 681
 - Fact 11.16.8, 694
 - Fact 11.18.5, 698
 - Fact 11.18.6, 698
- inverse matrix**
 - Proposition 11.2.8, 649
 - Fact 11.13.11, 679
- involutory matrix**
 - Fact 11.11.1, 671
- Jordan structure**
 - Corollary 11.4.4, 654
- Kronecker product**
 - Proposition 11.1.7, 645
 - Fact 11.14.37, 688
 - Fact 11.14.38, 688
- Kronecker sum**
 - Proposition 11.1.7, 645
 - Fact 11.14.36, 688
 - Fact 11.14.37, 688
- Laplace transform**
 - Proposition 11.2.2, 647
- Lie algebra**
 - Proposition 11.6.7, 659
- Lie algebra of a Lie group**
 - Proposition 11.6.3, 658
- Lie-Trotter formula**
 - Fact 11.14.7, 683
- Lie-Trotter product formula**

- Corollary 11.4.8, 656
- Fact 11.16.2, 692
- Fact 11.16.3, 692
- limit**
 - Fact 11.18.5, 698
 - Fact 11.18.6, 698
 - Fact 11.18.7, 699
- linear matrix equation**
 - Proposition 11.9.3, 667
- logarithm**
 - Fact 11.14.21, 685
- lower triangular matrix**
 - Fact 11.13.1, 677
 - Fact 11.13.16, 680
- Lyapunov equation**
 - Corollary 11.9.4, 668
 - Fact 11.18.18, 701
 - Fact 11.18.19, 701
- Lyapunov stability**
 - Proposition 11.8.2, 662
- Lyapunov-stable matrix**
 - Fact 11.18.6, 698
 - Fact 11.21.7, 713
- matrix cosine**
 - Fact 11.12.1, 677
- matrix differential equation**
 - Fact 11.13.3, 677
- matrix logarithm**
 - Theorem 11.5.1, 656
 - Proposition 11.4.2, 654
 - Fact 11.13.17, 680
 - Fact 11.14.31, 688
- matrix power**
 - Fact 11.13.19, 680
- matrix sine**
 - Fact 11.12.1, 677
- maximum eigenvalue**
 - Fact 11.16.4, 692
- maximum singular value**
 - Fact 11.15.1, 689
 - Fact 11.15.2, 689
 - Fact 11.15.5, 689
 - Fact 11.16.6, 694
 - Fact 11.16.10, 694
- nilpotent matrix**
 - Fact 11.11.1, 671
 - Fact 11.13.17, 680
- nondecreasing function**
 - Fact 8.10.44, 464
- norm**
 - Fact 11.16.9, 694
 - Fact 11.16.11, 694
 - Fact 11.16.12, 695
- norm bound**
 - Fact 11.18.10, 699
- normal matrix**
 - Proposition 11.2.8, 649
 - Fact 11.13.18, 680
 - Fact 11.14.5, 682
 - Fact 11.16.10, 694
- orthogonal matrix**
 - Proposition 11.6.7, 659
 - Fact 11.11.6, 673
 - Fact 11.11.7, 673
 - Fact 11.11.8, 674
 - Fact 11.11.9, 674
 - Fact 11.11.12, 674
 - Fact 11.11.13, 675
 - Fact 11.15.10, 692
- outer-product matrix**
 - Fact 11.11.1, 671
- partitioned matrix**
 - Fact 11.11.2, 672
 - Fact 11.14.1, 681
- Peierls-Bogoliubov inequality**
 - Fact 11.14.29, 687
- polar decomposition**
 - Fact 11.13.9, 679
- polynomial matrix**
 - Proposition 11.2.1, 646
- positive-definite matrix**
 - Proposition 11.2.8, 649
 - Proposition 11.2.9, 650
 - Fact 11.14.20, 685
 - Fact 11.14.22, 685
 - Fact 11.14.23, 686
 - Fact 11.15.1, 689
- positive-semidefinite matrix**
 - Fact 11.14.20, 685
 - Fact 11.14.35, 688
 - Fact 11.16.6, 694
 - Fact 11.16.16, 695
- quaternions**
 - Fact 11.11.15, 675
- rank-two matrix**
 - Fact 11.11.18, 676
- resolvent**
 - Proposition 11.2.2, 647
- Schur product**
 - Fact 11.14.21, 685
- semisimple matrix**
 - Proposition 11.2.7, 648
- semistability**
 - Proposition 11.8.2, 662
- semistable matrix**
 - Fact 11.18.5, 698
 - Fact 11.18.7, 699
 - Fact 11.21.7, 713
- series**
 - Proposition 11.4.7, 655
 - Fact 11.14.17, 684
- similar matrices**
 - Proposition 11.2.9, 650
- singular value**
 - Fact 11.15.5, 689
 - Fact 11.16.14, 695
 - Fact 11.16.15, 695
- skew-Hermitian matrix**
 - Proposition 11.2.8, 649
 - Proposition 11.2.9, 650
 - Fact 11.14.6, 683
 - Fact 11.14.33, 688
- skew-involutory matrix**
 - Fact 11.11.1, 671
- skew-symmetric matrix**
 - Example 11.3.6, 652
 - Fact 11.11.3, 672
 - Fact 11.11.6, 673
 - Fact 11.11.7, 673
 - Fact 11.11.8, 674
 - Fact 11.11.9, 674
 - Fact 11.11.15, 675
- Specht's ratio**
 - Fact 11.14.28, 687

spectral abscissa

Fact 11.13.2, 677

Fact 11.15.8, 691

Fact 11.15.9, 691

Fact 11.18.8, 699

Fact 11.18.9, 699

spectral radius

Fact 11.13.2, 677

spectrum

Proposition 11.2.3, 648

Corollary 11.2.6, 648

stable subspace

Proposition 11.8.8, 665

state equation

Proposition 12.1.1, 723

strong log**majorization**

Fact 11.16.4, 692

submultiplicative**norm**

Proposition 11.1.2, 644

Fact 11.15.8, 691

Fact 11.15.9, 691

Fact 11.16.7, 694

Fact 11.18.8, 699

Fact 11.18.9, 699

sum of integer**powers**

Fact 11.11.4, 672

symplectic matrix

Proposition 11.6.7, 659

thermodynamic**inequality**

Fact 11.14.31, 688

trace

Corollary 11.2.4, 648

Corollary 11.2.5, 648

Fact 8.14.18, 500

Fact 11.11.6, 673

Fact 11.14.3, 682

Fact 11.14.10, 683

Fact 11.14.28, 687

Fact 11.14.29, 687

Fact 11.14.30, 687

Fact 11.14.31, 688

Fact 11.14.36, 688

Fact 11.14.38, 688

Fact 11.15.4, 689

Fact 11.15.5, 689

Fact 11.16.1, 692

Fact 11.16.4, 692

transpose

Proposition 11.2.8, 649

unipotent matrix

Fact 11.13.17, 680

unitarily invariant**norm**

Fact 11.15.6, 690

Fact 11.16.4, 692

Fact 11.16.5, 694

Fact 11.16.13, 695

Fact 11.16.16, 695

Fact 11.16.17, 695

unitary matrix

Proposition 11.2.8, 649

Proposition 11.2.9, 650

Proposition 11.6.7, 659

Corollary 11.2.6, 648

Fact 11.14.6, 683

Fact 11.14.33, 688

Fact 11.14.34, 688

upper triangular**matrix**

Fact 11.11.4, 672

Fact 11.13.1, 677

Fact 11.13.16, 680

vibration equation

Example 11.3.7, 653

weak majorization

Fact 11.16.4, 692

Z-matrix

Fact 11.19.1, 706

Zassenhaus product**formula**

Fact 11.14.18, 685

matrix function**definition, 628****Lagrange-Hermite****interpolation****polynomial**

Theorem 10.5.2, 629

spectrum

Corollary 10.5.4, 629

matrix function defined**at a point****definition**

Definition 10.5.1, 628

matrix function**evaluation****identity theorem**

Theorem 10.5.3, 629

matrix inequality**matrix logarithm**

Proposition 8.6.4, 432

matrix inversion lemma**generalization**

Fact 2.16.21, 145

generalized inverse

Fact 6.4.4, 378

inverse matrix

Corollary 2.8.8, 108

matrix logarithm**chaotic order**

Fact 8.19.1, 522

complex matrix

Definition 11.4.1, 654

convergent series

Theorem 11.5.1, 656

convex function

Proposition 8.6.17, 437

determinant

Fact 8.18.30, 521

Fact 9.8.39, 578

Fact 11.14.24, 686

determinant and**derivative**

Proposition 10.7.3, 631

discrete-time**Lyapunov-stable****matrix**

Fact 11.14.19, 685

eigenvalues

Theorem 11.5.1, 656

exponential

Fact 11.14.26, 686

geometric mean

Fact 11.14.39, 689

Hamiltonian matrix

Fact 11.14.19, 685

Klein's inequality

Fact 11.14.25, 686

limit

Proposition 8.6.4, 432

matrix exponential

Theorem 11.5.1, 656

- Proposition 11.4.2, 654
- Fact 11.13.17, 680
- Fact 11.14.21, 685
- Fact 11.14.31, 688
- matrix inequality**
 - Proposition 8.6.4, 432
- maximum singular value**
 - Fact 8.18.30, 521
- nonsingular matrix**
 - Proposition 11.4.2, 654
- norm**
 - Theorem 11.5.1, 656
- positive-definite matrix**
 - Proposition 8.6.4, 432
 - Proposition 11.4.5, 654
 - Fact 8.9.43, 455
 - Fact 8.13.8, 486
 - Fact 8.18.29, 521
 - Fact 8.19.1, 522
 - Fact 8.19.2, 523
 - Fact 9.9.55, 590
 - Fact 11.14.24, 686
 - Fact 11.14.25, 686
 - Fact 11.14.26, 686
 - Fact 11.14.27, 686
- positive-semidefinite matrix**
 - Fact 9.9.54, 590
- quadratic form**
 - Fact 8.15.15, 502
- real matrix**
 - Proposition 11.4.3, 654
 - Fact 11.14.19, 685
- relative entropy**
 - Fact 11.14.25, 686
- Schur product**
 - Fact 8.21.47, 540
 - Fact 8.21.48, 540
- spectrum**
 - Theorem 11.5.1, 656
- symplectic matrix**
 - Fact 11.14.19, 685
- trace**
 - Fact 11.14.24, 686
 - Fact 11.14.25, 686
 - Fact 11.14.27, 686
 - Fact 11.14.31, 688
- unitarily invariant norm**
 - Fact 9.9.54, 590
- matrix measure properties**
 - Fact 11.15.7, 690
- matrix polynomial definition, 234**
- matrix power**
 - outer-product perturbation**
 - Fact 2.12.18, 127
- positive-definite matrix inequality**
 - Fact 8.10.51, 466
 - Fact 8.19.3, 523
- positive-semidefinite matrix**
 - Fact 8.12.30, 482
 - Fact 8.15.16, 502
- matrix product**
 - lower triangular matrix**
 - Fact 3.20.18, 221
 - normal matrix**
 - Fact 9.9.6, 580
 - strictly lower triangular matrix**
 - Fact 3.20.18, 221
 - strictly upper triangular matrix**
 - Fact 3.20.18, 221
 - unitarily invariant norm**
 - Fact 9.9.6, 580
 - upper triangular matrix**
 - Fact 3.20.18, 221
- matrix sign function**
 - convergent sequence**
 - Fact 5.15.21, 348
 - definition**
 - Definition 10.6.2, 630
 - partitioned matrix**
 - Fact 10.10.3, 637
 - positive-definite matrix**
 - Fact 10.10.4, 637
- properties**
 - Fact 10.10.2, 637
- square root**
 - Fact 5.15.21, 348
- matrix sine**
 - matrix cosine**
 - Fact 11.12.1, 677
 - matrix exponential**
 - Fact 11.12.1, 677
- maximal solution**
 - Riccati equation**
 - Definition 12.16.12, 780
 - Theorem 12.18.1, 787
 - Theorem 12.18.4, 787
 - Proposition 12.18.2, 787
 - Proposition 12.18.7, 789
- maximal solution of the Riccati equation**
 - closed-loop spectrum**
 - Proposition 12.18.2, 787
 - stabilizability**
 - Theorem 12.18.1, 787
- maximization**
 - continuous function**
 - Fact 10.11.4, 638
- maximum eigenvalue**
 - commutator**
 - Fact 9.9.30, 585
 - Fact 9.9.31, 585
 - Hermitian matrix**
 - Lemma 8.4.3, 425
 - Fact 5.11.5, 321
 - Fact 8.10.3, 456
 - matrix exponential**
 - Fact 11.16.4, 692
 - positive-semidefinite matrix**
 - Fact 8.18.11, 515
 - Fact 8.18.13, 516
 - Fact 8.18.14, 516
 - quadratic form**
 - Lemma 8.4.3, 425

- spectral abscissa
 - Fact 5.11.5, 321
- unitarily invariant norm
 - Fact 9.9.30, 585
 - Fact 9.9.31, 585
- maximum singular value**
 - absolute value
 - Fact 9.13.10, 603
 - block-diagonal matrix
 - Fact 5.11.33, 328
 - block-triangular matrix
 - Fact 5.11.32, 328
 - bound
 - Fact 5.11.35, 328
 - commutator
 - Fact 9.9.29, 584
 - Fact 9.14.9, 609
 - complex conjugate transpose
 - Fact 8.17.3, 508
 - Fact 8.18.11, 515
 - Fact 8.21.10, 533
 - Cordes inequality**
 - Fact 8.18.26, 520
 - derivative
 - Fact 11.15.2, 689
 - determinant
 - Fact 9.14.17, 611
 - Fact 9.14.18, 611
 - discrete-time
 - Lyapunov-stable matrix
 - Fact 11.21.17, 715
 - dissipative matrix
 - Fact 8.17.12, 511
 - eigenvalue of
 - Hermitian part
 - Fact 5.11.25, 326
 - eigenvalue
 - perturbation
 - Fact 9.12.4, 599
 - Fact 9.12.8, 601
 - elementary projector
 - Fact 9.14.1, 607
- equi-induced
 - self-adjoint norm
 - Fact 9.13.5, 602
- equi-induced
 - unitarily invariant norm
 - Fact 9.13.4, 602
- generalized inverse
 - Fact 9.14.8, 608
 - Fact 9.14.30, 615
- Hermitian matrix**
 - Fact 5.11.5, 321
 - Fact 9.9.41, 588
- Hölder-induced norm**
 - Fact 9.8.21, 575
- idempotent matrix
 - Fact 5.11.38, 328
 - Fact 5.11.39, 329
 - Fact 5.12.18, 336
- induced lower bound
 - Corollary 9.5.5, 560
- induced norm
 - Fact 9.8.24, 575
- inequality
 - Proposition 9.2.2, 548
 - Corollary 9.6.5, 562
 - Corollary 9.6.9, 562
 - Fact 9.9.32, 585
 - Fact 9.14.16, 611
- inverse matrix
 - Fact 9.14.8, 608
- Kreiss matrix theorem**
 - Fact 11.21.17, 715
- Kronecker product**
 - Fact 9.14.37, 617
- matrix difference
 - Fact 8.18.8, 515
 - Fact 9.9.32, 585
- matrix exponential
 - Fact 11.15.1, 689
 - Fact 11.15.2, 689
 - Fact 11.15.5, 689
 - Fact 11.16.6, 694
 - Fact 11.16.10, 694
- matrix logarithm
 - Fact 8.18.30, 521
- matrix power
 - Fact 8.18.26, 520
 - Fact 9.13.7, 603
 - Fact 9.13.9, 603
- normal matrix
 - Fact 5.14.15, 341
 - Fact 9.8.13, 573
 - Fact 9.12.8, 601
 - Fact 9.13.7, 603
 - Fact 9.13.8, 603
 - Fact 9.14.5, 608
 - Fact 11.16.10, 694
- outer-product matrix
 - Fact 5.11.16, 324
 - Fact 5.11.18, 324
 - Fact 9.7.26, 570
- partitioned matrix
 - Fact 8.17.3, 508
 - Fact 8.17.14, 512
 - Fact 8.18.1, 512
 - Fact 8.18.2, 513
 - Fact 9.10.1, 593
 - Fact 9.10.3, 594
 - Fact 9.10.4, 594
 - Fact 9.10.5, 595
 - Fact 9.14.12, 610
 - Fact 9.14.13, 610
 - Fact 9.14.14, 610
- positive-definite matrix
 - Fact 8.18.25, 520
- positive-semidefinite matrix
 - Fact 8.18.1, 512
 - Fact 8.18.2, 513
 - Fact 8.18.8, 515
 - Fact 8.18.12, 516
 - Fact 8.18.13, 516
 - Fact 8.18.14, 516
 - Fact 8.18.15, 517
 - Fact 8.18.16, 517
 - Fact 8.18.25, 520
 - Fact 8.18.26, 520
 - Fact 8.18.28, 521
 - Fact 8.18.30, 521
 - Fact 8.18.31, 522
 - Fact 8.20.9, 526
 - Fact 11.16.6, 694
- power

- Fact 11.21.17, 715
- product**
 - Fact 9.14.2, 607
- projector**
 - Fact 5.11.38, 328
 - Fact 5.12.17, 335
 - Fact 5.12.18, 336
 - Fact 9.14.1, 607
 - Fact 9.14.30, 615
- quadratic form**
 - Fact 9.13.1, 602
 - Fact 9.13.2, 602
- Schur product**
 - Fact 8.21.10, 533
 - Fact 9.14.31, 615
 - Fact 9.14.33, 616
 - Fact 9.14.35, 617
- spectral abscissa**
 - Fact 5.11.26, 326
- spectral radius**
 - Corollary 9.4.10, 556
 - Fact 5.11.5, 321
 - Fact 5.11.26, 326
 - Fact 8.18.25, 520
 - Fact 9.8.13, 573
 - Fact 9.13.9, 603
- square root**
 - Fact 8.18.14, 516
 - Fact 9.8.32, 576
 - Fact 9.14.15, 611
- sum of matrices**
 - Fact 9.14.15, 611
- trace**
 - Fact 5.12.7, 334
 - Fact 9.14.4, 608
- trace norm**
 - Corollary 9.3.8, 552
- unitarily invariant norm**
 - Fact 9.9.10, 581
 - Fact 9.9.29, 584
- maximum singular value bound**
- Frobenius norm**
 - Fact 9.13.13, 604
- minimum singular value bound**
 - Fact 9.13.14, 604
- polynomial root**
 - Fact 9.13.14, 604
- trace**
 - Fact 9.13.13, 604
- maximum singular value of a matrix difference**
- Kato**
 - Fact 9.9.32, 585
- maximum singular value of a partitioned matrix**
- Parrott's theorem**
 - Fact 9.14.13, 610
- Tomiyama**
 - Fact 9.14.12, 610
- McCarthy inequality**
 - positive-semidefinite matrix
 - Fact 8.12.29, 481
- McCoy**
 - simultaneous triangularization
 - Fact 5.17.5, 358
- McIntosh's inequality**
 - unitarily invariant norm
 - Fact 9.9.47, 589
- McLaughlin's inequality**
 - refined Cauchy-Schwarz inequality
 - Fact 1.16.17, 64
- McMillan degree**
 - Definition 4.7.10, 251
 - minimal realization
 - Theorem 12.9.16, 756
- mean**
 - inequality
 - Fact 1.16.18, 64
 - Laguerre-Samuelson inequality
 - Fact 1.15.12, 51
 - Fact 8.9.35, 454
 - variance inequality
 - Fact 1.15.12, 51
- Fact 8.9.35, 454
- mean-value inequality**
 - product of means
 - Fact 1.15.38, 57
 - Fact 1.15.44, 59
- Mercator's series**
 - infinite series
 - Fact 1.18.8, 73
- Mihet**
 - polynomial bound
 - Fact 11.20.13, 712
- Milne's inequality**
 - refined Cauchy-Schwarz inequality
 - Fact 1.16.15, 63
- Milnor**
 - simultaneous diagonalization of symmetric matrices
 - Fact 8.16.6, 507
- MIMO transfer function**
 - definition
 - Definition 12.9.1, 749
- minimal polynomial**
 - block-diagonal matrix
 - Lemma 5.2.7, 286
 - block-triangular matrix
 - Fact 4.10.12, 268
 - characteristic polynomial
 - Fact 4.9.24, 265
 - companion matrix
 - Proposition 5.2.1, 284
 - Corollary 5.2.4, 286
 - Corollary 5.2.5, 286
 - cyclic matrix
 - Proposition 5.5.15, 299
 - definition, 247
 - existence
 - Theorem 4.6.1, 247
 - index of an eigenvalue
 - Proposition 5.5.15, 299

- Jordan form**
 - Proposition 5.5.15, 299
- null space**
 - Corollary 11.8.6, 664
- partitioned matrix**
 - Fact 4.10.12, 268
- range**
 - Corollary 11.8.6, 664
- similar matrices**
 - Proposition 4.6.3, 248
 - Fact 11.23.3, 717
 - Fact 11.23.4, 717
 - Fact 11.23.5, 718
 - Fact 11.23.6, 719
 - Fact 11.23.7, 719
 - Fact 11.23.8, 720
 - Fact 11.23.9, 720
 - Fact 11.23.10, 721
 - Fact 11.23.11, 721
- spectrum**
 - Fact 4.10.8, 267
- stable subspace**
 - Proposition 11.8.5, 664
 - Fact 11.23.1, 716
 - Fact 11.23.2, 716
- upper**
 - block-triangular matrix**
 - Fact 4.10.12, 268
- minimal realization**
 - asymptotically stable matrix**
 - Definition 12.9.17, 757
 - asymptotically stable transfer function**
 - Proposition 12.9.18, 757
 - balanced realization**
 - Proposition 12.9.21, 757
 - block decomposition**
 - Proposition 12.9.10, 753
 - controllable pair**
 - Proposition 12.9.10, 753
 - Corollary 12.9.15, 756
 - definition**
 - Definition 12.9.14, 756
- Kalman**
 - decomposition**
 - Proposition 12.9.10, 753
- Lyapunov-stable matrix**
 - Definition 12.9.17, 757
- Lyapunov-stable transfer function**
 - Proposition 12.9.18, 757
- Markov**
 - block-Hankel matrix**
 - Proposition 12.9.12, 755
- McMillan degree**
 - Theorem 12.9.16, 756
- observable pair**
 - Proposition 12.9.10, 753
 - Corollary 12.9.15, 756
- pole**
 - Fact 12.22.2, 798
 - Fact 12.22.12, 800
- rational transfer function**
 - Fact 12.22.12, 800
- semistable matrix**
 - Definition 12.9.17, 757
- semistable transfer function**
 - Proposition 12.9.18, 757
- minimal-rank identity partitioned matrix**
 - Fact 6.5.7, 387
- minimum eigenvalue**
 - Hermitian matrix**
 - Lemma 8.4.3, 425
 - Fact 8.10.3, 456
 - nonnegative matrix**
 - Fact 4.11.9, 276
 - quadratic form**
 - Lemma 8.4.3, 425
 - Z-matrix**
 - Fact 4.11.9, 276
- minimum principle**
 - eigenvalue**
 - characterization**
 - Fact 8.17.15, 512
 - minimum singular value**
 - determinant**
 - Fact 9.14.18, 611
 - eigenvalue of Hermitian part**
 - Fact 5.11.25, 326
 - induced lower bound**
 - Corollary 9.5.5, 560
 - inequality**
 - Corollary 9.6.6, 562
 - Fact 9.13.6, 602
 - lower bound**
 - Fact 9.13.15, 604
 - Fact 9.13.21, 606
 - quadratic form**
 - Fact 9.13.1, 602
 - spectral abscissa**
 - Fact 5.11.26, 326
 - spectral radius**
 - Fact 5.11.26, 326
 - minimum singular value bound**
 - maximum singular value bound**
 - Fact 9.13.14, 604
 - polynomial root**
 - Fact 9.13.14, 604
 - Minkowski**
 - set-defined norm**
 - Fact 10.8.22, 633
 - Minkowski's**
 - determinant theorem**
 - positive-semidefinite matrix determinant**
 - Corollary 8.4.15, 429
 - Minkowski's inequality**
 - Hölder norm**
 - Lemma 9.1.3, 544
 - positive-semidefinite matrix**
 - Fact 8.12.29, 481
 - scalar case**
 - Fact 1.16.25, 66

minor, see
subdeterminant

Mircea's inequality
triangle
Fact 2.20.11, 156

Mirsky
singular value trace
bound
Fact 5.12.6, 334

Mirsky's theorem
singular value
perturbation
Fact 9.14.29, 614

MISO transfer function
definition
Definition 12.9.1, 749

mixed
arithmetic-geometric
mean inequality
arithmetic mean
Fact 1.15.39, 58

ML-matrix
definition, 230

Moler
regular pencil
Fact 5.17.3, 358

monic polynomial
definition, 231

monic polynomial
matrix
definition, 234

monotone norm
absolute norm
Proposition 9.1.2, 543
definition, 543

monotonicity
Callebaut
Fact 1.16.1, 60
power inequality
Fact 1.10.33, 36
power mean
inequality
Fact 1.15.30, 55
Riccati equation

Proposition 12.18.5,
788
Corollary 12.18.6, 788

monotonicity theorem
Hermitian matrix
eigenvalues
Theorem 8.4.9, 427
Fact 8.10.4, 456

Moore-Penrose
generalized inverse,
see generalized
inverse

Muirhead's theorem
Schur convex
function
Fact 1.15.25, 54
strong majorization
Fact 2.21.5, 162

multicompanion form
definition, 285
existence
Theorem 5.2.3, 285
similar matrices
Corollary 5.2.6, 286
similarity invariant
Corollary 5.2.6, 286

multigraph
definition, 8

multinomial theorem
power of sum
Fact 1.15.1, 48

multiple
definition, 233

multiplication
definition, 81
function composition
Theorem 2.1.3, 81
Kronecker product
Proposition 7.1.6, 400

multiplicative
commutator
realization
Fact 5.15.34, 351
reflector realization
Fact 5.15.35, 351

multiplicative
perturbation
small-gain theorem
Fact 9.13.23, 606

multiplicity of a root
definition, 232

multirelation
definition, 5

multiset
definition, 2

multispectrum
definition
Definition 4.4.4, 240
properties
Proposition 4.4.5, 241

N

N-matrix
almost nonnegative
matrix
Fact 11.19.3, 706
Fact 11.19.5, 707
asymptotically stable
matrix
Fact 11.19.5, 707
definition
Fact 11.19.3, 706
group-invertible
matrix
Fact 11.19.4, 706
Lyapunov-stable
matrix
Fact 11.19.4, 706
nonnegative matrix
Fact 11.19.3, 706

Nanjundiah
mixed arithmetic-
geometric mean
inequality
Fact 1.15.39, 58

natural frequency
definition, 654
Fact 5.14.35, 344

necessity

- definition, 1
- negation**
 - definition, 1
- negative-definite matrix**
 - asymptotically stable matrix
 - Fact 11.18.30, 704
 - definition
 - Definition 3.1.1, 165
- negative-semidefinite matrix**
 - definition
 - Definition 3.1.1, 165
 - Euclidean distance matrix
 - Fact 9.8.14, 573
- Nesbitt's inequality**
 - scalar inequality
 - Fact 1.11.21, 44
- Newcomb**
 - simultaneous cogredient diagonalization, 541
- Newton's identities**
 - elementary symmetric polynomial
 - Fact 4.8.2, 254
 - polynomial roots
 - Fact 4.8.2, 254
 - spectrum
 - Fact 4.10.6, 267
- Newton's inequality**
 - elementary symmetric polynomial
 - Fact 1.15.11, 50
- Newton-Raphson algorithm**
 - generalized inverse
 - Fact 6.3.35, 376
 - inverse matrix
 - Fact 2.16.29, 146
 - square root
 - Fact 5.15.21, 348
- Niculescu's inequality**
 - absolute-value function
 - Fact 1.10.19, 33
 - convex function
 - Fact 1.8.5, 22
 - square-root function
 - Fact 1.10.20, 33
- nilpotent matrix**
 - additive decomposition
 - Fact 5.9.3, 311
 - adjugate
 - Fact 6.3.6, 370
 - commutator
 - Fact 3.12.16, 200
 - Fact 3.17.11, 214
 - Fact 3.17.12, 214
 - Fact 3.17.13, 214
 - commuting matrices
 - Fact 3.17.9, 214
 - Fact 3.17.10, 214
 - defective matrix
 - Fact 5.14.18, 341
 - definition
 - Definition 3.1.1, 165
 - determinant
 - Fact 3.17.9, 214
 - example
 - Example 5.5.17, 299
 - factorization
 - Fact 5.15.29, 350
 - idempotent matrix
 - Fact 3.12.16, 200
 - identity perturbation
 - Fact 3.17.7, 214
 - Fact 3.17.8, 214
 - inertia
 - Fact 5.8.4, 307
 - Jordan-Chevalley decomposition
 - Fact 5.9.3, 311
 - Kronecker product
 - Fact 7.4.16, 406
 - Kronecker sum
 - Fact 7.5.3, 409
 - Fact 7.5.8, 409
 - lower triangular matrix
 - Fact 3.17.11, 214
 - matrix exponential
 - Fact 11.11.1, 671
 - Fact 11.13.17, 680
 - matrix sum
 - Fact 3.17.10, 214
 - null space
 - Fact 3.17.1, 213
 - Fact 3.17.2, 213
 - Fact 3.17.3, 213
 - outer-product matrix
 - Fact 5.14.3, 338
 - partitioned matrix
 - Fact 3.12.14, 200
 - Fact 5.10.23, 321
 - range
 - Fact 3.17.1, 213
 - Fact 3.17.2, 213
 - Fact 3.17.3, 213
 - rank
 - Fact 3.17.4, 213
 - Fact 3.17.5, 213
 - S-N decomposition
 - Fact 5.9.3, 311
 - similar matrices
 - Proposition 3.4.5, 174
 - Fact 5.10.23, 321
 - simultaneous triangularization
 - Fact 5.17.6, 358
 - spectrum
 - Proposition 5.5.21, 300
 - Toeplitz matrix
 - Fact 3.18.6, 216
 - trace
 - Fact 3.17.6, 214
 - triangular matrix
 - Fact 5.17.6, 358
 - unitarily similar matrices
 - Proposition 3.4.5, 174
 - upper triangular matrix
 - Fact 3.17.11, 214
- node**
 - definition, 8

**nondecreasing
function**

convex function
Lemma 8.6.16, 436

definition
Definition 8.6.12, 434

function composition
Lemma 8.6.16, 436

geometric mean
Fact 8.10.43, 461
Fact 8.10.44, 464

matrix exponential
Fact 8.10.44, 464

matrix functions
Proposition 8.6.13, 435

Schur complement
Proposition 8.6.13, 435

**nonderogatory
eigenvalue**

definition
Definition 5.5.4, 296

**nonderogatory matrix
definition**

Definition 5.5.4, 296

**nonempty set
definition, 2****nonincreasing function
concave function**

Lemma 8.6.16, 436

definition
Definition 8.6.12, 434

function composition
Lemma 8.6.16, 436

**nonnegative matrix
almost nonnegative
matrix**

Fact 11.19.1, 706

aperiodic graph
Fact 4.11.5, 273

companion matrix
Fact 4.11.13, 279

copositive matrix
Fact 8.15.37, 507

definition, 81
Definition 3.1.4, 168

difference equation
Fact 4.11.12, 277

eigenvalue
Fact 4.11.5, 273

Fibonacci numbers
Fact 4.11.12, 277

**limit of matrix
powers**

Fact 4.11.21, 280

Lucas numbers
Fact 4.11.12, 277

M-matrix
Fact 4.11.6, 275

matrix power
Fact 4.11.22, 281

minimum eigenvalue
Fact 4.11.9, 276

N-matrix
Fact 11.19.3, 706

spectral radius
Fact 4.11.5, 273
Fact 4.11.6, 275

Fact 4.11.16, 279

Fact 4.11.17, 280

Fact 7.6.13, 415

Fact 11.19.3, 706

**spectral radius
convexity**

Fact 4.11.19, 280

**spectral radius
monotonicity**

Fact 4.11.18, 280

trace
Fact 4.11.22, 281

**nonnegative matrix
eigenvalues**

**Perron-Frobenius
theorem**

Fact 4.11.5, 273

**nonnegative vector
definition, 79**

**linear system
solution**

Fact 4.11.14, 279

null space
Fact 4.11.15, 279

**nonsingular matrix
complex conjugate**

Proposition 2.6.8, 102

**complex conjugate
transpose**

Proposition 2.6.8, 102

Fact 2.16.30, 146

controllable subspace
Proposition 12.6.10,

740

cyclic matrix
Fact 5.14.9, 340

definition, 100

determinant
Corollary 2.7.4, 104
Lemma 2.8.6, 108

**determinant lower
bound**

Fact 4.10.18, 269

**diagonal dominance
theorem**

Fact 4.10.17, 269

Fact 4.10.18, 269

**diagonally dominant
matrix**

Fact 4.10.17, 269

dissipative matrix
Fact 3.20.13, 220

**distance to
singularity**

Fact 9.14.7, 608

elementary matrix
Fact 5.15.12, 347

factorization
Fact 5.15.12, 347
Fact 5.15.36, 351

group
Proposition 3.3.6, 172

idempotent matrix
Fact 3.12.11, 199
Fact 3.12.26, 203
Fact 3.12.28, 203
Fact 3.12.32, 205

inverse matrix
Fact 3.7.1, 178

matrix logarithm
Proposition 11.4.2, 654

norm
Fact 9.7.32, 571

normal matrix
Fact 3.7.1, 178

perturbation

- Fact 9.14.6, 608
- Fact 9.14.18, 611
- range-Hermitian matrix**
 - Proposition 3.1.6, 169
- similar matrices**
 - Fact 5.10.11, 318
- simplex**
 - Fact 2.20.4, 154
- skew Hermitian matrix**
 - Fact 3.7.1, 178
- spectral radius**
 - Fact 4.10.29, 272
- submultiplicative norm**
 - Fact 9.8.5, 572
- Sylvester's equation**
 - Fact 12.21.14, 796
- transpose**
 - Proposition 2.6.8, 102
- unitary matrix**
 - Fact 3.7.1, 178
- unobservable subspace**
 - Proposition 12.3.10, 730
- weak diagonal dominance theorem**
 - Fact 4.10.19, 270
- nonsingular matrix transformation**
 - Smith polynomial
 - Proposition 4.3.8, 238
- nonsingular polynomial matrix**
 - Definition 4.2.5, 235
 - regular polynomial matrix
 - Proposition 4.2.5, 235
- nonzero diagonal entries**
 - similar matrices
 - Fact 5.9.14, 313
- norm**
 - absolute
 - definition, 543
- absolute sum**
 - definition, 545
- column**
 - definition, 556
- compatible**
 - definition, 549
- complex conjugate transpose**
 - Fact 9.8.8, 572
- convex set**
 - Fact 9.7.23, 570
- Dunkl-Williams inequality**
 - Fact 9.7.10, 566
 - Fact 9.7.13, 567
- equi-induced**
 - Definition 9.4.1, 553
- equivalent**
 - Theorem 9.1.8, 546
- Euclidean**
 - definition, 545
- Euclidean-norm inequality**
 - Fact 9.7.4, 563
 - Fact 9.7.18, 569
- Frobenius**
 - definition, 547
- Hölder-norm inequality**
 - Fact 9.7.18, 569
- idempotent matrix**
 - Fact 11.16.12, 695
- induced**
 - Definition 9.4.1, 553
- induced norm**
 - Theorem 9.4.2, 553
- inequality**
 - Fact 9.7.2, 563
 - Fact 9.7.4, 563
 - Fact 9.7.10, 566
 - Fact 9.7.13, 567
 - Fact 9.7.16, 568
 - Fact 9.7.17, 569
- infinity**
 - definition, 545
- linear combination of norms**
 - Fact 9.7.31, 571
- linear system solution**
 - Fact 9.9.64, 592
 - Fact 9.9.65, 592
 - Fact 9.9.66, 592
- Maligranda inequality**
 - Fact 9.7.10, 566
 - Fact 9.7.13, 567
- Massera-Schaffer inequality**
 - Fact 9.7.10, 566
 - Fact 9.7.13, 567
- matrix**
 - Definition 9.2.1, 546
- matrix exponential**
 - Fact 11.16.9, 694
 - Fact 11.16.11, 694
 - Fact 11.16.12, 695
- matrix logarithm**
 - Theorem 11.5.1, 656
- monotone**
 - definition, 543
- nonsingular matrix**
 - Fact 9.7.32, 571
- normalized**
 - definition, 547
- partitioned matrix**
 - Fact 9.10.1, 593
 - Fact 9.10.2, 593
 - Fact 9.10.8, 596
- positive-definite matrix**
 - Fact 9.7.30, 571
- quadratic form**
 - Fact 9.7.30, 571
- row**
 - definition, 556
- self-adjoint**
 - definition, 547
- set-defined**
 - Fact 10.8.22, 633
- spectral**
 - definition, 549
- spectral radius**
 - Proposition 9.2.6, 549
- submultiplicative**
 - definition, 550
- trace**

- definition, 549
- triangle inequality**
 - Definition 9.1.1, 543
- unitarily invariant**
 - definition, 547
- vector**
 - Definition 9.1.1, 543
- weakly unitarily invariant**
 - definition, 547
- norm bound**
 - matrix exponential**
 - Fact 11.18.10, 699
- norm equality**
 - common eigenvector**
 - Fact 9.9.33, 585
 - Schatten norm**
 - Fact 9.9.33, 585
- norm identity**
 - Hlawka's equality**
 - Fact 9.7.4, 563
 - polarization identity**
 - Fact 9.7.4, 563
 - Pythagorean theorem**
 - Fact 9.7.4, 563
- norm inequality**
 - Aczel's inequality**
 - Fact 9.7.4, 563
 - Bessel's inequality**
 - Fact 9.7.4, 563
 - Buzano's inequality**
 - Fact 9.7.4, 563
 - convex combination**
 - Fact 9.7.15, 568
 - Hlawka's inequality**
 - Fact 9.7.4, 563
 - Hölder norm**
 - Fact 9.7.21, 569
 - orthogonal vectors**
 - Fact 9.7.25, 570
 - Parseval's inequality**
 - Fact 9.7.4, 563
 - polygonal inequalities**
 - Fact 9.7.4, 563
 - quadrilateral inequality**
 - Fact 9.7.4, 563
 - Schatten norm**
 - Fact 9.9.34, 586
 - Fact 9.9.36, 586
 - Fact 9.9.37, 586
 - Fact 9.9.38, 587
 - unitarily invariant norm**
 - Fact 9.9.47, 589
 - Fact 9.9.48, 589
 - Fact 9.9.49, 589
 - Fact 9.9.50, 589
 - vector inequality**
 - Fact 9.7.11, 567
 - Fact 9.7.12, 567
 - Fact 9.7.14, 568
 - Fact 9.7.15, 568
 - von Neumann–Jordan inequality**
 - Fact 9.7.11, 567
 - norm monotonicity**
 - power-sum inequality**
 - Fact 1.10.30, 35
 - Fact 1.15.34, 57
 - norm-compression inequality**
 - partitioned matrix**
 - Fact 9.10.1, 593
 - Fact 9.10.8, 596
 - positive-semidefinite matrix**
 - Fact 9.10.6, 595
 - normal matrix**
 - affine mapping**
 - Fact 3.7.14, 181
 - asymptotically stable matrix**
 - Fact 11.18.37, 705
 - block-diagonal matrix**
 - Fact 3.7.8, 179
 - characterizations**
 - Fact 3.7.12, 180
 - commutator**
 - Fact 3.8.6, 185
 - Fact 3.8.7, 185
 - Fact 9.9.31, 585
 - commuting matrices**
 - Fact 3.7.28, 183
 - Fact 3.7.29, 183
 - Fact 5.14.29, 342
 - Fact 5.17.7, 358
 - Fact 11.14.5, 682
 - complex conjugate transpose**
 - Fact 5.14.30, 343
 - Fact 6.3.16, 373
 - Fact 6.3.17, 373
 - Fact 6.6.10, 394
 - Fact 6.6.17, 397
 - congruence transformation**
 - Fact 5.10.17, 319
 - definition**
 - Definition 3.1.1, 165
 - determinant**
 - Fact 5.12.12, 335
 - discrete-time asymptotically stable matrix**
 - Fact 11.21.4, 712
 - discrete-time Lyapunov-stable matrix**
 - Fact 11.21.4, 712
 - dissipative matrix**
 - Fact 11.18.37, 705
 - eigenvalue**
 - Fact 5.14.15, 341
 - eigenvalue perturbation**
 - Fact 9.12.8, 601
 - eigenvector**
 - Proposition 4.5.4, 246
 - Lemma 4.5.3, 246
 - example**
 - Example 5.5.17, 299
 - Frobenius norm**
 - Fact 9.12.9, 601
 - generalized inverse**
 - Proposition 6.1.6, 365
 - Fact 6.3.16, 373
 - Fact 6.3.17, 373

- group generalized inverse**
 - Fact 6.6.10, 394
- group-invertible matrix**
 - Fact 6.6.17, 397
- Hermitian matrix**
 - Proposition 3.1.6, 169
- idempotent matrix**
 - Fact 3.13.3, 206
- inertia**
 - Fact 5.10.17, 319
- involutory matrix**
 - Fact 5.9.9, 312
 - Fact 5.9.10, 312
- Jordan form**
 - Fact 5.10.6, 317
- Kronecker product**
 - Fact 7.4.16, 406
- Kronecker sum**
 - Fact 7.5.8, 409
- Lyapunov-stable matrix**
 - Fact 11.18.37, 705
- matrix exponential**
 - Proposition 11.2.8, 649
 - Fact 11.13.18, 680
 - Fact 11.14.5, 682
 - Fact 11.16.10, 694
- matrix power**
 - Fact 9.13.7, 603
- matrix product**
 - Fact 9.9.6, 580
- maximum singular value**
 - Fact 5.14.15, 341
 - Fact 9.8.13, 573
 - Fact 9.12.8, 601
 - Fact 9.13.7, 603
 - Fact 9.13.8, 603
 - Fact 9.14.5, 608
 - Fact 11.16.10, 694
- orthogonal eigenvectors**
 - Corollary 5.4.8, 294
- partitioned matrix**
 - Fact 3.12.14, 200
 - Fact 8.11.12, 470
- polar decomposition**
 - Fact 11.13.9, 679
- positive-semidefinite matrix**
 - Fact 8.9.22, 452
 - Fact 8.10.11, 457
 - Fact 8.11.12, 470
- projector**
 - Fact 3.13.3, 206
 - Fact 3.13.20, 209
- Putnam-Fuglede theorem**
 - Fact 5.14.30, 343
- range-Hermitian matrix**
 - Proposition 3.1.6, 169
- reflector**
 - Fact 5.9.9, 312
 - Fact 5.9.10, 312
- Schatten norm**
 - Fact 9.9.27, 584
 - Fact 9.14.5, 608
- Schur decomposition**
 - Corollary 5.4.4, 293
 - Fact 5.10.6, 317
- Schur product**
 - Fact 9.9.63, 591
- semidissipative matrix**
 - Fact 11.18.37, 705
- semisimple matrix**
 - Proposition 5.5.11, 297
- shifted-unitary matrix**
 - Fact 3.11.34, 198
- similar matrices**
 - Proposition 5.5.11, 297
 - Fact 5.9.9, 312
 - Fact 5.9.10, 312
 - Fact 5.10.7, 317
- similarity transformation**
 - Fact 5.15.3, 345
- singular value**
 - Fact 5.14.15, 341
- skew-Hermitian matrix**
 - Proposition 3.1.6, 169
- spectral decomposition**
 - Fact 5.14.14, 340
- spectral radius**
 - Fact 5.14.15, 341
- spectral variation**
 - Fact 9.12.5, 600
 - Fact 9.12.6, 600
- spectrum**
 - Fact 4.10.24, 271
 - Fact 8.14.7, 496
 - Fact 8.14.8, 497
- square root**
 - Fact 8.9.27, 453
 - Fact 8.9.28, 453
 - Fact 8.9.29, 453
- trace**
 - Fact 3.7.12, 180
 - Fact 8.12.5, 476
- trace of product**
 - Fact 5.12.4, 333
- transpose**
 - Fact 5.9.9, 312
 - Fact 5.9.10, 312
- unitarily invariant norm**
 - Fact 9.9.6, 580
- unitarily similar matrices**
 - Proposition 3.4.5, 174
 - Corollary 5.4.4, 293
 - Fact 5.10.6, 317
 - Fact 5.10.7, 317
- unitary matrix**
 - Proposition 3.1.6, 169
 - Fact 3.11.4, 189
 - Fact 5.15.1, 345
- normal rank**
 - definition for a polynomial matrix**
 - Definition 4.2.4, 235
 - definition for a rational transfer function**
 - Definition 4.7.4, 249
- rational transfer function, 281**
- normalized norm**
 - definition, 547**
 - equi-induced norm**

Theorem 9.4.2, 553

**normalized
submultiplicative
norm**

inverse matrix

Fact 9.8.44, 579

Fact 9.9.56, 590

Fact 9.9.57, 590

Fact 9.9.58, 591

Fact 9.9.59, 591

null space

adjugate

Fact 2.16.7, 143

definition, 94

Drazin generalized

inverse

Proposition 6.2.2, 368

generalized inverse

Proposition 6.1.6, 365

Fact 6.3.24, 374

group generalized

inverse

Proposition 6.2.3, 369

group-invertible

matrix

Fact 3.6.1, 177

idempotent matrix

Fact 3.12.3, 199

Fact 3.15.4, 200

Fact 6.3.24, 374

identity

Fact 2.10.20, 117

inclusion

Fact 2.10.5, 116

Fact 2.10.7, 116

inclusion for a

matrix power

Corollary 2.4.2, 94

inclusion for a

matrix product

Lemma 2.4.1, 94

Fact 2.10.2, 115

intersection

Fact 2.10.9, 116

involutory matrix

Fact 3.15.4, 212

left-equivalent

matrices

Proposition 5.1.3, 283

Lyapunov equation

Fact 12.21.15, 797

matrix sum

Fact 2.10.10, 116

minimal polynomial

Corollary 11.8.6, 664

nilpotent matrix

Fact 3.17.1, 213

Fact 3.17.2, 213

Fact 3.17.3, 213

outer-product

matrix

Fact 2.10.11, 116

partitioned matrix

Fact 2.11.3, 121

positive-semidefinite

matrix

Fact 8.7.3, 443

Fact 8.7.5, 443

Fact 8.15.1, 500

quadratic form

Fact 8.15.1, 500

range

Corollary 2.5.6, 97

Fact 2.10.1, 115

range inclusions

Theorem 2.4.3, 94

range-Hermitian

matrix

Fact 3.6.3, 177

semisimple

eigenvalue

Proposition 5.5.8, 296

skew-Hermitian

matrix

Fact 8.7.3, 443

symmetric matrix

Fact 3.7.4, 178

nullity, see defect

nullity theorem

defect of a submatrix

Fact 2.11.20, 125

partitioned matrix

Fact 9.14.11, 609

numerical radius

weakly unitarily

invariant norm

Fact 9.8.38, 577

numerical range

spectrum of convex

hull

Fact 8.14.7, 496

Fact 8.14.8, 497

O

**oblique projector, see
idempotent matrix**

observability

closed-loop spectrum

Lemma 12.16.17, 781

Riccati equation

Lemma 12.16.18, 781

Sylvester's equation

Fact 12.21.14, 796

observability Gramian

asymptotically stable

matrix

Corollary 12.4.10, 734

H₂ norm

Corollary 12.11.4, 767

L₂ norm

Theorem 12.11.1, 765

observably

asymptotically

stable

Proposition 12.4.3, 732

Proposition 12.4.4, 733

Proposition 12.4.5, 733

Proposition 12.4.6, 734

Proposition 12.4.7, 734

observability matrix

definition, 728

generalized inverse

Fact 12.20.19, 793

Lyapunov equation

Fact 12.21.15, 797

observable pair

Theorem 12.3.18, 732

Fact 12.20.19, 793

rank

Corollary 12.3.3, 729

Sylvester's equation

Fact 12.21.13, 796

observability pencil**definition**

Definition 12.3.12, 731

Smith form

Proposition 12.3.15, 731

Smith zeros

Proposition 12.3.16, 731

unobservable**eigenvalue**

Proposition 12.3.13, 731

unobservable**spectrum**

Proposition 12.3.16, 731

observable canonical form**definition, 750****equivalent****realizations**

Corollary 12.9.9, 752

realization

Proposition 12.9.3, 750

observable dynamics**block-triangular****matrix**

Theorem 12.3.8, 730

orthogonal matrix

Theorem 12.3.8, 730

observable eigenvalue**closed-loop spectrum**

Lemma 12.16.16, 781

observable subspace

Proposition 12.3.17, 732

observable pair**asymptotically stable****matrix**

Proposition 12.4.9, 734

Corollary 12.4.10, 734

eigenvalue placement

Proposition 12.3.19, 732

equivalent**realizations**

Proposition 12.9.8, 752

invariant zero

Corollary 12.10.12, 765

Markov**block-Hankel****matrix**

Proposition 12.9.11, 754

minimal realization

Proposition 12.9.10, 753

Corollary 12.9.15, 756

observability matrix

Theorem 12.3.18, 732

Fact 12.20.19, 793

positive-definite**matrix**

Theorem 12.3.18, 732

observable subspace**observable****eigenvalue**

Proposition 12.3.17, 732

observably**asymptotically stable****asymptotically stable****matrix**

Proposition 11.9.5, 735

Proposition 12.5.5, 735

block-triangular**matrix**

Proposition 12.4.3, 732

definition

Definition 12.4.1, 732

detectability

Proposition 12.5.5, 735

Lyapunov equation

Proposition 12.4.3, 732

observability**Gramian**

Proposition 12.4.3, 732

Proposition 12.4.4, 733

Proposition 12.4.5, 733

Proposition 12.4.6, 734

Proposition 12.4.7, 734

orthogonal matrix

Proposition 12.4.3, 732

output injection

Proposition 12.4.2, 732

rank

Proposition 12.4.4, 733

stabilizability

Proposition 11.9.5, 735

octahedral group**group**

Fact 3.21.7, 223

octonions**inequality**

Fact 1.14.1, 47

real matrix**representation**

Fact 3.22.1, 225

odd polynomial**asymptotically stable****polynomial**

Fact 11.17.6, 696

definition, 232**off-diagonal entries****definition, 80****off-diagonally located****block****definition, 80****OLHP****open left half plane****definition, 77****one-sided cone****definition, 89****induced by****antisymmetric****relation**

Proposition 2.3.6, 93

positive-semidefinite**matrix, 417****quadratic form**

Fact 8.14.14, 498

one-sided directional**differential****convex function**

Proposition 10.4.1, 626

definition, 625**example**

Fact 10.11.16, 639

homogeneity

- Fact 10.11.15, 639
- one-to-one**
 - definition, 3
 - inverse function
 - Theorem 1.2.2, 4
- one-to-one function**
 - composition of functions
 - Fact 1.5.16, 13
 - equivalent conditions
 - Fact 1.5.14, 12
 - finite domain
 - Fact 1.5.13, 12
- one-to-one matrix**
 - equivalent properties
 - Theorem 2.6.1, 98
 - nonsingular equivalence
 - Corollary 2.6.6, 101
- ones matrix**
 - definition, 84
 - rank
 - Fact 2.10.18, 117
- onto**
 - definition, 3
 - inverse function
 - Theorem 1.2.2, 4
- onto function**
 - composition of functions
 - Fact 1.5.16, 13
 - equivalent conditions
 - Fact 1.5.15, 13
 - finite domain
 - Fact 1.5.13, 12
- onto matrix**
 - equivalent properties
 - Theorem 2.6.1, 98
 - nonsingular equivalence
 - Corollary 2.6.6, 101
- open ball**
 - bounded set
 - Fact 10.8.2, 632
 - completely solid set
 - Fact 10.8.1, 632
- convex set**
 - Fact 10.8.1, 632
- inner product**
 - Fact 9.7.24, 570
- open ball of radius ε**
 - definition, 621
- open half space**
 - affine open half space
 - Fact 2.9.6, 111
 - definition, 91
- open mapping theorem**
 - open set image
 - Theorem 10.3.6, 624
- open relative to a set**
 - continuous function
 - Theorem 10.3.4, 624
 - definition
 - Definition 10.1.2, 621
- open set**
 - complement
 - Fact 10.8.4, 632
 - continuous function
 - Theorem 10.3.7, 624
 - Corollary 10.3.5, 624
 - convex hull
 - Fact 10.8.14, 633
 - definition
 - Definition 10.1.1, 621
 - intersection
 - Fact 10.9.9, 635
 - invariance of domain
 - Theorem 10.3.7, 624
 - right-invertible matrix
 - Theorem 10.3.6, 624
 - union
 - Fact 10.9.9, 635
- Oppenheim's inequality**
 - determinant inequality
 - Fact 8.21.19, 534
- optimal 2-uniform convexity**
 - powers
 - Fact 1.10.15, 33
 - Fact 9.9.35, 586
- order**
 - definition, 79
 - Definition 12.9.2, 749
- ORHP**
 - open right half plane
 - definition, 77
- Orlicz**
 - Hölder-induced norm
 - Fact 9.8.18, 574
- orthogonal**
 - complement
 - definition, 91
 - intersection
 - Fact 2.9.15, 112
 - projector
 - Proposition 3.5.2, 175
 - subspace
 - Proposition 3.5.2, 175
 - Fact 2.9.16, 112
 - Fact 2.9.27, 114
 - sum
 - Fact 2.9.15, 112
- orthogonal**
 - eigenvectors
 - normal matrix
 - Corollary 5.4.8, 294
- orthogonal matrices and matrix exponentials**
 - Davenport
 - Fact 11.11.13, 675
- orthogonal matrix, see unitary matrix**
 - 2×2
 - parameterization
 - Fact 3.11.6, 190
 - 3×3 skew-symmetric matrix
 - Fact 11.11.10, 674
 - Fact 11.11.11, 674
- additive**
 - decomposition
 - Fact 5.19.2, 360

- algebraic multiplicity**
 - Fact 5.11.2, 321
- Cayley transform**
 - Fact 3.11.8, 190
 - Fact 3.11.28, 196
 - Fact 3.11.30, 197
 - Fact 3.11.31, 198
- controllable dynamics**
 - Theorem 12.6.8, 739
- controllable subspace**
 - Proposition 12.6.9, 739
- controllably asymptotically stable**
 - Proposition 12.7.3, 743
- convex combination**
 - Fact 5.19.3, 360
- cross product**
 - Fact 3.10.2, 189
 - Fact 3.10.3, 189
 - Fact 3.11.8, 190
- cross-product matrix**
 - Fact 11.11.12, 674
 - Fact 11.11.13, 675
- definition**
 - Definition 3.1.1, 165
- detectability**
 - Proposition 12.5.4, 735
- determinant**
 - Fact 3.11.21, 196
 - Fact 3.11.22, 196
- direction cosines**
 - Fact 3.11.10, 192
- eigenvalue**
 - Fact 5.11.2, 321
- elementary reflector**
 - Fact 5.15.15, 347
- Euler parameters**
 - Fact 3.11.10, 192
 - Fact 3.11.11, 193
- existence of transformation**
 - Fact 3.9.5, 186
- factorization**
 - Fact 5.15.15, 347
 - Fact 5.15.16, 347
 - Fact 5.15.31, 350
 - Fact 5.15.35, 351
- group**
 - Proposition 3.3.6, 172
- Hamiltonian matrix**
 - Fact 3.19.13, 217
- Kronecker permutation matrix**
 - Fact 7.4.29, 407
- Kronecker product**
 - Fact 7.4.16, 406
- logarithm**
 - Fact 11.15.10, 692
- matrix exponential**
 - Proposition 11.6.7, 659
 - Fact 11.11.6, 673
 - Fact 11.11.7, 673
 - Fact 11.11.8, 674
 - Fact 11.11.9, 674
 - Fact 11.11.10, 674
 - Fact 11.11.11, 674
 - Fact 11.11.12, 674
 - Fact 11.11.13, 675
 - Fact 11.15.10, 692
- observable dynamics**
 - Theorem 12.3.8, 730
- observably asymptotically stable**
 - Proposition 12.4.3, 732
- orthosymplectic matrix**
 - Fact 3.19.13, 217
- parameterization**
 - Fact 3.11.9, 191
 - Fact 3.11.10, 192
- partitioned matrix**
 - Fact 3.11.27, 196
- permutation matrix**
 - Proposition 3.1.6, 169
- quaternions**
 - Fact 3.11.10, 192
- reflector**
 - Fact 3.11.9, 191
 - Fact 5.15.31, 350
 - Fact 5.15.35, 351
- Rodrigues**
 - Fact 3.11.10, 192
- Rodrigues's formulas**
 - Fact 3.11.11, 193
- rotation matrix**
 - Fact 3.11.9, 191
 - Fact 3.11.10, 192
 - Fact 3.11.11, 193
 - Fact 3.11.12, 194
 - Fact 3.11.31, 198
- skew-symmetric matrix**
 - Fact 3.11.28, 196
 - Fact 3.11.30, 197
 - Fact 3.11.31, 198
- SO(3)**
 - Fact 3.11.7, 190
- square root**
 - Fact 8.9.26, 453
- stabilizability**
 - Proposition 12.8.4, 747
- subspace**
 - Fact 3.11.1, 189
 - Fact 3.11.2, 189
- trace**
 - Fact 3.11.17, 195
 - Fact 3.11.18, 195
 - Fact 5.12.9, 334
 - Fact 5.12.10, 334
- unobservable subspace**
 - Proposition 12.3.9, 730
- orthogonal projector, see projector**
- orthogonal vectors**
 - norm inequality
 - Fact 9.7.25, 570
- unitary matrix**
 - Fact 3.11.14, 194
- vector sum and difference**
 - Fact 2.12.2, 126
- orthogonality**
 - single complex matrix
 - Lemma 2.2.4, 87
 - single complex vector
 - Lemma 2.2.2, 85
 - single real matrix
 - Lemma 2.2.3, 86
 - single real vector
 - Lemma 2.2.1, 85

orthogonality of complex matrices
definition, 87

orthogonality of complex vectors
definition, 85

orthogonality of real matrices
definition, 86

orthogonality of real vectors
definition, 85

orthogonally complementary subspaces
definition, 91
orthogonal complement
Proposition 2.3.3, 91

orthogonally similar matrices
definition
Definition 3.4.4, 174

diagonal matrix
Fact 5.9.15, 313

skew-symmetric matrix
Fact 5.14.33, 343

symmetric matrix
Fact 5.9.15, 313

upper block-triangular matrix
Corollary 5.4.2, 293
upper triangular matrix
Corollary 5.4.3, 293

orthosymplectic matrix group
Proposition 3.3.6, 172
Hamiltonian matrix
Fact 3.19.13, 217
orthogonal matrix
Fact 3.19.13, 217

oscillator
companion matrix

Fact 5.14.35, 344
definition, 654

Ostrowski
inertia of a
Hermitian matrix
Fact 12.21.5, 794
quantitative form of
Sylvester's law of
inertia
Fact 5.8.17, 310

Ostrowski-Taussky inequality
determinant
Fact 8.13.2, 485

OD
open unit disk
definition, 670

outbound Laplacian matrix
adjacency matrix
Theorem 3.2.2, 170
definition
Definition 3.2.1, 170

outdegree
graph
Definition 1.4.3, 9

outdegree matrix
definition
Definition 3.2.1, 170

outer-product matrix
algebraic multiplicity
Fact 5.14.3, 338
characteristic polynomial
Fact 4.9.16, 262
Fact 4.9.18, 263

cross product
Fact 3.11.8, 190

defective matrix
Fact 5.14.3, 338
definition, 86
Definition 3.1.2, 166

doublet
Fact 2.10.24, 118
Fact 2.12.6, 126
Euclidean norm

Fact 9.7.27, 570
existence of transformation
Fact 3.9.1, 185

Frobenius norm
Fact 9.7.26, 570

generalized inverse
Fact 6.3.2, 370

group-invertible matrix
Fact 5.14.3, 338

Hermitian matrix
Fact 3.7.18, 181
Fact 3.9.2, 185

idempotent matrix
Fact 3.7.18, 181
Fact 3.12.6, 199

identity
Fact 2.12.3, 126
Fact 2.12.5, 126

index of a matrix
Fact 5.14.3, 338

Kronecker product
Proposition 7.1.8, 401
linear independence

Fact 2.12.4, 126
Fact 2.12.8, 126

matrix exponential
Fact 11.11.1, 671

matrix power
Fact 2.12.7, 126

maximum singular value

Fact 5.11.16, 324
Fact 5.11.18, 324
Fact 9.7.26, 570

nilpotent matrix
Fact 5.14.3, 338

null space
Fact 2.10.11, 116

partitioned matrix
Fact 4.9.18, 263

positive-definite matrix
Fact 3.9.3, 186

positive-semidefinite matrix
Fact 8.9.2, 450
Fact 8.9.3, 450

Fact 8.9.4, 450
 Fact 8.15.2, 500
 Fact 8.15.3, 500
quadratic form
 Fact 9.13.3, 602
range
 Fact 2.10.11, 116
rank
 Fact 2.10.19, 117
 Fact 2.10.24, 118
 Fact 3.7.17, 181
 Fact 3.12.6, 199
semisimple matrix
 Fact 5.14.3, 338
singular value
 Fact 5.11.17, 324
skew-Hermitian matrix
 Fact 3.7.17, 181
 Fact 3.9.4, 186
spectral abscissa
 Fact 5.11.13, 323
spectral radius
 Fact 5.11.13, 323
spectrum
 Fact 5.11.13, 323
 Fact 5.14.1, 338
sum
 Fact 2.10.24, 118
trace
 Fact 5.14.3, 338
unitarily invariant norm
 Fact 9.8.40, 578

outer-product perturbation
 adjugate
 Fact 2.16.3, 141
 determinant
 Fact 2.16.3, 141
 elementary matrix
 Fact 3.7.19, 181
 generalized inverse
 Fact 6.4.2, 377
 inverse matrix
 Fact 2.16.3, 141
 matrix power
 Fact 2.12.18, 127
 rank

Fact 2.10.25, 118
 Fact 6.4.2, 377
unitary matrix
 Fact 3.11.15, 194
output convergence
 detectability
 Fact 12.20.2, 791
output equation
 definition, 725
output feedback
 characteristic polynomial
 Fact 12.22.13, 800
 determinant
 Fact 12.22.13, 800

output injection
 detectability
 Proposition 12.5.2, 734
 observably asymptotically stable
 Proposition 12.4.2, 732

ovals of Cassini
 spectrum bounds
 Fact 4.10.21, 271

Ozeki's inequality
 reversed
 Cauchy-Schwarz inequality
 Fact 1.16.23, 65

P

parallel affine subspaces
 definition, 89

parallel interconnection
 definition, 770
 transfer function
 Proposition 12.13.2, 770

parallel sum
 definition
 Fact 8.20.18, 528

parallelepiped volume
 Fact 2.20.16, 160
 Fact 2.20.17, 160

parallelogram
 area
 Fact 2.20.17, 160
 Fact 9.7.5, 565
 bivector
 Fact 9.7.5, 565
 cross product
 Fact 9.7.5, 565

parallelogram law
 complex numbers
 Fact 1.18.2, 69
 vector identity
 Fact 9.7.4, 563

parent
 Definition 1.4.2, 8

Parker
 equal diagonal entries by unitary similarity
 Fact 5.9.17, 313

Parodi
 polynomial root bound
 Fact 11.20.9, 710

Parrott's theorem
 maximum singular value of a partitioned matrix
 Fact 9.14.13, 610

Parseval's inequality
 norm inequality
 Fact 9.7.4, 563

Parseval's theorem
 Fourier transform
 Fact 12.22.1, 798
 H_2 norm
 Theorem 12.11.3, 766

partial derivative
 definition, 625

partial isometry

- generalized inverse**
 - Fact 6.3.28, 375
- partial ordering**
 - definition**
 - Definition 1.3.8, 7
 - generalized Löwner ordering**
 - Fact 8.19.10, 524
 - planar case**
 - Fact 1.5.7, 12
 - positive-semidefinite matrix**
 - Proposition 8.1.1, 417
 - rank subtractivity**
 - Fact 2.10.32, 119
- partition**
 - definition, 3**
 - equivalence relation**
 - Theorem 1.3.7, 7
- partitioned matrix**
 - adjugate**
 - Fact 2.14.27, 139
 - characteristic polynomial**
 - Fact 4.9.14, 262
 - Fact 4.9.15, 262
 - Fact 4.9.17, 263
 - Fact 4.9.18, 263
 - Fact 4.9.22, 264
 - Fact 4.9.23, 264
 - column norm**
 - Fact 9.8.11, 572
 - complementary subspaces**
 - Fact 3.12.33, 205
 - complex conjugate**
 - Fact 2.19.9, 153
 - complex conjugate transpose**
 - Proposition 2.8.1, 106
 - Fact 6.5.3, 386
 - complex matrix**
 - Fact 2.19.4, 152
 - Fact 2.19.5, 152
 - Fact 2.19.6, 152
 - Fact 2.19.7, 153
 - Fact 3.11.27, 196
 - contractive matrix**
 - Fact 8.11.24, 473
 - damping matrix**
 - Fact 5.12.21, 337
 - defect**
 - Fact 2.11.3, 121
 - Fact 2.11.8, 122
 - Fact 2.11.11, 123
 - definition, 80**
 - determinant**
 - Proposition 2.8.1, 106
 - Corollary 2.8.5, 107
 - Lemma 8.2.6, 421
 - Fact 2.14.2, 133
 - Fact 2.14.3, 133
 - Fact 2.14.4, 133
 - Fact 2.14.5, 134
 - Fact 2.14.6, 134
 - Fact 2.14.7, 134
 - Fact 2.14.9, 134
 - Fact 2.14.10, 135
 - Fact 2.14.11, 135
 - Fact 2.14.13, 135
 - Fact 2.14.14, 136
 - Fact 2.14.15, 136
 - Fact 2.14.16, 136
 - Fact 2.14.17, 136
 - Fact 2.14.18, 137
 - Fact 2.14.19, 137
 - Fact 2.14.20, 137
 - Fact 2.14.21, 137
 - Fact 2.14.22, 138
 - Fact 2.14.23, 138
 - Fact 2.14.24, 138
 - Fact 2.14.25, 138
 - Fact 2.14.26, 139
 - Fact 2.14.28, 139
 - Fact 2.17.5, 147
 - Fact 2.19.3, 151
 - Fact 2.19.9, 153
 - Fact 5.12.21, 337
 - Fact 6.5.26, 392
 - Fact 6.5.27, 392
 - Fact 6.5.28, 393
 - Fact 8.13.35, 492
 - Fact 8.13.36, 492
 - Fact 8.13.38, 492
 - Fact 8.13.39, 493
 - Fact 8.13.40, 493
 - Fact 8.13.41, 493
 - Fact 8.13.42, 493**
 - determinant of block 2×2**
 - Proposition 2.8.3, 107
 - Proposition 2.8.4, 107
 - discrete-time asymptotically stable matrix**
 - Fact 11.21.9, 713
 - Drazin generalized inverse**
 - Fact 6.6.1, 393
 - Fact 6.6.2, 393
 - eigenvalue**
 - Proposition 5.6.6, 303
 - Fact 5.12.20, 337
 - Fact 5.12.21, 337
 - Fact 5.12.22, 338
 - eigenvalue perturbation**
 - Fact 4.10.27, 272
 - factorization, 420**
 - Fact 2.14.9, 134
 - Fact 2.16.2, 141
 - Fact 2.17.3, 147
 - Fact 2.17.4, 147
 - Fact 2.17.5, 147
 - Fact 6.5.25, 392
 - Fact 8.11.25, 473
 - Fact 8.11.26, 473
 - factorization of block 2×2**
 - Proposition 2.8.3, 107
 - Proposition 2.8.4, 107
 - generalized inverse**
 - Fact 6.3.30, 376
 - Fact 6.5.1, 385
 - Fact 6.5.2, 386
 - Fact 6.5.3, 386
 - Fact 6.5.4, 386
 - Fact 6.5.13, 388
 - Fact 6.5.17, 390
 - Fact 6.5.18, 390
 - Fact 6.5.19, 390
 - Fact 6.5.20, 391
 - Fact 6.5.21, 391
 - Fact 6.5.22, 391
 - Fact 6.5.23, 391
 - Fact 6.5.24, 391

- Fact 8.20.22, 530
- geometric multiplicity**
 - Proposition 5.5.14, 298
- Hamiltonian matrix**
 - Proposition 3.1.7, 169
 - Fact 3.19.6, 216
 - Fact 3.19.8, 217
 - Fact 4.9.22, 264
 - Fact 5.12.21, 337
- Hermitian matrix**
 - Fact 3.7.27, 182
 - Fact 4.10.27, 272
 - Fact 5.8.19, 310
 - Fact 5.12.1, 333
 - Fact 6.5.5, 386
- Hölder-induced norm**
 - Fact 9.8.11, 572
- idempotent matrix**
 - Fact 3.12.14, 200
 - Fact 3.12.20, 201
 - Fact 3.12.33, 205
 - Fact 5.10.22, 320
- index of a matrix**
 - Fact 5.14.32, 343
 - Fact 6.6.13, 395
- inertia**
 - Fact 5.8.19, 310
 - Fact 5.8.20, 310
 - Fact 5.8.21, 311
 - Fact 5.12.1, 333
 - Fact 6.5.5, 386
- inverse matrix**
 - Fact 2.16.4, 142
 - Fact 2.17.2, 146
 - Fact 2.17.3, 147
 - Fact 2.17.4, 147
 - Fact 2.17.5, 147
 - Fact 2.17.6, 148
 - Fact 2.17.8, 148
 - Fact 2.17.9, 148
 - Fact 5.12.21, 337
- inverse of block 2×2**
 - Proposition 2.8.7, 108
 - Corollary 2.8.9, 109
- involutory matrix**
 - Fact 3.15.5, 212
- Kronecker product**
 - Fact 7.4.18, 406
 - Fact 7.4.19, 406
 - Fact 7.4.24, 407
- mass matrix**
 - Fact 5.12.21, 337
- matricial norm**
 - Fact 9.10.1, 593
- matrix exponential**
 - Fact 11.11.2, 672
 - Fact 11.14.1, 681
- matrix sign function**
 - Fact 10.10.3, 637
- maximum eigenvalue**
 - Fact 5.12.20, 337
- maximum singular value**
 - Fact 8.17.3, 508
 - Fact 8.17.14, 512
 - Fact 8.18.1, 512
 - Fact 8.18.2, 513
 - Fact 9.10.1, 593
 - Fact 9.10.3, 594
 - Fact 9.10.4, 594
 - Fact 9.10.5, 595
 - Fact 9.14.12, 610
 - Fact 9.14.13, 610
 - Fact 9.14.14, 610
- minimal polynomial**
 - Fact 4.10.12, 268
- minimal-rank identity**
 - Fact 6.5.7, 387
- minimum eigenvalue**
 - Fact 5.12.20, 337
- multiplicative identities, 82**
- nilpotent matrix**
 - Fact 3.12.14, 200
 - Fact 5.10.23, 321
- norm**
 - Fact 9.10.1, 593
 - Fact 9.10.2, 593
 - Fact 9.10.8, 596
- norm-compression inequality**
 - Fact 9.10.1, 593
 - Fact 9.10.8, 596
- normal matrix**
 - Fact 3.12.14, 200
- Fact 8.11.12, 470
- null space**
 - Fact 2.11.3, 121
- orthogonal matrix**
 - Fact 3.11.27, 196
- outer-product matrix**
 - Fact 4.9.18, 263
- polynomial**
 - Fact 4.10.10, 267
- positive-definite matrix**
 - Proposition 8.2.4, 420
 - Proposition 8.2.5, 420
 - Lemma 8.2.6, 421
 - Fact 8.9.18, 452
 - Fact 8.11.1, 467
 - Fact 8.11.2, 467
 - Fact 8.11.5, 468
 - Fact 8.11.8, 469
 - Fact 8.11.10, 469
 - Fact 8.11.13, 470
 - Fact 8.11.29, 474
 - Fact 8.11.30, 475
 - Fact 8.13.21, 488
 - Fact 8.17.14, 512
 - Fact 8.21.6, 532
 - Fact 11.21.9, 713
- positive-semidefinite matrix**
 - Proposition 8.2.3, 420
 - Proposition 8.2.4, 420
 - Corollary 8.2.2, 419
 - Lemma 8.2.1, 419
 - Lemma 8.2.6, 421
 - Fact 5.12.22, 338
 - Fact 8.7.6, 443
 - Fact 8.9.18, 452
 - Fact 8.11.1, 467
 - Fact 8.11.2, 467
 - Fact 8.11.5, 468
 - Fact 8.11.6, 469
 - Fact 8.11.7, 469
 - Fact 8.11.8, 469
 - Fact 8.11.9, 469
 - Fact 8.11.11, 469
 - Fact 8.11.12, 470
 - Fact 8.11.13, 470
 - Fact 8.11.14, 470

Fact 8.11.15, 470
 Fact 8.11.17, 471
 Fact 8.11.18, 471
 Fact 8.11.19, 471
 Fact 8.11.20, 472
 Fact 8.11.21, 472
 Fact 8.11.30, 475
 Fact 8.11.31, 475
 Fact 8.12.36, 483
 Fact 8.12.39, 484
 Fact 8.12.40, 484
 Fact 8.12.41, 484
 Fact 8.13.21, 488
 Fact 8.13.35, 492
 Fact 8.13.36, 492
 Fact 8.13.38, 492
 Fact 8.13.39, 493
 Fact 8.13.40, 493
 Fact 8.13.41, 493
 Fact 8.13.42, 493
 Fact 8.15.4, 500
 Fact 8.17.14, 512
 Fact 8.18.1, 512
 Fact 8.18.2, 513
 Fact 8.18.28, 521
 Fact 8.20.22, 530
 Fact 8.21.39, 539
 Fact 8.21.40, 539
 Fact 8.21.43, 540
 Fact 8.21.44, 540
 Fact 9.8.33, 576
 Fact 9.10.6, 595
 Fact 9.10.7, 596

power

Fact 2.12.21, 128

product

Fact 2.12.22, 128

projector

Fact 3.13.12, 208
 Fact 3.13.22, 210
 Fact 3.13.23, 210
 Fact 6.5.13, 388

quadratic form

Fact 8.15.5, 500
 Fact 8.15.6, 501

range

Fact 2.11.1, 120
 Fact 2.11.2, 121
 Fact 6.5.3, 386

rank

Corollary 2.8.5, 107
 Fact 2.11.6, 121
 Fact 2.11.8, 122
 Fact 2.11.9, 122
 Fact 2.11.10, 122
 Fact 2.11.11, 123
 Fact 2.11.12, 123
 Fact 2.11.13, 123
 Fact 2.11.14, 123
 Fact 2.11.15, 124
 Fact 2.11.16, 124
 Fact 2.11.18, 124
 Fact 2.11.19, 125
 Fact 2.14.4, 133
 Fact 2.14.5, 134
 Fact 2.14.11, 135
 Fact 2.17.5, 147
 Fact 2.17.10, 149
 Fact 3.12.20, 201
 Fact 3.13.12, 208
 Fact 3.13.22, 210
 Fact 5.12.21, 337
 Fact 6.3.30, 376
 Fact 6.5.6, 386
 Fact 6.5.7, 387
 Fact 6.5.8, 387
 Fact 6.5.9, 387
 Fact 6.5.10, 388
 Fact 6.5.12, 388
 Fact 6.5.13, 388
 Fact 6.5.14, 388
 Fact 6.5.15, 389
 Fact 6.6.2, 393
 Fact 8.7.6, 443
 Fact 8.7.7, 444

rank of block 2×2

Proposition 2.8.3, 107
 Proposition 2.8.4, 107

row norm

Fact 9.8.11, 572

Schatten norm

Fact 9.10.2, 593
 Fact 9.10.3, 594
 Fact 9.10.4, 594
 Fact 9.10.5, 595
 Fact 9.10.6, 595
 Fact 9.10.7, 596
 Fact 9.10.8, 596

Schur complement

Fact 6.5.4, 386
 Fact 6.5.5, 386
 Fact 6.5.6, 386
 Fact 6.5.8, 387
 Fact 6.5.12, 388
 Fact 6.5.29, 393
 Fact 8.21.39, 539

Schur product

Fact 8.21.6, 532
 Fact 8.21.39, 539
 Fact 8.21.40, 539

semicontractive matrix

Fact 8.11.6, 469
 Fact 8.11.22, 473

similar matrices

Fact 5.10.21, 320
 Fact 5.10.22, 320
 Fact 5.10.23, 321

singular value

Proposition 5.6.6, 303
 Fact 9.14.11, 609
 Fact 9.14.24, 613

skew-Hermitian matrix

Fact 3.7.27, 182

skew-symmetric matrix

Fact 3.11.27, 196

spectrum

Fact 2.19.3, 151
 Fact 4.10.25, 271
 Fact 4.10.26, 271

stability

Fact 11.18.38, 705

stiffness matrix

Fact 5.12.21, 337

Sylvester's equation

Fact 5.10.20, 320
 Fact 5.10.21, 320
 Fact 6.5.7, 387

symmetric matrix

Fact 3.11.27, 196

symplectic matrix

Fact 3.19.9, 217

trace

Proposition 2.8.1, 106
 Fact 8.12.36, 483

- Fact 8.12.39, 484
- Fact 8.12.40, 484
- Fact 8.12.41, 484
- Fact 8.12.42, 484
- transpose**
 - Proposition 2.8.1, 106
- unitarily invariant norm**
 - Fact 9.8.33, 576
- unitarily similar matrices**
 - Fact 5.9.23, 314
- unitary matrix**
 - Fact 3.11.24, 196
 - Fact 3.11.26, 196
 - Fact 3.11.27, 196
 - Fact 8.11.22, 473
 - Fact 8.11.23, 473
 - Fact 8.11.24, 473
 - Fact 9.14.11, 609
- partitioned**
 - positive-semidefinite matrix**
 - determinant**
 - Proposition 8.2.3, 420
 - rank**
 - Proposition 8.2.3, 420
- partitioned transfer function**
- H_2 norm**
 - Fact 12.22.16, 801
 - Fact 12.22.17, 801
- realization**
 - Proposition 12.13.3, 771
 - Fact 12.22.7, 799
- transfer function**
 - Fact 12.22.7, 799
- Pascal matrix**
 - positive-semidefinite matrix**
 - Fact 8.8.5, 447
 - Vandermonde matrix**
 - Fact 5.16.3, 354
- path**
 - definition**
 - Definition 1.4.3, 9
- pathwise connected**
 - continuous function**
 - Fact 10.11.5, 638
 - definition**
 - Definition 10.3.12, 625
 - group**
 - Proposition 11.6.8, 660
- Pauli spin matrices**
 - quaternions**
 - Fact 3.22.6, 227
- PBH test**
 - definition, 805**
- Pecaric**
 - Euclidean norm inequality**
 - Fact 9.7.8, 566
- Pedersen**
 - trace of a convex function**
 - Fact 8.12.33, 482
- Peierls-Bogoliubov inequality**
- matrix exponential**
 - Fact 11.14.29, 687
- pencil**
 - definition, 304**
 - deflating subspace**
 - Fact 5.13.1, 338
 - generalized eigenvalue**
 - Proposition 5.7.3, 305
 - Proposition 5.7.4, 306
 - invariant zero**
 - Corollary 12.10.4, 759
 - Corollary 12.10.5, 760
 - Corollary 12.10.6, 761
 - Kronecker canonical form**
 - Theorem 5.7.1, 304
 - Weierstrass canonical form**
 - Proposition 5.7.3, 305
- Penrose**
 - generalized inverse of a matrix sum**
 - Fact 6.4.34, 383
- period**
 - definition**
 - Definition 1.4.3, 9
 - graph**
 - Definition 1.4.3, 9
- permutation**
 - definition, 103**
- permutation group**
 - group**
 - Fact 3.21.7, 223
- permutation matrix**
 - circulant matrix**
 - Fact 5.16.8, 357
 - definition**
 - Definition 3.1.1, 165
 - determinant**
 - Fact 2.13.9, 130
 - irreducible matrix**
 - Fact 3.20.3, 217
 - orthogonal matrix**
 - Proposition 3.1.6, 169
 - spectrum**
 - Fact 5.16.8, 357
 - transposition matrix**
 - Fact 3.21.6, 222
- Perron-Frobenius theorem**
 - nonnegative matrix eigenvalues**
 - Fact 4.11.5, 273
- perturbation**
 - asymptotically stable matrix**
 - Fact 11.18.16, 700
 - inverse matrix**
 - Fact 9.9.60, 591
 - nonsingular matrix**
 - Fact 9.14.18, 611
- perturbed matrix**
 - spectrum**
 - Fact 4.10.3, 266
- Pesonen**
 - simultaneous diagonalization of symmetric matrices**
 - Fact 8.16.6, 507

Petrovich

complex inequality
Fact 1.18.2, 69

Pfaff's theorem

determinant of a skew-symmetric matrix
Fact 4.8.14, 259

Pfaffian

skew-symmetric matrix
Fact 4.8.14, 259

Pick matrix

positive-semidefinite matrix
Fact 8.8.17, 449

plane rotation

orthogonal matrix
Fact 5.15.16, 347

Poincaré separation theorem

eigenvalue inequality
Fact 8.17.16, 512

pointed cone

definition, 89
induced by reflexive relation
Proposition 2.3.6, 93
positive-semidefinite matrix, 417

polar

closed set
Fact 2.9.4, 110
convex cone
Fact 2.9.4, 110
definition, 91

polar cone

definition, 164

polar decomposition

existence
Corollary 5.6.5, 303

Frobenius norm

Fact 9.9.42, 588

matrix exponential

Fact 11.13.9, 679

normal matrix

Fact 5.18.8, 360
Fact 11.13.9, 679

uniqueness

Fact 5.18.2, 359
Fact 5.18.3, 359
Fact 5.18.4, 359
Fact 5.18.5, 359
Fact 5.18.6, 359
Fact 5.18.7, 359

unitarily invariant norm

Fact 9.9.42, 588

unitary matrix

Fact 5.18.8, 360

polarization identity

complex numbers
Fact 1.18.2, 69
norm identity
Fact 9.7.4, 563
vector identity
Fact 9.7.4, 563

polarized**Cayley-Hamilton theorem**

trace
Fact 4.9.3, 260

triple product identity

Fact 4.9.4, 260
Fact 4.9.6, 261

pole**minimal realization**

Fact 12.22.2, 798
Fact 12.22.12, 800

rational transfer function

Definition 4.7.4, 249

Smith-McMillan form

Proposition 4.7.11, 251

pole of a rational function

definition
Definition 4.7.1, 249

pole of a transfer function**definition**

Definition 4.7.10, 251

Polya's inequality

logarithmic mean
Fact 1.10.36, 37

Polya-Szego inequality

reversed
Cauchy-Schwarz inequality
Fact 1.16.21, 64

polygon

area
Fact 2.20.14, 159

polygonal inequalities

Euclidean norm
Fact 9.7.4, 563
Fact 9.7.7, 565

polyhedral convex cone

definition, 90

polynomial

asymptotically stable
Definition 11.8.3, 663

Bezout matrix

Fact 4.8.6, 255
Fact 4.8.8, 257

bound

Fact 11.20.13, 712

continuity of roots

Fact 10.11.2, 638

coprime

Fact 4.8.3, 254
Fact 4.8.4, 254
Fact 4.8.5, 255

definition, 231**Descartes rule of signs**

Fact 11.17.1, 695

discrete-time

asymptotically stable
Definition 11.10.3, 671

discrete-time

Lyapunov stable
Definition 11.10.3, 671

- discrete-time semistable**
 - Definition 11.10.3, 671
- Fujiwara's bound**
 - Fact 11.20.8, 710
- greatest common divisor**
 - Fact 4.8.5, 255
- interpolation**
 - Fact 4.8.11, 259
- Kojima's bound**
 - Fact 11.20.8, 710
- least common multiple**
 - Fact 4.8.3, 254
- Lyapunov stable**
 - Definition 11.8.3, 663
- partitioned matrix**
 - Fact 4.10.10, 267
- root bound**
 - Fact 11.20.4, 709
 - Fact 11.20.5, 709
 - Fact 11.20.6, 709
 - Fact 11.20.7, 710
 - Fact 11.20.8, 710
 - Fact 11.20.9, 710
 - Fact 11.20.10, 711
- root bounds**
 - Fact 11.20.11, 711
 - Fact 11.20.12, 712
- roots**
 - Fact 4.8.1, 253
 - Fact 4.8.2, 254
- roots of derivative**
 - Fact 10.11.3, 638
- semistable**
 - Definition 11.8.3, 663
- spectrum**
 - Fact 4.10.9, 267
 - Fact 4.10.10, 267
- Vandermonde matrix**
 - Fact 5.16.6, 355
- polynomial bound**
 - Huygens**
 - Fact 11.20.13, 712
 - Mihet**
 - Fact 11.20.13, 712
- polynomial coefficients**
- asymptotically stable polynomial**
 - Fact 11.17.2, 696
 - Fact 11.17.3, 696
 - Fact 11.17.7, 697
 - Fact 11.17.8, 697
 - Fact 11.17.10, 697
 - Fact 11.17.11, 697
 - Fact 11.17.12, 697
- discrete-time asymptotically stable polynomial**
 - Fact 11.20.1, 708
 - Fact 11.20.2, 709
 - Fact 11.20.3, 709
- polynomial division**
 - quotient and remainder**
 - Lemma 4.1.2, 233
- polynomial matrix**
 - definition, 234**
 - matrix exponential**
 - Proposition 11.2.1, 646
 - Smith form**
 - Proposition 4.3.4, 237
- polynomial matrix division**
 - linear divisor**
 - Corollary 4.2.3, 235
 - Lemma 4.2.2, 235
- polynomial multiplication**
 - Toeplitz matrix**
 - Fact 4.8.10, 258
- polynomial representation**
 - commuting matrices**
 - Fact 5.14.22, 341
 - Fact 5.14.23, 342
 - Fact 5.14.24, 342
 - inverse matrix**
 - Fact 4.8.13, 259
- polynomial root**
 - maximum singular value bound**
 - Fact 9.13.14, 604
- minimum singular value bound**
 - Fact 9.13.14, 604
- polynomial root bound**
 - Bourbaki**
 - Fact 11.20.4, 709
 - Carmichael**
 - Fact 11.20.10, 711
 - Fujii-Kubo**
 - Fact 11.20.9, 710
 - Joyal**
 - Fact 11.20.7, 710
 - Labelle**
 - Fact 11.20.7, 710
 - Linden**
 - Fact 11.20.9, 710
 - Mason**
 - Fact 11.20.10, 711
 - Parodi**
 - Fact 11.20.9, 710
 - Rahman**
 - Fact 11.20.7, 710
 - Walsh**
 - Fact 11.20.5, 709
- polynomial root bounds**
 - Berwald**
 - Fact 11.20.11, 711
 - Cauchy**
 - Fact 11.20.11, 711
 - Cohn**
 - Fact 11.20.11, 711
- polynomial root locations**
 - Enestrom-Kakeya theorem**
 - Fact 11.20.3, 709
- polynomial roots**
 - Bezout matrix**
 - Fact 4.8.9, 258
 - Newton's identities**
 - Fact 4.8.2, 254
- polytope**
 - definition, 90**
- Popoviciu**

- arithmetic-mean–geometric-mean inequality
 - Fact 1.15.29, 55
- Popoviciu's inequality**
 - convex function
 - Fact 1.8.6, 22
- positive diagonal**
 - upper triangular matrix
 - Fact 5.15.9, 346
- positive matrix**
 - almost nonnegative matrix
 - Fact 11.19.2, 706
 - definition, 81
 - Definition 3.1.4, 168
 - eigenvalue
 - Fact 4.11.20, 280
 - Kronecker sum
 - Fact 7.5.8, 409
 - Schur product
 - Fact 7.6.13, 415
 - Fact 7.6.14, 415
 - spectral radius
 - Fact 7.6.14, 415
 - spectrum
 - Fact 5.11.12, 323
 - unstable matrix
 - Fact 11.18.20, 701
- positive vector**
 - definition, 79
 - null space
 - Fact 4.11.15, 279
- positive-definite matrix**
 - arithmetic mean
 - Fact 8.10.34, 460
 - arithmetic-mean–geometric-mean inequality
 - Fact 8.13.8, 486
 - asymptotically stable matrix
 - Proposition 11.9.5, 668
 - Proposition 12.4.9, 734
 - Corollary 11.9.7, 669
 - Fact 11.18.21, 701
 - Cauchy matrix
 - Fact 8.8.16, 449
 - Fact 12.21.18, 797
 - Cayley transform
 - Fact 8.9.30, 453
 - cogredient
 - diagonalization
 - Theorem 8.3.1, 423
 - Fact 8.16.5, 507
 - commuting matrices
 - Fact 8.9.40, 455
 - complex conjugate transpose
 - Fact 8.9.39, 455
 - complex matrix
 - Fact 3.7.9, 179
 - congruent matrices
 - Proposition 3.4.5, 174
 - Corollary 8.1.3, 419
 - contractive matrix
 - Fact 8.11.13, 470
 - contragredient
 - diagonalization
 - Theorem 8.3.2, 423
 - Corollary 8.3.3, 423
 - controllable pair
 - Theorem 12.6.18, 742
 - convex function
 - Fact 8.14.17, 499
 - definition
 - Definition 3.1.1, 165
 - determinant
 - Proposition 8.4.14, 429
 - Fact 8.12.1, 475
 - Fact 8.13.6, 486
 - Fact 8.13.7, 486
 - Fact 8.13.8, 486
 - Fact 8.13.9, 486
 - Fact 8.13.10, 487
 - Fact 8.13.12, 487
 - Fact 8.13.13, 487
 - Fact 8.13.14, 487
 - Fact 8.13.15, 488
 - Fact 8.13.17, 488
 - Fact 8.13.19, 488
 - Fact 8.13.21, 488
 - Fact 8.13.23, 489
 - discrete-time
 - asymptotically stable matrix
 - Proposition 11.10.5, 671
 - Fact 11.21.9, 713
 - Fact 11.21.15, 714
 - discrete-time
 - Lyapunov-stable matrix
 - Proposition 11.10.6, 671
 - dissipative matrix
 - Fact 8.17.12, 511
 - Fact 11.18.21, 701
 - eigenvalue
 - Fact 8.10.24, 458
 - Fact 8.15.20, 503
 - Fact 8.15.29, 505
 - Fact 8.15.30, 505
 - Fact 8.18.29, 521
 - Fact 8.21.21, 535
 - ellipsoid
 - Fact 3.7.35, 184
 - exponential
 - Fact 11.14.26, 686
 - factorization
 - Fact 5.15.26, 349
 - Fact 5.18.4, 359
 - Fact 5.18.5, 359
 - Fact 5.18.6, 359
 - Fact 5.18.8, 360
 - Furuta inequality
 - Fact 8.10.50, 465
 - generalized
 - geometric mean
 - Fact 8.10.45, 464
 - generalized inverse
 - Proposition 6.1.6, 365
 - Fact 6.4.7, 379
 - geometric mean
 - Fact 8.10.43, 461
 - Fact 8.10.46, 464
 - Fact 8.21.51, 541
 - group-invertible matrix
 - Fact 8.10.12, 457
 - Hermitian matrix
 - Fact 5.15.41, 351

- Fact 8.10.13, 457
- Fact 8.13.7, 486
- Hilbert matrix**
 - Fact 3.18.4, 215
- idempotent matrix**
 - Fact 5.15.30, 350
- identity**
 - Fact 8.10.6, 456
 - Fact 8.10.7, 456
- increasing function**
 - Fact 8.10.53, 466
- inequality**
 - Fact 8.9.41, 455
 - Fact 8.9.42, 455
 - Fact 8.10.8, 456
 - Fact 8.10.9, 457
 - Fact 8.10.19, 458
 - Fact 8.10.20, 458
 - Fact 8.10.21, 458
 - Fact 8.10.22, 458
 - Fact 8.10.28, 459
 - Fact 8.10.40, 461
 - Fact 8.10.48, 465
 - Fact 8.10.51, 466
 - Fact 8.11.27, 474
 - Fact 8.15.21, 503
 - Fact 8.15.22, 503
 - Fact 8.19.3, 523
 - Fact 8.21.42, 539
- inertia**
 - Fact 5.8.10, 308
- inertia matrix**
 - Fact 8.9.5, 451
- inner-product minimization**
 - Fact 8.15.12, 502
- integral**
 - Fact 8.15.32, 505
 - Fact 8.15.33, 506
 - Fact 8.15.34, 506
 - Fact 8.15.35, 506
- inverse**
 - Fact 8.11.10, 469
- inverse matrix**
 - Proposition 8.6.6, 432
 - Lemma 8.6.5, 432
 - Fact 8.9.17, 452
 - Fact 8.9.41, 455
- Kronecker product**
 - Fact 7.4.16, 406
- left inverse**
 - Fact 3.7.25, 182
- limit**
 - Fact 8.10.47, 465
- Lyapunov equation**
 - Fact 12.21.16, 797
 - Fact 12.21.18, 797
- Lyapunov-stable matrix**
 - Proposition 11.9.6, 669
 - Corollary 11.9.7, 669
- matrix exponential**
 - Proposition 11.2.8, 649
 - Proposition 11.2.9, 650
 - Fact 11.14.20, 685
 - Fact 11.14.22, 685
 - Fact 11.14.23, 686
 - Fact 11.15.1, 689
- matrix logarithm**
 - Proposition 8.6.4, 432
 - Proposition 11.4.5, 654
 - Fact 8.9.43, 455
 - Fact 8.13.8, 486
 - Fact 8.18.29, 521
 - Fact 8.19.1, 522
 - Fact 8.19.2, 523
 - Fact 9.9.55, 590
 - Fact 11.14.24, 686
 - Fact 11.14.25, 686
 - Fact 11.14.26, 686
 - Fact 11.14.27, 686
- matrix power**
 - Fact 8.10.41, 461
 - Fact 8.10.42, 461
- matrix product**
 - Corollary 8.3.6, 424
- matrix sign function**
 - Fact 10.10.4, 637
- maximum singular value**
 - Fact 8.18.8, 515
 - Fact 8.18.25, 520
- norm**
 - Fact 9.7.30, 571
- observable pair**
 - Theorem 12.3.18, 732
- outer-product matrix**
 - Fact 3.9.3, 186
- partitioned matrix**
 - Proposition 8.2.4, 420
 - Proposition 8.2.5, 420
 - Lemma 8.2.6, 421
 - Fact 8.9.18, 452
 - Fact 8.11.1, 467
 - Fact 8.11.2, 467
 - Fact 8.11.5, 468
 - Fact 8.11.8, 469
 - Fact 8.11.10, 469
 - Fact 8.11.13, 470
 - Fact 8.11.29, 474
 - Fact 8.11.30, 475
 - Fact 8.13.21, 488
 - Fact 8.17.14, 512
 - Fact 8.21.6, 532
 - Fact 11.21.9, 713
- positive-semidefinite matrix**
 - Fact 8.8.13, 448
 - Fact 8.8.14, 449
 - Fact 8.10.27, 458
 - Fact 8.12.25, 481
- power**
 - Fact 8.9.42, 455
 - Fact 8.10.38, 461
 - Fact 8.10.39, 461
 - Fact 8.10.48, 465
- power inequality**
 - Fact 8.10.52, 466
- properties of $<$ and $\leq$**
 - Proposition 8.1.2, 418
- quadratic form**
 - Fact 8.15.24, 504
 - Fact 8.15.25, 504
 - Fact 8.15.26, 504
 - Fact 8.15.29, 505
 - Fact 8.15.30, 505
 - Fact 9.8.37, 577
- quadratic form inequality**
 - Fact 8.15.4, 500
- regularized Tikhonov inverse**
 - Fact 8.9.40, 455
- Riccati equation**
 - Fact 12.23.4, 802

Schur product

- Fact 8.21.4, 531
- Fact 8.21.5, 532
- Fact 8.21.6, 532
- Fact 8.21.7, 533
- Fact 8.21.13, 533
- Fact 8.21.14, 534
- Fact 8.21.15, 534
- Fact 8.21.21, 535
- Fact 8.21.33, 538
- Fact 8.21.34, 538
- Fact 8.21.36, 538
- Fact 8.21.38, 539
- Fact 8.21.42, 539
- Fact 8.21.47, 540
- Fact 8.21.49, 541
- Fact 8.21.50, 541
- Fact 8.21.51, 541

simultaneous**diagonalization**

- Fact 8.16.5, 507

skew-Hermitian**matrix**

- Fact 8.13.6, 486
- Fact 11.18.12, 700

spectral order

- Fact 8.19.4, 523

spectral radius

- Fact 8.10.5, 456
- Fact 8.18.25, 520

spectrum

- Proposition 5.5.21, 300

strictly convex**function**

- Fact 8.14.15, 499
- Fact 8.14.16, 499

subdeterminant

- Proposition 8.2.8, 422
- Fact 8.13.17, 488

submatrix

- Proposition 8.2.8, 422
- Fact 8.11.28, 474

Toeplitz matrix

- Fact 8.13.13, 487

trace

- Proposition 8.4.14, 429
- Fact 8.9.16, 452
- Fact 8.10.46, 464
- Fact 8.11.10, 469

- Fact 8.12.1, 475

- Fact 8.12.2, 475

- Fact 8.12.24, 480

- Fact 8.12.27, 481

- Fact 8.12.37, 483

- Fact 8.13.12, 487

- Fact 11.14.24, 686

- Fact 11.14.25, 686

- Fact 11.14.27, 686

tridiagonal matrix

- Fact 8.8.18, 450

**unitarily similar
matrices**

- Proposition 3.4.5, 174
- Proposition 5.6.3, 302

upper bound

- Fact 8.10.31, 459

positive-definite matrix**product****inequality**

- Fact 8.10.43, 461
- Fact 8.10.45, 464

**positive-definite
solution****Riccati equation**

- Theorem 12.17.2, 782
- Proposition 12.19.3, 790
- Corollary 12.19.2, 790

**positive-semidefinite
function****positive-semidefinite
matrix**

- Fact 8.8.1, 444

**positive-semidefinite
matrix****absolute-value
matrix**

- Fact 8.9.1, 450

**antisymmetric
relation**

- Proposition 8.1.1, 417

average

- Fact 5.19.5, 360

Brownian motion

- Fact 8.8.4, 446

Cartesian**decomposition**

- Fact 9.9.40, 587

Cauchy matrix

- Fact 8.8.7, 447
- Fact 8.8.9, 448
- Fact 12.21.19, 797

**Cauchy-Schwarz
inequality**

- Fact 8.11.14, 470
- Fact 8.11.15, 470
- Fact 8.15.8, 501

closed set

- Fact 10.8.18, 633

cogredient**diagonalization**

- Theorem 8.3.4, 423

commuting matrices

- Fact 8.19.5, 467, 523

completely solid set

- Fact 10.8.18, 633

complex matrix

- Fact 3.7.9, 179

congruent matrices

- Proposition 3.4.5, 174
- Corollary 8.1.3, 419

contragredient**diagonalization**

- Theorem 8.3.5, 424
- Corollary 8.3.7, 424

controllability

- Fact 12.20.6, 791

convex combination

- Fact 5.19.6, 360
- Fact 8.13.16, 488

convex cone, 417**convex function**

- Fact 8.14.15, 499
- Fact 8.20.20, 530

convex set

- Fact 8.14.2, 494
- Fact 8.14.3, 495
- Fact 8.14.4, 495
- Fact 8.14.5, 495
- Fact 8.14.6, 496

copositive matrix

- Fact 8.15.37, 507

cosines

- Fact 8.8.15, 449

definition

Definition 3.1.1, 165

determinant

Corollary 8.4.15, 429

Fact 8.13.16, 488

Fact 8.13.18, 488

Fact 8.13.20, 488

Fact 8.13.21, 488

Fact 8.13.24, 489

Fact 8.13.29, 490

Fact 8.13.35, 492

Fact 8.13.36, 492

Fact 8.13.38, 492

Fact 8.13.39, 493

Fact 8.13.40, 493

Fact 8.13.41, 493

Fact 8.17.11, 511

Fact 8.18.30, 521

Fact 8.21.8, 533

Fact 8.21.19, 534

Fact 8.21.20, 535

Fact 9.8.39, 578

diagonal entries

Fact 8.9.8, 451

Fact 8.9.9, 451

diagonal entry

Fact 8.10.16, 457

Fact 8.12.3, 476

discrete-time**Lyapunov-stable matrix**

Fact 11.21.15, 714

Drazin generalized**inverse**

Fact 8.20.2, 525

eigenvalue

Fact 8.12.3, 476

Fact 8.15.11, 501

Fact 8.18.6, 514

Fact 8.18.19, 518

Fact 8.18.20, 518

Fact 8.18.22, 519

Fact 8.18.23, 519

Fact 8.18.24, 520

Fact 8.18.27, 521

Fact 8.20.17, 528

Fact 8.21.18, 534

Fact 8.21.20, 535

Euler totient**function**

Fact 8.8.5, 447

factorization

Fact 5.15.22, 349

Fact 5.15.26, 349

Fact 5.18.2, 359

Fact 5.18.3, 359

Fact 5.18.7, 359

Fact 8.9.36, 454

Fact 8.9.37, 454

Fejer's theorem

Fact 8.21.35, 538

Frobenius norm

Fact 9.8.39, 578

Fact 9.9.12, 581

Fact 9.9.15, 582

Fact 9.9.27, 584

Furuta inequality

Proposition 8.6.7, 433

generalized inverse

Proposition 6.1.6, 365

Fact 6.4.4, 378

Fact 8.20.1, 525

Fact 8.20.2, 525

Fact 8.20.3, 525

Fact 8.20.4, 525

Fact 8.20.6, 526

Fact 8.20.7, 526

Fact 8.20.8, 526

Fact 8.20.9, 526

Fact 8.20.10, 526

Fact 8.20.11, 527

Fact 8.20.13, 527

Fact 8.20.15, 527

Fact 8.20.16, 527

Fact 8.20.17, 528

Fact 8.20.18, 528

Fact 8.20.19, 530

Fact 8.20.20, 530

Fact 8.20.22, 530

Fact 8.20.23, 531

geometric mean

Fact 8.10.43, 461

group generalized**inverse**

Fact 8.20.1, 525

group-invertible**matrix**

Fact 8.10.12, 457

Hadamard-Fischer inequality

Fact 8.13.36, 492

Hermitian matrix

Fact 5.15.41, 351

Fact 8.9.11, 452

Fact 8.10.13, 457

Hölder's inequality

Fact 8.12.11, 477

Fact 8.12.12, 477

Hua's inequalities

Fact 8.11.21, 472

Hua's matrix**equality**

Fact 8.11.21, 472

idempotent matrix

Fact 5.15.30, 350

identity

Fact 8.11.16, 470

Fact 8.19.6, 523

increasing sequence

Proposition 8.6.3, 432

inequality

Proposition 8.6.7, 433

Corollary 8.6.8, 433

Corollary 8.6.9, 433

Fact 8.9.10, 451

Fact 8.9.19, 452

Fact 8.9.21, 452

Fact 8.9.38, 455

Fact 8.10.19, 458

Fact 8.10.20, 458

Fact 8.10.21, 458

Fact 8.10.28, 459

Fact 8.10.29, 459

Fact 8.10.30, 459

Fact 8.15.21, 503

Fact 8.15.22, 503

Fact 8.21.42, 539

Fact 9.14.22, 612

inertia

Fact 5.8.9, 308

Fact 5.8.10, 308

Fact 12.21.9, 796

integral

Proposition 8.6.10, 433

inverse matrix

Fact 8.10.37, 461

Kantorovich**inequality**

Fact 8.15.9, 501

kernel function

Fact 8.8.1, 444

Fact 8.8.2, 445

Kronecker product

Fact 7.4.16, 406

Fact 8.21.16, 534

Fact 8.21.22, 536

Fact 8.21.23, 536

Fact 8.21.24, 536

Fact 8.21.26, 536

Fact 8.21.27, 536

Fact 8.21.29, 536

Kronecker sum

Fact 7.5.8, 409

lattice

Fact 8.10.32, 459

Fact 8.10.33, 459

left-equivalent**matrices**

Fact 5.10.19, 319

Lehmer matrix

Fact 8.8.5, 447

limit

Proposition 8.6.3, 432

Fact 8.10.47, 465

linear combination

Fact 8.13.18, 488

log majorization

Fact 8.11.9, 469

Lyapunov equation

Fact 12.21.15, 797

Fact 12.21.19, 797

matrix exponential

Fact 11.14.20, 685

Fact 11.14.35, 688

Fact 11.16.6, 694

Fact 11.16.16, 695

matrix logarithm

Fact 9.9.54, 590

matrix power

Corollary 8.6.11, 434

Fact 8.9.14, 452

Fact 8.10.36, 461

Fact 8.10.49, 465

Fact 8.12.30, 482

Fact 8.15.13, 502

Fact 8.15.14, 502

Fact 8.15.15, 502

Fact 8.15.16, 502

Fact 9.9.17, 582

matrix product

Corollary 8.3.6, 424

maximum eigenvalue

Fact 8.18.14, 516

maximum singular**value**

Fact 8.18.1, 512

Fact 8.18.2, 513

Fact 8.18.11, 515

Fact 8.18.12, 516

Fact 8.18.13, 516

Fact 8.18.14, 516

Fact 8.18.15, 517

Fact 8.18.16, 517

Fact 8.18.25, 520

Fact 8.18.26, 520

Fact 8.18.28, 521

Fact 8.18.30, 521

Fact 8.18.31, 522

Fact 8.20.9, 526

Fact 11.16.6, 694

McCarthy inequality

Fact 8.12.29, 481

Minkowski's**inequality**

Fact 8.12.29, 481

norm-compression**inequality**

Fact 9.10.6, 595

normal matrix

Fact 8.9.22, 452

Fact 8.10.11, 457

Fact 8.11.12, 470

null space

Fact 8.7.3, 443

Fact 8.7.5, 443

Fact 8.15.1, 500

Fact 8.15.23, 504

one-sided cone, 417**outer-product**

Fact 8.9.3, 450

outer-product**matrix**

Fact 8.9.2, 450

Fact 8.9.4, 450

Fact 8.15.2, 500

Fact 8.15.3, 500

partial ordering

Proposition 8.1.1, 417

Fact 8.19.9, 524

partitioned matrix

Proposition 8.2.3, 420

Proposition 8.2.4, 420

Corollary 8.2.2, 419

Lemma 8.2.1, 419

Lemma 8.2.6, 421

Fact 5.12.22, 338

Fact 8.7.6, 443

Fact 8.9.18, 452

Fact 8.11.1, 467

Fact 8.11.2, 467

Fact 8.11.5, 468

Fact 8.11.6, 469

Fact 8.11.7, 469

Fact 8.11.8, 469

Fact 8.11.9, 469

Fact 8.11.11, 469

Fact 8.11.12, 470

Fact 8.11.13, 470

Fact 8.11.14, 470

Fact 8.11.15, 470

Fact 8.11.17, 471

Fact 8.11.18, 471

Fact 8.11.19, 471

Fact 8.11.20, 472

Fact 8.11.21, 472

Fact 8.11.30, 475

Fact 8.11.31, 475

Fact 8.12.36, 483

Fact 8.12.39, 484

Fact 8.12.40, 484

Fact 8.12.41, 484

Fact 8.13.21, 488

Fact 8.13.35, 492

Fact 8.13.36, 492

Fact 8.13.38, 492

Fact 8.13.39, 493

Fact 8.13.40, 493

Fact 8.13.41, 493

Fact 8.13.42, 493

Fact 8.15.4, 500

Fact 8.17.14, 512

Fact 8.18.1, 512

Fact 8.18.2, 513

- Fact 8.18.28, 521
- Fact 8.20.22, 530
- Fact 8.21.39, 539
- Fact 8.21.40, 539
- Fact 8.21.43, 540
- Fact 8.21.44, 540
- Fact 9.8.33, 576
- Fact 9.10.6, 595
- Fact 9.10.7, 596
- Pascal matrix**
 - Fact 8.8.5, 447
- Pick matrix**
 - Fact 8.8.17, 449
- pointed cone, 417**
- positive-definite matrix**
 - Fact 8.8.13, 448
 - Fact 8.8.14, 449
 - Fact 8.10.27, 458
 - Fact 8.12.25, 481
- positive-semidefinite function**
 - Fact 8.8.1, 444
- power**
 - Fact 8.10.38, 461
 - Fact 8.10.39, 461
- projector**
 - Fact 3.13.4, 207
- properties of $<$ and $\leq$**
 - Proposition 8.1.2, 418
- quadratic form inequality**
 - Fact 8.15.4, 500
 - Fact 8.15.7, 501
- range**
 - Theorem 8.6.2, 431
 - Corollary 8.2.2, 419
 - Fact 8.7.1, 443
 - Fact 8.7.2, 443
 - Fact 8.7.3, 443
 - Fact 8.7.4, 443
 - Fact 8.7.5, 443
 - Fact 8.10.2, 456
 - Fact 8.20.7, 526
 - Fact 8.20.8, 526
 - Fact 8.20.10, 526
 - Fact 8.20.11, 527
- range-Hermitian matrix**
 - Fact 8.20.21, 530
- rank**
 - Fact 5.8.9, 308
 - Fact 8.7.1, 443
 - Fact 8.7.5, 443
 - Fact 8.7.6, 443
 - Fact 8.7.7, 444
 - Fact 8.10.2, 456
 - Fact 8.10.14, 457
 - Fact 8.20.11, 527
 - Fact 8.21.16, 534
- rank subtractivity**
- partial ordering**
 - Fact 8.19.5, 523
 - Fact 8.20.7, 526
 - Fact 8.20.8, 526
- real eigenvalues**
 - Fact 5.14.13, 340
- reflexive relation**
 - Proposition 8.1.1, 417
- reproducing kernel space**
 - Fact 8.8.2, 445
- right inverse**
 - Fact 3.7.26, 182
- Schatten norm**
 - Fact 9.9.22, 583
 - Fact 9.9.39, 587
 - Fact 9.9.40, 587
 - Fact 9.10.6, 595
- Fact 9.10.7, 596
- Schur complement**
 - Corollary 8.6.18, 442
 - Fact 8.11.3, 468
 - Fact 8.11.4, 468
 - Fact 8.11.17, 471
 - Fact 8.11.18, 471
 - Fact 8.11.19, 471
 - Fact 8.11.20, 472
 - Fact 8.11.27, 474
 - Fact 8.20.19, 530
 - Fact 8.21.11, 533
- Schur inverse**
 - Fact 8.21.1, 531
- Schur power**
 - Fact 8.21.2, 531
 - Fact 8.21.3, 531
 - Fact 8.21.25, 536
- Schur product**
 - Fact 8.21.4, 531
 - Fact 8.21.7, 533
 - Fact 8.21.11, 533
 - Fact 8.21.12, 533
 - Fact 8.21.14, 534
 - Fact 8.21.17, 534
 - Fact 8.21.18, 534
 - Fact 8.21.20, 535
 - Fact 8.21.22, 536
 - Fact 8.21.23, 536
 - Fact 8.21.31, 537
 - Fact 8.21.35, 538
 - Fact 8.21.37, 538
 - Fact 8.21.39, 539
 - Fact 8.21.40, 539
 - Fact 8.21.41, 539
 - Fact 8.21.42, 539
 - Fact 8.21.43, 540
 - Fact 8.21.44, 540
 - Fact 8.21.45, 540
 - Fact 8.21.46, 540
- semicontractive matrix**
 - Fact 8.11.6, 469
 - Fact 8.11.13, 470
- semisimple matrix**
 - Corollary 8.3.6, 424
- shorted operator**
 - Fact 8.20.19, 530
- signature**

- Fact 5.8.9, 308
- singular value**
 - Fact 8.18.7, 514
 - Fact 9.14.27, 613
- singular values**
 - Fact 8.11.9, 469
- skew-Hermitian matrix**
 - Fact 8.9.12, 452
- spectral order**
 - Fact 8.19.4, 523
- spectral radius**
 - Fact 8.18.25, 520
 - Fact 8.20.8, 526
- spectrum**
 - Proposition 5.5.21, 300
 - Fact 8.20.16, 527
- square root**
 - Fact 8.9.6, 451
 - Fact 8.10.18, 458
 - Fact 8.10.26, 458
 - Fact 8.21.29, 536
 - Fact 9.8.32, 576
- stabilizability**
 - Fact 12.20.6, 791
- star partial ordering**
 - Fact 8.19.8, 524
 - Fact 8.20.8, 526
- structured matrix**
 - Fact 8.8.2, 445
 - Fact 8.8.3, 446
 - Fact 8.8.4, 446
 - Fact 8.8.5, 447
 - Fact 8.8.6, 447
 - Fact 8.8.7, 447
 - Fact 8.8.8, 447
 - Fact 8.8.9, 448
 - Fact 8.8.10, 448
 - Fact 8.8.11, 448
 - Fact 8.8.12, 448
- subdeterminant**
 - Proposition 8.2.7, 421
- submatrix**
 - Proposition 8.2.7, 421
 - Fact 8.7.7, 444
 - Fact 8.13.36, 492
- submultiplicative norm**
 - Fact 9.9.7, 580
- Szasz's inequality**
 - Fact 8.13.36, 492
- trace**
 - Proposition 8.4.13, 428
 - Fact 2.12.16, 127
 - Fact 8.12.3, 476
 - Fact 8.12.9, 477
 - Fact 8.12.10, 477
 - Fact 8.12.11, 477
 - Fact 8.12.12, 477
 - Fact 8.12.13, 477
 - Fact 8.12.17, 478
 - Fact 8.12.18, 478
 - Fact 8.12.19, 479
 - Fact 8.12.20, 479
 - Fact 8.12.21, 480
 - Fact 8.12.22, 480
 - Fact 8.12.23, 480
 - Fact 8.12.24, 480
 - Fact 8.12.26, 481
 - Fact 8.12.28, 481
 - Fact 8.12.29, 481
 - Fact 8.12.34, 483
 - Fact 8.12.35, 483
 - Fact 8.12.36, 483
 - Fact 8.12.38, 483
 - Fact 8.12.39, 484
 - Fact 8.12.40, 484
 - Fact 8.12.41, 484
 - Fact 8.13.20, 488
 - Fact 8.18.16, 517
 - Fact 8.18.20, 518
 - Fact 8.20.3, 525
 - Fact 8.20.17, 528
- trace norm**
 - Fact 9.9.15, 582
- transitive relation**
 - Proposition 8.1.1, 417
- triangle inequality**
 - Fact 9.9.21, 583
- unitarily invariant norm**
 - Fact 9.9.7, 580
 - Fact 9.9.14, 582
 - Fact 9.9.15, 582
 - Fact 9.9.16, 582
 - Fact 9.9.17, 582
 - Fact 9.9.27, 584
 - Fact 9.9.46, 588
- Fact 9.9.51, 589
- Fact 9.9.52, 590
- Fact 9.9.53, 590
- Fact 9.9.54, 590
- Fact 11.16.16, 695
- Fact 11.16.17, 695
- unitarily**
 - left-equivalent matrices**
 - Fact 5.10.18, 319
 - Fact 5.10.19, 319
 - unitarily**
 - right-equivalent matrices**
 - Fact 5.10.18, 319
 - unitarily similar matrices**
 - Proposition 3.4.5, 174
 - Proposition 5.6.3, 302
- upper bound**
 - Fact 8.10.35, 460
- upper triangular matrix**
 - Fact 8.9.37, 454
- weak majorization**
 - Fact 8.18.6, 514
- Young's inequality**
 - Fact 8.12.12, 477
- zero matrix**
 - Fact 8.10.10, 457
- positive-semidefinite matrix determinant**
- Fischer's inequality**
 - Fact 8.13.35, 492
 - Fact 8.13.36, 492
- Minkowski's determinant theorem**
 - Corollary 8.4.15, 429
- reverse Fischer inequality**
 - Fact 8.13.41, 493
- positive-semidefinite matrix inequality**
- Araki**
 - Fact 8.12.21, 480
- Araki-Lieb-Thirring inequality**

Fact 8.12.20, 479

**positive-semidefinite
matrix root**
definition, 431

**positive-semidefinite
matrix square root**
definition, 431

**positive-semidefinite
solution**

Riccati equation

Theorem 12.17.2, 782

Theorem 12.18.4, 787

Proposition 12.17.1,
782

Proposition 12.19.1,
789

Corollary 12.17.3, 783

Corollary 12.18.8, 789

Corollary 12.19.2, 790

**positive-semidefinite
square root**
definition, 431

positivity

quadratic form on a
subspace

Fact 8.15.27, 504

Fact 8.15.28, 504

power

adjugate

Fact 4.9.8, 261

cyclic matrix

Fact 5.14.9, 340

derivative

Proposition 10.7.2, 630

discrete-time

asymptotically

stable matrix

Fact 11.21.2, 712

discrete-time

dynamics

Fact 11.21.3, 712

discrete-time

Lyapunov-stable

matrix

Fact 11.21.10, 714

discrete-time

semistable matrix

Fact 11.21.2, 712

group-invertible
matrix

Fact 3.6.2, 177

Fact 6.6.19, 398

idempotent matrix

Fact 3.12.3, 198

identities

Fact 7.6.11, 414

inequality

Fact 1.9.7, 24

Fact 1.10.12, 32

Fact 1.10.31, 36

Fact 1.15.2, 48

Fact 1.15.4, 48

Fact 1.15.5, 49

Fact 1.15.6, 49

Fact 1.15.7, 49

Fact 1.15.8, 49

Fact 1.15.9, 49

Fact 1.15.11, 50

Fact 1.15.22, 53

Kronecker product

Fact 7.4.4, 405

Fact 7.4.10, 405

Fact 7.4.21, 406

Kronecker sum

Fact 7.5.1, 409

lower triangular

matrix

Fact 3.18.7, 216

matrix classes

Fact 3.7.32, 183

matrix exponential

Fact 11.13.19, 680

maximum singular

value

Fact 8.18.26, 520

Fact 9.13.7, 603

Fact 9.13.9, 603

Fact 11.21.17, 715

nonnegative matrix

Fact 4.11.22, 281

normal matrix

Fact 9.13.7, 603

outer-product

matrix

Fact 2.12.7, 126

**positive-definite
matrix**

Fact 8.10.41, 461

Fact 8.10.42, 461

**positive-semidefinite
matrix**

Corollary 8.6.11, 434

Fact 8.9.14, 452

Fact 8.10.36, 461

Fact 8.10.49, 465

Fact 9.9.17, 582

scalar inequality

Fact 1.9.3, 23

Fact 1.9.4, 23

Fact 1.9.5, 23

Fact 1.9.8, 24

Fact 1.9.9, 24

Fact 1.9.10, 24

Fact 1.10.18, 33

Fact 1.11.5, 39

Schur product

Fact 7.6.11, 414

similar matrices

Fact 5.9.1, 311

singular value

inequality

Fact 9.13.19, 605

Fact 9.13.20, 605

skew-Hermitian

matrix

Fact 8.9.14, 452

strictly lower

triangular matrix

Fact 3.18.7, 216

strictly upper

triangular matrix

Fact 3.18.7, 216

symmetric matrix

Fact 3.7.4, 178

trace

Fact 2.12.13, 127

Fact 2.12.17, 127

Fact 4.10.22, 271

Fact 4.11.22, 281

Fact 5.11.9, 322

Fact 5.11.10, 322

Fact 8.12.4, 476

Fact 8.12.5, 476

unitarily invariant
norm

Fact 9.9.17, 582

upper triangular
matrix

Fact 3.18.7, 216

power difference

expansion

Fact 2.12.20, 128

power function

scalar inequalities

Fact 1.10.23, 34

power inequality

Lehmer mean

Fact 1.10.35, 36

monotonicity

Fact 1.10.33, 36

Fact 1.10.34, 36

Fact 1.10.35, 36

positive-definite
matrix

Fact 8.10.52, 466

scalar case

Fact 1.9.11, 24

Fact 1.9.12, 24

Fact 1.10.41, 38

sum inequality

Fact 1.16.28, 66

Fact 1.16.29, 66

two-variable

Fact 1.10.21, 33

Fact 1.10.22, 34

power mean

monotonicity

Fact 1.15.30, 55

power of a

positive-semidefinite
matrix

Bessis-Moussa-
Villani trace

conjecture

Fact 8.12.30, 482

power-sum inequality

Hölder norm

Fact 1.15.34, 57

norm monotonicity

Fact 1.10.30, 35

Fact 1.15.34, 57

Powers

Schatten norm for
positive-

semidefinite

matrices

Fact 9.9.22, 583

powers

Beckner's two-point

inequality

Fact 1.10.15, 33

Fact 9.9.35, 586

inequality

Fact 1.10.8, 31

Fact 1.10.9, 32

Fact 1.10.10, 32

Fact 1.10.14, 33

Fact 1.10.15, 33

Fact 1.10.16, 33

Fact 9.7.20, 569

Fact 9.9.35, 586

optimal 2-uniform

convexity

Fact 1.10.15, 33

Fact 9.9.35, 586

primary circulant

circulant matrix

Fact 5.16.7, 355

irreducible matrix

Fact 3.20.3, 217

prime numbers

Euler product

formula

Fact 1.7.8, 19

factorization

involving π

Fact 1.7.8, 19

primitive matrix

definition

Fact 4.11.5, 273

principal angle

gap topology

Fact 10.9.18, 636

subspace

Fact 2.9.19, 112

Fact 5.11.39, 329

Fact 5.12.17, 335

Fact 10.9.18, 636

**principal angle and
subspaces**

Ljance

Fact 5.11.39, 329

principal branch

logarithm function

Fact 1.18.7, 72

principal square root
definition, 630

integral formula

Fact 10.10.1, 637

square root

Theorem 10.6.1, 629

principal submatrix

definition, 80

problem

absolute value

inequality

Fact 1.11.1, 39

Fact 1.11.12, 43

Fact 1.12.3, 46

adjoint norm

Fact 9.8.8, 572

adjugate of a normal
matrix

Fact 3.7.10, 179

asymptotic stability
of a compartmental

matrix

Fact 11.19.6, 707

bialternate product
and compound

matrix

Fact 7.5.17, 411

Cayley transform of
a Lyapunov-stable

matrix

Fact 11.21.8, 713

commutator

realization

Fact 3.8.2, 184

commuting

projectors

Fact 3.13.20, 209

- convergence of the Baker-Campbell-Hausdorff series
 - Fact 11.14.6, 683
- convergent sequence for the generalized inverse
 - Fact 6.3.35, 376
- cross product of complex vectors
 - Fact 3.10.1, 186
- determinant lower bound
 - Fact 8.13.31, 491
- determinant of partitioned matrix
 - Fact 2.14.13, 135
- determinant of the geometric mean
 - Fact 8.21.19, 534
- dimension of the centralizer
 - Fact 7.5.2, 409
- discrete-time Lyapunov-stable matrix and the matrix exponential
 - Fact 11.21.4, 712
- entries of an orthogonal matrix
 - Fact 3.11.9, 191
- equality in the triangle inequality
 - Fact 9.7.3, 563
- exponential representation of a discrete-time Lyapunov-stable matrix
 - Fact 11.21.7, 713
- factorization of a partitioned matrix
 - Fact 6.5.25, 392
- factorization of a unitary matrix
 - Fact 5.15.16, 347
- factorization of an orthogonal matrix by reflectors
 - Fact 5.15.31, 350
- factorization of nonsingular matrix by elementary matrices
 - Fact 5.15.12, 347
- Frobenius norm lower bound
 - Fact 9.9.11, 581
 - Fact 9.9.15, 582
- generalized inverse least squares solution
 - Fact 9.15.2, 618
- generalized inverse of a partitioned matrix
 - Fact 6.5.24, 391
- geometric mean and generalized inverses
 - Fact 8.10.43, 461
- Hahn-Banach theorem interpretation
 - Fact 10.9.12, 635
- Hölder-induced norm inequality
 - Fact 9.8.21, 575
- Hurwitz stability test
 - Fact 11.18.23, 702
- inequalities involving the trace and Frobenius norm
 - Fact 9.11.3, 597
- inverse image of a subspace
 - Fact 2.9.30, 114
- inverse matrix
 - Fact 2.17.8, 148
- Kronecker product of positive-semidefinite matrices
 - Fact 8.21.22, 536
- least squares and unitary biequivalence
 - Fact 9.15.6, 619
- Lie algebra of upper triangular Lie groups
 - Fact 11.22.1, 715
- lower bounds for the difference of complex numbers
 - Fact 1.18.2, 69
- Lyapunov-stable matrix and the matrix exponential
 - Fact 11.18.37, 705
- majorization and singular values
 - Fact 8.17.5, 509
- matrix exponential
 - Fact 11.14.2, 681
- matrix exponential and proper rotation
 - Fact 11.11.7, 673
 - Fact 11.11.8, 674
 - Fact 11.11.9, 674
- matrix exponential formula
 - Fact 11.14.34, 688
- maximum eigenvalue of the difference of positive-semidefinite matrices
 - Fact 8.18.14, 516
- maximum singular value of an idempotent matrix
 - Fact 5.11.38, 328
- modification of a positive-semidefinite matrix
 - Fact 8.8.13, 448
- orthogonal complement
 - Fact 2.9.15, 112
- orthogonal matrix
 - Fact 3.9.5, 186

polar decomposition
of a matrix
exponential
 Fact 11.13.9, 679

Popoviciu's
inequality and
Hlawka's inequality
 Fact 1.8.6, 22

positive-definite
matrix
 Fact 8.8.9, 448

positive-semidefinite
matrix trace upper
bound
 Fact 8.12.20, 479

power inequality
 Fact 1.9.2, 23
 Fact 1.10.41, 38
 Fact 1.15.7, 49

quadrilateral with an
inscribed circle
 Fact 2.20.13, 159

rank of a positive-
semidefinite
matrix
 Fact 8.8.2, 445

reflector
 Fact 3.14.7, 211

reverse triangle
inequality
 Fact 9.7.6, 565

simisimple imaginary
eigenvalues of a
partitioned matrix
 Fact 5.12.21, 337

singular value of a
partitioned matrix
 Fact 9.14.14, 610

singular values of a
normal matrix
 Fact 9.11.2, 597

special orthogonal
group and matrix
exponentials
 Fact 11.11.13, 675

spectrum of a
partitioned
positive-

semidefinite
matrix
 Fact 5.12.22, 338

spectrum of a sum of
outer products
 Fact 5.11.13, 323

spectrum of the
Laplacian matrix
 Fact 4.11.11, 277

sum of commutators
 Fact 2.18.12, 151

trace of a
positive-definite
matrix
 Fact 8.12.27, 481

upper bounds for the
trace of a product
of matrix
exponentials
 Fact 11.16.4, 692

product

adjugate
 Fact 2.16.10, 143

characteristic
polynomial
 Corollary 4.4.11, 245

compound matrix
 Fact 7.5.17, 411

Drazin generalized
inverse
 Fact 6.6.3, 393
 Fact 6.6.4, 394

generalized inverse
 Fact 6.4.5, 378
 Fact 6.4.6, 378
 Fact 6.4.8, 379
 Fact 6.4.9, 379

Fact 6.4.10, 379
 Fact 6.4.11, 379
 Fact 6.4.12, 379
 Fact 6.4.13, 380
 Fact 6.4.14, 380

Fact 6.4.16, 380
 Fact 6.4.17, 380
 Fact 6.4.21, 381
 Fact 6.4.22, 381
 Fact 6.4.23, 381

Fact 6.4.30, 382

idempotent matrix

Fact 3.12.29, 203

identities
 Fact 2.12.19, 127

induced lower bound
 Proposition 9.5.3, 559

left inverse
 Fact 2.15.5, 141

left-invertible matrix
 Fact 2.10.3, 115

maximum singular
value
 Fact 9.14.2, 607

positive-definite
matrix
 Corollary 8.3.6, 424

positive-semidefinite
matrix
 Corollary 8.3.6, 424

projector
 Fact 3.13.18, 209
 Fact 3.13.20, 209
 Fact 3.13.21, 209

Fact 6.4.16, 380
 Fact 6.4.17, 380

Fact 6.4.21, 381
 Fact 6.4.23, 381
 Fact 8.10.23, 458

quadruple
 Fact 2.16.11, 143

rank
 Fact 3.7.30, 183

right inverse
 Fact 2.15.6, 141

right-invertible
matrix
 Fact 2.10.3, 115

singular value
 Proposition 9.6.1, 560
 Proposition 9.6.2, 561
 Proposition 9.6.3, 561
 Proposition 9.6.4, 561
 Fact 8.18.21, 519
 Fact 9.14.26, 613

singular value
inequality
 Fact 9.13.17, 604
 Fact 9.13.18, 605

skew-symmetric
matrix

- Fact 5.15.37, 351
- trace**
 - Fact 5.12.6, 334
 - Fact 5.12.7, 334
 - Fact 8.12.14, 478
 - Fact 8.12.15, 478
 - Fact 9.14.3, 607
 - Fact 9.14.4, 608
- vec**
 - Fact 7.4.6, 405
- product identity**
 - Lagrange identity**
 - Fact 1.16.8, 61
- product of matrices**
 - definition, 81**
- product of projectors**
 - Crimmins**
 - Fact 6.3.32, 376
- product of sums**
 - inequality**
 - Fact 1.16.10, 62
- projector**
 - commutator**
 - Fact 3.13.23, 210
 - Fact 9.9.9, 581
 - commuting matrices**
 - Fact 6.4.33, 383
 - Fact 8.10.23, 458
 - Fact 8.10.25, 458
 - complementary subspaces**
 - Fact 3.13.24, 210
 - complex conjugate transpose**
 - Fact 3.13.1, 206
 - controllable subspace**
 - Lemma 12.6.6, 738
 - definition**
 - Definition 3.1.1, 165
 - difference**
 - Fact 3.13.24, 210
 - Fact 5.12.17, 335
 - Fact 6.4.33, 383
 - elementary reflector**
 - Fact 5.15.13, 347
 - Euclidean norm**
 - Fact 9.8.2, 571
 - Fact 9.8.3, 571
 - Fact 10.9.17, 636
 - factorization**
 - Fact 5.15.13, 347
 - Fact 5.15.17, 348
 - Fact 6.3.32, 376
 - generalized inverse**
 - Fact 6.3.3, 370
 - Fact 6.3.4, 370
 - Fact 6.3.5, 370
 - Fact 6.3.26, 375
 - Fact 6.3.27, 375
 - Fact 6.3.32, 376
 - Fact 6.4.15, 380
 - Fact 6.4.16, 380
 - Fact 6.4.17, 380
 - Fact 6.4.18, 381
 - Fact 6.4.19, 381
 - Fact 6.4.21, 381
 - Fact 6.4.23, 381
 - Fact 6.4.24, 381
 - Fact 6.4.25, 381
 - Fact 6.4.33, 383
 - Fact 6.4.41, 384
 - Fact 6.4.46, 385
 - Fact 6.5.10, 388
 - greatest lower bound**
 - Fact 6.4.41, 384
 - group-invertible matrix**
 - Fact 3.13.21, 209
 - Hermitian matrix**
 - Fact 3.13.2, 206
 - Fact 3.13.13, 208
 - Fact 3.13.20, 209
 - Fact 5.15.17, 348
 - idempotent matrix**
 - Fact 3.13.3, 206
 - Fact 3.13.13, 208
 - Fact 3.13.20, 209
 - Fact 3.13.24, 210
 - Fact 5.10.13, 319
 - Fact 5.12.18, 336
 - Fact 6.3.26, 375
 - Fact 6.4.18, 381
 - Fact 6.4.19, 381
 - Fact 6.4.20, 381
 - Fact 6.4.25, 381
 - identity**
 - Fact 3.13.9, 207
 - inequality**
 - Fact 8.9.23, 452
 - intersection of ranges**
 - Fact 6.4.41, 384
 - Kronecker product**
 - Fact 7.4.16, 406
 - least upper bound**
 - Fact 6.4.41, 385
 - matrix difference**
 - Fact 3.13.24, 210
 - Fact 6.4.20, 381
 - matrix limit**
 - Fact 6.4.41, 384
 - Fact 6.4.46, 385
 - matrix product**
 - Fact 3.13.18, 209
 - Fact 3.13.20, 209
 - Fact 3.13.21, 209
 - Fact 6.4.16, 380
 - Fact 6.4.17, 380
 - Fact 6.4.21, 381
 - Fact 6.4.23, 381
 - matrix sum**
 - Fact 5.19.4, 360
 - maximum singular value**
 - Fact 5.11.38, 328
 - Fact 5.12.17, 335
 - Fact 5.12.18, 336
 - Fact 9.14.1, 607
 - Fact 9.14.30, 615
 - normal matrix**
 - Fact 3.13.3, 206
 - Fact 3.13.20, 209
 - onto a subspace**
 - definition, 175**
 - orthogonal complement**
 - Proposition 3.5.2, 175
 - partitioned matrix**
 - Fact 3.13.12, 208
 - Fact 3.13.22, 210
 - Fact 3.13.23, 210
 - Fact 6.5.13, 388
 - positive-semidefinite matrix**
 - Fact 3.13.4, 207

product

- Fact 3.13.24, 210
- Fact 5.12.16, 335
- Fact 6.4.19, 381
- Fact 8.10.23, 458

quadratic form

- Fact 3.13.10, 208
- Fact 3.13.11, 208

range

- Proposition 3.5.1, 175
- Fact 3.13.5, 207
- Fact 3.13.14, 208
- Fact 3.13.15, 208
- Fact 3.13.17, 208
- Fact 3.13.18, 209
- Fact 3.13.19, 209
- Fact 3.13.20, 209
- Fact 6.4.41, 384
- Fact 6.4.45, 385
- Fact 6.4.46, 385

range-Hermitian matrix

- Fact 3.13.3, 206
- Fact 3.13.20, 209

rank

- Fact 3.13.9, 207
- Fact 3.13.12, 208
- Fact 3.13.22, 210
- Fact 3.13.23, 210
- Fact 5.12.17, 335

reflector

- Fact 3.13.16, 208
- Fact 3.14.1, 211

right inverse

- Fact 3.13.6, 207

similar matrices

- Corollary 5.5.22, 301
- Fact 5.10.13, 319

simultaneous

- triangularization**
- Fact 5.17.6, 358

skew-Hermitian matrix

- Fact 9.9.9, 581

spectrum

- Proposition 5.5.21, 300
- Fact 5.12.15, 335
- Fact 5.12.16, 335

square root

- Fact 8.10.25, 458

subspace

- Proposition 3.5.2, 175
- Fact 9.8.3, 571
- Fact 10.9.17, 636

sum

- Fact 3.13.23, 210
- Fact 5.12.17, 335

trace

- Fact 5.8.11, 309

tripotent matrix

- Fact 6.4.33, 383

union of ranges

- Fact 6.4.41, 385

unitarily similar matrices

- Fact 5.10.12, 319

unobservable

- subspace**
- Lemma 12.3.6, 729

projector onto a subspace

- definition, 175**

proper rational function

- definition**
- Definition 4.7.1, 249

proper rational transfer function

- definition**
- Definition 4.7.2, 249

realization

- Theorem 12.9.4, 751

proper rotation

- matrix exponential**
- Fact 11.11.7, 673
- Fact 11.11.8, 674
- Fact 11.11.9, 674

proper separation theorem

- convex sets**
- Fact 10.9.14, 635

proper subset

- definition, 3**

proposition

- definition, 1**

Ptak

- maximum singular value**
- Fact 9.13.9, 603

Ptolemy's inequality

- quadrilateral**
- Fact 2.20.13, 159

Ptolemy's theorem

- quadrilateral**
- Fact 2.20.13, 159

Purves

- similar matrices and nonzero diagonal entries**
- Fact 5.9.14, 313

Putnam-Fuglede theorem

- normal matrix**
- Fact 5.14.30, 343

Pythagorean theorem

- norm identity**
- Fact 9.7.4, 563
- vector identity**
- Fact 9.7.4, 563

Pythagorean triples

- quadratic identity**
- Fact 1.10.1, 30

Q**QR decomposition**

- existence**
- Fact 5.15.8, 346

quadratic identity

- Fact 1.11.2, 39

inequality

- Fact 1.10.4, 31
- Fact 1.10.5, 31
- Fact 1.10.6, 31
- Fact 1.10.7, 31
- Fact 1.11.3, 39
- Fact 1.11.4, 39

quadratic form

- cone**

- Fact 8.14.11, 498
- Fact 8.14.13, 498
- Fact 8.14.14, 498
- convex cone**
 - Fact 8.14.11, 498
 - Fact 8.14.13, 498
 - Fact 8.14.14, 498
- convex set**
 - Fact 8.14.2, 494
 - Fact 8.14.3, 495
 - Fact 8.14.4, 495
 - Fact 8.14.5, 495
 - Fact 8.14.6, 496
 - Fact 8.14.9, 497
 - Fact 8.14.11, 498
 - Fact 8.14.12, 498
 - Fact 8.14.13, 498
 - Fact 8.14.14, 498
- copositive matrix**
 - Fact 8.15.37, 507
- definition, 166**
- dual norm**
 - Fact 9.8.34, 577
- eigenvalue**
 - Lemma 8.4.3, 425
 - Fact 8.15.20, 503
- field**
 - Fact 3.7.7, 179
- Hermitian matrix**
 - Fact 3.7.6, 178
 - Fact 3.7.7, 179
 - Fact 8.15.24, 504
 - Fact 8.15.25, 504
 - Fact 8.15.26, 504
 - Fact 8.15.31, 505
- hidden convexity**
 - Fact 8.14.11, 498
- Hölder-induced norm**
 - Fact 9.8.35, 577
 - Fact 9.8.36, 577
- idempotent matrix**
 - Fact 3.13.11, 208
- induced norm**
 - Fact 9.8.34, 577
- inequality**
 - Fact 8.15.7, 501
 - Fact 8.15.8, 501
 - Fact 8.15.13, 502
- Fact 8.15.14, 502
- Fact 8.15.15, 502
- Fact 8.15.16, 502
- Fact 8.15.18, 503
- Fact 8.15.19, 503
- Fact 8.15.21, 503
- Fact 8.15.22, 503
- integral**
 - Fact 8.15.34, 506
 - Fact 8.15.35, 506
- Kantorovich inequality**
 - Fact 8.15.9, 501
- Laplacian matrix**
 - Fact 8.15.36, 506
- linear constraint**
 - Fact 8.14.10, 497
- matrix logarithm**
 - Fact 8.15.15, 502
- maximum eigenvalue**
 - Lemma 8.4.3, 425
- maximum singular value**
 - Fact 9.13.1, 602
 - Fact 9.13.2, 602
- minimum eigenvalue**
 - Lemma 8.4.3, 425
- minimum singular value**
 - Fact 9.13.1, 602
- norm**
 - Fact 9.7.30, 571
- null space**
 - Fact 8.15.1, 500
 - Fact 8.15.23, 504
- one-sided cone**
 - Fact 8.14.14, 498
- outer-product matrix**
 - Fact 9.13.3, 602
- partitioned matrix**
 - Fact 8.15.5, 500
 - Fact 8.15.6, 501
- positive-definite matrix**
 - Fact 8.15.24, 504
 - Fact 8.15.25, 504
 - Fact 8.15.26, 504
 - Fact 8.15.29, 505
- Fact 8.15.30, 505
- Fact 9.8.37, 577
- positive-definite matrix inequality**
 - Fact 8.15.4, 500
- positive-semidefinite matrix**
 - Fact 8.14.2, 494
 - Fact 8.14.3, 495
 - Fact 8.14.4, 495
 - Fact 8.14.5, 495
 - Fact 8.14.6, 496
 - Fact 8.15.1, 500
 - Fact 8.15.9, 501
 - Fact 8.15.10, 501
 - Fact 8.15.11, 501
 - Fact 8.15.13, 502
 - Fact 8.15.14, 502
 - Fact 8.15.15, 502
 - Fact 8.15.16, 502
 - Fact 8.15.17, 503
 - Fact 8.15.18, 503
- positive-semidefinite matrix inequality**
 - Fact 8.15.4, 500
 - Fact 8.15.7, 501
- projector**
 - Fact 3.13.10, 208
 - Fact 3.13.11, 208
- quadratic minimization lemma**
 - Fact 8.14.15, 499
- Rayleigh quotient**
 - Lemma 8.4.3, 425
- Reid's inequality**
 - Fact 8.15.18, 503
- skew-Hermitian matrix**
 - Fact 3.7.6, 178
- skew-symmetric matrix**
 - Fact 3.7.5, 178
- spectrum**
 - Fact 8.14.7, 496
 - Fact 8.14.8, 497
- subspace**
 - Fact 8.15.27, 504
 - Fact 8.15.28, 504

symmetric matrix
Fact 3.7.5, 178
vector derivative
Proposition 10.7.1, 630

quadratic form inequality

Marcus
Fact 8.15.19, 503

quadratic form on a subspace

positivity
Fact 8.15.27, 504
Fact 8.15.28, 504

quadratic formula
complex numbers
Fact 1.18.3, 70

quadratic inequality
Aczel's inequality
Fact 1.16.19, 64

sum
Fact 1.10.17, 33
sum of squares
Fact 1.12.4, 47
Fact 1.14.1, 47

quadratic matrix equation
spectrum

Fact 5.11.3, 321
Fact 5.11.4, 321

quadratic minimization lemma

quadratic form
Fact 8.14.15, 499

quadratic performance measure

definition, 775
H₂ norm
Proposition 12.15.1, 776

quadrilateral

Brahmagupta's formula
Fact 2.20.13, 159
Ptolemy's inequality
Fact 2.20.13, 159

Ptolemy's theorem
Fact 2.20.13, 159
semiperimeter
Fact 2.20.13, 159

quadrilateral inequality

Euclidean norm
Fact 9.7.4, 563

quadruple product

trace
Fact 7.4.9, 405
vec
Fact 7.4.9, 405

quantum information

matrix logarithm
Fact 11.14.27, 686

quartic

arithmetic-mean–geometric-mean inequality
Fact 1.12.5, 47
identity
Fact 1.10.3, 30
inequality
Fact 1.17.1, 67

quaternion group

symplectic group
Fact 3.22.4, 227

quaternions

2 × 2 matrix representation
Fact 3.22.6, 227
4 × 4 matrix representation
Fact 3.22.3, 227
angular velocity vector
Fact 11.11.15, 675
complex decomposition
Fact 3.22.2, 227
complex matrix representation
Fact 3.22.7, 229
inequality
Fact 1.14.1, 47
matrix exponential

Fact 11.11.15, 675
orthogonal matrix
Fact 3.11.10, 192
Pauli spin matrices
Fact 3.22.6, 227
real matrix representation
Fact 3.22.1, 225
Fact 3.22.8, 229
Rodrigues's formulas
Fact 3.11.11, 193
unitary matrix
Fact 3.22.9, 229

quintic

inequality
Fact 1.10.11, 32

quintic polynomial

Abel
Fact 3.21.7, 223
Galois
Fact 3.21.7, 223

quotient

definition, 233

R

Rado

arithmetic-mean–geometric-mean inequality
Fact 1.15.29, 55
convex hull interpretation of strong majorization
Fact 2.21.7, 163

Radstrom

set cancellation
Fact 10.9.7, 635

Rahman

polynomial root bound
Fact 11.20.7, 710

Ramanujan

cube identity
Fact 2.12.24, 128

Ramus

fundamental triangle inequality
Fact 2.20.11, 156

range

adjugate
Fact 2.16.7, 143

complex conjugate transpose
Fact 6.5.3, 386
Fact 8.7.2, 443

controllability
Fact 12.20.7, 791

Drazin generalized inverse
Proposition 6.2.2, 368

factorization
Theorem 8.6.2, 431

generalized inverse
Proposition 6.1.6, 365
Fact 6.3.24, 374
Fact 6.4.42, 384
Fact 6.4.43, 385
Fact 6.5.3, 386

group generalized inverse
Proposition 6.2.3, 369

group-invertible matrix
Fact 3.6.1, 177
Fact 5.14.4, 339

Hermitian matrix
Lemma 8.6.1, 431

idempotent matrix
Fact 3.12.3, 199
Fact 3.12.4, 199
Fact 3.15.4, 200
Fact 6.3.24, 374

identity
Fact 2.10.8, 116
Fact 2.10.12, 116
Fact 2.10.20, 117

inclusion
Fact 2.10.7, 116
Fact 2.10.8, 116

inclusion for a matrix power
Corollary 2.4.2, 94

inclusion for a matrix product
Lemma 2.4.1, 94
Fact 2.10.2, 115

index of a matrix
Fact 5.14.4, 339

involutory matrix
Fact 3.15.4, 212

Kronecker product
Fact 7.4.22, 407

minimal polynomial
Corollary 11.8.6, 664

nilpotent matrix
Fact 3.17.1, 213
Fact 3.17.2, 213
Fact 3.17.3, 213

null space
Corollary 2.5.6, 97
Fact 2.10.1, 115

null space inclusions
Theorem 2.4.3, 94

outer-product matrix
Fact 2.10.11, 116

partitioned matrix
Fact 2.11.1, 120
Fact 2.11.2, 121
Fact 6.5.3, 386

positive-semidefinite matrix
Theorem 8.6.2, 431
Corollary 8.2.2, 419
Fact 8.7.1, 443
Fact 8.7.2, 443
Fact 8.7.3, 443
Fact 8.7.4, 443
Fact 8.7.5, 443
Fact 8.10.2, 456
Fact 8.20.7, 526
Fact 8.20.8, 526
Fact 8.20.10, 526
Fact 8.20.11, 527

projector
Proposition 3.5.1, 175
Fact 3.13.14, 208
Fact 3.13.15, 208
Fact 3.13.17, 208
Fact 3.13.18, 209
Fact 3.13.19, 209

Fact 3.13.20, 209
Fact 6.4.41, 384
Fact 6.4.45, 385
Fact 6.4.46, 385

rank
Fact 2.11.5, 121

right-equivalent matrices
Proposition 5.1.3, 283

Schur product
Fact 7.6.5, 413

skew-Hermitian matrix
Fact 8.7.3, 443

square root
Fact 8.7.2, 443

stabilizability
Fact 12.20.7, 791

subspace
Fact 2.9.24, 113

symmetric matrix
Fact 3.7.4, 178

range of a function
definition, 3

range of a matrix
definition, 93

range-Hermitian matrix
commuting matrices
Fact 6.4.26, 382
Fact 6.4.27, 382

complex conjugate transpose
Fact 3.6.4, 178
Fact 6.3.10, 371
Fact 6.6.16, 396

congruent matrices
Proposition 3.4.5, 174
Fact 5.9.6, 312

definition
Definition 3.1.1, 165

dissipative matrix
Fact 5.14.31, 343

generalized inverse
Proposition 6.1.6, 365
Fact 6.3.10, 371
Fact 6.3.11, 372
Fact 6.3.12, 372
Fact 6.3.16, 373

- Fact 6.3.17, 373
- Fact 6.4.26, 382
- Fact 6.4.27, 382
- Fact 6.4.28, 382
- Fact 6.4.29, 382
- generalized projector**
 - Fact 3.6.4, 178
- group generalized**
 - inverse**
 - Fact 6.6.8, 394
- group-invertible matrix**
 - Proposition 3.1.6, 169
 - Fact 6.6.16, 396
- idempotent matrix**
 - Fact 3.13.3, 206
 - Fact 6.3.27, 375
- Kronecker product**
 - Fact 7.4.16, 406
- Kronecker sum**
 - Fact 7.5.8, 409
- nonsingular matrix**
 - Proposition 3.1.6, 169
- normal matrix**
 - Proposition 3.1.6, 169
- null space**
 - Fact 3.6.3, 177
- positive-semidefinite matrix**
 - Fact 8.20.21, 530
- product**
 - Fact 6.4.29, 382
- projector**
 - Fact 3.13.3, 206
 - Fact 3.13.20, 209
- rank**
 - Fact 3.6.3, 177
 - Fact 3.6.5, 178
- right-equivalent matrices**
 - Fact 3.6.3, 177
- Schur decomposition**
 - Corollary 5.4.4, 293
- unitarily similar matrices**
 - Proposition 3.4.5, 174
 - Corollary 5.4.4, 293
- rank**
 - additivity**
 - Fact 2.11.4, 121
 - Fact 6.4.32, 382
 - adjugate**
 - Fact 2.16.7, 143
 - Fact 2.16.8, 143
 - biequivalent matrices**
 - Proposition 5.1.3, 283
 - commutator**
 - Fact 3.12.31, 204
 - Fact 3.13.23, 210
 - Fact 5.17.5, 358
 - Fact 6.3.9, 371
 - complex conjugate transpose**
 - Fact 2.10.21, 117
 - complex matrix**
 - Fact 2.19.3, 151
 - controllability matrix**
 - Corollary 12.6.3, 737
 - controllable pair**
 - Fact 5.14.10, 340
 - controllably**
 - asymptotically stable**
 - Proposition 12.7.4, 745
 - Proposition 12.7.5, 746
 - cyclic matrix**
 - Fact 5.11.1, 321
 - definition, 95**
 - diagonal dominance**
 - Fact 4.10.23, 271
 - difference**
 - Fact 2.10.31, 119
 - dimension inequality**
 - Fact 2.10.4, 115
 - factorization**
 - Fact 5.15.40, 351
 - Frobenius norm**
 - Fact 9.11.4, 598
 - Fact 9.14.28, 614
 - Fact 9.15.4, 618
 - generalized inverse**
 - Fact 6.3.9, 371
 - Fact 6.3.22, 374
 - Fact 6.3.36, 377
 - Fact 6.4.2, 377
 - Fact 6.4.32, 382
 - Fact 6.4.44, 385
 - Fact 6.5.6, 386**
 - Fact 6.5.8, 387**
 - Fact 6.5.9, 387**
 - Fact 6.5.12, 388**
 - Fact 6.5.13, 388**
 - Fact 6.5.14, 388**
 - geometric multiplicity**
 - Proposition 4.5.2, 246
 - group-invertible matrix**
 - Fact 3.6.1, 177
 - Fact 5.8.5, 308
 - Fact 5.14.4, 339
 - Hermitian matrix**
 - Fact 3.7.22, 182
 - Fact 3.7.30, 183
 - Fact 5.8.6, 308
 - Fact 5.8.7, 308
 - Fact 8.9.7, 451
 - idempotent matrix**
 - Fact 3.12.6, 199
 - Fact 3.12.9, 199
 - Fact 3.12.19, 201
 - Fact 3.12.20, 201
 - Fact 3.12.22, 201
 - Fact 3.12.24, 202
 - Fact 3.12.25, 202
 - Fact 3.12.27, 203
 - Fact 3.12.31, 204
 - Fact 5.8.1, 307
 - Fact 5.11.7, 322
 - identities with defect**
 - Corollary 2.5.1, 96
 - identities with transpose**
 - Corollary 2.5.3, 96
 - identity**
 - Fact 2.10.12, 116
 - Fact 2.10.13, 116
 - Fact 2.10.17, 117
 - Fact 2.10.20, 117
 - Fact 2.10.23, 118
 - identity with defect**
 - Corollary 2.5.5, 97
 - identity with powers**
 - Proposition 2.5.8, 97
 - inequality**
 - Fact 2.10.22, 118

- inertia**
 - Fact 5.8.5, 308
 - Fact 5.8.18, 310
- inverse**
 - Fact 2.11.21, 125
 - Fact 2.11.22, 125
- inverse matrix**
 - Fact 2.17.10, 149
 - Fact 6.5.11, 388
- Kronecker product**
 - Fact 7.4.23, 407
 - Fact 7.4.24, 407
 - Fact 7.4.25, 407
 - Fact 8.21.16, 534
- Kronecker sum**
 - Fact 7.5.2, 409
 - Fact 7.5.9, 409
 - Fact 7.5.10, 410
- linear matrix equation**
 - Fact 2.10.16, 117
- linear system solution**
 - Theorem 2.6.4, 100
 - Corollary 2.6.7, 101
- lower bound for product**
 - Proposition 2.5.9, 97
 - Corollary 2.5.10, 98
- M-matrix**
 - Fact 8.7.7, 444
- matrix difference**
 - Fact 2.10.27, 118
 - Fact 2.10.30, 119
- matrix power**
 - Fact 2.10.22, 118
- matrix powers**
 - Corollary 2.5.7, 97
 - Fact 3.17.5, 213
- matrix sum**
 - Fact 2.10.27, 118
 - Fact 2.10.28, 118
 - Fact 2.10.29, 119
 - Fact 2.11.4, 121
- nilpotent matrix**
 - Fact 3.17.4, 213
 - Fact 3.17.5, 213
- nonsingular submatrices**
 - Proposition 2.7.7, 106
- observability matrix**
 - Corollary 12.3.3, 729
- observably asymptotically stable**
 - Proposition 12.4.4, 733
- ones matrix**
 - Fact 2.10.18, 117
- outer-product matrix**
 - Fact 2.10.19, 117
 - Fact 2.10.24, 118
 - Fact 3.12.6, 199
- outer-product perturbation**
 - Fact 2.10.25, 118
 - Fact 6.4.2, 377
- partitioned matrix**
 - Corollary 2.8.5, 107
 - Fact 2.11.7, 121
 - Fact 2.11.8, 122
 - Fact 2.11.9, 122
 - Fact 2.11.10, 122
 - Fact 2.11.11, 123
 - Fact 2.11.12, 123
 - Fact 2.11.13, 123
 - Fact 2.11.14, 123
 - Fact 2.11.15, 124
 - Fact 2.11.16, 124
 - Fact 2.11.18, 124
 - Fact 2.11.19, 125
 - Fact 2.14.4, 133
 - Fact 2.14.5, 134
 - Fact 2.14.11, 135
 - Fact 2.17.5, 147
 - Fact 2.17.10, 149
 - Fact 3.12.20, 201
 - Fact 3.13.12, 208
 - Fact 3.13.22, 210
 - Fact 5.12.21, 337
 - Fact 6.3.30, 376
 - Fact 6.5.6, 386
 - Fact 6.5.7, 387
 - Fact 6.5.8, 387
 - Fact 6.5.9, 387
 - Fact 6.5.10, 388
 - Fact 6.5.12, 388
 - Fact 6.5.13, 388
- Fact 6.5.14, 388**
- Fact 6.5.15, 389**
- Fact 6.6.2, 393**
- Fact 8.7.6, 443**
- Fact 8.7.7, 444**
- partitioned positive-semidefinite matrix**
 - Proposition 8.2.3, 420
- positive-semidefinite matrix**
 - Fact 5.8.9, 308
 - Fact 8.7.1, 443
 - Fact 8.7.5, 443
 - Fact 8.7.6, 443
 - Fact 8.7.7, 444
 - Fact 8.10.2, 456
 - Fact 8.10.14, 457
 - Fact 8.20.11, 527
 - Fact 8.21.16, 534
- product**
 - Proposition 2.6.3, 99
 - Fact 3.7.30, 183
- product of matrices**
 - Fact 2.10.14, 116
 - Fact 2.10.26, 118
- projector**
 - Fact 3.13.9, 207
 - Fact 3.13.12, 208
 - Fact 3.13.22, 210
 - Fact 3.13.23, 210
 - Fact 5.12.17, 335
- range**
 - Fact 2.11.5, 121
- range-Hermitian matrix**
 - Fact 3.6.3, 177
 - Fact 3.6.5, 178
- rational transfer function**
 - Definition 4.7.4, 249
- Riccati equation**
 - Proposition 12.19.4, 790
- Rosenbrock system matrix**
 - Proposition 12.10.3, 759

Proposition 12.10.11, 764

Schur complement

Proposition 8.2.3, 420

Fact 6.5.6, 386

Fact 6.5.8, 387

Fact 6.5.11, 388

Schur product

Fact 7.6.6, 413

Fact 8.21.16, 534

simple matrix

Fact 5.11.1, 321

singular value

Proposition 5.6.2, 302

Fact 9.14.28, 614

Fact 9.15.4, 618

skew-Hermitian matrix

Fact 3.7.17, 181

Fact 3.7.30, 183

Smith form

Proposition 4.3.5, 237

Proposition 4.3.6, 238

Smith-McMillan form

Proposition 4.7.7, 250

Proposition 4.7.8, 250

subapce dimension theorem

Fact 2.11.10, 122

submatrix

Proposition 4.3.5, 237

Proposition 4.7.7, 250

Fact 2.11.6, 121

Fact 2.11.17, 124

Fact 2.11.20, 125

Fact 2.11.21, 125

Fact 2.11.22, 125

Fact 3.20.5, 218

subspace dimension theorem

Fact 2.11.9, 122

subtractivity

Fact 2.10.30, 119

Fact 2.10.31, 119

Sylvester's equation

Fact 12.21.13, 796

totally positive matrix

Fact 8.7.7, 444

trace

Fact 5.11.10, 322

Fact 9.11.4, 598

tripotent matrix

Fact 2.10.23, 118

Fact 3.16.3, 213

Fact 3.16.4, 213

unitarily invariant norm

Fact 9.14.28, 614

upper bound for product

Corollary 2.5.10, 98

upper bound on rank of a product

Lemma 2.5.2, 96

upper bound with dimensions

Corollary 2.5.4, 97

rank of a polynomial matrix

definition

Definition 4.2.4, 235

Definition 4.3.3, 237

submatrix

Proposition 4.2.7, 236

rank of a rational function

linearly independent columns

Proposition 4.7.6, 250

Proposition 4.7.9, 251

rank subtractivity equivalent conditions

Fact 2.10.30, 119

transitivity

Fact 2.10.31, 119

rank subtractivity partial ordering commuting matrices

Fact 8.19.5, 523

definition

Fact 2.10.32, 119

generalized inverse

Fact 6.5.30, 393

positive-semidefinite matrix

Fact 8.19.5, 523

Fact 8.19.9, 524

Fact 8.20.7, 526

Fact 8.20.8, 526

rank-deficient matrix determinant

Fact 2.13.3, 129

rank-two matrix matrix exponential

Fact 11.11.18, 676

ratio of powers scalar inequalities

Fact 1.10.39, 38

rational canonical form, see multicompanion form or elementary multicompanion form

rational function complex conjugate

Fact 4.8.17, 260

definition

Definition 4.7.1, 249

Hankel matrix

Fact 4.8.8, 257

imaginary part

Fact 4.8.17, 260

spectrum

Fact 5.11.15, 324

rational transfer function

blocking zero

Definition 4.7.4, 249

definition

Definition 4.7.2, 249

Markov

block-Hankel matrix

Proposition 12.9.11,

754

Proposition 12.9.12,

755

Proposition 12.9.13,

755

- Markov parameter**
 - Proposition 12.9.7, 751
- minimal realization**
 - Fact 12.22.12, 800
- normal rank**
 - Definition 4.7.4, 249
- poles**
 - Definition 4.7.4, 249
- rank**
 - Definition 4.7.4, 249
- realization**
 - Fact 12.22.11, 800
- Rayleigh quotient**
 - Hermitian matrix
 - Lemma 8.4.3, 425
 - quadratic form
 - Lemma 8.4.3, 425
- real eigenvalues**
 - positive-semidefinite matrix
 - Fact 5.14.13, 340
- real hypercompanion form**
 - definition, 289
- real Jordan form**
 - existence
 - Theorem 5.3.5, 290
 - hypercompanion matrix
 - Fact 5.10.1, 316
 - Jordan form
 - Fact 5.10.2, 317
 - similarity transformation
 - Fact 5.10.1, 316
 - Fact 5.10.2, 317
- real Jordan matrix**
 - definition, 289
- real normal form**
 - existence
 - Corollary 5.4.9, 295
- real part**
 - frequency response
 - Fact 12.22.5, 799
 - transfer function
 - Fact 12.22.5, 799
- real Schur decomposition**
 - definition, 293
 - existence
 - Corollary 5.4.2, 293
 - Corollary 5.4.3, 293
- real symplectic group**
 - special orthogonal group
 - Fact 3.22.5, 227
- real vector**
 - definition, 85
- realization**
 - controllable canonical form
 - Proposition 12.9.3, 750
 - definition
 - Definition 12.9.2, 749
 - derivative
 - Fact 12.22.6, 799
 - feedback
 - interconnection
 - Proposition 12.13.4, 772
 - Proposition 12.14.1, 774
 - Fact 12.22.8, 799
 - observable canonical form
 - Proposition 12.9.3, 750
 - partitioned transfer function
 - Proposition 12.13.3, 771
 - Fact 12.22.7, 799
 - proper rational transfer function
 - Theorem 12.9.4, 751
 - rational transfer function
 - Fact 12.22.11, 800
 - similar matrices
 - Proposition 12.9.5, 751
 - transfer function
 - Proposition 12.13.1, 770
 - Fact 12.22.3, 798
 - Fact 12.22.4, 799
- rearrangement inequality**
 - Chebyshev's inequality
 - Fact 1.16.3, 60
 - product of sums
 - Fact 1.16.4, 60
 - reverse inequality
 - Fact 1.16.6, 61
 - sum of differences
 - Fact 1.16.4, 60
 - sum of products
 - Fact 1.16.4, 60
 - sum of products inequality
 - Fact 1.16.5, 60
- reciprocal**
 - scalar inequality
 - Fact 1.11.13, 43
 - Fact 1.11.18, 44
 - Fact 1.11.19, 44
 - Fact 1.11.20, 44
- reciprocal argument**
 - transfer function
 - Fact 12.22.4, 799
- reciprocal powers inequality**
 - Fact 1.16.26, 66
 - Fact 1.16.27, 66
- reciprocals**
 - scalar inequality
 - Fact 1.11.23, 45
 - Fact 1.11.26, 45
 - Walker's inequality
 - Fact 1.11.22, 45
- reducible matrix**
 - absolute value
 - Fact 3.20.4, 218
 - definition
 - Definition 3.1.1, 165
 - zero entry
 - Fact 3.20.1, 217
 - Fact 3.20.2, 217

redundant
assumptions
 definition, 2

reflection theorem
 elementary reflector
 Fact 3.14.4, 211

reflector
 definition
 Definition 3.1.1, 165
 elementary reflector
 Fact 5.15.14, 347
factorization
 Fact 5.15.14, 347
Hermitian matrix
 Fact 3.14.2, 211

identity
 Fact 3.14.8, 212

involutory matrix
 Fact 3.14.2, 211

Kronecker product
 Fact 7.4.16, 406

normal matrix
 Fact 5.9.9, 312
 Fact 5.9.10, 312

orthogonal matrix
 Fact 3.11.9, 191
 Fact 5.15.31, 350
 Fact 5.15.35, 351

projector
 Fact 3.13.16, 208
 Fact 3.14.1, 211

rotation matrix
 Fact 3.11.9, 191

similar matrices
 Corollary 5.5.22, 301

skew reflector
 Fact 3.14.7, 211

spectrum
 Proposition 5.5.21, 300

trace
 Fact 5.8.11, 309

tripotent matrix
 Proposition 3.1.6, 169

unitary matrix
 Fact 3.14.2, 211

reflexive hull
 definition
 Definition 1.3.4, 5

relation
 Proposition 1.3.5, 6

reflexive relation
 definition

Definition 1.3.2, 5

graph
 Definition 1.4.1, 8

intersection
 Proposition 1.3.3, 5

pointed cone
 induced by
 Proposition 2.3.6, 93

positive-semidefinite matrix
 Proposition 8.1.1, 417

regular pencil
 definition, 304

generalized eigenvalue
 Proposition 5.7.3, 305
 Proposition 5.7.4, 306

invariant zero
 Corollary 12.10.4, 759
 Corollary 12.10.5, 760
 Corollary 12.10.6, 761

Kronecker canonical form
 Proposition 5.7.2, 305

Moler
 Fact 5.17.3, 358

simultaneous triangularization
 Fact 5.17.2, 358

Stewart
 Fact 5.17.3, 358

upper Hessenberg
 Fact 5.17.3, 358

upper triangular
 Fact 5.17.3, 358

regular polynomial matrix
 definition, 234
 nonsingular
 polynomial matrix
 Proposition 4.2.5, 235

regularized Tikhonov inverse

positive-definite matrix
 Fact 8.9.40, 455

Reid's inequality
 quadratic form
 Fact 8.15.18, 503

relation
 definition, 5
 function
 Proposition 1.3.1, 5

relative complement
 definition, 2

relative degree
 definition
 Definition 4.7.1, 249
 Definition 4.7.3, 249

relative entropy
 matrix logarithm
 Fact 11.14.25, 686

relative gain array
 definition
 Fact 8.21.4, 531

relatively closed set
 complement
 Fact 10.8.5, 632

relatively open set
 complement
 Fact 10.8.5, 632

remainder
 definition, 233

representation
 groups
 Fact 3.21.8, 224

reproducing kernel space
 positive-semidefinite matrix
 Fact 8.8.2, 445

resolvent
 definition, 243
 Laplace transform
 Proposition 11.2.2, 647
 matrix exponential

Proposition 11.2.2, 647

resultant

coprime polynomials
Fact 4.8.4, 254

reversal of a graph

Definition 1.4.1, 8

reversal of a relation

definition
Definition 1.3.4, 5

reverse

arithmetic-mean–
geometric-mean
inequality

Specht
Fact 1.15.19, 52

Specht's ratio
Fact 1.15.19, 52

reverse complex

conjugate transpose
definition, 88

reverse identity matrix

definition, 84
determinant
Fact 2.13.1, 128

spectrum
Fact 5.9.24, 314

symplectic matrix
Fact 3.19.3, 216

reverse inequality

arithmetic-mean–
geometric-mean
inequality
Fact 1.15.18, 52
Fact 1.15.19, 52

Euclidean norm
triangle inequality
Fact 9.7.6, 565

Fischer's inequality
Fact 8.13.41, 493

Hölder norm triangle
inequality
Fact 9.7.19, 569

Young inequality
Fact 1.10.22, 34

reverse transpose

definition, 88
similar matrices
Fact 5.9.11, 313

reverse-diagonal entries

definition, 80

reverse-diagonal matrix

definition
Definition 3.1.3, 167
semisimple matrix
Fact 5.14.12, 340

reverse-Hermitian matrix

definition
Definition 3.1.1, 165

reverse-symmetric matrix

definition
Definition 3.1.1, 165

factorization
Fact 5.9.12, 313

similar matrices
Fact 5.9.11, 313

Toeplitz matrix
Fact 3.18.5, 215

reversed relation

relation
Proposition 1.3.5, 6

Riccati differential equation

matrix differential
equation
Fact 12.23.5, 803

Riccati equation

closed-loop spectrum
Proposition 12.16.14,
781
Proposition 12.18.2,
787
Proposition 12.18.3,
787
Proposition 12.18.7,
789

detectability

Corollary 12.17.3, 783
Corollary 12.19.2, 790

existence

Fact 12.23.3, 802

geometric mean

Fact 12.23.4, 802

golden mean

Fact 12.23.4, 802

golden ratio

Fact 12.23.4, 802

Hamiltonian

Theorem 12.17.9, 784
Proposition 12.16.14,
781
Corollary 12.16.15, 781

inertia

Lemma 12.16.18, 781

linear-quadratic control problem

Theorem 12.15.2, 776

maximal solution

Definition 12.16.12,
780
Theorem 12.18.1, 787
Theorem 12.18.4, 787
Proposition 12.18.2,
787
Proposition 12.18.7,
789

monotonicity

Proposition 12.18.5,
788
Corollary 12.18.6, 788

observability

Lemma 12.16.18, 781

positive-definite matrix

Fact 12.23.4, 802

positive-definite solution

Theorem 12.17.2, 782
Proposition 12.19.3,
790

Corollary 12.19.2, 790

positive-semidefinite solution

Theorem 12.17.2, 782
Theorem 12.18.4, 787
Proposition 12.17.1,
782

- Proposition 12.19.1, 789
 - Corollary 12.17.3, 783
 - Corollary 12.18.8, 789
 - Corollary 12.19.2, 790
 - rank**
 - Proposition 12.19.4, 790
 - solution**
 - Definition 12.16.12, 780
 - Fact 12.23.2, 802
 - stabilizability**
 - Theorem 12.17.9, 784
 - Theorem 12.18.1, 787
 - Corollary 12.19.2, 790
 - stabilizing solution**
 - Definition 12.16.12, 780
 - Theorem 12.17.2, 782
 - Theorem 12.17.9, 784
 - Theorem 12.18.4, 787
 - Proposition 12.17.1, 782
 - Proposition 12.18.3, 787
 - Proposition 12.19.4, 790
 - Corollary 12.16.15, 781
 - right divides**
 - definition, 234
 - right coprime**
 - polynomial matrices**
 - Bezout identity
 - Theorem 4.7.14, 252
 - right equivalence**
 - equivalence relation
 - Fact 5.10.3, 317
 - right inverse**
 - (1)-inverse
 - Proposition 6.1.2, 364
 - definition, 4
 - generalized inverse
 - Corollary 6.1.4, 364
 - idempotent matrix
 - Fact 3.12.10, 199
 - linear system
 - Fact 6.3.1, 369
 - matrix product**
 - Fact 2.15.6, 141
 - positive-semidefinite matrix**
 - Fact 3.7.26, 182
 - projector**
 - Fact 3.13.6, 207
 - representation**
 - Fact 2.15.4, 140
 - right-inner matrix**
 - Fact 3.11.5, 190
 - transfer function**
 - Fact 12.22.9, 799
 - uniqueness**
 - Theorem 1.2.2, 4
 - right-equivalent matrices**
 - definition
 - Definition 3.4.3, 174
 - group-invertible matrix**
 - Fact 3.6.1, 177
 - Kronecker product**
 - Fact 7.4.11, 405
 - range**
 - Proposition 5.1.3, 283
 - range-Hermitian matrix**
 - Fact 3.6.3, 177
 - right-inner matrix**
 - definition
 - Definition 3.1.2, 166
 - generalized inverse**
 - Fact 6.3.8, 371
 - right inverse**
 - Fact 3.11.5, 190
 - right-invertible function**
 - definition, 4
 - right-invertible matrix**
 - definition, 98
 - equivalent properties
 - Theorem 2.6.1, 98
 - generalized inverse
 - Proposition 6.1.5, 364
 - inverse
 - Proposition 2.6.5, 101
 - linear system**
 - solution
 - Fact 2.13.7, 129
 - matrix product**
 - Fact 2.10.3, 115
 - nonsingular equivalence**
 - Corollary 2.6.6, 101
 - open set**
 - Theorem 10.3.6, 624
 - unique right inverse**
 - Proposition 2.6.2, 99
- rigid body**
 - inertia matrix**
 - Fact 8.9.5, 451
- rigid-body rotation**
 - matrix exponential**
 - Fact 11.11.6, 673
- Rodrigues**
 - orthogonal matrix**
 - Fact 3.11.10, 192
- Rodrigues's formulas**
 - Euler parameters**
 - Fact 3.11.11, 193
 - orthogonal matrix**
 - Fact 3.11.11, 193
 - quaternions**
 - Fact 3.11.11, 193
- Rogers-Hölder inequality**
 - scalar case
 - Fact 1.16.12, 62
- root**
 - Definition 1.4.2, 8
 - polynomial**
 - Fact 11.20.4, 709
 - Fact 11.20.5, 709
 - Fact 11.20.6, 709
 - Fact 11.20.7, 710
 - Fact 11.20.8, 710
 - Fact 11.20.9, 710
 - Fact 11.20.10, 711
- root bounds**
 - polynomial**
 - Fact 11.20.11, 711
 - Fact 11.20.12, 712

root locus
 eigenvalue
 Fact 4.10.28, 272

roots
 polynomial
 Fact 4.8.1, 253
 Fact 4.8.2, 254

roots of polynomial
 convex hull
 Fact 10.11.3, 638

Rosenbrock system
 matrix
 definition
 Definition 12.10.1, 757
 rank
 Proposition 12.10.3,
 759
 Proposition 12.10.11,
 764

rotation
 vector
 Fact 3.11.13, 194

rotation matrix
 definition, 172
 logarithm
 Fact 11.15.10, 692
 orthogonal matrix
 Fact 3.11.9, 191
 Fact 3.11.10, 192
 Fact 3.11.11, 193
 Fact 3.11.12, 194
 Fact 3.11.31, 198

reflector
 Fact 3.11.9, 191
 trace
 Fact 3.11.17, 195

rotation-dilation
 factorization
 Fact 2.19.2, 151

Roth
 solutions of
 Sylvester's equation
 Fact 5.10.20, 320
 Fact 5.10.21, 320

Roup

positive-definite
 matrix
 Fact 8.8.14, 449

Routh criterion
 asymptotically stable
 polynomial
 Fact 11.17.2, 696

Routh form
 tridiagonal matrix
 Fact 11.18.27, 703

row
 definition, 79

row norm
 column norm
 Fact 9.8.10, 572
 definition, 556
 Hölder-induced
 norm
 Fact 9.8.21, 575
 Fact 9.8.23, 575
 Kronecker product
 Fact 9.9.61, 591
 partitioned matrix
 Fact 9.8.11, 572
 spectral radius
 Corollary 9.4.10, 556

S

S-N decomposition
 diagonalizable
 matrix
 Fact 5.9.3, 311
 nilpotent matrix
 Fact 5.9.3, 311

scalar inequality
 arithmetic mean
 Fact 1.11.6, 39
 Bernoulli's
 inequality
 Fact 1.9.1, 22
 Cauchy-Schwarz
 inequality
 Fact 1.16.9, 62
 exponential function
 Fact 1.9.14, 25

Fact 1.9.15, 25
 Fact 1.9.16, 25
 Fact 1.9.17, 26
 geometric mean
 Fact 1.11.6, 39
 Hölder's inequality
 Fact 1.16.11, 62
 Fact 1.16.12, 62
 Hua's inequality
 Fact 1.15.13, 51
 Kantorovich
 inequality
 Fact 1.15.36, 57
 logarithm
 Fact 1.15.45, 59
 Fact 1.15.46, 59
 Fact 1.15.47, 59
 Minkowski's
 inequality
 Fact 1.16.25, 66
 rearrangement
 inequality
 Fact 1.16.7, 61
 reciprocal powers
 Fact 1.16.26, 66
 Fact 1.16.27, 66
 reversal of Hölder's
 inequality
 Fact 1.16.22, 65
 Rogers-Hölder
 inequality
 Fact 1.16.12, 62
 Schweitzer's
 inequality
 Fact 1.15.37, 57
 Wang's inequality
 Fact 1.15.13, 51
 Young inequality
 Fact 1.10.21, 33
 Young's inequality
 Fact 1.10.32, 36
 Fact 1.15.31, 56

Schatten norm
 absolute value
 Fact 9.13.11, 603
 Cartesian
 decomposition
 Fact 9.9.37, 586
 Fact 9.9.39, 587

Fact 9.9.40, 587
Clarkson inequalities
 Fact 9.9.34, 586
commutator
 Fact 9.9.27, 584
compatible norms
 Proposition 9.3.6, 551
 Corollary 9.3.7, 552
 Corollary 9.3.8, 552
definition
 Proposition 9.2.3, 548
eigenvalue
 Fact 9.11.6, 598
equality
 Fact 9.9.33, 585
Frobenius norm
 Fact 9.8.20, 575
Hanner inequality
 Fact 9.9.36, 586
Hermitian matrix
 Fact 9.9.27, 584
 Fact 9.9.39, 587
Hölder matrix norm
 Fact 9.11.6, 598
Hölder norm
 Proposition 9.2.5, 549
inequality
 Fact 9.9.34, 586
 Fact 9.9.36, 586
 Fact 9.9.37, 586
 Fact 9.9.38, 587
 Fact 9.9.45, 588
Kronecker product
 Fact 9.14.37, 617
matrix difference
 Fact 9.9.23, 584
monotonicity
 Proposition 9.2.4, 549
normal matrix
 Fact 9.9.27, 584
 Fact 9.14.5, 608
partitioned matrix
 Fact 9.10.2, 593
 Fact 9.10.3, 594
 Fact 9.10.4, 594
 Fact 9.10.5, 595
 Fact 9.10.6, 595
 Fact 9.10.7, 596
 Fact 9.10.8, 596

positive-semidefinite matrix
 Fact 9.9.22, 583
 Fact 9.9.39, 587
 Fact 9.9.40, 587
 Fact 9.10.6, 595
 Fact 9.10.7, 596
Schur product
 Fact 9.14.34, 616
trace
 Fact 9.12.1, 599
unitarily invariant norm
 Fact 9.8.9, 572

Schauder fixed-point theorem
 image of a
 continuous function
 Theorem 10.3.10, 625

Schinzel
 determinant upper
 bound
 Fact 2.13.15, 131

Schmidt-Mirsky theorem
 fixed-rank
 approximation
 Fact 9.14.28, 614

Schneider
 inertia of a
 Hermitian matrix
 Fact 12.21.4, 794
 Fact 12.21.5, 794

Schoenberg
 Euclidean distance
 matrix
 Fact 9.8.14, 573

Schott's theorem
 Schur product of
 positive-
 semidefinite
 matrices
 Fact 8.21.12, 533

Schur
 dimension of the
 algebra generated

by commuting
 matrices
 Fact 5.10.15, 319

Schur complement
convex function
 Proposition 8.6.17, 437
 Lemma 8.6.16, 436

definition
 Definition 6.1.8, 367

determinant
 Proposition 8.2.3, 420

increasing function
 Proposition 8.6.13, 435

inequality
 Fact 8.11.17, 471

inertia
 Fact 6.5.5, 386

nondecreasing function
 Proposition 8.6.13, 435

partitioned matrix

Fact 6.5.4, 386
 Fact 6.5.5, 386
 Fact 6.5.6, 386
 Fact 6.5.8, 387
 Fact 6.5.12, 388
 Fact 6.5.29, 393
 Fact 8.21.39, 539

positive-semidefinite matrix

Corollary 8.6.18, 442
 Fact 8.11.3, 468
 Fact 8.11.4, 468
 Fact 8.11.18, 471
 Fact 8.11.19, 471
 Fact 8.11.20, 472
 Fact 8.11.27, 474
 Fact 8.20.19, 530
 Fact 8.21.11, 533

rank
 Proposition 8.2.3, 420
 Fact 6.5.6, 386
 Fact 6.5.8, 387
 Fact 6.5.11, 388

Schur product
 Fact 8.21.11, 533

Schur concave function
 definition

- Definition 2.1.2, 78
- elementary symmetric function**
 - Fact 1.15.20, 53
- entropy**
 - Fact 2.21.6, 162
- strong majorization**
 - Fact 2.21.6, 162
- Schur convex function**
 - definition**
 - Definition 2.1.2, 78
 - Muirhead's theorem**
 - Fact 1.15.25, 54
 - strong majorization**
 - Fact 2.21.4, 162
 - Fact 2.21.5, 162
- Schur decomposition**
 - Hermitian matrix**
 - Corollary 5.4.5, 294
 - Jordan form**
 - Fact 5.10.6, 317
 - normal matrix**
 - Corollary 5.4.4, 293
 - Fact 5.10.6, 317
 - range-Hermitian matrix**
 - Corollary 5.4.4, 293
- Schur inverse**
 - positive-semidefinite matrix**
 - Fact 8.21.1, 531
- Schur power**
 - definition, 404**
 - Lyapunov equation**
 - Fact 8.8.16, 449
 - positive-semidefinite matrix**
 - Fact 8.21.2, 531
 - Fact 8.21.3, 531
 - Fact 8.21.25, 536
- Schur product**
 - associative identities, 404**
 - commutative identities, 404**
 - complex conjugate transpose**
 - Fact 8.21.9, 533
 - definition, 404**
 - distributive identities, 404**
 - eigenvalue**
 - Fact 8.21.18, 534
 - Frobenius norm**
 - Fact 9.14.34, 616
 - geometric mean**
 - Fact 8.21.51, 541
 - Hermitian matrix**
 - Fact 8.21.28, 536
 - Fact 8.21.32, 537
 - Kronecker product**
 - Proposition 7.3.1, 404
 - lower bound**
 - Fact 8.21.14, 534
 - M-matrix**
 - Fact 7.6.15, 415
 - matrix exponential**
 - Fact 11.14.21, 685
 - matrix identity**
 - Fact 7.6.3, 413
 - Fact 7.6.4, 413
 - Fact 7.6.10, 414
 - matrix logarithm**
 - Fact 8.21.47, 540
 - Fact 8.21.48, 540
 - matrix power**
 - Fact 7.6.11, 414
 - matrix-vector identity**
 - Fact 7.6.9, 414
 - maximum singular value**
 - Fact 8.21.10, 533
 - Fact 9.14.31, 615
 - Fact 9.14.33, 616
 - Fact 9.14.35, 617
 - nonnegative matrix**
 - Fact 7.6.13, 415
 - normal matrix**
 - Fact 9.9.63, 591
 - partitioned matrix**
 - Fact 8.21.6, 532
 - Fact 8.21.39, 539
 - Fact 8.21.40, 539
 - positive matrix**
 - Fact 7.6.14, 415
 - positive-definite matrix**
 - Fact 8.21.4, 531
 - Fact 8.21.5, 532
 - Fact 8.21.6, 532
 - Fact 8.21.7, 533
 - Fact 8.21.13, 533
 - Fact 8.21.14, 534
 - Fact 8.21.15, 534
 - Fact 8.21.21, 535
 - Fact 8.21.33, 538
 - Fact 8.21.34, 538
 - Fact 8.21.36, 538
 - Fact 8.21.38, 539
 - Fact 8.21.42, 539
 - Fact 8.21.47, 540
 - Fact 8.21.49, 541
 - Fact 8.21.50, 541
 - Fact 8.21.51, 541
 - positive-semidefinite matrix**
 - Fact 8.21.4, 531
 - Fact 8.21.7, 533
 - Fact 8.21.11, 533
 - Fact 8.21.12, 533
 - Fact 8.21.14, 534
 - Fact 8.21.17, 534
 - Fact 8.21.18, 534
 - Fact 8.21.20, 535
 - Fact 8.21.22, 536
 - Fact 8.21.23, 536
 - Fact 8.21.31, 537
 - Fact 8.21.35, 538
 - Fact 8.21.37, 538
 - Fact 8.21.39, 539
 - Fact 8.21.40, 539
 - Fact 8.21.41, 539
 - Fact 8.21.42, 539
 - Fact 8.21.43, 540
 - Fact 8.21.44, 540
 - Fact 8.21.45, 540
 - Fact 8.21.46, 540
 - quadratic form**
 - Fact 7.6.7, 413
 - range**
 - Fact 7.6.5, 413
 - rank**
 - Fact 7.6.6, 413
 - Fact 8.21.16, 534

- Schatten norm**
 - Fact 9.14.34, 616
- Schur complement**
 - Fact 8.21.11, 533
- singular value**
 - Fact 9.14.31, 615
 - Fact 9.14.32, 615
 - Fact 9.14.33, 616
- spectral radius**
 - Fact 7.6.13, 415
 - Fact 7.6.14, 415
 - Fact 7.6.16, 416
 - Fact 7.6.17, 416
 - Fact 9.14.33, 616
- submultiplicative norm**
 - Fact 9.8.41, 578
- trace**
 - Fact 7.6.8, 413
 - Fact 8.21.17, 534
 - Fact 9.14.32, 615
- transpose**
 - Fact 7.6.12, 414
- unitarily invariant norm**
 - Fact 9.8.41, 578
 - Fact 9.9.62, 591
 - Fact 9.9.63, 591
 - Fact 9.14.36, 617
- vector identity**
 - Fact 7.6.1, 413
 - Fact 7.6.2, 413
- weak majorization**
 - Fact 9.14.31, 615
- Schur product of polynomials**
 - asymptotically stable polynomial
 - Fact 11.17.9, 697
- Schur's formulas**
 - determinant of partitioned matrix
 - Fact 2.14.13, 135
- Schur's inequality**
 - eigenvalue
 - Fact 8.17.5, 509
 - eigenvalues and the Frobenius norm
 - Fact 9.11.3, 597
- Schur's theorem**
 - eigenvalue inequality
 - Fact 8.17.8, 510
- Schur product of positive-semidefinite matrices**
 - Fact 8.21.12, 533
- Schur-Cohn criterion**
 - discrete-time asymptotically stable polynomial
 - Fact 11.20.1, 708
- Schur-Horn theorem**
 - diagonal entries of a unitary matrix
 - Fact 3.11.19, 195
 - Fact 8.17.10, 511
- Schwarz form**
 - tridiagonal matrix
 - Fact 11.18.25, 702
 - Fact 11.18.26, 702
- Schweitzer's inequality**
 - scalar inequality
 - Fact 1.15.37, 57
- secant condition**
 - asymptotically stable matrix
 - Fact 11.18.29, 704
- second derivative**
 - definition, 627
- Seiler**
 - determinant inequality
 - Fact 8.13.30, 490
- self-adjoint norm**
 - definition, 547
 - unitarily invariant norm
 - Fact 9.8.7, 572
- self-conjugate set**
 - definition, 232
- semicontractive matrix**
 - complex conjugate transpose
 - Fact 3.20.12, 220
 - definition
 - Definition 3.1.2, 166
 - discrete-time Lyapunov-stable matrix
 - Fact 11.21.4, 712
 - partitioned matrix
 - Fact 8.11.6, 469
 - Fact 8.11.22, 473
 - positive-semidefinite matrix
 - Fact 8.11.6, 469
 - Fact 8.11.13, 470
 - unitary matrix
 - Fact 8.11.22, 473
- semidissipative matrix**
 - definition
 - Definition 3.1.1, 165
- determinant**
 - Fact 8.13.3, 485
 - Fact 8.13.4, 485
 - Fact 8.13.11, 486, 487
- discrete-time**
 - Lyapunov-stable matrix
 - Fact 11.21.4, 712
- dissipative matrix**
 - Fact 8.13.31, 491
- Kronecker sum**
 - Fact 7.5.8, 409
- Lyapunov-stable matrix**
 - Fact 11.18.37, 705
- normal matrix**
 - Fact 11.18.37, 705
- semiperimeter**
 - quadrilateral
 - Fact 2.20.13, 159
 - triangle
 - Fact 2.20.11, 156
- semisimple eigenvalue**
 - cyclic eigenvalue
 - Proposition 5.5.5, 296
 - defect
 - Proposition 5.5.8, 296

- definition**
 - Definition 5.5.4, 296
- index of an eigenvalue**
 - Proposition 5.5.8, 296
- null space**
 - Proposition 5.5.8, 296
- simple eigenvalue**
 - Proposition 5.5.5, 296
- semisimple matrix**
- cyclic matrix**
 - Fact 5.14.11, 340
- definition**
 - Definition 5.5.4, 296
- elementary matrix**
 - Fact 5.14.17, 341
- idempotent matrix**
 - Fact 5.14.21, 341
- identity perturbation**
 - Fact 5.14.16, 341
- involutory matrix**
 - Fact 5.14.19, 341
- Kronecker product**
 - Fact 7.4.16, 406
- matrix exponential**
 - Proposition 11.2.7, 648
- normal matrix**
 - Proposition 5.5.11, 297
- outer-product matrix**
 - Fact 5.14.3, 338
- positive-semidefinite matrix**
 - Corollary 8.3.6, 424
- reverse-diagonal matrix**
 - Fact 5.14.12, 340
- similar matrices**
 - Proposition 5.5.11, 297
 - Fact 5.9.4, 312
 - Fact 5.10.5, 317
- simple matrix**
 - Fact 5.14.11, 340
- skew-involutory matrix**
 - Fact 5.14.19, 341
- semistability eigenvalue**
 - Proposition 11.8.2, 662
- linear dynamical system**
 - Proposition 11.8.2, 662
- Lyapunov equation**
 - Corollary 11.9.1, 666
- matrix exponential**
 - Proposition 11.8.2, 662
- semistable matrix**
- compartmental matrix**
 - Fact 11.19.6, 707
- definition**
 - Definition 11.8.1, 662
- group-invertible matrix**
 - Fact 11.18.3, 698
- Kronecker sum**
 - Fact 11.18.32, 704
 - Fact 11.18.33, 704
- limit**
 - Fact 11.18.7, 699
- Lyapunov equation**
 - Fact 12.21.15, 797
- Lyapunov-stable matrix**
 - Fact 11.18.1, 698
- matrix exponential**
 - Fact 11.18.5, 698
 - Fact 11.18.7, 699
 - Fact 11.21.7, 713
- minimal realization**
 - Definition 12.9.17, 757
- semistable polynomial**
 - Proposition 11.8.4, 663
- similar matrices**
 - Fact 11.18.4, 698
- unstable subspace**
 - Proposition 11.8.8, 665
- semistable polynomial**
- definition**
 - Definition 11.8.3, 663
- reciprocal argument**
 - Fact 11.17.5, 696
- semistable matrix**
 - Proposition 11.8.4, 663
- semistable transfer function**
- minimal realization**
 - Proposition 12.9.18, 757
- SISO entries**
 - Proposition 12.9.19, 757
- separation theorem**
- convex cone**
 - Fact 10.9.13, 635
- inner product**
 - Fact 10.9.13, 635
 - Fact 10.9.14, 635
- sequence**
- definition**
 - Definition 10.2.1, 622
- generalized inverse**
 - Fact 6.3.36, 377
- series**
- commutator**
 - Fact 11.14.17, 684
- definition**
 - Definition 10.2.6, 623
 - Definition 10.2.8, 623
- inverse matrix**
 - Proposition 9.4.13, 557
- matrix exponential**
 - Fact 11.14.17, 684
- set**
- definition, 2**
- distance from a point**
 - Fact 10.9.15, 636
 - Fact 10.9.16, 636
- set cancellation**
- convex set**
 - Fact 10.9.7, 635
- Radstrom**
 - Fact 10.9.7, 635
- set identities**
- intersection**
 - Fact 1.5.6, 11
- union**
 - Fact 1.5.6, 11
- sextic**

- arithmetic-mean–geometric-mean inequality
 - Fact 1.13.1, 47
- Shannon’s inequality**
 - logarithm
 - Fact 1.16.30, 67
- shear**
 - factorization
 - Fact 5.15.11, 346
- Shemesh**
 - common eigenvector
 - Fact 5.14.27, 342
- Sherman-Morrison-Woodbury formula**
 - determinant of an outer-product perturbation
 - Fact 2.16.3, 141
- shift**
 - controllability
 - Fact 12.20.10, 792
 - stabilizability
 - Fact 12.20.11, 792
- shifted argument**
 - transfer function
 - Fact 12.22.3, 798
- shifted-orthogonal matrix**
 - definition
 - Definition 3.1.1, 165
- shifted-unitary matrix**
 - block-diagonal matrix
 - Fact 3.11.25, 196
 - definition
 - Definition 3.1.1, 165
 - normal matrix
 - Fact 3.11.34, 198
 - spectrum
 - Proposition 5.5.21, 300
 - unitary matrix
 - Fact 3.11.33, 198
- Shoda**
 - factorization
 - Fact 5.15.7, 346
 - Fact 5.15.34, 351
 - Shoda’s theorem**
 - commutator realization
 - Fact 5.9.18, 313
 - zero trace
 - Fact 5.9.18, 313
 - shortcut of a relation**
 - definition
 - Definition 1.3.4, 5
 - shorted operator**
 - definition
 - Fact 8.20.19, 530
 - positive-semidefinite matrix
 - Fact 8.20.19, 530
 - sign**
 - matrix, 89
 - vector, 89
 - sign of entries**
 - asymptotically stable matrix
 - Fact 11.19.5, 708
 - sign stability**
 - asymptotically stable matrix
 - Fact 11.19.5, 708
 - signature**
 - definition, 245
 - Hermitian matrix
 - Fact 5.8.6, 308
 - Fact 5.8.7, 308
 - Fact 8.10.17, 457
 - involutory matrix
 - Fact 5.8.2, 307
 - positive-semidefinite matrix
 - Fact 5.8.9, 308
 - tripotent matrix
 - Fact 5.8.3, 307
 - signed volume**
 - simplex
 - Fact 2.20.15, 160
 - similar matrices**
 - asymptotically stable matrix
 - Fact 11.18.4, 698
 - biequivalent matrices
 - Proposition 3.4.5, 174
 - block-diagonal matrix
 - Theorem 5.3.2, 288
 - Theorem 5.3.3, 289
 - campanion matrix
 - Fact 5.16.5, 354
 - characteristic polynomial
 - Fact 4.9.10, 262
 - complex conjugate
 - Fact 5.9.31, 316
 - cyclic matrix
 - Fact 5.16.5, 354
 - definition
 - Definition 3.4.4, 174
 - diagonal entries
 - Fact 5.9.13, 313
 - diagonalizable over $\mathbb{R}$
 - Proposition 5.5.12, 297
 - Corollary 5.5.22, 301
 - discrete-time
 - asymptotically stable matrix
 - Fact 11.18.4, 698
 - discrete-time Lyapunov-stable matrix
 - Fact 11.18.4, 698
 - discrete-time semistable matrix
 - Fact 11.18.4, 698
 - equivalence class
 - Fact 5.10.4, 317
 - equivalent realizations
 - Definition 12.9.6, 751
 - example
 - Example 5.5.20, 300
 - factorization
 - Fact 5.15.6, 346
 - geometric multiplicity
 - Proposition 5.5.10, 297

- group-invertible matrix**
 - Proposition 3.4.5, 174
 - Fact 5.9.5, 312
- Hermitian matrix**
 - Proposition 5.5.12, 297
- idempotent matrix**
 - Proposition 3.4.5, 174
 - Proposition 5.6.3, 302
 - Corollary 5.5.22, 301
 - Fact 5.10.9, 318
 - Fact 5.10.13, 319
 - Fact 5.10.14, 319
 - Fact 5.10.22, 320
- inverse matrix**
 - Fact 5.15.31, 350
- involutory matrix**
 - Proposition 3.4.5, 174
 - Corollary 5.5.22, 301
 - Fact 5.15.31, 350
- Kronecker product**
 - Fact 7.4.12, 406
- Kronecker sum**
 - Fact 7.5.9, 409
- lower triangular matrix**
 - Fact 5.9.2, 311
- Lyapunov-stable matrix**
 - Fact 11.18.4, 698
- matrix classes**
 - Proposition 3.4.5, 174
- matrix exponential**
 - Proposition 11.2.9, 650
- matrix power**
 - Fact 5.9.1, 311
- minimal polynomial**
 - Proposition 4.6.3, 248
 - Fact 11.23.3, 717
 - Fact 11.23.4, 717
 - Fact 11.23.5, 718
 - Fact 11.23.6, 719
 - Fact 11.23.7, 719
 - Fact 11.23.8, 720
 - Fact 11.23.9, 720
 - Fact 11.23.10, 721
 - Fact 11.23.11, 721
- multicompanion form**
 - Corollary 5.2.6, 286
- nilpotent matrix**
 - Proposition 3.4.5, 174
 - Fact 5.10.23, 321
- nonsingular matrix**
 - Fact 5.10.11, 318
- nonzero diagonal entries**
 - Fact 5.9.14, 313
- normal matrix**
 - Proposition 5.5.11, 297
 - Fact 5.9.9, 312
 - Fact 5.9.10, 312
 - Fact 5.10.7, 317
- partitioned matrix**
 - Fact 5.10.21, 320
 - Fact 5.10.22, 320
 - Fact 5.10.23, 321
- projector**
 - Corollary 5.5.22, 301
 - Fact 5.10.13, 319
- realization**
 - Proposition 12.9.5, 751
- reflector**
 - Corollary 5.5.22, 301
- reverse transpose**
 - Fact 5.9.11, 313
- reverse-symmetric matrix**
 - Fact 5.9.11, 313
- semisimple matrix**
 - Proposition 5.5.11, 297
 - Fact 5.9.4, 312
 - Fact 5.10.5, 317
- semistable matrix**
 - Fact 11.18.4, 698
- similarity invariant**
 - Theorem 4.3.10, 239
 - Corollary 5.2.6, 286
- simultaneous diagonalization**
 - Fact 5.17.8, 358
- skew-Hermitian matrix**
 - Fact 5.9.4, 312
 - Fact 11.18.12, 700
- skew-idempotent matrix**
 - Corollary 5.5.22, 301
- skew-involutory matrix**
 - Proposition 3.4.5, 174
- skew-symmetric matrix**
 - Fact 5.15.39, 351
- Sylvester's equation**
 - Corollary 7.2.5, 404
 - Fact 7.5.14, 410
- symmetric matrix**
 - Fact 5.15.39, 351
- transpose**
 - Proposition 5.5.12, 297
 - Corollary 4.3.11, 239
 - Corollary 5.3.8, 291
 - Corollary 5.5.22, 301
 - Fact 5.9.9, 312
 - Fact 5.9.10, 312
- tripotent matrix**
 - Proposition 3.4.5, 174
 - Corollary 5.5.22, 301
- unitarily invariant norm**
 - Fact 9.8.31, 576
- unitarily similar matrices**
 - Fact 5.10.7, 317
- upper triangular matrix**
 - Fact 5.9.2, 311
- Vandermonde matrix**
 - Fact 5.16.5, 354
- similarity**
 - equivalence relation**
 - Fact 5.10.3, 317
- similarity invariant**
 - characteristic polynomial**
 - Proposition 4.4.2, 240
 - Proposition 4.6.2, 248
 - definition**
 - Definition 4.3.9, 239
 - multicompanion form**
 - Corollary 5.2.6, 286
 - similar matrices**
 - Theorem 4.3.10, 239
 - Corollary 5.2.6, 286

similarity**transformation****complex conjugate****transpose**

Fact 5.9.8, 312

Fact 5.15.4, 345

complex symmetric**Jordan form**

Fact 5.15.2, 345

Fact 5.15.3, 345

eigenvector

Fact 5.14.6, 339

Fact 5.14.7, 339

hypercompanion**matrix**

Fact 5.10.1, 316

inverse matrix

Fact 5.15.4, 345

normal matrix

Fact 5.15.3, 345

real Jordan form

Fact 5.10.1, 316

Fact 5.10.2, 317

symmetric matrix

Fact 5.15.2, 345

Fact 5.15.3, 345

SIMO transfer function**definition**

Definition 12.9.1, 749

Simon**determinant****inequality**

Fact 8.13.30, 490

normal product and**Schatten norm**

Fact 9.14.5, 608

simple eigenvalue**cyclic eigenvalue**

Proposition 5.5.5, 296

definition

Definition 5.5.4, 296

semisimple**eigenvalue**

Proposition 5.5.5, 296

simple graph**definition**

Definition 1.4.3, 9

simple matrix**commuting matrices**

Fact 5.14.23, 342

cyclic matrix

Fact 5.14.11, 340

definition

Definition 5.5.4, 296

identity perturbation

Fact 5.14.16, 341

rank

Fact 5.11.1, 321

semisimple matrix

Fact 5.14.11, 340

simplex**convex hull**

Fact 2.20.4, 154

definition, 90**interior**

Fact 2.20.4, 154

nonsingular matrix

Fact 2.20.4, 154

signed volume

Fact 2.20.15, 160

volume

Fact 2.20.19, 160

simultaneous**diagonalization****cogredient****transformation**

Fact 8.16.4, 507

Fact 8.16.6, 507

commuting matrices

Fact 8.16.1, 507

definition, 422**diagonalizable****matrix**

Fact 8.16.2, 507

Fact 8.16.3, 507

Hermitian matrix

Fact 8.16.1, 507

Fact 8.16.4, 507

Fact 8.16.6, 507

positive-definite**matrix**

Fact 8.16.5, 507

similar matrices

Fact 5.17.8, 358

unitarily similar**matrices**

Fact 5.17.7, 358

unitary matrix

Fact 8.16.1, 507

simultaneous**diagonalization of
symmetric matrices****Milnor**

Fact 8.16.6, 507

Pesonen

Fact 8.16.6, 507

simultaneous**orthogonal
biequivalence
transformation****upper Hessenberg**

Fact 5.17.3, 358

upper triangular

Fact 5.17.3, 358

simultaneous**triangularization****cogredient****transformation**

Fact 5.17.9, 358

common eigenvector

Fact 5.17.1, 358

commutator

Fact 5.17.5, 358

Fact 5.17.6, 358

commuting matrices

Fact 5.17.4, 358

nilpotent matrix

Fact 5.17.6, 358

projector

Fact 5.17.6, 358

regular pencil

Fact 5.17.2, 358

simultaneous unitary**biequivalence****transformation**

Fact 5.17.2, 358

unitarily similar**matrices**

Fact 5.17.4, 358

Fact 5.17.6, 358

- simultaneous unitary biequivalence transformation**
 - simultaneous triangularization
 - Fact 5.17.2, 358
- sine rule**
 - triangle
 - Fact 2.20.11, 156
- singular matrix**
 - definition, 101
 - Kronecker product
 - Fact 7.4.27, 407
 - spectrum
 - Proposition 5.5.21, 300
- singular pencil**
 - definition, 304
 - generalized eigenvalue
 - Proposition 5.7.3, 305
- singular polynomial matrix**
 - Definition 4.2.5, 235
- singular value**
 - 2×2 matrix
 - Fact 5.11.31, 328
 - adjugate
 - Fact 5.11.36, 328
 - bidiagonal matrix
 - Fact 5.11.47, 332
 - block-diagonal matrix
 - Fact 8.18.9, 515
 - Fact 8.18.10, 515
 - Fact 9.14.21, 612
 - Fact 9.14.25, 613
 - Cartesian decomposition
 - Fact 8.18.7, 514
 - companion matrix
 - Fact 5.11.30, 327
 - complex conjugate transpose
 - Fact 5.11.20, 324
 - Fact 5.11.34, 328
 - convex function
 - Fact 11.16.14, 695
 - Fact 11.16.15, 695
 - definition
 - Definition 5.6.1, 301
 - determinant
 - Fact 5.11.28, 326
 - Fact 5.11.29, 327
 - Fact 8.13.1, 485
 - Fact 9.13.23, 606
 - eigenvalue
 - Fact 8.17.5, 509
 - Fact 8.17.6, 509
 - Fact 9.13.22, 606
 - eigenvalue of Hermitian part
 - Fact 5.11.27, 326
 - Fact 8.17.4, 509
 - Fan dominance theorem
 - Fact 9.14.19, 611
 - fixed-rank approximation
 - Fact 9.14.28, 614
 - Fact 9.15.4, 618
 - Frobenius
 - Corollary 9.6.7, 562
 - generalized inverse
 - Fact 6.3.29, 376
 - homogeneity
 - Fact 5.11.19, 324
 - idempotent matrix
 - Fact 5.11.38, 328
 - induced lower bound
 - Proposition 9.5.4, 560
 - inequality
 - Proposition 9.2.2, 548
 - Corollary 9.6.5, 562
 - Fact 9.14.23, 612
 - Fact 9.14.24, 613
 - interlacing
 - Fact 9.14.10, 609
 - matrix difference
 - Fact 8.18.9, 515
 - Fact 8.18.10, 515
 - matrix exponential
 - Fact 11.15.5, 689
 - Fact 11.16.14, 695
 - Fact 11.16.15, 695
 - matrix power
 - Fact 9.13.19, 605
 - Fact 9.13.20, 605
 - matrix product
 - Proposition 9.6.1, 560
 - Proposition 9.6.2, 561
 - Proposition 9.6.3, 561
 - Proposition 9.6.4, 561
 - Fact 8.18.21, 519
 - Fact 9.13.17, 604
 - Fact 9.13.18, 605
 - Fact 9.14.26, 613
 - matrix sum
 - Proposition 9.6.8, 562
 - Fact 9.14.20, 612
 - Fact 9.14.21, 612
 - Fact 9.14.25, 613
 - normal matrix
 - Fact 5.14.15, 341
 - outer-product matrix
 - Fact 5.11.17, 324
 - partitioned matrix
 - Proposition 5.6.6, 303
 - Fact 9.14.11, 609
 - Fact 9.14.24, 613
 - perturbation
 - Fact 9.14.6, 608
 - positive-semidefinite matrix
 - Fact 8.18.7, 514
 - Fact 9.14.27, 613
 - rank
 - Proposition 5.6.2, 302
 - Fact 9.14.28, 614
 - Fact 9.15.4, 618
 - Schur product
 - Fact 9.14.31, 615
 - Fact 9.14.32, 615
 - Fact 9.14.33, 616
 - strong log majorization
 - Fact 9.13.19, 605
 - submatrix
 - Fact 9.14.10, 609
 - trace
 - Fact 5.12.6, 334
 - Fact 8.17.2, 508
 - Fact 9.12.1, 599
 - Fact 9.13.16, 604
 - Fact 9.14.3, 607

- Fact 9.14.32, 615
- unitarily invariant norm**
 - Fact 9.14.28, 614
- unitary matrix**
 - Fact 5.11.37, 328
 - Fact 9.14.11, 609
- weak log majorization**
 - Proposition 9.6.2, 561
- weak majorization**
 - Proposition 9.2.2, 548
 - Proposition 9.6.3, 561
 - Fact 5.11.27, 326
 - Fact 8.17.5, 509
 - Fact 8.18.7, 514
 - Fact 8.18.21, 519
 - Fact 9.13.17, 604
 - Fact 9.13.18, 605
 - Fact 9.13.20, 605
 - Fact 9.14.19, 611
 - Fact 9.14.20, 612
 - Fact 9.14.31, 615
- Weyl majorant theorem**
 - Fact 9.13.20, 605
- singular value decomposition**
 - existence
 - Theorem 5.6.4, 302
- generalized inverse**
 - Fact 6.3.15, 373
- group generalized inverse**
 - Fact 6.6.15, 395
- least squares**
 - Fact 9.14.28, 614
 - Fact 9.15.4, 618
 - Fact 9.15.5, 618
 - Fact 9.15.6, 619
- unitary similarity**
 - Fact 5.9.28, 315
 - Fact 6.3.15, 373
 - Fact 6.6.15, 395
- singular value perturbation**
 - unitarily invariant norm**
 - Fact 9.14.29, 614
- singular values**
 - determinant**
 - Fact 5.12.13, 335
 - positive-semidefinite matrix**
 - Fact 8.11.9, 469
 - unitarily biequivalent matrices**
 - Fact 5.10.18, 319
- SISO transfer function**
 - definition**
 - Definition 12.9.1, 749
- size**
 - definition, 79**
- skew reflector**
 - Hamiltonian matrix**
 - Fact 3.19.3, 216
 - reflector**
 - Fact 3.14.7, 211
- skew-Hermitian matrix**
 - Fact 3.14.6, 211
- skew-involutory matrix**
 - Fact 3.14.6, 211
- spectrum**
 - Proposition 5.5.21, 300
- unitary matrix**
 - Fact 3.14.6, 211
- skew-Hermitian matrix, see skew-symmetric matrix**
 - adjugate**
 - Fact 3.7.10, 179
 - Fact 3.7.11, 179
 - asymptotically stable matrix**
 - Fact 11.18.30, 704
 - block-diagonal matrix**
 - Fact 3.7.8, 179
 - Cartesian decomposition**
 - Fact 3.7.27, 182
 - Fact 3.7.28, 183
 - Fact 3.7.29, 183**
 - Cayley transform**
 - Fact 3.11.28, 196
 - characteristic polynomial**
 - Fact 4.9.13, 262
 - commutator**
 - Fact 3.8.1, 184
 - Fact 3.8.4, 185
 - complex conjugate**
 - Fact 3.12.8, 199
 - congruent matrices**
 - Proposition 3.4.5, 174
 - definition**
 - Definition 3.1.1, 165
 - determinant**
 - Fact 3.7.11, 179
 - Fact 3.7.16, 181
 - Fact 8.13.6, 486
 - eigenvalue**
 - Fact 5.11.6, 321
 - existence of transformation**
 - Fact 3.9.4, 186
 - Hermitian matrix**
 - Fact 3.7.9, 179
 - Fact 3.7.28, 183
 - inertia**
 - Fact 5.8.4, 307
 - Kronecker product**
 - Fact 7.4.17, 406
 - Kronecker sum**
 - Fact 7.5.8, 409
 - Lyapunov equation**
 - Fact 11.18.12, 700
 - matrix exponential**
 - Proposition 11.2.8, 649
 - Proposition 11.2.9, 650
 - Fact 11.14.6, 683
 - Fact 11.14.33, 688
 - matrix power**
 - Fact 8.9.14, 452
 - normal matrix**
 - Proposition 3.1.6, 169
 - null space**
 - Fact 8.7.3, 443
 - outer-product matrix**
 - Fact 3.7.17, 181

- Fact 3.9.4, 186
- partitioned matrix**
 - Fact 3.7.27, 182
- positive-definite matrix**
 - Fact 8.13.6, 486
 - Fact 11.18.12, 700
- positive-semidefinite matrix**
 - Fact 8.9.12, 452
- projector**
 - Fact 9.9.9, 581
- quadratic form**
 - Fact 3.7.6, 178
- range**
 - Fact 8.7.3, 443
- rank**
 - Fact 3.7.17, 181
 - Fact 3.7.30, 183
- similar matrices**
 - Fact 5.9.4, 312
 - Fact 11.18.12, 700
- skew reflector**
 - Fact 3.14.6, 211
- skew-involutory matrix**
 - Fact 3.14.6, 211
- skew-symmetric matrix**
 - Fact 3.7.9, 179
- spectrum**
 - Proposition 5.5.21, 300
- symmetric matrix**
 - Fact 3.7.9, 179
- trace**
 - Fact 3.7.24, 182
- trace of a product**
 - Fact 8.12.6, 476
- unitarily similar matrices**
 - Proposition 3.4.5, 174
 - Proposition 5.6.3, 302
- unitary matrix**
 - Fact 3.11.28, 196
 - Fact 3.14.6, 211
 - Fact 11.14.33, 688
- skew-idempotent matrix**
- idempotent matrix**
 - Fact 3.12.5, 199
- similar matrices**
 - Corollary 5.5.22, 301
- skew-involutory matrix**
- definition**
 - Definition 3.1.1, 165
- Hamiltonian matrix**
 - Fact 3.19.2, 216
 - Fact 3.19.3, 216
- inertia**
 - Fact 5.8.4, 307
- matrix exponential**
 - Fact 11.11.1, 671
- semisimple matrix**
 - Fact 5.14.19, 341
- similar matrices**
 - Proposition 3.4.5, 174
- size**
 - Fact 3.15.6, 212
- skew reflector**
 - Fact 3.14.6, 211
- skew-Hermitian matrix**
 - Fact 3.14.6, 211
- skew-symmetric matrix**
 - Fact 3.19.3, 216
- spectrum**
 - Proposition 5.5.21, 300
- symplectic matrix**
 - Fact 3.19.2, 216
- unitarily similar matrices**
 - Proposition 3.4.5, 174
- unitary matrix**
 - Fact 3.14.6, 211
- skew-symmetric matrix, see skew-Hermitian matrix**
- adjugate**
 - Fact 4.9.20, 263
- Cayley transform**
 - Fact 3.11.8, 190
 - Fact 3.11.28, 196
 - Fact 3.11.30, 197
 - Fact 3.11.31, 198
- characteristic polynomial**
 - Fact 4.9.12, 262
 - Fact 4.9.19, 263
 - Fact 4.9.20, 263
 - Fact 5.14.34, 343
- commutator**
 - Fact 3.8.5, 185
- congruent matrices**
 - Fact 3.7.34, 184
 - Fact 5.9.16, 313
- controllability**
 - Fact 12.20.5, 791
- definition**
 - Definition 3.1.1, 165
- determinant**
 - Fact 3.7.15, 181
 - Fact 3.7.33, 184
 - Fact 4.8.14, 259
 - Fact 4.9.20, 263
 - Fact 4.10.2, 266
- eigenvalue**
 - Fact 4.10.2, 266
- factorization**
 - Fact 5.15.37, 351
 - Fact 5.15.38, 351
- Hamiltonian matrix**
 - Fact 3.7.34, 184
 - Fact 3.19.3, 216
 - Fact 3.19.8, 217
- Hermitian matrix**
 - Fact 3.7.9, 179
- linear matrix equation**
 - Fact 3.7.3, 178
- matrix exponential**
 - Example 11.3.6, 652
 - Fact 11.11.3, 672
 - Fact 11.11.6, 673
 - Fact 11.11.7, 673
 - Fact 11.11.8, 674
 - Fact 11.11.9, 674
 - Fact 11.11.10, 674
 - Fact 11.11.11, 674
 - Fact 11.11.14, 675
 - Fact 11.11.15, 675
 - Fact 11.11.16, 676
 - Fact 11.11.17, 676
- matrix product**

Fact 5.15.37, 351
orthogonal matrix
 Fact 3.11.28, 196
 Fact 3.11.30, 197
 Fact 3.11.31, 198
 Fact 11.11.10, 674
 Fact 11.11.11, 674
orthogonally similar matrices
 Fact 5.14.33, 343
partitioned matrix
 Fact 3.11.27, 196
Pfaffian
 Fact 4.8.14, 259
quadratic form
 Fact 3.7.5, 178
similar matrices
 Fact 5.15.39, 351
skew-Hermitian matrix
 Fact 3.7.9, 179
skew-involutory matrix
 Fact 3.19.3, 216
spectrum
 Fact 4.9.20, 263
 Fact 4.10.2, 266
 Fact 5.14.33, 343
symmetric matrix
 Fact 5.9.16, 313
 Fact 5.15.39, 351
trace
 Fact 3.7.23, 182
 Fact 3.7.31, 183
unit imaginary matrix
 Fact 3.7.34, 184
small-gain theorem
 multiplicative
 perturbation
 Fact 9.13.23, 606
Smith form
 biequivalent matrices
 Theorem 5.1.1, 283
 Corollary 5.1.2, 283
 controllability pencil
 Proposition 12.6.15, 741

existence
 Theorem 4.3.2, 237
observability pencil
 Proposition 12.3.15, 731
polynomial matrix
 Proposition 4.3.4, 237
rank
 Proposition 4.3.5, 237
 Proposition 4.3.6, 238
submatrix
 Proposition 4.3.5, 237
unimodular matrix
 Proposition 4.3.7, 238

Smith polynomial
 nonsingular matrix
 transformation
 Proposition 4.3.8, 238

Smith polynomials
 definition
 Definition 4.3.3, 237

Smith zeros
 controllability pencil
 Proposition 12.6.16, 741
 definition
 Definition 4.3.3, 237
 observability pencil
 Proposition 12.3.16, 731
 uncontrollable
 spectrum
 Proposition 12.6.16, 741
 unobservable
 spectrum
 Proposition 12.3.16, 731

Smith's method
 finite-sum solution of
 Lyapunov equation
 Fact 12.21.17, 797

Smith-McMillan form
 blocking zero
 Proposition 4.7.11, 251
 coprime polynomials
 Fact 4.8.15, 259

coprime right polynomial fraction description
 Proposition 4.7.16, 253
existence
 Theorem 4.7.5, 250
poles
 Proposition 4.7.11, 251
rank
 Proposition 4.7.7, 250
 Proposition 4.7.8, 250
submatrix
 Proposition 4.7.7, 250

SO(2)
 parameterization
 Fact 3.11.6, 190

solid angle
 circular cone
 Fact 2.20.22, 161
 Fact 2.20.23, 161
 cone
 Fact 2.20.21, 161

solid set
 completely solid set
 Fact 10.8.9, 632
convex hull
 Fact 10.8.10, 632
convex set
 Fact 10.8.9, 632
definition, 622
dimension
 Fact 10.8.16, 633
image
 Fact 10.8.17, 633

solution
 Riccati equation
 Definition 12.16.12, 780

span
 affine subspace
 Fact 2.9.7, 111
 Fact 2.20.4, 154
 Fact 10.8.12, 633
constructive
 characterization
 Theorem 2.3.5, 91
 convex conical hull

- Fact 2.9.3, 110
- definition, 90**
- intersection**
 - Fact 2.9.12, 111
- union**
 - Fact 2.9.12, 111
- spanning path graph**
 - Fact 1.6.6, 14
- tournament**
 - Fact 1.6.6, 14
- spanning subgraph**
 - Definition 1.4.3, 9
- Specht**
 - reverse
 - arithmetic-mean–geometric-mean inequality
 - Fact 1.15.19, 52
- Specht's ratio**
 - matrix exponential
 - Fact 11.14.28, 687
 - power of a
 - positive-definite matrix
 - Fact 11.14.22, 685
 - Fact 11.14.23, 686
 - reverse
 - arithmetic-mean–geometric-mean inequality
 - Fact 1.15.19, 52
 - reverse Young inequality
 - Fact 1.10.22, 34
- special orthogonal group**
 - real symplectic group
 - Fact 3.22.5, 227
- spectral abscissa**
 - definition, 245
 - eigenvalue
 - Fact 5.11.24, 325
 - Hermitian matrix
 - Fact 5.11.5, 321
- Kronecker sum**
 - Fact 7.5.6, 409
- matrix exponential**
 - Fact 11.13.2, 677
 - Fact 11.15.8, 691
 - Fact 11.15.9, 691
 - Fact 11.18.8, 699
 - Fact 11.18.9, 699
- maximum eigenvalue**
 - Fact 5.11.5, 321
- maximum singular value**
 - Fact 5.11.26, 326
- minimum singular value**
 - Fact 5.11.26, 326
- outer-product matrix**
 - Fact 5.11.13, 323
- spectral radius**
 - Fact 4.10.4, 266
 - Fact 11.13.2, 677
- spectral decomposition**
 - normal matrix
 - Fact 5.14.14, 340
- spectral factorization**
 - definition, 232
 - Hamiltonian
 - Proposition 12.16.13, 780
 - polynomial roots
 - Proposition 4.1.1, 232
- spectral norm**
 - definition, 549
- spectral order**
 - positive-definite matrix
 - Fact 8.19.4, 523
 - positive-semidefinite matrix
 - Fact 8.19.4, 523
- spectral radius**
 - bound
 - Fact 4.10.22, 271
 - column norm
 - Corollary 9.4.10, 556
 - commuting matrices
 - Fact 5.12.11, 334
- convergent sequence**
 - Fact 4.10.5, 266
 - Fact 9.8.4, 572
- convexity for nonnegative matrices**
 - Fact 4.11.19, 280
- definition, 245**
- equi-induced norm**
 - Corollary 9.4.5, 554
- Frobenius norm**
 - Fact 9.13.12, 603
- Hermitian matrix**
 - Fact 5.11.5, 321
- induced norm**
 - Corollary 9.4.5, 554
 - Corollary 9.4.10, 556
- infinite series**
 - Fact 10.11.24, 641
- inverse matrix**
 - Proposition 9.4.13, 557
- Kronecker product**
 - Fact 7.4.14, 406
- lower bound**
 - Fact 9.13.12, 603
- matrix exponential**
 - Fact 11.13.2, 677
- matrix sum**
 - Fact 5.12.2, 333
 - Fact 5.12.3, 333
- maximum singular value**
 - Corollary 9.4.10, 556
 - Fact 5.11.5, 321
 - Fact 5.11.26, 326
 - Fact 8.18.25, 520
 - Fact 9.8.13, 573
 - Fact 9.13.9, 603
- minimum singular value**
 - Fact 5.11.26, 326
- monotonicity for nonnegative matrices**
 - Fact 4.11.18, 280
- nonnegative matrix**
 - Fact 4.11.6, 275
 - Fact 4.11.16, 279

- Fact 4.11.17, 280
- Fact 7.6.13, 415
- Fact 11.19.3, 706
- nonsingular matrix**
 - Fact 4.10.29, 272
- norm**
 - Proposition 9.2.6, 549
- normal matrix**
 - Fact 5.14.15, 341
- outer-product matrix**
 - Fact 5.11.13, 323
- perturbation**
 - Fact 9.14.6, 608
- positive matrix**
 - Fact 7.6.14, 415
- positive-definite matrix**
 - Fact 8.10.5, 456
 - Fact 8.18.25, 520
- positive-semidefinite matrix**
 - Fact 8.18.25, 520
 - Fact 8.20.8, 526
- row norm**
 - Corollary 9.4.10, 556
- Schur product**
 - Fact 7.6.13, 415
 - Fact 7.6.14, 415
 - Fact 7.6.16, 416
 - Fact 7.6.17, 416
 - Fact 9.14.33, 616
- spectral abscissa**
 - Fact 4.10.4, 266
 - Fact 11.13.2, 677
- submultiplicative norm**
 - Proposition 9.3.2, 550
 - Proposition 9.3.3, 550
 - Corollary 9.3.4, 550
 - Fact 9.8.4, 572
 - Fact 9.9.3, 580
- trace**
 - Fact 4.10.22, 271
 - Fact 5.11.46, 332
 - Fact 9.13.12, 603
- spectral radius of a product**
- Bourin**
 - Fact 8.18.25, 520
- spectral variation**
- Hermitian matrix**
 - Fact 9.12.5, 600
 - Fact 9.12.7, 601
- normal matrix**
 - Fact 9.12.5, 600
 - Fact 9.12.6, 600
- spectrum**
- adjugate**
 - Fact 4.10.7, 267
- asymptotic eigenvalue**
 - Fact 4.10.28, 272
- asymptotically stable matrix**
 - Fact 11.18.13, 700
- block-triangular matrix**
 - Proposition 5.5.13, 298
- bounds**
 - Fact 4.10.16, 269
 - Fact 4.10.20, 270
 - Fact 4.10.21, 271
- Cartesian decomposition**
 - Fact 5.11.21, 325
- circulant matrix**
 - Fact 5.16.7, 355
- commutator**
 - Fact 5.12.14, 335
- commuting matrices**
 - Fact 5.12.14, 335
- continuity**
 - Fact 10.11.8, 638
 - Fact 10.11.9, 639
- convex hull**
 - Fact 8.14.7, 496
 - Fact 8.14.8, 497
- cross-product matrix**
 - Fact 4.9.19, 263
- definition**
 - Definition 4.4.4, 240
- dissipative matrix**
 - Fact 8.13.31, 491
- doublet**
 - Fact 5.11.13, 323
- elementary matrix**
 - Proposition 5.5.21, 300
- elementary projector**
 - Proposition 5.5.21, 300
- elementary reflector**
 - Proposition 5.5.21, 300
- group-invertible matrix**
 - Proposition 5.5.21, 300
- Hamiltonian**
 - Theorem 12.17.9, 784
 - Proposition 12.16.13, 780
 - Proposition 12.17.5, 783
 - Proposition 12.17.7, 784
 - Proposition 12.17.8, 784
 - Lemma 12.17.4, 783
 - Lemma 12.17.6, 783
- Hamiltonian matrix**
 - Proposition 5.5.21, 300
- Hermitian matrix**
 - Proposition 5.5.21, 300
 - Lemma 8.4.8, 427
- idempotent matrix**
 - Proposition 5.5.21, 300
 - Fact 5.11.7, 322
- identity perturbation**
 - Fact 4.10.13, 268
 - Fact 4.10.14, 269
- inverse matrix**
 - Fact 5.11.14, 324
- involutory matrix**
 - Proposition 5.5.21, 300
- Laplacian matrix**
 - Fact 11.19.7, 708
- mass-spring system**
 - Fact 5.12.21, 337
- matrix exponential**
 - Proposition 11.2.3, 648
 - Corollary 11.2.6, 648
- matrix function**
 - Corollary 10.5.4, 629
- matrix logarithm**
 - Theorem 11.5.1, 656
- minimal polynomial**
 - Fact 4.10.8, 267
- nilpotent matrix**

- Proposition 5.5.21, 300
- normal matrix**
 - Fact 4.10.24, 271
 - Fact 8.14.7, 496
 - Fact 8.14.8, 497
- outer-product matrix**
 - Fact 5.11.13, 323
 - Fact 5.14.1, 338
- partitioned matrix**
 - Fact 2.19.3, 151
 - Fact 4.10.25, 271
 - Fact 4.10.26, 271
- permutation matrix**
 - Fact 5.16.8, 357
- perturbed matrix**
 - Fact 4.10.3, 266
- polynomial**
 - Fact 4.10.9, 267
 - Fact 4.10.10, 267
- positive matrix**
 - Fact 5.11.12, 323
- positive-definite matrix**
 - Proposition 5.5.21, 300
- positive-semidefinite matrix**
 - Proposition 5.5.21, 300
 - Fact 8.20.16, 527
- projector**
 - Proposition 5.5.21, 300
 - Fact 5.12.15, 335
 - Fact 5.12.16, 335
- properties**
 - Proposition 4.4.5, 241
- quadratic form**
 - Fact 8.14.7, 496
 - Fact 8.14.8, 497
- quadratic matrix equation**
 - Fact 5.11.3, 321
 - Fact 5.11.4, 321
- rational function**
 - Fact 5.11.15, 324
- reflector**
 - Proposition 5.5.21, 300
- reverse identity matrix**
 - Fact 5.9.24, 314
- shifted-unitary matrix**
 - Proposition 5.5.21, 300
- singular matrix**
 - Proposition 5.5.21, 300
- skew reflector**
 - Proposition 5.5.21, 300
- skew-Hermitian matrix**
 - Proposition 5.5.21, 300
- skew-involutory matrix**
 - Proposition 5.5.21, 300
- skew-symmetric matrix**
 - Fact 4.9.20, 263
 - Fact 4.10.2, 266
 - Fact 5.14.33, 343
- subspace decomposition**
 - Proposition 5.5.7, 296
- Sylvester's equation**
 - Corollary 7.2.5, 404
 - Fact 7.5.14, 410
- symplectic matrix**
 - Proposition 5.5.21, 300
- Toeplitz matrix**
 - Fact 4.10.15, 269
 - Fact 5.11.43, 331
 - Fact 5.11.44, 331
 - Fact 8.9.34, 454
- trace**
 - Fact 4.10.6, 267
- tridiagonal matrix**
 - Fact 5.11.40, 329
 - Fact 5.11.41, 329
 - Fact 5.11.42, 330
 - Fact 5.11.43, 331
 - Fact 5.11.44, 331
- tripotent matrix**
 - Proposition 5.5.21, 300
- unipotent matrix**
 - Proposition 5.5.21, 300
- unit imaginary matrix**
 - Fact 5.9.25, 315
- unitary matrix**
 - Proposition 5.5.21, 300
- spectrum bounds**
- Brauer**
 - Fact 4.10.21, 271
- ovals of Cassini**
 - Fact 4.10.21, 271
- spectrum of convex hull**
 - field of values**
 - Fact 8.14.7, 496
 - Fact 8.14.8, 497
 - numerical range**
 - Fact 8.14.7, 496
 - Fact 8.14.8, 497
- sphere of radius ε**
 - definition, 621**
- spin group**
 - double cover**
 - Fact 3.11.10, 192
- spread**
 - commutator**
 - Fact 9.9.30, 585
 - Fact 9.9.31, 585
 - Hermitian matrix**
 - Fact 8.15.31, 505
- square**
 - definition, 79**
 - trace**
 - Fact 8.17.7, 510
- square root**
 - 2×2 positive-semidefinite matrix**
 - Fact 8.9.6, 451
 - asymptotically stable matrix**
 - Fact 11.18.36, 705
 - commuting matrices**
 - Fact 5.18.1, 359
 - Fact 8.10.25, 458
 - convergent sequence**
 - Fact 5.15.21, 348
 - Fact 8.9.32, 454
 - definition, 431**
 - generalized inverse**
 - Fact 8.20.4, 525
 - group-invertible matrix**

- Fact 5.15.20, 348
- identity**
 - Fact 8.9.24, 453
 - Fact 8.9.25, 453
- Jordan form**
 - Fact 5.15.19, 348
- Kronecker product**
 - Fact 8.21.29, 536
 - Fact 8.21.30, 537
- matrix sign function**
 - Fact 5.15.21, 348
- maximum singular value**
 - Fact 8.18.14, 516
 - Fact 9.8.32, 576
 - Fact 9.14.15, 611
- Newton-Raphson algorithm**
 - Fact 5.15.21, 348
- normal matrix**
 - Fact 8.9.27, 453
 - Fact 8.9.28, 453
 - Fact 8.9.29, 453
- orthogonal matrix**
 - Fact 8.9.26, 453
- positive-semidefinite matrix**
 - Fact 8.10.18, 458
 - Fact 8.10.26, 458
 - Fact 8.21.29, 536
 - Fact 9.8.32, 576
- principal square root**
 - Theorem 10.6.1, 629
- projector**
 - Fact 8.10.25, 458
- range**
 - Fact 8.7.2, 443
- scalar inequality**
 - Fact 1.9.6, 24
 - Fact 1.12.1, 46
 - Fact 1.12.2, 46
- submultiplicative norm**
 - Fact 9.8.32, 576
- sum of squares**
 - Fact 2.18.8, 150
- unitarily invariant norm**
 - Fact 9.9.18, 583
- Fact 9.9.19, 583
- unitary matrix**
 - Fact 8.9.26, 453
- square-root function**
 - Niculescu's inequality
 - Fact 1.10.20, 33
- squares**
 - scalar inequality
 - Fact 1.11.21, 44
- stability**
 - mass-spring system
 - Fact 11.18.38, 705
 - partitioned matrix
 - Fact 11.18.38, 705
- stability radius**
 - asymptotically stable matrix
 - Fact 11.18.17, 700
- stabilizability**
 - asymptotically stable matrix
 - Proposition 11.9.5, 735
 - Proposition 12.8.3, 747
 - Proposition 12.8.5, 748
 - Corollary 12.8.6, 749
- block-triangular matrix**
 - Proposition 12.8.4, 747
- controllably asymptotically stable**
 - Proposition 12.8.3, 747
 - Proposition 12.8.5, 748
- definition**
 - Definition 12.8.1, 747
- full-state feedback**
 - Proposition 12.8.2, 747
- Hamiltonian**
 - Fact 12.23.1, 802
- input matrix**
 - Fact 12.20.15, 792
- Lyapunov equation**
 - Corollary 12.8.6, 749
- maximal solution of the Riccati equation**
 - Theorem 12.18.1, 787
- observably asymptotically stable**
 - Proposition 11.9.5, 735
- orthogonal matrix**
 - Proposition 12.8.4, 747
- positive-semidefinite matrix**
 - Fact 12.20.6, 791
- positive-semidefinite ordering**
 - Fact 12.20.8, 791
- range**
 - Fact 12.20.7, 791
- Riccati equation**
 - Theorem 12.17.9, 784
 - Theorem 12.18.1, 787
 - Corollary 12.19.2, 790
- shift**
 - Fact 12.20.11, 792
- stabilization**
 - controllability
 - Fact 12.20.17, 792
 - Gramian
 - Fact 12.20.17, 792
- stabilizing solution**
 - Hamiltonian
 - Corollary 12.16.15, 781
 - Riccati equation
 - Definition 12.16.12, 780
 - Theorem 12.17.2, 782
 - Theorem 12.17.9, 784
 - Theorem 12.18.4, 787
 - Proposition 12.17.1, 782
 - Proposition 12.18.3, 787
 - Proposition 12.19.4, 790
 - Corollary 12.16.15, 781
- stable subspace**
 - complementary subspaces
 - Proposition 11.8.8, 665
- group-invertible matrix**

- Proposition 11.8.8, 665
- idempotent matrix**
 - Proposition 11.8.8, 665
- invariant subspace**
 - Proposition 11.8.8, 665
- matrix exponential**
 - Proposition 11.8.8, 665
- minimal polynomial**
 - Proposition 11.8.5, 664
 - Fact 11.23.1, 716
 - Fact 11.23.2, 716
- standard control problem**
 - definition, 774
- standard nilpotent matrix**
 - definition, 166
- star partial ordering**
 - characterization
 - Fact 6.4.47, 385
 - commuting matrices
 - Fact 2.10.36, 120
 - definition
 - Fact 2.10.35, 120
 - Fact 8.19.7, 524
 - generalized inverse
 - Fact 8.19.8, 524
 - positive-semidefinite matrix
 - Fact 8.19.8, 524
 - Fact 8.19.9, 524
 - Fact 8.20.8, 526
- star-dagger matrix**
 - generalized inverse
 - Fact 6.3.13, 372
- state convergence**
 - detectability
 - Fact 12.20.2, 791
 - discrete-time
 - time-varying
 - system
 - Fact 11.21.16, 715
- state equation**
 - definition, 723
 - matrix exponential
 - Proposition 12.1.1, 723
- variation of constants formula**
 - Proposition 12.1.1, 723
- state transition matrix**
 - time-varying
 - dynamics
 - Fact 11.13.5, 678
- statement**
 - definition, 1
- Stein equation**
 - discrete-time
 - Lyapunov equation
 - Fact 11.21.15, 714
- step function, 724**
- step response**
 - definition, 725
 - Lyapunov-stable matrix
 - Fact 12.20.1, 790
- step-down matrix**
 - resultant
 - Fact 4.8.4, 254
- Stephanos**
 - eigenvector of a
 - Kronecker product
 - Fact 7.4.21, 406
- Stewart**
 - regular pencil
 - Fact 5.17.3, 358
- stiffness**
 - definition, 654
- stiffness matrix**
 - partitioned matrix
 - Fact 5.12.21, 337
- Stirling matrix**
 - Vandermonde matrix
 - Fact 5.16.3, 354
- Stirling's formula**
 - factorial
 - Fact 1.9.19, 26
- Storey**
- asymptotic stability of a tridiagonal matrix**
 - Fact 11.18.24, 702
- Stormer**
 - Schatten norm for
 - positive-semidefinite matrices
 - Fact 9.9.22, 583
- strengthening**
 - definition, 2
- strictly concave function**
 - definition
 - Definition 8.6.14, 436
- strictly convex function**
 - definition
 - Definition 8.6.14, 436
- positive-definite matrix**
 - Fact 8.14.15, 499
 - Fact 8.14.16, 499
- trace**
 - Fact 8.14.16, 499
- transformation**
 - Fact 1.8.2, 21
- strictly dissipative matrix**
 - dissipative matrix
 - Fact 8.9.31, 453
- strictly lower triangular matrix**
 - definition
 - Definition 3.1.3, 167
- matrix power**
 - Fact 3.18.7, 216
- matrix product**
 - Fact 3.20.18, 221
- strictly proper rational function**
 - definition
 - Definition 4.7.1, 249
- strictly proper rational transfer function**

- definition
 - Definition 4.7.2, 249
- strictly upper triangular matrix**
 - definition
 - Definition 3.1.3, 167
- Lie algebra**
 - Fact 3.21.4, 222
 - Fact 11.22.1, 715
- matrix power**
 - Fact 3.18.7, 216
- matrix product**
 - Fact 3.20.18, 221
- strong Kronecker product**
 - Kronecker product, 416
- strong log majorization**
 - convex function
 - Fact 2.21.9, 163
 - definition
 - Definition 2.1.1, 78
 - matrix exponential
 - Fact 11.16.4, 692
 - singular value inequality
 - Fact 9.13.19, 605
- strong majorization**
 - convex function
 - Fact 2.21.8, 163
 - Fact 2.21.11, 163
 - convex hull
 - Fact 2.21.7, 163
 - definition
 - Definition 2.1.1, 78
 - diagonal entry
 - Fact 8.17.8, 510
 - doubly stochastic matrix
 - Fact 2.21.7, 163
 - eigenvalue
 - Corollary 8.6.19, 442
 - Fact 8.18.4, 513
 - Fact 8.18.29, 521
 - entropy
 - Fact 2.21.6, 162
 - Hermitian matrix
 - Fact 8.17.8, 510
 - Muirhead's theorem
 - Fact 2.21.5, 162
 - ones vector
 - Fact 2.21.1, 162
 - Schur concave function
 - Fact 2.21.6, 162
 - Schur convex function
 - Fact 2.21.4, 162
 - Fact 2.21.5, 162
 - strongly decreasing function**
 - definition
 - Definition 8.6.12, 434
 - strongly increasing function**
 - definition
 - Definition 8.6.12, 434
 - determinant**
 - Proposition 8.6.13, 435
 - matrix functions**
 - Proposition 8.6.13, 435
 - structured matrix**
 - positive-semidefinite matrix
 - Fact 8.8.2, 445
 - Fact 8.8.3, 446
 - Fact 8.8.4, 446
 - Fact 8.8.5, 447
 - Fact 8.8.6, 447
 - Fact 8.8.7, 447
 - Fact 8.8.8, 447
 - Fact 8.8.9, 448
 - Fact 8.8.10, 448
 - Fact 8.8.11, 448
 - Fact 8.8.12, 448
 - Styan**
 - difference of idempotent matrices
 - Fact 5.12.19, 337
 - Hermitian matrix**
 - inertia identity
 - Fact 8.10.15, 457
 - rank of a tripotent matrix
 - Fact 2.10.23, 118
 - rank of an idempotent matrix
 - Fact 3.12.27, 203
 - SU(2)**
 - quaternions
 - Fact 3.22.6, 227
 - subdeterminant**
 - asymptotically stable matrix
 - Fact 11.19.1, 707
 - asymptotically stable polynomial
 - Fact 11.18.23, 702
 - definition, 105
 - inverse
 - Fact 2.13.5, 129
 - Lyapunov-stable polynomial**
 - Fact 11.18.23, 702
 - positive-definite matrix**
 - Proposition 8.2.8, 422
 - Fact 8.13.17, 488
 - positive-semidefinite matrix**
 - Proposition 8.2.7, 421
 - subdiagonal**
 - definition, 80
 - subdifferential**
 - convex function
 - Fact 10.11.14, 639
 - subgraph**
 - Definition 1.4.3, 9
 - sublevel set**
 - convex set
 - Fact 8.14.1, 494
 - submatrix**
 - complementary
 - Fact 2.11.20, 125
 - defect**
 - Fact 2.11.20, 125
 - definition, 80**
 - determinant**

- Fact 2.14.1, 132
- Hermitian matrix**
 - Theorem 8.4.5, 426
 - Corollary 8.4.6, 426
 - Lemma 8.4.4, 425
 - Fact 5.8.8, 308
- inertia**
 - Fact 5.8.8, 308
- Kronecker product**
 - Proposition 7.3.1, 404
- M-matrix**
 - Fact 4.11.7, 276
- positive-definite matrix**
 - Proposition 8.2.8, 422
- positive-semidefinite matrix**
 - Proposition 8.2.7, 421
 - Fact 8.7.7, 444
 - Fact 8.13.36, 492
- rank**
 - Proposition 4.3.5, 237
 - Proposition 4.7.7, 250
 - Fact 2.11.6, 121
 - Fact 2.11.17, 124
 - Fact 2.11.20, 125
 - Fact 2.11.21, 125
 - Fact 2.11.22, 125
 - Fact 3.20.5, 218
- singular value**
 - Fact 9.14.10, 609
- Smith form**
 - Proposition 4.3.5, 237
- Smith-McMillan form**
 - Proposition 4.7.7, 250
- Z-matrix**
 - Fact 4.11.7, 276
- submultiplicative norm**
 - commutator**
 - Fact 9.9.8, 580
 - compatible norms**
 - Proposition 9.3.1, 550
 - equi-induced norm**
 - Corollary 9.4.4, 554
 - Fact 9.8.45, 579
 - H_2 norm**
 - Fact 12.22.20, 801
 - Hölder norm**
 - Fact 9.9.20, 583
- idempotent matrix**
 - Fact 9.8.6, 572
- infinity norm**
 - Fact 9.9.1, 580
 - Fact 9.9.2, 580
- matrix exponential**
 - Proposition 11.1.2, 644
 - Fact 11.15.8, 691
 - Fact 11.15.9, 691
 - Fact 11.16.7, 694
 - Fact 11.18.8, 699
 - Fact 11.18.9, 699
- matrix norm**
 - Fact 9.9.4, 580
- nonsingular matrix**
 - Fact 9.8.5, 572
- positive-semidefinite matrix**
 - Fact 9.9.7, 580
- Schur product**
 - Fact 9.8.41, 578
- spectral radius**
 - Proposition 9.3.2, 550
 - Proposition 9.3.3, 550
 - Corollary 9.3.4, 550
 - Fact 9.8.4, 572
 - Fact 9.9.3, 580
- square root**
 - Fact 9.8.32, 576
- unitarily invariant norm**
 - Fact 9.8.41, 578
 - Fact 9.9.7, 580
- submultiplicative norms**
 - definition, 550**
- subset**
 - closure**
 - Fact 10.9.1, 634
 - definition, 2**
 - interior**
 - Fact 10.9.1, 634
- subset operation**
 - induced partial ordering**
 - Fact 1.5.17, 13
 - transitivity**
 - Fact 1.5.17, 13
- subspace**
 - affine**
 - definition, 89
 - affine subspace**
 - Fact 2.9.8, 111
 - closed set**
 - Fact 10.8.21, 633
 - common eigenvector**
 - Fact 5.14.27, 342
 - complementary**
 - Fact 2.9.18, 112
 - Fact 2.9.23, 113
 - complex conjugate transpose**
 - Fact 2.9.28, 114
 - definition, 89**
 - dimension**
 - Fact 2.9.20, 112
 - Fact 2.9.21, 113
 - Fact 2.9.22, 113
 - dimension inequality**
 - Fact 2.10.4, 115
 - gap topology**
 - Fact 10.9.18, 636
 - image under linear mapping**
 - Fact 2.9.26, 113
 - inclusion**
 - Fact 2.9.11, 111
 - Fact 2.9.14, 112
 - Fact 2.9.28, 114
 - inclusion and dimension ordering**
 - Lemma 2.3.4, 91
 - inner product**
 - Fact 10.9.12, 635
 - intersection**
 - Fact 2.9.9, 111
 - Fact 2.9.16, 112
 - Fact 2.9.17, 112
 - Fact 2.9.29, 114
 - Fact 2.9.30, 114
 - left inverse**
 - Fact 2.9.26, 113
 - minimal principal angle**
 - Fact 5.11.39, 329
 - Fact 5.12.17, 335

Fact 10.9.18, 636

orthogonal

complement

Proposition 3.5.2, 175

Fact 2.9.14, 112

Fact 2.9.16, 112

Fact 2.9.18, 112

Fact 2.9.27, 114

orthogonal matrix

Fact 3.11.1, 189

Fact 3.11.2, 189

principal angle

Fact 2.9.19, 112

projector

Proposition 3.5.1, 175

Proposition 3.5.2, 175

Fact 9.8.3, 571

Fact 10.9.17, 636

quadratic form

Fact 8.15.27, 504

Fact 8.15.28, 504

range

Proposition 3.5.1, 175

Fact 2.9.24, 113

span

Fact 2.9.13, 111

span of image

Fact 2.9.26, 113

sum

Fact 2.9.9, 111

Fact 2.9.13, 111

Fact 2.9.16, 112

Fact 2.9.17, 112

Fact 2.9.29, 114

Fact 2.9.30, 114

union

Fact 2.9.11, 111

Fact 2.9.13, 111

unitary matrix

Fact 3.11.1, 189

Fact 3.11.2, 189

subspace

decomposition

spectrum

Proposition 5.5.7, 296

subspace dimension

theorem

dimension

Theorem 2.3.1, 90

rank

Fact 2.11.9, 122

Fact 2.11.10, 122

subspace intersection

inverse image

Fact 2.9.30, 114

subspace sum

inverse image

Fact 2.9.30, 114

sufficiency

definition, 1

sum

Drazin generalized

inverse

Fact 6.6.5, 394

eigenvalue

Fact 5.12.2, 333

Fact 5.12.3, 333

generalized inverse

Fact 6.4.34, 383

Fact 6.4.35, 383

Fact 6.4.36, 383

Hamiltonian matrix

Fact 3.19.5, 216

outer-product

matrix

Fact 2.10.24, 118

projector

Fact 5.12.17, 335

singular value

Fact 9.14.20, 612

Fact 9.14.21, 612

Fact 9.14.25, 613

spectral radius

Fact 5.12.2, 333

Fact 5.12.3, 333

sum inequality

power inequality

Fact 1.16.28, 66

Fact 1.16.29, 66

sum of integer powers

inequality

Fact 1.9.31, 30

matrix exponential

Fact 11.11.4, 672

sum of matrices

determinant

Fact 5.12.12, 335

Fact 9.14.18, 611

idempotent matrix

Fact 3.12.22, 201

Fact 3.12.26, 203

Fact 5.19.7, 361

Fact 5.19.8, 361

Fact 5.19.9, 361

inverse matrix

Corollary 2.8.10, 110

Kronecker product

Proposition 7.1.4, 400

nilpotent matrix

Fact 3.17.10, 214

projector

Fact 3.13.23, 210

Fact 5.19.4, 360

sum of orthogonal

matrices

determinant

Fact 3.11.22, 196

sum of powers

Carlson inequality

Fact 1.15.41, 58

Copson inequality

Fact 1.15.43, 59

Hardy inequality

Fact 1.15.42, 58

sum of products

Hardy-Hilbert

inequality

Fact 1.16.13, 63

Fact 1.16.14, 63

inequality

Fact 1.15.20, 53

sum of products

inequality

Hardy-Littlewood

rearrangement

inequality

Fact 1.16.4, 60

Fact 1.16.5, 60

sum of sets

convex set

Fact 2.9.1, 110

- Fact 2.9.2, 110
- Fact 10.9.4, 634
- Fact 10.9.5, 634
- Fact 10.9.7, 635
- dual cone**
 - Fact 2.9.5, 111
- sum of squares**
 - square root**
 - Fact 2.18.8, 150
- sum of subspaces**
 - subspace dimension theorem**
 - Theorem 2.3.1, 90
- sum of transfer functions**
 - H_2 norm
 - Proposition 12.11.6, 767
- sum-of-squares inequality**
 - square-of-sum inequality**
 - Fact 1.15.14, 48
- summation**
 - identity**
 - Fact 1.7.3, 17
 - Fact 1.7.4, 18
 - Fact 1.7.5, 18
- superdiagonal**
 - definition, 80**
- supermultiplicativity**
 - induced lower bound**
 - Proposition 9.5.6, 560
- support of a relation**
 - definition**
 - Definition 1.3.4, 5
- surjective function**
 - definition, 76**
- Sylvester matrix**
 - coprime polynomials**
 - Fact 4.8.4, 254
- Sylvester's equation**
 - controllability**
 - Fact 12.21.14, 796
- controllability matrix**
 - Fact 12.21.13, 796
- linear matrix equation**
 - Proposition 7.2.4, 403
 - Proposition 11.9.3, 667
 - Fact 5.10.20, 320
 - Fact 5.10.21, 320
 - Fact 6.5.7, 387
- nonsingular matrix**
 - Fact 12.21.14, 796
- observability**
 - Fact 12.21.14, 796
- observability matrix**
 - Fact 12.21.13, 796
- partitioned matrix**
 - Fact 5.10.20, 320
 - Fact 5.10.21, 320
 - Fact 6.5.7, 387
- rank**
 - Fact 12.21.13, 796
- similar matrices**
 - Corollary 7.2.5, 404
 - Fact 7.5.14, 410
- spectrum**
 - Corollary 7.2.5, 404
 - Fact 7.5.14, 410
- Sylvester's identity**
 - determinant**
 - Fact 2.14.1, 132
- Sylvester's inequality**
 - rank of a product, 97**
- Sylvester's law of inertia**
 - definition, 294**
 - Ostrowski**
 - Fact 5.8.17, 310
- Sylvester's law of nullity**
 - defect**
 - Fact 2.10.15, 117
- symmetric cone**
 - induced by symmetric relation**
 - Proposition 2.3.6, 93
- symmetric gauge function**
 - unitarily invariant norm**
 - Fact 9.8.42, 579
 - weak majorization**
 - Fact 2.21.14, 164
- symmetric graph**
 - adjacency matrix**
 - Fact 4.11.1, 272
 - cycle**
 - Fact 1.6.5, 14
 - degree matrix**
 - Fact 4.11.1, 272
 - forest**
 - Fact 1.6.5, 14
 - Laplacian**
 - Fact 4.11.1, 272
 - Laplacian matrix**
 - Fact 8.15.36, 506
- symmetric hull**
 - definition**
 - Definition 1.3.4, 5
 - relation**
 - Proposition 1.3.5, 6
- symmetric matrix**
 - congruent matrices**
 - Fact 5.9.16, 313
 - definition**
 - Definition 3.1.1, 165
 - eigenvalue**
 - Fact 4.10.1, 265
 - factorization**
 - Corollary 5.3.9, 292
 - Fact 5.15.24, 349
 - Hankel matrix**
 - Fact 3.18.2, 215
 - Hermitian matrix**
 - Fact 3.7.9, 179
 - involutory matrix**
 - Fact 5.15.36, 351
 - linear matrix equation**
 - Fact 3.7.3, 178
 - matrix power**
 - Fact 3.7.4, 178
 - matrix transpose**
 - Fact 3.7.2, 178

- maximum eigenvalue
 - Fact 5.12.20, 337
- minimum eigenvalue
 - Fact 5.12.20, 337
- orthogonally similar matrices
 - Fact 5.9.15, 313
- partitioned matrix
 - Fact 3.11.27, 196
- quadratic form
 - Fact 3.7.5, 178
- similar matrices
 - Fact 5.15.39, 351
- similarity transformation
 - Fact 5.15.2, 345
 - Fact 5.15.3, 345
- skew-Hermitian matrix
 - Fact 3.7.9, 179
- skew-symmetric matrix
 - Fact 5.9.16, 313
 - Fact 5.15.39, 351
- trace
 - Fact 5.12.8, 334
- symmetric relation**
 - definition
 - Definition 1.3.2, 5
 - graph
 - Definition 1.4.1, 8
 - intersection
 - Proposition 1.3.3, 5
 - symmetric cone induced by
 - Proposition 2.3.6, 93
- symmetric set**
 - definition, 89
- symmetry group**
 - group
 - Fact 3.21.7, 223
- symplectic group**
 - determinant
 - Fact 3.19.11, 217
 - quaternion group
 - Fact 3.22.4, 227
 - special orthogonal group
 - Fact 3.22.5, 227
 - unitary group
 - Fact 3.21.3, 222
- symplectic matrix**
 - Cayley transform
 - Fact 3.19.12, 217
 - definition
 - Definition 3.1.5, 169
 - determinant
 - Fact 3.19.10, 217
 - Fact 3.19.11, 217
 - group
 - Proposition 3.3.6, 172
 - Hamiltonian matrix
 - Fact 3.19.2, 216
 - Fact 3.19.12, 217
 - Fact 3.19.13, 217
 - identity
 - Fact 3.19.1, 216
 - identity matrix
 - Fact 3.19.3, 216
 - matrix exponential
 - Proposition 11.6.7, 659
 - matrix logarithm
 - Fact 11.14.19, 685
 - partitioned matrix
 - Fact 3.19.9, 217
 - reverse identity matrix
 - Fact 3.19.3, 216
 - skew-involutory matrix
 - Fact 3.19.2, 216
 - spectrum
 - Proposition 5.5.21, 300
 - unit imaginary matrix
 - Fact 3.19.3, 216
- symplectic similarity**
 - Hamiltonian matrix
 - Fact 3.19.4, 216
- Szasz's inequality**
 - positive-semidefinite matrix
 - Fact 8.13.36, 492

T

- T-congruence**
 - complex-symmetric matrix
 - Fact 5.9.22, 314
- T-congruent diagonalization**
 - complex-symmetric matrix
 - Fact 5.9.22, 314
- T-congruent matrices**
 - definition
 - Definition 3.4.4, 174
- Tao**
 - Hölder-induced norm
 - Fact 9.8.19, 575
- Taussky-Todd factorization**
 - Fact 5.15.7, 346
- tautology**
 - definition, 1
- tetrahedral group**
 - group
 - Fact 3.21.7, 223
- tetrahedron**
 - volume
 - Fact 2.20.15, 160
- theorem**
 - definition, 1
- thermodynamic inequality**
 - matrix exponential
 - Fact 11.14.31, 688
 - relative entropy
 - Fact 11.14.25, 686
- Tian**
 - idempotent matrix and similar matrices
 - Fact 5.10.22, 320
 - range of a partitioned matrix

Fact 6.5.3, 386

**Tikhonov inverse
positive-definite
matrix**

Fact 8.9.40, 455

time-varying dynamics

commuting matrices

Fact 11.13.4, 678

determinant

Fact 11.13.4, 678

**matrix differential
equation**

Fact 11.13.4, 678

Fact 11.13.5, 678

**state transition
matrix**

Fact 11.13.5, 678

trace

Fact 11.13.4, 678

Toeplitz matrix

**block-Toeplitz
matrix**

Fact 3.18.3, 215

definition

Definition 3.1.3, 167

determinant

Fact 2.13.13, 131

Fact 3.20.7, 219

Hankel matrix

Fact 3.18.1, 215

**lower triangular
matrix**

Fact 3.18.7, 216

Fact 11.13.1, 677

nilpotent matrix

Fact 3.18.6, 216

polynomial

multiplication

Fact 4.8.10, 258

**positive-definite
matrix**

Fact 8.13.13, 487

**reverse-symmetric
matrix**

Fact 3.18.5, 215

spectrum

Fact 4.10.15, 269

Fact 5.11.43, 331

Fact 5.11.44, 331

Fact 8.9.34, 454

tridiagonal matrix

Fact 3.20.7, 219

Fact 5.11.43, 331

Fact 5.11.44, 331

**upper triangular
matrix**

Fact 3.18.7, 216

Fact 11.13.1, 677

Tomiyama

**maximum singular
value of a
partitioned matrix**

Fact 9.14.12, 610

total ordering

definition

Definition 1.3.9, 7

dictionary ordering

Fact 1.5.8, 12

lexicographic

ordering

Fact 1.5.8, 12

planar example

Fact 1.5.8, 12

total response, 725

**totally nonnegative
matrix**

definition

Fact 11.18.23, 702

totally positive matrix

rank

Fact 8.7.7, 444

tournament

graph

Fact 1.6.6, 14

Hamiltonian cycle

Fact 1.6.6, 14

spanning path

Fact 1.6.6, 14

trace

2×2 matrices

Fact 2.12.9, 126

2×2 matrix identity

Fact 4.9.3, 260

Fact 4.9.4, 260

3×3 matrix identity

Fact 4.9.5, 261

Fact 4.9.6, 261

adjugate

Fact 4.9.8, 261

**asymptotically stable
matrix**

Fact 11.18.31, 704

commutator

Fact 2.18.1, 149

Fact 2.18.2, 149

Fact 5.9.18, 313

**complex conjugate
transpose**

Fact 8.12.4, 476

Fact 8.12.5, 476

Fact 9.13.16, 604

convex function

Proposition 8.6.17, 437

Fact 8.14.17, 499

definition, 86

derivative

Proposition 10.7.4, 631

Fact 11.14.3, 682

determinant

Proposition 8.4.14, 429

Corollary 11.2.4, 648

Corollary 11.2.5, 648

Fact 2.13.16, 132

Fact 8.12.1, 475

Fact 8.13.20, 488

Fact 11.14.20, 685

dimension

Fact 2.18.11, 150

eigenvalue

Proposition 8.4.13, 428

Fact 5.11.11, 322

Fact 8.17.5, 509

Fact 8.18.18, 518

eigenvalue bound

Fact 5.11.45, 331

elementary projector

Fact 5.8.11, 309

elementary reflector

Fact 5.8.11, 309

Frobenius norm

Fact 9.11.3, 597

Fact 9.11.4, 598

Fact 9.11.5, 598

- Fact 9.12.2, 599
- generalized inverse**
 - Fact 6.3.22, 374
- group generalized inverse**
 - Fact 6.6.6, 394
- Hamiltonian matrix**
 - Fact 3.19.7, 216
- Hermitian matrix**
 - Proposition 8.4.13, 428
 - Corollary 8.4.10, 427
 - Lemma 8.4.12, 428
 - Fact 3.7.13, 180
 - Fact 3.7.22, 182
 - Fact 8.12.38, 483
- Hermitian matrix product**
 - Fact 5.12.4, 333
 - Fact 5.12.5, 333
 - Fact 8.12.6, 476
 - Fact 8.12.7, 477
 - Fact 8.12.8, 477
 - Fact 8.12.16, 478
 - Fact 8.18.18, 518
- idempotent matrix**
 - Fact 5.8.1, 307
 - Fact 5.11.7, 322
- identities, 86**
- inequality**
 - Fact 5.12.9, 334
- involutory matrix**
 - Fact 5.8.2, 307
- Klein's inequality**
 - Fact 11.14.25, 686
- Kronecker**
 - permutation matrix**
 - Fact 7.4.29, 407
- Kronecker product**
 - Proposition 7.1.12, 402
 - Fact 11.14.38, 688
- Kronecker sum**
 - Fact 11.14.36, 688
- matrix exponential**
 - Corollary 11.2.4, 648
 - Corollary 11.2.5, 648
 - Fact 8.14.18, 500
 - Fact 11.11.6, 673
 - Fact 11.14.3, 682
 - Fact 11.14.10, 683
- Fact 11.14.28, 687
- Fact 11.14.29, 687
- Fact 11.14.30, 687
- Fact 11.14.31, 688
- Fact 11.14.36, 688
- Fact 11.14.38, 688
- Fact 11.15.4, 689
- Fact 11.15.5, 689
- Fact 11.16.1, 692
- Fact 11.16.4, 692
- matrix logarithm**
 - Fact 11.14.24, 686
 - Fact 11.14.25, 686
 - Fact 11.14.27, 686
 - Fact 11.14.31, 688
- matrix power**
 - Fact 2.12.13, 127
 - Fact 2.12.17, 127
 - Fact 4.10.22, 271
 - Fact 4.11.22, 281
 - Fact 5.11.9, 322
 - Fact 5.11.10, 322
 - Fact 8.12.4, 476
 - Fact 8.12.5, 476
- matrix product**
 - Fact 2.12.16, 127
 - Fact 5.12.6, 334
 - Fact 5.12.7, 334
 - Fact 8.12.14, 478
 - Fact 8.12.15, 478
 - Fact 9.14.3, 607
 - Fact 9.14.4, 608
- matrix squared**
 - Fact 5.11.9, 322
 - Fact 5.11.10, 322
- maximum singular value**
 - Fact 5.12.7, 334
 - Fact 9.14.4, 608
- maximum singular value bound**
 - Fact 9.13.13, 604
- nilpotent matrix**
 - Fact 3.17.6, 214
- nonnegative matrix**
 - Fact 4.11.22, 281
- normal matrix**
 - Fact 3.7.12, 180
 - Fact 8.12.5, 476
- normal matrix product**
 - Fact 5.12.4, 333
- orthogonal matrix**
 - Fact 3.11.17, 195
 - Fact 3.11.18, 195
 - Fact 5.12.9, 334
 - Fact 5.12.10, 334
- outer-product matrix**
 - Fact 5.14.3, 338
- partitioned matrix**
 - Fact 8.12.36, 483
 - Fact 8.12.39, 484
 - Fact 8.12.40, 484
 - Fact 8.12.41, 484
 - Fact 8.12.42, 484
- polarized**
 - Cayley-Hamilton theorem**
 - Fact 4.9.3, 260
- positive-definite matrix**
 - Proposition 8.4.14, 429
 - Fact 8.9.16, 452
 - Fact 8.10.46, 464
 - Fact 8.11.10, 469
 - Fact 8.12.1, 475
 - Fact 8.12.2, 475
 - Fact 8.12.24, 480
 - Fact 8.12.27, 481
 - Fact 8.12.37, 483
 - Fact 8.13.12, 487
 - Fact 11.14.24, 686
 - Fact 11.14.25, 686
 - Fact 11.14.27, 686
- positive-semidefinite matrix**
 - Proposition 8.4.13, 428
 - Fact 8.12.3, 476
 - Fact 8.12.9, 477
 - Fact 8.12.10, 477
 - Fact 8.12.11, 477
 - Fact 8.12.12, 477
 - Fact 8.12.13, 477
 - Fact 8.12.17, 478
 - Fact 8.12.18, 478
 - Fact 8.12.19, 479
 - Fact 8.12.20, 479

- Fact 8.12.21, 480
- Fact 8.12.22, 480
- Fact 8.12.23, 480
- Fact 8.12.24, 480
- Fact 8.12.26, 481
- Fact 8.12.28, 481
- Fact 8.12.29, 481
- Fact 8.12.34, 483
- Fact 8.12.35, 483
- Fact 8.12.36, 483
- Fact 8.12.38, 483
- Fact 8.12.39, 484
- Fact 8.12.40, 484
- Fact 8.12.41, 484
- Fact 8.13.20, 488
- Fact 8.18.16, 517
- Fact 8.18.20, 518
- Fact 8.20.3, 525
- Fact 8.20.17, 528
- projector**
 - Fact 5.8.11, 309
- quadruple product**
 - Fact 7.4.9, 405
- rank**
 - Fact 5.11.10, 322
 - Fact 9.11.4, 598
- reflector**
 - Fact 5.8.11, 309
- rotation matrix**
 - Fact 3.11.17, 195
- Schatten norm**
 - Fact 9.12.1, 599
- Schur product**
 - Fact 8.21.17, 534
 - Fact 9.14.32, 615
- singular value**
 - Fact 5.12.6, 334
 - Fact 8.17.2, 508
 - Fact 9.12.1, 599
 - Fact 9.13.16, 604
 - Fact 9.14.3, 607
 - Fact 9.14.32, 615
- skew-Hermitian matrix**
 - Fact 3.7.24, 182
- skew-Hermitian matrix product**
 - Fact 8.12.6, 476
- skew-symmetric matrix**
 - Fact 3.7.23, 182
 - Fact 3.7.31, 183
- spectral radius**
 - Fact 4.10.22, 271
 - Fact 5.11.46, 332
 - Fact 9.13.12, 603
- spectrum**
 - Fact 4.10.6, 267
- square**
 - Fact 8.17.7, 510
- strictly convex function**
 - Fact 8.14.16, 499
- symmetric matrix**
 - Fact 5.12.8, 334
- time-varying dynamics**
 - Fact 11.13.4, 678
- trace norm**
 - Fact 9.11.2, 597
- triple product**
 - Fact 2.12.11, 127
 - Fact 7.4.7, 405
- tripotent matrix**
 - Fact 3.16.4, 213
 - Fact 5.8.3, 307
- unitarily similar matrices**
 - Fact 5.10.8, 318
- unitary matrix**
 - Fact 3.11.16, 194
 - Fact 3.11.32, 198
- vec**
 - Proposition 7.1.1, 399
 - Fact 7.4.7, 405
 - Fact 7.4.9, 405
- zero matrix**
 - Fact 2.12.14, 127
 - Fact 2.12.15, 127
- trace and singular value**
 - von Neumann's trace inequality
 - Fact 9.12.1, 599
- trace norm**
 - compatible norms
- Corollary 9.3.8, 552
- definition, 549**
- Frobenius norm**
 - Fact 9.9.11, 581
- matrix difference**
 - Fact 9.9.24, 584
- maximum singular value**
 - Corollary 9.3.8, 552
- positive-semidefinite matrix**
 - Fact 9.9.15, 582
- trace**
 - Fact 9.11.2, 597
- trace of a convex function**
 - Berezin
 - Fact 8.12.33, 482
 - Brown
 - Fact 8.12.33, 482
 - Hansen
 - Fact 8.12.33, 482
 - Kosaki
 - Fact 8.12.33, 482
 - Pedersen
 - Fact 8.12.33, 482
- trace of a Hermitian matrix product**
 - Fan
 - Fact 5.12.4, 333
- trace of a product**
 - Fan
 - Fact 5.12.10, 334
- traceable graph**
 - definition
 - Definition 1.4.3, 9
- Tracy-Singh product**
 - Kronecker product, 416
- trail**
 - definition
 - Definition 1.4.3, 9
- transfer function**
 - cascade
 - interconnection

Proposition 12.13.2,
770

derivative

Fact 12.22.6, 799

feedback

interconnection

Fact 12.22.8, 799

frequency response

Fact 12.22.5, 799

H_2 norm

Fact 12.22.16, 801

Fact 12.22.17, 801

Fact 12.22.18, 801

Fact 12.22.19, 801

imaginary part

Fact 12.22.5, 799

Jordan form

Fact 12.22.10, 800

parallel

interconnection

Proposition 12.13.2,
770

partitioned transfer function

Fact 12.22.7, 799

real part

Fact 12.22.5, 799

realization

Fact 12.22.3, 798

Fact 12.22.4, 799

Fact 12.22.6, 799

Fact 12.22.7, 799

Fact 12.22.8, 799

realization of inverse

Proposition 12.13.1,
770

realization of parahermitian conjugate

Proposition 12.13.1,
770

realization of

transpose

Proposition 12.13.1,
770

reciprocal argument

Fact 12.22.4, 799

right inverse

Fact 12.22.9, 799

shifted argument

Fact 12.22.3, 798

transitive hull

definition

Definition 1.3.4, 5

relation

Proposition 1.3.5, 6

transitive relation

convex cone induced by

Proposition 2.3.6, 93

definition

Definition 1.3.2, 5

graph

Definition 1.4.1, 8

intersection

Proposition 1.3.3, 5

positive-semidefinite matrix

Proposition 8.1.1, 417

transmission zero

definition

Definition 4.7.10, 251

Definition 4.7.13, 252

invariant zero

Theorem 12.10.8, 762

Theorem 12.10.9, 762

null space

Fact 4.8.16, 260

rank

Proposition 4.7.12, 251

transpose

controllability

Fact 12.20.16, 792

diagonalizable matrix

Fact 5.14.5, 339

involutory matrix

Fact 5.9.7, 312

Kronecker

permutation matrix

Proposition 7.1.13, 402

Kronecker product

Proposition 7.1.3, 400

matrix exponential

Proposition 11.2.8, 649

normal matrix

Fact 5.9.9, 312

Fact 5.9.10, 312

similar matrices

Proposition 5.5.12, 297

Corollary 4.3.11, 239

Corollary 5.3.8, 291

Corollary 5.5.22, 301

Fact 5.9.9, 312

Fact 5.9.10, 312

transpose of a matrix definition, 86

transpose of a vector definition, 84

transposition matrix permutation matrix Fact 3.21.6, 222

triangle

area

Fact 2.20.7, 155

Fact 2.20.8, 156

Fact 2.20.10, 156

Bandila's inequality

Fact 2.20.11, 156

cosine rule

Fact 2.20.11, 156

Euler's inequality

Fact 2.20.11, 156

fundamental triangle inequality

Fact 2.20.11, 156

Heron's formula

Fact 2.20.11, 156

inequality

Fact 1.11.17, 43

Klamkin's inequality

Fact 2.20.11, 156

Mircea's inequality

Fact 2.20.11, 156

semiperimeter

Fact 2.20.11, 156

sine rule

Fact 2.20.11, 156

triangle inequality

Blundon

Fact 2.20.11, 156

definition

Definition 9.1.1, 543

equality

Fact 9.7.3, 563

Frobenius norm

Fact 9.9.13, 582

linear dependence

Fact 9.7.3, 563

positive-semidefinite matrix

Fact 9.9.21, 583

reverse Hölder norm inequality

Fact 9.7.19, 569

Satnoianu

Fact 2.20.11, 156

triangular matrix**nilpotent matrix**

Fact 5.17.6, 358

triangularization**commutator**

Fact 5.17.5, 358

commuting matrices

Fact 5.17.4, 358

tridiagonal matrix**asymptotically stable matrix**

Fact 11.18.24, 702

Fact 11.18.25, 702

Fact 11.18.26, 702

Fact 11.18.27, 703

Fact 11.18.28, 703

cyclic matrix

Fact 11.18.25, 702

definition

Definition 3.1.3, 167

determinant

Fact 3.20.6, 218

Fact 3.20.7, 219

Fact 3.20.8, 219

Fact 3.20.9, 219

Fact 3.20.11, 220

inverse matrix

Fact 3.20.9, 219

Fact 3.20.10, 219

Fact 3.20.11, 220

positive-definite matrix

Fact 8.8.18, 450

Routh form

Fact 11.18.27, 703

Schwarz form

Fact 11.18.25, 702

Fact 11.18.26, 702

spectrum

Fact 5.11.40, 329

Fact 5.11.41, 329

Fact 5.11.42, 330

Fact 5.11.43, 331

Fact 5.11.44, 331

Toeplitz matrix

Fact 3.20.7, 219

Fact 5.11.43, 331

Fact 5.11.44, 331

trigonometric identities

Fact 1.19.1, 74

trigonometric inequality**Huygens's inequality**

Fact 1.9.29, 28

Jordan's inequality

Fact 1.9.29, 28

scalar

Fact 1.9.29, 28

Fact 1.9.30, 29

Fact 1.10.29, 35

triple product**identity**

Fact 2.12.10, 126

Kronecker product

Proposition 7.1.5, 400

Fact 7.4.7, 405

trace

Fact 4.9.4, 260

Fact 4.9.6, 261

Fact 7.4.7, 405

vec

Proposition 7.1.9, 401

tripotent matrix**definition**

Definition 3.1.1, 165

Drazin generalized**inverse**

Proposition 6.2.2, 368

group-invertible**matrix**

Proposition 3.1.6, 169

Hermitian matrix

Fact 3.16.3, 213

idempotent matrix

Fact 3.16.1, 212

Fact 3.16.5, 213

inertia

Fact 5.8.3, 307

involutory matrix

Fact 3.16.2, 212

Kronecker product

Fact 7.4.16, 406

projector

Fact 6.4.33, 383

rank

Fact 2.10.23, 118

Fact 3.16.3, 213

Fact 3.16.4, 213

reflector

Proposition 3.1.6, 169

signature

Fact 5.8.3, 307

similar matrices

Proposition 3.4.5, 174

Corollary 5.5.22, 301

spectrum

Proposition 5.5.21, 300

trace

Fact 3.16.4, 213

Fact 5.8.3, 307

unitarily similar**matrices**

Proposition 3.4.5, 174

tuple

definition, 3

Turan's inequalities**spectral radius****bound**

Fact 4.10.22, 271

two-sided directional differential

definition, 625

U**ULU decomposition****factorization**

Fact 5.15.11, 346

Umegaki

relative entropy
Fact 11.14.25, 686

**uncontrollable
eigenvalue**

controllability pencil
Proposition 12.6.13,
740

full-state feedback
Proposition 12.6.14,
740

Hamiltonian

Proposition 12.17.7,
784
Proposition 12.17.8,
784
Lemma 12.17.4, 783
Lemma 12.17.6, 783

uncontrollable

multispectrum
definition
Definition 12.6.11, 740

**uncontrollable
spectrum**

controllability pencil
Proposition 12.6.16,
741

definition
Definition 12.6.11, 740

invariant zero
Theorem 12.10.9, 762

Smith zeros

Proposition 12.6.16,
741

**uncontrollable-
unobservable
spectrum**

invariant zero
Theorem 12.10.9, 762

unimodular matrix

coprime right
polynomial fraction
description
Proposition 4.7.15, 253
definition
Definition 4.3.1, 236
determinant

Proposition 4.3.7, 238

Smith form

Proposition 4.3.7, 238

union

boundary
Fact 10.9.2, 634

cardinality
Fact 1.5.5, 11

closed set
Fact 10.9.10, 635

closure
Fact 10.9.2, 634

convex cone
Fact 2.9.10, 111

convex set
Fact 10.9.7, 634

definition, 2
interior

Fact 10.9.2, 634
Fact 10.9.3, 634

open set
Fact 10.9.9, 635

span
Fact 2.9.12, 111

union of ranges

projector
Fact 6.4.41, 385

unipotent matrix

definition
Definition 3.1.1, 165

group
Fact 3.21.5, 222
Fact 11.22.1, 715

Heisenberg group
Fact 3.21.5, 222
Fact 11.22.1, 715

matrix exponential
Fact 11.13.17, 680
spectrum
Proposition 5.5.21, 300

unit imaginary matrix

congruent matrices
Fact 3.7.34, 184
definition, 169
Hamiltonian matrix
Fact 3.19.3, 216

**skew-symmetric
matrix**

Fact 3.7.34, 184

spectrum

Fact 5.9.25, 315

symplectic matrix
Fact 3.19.3, 216

unit impulse function

definition, 724

unit sphere

group
Fact 3.21.2, 221

unit-length quaternions

Sp(1)
Fact 3.22.1, 225

**unitarily biequivalent
matrices**

definition
Definition 3.4.3, 174
singular values
Fact 5.10.18, 319

unitarily invariant norm

commutator
Fact 9.9.29, 584
Fact 9.9.30, 585
Fact 9.9.31, 585

**complex conjugate
transpose**

Fact 9.8.30, 576

definition, 547

fixed-rank
approximation
Fact 9.14.28, 614

Frobenius norm
Fact 9.14.34, 616

Heinz inequality
Fact 9.9.49, 589

Hermitian matrix
Fact 9.9.5, 580
Fact 9.9.41, 588
Fact 9.9.43, 588
Fact 11.16.13, 695

Hermitian

perturbation
Fact 9.12.4, 599

inequality
Fact 9.9.11, 581

- Fact 9.9.44, 588
- Fact 9.9.47, 589
- Fact 9.9.48, 589
- Fact 9.9.49, 589
- Fact 9.9.50, 589
- matrix exponential**
 - Fact 11.15.6, 690
 - Fact 11.16.4, 692
 - Fact 11.16.5, 694
 - Fact 11.16.13, 695
 - Fact 11.16.16, 695
 - Fact 11.16.17, 695
- matrix logarithm**
 - Fact 9.9.54, 590
- matrix power**
 - Fact 9.9.17, 582
- matrix product**
 - Fact 9.9.6, 580
- maximum eigenvalue**
 - Fact 9.9.30, 585
 - Fact 9.9.31, 585
- maximum singular value**
 - Fact 9.9.10, 581
 - Fact 9.9.29, 584
- McIntosh's inequality**
 - Fact 9.9.47, 589
- normal matrix**
 - Fact 9.9.6, 580
- outer-product matrix**
 - Fact 9.8.40, 578
- partitioned matrix**
 - Fact 9.8.33, 576
- polar decomposition**
 - Fact 9.9.42, 588
- positive-semidefinite matrix**
 - Fact 9.9.7, 580
 - Fact 9.9.14, 582
 - Fact 9.9.15, 582
 - Fact 9.9.16, 582
 - Fact 9.9.17, 582
 - Fact 9.9.27, 584
 - Fact 9.9.46, 588
 - Fact 9.9.51, 589
 - Fact 9.9.52, 590
 - Fact 9.9.53, 590
 - Fact 9.9.54, 590
 - Fact 11.16.16, 695
 - Fact 11.16.17, 695
- properties**
 - Fact 9.8.41, 578
- rank**
 - Fact 9.14.28, 614
- Schatten norm**
 - Fact 9.8.9, 572
- Schur product**
 - Fact 9.8.41, 578
 - Fact 9.9.62, 591
 - Fact 9.9.63, 591
 - Fact 9.14.36, 617
- self-adjoint norm**
 - Fact 9.8.7, 572
- similar matrices**
 - Fact 9.8.31, 576
- singular value**
 - Fact 9.14.28, 614
- singular value perturbation**
 - Fact 9.14.29, 614
- square root**
 - Fact 9.9.18, 583
 - Fact 9.9.19, 583
- submultiplicative norm**
 - Fact 9.8.41, 578
 - Fact 9.9.7, 580
- symmetric gauge function**
 - Fact 9.8.42, 579
- unitarily left-equivalent matrices**
 - complex conjugate transpose**
 - Fact 5.10.18, 319
 - Fact 5.10.19, 319
 - definition**
 - Definition 3.4.3, 174
 - positive-semidefinite matrix**
 - Fact 5.10.18, 319
 - Fact 5.10.19, 319
- unitarily right-equivalent matrices**
 - complex conjugate transpose**
 - Fact 5.10.18, 319
 - definition**
 - Definition 3.4.3, 174
 - positive-semidefinite matrix**
 - Fact 5.10.18, 319
 - Fact 5.10.19, 319
- complex conjugate transpose**
 - Fact 5.10.18, 319
- definition**
 - Definition 3.4.3, 174
- positive-semidefinite matrix**
 - Fact 5.10.18, 319
- unitarily similar matrices**
 - biequivalent matrices**
 - Proposition 3.4.5, 174
 - complex conjugate transpose**
 - Fact 5.9.20, 314
 - Fact 5.9.21, 314
 - definition**
 - Definition 3.4.4, 174
 - diagonal entries**
 - Fact 5.9.17, 313
 - Fact 5.9.19, 313
 - elementary matrix**
 - Proposition 5.6.3, 302
 - elementary projector**
 - Proposition 5.6.3, 302
 - elementary reflector**
 - Proposition 5.6.3, 302
 - group-invertible matrix**
 - Proposition 3.4.5, 174
 - Hermitian matrix**
 - Proposition 3.4.5, 174
 - Proposition 5.6.3, 302
 - Corollary 5.4.5, 294
 - idempotent matrix**
 - Proposition 3.4.5, 174
 - Fact 5.9.21, 314
 - Fact 5.9.26, 315
 - Fact 5.9.27, 315
 - Fact 5.10.10, 318
 - involutory matrix**
 - Proposition 3.4.5, 174
 - Kronecker product**
 - Fact 7.4.12, 406
 - matrix classes**
 - Proposition 3.4.5, 174
 - nilpotent matrix**
 - Proposition 3.4.5, 174
 - normal matrix**

- Proposition 3.4.5, 174
- Corollary 5.4.4, 293
- Fact 5.10.6, 317
- Fact 5.10.7, 317
- partitioned matrix**
 - Fact 5.9.23, 314
- positive-definite matrix**
 - Proposition 3.4.5, 174
 - Proposition 5.6.3, 302
- positive-semidefinite matrix**
 - Proposition 3.4.5, 174
 - Proposition 5.6.3, 302
- projector**
 - Fact 5.10.12, 319
- range-Hermitian matrix**
 - Proposition 3.4.5, 174
 - Corollary 5.4.4, 293
- similar matrices**
 - Fact 5.10.7, 317
- simultaneous diagonalization**
 - Fact 5.17.7, 358
- simultaneous triangularization**
 - Fact 5.17.4, 358
 - Fact 5.17.6, 358
- skew-Hermitian matrix**
 - Proposition 3.4.5, 174
 - Proposition 5.6.3, 302
- skew-involutory matrix**
 - Proposition 3.4.5, 174
- trace**
 - Fact 5.10.8, 318
- tripotent matrix**
 - Proposition 3.4.5, 174
- unitary matrix**
 - Proposition 3.4.5, 174
 - Proposition 5.6.3, 302
- upper triangular matrix**
 - Theorem 5.4.1, 292
- unitary biequivalence equivalence relation**
 - Fact 5.10.3, 317
- unitary group**
 - symplectic group
 - Fact 3.21.3, 222
- unitary left equivalence equivalence relation**
 - Fact 5.10.3, 317
- unitary matrix, see orthogonal matrix**
- additive decomposition**
 - Fact 5.19.1, 360
- block-diagonal matrix**
 - Fact 3.11.25, 196
- Cayley transform**
 - Fact 3.11.28, 196
- cogredient diagonalization**
 - Fact 8.16.1, 507
- complex-symmetric matrix**
 - Fact 5.9.22, 314
- convergent sequence**
 - Fact 8.9.33, 454
- CS decomposition**
 - Fact 5.9.29, 316
- definition**
 - Definition 3.1.1, 165
- determinant**
 - Fact 3.11.15, 194
 - Fact 3.11.20, 196
 - Fact 3.11.23, 196
 - Fact 3.11.24, 196
- diagonal entries**
 - Fact 3.11.19, 195
 - Fact 8.17.10, 511
- diagonal matrix**
 - Theorem 5.6.4, 302
- discrete-time Lyapunov-stable matrix**
 - Fact 11.21.13, 714
- dissipative matrix**
 - Fact 8.9.31, 453
- factorization**
 - Fact 5.15.8, 346
 - Fact 5.18.6, 359
- Frobenius norm**
 - Fact 9.9.42, 588
- geometric-mean decomposition**
 - Fact 5.9.30, 316
- group**
 - Proposition 3.3.6, 172
- group generalized inverse**
 - Fact 6.3.34, 376
- Hermitian matrix**
 - Fact 3.11.29, 197
 - Fact 8.16.1, 507
 - Fact 11.14.34, 688
- identities**
 - Fact 3.11.3, 189
- Kronecker product**
 - Fact 7.4.16, 406
- matrix exponential**
 - Proposition 11.2.8, 649
 - Proposition 11.2.9, 650
 - Proposition 11.6.7, 659
 - Corollary 11.2.6, 648
 - Fact 11.14.6, 683
 - Fact 11.14.33, 688
 - Fact 11.14.34, 688
- matrix limit**
 - Fact 6.3.34, 376
- normal matrix**
 - Proposition 3.1.6, 169
 - Fact 3.11.4, 189
 - Fact 5.15.1, 345
- orthogonal vectors**
 - Fact 3.11.14, 194
- outer-product perturbation**
 - Fact 3.11.15, 194
- partitioned matrix**
 - Fact 3.11.24, 196
 - Fact 3.11.26, 196
 - Fact 3.11.27, 196
 - Fact 8.11.22, 473
 - Fact 8.11.23, 473
 - Fact 8.11.24, 473
 - Fact 9.14.11, 609
- polar decomposition**
 - Fact 5.18.8, 360
- quaternions**
 - Fact 3.22.9, 229
- reflector**

Fact 3.14.2, 211
semicontractive matrix
 Fact 8.11.22, 473
shifted-unitary matrix
 Fact 3.11.33, 198
simultaneous diagonalization
 Fact 8.16.1, 507
singular value
 Fact 5.11.37, 328
 Fact 9.14.11, 609
skew reflector
 Fact 3.14.6, 211
skew-Hermitian matrix
 Fact 3.11.28, 196
 Fact 3.14.6, 211
 Fact 11.14.33, 688
skew-involutory matrix
 Fact 3.14.6, 211
spectrum
 Proposition 5.5.21, 300
square root
 Fact 8.9.26, 453
subspace
 Fact 3.11.1, 189
 Fact 3.11.2, 189
sum
 Fact 3.11.23, 196
trace
 Fact 3.11.16, 194
 Fact 3.11.32, 198
unitarily similar matrices
 Proposition 3.4.5, 174
 Proposition 5.6.3, 302
upper triangular matrix
 Fact 5.15.8, 346
unitary right equivalence
 equivalence relation
 Fact 5.10.3, 317
unitary similarity
 equivalence relation

Fact 5.10.3, 317
singular value decomposition
 Fact 5.9.28, 315
 Fact 6.3.15, 373
 Fact 6.6.15, 395
universal statement
 definition, 2
 logical equivalents
 Fact 1.5.4, 11
unobservable eigenvalue
 definition
 Definition 12.3.11, 730
full-state feedback
 Proposition 12.3.14, 731
Hamiltonian
 Proposition 12.17.7, 784
 Proposition 12.17.8, 784
 Lemma 12.17.4, 783
 Lemma 12.17.6, 783
invariant zero
 Proposition 12.10.11, 764
observability pencil
 Proposition 12.3.13, 731
unobservable multispectrum
 definition
 Definition 12.3.11, 730
unobservable spectrum
 definition
 Definition 12.3.11, 730
invariant zero
 Theorem 12.10.9, 762
observability pencil
 Proposition 12.3.16, 731
Smith zeros
 Proposition 12.3.16, 731
unobservable subspace

block-triangular matrix
 Proposition 12.3.9, 730
 Proposition 12.3.10, 730
definition
 Definition 12.3.1, 728
equivalent expressions
 Lemma 12.3.2, 728
full-state feedback
 Proposition 12.3.5, 729
identity shift
 Lemma 12.3.7, 730
invariant subspace
 Corollary 12.3.4, 729
nonsingular matrix
 Proposition 12.3.10, 730
orthogonal matrix
 Proposition 12.3.9, 730
projector
 Lemma 12.3.6, 729
unstable equilibrium
 definition
 Definition 11.7.1, 660
unstable matrix
positive matrix
 Fact 11.18.20, 701
unstable subspace
 complementary subspaces
 Proposition 11.8.8, 665
definition, 665
idempotent matrix
 Proposition 11.8.8, 665
invariant subspace
 Proposition 11.8.8, 665
semistable matrix
 Proposition 11.8.8, 665
upper block-triangular matrix
 characteristic polynomial
 Fact 4.10.11, 267
definition
 Definition 3.1.3, 167

- inverse matrix
 - Fact 2.17.7, 148
 - Fact 2.17.9, 148
- minimal polynomial
 - Fact 4.10.12, 268
- orthogonally similar matrices
 - Corollary 5.4.2, 293
- power
 - Fact 2.12.21, 128
- upper bound
 - positive-definite matrix
 - Fact 8.10.31, 459
- upper bound for a partial ordering
 - definition
 - Definition 1.3.9, 7
- upper Hessenberg
 - regular pencil
 - Fact 5.17.3, 358
 - simultaneous
 - orthogonal
 - biequivalence transformation
 - Fact 5.17.3, 358
- upper Hessenberg matrix
 - definition
 - Definition 3.1.3, 167
- upper triangular
 - regular pencil
 - Fact 5.17.3, 358
 - simultaneous
 - orthogonal
 - biequivalence transformation
 - Fact 5.17.3, 358
- upper triangular matrix
 - commutator
 - Fact 3.17.11, 214
 - definition
 - Definition 3.1.3, 167
 - factorization
 - Fact 5.15.8, 346
 - Fact 5.15.10, 346
- group
 - Fact 3.21.5, 222
 - Fact 11.22.1, 715
- Heisenberg group
 - Fact 3.21.5, 222
 - Fact 11.22.1, 715
- invariant subspace
 - Fact 5.9.2, 311
- Kronecker product
 - Fact 7.4.3, 405
- Lie algebra
 - Fact 3.21.4, 222
 - Fact 11.22.1, 715
- matrix exponential
 - Fact 11.11.4, 672
 - Fact 11.13.1, 677
 - Fact 11.13.16, 680
- matrix power
 - Fact 3.18.7, 216
- matrix product
 - Fact 3.20.18, 221
- nilpotent matrix
 - Fact 3.17.11, 214
- orthogonally similar matrices
 - Corollary 5.4.3, 293
- positive diagonal
 - Fact 5.15.9, 346
- positive-semidefinite matrix
 - Fact 8.9.37, 454
- similar matrices
 - Fact 5.9.2, 311
- Toeplitz matrix
 - Fact 3.18.7, 216
 - Fact 11.13.1, 677
- unitarily similar matrices
 - Theorem 5.4.1, 292
- unitary matrix
 - Fact 5.15.8, 346
- Urquhart
 - generalized inverse
 - Fact 6.3.14, 372
- companion matrix
 - Fact 5.16.4, 354
- determinant
 - Fact 5.16.3, 354
- Fourier matrix
 - Fact 5.16.7, 355
- polynomial
 - Fact 5.16.6, 355
- similar matrices
 - Fact 5.16.5, 354
- variance
 - Laguerre-Samuelson inequality
 - Fact 1.15.12, 51
 - Fact 8.9.35, 454
- variance inequality
 - mean
 - Fact 1.15.12, 51
 - Fact 8.9.35, 454
- variation of constants
 - formula
 - state equation
 - Proposition 12.1.1, 723
- variational cone
 - definition, 625
 - dimension
 - Fact 10.8.20, 633
- vec
 - definition, 399
- Kronecker
 - permutation matrix
 - Fact 7.4.29, 407
- Kronecker product
 - Fact 7.4.5, 405
 - Fact 7.4.6, 405
 - Fact 7.4.8, 405
- matrix product
 - Fact 7.4.6, 405
- quadruple product
 - Fact 7.4.9, 405
- trace
 - Proposition 7.1.1, 399
 - Fact 7.4.7, 405
 - Fact 7.4.9, 405
- triple product
 - Proposition 7.1.9, 401

V

Vandermonde matrix

vector

- definition, 78
- Hölder norm
 - Fact 9.7.34, 571

vector derivative

- quadratic form
 - Proposition 10.7.1, 630

vector identity

- cosine law
 - Fact 9.7.4, 563
- parallelogram law
 - Fact 9.7.4, 563
- polarization identity
 - Fact 9.7.4, 563
- Pythagorean theorem
 - Fact 9.7.4, 563

vector inequality

- Hölder's inequality
 - Proposition 9.1.6, 545
- norm inequality
 - Fact 9.7.11, 567
 - Fact 9.7.12, 567
 - Fact 9.7.14, 568
 - Fact 9.7.15, 568

vibration equation

- matrix exponential
 - Example 11.3.7, 653

volume

- convex polyhedron
 - Fact 2.20.20, 160
- ellipsoid
 - Fact 3.7.35, 184
- hyperellipsoid
 - Fact 3.7.35, 184
- parallelepiped
 - Fact 2.20.16, 160
 - Fact 2.20.17, 160
- simplex
 - Fact 2.20.19, 160
- tetrahedron
 - Fact 2.20.15, 160
- transformed set
 - Fact 2.20.18, 160

von Neumann

- symmetric gauge function and unitarily invariant norm
 - Fact 9.8.42, 579

von Neumann's trace inequality

- trace and singular value
 - Fact 9.12.1, 599

von Neumann–Jordan inequality

- norm inequality
 - Fact 9.7.11, 567

W**walk**

- connected graph
 - Fact 4.11.4, 273
- definition
 - Definition 1.4.3, 9
- graph
 - Fact 4.11.3, 273

Walker's inequality

- scalar inequality
 - Fact 1.11.22, 45

Walsh

- polynomial root bound
 - Fact 11.20.5, 709

Wang's inequality

- scalar inequality
 - Fact 1.15.13, 51

weak diagonal

- dominance theorem
- nonsingular matrix
 - Fact 4.10.19, 270

weak log majorization

- definition
 - Definition 2.1.1, 78
- eigenvalue
 - Fact 8.18.27, 521
- singular value
 - Proposition 9.6.2, 561

- weak majorization
 - Fact 2.21.13, 164

weak majorization**convex function**

- Fact 2.21.8, 163
- Fact 2.21.9, 163
- Fact 2.21.10, 163
- Fact 2.21.11, 163
- Fact 8.18.5, 513

definition

- Definition 2.1.1, 78

eigenvalue

- Fact 8.17.5, 509
- Fact 8.18.5, 513
- Fact 8.18.6, 514
- Fact 8.18.27, 521

eigenvalue of**Hermitian part**

- Fact 5.11.27, 326

increasing function

- Fact 2.21.10, 163

matrix exponential

- Fact 11.16.4, 692

positive-semidefinite**matrix**

- Fact 8.18.6, 514

powers

- Fact 2.21.14, 164

scalar inequality

- Fact 2.21.2, 162
- Fact 2.21.3, 162

Schur product

- Fact 9.14.31, 615

singular value

- Proposition 9.2.2, 548
- Proposition 9.6.3, 561
- Fact 5.11.27, 326
- Fact 8.17.5, 509
- Fact 8.18.7, 514
- Fact 9.14.19, 611
- Fact 9.14.20, 612

singular value**inequality**

- Fact 8.18.21, 519
- Fact 9.13.17, 604
- Fact 9.13.18, 605
- Fact 9.13.20, 605
- Fact 9.14.31, 615

symmetric gauge function

Fact 2.21.14, 164

weak log

majorization

Fact 2.21.13, 164

Weyl majorant theorem

Fact 9.13.20, 605

Weyl's inequalities

Fact 8.17.5, 509

weakly unitarily

invariant norm

definition, 547

matrix power

Fact 9.8.38, 577

numerical radius

Fact 9.8.38, 577

Wei-Norman expansion

time-varying

dynamics

Fact 11.13.4, 678

Weierstrass

cogredient

diagonalization of

positive-definite

matrices

Fact 8.16.2, 507

Weierstrass canonical form

pencil

Proposition 5.7.3, 305

weighted

arithmetic-mean–

geometric-mean

inequality

arithmetic-mean–

geometric-mean

inequality

Fact 1.15.32, 56

Weyl, 428

singular value

inequality

Fact 5.11.28, 326

singular values and strong log

majorization

Fact 9.13.19, 605

Weyl majorant theorem

singular values and

weak majorization

Fact 9.13.20, 605

Weyl's inequalities

weak majorization

and singular values

Fact 8.17.5, 509

Weyl's inequality

Hermitian matrix

eigenvalues

Theorem 8.4.9, 427

Fact 8.10.4, 456

Wielandt

eigenvalue

perturbation

Fact 9.12.9, 601

positive power of a

primitive matrix

Fact 4.11.5, 273

Wielandt inequality

quadratic form

inequality

Fact 8.15.29, 505

X

Xie

asymptotically stable

polynomial

Fact 11.17.7, 697

Y

Yamamoto

singular value limit

Fact 9.13.22, 606

Young inequality

positive-definite

matrix

Fact 8.9.42, 455

Fact 8.10.46, 464

reverse inequality

Fact 1.10.22, 34

scalar inequality

Fact 1.10.21, 33

Specht's ratio

Fact 1.10.22, 34

Young's inequality

positive-semidefinite

matrix

Fact 8.12.12, 477

positive-semidefinite

matrix inequality

Fact 9.14.22, 612

scalar case

Fact 1.10.32, 36

Fact 1.15.31, 56

Z

Z-matrix

definition

Definition 3.1.4, 168

M-matrix

Fact 4.11.6, 275

Fact 4.11.8, 276

M-matrix inequality

Fact 4.11.8, 276

matrix exponential

Fact 11.19.1, 706

minimum eigenvalue

Fact 4.11.9, 276

submatrix

Fact 4.11.7, 276

Zassenhaus expansion

time-varying

dynamics

Fact 11.13.4, 678

Zassenhaus product formula

matrix exponential

Fact 11.14.18, 685

zero

blocking

Definition 4.7.10, 251

invariant

Definition 12.10.1, 757

invariant and determinant

Fact 12.22.14, 800

invariant and equivalent realizations

Proposition 12.10.10, 764

invariant and**full-state feedback**

Proposition 12.10.10, 764

Fact 12.22.14, 800

invariant and**observable pair**

Corollary 12.10.12, 765

invariant and**transmission**

Theorem 12.10.8, 762

invariant and**unobservable****eigenvalue**

Proposition 12.10.11, 764

transmission

Definition 4.7.10, 251

Proposition 4.7.12, 251

transmission and**invariant**

Theorem 12.10.8, 762

zero diagonal**commutator**

Fact 3.8.2, 184

zero entry**reducible matrix**

Fact 3.20.1, 217

Fact 3.20.2, 217

zero matrix**definition, 83****positive-semidefinite****matrix**

Fact 8.10.10, 457

trace

Fact 2.12.14, 127

Fact 2.12.15, 127

zero of a rational function**definition**

Definition 4.7.1, 249

zero trace**Shoda's theorem**

Fact 5.9.18, 313

zeros matrix**maximal null space**

Fact 2.12.12, 127

zeta function**Euler product****formula**

Fact 1.7.8, 19

MATRIX MATHEMATICS

Second Edition

Theory, Facts, and Formulas

When first published in 2005, *Matrix Mathematics* quickly became the essential reference book for users of matrices in all branches of engineering, science, and applied mathematics. In this fully updated and expanded edition, the author brings together the latest results on matrix theory to make this the most complete, current, and easy-to-use book on matrices.

Each chapter describes relevant background theory followed by specialized results. Hundreds of identities, inequalities, and matrix facts are stated clearly and rigorously with cross references, citations to the literature, and illuminating remarks. Beginning with preliminaries on sets, functions, and relations, *Matrix Mathematics* covers all of the major topics in matrix theory, including matrix transformations; polynomial matrices; matrix decompositions; generalized inverses; Kronecker and Schur algebra; positive-semidefinite matrices; vector and matrix norms; the matrix exponential and stability theory; and linear systems and control theory. Also included are a detailed list of symbols, a summary of notation and conventions, an extensive bibliography and author index with page references, and an exhaustive subject index. This significantly expanded edition of *Matrix Mathematics* features a wealth of new material on graphs, scalar identities and inequalities, alternative partial orderings, matrix pencils, finite groups, zeros of multivariable transfer functions, roots of polynomials, convex functions, and matrix norms.

- Covers hundreds of important and useful results on matrix theory, many never before available in any book
- Provides a list of symbols and a summary of conventions for easy use
- Includes an extensive collection of scalar identities and inequalities
- Features a detailed bibliography and author index with page references
- Includes an exhaustive subject index with cross-referencing

DENNIS S. BERNSTEIN is professor of aerospace engineering at the University of Michigan.

PRAISE FOR THE FIRST EDITION OF MATRIX MATHEMATICS

"When a matrix question is thrown my way, I will now refer my correspondents . . . to Bernstein's handbook."

—Philip J. Davis, *SIAM News*

"*Matrix Mathematics* contains an impressive collection of definitions, relations, properties, equations, inequalities, and facts centered around matrices and their use in systems and control. The amount of material that is covered is quite impressive and well structured. . . . I highly recommend the book as a source for retrieving matrix results that one would otherwise have to search for in the extensive literature on matrix theory."

—Paul Van Dooren,
IEEE Control Systems Magazine

"The author was very successful in collecting the enormous amount of results in matrix theory in a single source. . . . A beautiful work and an admirable performance!"

—H. Mitsch,
Monatshefte für Mathematik

Front cover concept by the author

 PRINCETON
press.princeton.edu

ISBN: 978-0-691-14039-1

90000
9 780691 140391