



Ethiopian TVET-System



MEDICAL LABORATORY

Level -III

Based on Apr.2018G.C. Occupational Standard

**Module Title: Apply Infection Prevention
Techniques and Workplace OHS**
TTLM Code: HLT MLT3 TTLM 0919v1

**This module includes the following Learning
Guides**

LG06: Follow infection control guidelines

LG07: Identify and respond to infection risks

LG08: Maintain personal hygiene

LG09: Use PPE

LG10: Limit contamination

**LG11: Handle, package, label, store, transport and dispose of
clinical and other waste**

LG12: Clean environmental surfaces



Instruction Sheet

LG06: Follow infection control guidelines

This learning guide is developed to provide you the necessary information regarding the following **content coverage and topics** –

Follow infection control guidelines

- preventing spread of infection
- using additional precautions if there is no sufficient standard precautions to prevent transmission of infection
- minimizing Contamination of materials, equipment and instruments by aerosols and splatter

This guide will also assist you to attain the learning outcome stated in the cover page. Specifically, upon completion of this Learning Guide, you will be able to –

- Demonstrate the application of standard precautions to prevent the spread of infection in accordance with organization requirements
- The application of *additional precautions* is demonstrated when standard precautions alone may not be sufficient to prevent transmission of infection
- *minimized* contamination of materials, equipment and instruments by aerosols and splatter.

Learning Instructions:

1. Read the specific objectives of this Learning Guide.
2. Follow the instructions described in **number 3 to 16**.
3. Read the information written in the “Information Sheets 1”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
4. Accomplish the “Self-check 1” in **page 8**.
5. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 1).
6. If you earned a satisfactory evaluation proceed to “Information Sheet 2”. However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 1.
7. Submit your accomplished Self-check. This will form part of your training portfolio.
8. Read the information written in the “Information Sheet 2”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
9. Accomplish the “Self-check 2” **in page 16**.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 2 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	---------------



10. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 2).
11. Read the information written in the “Information **Sheets 3**”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
12. Accomplish the “Self-check 3” in **page 19**.
13. Ask your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 3).
14. If you earned a satisfactory evaluation proceed to “Operation Sheet 1” **in page 21**; However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 3.
15. Read the “Operation Sheet 1 and try to understand the procedures discussed.
16. Do the “LAP test” in **page 22** (if you are ready). Request your trainer to evaluate your performance and outputs. Your trainer will give you feedback and the evaluation will be either satisfactory or unsatisfactory. If unsatisfactory, your trainer shall advice you on additional work. But if satisfactory you can proceed to Learning Guide #2

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 3 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	---------------



1. Preventing spread of infection

1.1. Introduction

1.1.1 Definition of terms

Asepsis and aseptic Technique: A combination of efforts made to prevent entry of microorganisms into any area of the body

- The goal of aseptic technique is to reduce or eliminate the number of microorganisms on both animate (living) surfaces (skin and mucous membranes) and inanimate objects (surgical instruments and other items) to an infection free

Antiseptic: A germicide that is used on skin or living tissue for the purpose of inhibiting or destroying microorganisms. Examples include alcohols, chlorhexidine, chlorine, hexachlorophene, and iodine

Antisepsis: a process of reducing the number of microorganisms on the skin, mucous membranes or other body tissue by applying an antimicrobial (antiseptic) agent

Antimicrobial agents: A general term for the drugs, chemicals, or other substances that either kill or slow the growth of microbes. Among the antimicrobial agents in use today are antibacterial drugs (which kill bacteria), antiviral agents (which kill viruses), antifungal agents (which kill fungi), and antiparasitic drugs (which kill parasites).

Cleaning: - a process that physically removes all visible dust, soil blood or other body fluids from inanimate object as well as removing sufficient number of microorganisms to reduce risks for those who touch the skin or handle the object.

- It consists of thorough washing with soap or detergent and water, rinsing with clean water and drying

One of the most basic measures for the maintenance of hygiene, and one that is particularly important in the hospital environment, is cleaning. The principal aim of cleaning is to remove visible dirt. It is essentially a mechanical process: the dirt is dissolved by water, diluted until it is no longer visible, and rinsed off. Soaps and detergents act as solubility promoting agents

Colonization: the presence of pathogenic (illness or disease causing) organisms in a person or animal in abundance (pathogens can be detected by cultures or other tests) usually without causing symptoms or clinical findings (they do not invade tissues, cause cellular changes or cause damages).

- It also defined as the presence or increased number of a particular invasive bacterial species in the resident micro flora.

Colonized person: can be a major source of transfer of pathogens to other persons.

- For instance
 - Neisseria meningitides colonizes nasal cavity and oropharynx with or without causing subsequent infections
 - Entamoeba histolytica can colonize the large bowel without any harm to the host but are often shade in the stool as infectious cysts which may cause dysentery



Contamination: The presence of an infectious agent on a body surface or on clothes, gowns, gloves, bedding, furniture, computer keyboards, or other inanimate objects that may be capable of producing disease or infection

Contact: An exposed individual who might have been infected through transmission from another host or the environment

Decontamination: a process that makes inanimate objects safer for the staff to handle them before cleaning (i.e. inactivates HBV, HCV and HIV and reduce, but does not eliminate, the number of other contaminating microorganisms)

- It is also defined as a process or treatment that renders a medical device, instrument, or environmental surface safe to handle because it is no longer capable of transmitting particles of infectious material.

Disinfectant: A chemical agent used on inanimate (non-living) objects to destroy virtually all recognized pathogenic microorganisms, but not necessarily all microbial forms (e.g., bacterial spores).

Disinfection: The destruction of pathogenic and other kinds of microorganisms by physical or chemical means. Disinfection is less lethal than sterilization, because it destroys most recognized pathogenic microorganisms, but not necessarily all microbial forms, such as bacterial spores.

Disease: any deviation from being healthy or interruption of the normal structure or function of any body part, organ, or system manifested by a characteristics set of symptoms and signs whose etiology, pathology, and prognosis may be known or unknown

Endogenous infection: Micro-organisms originating from the service user's own body which cause harm in another body site.

Exogenous infection: Micro-organisms originating from a source or reservoir which are transmitted by any mechanism to a person, i.e. contact or airborne routes.

High level disinfection (HLD): a process of eliminating all microorganisms except some bacterial endospores from inanimate objects by boiling, steaming or using chemical disinfectants

Infection: is an invasion and multiplication of microorganisms in body tissues which may clinically be apparent or result in local cellular injury due to competitive metabolism, toxins, intracellular replication or antigen antibody response.

Infectious Microorganisms: are microorganisms capable of producing disease in the appropriate hosts

Infection prevention: is a systematic effort or process of placing barriers between a susceptible host (person lacking effective natural or acquired protection) and the microorganisms

Microorganisms: are causative agents of infections such as bacteria, viruses, fungi and parasite

Nosocomial Infection: is a term used interchangeably with "healthcare facility acquired infection" or "healthcare associated infections (HAIs)" and is defined as a situation in which patients coming to health institutions seeking treatments acquire an infection/s in healthcare facility afterwards other than disease/health problems they had

- It also defined as infections acquired while a patient is under hospital or any health facility care which are not present or incubating at time of admission
- It is a time related criterion which refers to infectious occurring more than 48 hours after admission

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 5 of 139
---------------------------	----------------------	---------------------------------------	-----------------------	---------------



Nosocomial infections—known also as hospital-acquired infections, hospital-associated infections, and hospital infections—are infections that are not present in the patient at the time of admission to hospital but develop during the course of the stay in hospital. There are two forms:

- **Endogenous infection, self-infection, or auto-infection.** The causative agent of the infection is present in the patient at the time of admission to hospital but there are no signs of infection. The infection develops during the stay in hospital as a result of the patient's altered resistance.

- **Cross-contamination followed by cross-infection.** During the stay in hospital the patient comes into contact with new infective agents, becomes contaminated, and subsequently develops an infection.

Patient Safety: is an issue focusing at reduction or aversion of unsafe acts/circumstances within the healthcare system through the use of best practices leading to optimal patient outcomes

Protective Barriers: are physical, mechanical or chemical processes that help to prevent the spread of infectious microorganisms from person to person (patient, healthcare client or health workers) and /or equipment, instruments and environmental surface to people

Sterilization: a process of eliminating all microorganisms (bacteria, viruses, fungi and parasites) including bacterial endospores from inanimate object by high-pressure steam (autoclave), dry heat (oven), chemical sterilization or radiation

1.1.2 General principles of infection prevention

Good infection prevention and control are essential to ensure that people who use health and social care services receive safe and effective care. Effective prevention and control of infection must be part of everyday practice and be applied consistently by everyone

Adherence to good practice in relation to infection prevention and control has been shown to reduce the risk of infection to residents and care workers. Providers should have relevant policies in place as identified by local risk assessment, and having taken account of the Code of Practice. Staff training is important and will improve compliance with policies, which should be regularly audited, updated and clearly marked with a review date.

It is not always possible to identify people who may spread infection to others and precautions for the care and treatment of all residents are recommended to prevent this spread of infection. Basic infection prevention and control practices are designed to reduce the risk of cross infection from both recognized and unrecognized sources of infection and should be applied at all times by all care workers. All those who provide care should be trained in understanding the basic infection prevention and control practices.

Standard principles

Some of the general advice includes

Everyone involved in providing care should be:

- educated about the standard principles of infection prevention and control **and** trained in hand decontamination, the use of personal protective equipment, and the safe use and disposal of sharps.

Wherever care is delivered, healthcare workers must have available appropriate supplies of:

- materials for hand decontamination
- sharps containers
- personal protective equipment.

Educate patients and care givers about:

- the benefits of effective hand decontamination

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 6 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	---------------



- the correct techniques and timing of hand decontamination
- when it is appropriate to use liquid soap and water or hand rub
- the availability of hand decontamination facilities
- their role in maintaining standards of healthcare workers' hand decontamination.

Waste disposal

1.1.3 Risk of disease transmission in the health care facility

Healthcare personnel including the support staff (e.g. housekeeping, laundry staff and maintenance), who work in healthcare settings are at risk of exposure to serious potentially life threatening infectious such as HIV, HBV, HCV. Direct contact with blood and other body fluids is the most common or frequent risk healthcare workers encounter while caring of patients. Studies in the United States have shown that the risk of acquiring HBV after being stuck with a needle from an HBV+ client ranges 27 to 37%. In addition, the risk of acquiring HCV and HIV after being stuck with a needle from an infected person is 3 to 10% and 0.2% to 0.4% respectively (Gerberding, 1990; Gershon et al., 1995 & Landpher, 1994). Among these, the efficiency for transmission of hepatitis B is high. For example, an accidental splash in the eye of as little as 10^{-8} ml (0.0000001m) of infected blood can transmit HBV to a susceptible host (Bond et al., 1982).

Each year, In United States, 800,000 sharp injuries are reported by healthcare workers (Rogers, 1997). However, it is anticipated that most healthcare workers do not report needle stick or other sharps injuries as often as they should. Hence, the number of needle stick injuries is under reported in most cases

Similarly, a survey conducted in the year 2003 and 2004 on about 40 health facilities in Ethiopia, reported that 32% of the healthcare workers in these institution sustained needle stick injuries in 12-months' time.

The Disease Transmission Cycle

Microorganisms live everywhere in our environment. Human normally carry them on their skin, upper respiratory, intestinal and genital tracts. Generally, microorganisms live in animals, plants, soil, air and water. Not all, but some among these microorganism are pathogenic (likely to cause disease) in varied degrees. When they get favorable conditions, most of these microorganisms may cause infections if transmitted to immune compromised people such as patients with AIDS (Burke, 1977)

Microorganisms can be classified as bacteria, viruses, fungi, and protozoa. Bacteria can be further divided into three categories: vegetative (e.g., Staphylococcus), mycobacteria (e.g., tuberculosis [TB]), and endospores (e.g., tetanus). Of all the common infectious agents, endospores are the most difficult to kill due to their protective coating

All humans are susceptible to bacterial infections and also to most viral agents. The number (dose) of organisms necessary to produce infection in a susceptible host varies with the location.

The essential factor for the transmission of disease-causing microorganisms from person to person comprises the following six components:

The disease transmission cycle, shown in Figure 1.1, describes how infections are transmitted from one person to another.

1. **Agent:** The microorganism that can cause infection or disease. The infectious agent can include bacteria, viruses, fungi, and parasites.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 7 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	---------------



2. **Host or reservoir:** The reservoir of an agent is the habitat in which an infectious agent normally lives, grows, and/or multiplies. Reservoirs include humans, animals, and the environment (plants, soil, air, water, etc.). Solutions, instruments, and other items used in clinical procedures can also serve as reservoirs for potentially infectious microorganisms.
3. **Portal of exit:** The gateway through which the agent leaves the host or reservoir. The infectious agent can leave the reservoir through the bloodstream, broken skin (e.g., puncture, cut, surgical site, rash), mucous membranes (e.g., eyes, nose, mouth), respiratory tract (e.g., upper respiratory), genitourinary tract (e.g., vagina, penis), gastrointestinal tract (e.g., mouth, anus), or placenta by means of blood, excretions, secretions, or droplets that come from these places. However, the agent must have the right environment in which to survive until it infects another person. For example, the bacteria that cause TB can survive in sputum for weeks but will be killed by sunlight within a few hours.
4. **Method of transmission:** After an agent exits its natural reservoir, it may be transmitted to a susceptible host in numerous ways. These modes of transmission are classified as: -
 - a. **Direct transmission:** An immediate transfer of the agent from a reservoir to a susceptible host by direct contact or droplet spread.
 - i. **Direct contact:** This occurs through kissing, skin-to-skin contact, and sexual intercourse. Direct contact refers also to contact with soil or vegetation harboring infectious organisms. Thus, infectious mononucleosis (“kissing disease”) and gonorrhea are spread from person to person by direct contact. Hookworm is spread by direct contact with contaminated soil.
 - ii. **Droplet spread:** This refers to spray with relatively large, short-range aerosols produced by sneezing, coughing, or even talking. Droplet spread is classified as direct because transmission is by direct spray over a few feet, before the droplets fall to the ground (e.g., influenza, TB).




Fig1.1. disease transmission cycle

b. **Indirect transmission:** In indirect transmission, an agent is carried from a reservoir to a susceptible host by suspended air particles (airborne) or by animate (vector) or inanimate (vehicle-borne) intermediaries.

- i. **Airborne:** The infectious agent can be carried by air currents (e.g., measles, TB).
- ii. **Vehicle-borne:** The infectious agent can be transmitted indirectly from the reservoir to a susceptible host by material that maintains the life of the infectious agent. Such vehicles include food (e.g., Salmonella), blood (e.g., HBV, HIV), water (e.g., cholera, Shigella), or instruments and other items.

iii. **Vector-borne:**

1. **Biological:** The infectious agent can be transmitted to a susceptible host through insects and other invertebrate animals (e.g., mosquitoes can transmit malaria and yellow fever; fleas can transmit plague).
2. **Mechanical:** The infectious agent is carried by certain animals without having any biological change (flies can carry and transmit Vibrio cholera).

Infection prevention in a health care setting primarily deals with preventing the spread of infectious diseases through the air, blood or body fluids, and contact, including droplet, fecaloral, and food-borne routes.

5. **Portal of entry:** The gateway through which an infectious agent enters into the susceptible host. These portals of entry could be mouth, nose, skin, or other (including the respiratory system, vascular system, genitourinary system, and placenta).

6. **Susceptible host:** An organism that catches the infectious agent/pathogen. People are exposed to disease-causing agents every day but do not always get sick. For the purpose of this training, susceptible hosts include clients, service providers, support staff, and members of the community.



Preventing Infectious Diseases

To prevent the transmission of infections, the disease transmission cycle needs to be broken at some point. Hence, understanding the disease transmission cycle is important if health care workers are to:

- Prevent transmission of microorganisms from patient to patient, from patient to provider or from provider to patient during medical and surgical procedures as well as from health facilities to the general community.
- Teach others the factors required for transmission to occur.
- Most importantly, teach others how to break the disease transmission cycle.

In a health care facility, this can be accomplished by following proper Infection Prevention and patient safety (IPPS) practice, such as the following:

- Reducing the number of microorganisms present (e.g., hand washing, cleaning of instruments)
- Killing, inhibiting, or inactivating microorganisms (e.g., hand washing with a waterless alcohol preparation, decontamination, sterilization, or high-level disinfection [HLD])
- Creating barriers to prevent infectious agents from spreading (e.g., wearing gloves or personal protective equipment [PPE])
- Reducing or eliminating risky practices (e.g., by using hands-free technique in the operation room, using disposable gloves and syringes, etc.)
- Making sure that people, especially health care workers, are immune or vaccinated
- Reducing adverse events (e.g., by improving data collection, epidemiological surveys of adverse events, training on prevention of adverse events)

Spaulding Categories of Potential Infection Risk

In 1968, Spaulding proposed three categories of potential infection risk to serve as the basis for selecting the prevention practice or process to use (e.g., sterilization or medical instruments, gloves, and other items) when caring for patients

This classification has stood the test of time and still serves as a good basis for setting priorities for any infection prevention program. The Spaulding categories are summarized below:

- **Critical:** These items and practices affect normally sterile tissues or the blood system and represent the highest level of infection risk. Failure to provide management of sterile or, where appropriate, high-level disinfected items is most likely to result in infections that are most serious.
 - ✓ Critical items confer a high risk for infection if they are contaminated with any microorganism as they are entering sterile tissue or the vascular system so they must be sterile because any microbial contamination could be a source to transmit diseases. That class includes medical devices e.g. Implants, invasive catheters , Implants and any device that touch sterile body cavities. Utmost of the articles in this class should be purchased as sterile or be sterilized before usage.
- **Semi-critical:** These items and practices are second in importance and affect mucous membranes and small areas of non-intact skin. Management needs are considerable and require knowledge and skills in handling many invasive devices (e.g., gastrointestinal endoscopes and vaginal specula), performing decontamination, cleaning and high-level disinfection, and gloving for personnel who touch mucous membranes and non-intact skin.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 10 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



- ✓ This class includes metabolic process medical care and physiological state instrumentality, some endoscopes, medical instrument blades, passageway manometry probes, cystoscopes, body part
- **Noncritical:** Management of items and practices that involve intact skin and represent the lowest level of risk.
 - ✓ Poor management of noncritical items such as overuse of examination gloves often consumes a major share of resources while providing only limited benefits. (Spaulding 1968).
 - ✓ Noncritical things, which contact with intact skin, however, not secretion membranes. Intact skin acts as an honest barrier to most microorganisms; therefore, the sterility of things returning in touch with intact skin is “not essential.”
 - ✓ Objects That contact Intact skin but not Mucous membranes, and Require low level disinfection

1.1.4 Common health facility-acquired infections (HCFs)

These infections earlier called ‘**nosocomial infections**’ (NI) or simply ‘**hospital infections**’ are infections occurring during a stay in hospital that were neither present nor incubating at the time of hospital admission. Mostly, nosocomial infections only appear in patients hospitalized for 48 hours or longer

Healthcare-associated infections:

- are a consistent issue for both hospital patients and healthcare providers
- occur in both adult and pediatric patients. Bloodstream infections, followed by pneumonia and urinary tract infections are the most infections in children, urinary tract infections are the most common healthcare – associated infections in adults.

Infections in HCFs have become a major health problem especially in the health institutions located in developing countries. Therefore, identification of these infections, their source and some factors responsible for their acquisition is very important.

There are many microbes responsible for various forms of infection in health care facilities. For simplicity the most common infections and their etiologies are discussed in the following categories

I. Bacterial infections

Bacteria are the major cause of infections in HCFs. The Gram-negative bacteria are most commonly isolated pathogens from all sites of infection. This is mainly because of the abundant anti-microbial that affects gram-negative bacteria for which the bacteria develop resistance.

The following are the most important bacterial infections that deserve every health professional’s awareness in HCFs:

1. Urinary tract infections

Urinary tract infections include infections of the urethra, bladder, the ureters and the kidneys. Most cases of UTI are caused by gram-negative bacteria especially E. coli. Escherichia coli is a normal flora in the gastro-intestinal system of humans but under certain conditions it might result in infections like urinary tract infection.

Most patients within health care facilities acquired UTIs after catheterization, instrumentation and/or operation in the lower urinary tract. Host factors also contribute for the risk of developing UTI in health care facilities.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 11 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



2. Wound infection and abscess

Wounds can be broadly classified as traumatic (accidental) wounds and surgical wounds. In developing countries such as Ethiopia where sporadic disputes and poverty prevail traumatic wounds are the more common ones. These traumatic wounds are most of the time infected (dirty) wounds. Dirty wounds are more prone to multiple micro-organism contamination.

Commonly isolated bacteria include:

- **Gram positive (both aerobic and anaerobic)**
 - Staphylococcus species
 - Streptococcus especially in deep abscesses
 - Clostridium species (Clostridium tetani, Clostridium perfringens)
- **Gram negative**
 - Escherichia coli
 - Pseudomonas aeruginosa

Special attention should be given to unclean burn wounds where microorganisms like clostridium species and pseudomonas are mostly isolated.

3. Air-borne infections

These are infections transmitted by inhalation of pathogenic (disease causing) micro-organisms. Pneumonia, which literally means acute infection of the lung, is one the leading causes of death, especially in children, in the developing world.

Though Streptococcus pneumoniae is a common cause at the community level, Staphylococcus aureus is known for its aggressiveness in health care facilities.

Tuberculosis is another common bacterial infection of the lung in health care facilities. This chronic infection of the lung is caused by Mycobacterium tuberculosis. With the advent of HIV, multi-drug resistance tuberculosis is becoming a problem that needs attention by all health workers in health care facilities.

4. Relapsing fever

This is a common acute febrile illness in situations where hygiene is poor (e.g. refugee camps) and is caused by spirochete called Borrelia recurrentis. Relapsing fever, if not identified early, can result in death. It needs joint management by all health worker teams in health care facilities.

II. Non-Bacterial causes of infections

Previously it was mentioned that bacteria play a major role in nosocomial infections. Though not as common as bacteria other micro-organisms like viruses, fungi, protozoas, helminthes and rickettsias also play a major role in causing infection in health care facilities.

The most common ones include:

1. HIV/AIDS

Acquired immunodeficiency syndrome (AIDS) is a systemic viral disease caused by a virus called HIV (Human Immuno Deficiency Virus). Though it is a chronic illness with long periods of latency (3-12yrs), initial acquisition of the virus can occur in health care facilities due to negligence and accidents in handling contaminated specimens and equipment.

2. Hepatitis

Viral Hepatitis is inflammation (infection) of the liver caused by viruses. The most common causes are viral hepatitis - hepatitis A, B, C, D, E, and G. Feco-Oral route is the commonest

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 12 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



way of transmission in cases of hepatitis A and E infection therefore they pose major problems in HCFs where sanitary conditions are very poor. On the other hand, contact with contaminated blood and blood products is a major way of transmission of hepatitis B, C, D, and G in HCF so they pose major problems in facilities where universal precaution is not properly followed.

3. Malaria

Malaria is an acute febrile illness most commonly found in the lowlands of Ethiopia. It is caused by the plasmodium species. Plasmodium falciparum and plasmodium vivax are the two most common etiologic agents in our country.

Because of the high prevalence of malaria in our country identification and management of cases, as well as the setting up of preventive measures, are important issues to be performed by members of the health care team. Health care facilities, which are located in malaria endemic areas, should follow general preventive measures (like window iron mesh, bed nets, draining of marshy areas in the vicinity of the HCF...) to control malaria infection in the HCF.

4. Fungal infections

Dermatophytes (tinea), Candida and aspergillus species are the most frequently encountered fungal pathogens in health care facilities. “Ringworm” infections of the skin caused by tinea species are common in poor hygienic set ups in health care facilities.

Candida albicans is the candida species most commonly identified in health care facilities. It is the normal flora of the skin and gastro intestinal tract.

Unsupervised or prolonged use of medication, which perhaps is due to a lack of awareness by health professionals, can result in the colonization of this pathogen in all body areas.

Aspergillus species are also common in the environment. Air is the principal route of transmission of aspergillus. Acquisition of this fungal species might result in chronic respiratory infections.

5. Helmenthic infection

Poor handling of stool specimens especially in the laboratory set-up might result in the common helmenthic infection. These common helmenthic infections include ascariasis, taeniasis, and hookworm infections. Infection due to entrobilus vermicuclaris, which is highly contagious, is also possible in health care facilities where hygienic care is poor.

6. Ricketsial disease

One important cause of acute febrile illness in health care facilities which arises due to poor sanitary conditions is typhus (caused by rickettsia typhi). The health care worker has to give due attention to hygienic control of the disease (like proper washing and ironing of bed sheets, pajamas, fumigation of wards...).

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 13 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Self-Check 1	Written Test
---------------------	---------------------

Directions: Answer all the questions listed below. Use the Answer sheet provided in the next page:(2 point each)

- A procedure made to prevent entry of microorganisms into any area of the body
 A. Asepsis
 B. Hand washing
 C. Sterilization
 D. Prevention
- A germicide that is used on skin or living tissue for the purpose of inhibiting or destroying microorganisms.
 A. Treatment
 B. Standard precaution
 C. Infection prevention
 D. Antiseptic
- A process that physically removes all visible dust, soil, blood or other body fluids from inanimate object as well as removing sufficient number of microorganisms
 A. Cleaning
 B. Colonization
 C. Contamination
 D. Boiling
- The most common or frequent risk that health care personnel exposed is from
 A. Tuberculosis
 B. Direct contact with blood and other body fluids
 C. Multidrug resistance microorganisms
 D. None of the above
- An agent is carried from a reservoir to a susceptible host by suspended air particles or by animate or inanimate intermediaries. This type of disease transmission is said to be
 A. Direct transmission
 B. Blood born disease
 C. Indirect transmission
 D. Direct contact

Note: Satisfactory rating - 5 points & Above Unsatisfactory - below 5 points

You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = <u> 5 </u>
Rating: <u> 10 </u>

Name: _____

Date: _____

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 14 of 139
---------------------------	----------------------	---------------------------------------	-----------------------	----------------



2. Standard precautions to prevent transmission of infection

2.1. Standard precaution

The use of standard precautions is the primary strategy for minimizing the transmission of healthcare-associated infections.

Because most people with blood borne viral infections such as HIV and HBV do not have symptoms, nor can they easily be recognized as such, Standard precautions are designed for the care of all person-patient, clients and staff regardless of whether or not they are actually infected.

Standard precautions combined the major features of Universal Precautions (UP) and body substances Isolation (BSI) and are based on the principle that all blood, body fluids, secretions, excretions except sweat, non-intact skin, and mucous membrane may contain transmissible infectious agents.

- Regarding precautions two-tier approach is currently employed. This includes
 - ✓ **STANDARD PRECAUTIONS** (which applies to all patients) and
 - ✓ **TRANSMISSION-BASED PRECAUTIONS** (which apply to patients with documented or suspected infection or colonization with certain micro-organisms).

The application of Standard Precautions during patient care is determined by the nature of the healthcare worker-patient interaction and the extent of anticipated blood, body fluid, or pathogen exposure.

For some interactions: for example:

- For performing vein puncture, only gloves may be needed
- For intubations use of gloves, gown, and face shield or mask and goggles is necessary

Standard Precautions

- are formerly known as universal precautions underpin routine safe practice, protecting both staff and clients from infection.
- are designed to reduce the risk of transmission of micro-organisms from both recognized and unrecognized sources of infection in the hospital.
- applies to all patients regardless of their diagnosis.
- shall be implemented when contact with any of the following are anticipated
 - ✓ Blood - All body fluids, secretions and excretions, with the exception of sweat regardless of whether or not they contain visible blood.
 - ✓ Non-intact skin (this includes rashes)
 - ✓ Mucous membranes

By applying standard precautions at all times and to all patients, best practice becomes second nature and the risks of infection are minimized. They include:

1. achieving optimum hand hygiene
2. using personal protective equipment
3. safe handling and disposal of sharps



4. safe handling and disposal of clinical waste
5. managing blood and bodily fluids
6. decontaminating equipment
7. achieving and maintaining a clean clinical environment
8. appropriate use of indwelling devices
9. managing accidents
10. good communication – with other health care workers, patients and visitors
11. training/education.

It is essential that standard precautions are applied at all times. This is because:

- people may be placed at risk of infection from others who carry infectious agents
- people may be infectious before signs or symptoms of disease are recognized or detected, or before laboratory tests are confirmed in time to contribute to care;
- people may be at risk from infectious agents present in the surrounding environment including environmental surfaces or from equipment
- there may be an increased risk of transmission associated with specific procedures and practices.

The three new areas of practices that have been added in Standard precautions are

1. respiratory hygiene and cough etiquette
2. safe injection practice and
3. use of masks for insertion of catheters or injection of material into spinal or epidural spaces via lumbar puncture procedures

Components of Standard Precautions:

- Hand hygiene, before and after every episode of patient contact
 - ✓ After touching blood, body fluids, secretions, excretions and contaminated items
 - ✓ Immediately after removing gloves
 - ✓ Between patient contacts
- the use of personal protective equipment
 - ✓ Appropriate use of gown/apron, mask, goggles and face shield
 - Use gloves for contact with blood, body fluids, secretions/excretions or contaminated items
 - Use gloves for contact with mucus membranes and non-intact skin
 - Use gown/apron to protect skin from blood or body fluid contact
 - Use gown/apron to prevent soiling of clothing during procedures that may involve contact with blood or any body fluids (secretions/excretions)
 - Use mask to protect mucous membranes of eyes, nose and mouth when contact with blood and body fluids is likely or possible
 - ✓ *During aerosol-generating processes in patients with suspected or confirmed infections transmitted by respiratory aerosols (e.g SARS), a fit-tested N95 or higher respiratory should be worn in addition to gloves, gown and face/eye protection*
- the safe use of needles and disposal of sharps
 - ✓ avoid recapping, bending, breaking, or hand manipulate used needles; if recapping is required, use a one-handed scoop technique only
 - ✓ Avoid removing used needles from disposable syringes

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 16 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



- ✓ Place used sharps in puncture-resistant container at point of use
- Patient resuscitation
 - ✓ Use mouthpiece, resuscitation bags or other ventilation devices to avoid mouth-to-mouth resuscitation
- Patient placement
 - ✓ Place patients who contaminate the environment or cannot maintain appropriate hygiene in private rooms
 - ✓ Place patients on airborne, droplet, contact precautions in appropriate rooms
- soiled patient-care equipment
 - ✓ handle soiled equipment in a manner to prevent contact with skin or mucus membranes and to prevent contamination of clothing or the environment
 - ✓ Clean reusable equipment prior to reuse
- routine environmental cleaning
 - ✓ Develop procedures for routine care, cleaning and disinfection of equipment and environmental surfaces, especially frequently touched surfaces in patient care area
- waste management
- Textile and laundry
 - ✓ Handle in a manner that prevents transfer of microorganisms to others and to the environment
- appropriate handling of linen.
 - ✓ Handle soiled linen to prevent touching of skin or mucous membranes
 - ✓ Do not pre rinse soiled linens in patient care areas
- Respiratory hygiene/cough etiquette
 - ✓ Instruct symptomatic persons to cover mouth/nose when sneezing /coughing or use tissue papers and dispose in no-touch receptacle
 - ✓ Observe hand hygiene after soiling of hands with respiratory secretions
 - ✓ Wear separation, >3feet if possible

Standard precautions should be used in the handling of: blood (including dried blood); all other body substances, secretions and excretions (excluding sweat), regardless of whether they contain visible blood; non-intact skin; and mucous membranes.

2.2. Transmission based precautions

Any infection prevention and control strategy should be based on the use of standard precautions as a minimum level of control. Transmission-based precautions are recommended as extra work practices in situations where standard precautions alone may be insufficient to prevent transmission. Transmission-based precautions are also used in the event of an outbreak (e.g. gastroenteritis); to assist in containing the outbreak and preventing further infection.

- Transmission-based precautions are used in addition to standard precautions, where the suspected or confirmed presence of infectious agents represents an increased risk of transmission.
- The application of transmission-based precautions is particularly important in containing multi-resistant organisms (MROs) and in outbreak management.
- Medical and dental procedures increase the risk of transmission of infectious agents. Effective work practices to minimize risk of transmission of infection related to

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 17 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



procedures require consideration of the specific situation, as well as appropriate use of standard and transmission-based precautions.

Transmission-based precautions should be tailored to the particular infectious agent involved and its mode of transmission. This may involve a combination of practices

Strategies for implementing transmission-based precautions

Transmission-based precautions may include one or any combination of the following:

- Allocating a single room with closing door to patient with a suspected or confirmed infection (isolation)
- Wearing specific personal protective equipment
- providing patient-dedicated equipment
- using a TGA-registered disinfectant with label claims specifying its effectiveness against specific infectious organisms
- Using specific air handling techniques
- Restricting movement both of patients and healthcare workers.

Contact precautions are used when there is known or suspected risk of direct or indirect contact transmission of infectious agents that are not effectively contained by standard precautions alone.

Droplet precautions are used for patients known or suspected to be infected with agents transmitted over short distances by large respiratory droplets.

Airborne precautions are used for patients known or suspected to be infected with agents transmitted person-to-person by the airborne route.



Self-Check 2	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

- The primary strategy for minimizing the transmission of healthcare-associated infections is the use of
 - Advanced treatment
 - Secondary prevention
 - Hand washing
 - Standard precautions**
- According to transmission based precaution, which one of the following is/are are used when there is known or suspected risk of direct or indirect contact transmission of infectious agents that are not effectively contained by standard precautions alone.
 - Droplet precautions
 - Contact precautions**
 - Airborne precaution
 - All of the above

Note: Satisfactory rating – 2.5 points points

Unsatisfactory – below 2

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Answer

NO	
1	
2	
3	
4	



Information Sheet 3	Minimizing Contamination of materials, equipment and instruments
----------------------------	--

1. Minimizing contamination of materials, equipment and instruments

The recommended infection prevention practices for minimizing microbial contamination of specific areas in healthcare facilities are

Procedure Area

- Limit traffic to authorized staff and patients at all times.
- Permit **only** the patient and staff performing and assisting with procedures in the procedure room (family members should be limited with obstetrical procedures).
- Patients can wear their own clean clothing.
- Staff should wear attire and personal protective equipment (PPE) according to procedures performed.
- Have a covered container filled with a 0.5% chlorine solution for immediate decontamination of instruments and other items once they are no longer needed.
- Have a leak proof, covered waste container for disposal of contaminated waste items (cotton, gauze, dressings) at point of use.
- Have a puncture-resistant container for safe disposal of sharps (e.g., used suture needles, hypodermic needles and syringes, and disposable scalpel blades) at point of use.
- Have storage space in procedure rooms for clean, high-level disinfected and sterile supplies. (Storage shelves should be enclosed to minimize dust and debris collecting on stored items.)

Surgical Unit

The surgical unit is often divided into four designated areas, which are defined by the activities performed in each—unrestricted area, transition zone, semi restricted area and restricted area. Environmental controls and use of surgical attire increase as one moves from unrestricted to restricted areas. Moreover, staff with respiratory or skin infections and uncovered open sores should not be allowed in the surgical unit

Unrestricted Area

This area is the entrance from the main corridor and is isolated from other areas of the surgical unit. This is the point through which staff, patients and materials enter the surgical unit.

Transition Zone

This area consists primarily of dressing rooms and lockers. It is where staff put on surgical attire that allows them to move from unrestricted to semi restricted or restricted areas in the surgical unit. Only authorized staff should enter this area.

Semi restricted Area

This is the peripheral support area of the surgical unit and includes preoperative and recovery rooms, storage space for sterile and high-level disinfected items, and corridors leading to the restricted area. Support activities (e.g., instrument processing and storage) for the operating room occur here.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 20 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



- Limit traffic to authorized staff and patients at all times.
- Have a work area for processing of clean instruments.
- Have storage space for clean and sterile or high-level disinfected supplies with enclosed shelves to minimize dust and debris collecting on stored items.
- Have doors limiting access to the restricted area of the surgical unit.
- Staff who work in this area should wear surgical attire and a cap.
- Staff should wear clean, closed shoes that will protect their feet from fluids and dropped items

Restricted Area

This area consists of the operating room(s) and scrub sink areas.

- Limit traffic to authorized staff and patients at all times.
- Keep the door closed at all times, except during movement of staff, patients, supplies and equipment.
- Scrubbed staff must wear full surgical attire and cover head and facial hair with a cap and mask.
- Staff should wear clean, closed shoes that will protect their feet from fluids and dropped items.
- Masks are required when sterile supplies are open and scrubbed staff are operating.
- Patients entering the surgical unit should wear clean gowns or be covered with clean linen, and have their hair covered.
- Patients do not need to wear masks during transport (unless they require airborne precautions).

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 21 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Self-Check 3	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. Procedure area limit traffic to authorized staff and patients at all times **True**
False(2 point each)
2. The surgical unit is often divided into four designated areas, which are defined by the activities performed in each **True** False(2 point each)
3. Unrestricted area is the entrance from the main corridor and is isolated from other areas of the surgical unit. **True** False(2 point each)
4. Semi restricted area is the peripheral support area of the surgical unit and includes preoperative and recovery **True** False(2 point each)
5. Restricted area consists of the operating room(s) and scrub sink areas. **True**
False(2 point each)
- 6.

Note: Satisfactory rating - 12 points Unsatisfactory – below 12 points
You can ask your trainer for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Answer

No	
1	True
2	True
3	True
4	True
5	True



Operation Sheet 2	Standard precautions
--------------------------	----------------------

1. Discuss and apply standard precautions at all times and to all patients, best practice becomes second nature and the risks of infection are minimized. They include:



LAP Test	Practical Demonstration
-----------------	--------------------------------

Name: _____ Date: _____

Time started: _____ Time finished: _____

Instructions: Given necessary templates, workshop, tools and materials you are required to perform the following tasks.

Task 1: Identify standard precautions to prevent transmission of infection



This learning guide is developed to provide you the necessary information regarding the following **content coverage and topics** –

Identify and respond to infection risks

- identification of Infection risks
- documenting and reporting any Activities that put clients or workers in to risk
- giving response to situations that pose an infection risk
- following risk control Procedures
- precaution exposure to blood or other body fluids
- placing signs when and where appropriate
- removing Spills in accordance with the policies and procedure

This guide will also assist you to attain the learning outcome stated in the cover page. Specifically, upon completion of this Learning Guide, you **will be able to** –

- identify **Infection risks** and an appropriate response implemented within own role and responsibility
- document and report activities and tasks that put clients and/or other workers at risk are
- given appropriately response to situations that pose an infection risk in accordance with the policies and procedures of the organization
- follow **procedures for risk control** and risk containment for specific risks
- **followed protocols for care following exposure to blood or other body fluids as required**
- **placed signs are when and where appropriate**
- **remove spills in accordance with the policies and procedures of the organization**

Learning Instructions:

1. Read the specific objectives of this Learning Guide.
2. Follow the instructions described in number **3 to 16**.
3. Read the information written in the “Information Sheets 1”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
4. Accomplish the “Self-check 1” **in page 8**.
5. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 1).
6. If you earned a satisfactory evaluation proceed to “Information Sheet 2”. However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 1.
7. Submit your accomplished Self-check. This will form part of your training portfolio.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 25 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



8. Read the information written in the “Information Sheet 2”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
9. Accomplish the “Self-check 2” in page 16.
10. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 2).
11. Read the information written in the “Information Sheets 7”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
12. Accomplish the “Self-check 3” in page 19.
13. Ask your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 3).
14. If you earned a satisfactory evaluation proceed to “Operation Sheet 1” in page 21; However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 3.
15. Read the “Operation Sheet 1 and try to understand the procedures discussed.
16. Do the “LAP test” in page 22 (if you are ready). Request your trainer to evaluate your performance and outputs. Your trainer will give you feedback and the evaluation will be either satisfactory or unsatisfactory. If unsatisfactory, your trainer shall advice you on additional work. But if satisfactory you can proceed to Learning Guide #2.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 26 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Information Sheet-1	identification of Infection risks
----------------------------	-----------------------------------

1. Identification *Infection risks*

1.1. Infection risks

RISK ASSESSMENT

A **risk** is the chance, high or low, that a hazard will cause harm, injury or ill health, or the likelihood, or possibility, which harm (injury, illness, death, damage etc) may occur from exposure to a hazard

A **hazard** is a situation or item that could cause harm. Risks and hazards should be monitored so they are minimized, protecting the health and wellbeing of all workers and clients.

Risk Assessment: Is defined as the process of assessing the risks associated with each of the hazards identified so the nature of the risk can be understood. This includes the nature of the harm that may result from the hazard, the severity of that harm and the likelihood of this occurring.

Risk Control: Taking actions to eliminate health and safety risks so far as is reasonably practicable. Where risks cannot be eliminated, then implementation of control measures is required, to minimize risks as far as is reasonably practicable. A hierarchy of controls has been developed and is described below to assist in selection of the most appropriate risk control measure/s.

Monitoring and Review: This involves ongoing monitoring of the hazards identified, risks assessed and risk control processes and reviewing them to make sure they are working effectively.

Health care facilities are ideal settings for the transmission of nosocomial infections in the following ways:

1. Invasive procedures have the potential to introduce microorganisms.
2. Service providers and support staff are constantly performing clinical procedures or other activities (susceptible host).
3. Clients receiving services may be harboring microorganisms.

Who Is at Risk of Infections?

Service providers and support staff: Health care personnel, including support staff (e.g., housekeeping, laundry staff, and maintenance), who work in health care settings are at risk of exposure to serious, potentially life-threatening infections such as HIV, hepatitis B virus (HBV), and hepatitis C virus (HCV).

All workplaces are legally obliged to have processes in place to identify infection risks, as well as policies and procedures to provide workers with guidance on how they should respond to such risks.

Every person in the workplace, from trainee personal care workers through to senior management, has work health and safety (WHS) obligations. These obligations include taking all reasonable steps to prevent the spread of infection. Knowledge of infection risks and appropriate responses is an essential part of meeting WHS requirements.

Transmission of diseases in health care settings can occur from patient to health care worker, between patients, or from health care worker to patients.

Clients: Clients are at risk of infections when:

- Service providers do not wash their hands before and after providing care to each client and before and after every procedure (cross-contamination).
- Service providers do not adequately prepare clients before clinical procedure.
- Service providers do not correctly process instruments and other items used in clinical procedures.
- Medical waste is disposed of inappropriately.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 27 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



The community: Members of the community are also at risk of infections, particularly from inappropriate disposal of medical waste, such as contaminated sharps materials

Self-Check 1	Written Test
---------------------	---------------------

Directions: Answer all the questions listed below. Use the Answer sheet provided in the next page:

1.the chance, high or low, that a hazard will cause harm, injury or ill health, or the likelihood, or possibility, which harm(2 point each).
A. **Risk** B. Hazard C. Risk Assessment D. Risk control
2. Of the following who is/are at risk of infection (2 point each).
A. Client
B. Health worker
C. Community
D. All of the above
3. Which of the following is the way for the transmission of nosocomial infections(2 point each).
A. Invasive procedures have the potential to introduce microorganisms.
B. Service providers and support staff are constantly performing clinical procedures
C. Clients receiving services may be harboring microorganisms
D. ALL

Note: Satisfactory rating - 16 points Unsatisfactory - below 16 points

You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Answer

No	
1	A
2	D
3	D
4	



Information Sheet 2	Document and report activities and tasks that put clients and/or other workers at risk are
----------------------------	--

2. Document and report activities and tasks that put clients and/or other workers at risk

2.1. Activities and tasks that put clients and/or other workers at risk

Although healthcare workers (HCWs) are essential to the health of the world's population, they, themselves, are often put in physical jeopardy. Globally, HCWs are exposed each day to a variety of health and safety hazards, including:

- Biological, (e.g., pathogens such as Human Immunodeficiency Virus (HIV), Hepatitis B Virus (HBV), Hepatitis C Virus (HCV), Ebola, Mycobacterium Tuberculosis (MTB), SARS virus and Neisseria Meningitis)
- Sharp's injuries
- Ergonomic, (e.g., heavy lifting)
- Physical, (e.g., slips, trips and falls)
- Psychosocial, (e.g., violence and stress)
- Chemical, (e.g., chlorine, gutaraldehyde, ethylene oxide)
- Radiological and nuclear

Safe work leads to worker well-being and retention, increased productivity and economic best outcomes. Risks and hazards are fluid and need monitoring and adjustments made to the appropriate safety plans and processes

The risk to staff arises:

- from sharps and hollow needles;
- splashing of conjunctivae and mucous membranes with contaminated blood and body fluids;
- heavy contamination of broken skin, e.g. cuts, dermatitis etc.;
- handling of large quantities of blood and body fluids without protective clothing.

The risk to patients arises from: -

- use of recycled hollow needles and syringes;
- contaminated blood transfusion;
- heavy soiling of the environment;
- poor ward facilities and cleaning



Self-Check 2	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. The **risk to staff** arises from
 - A. sharps and hollow needles
 - B. splashing of conjunctivae and mucous membranes with contaminated blood and body fluids
 - C. heavy contamination of broken skin
 - D. handling of large quantities of blood and body fluids without protective clothing
 - E. ALL**

2. The risk to patients arises from: -
 - A. use of recycled hollow needles and syringes
 - B. contaminated blood transfusion
 - C. heavy soiling of the environment
 - D. poor ward facilities and cleaning
 - E. ALL**

Note: Satisfactory rating - 12 points Unsatisfactory – below 12 points
 You can ask your trainer for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Answer

No	
1	
2	
4	



Information Sheet 3	Given appropriately response to situations that pose an infection risk in accordance with the policies and procedures of the organization
----------------------------	---

3. Given appropriately response to situations that pose an infection risk in accordance with the policies and procedures of the organization

3.1. Response to infection risks

To successfully identify and respond to infection risks we must understand and follow safe work practices that prevent the transmission of infections.

As a health worker it is also your responsibility to follow recommended procedures in your workplace and take adequate precautions to protect yourself from injury and infection.

Risk management is the process of making health care safer for the patient, staff and visitors by identifying hazards in the workplace and taking action to minimize their harm wherever possible. There are a number of steps in the risk management process:

- *identifying the hazard*
- *assessing the risks*
- *using control measures.*

Identifying a hazard

- A hazard is anything with the potential to cause harm to you, the patients, your co-workers or visitors to the work area. In the sterilization setting this includes chemicals, sharps such as needles, soiled instruments, power, water, steam, noise, and heat.
- In developing procedures or buying new equipment, identify these risks early so that work practices can be developed that ensure the hazard is eliminated as much as possible. Regular safety inspections and audits can help identify and manage hazards.
- All employees, patients, volunteers, contractors and visitors that enter the work place have a responsibility to behave in a safe and responsible manner and report any hazards or near accidents.

Assessing the risks

- It is important to assess the risks associated with each hazard to determine how it can be eliminated.
 - ✓ Is there a high risk of injury or is the hazard a result of a combination of unusual circumstances that may never re-occur?
- Budgets are limited in health care settings, so it is important to look at all the options for dealing with a hazard. You should also document the process to seek additional support for action.

Control measures

- The more serious the consequence, the more urgent it is for the risk to be dealt with and eliminated immediately. If the risk cannot be eliminated it may be possible to circumvent the risk of injury by changing practice. The last alternative is to use some form of personal protection when exposed to the hazard.
- When deciding on control measures this should be a team effort so that management and staff work together. The control measures should not impose another risk.

Monitoring control measures

Once control measures have been implemented it is important to monitor and re-evaluate practice to ensure compliance with new practice

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 31 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Self-Check 3	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page

6. The first step in risk management process is
- A. Identifying the hazard
 - B. Assessing the risks
 - C. Using control measures
 - D. None of the above

Note: Satisfactory rating - 09 points Unsatisfactory - below 09 points
 You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Answer

1. _____



Information Sheet 4

Follow **procedure for risk control** and risk containment for specific risks

2.2. Procedures for risk control

Strategies to identify risks

Strategies for identifying risks vary. Risk identification can be proactive or reactive. The following contains information about proactive and reactive strategies that can help management and workers identify hazards that present risks to health and safety.

Proactive strategies

- A proactive strategy is one carried out to prevent an accident or incident; for example, implementing processes to identify hazards and risks. Two examples are a job safety analysis (JSA) and an audit.
- A JSA contains information about how a job should be carried out, types of risks and control measures.
- Providers should carry out regular internal audits to check that the control measures for infection and other risks are being implemented. External bodies such as state and territory WHS authorities can also carry out audits to check that safety controls are appropriate.

Reactive strategies

- A reactive approach to risk identification involves reviewing accidents and incidents through measures such as report forms and data, as well as establishing consultation processes such as workplace health and safety committees (HSCs).
- Incident and accident report forms are filled out after any incident or accident. Data from these forms is used by HSCs, WHS officers and managers to identify hazards.
- Committees, team meetings and other forums give staff the chance to discuss infection control risks and provide suggestions for policy and procedure improvements.

Carrying out a risk assessment

Once a hazard has been identified, you need to conduct an assessment of the risk of injury, harm or damage. An example of a risk is the likelihood of a hazard resulting in an injury or disease, together with the seriousness of the injury or disease.

The five steps in carrying out a risk assessment are shown here.

Risk assessment steps

1. Evaluate the likelihood of an injury or illness occurring and the likely severity of any injury or illness
2. Review health and safety information relevant to the hazard such as incident reports, SDSs, results of workplace monitoring and inspections and supplier information
3. Identify factors that contribute to the risk such as the physical layout of the workplace, the knowledge, skills and experience of workers, and existing work practices
4. Identify actions necessary to eliminate or control the risk
5. Complete any relevant records

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 33 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------

**Self-Check 4****Written Test**

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page

1. A proactive strategy is one carried out to prevent an accident or incident. **True** False(2 point each)
2. A reactive approach to risk identification involves reviewing accidents and incidents through measures. **True** False(2 point each)
- 3.

Note: Satisfactory rating - 09 points Unsatisfactory - below 09 points
You can ask you teacher for the copy of the correct answers.



Information Sheet 5

Protocols are followed for care following exposure to blood or other body fluids as required

2.3. Protocols for care of blood borne exposures

Definition: Blood borne Pathogen Exposure - a percutaneous injury (e.g., a needle stick or cut with a sharp object) or contact of mucous membrane or non-intact skin (e.g., exposed skin that is chapped, abraded, or afflicted with dermatitis) with blood, tissue, or other body fluids that are potentially infectious. In addition to blood and body fluids containing visible blood- semen, vaginal secretions, cerebrospinal fluid, synovial fluid, pleural fluid, peritoneal fluid, pericardial fluid, and amniotic fluid are also considered potentially infectious. Urine or gastric contents without visible blood are not considered potentially infectious.

Blood is the most infectious body fluid for the transmission of HIV, HBV and HCV. If the exposure incident involved a body fluid capable of transmitting any of the viruses (HIV, HBV or HCV

Exposure to blood or body substances may be defined as direct contact with blood or other body substances through broken skin, mucous membranes (eyes, nose or mouth) or needle stick injury

Health care workers (HCW) are at risk of acquiring infection through occupational exposure. Hospital employees can also transmit infections to patients and other employees. Thus, an employee's health programme must be in place to prevent and manage infections in hospital staff

Occupational injuries may be divided into:

- a) Percutaneous exposure (from needles, instruments, bone fragments, human bite which penetrates the skin layer, etc.);
- b) Exposure via broken skin (exposed skin that is chapped, abraded, or afflicted with dermatitis etc.) with blood, tissue, or other body fluids that are potentially infectious; and
- c) Exposure via mucous membranes including the eye

Specific post-exposure policies must be developed, and compliance ensured for a number of infectious diseases for example: human immunodeficiency virus (HIV), viral hepatitis, severe acute respiratory syndrome (SARS), varicella, rubella and tuberculosis. Health care workers with infections should report their illnesses/incident to staff clinics for further evaluation and management

Hepatitis B virus (HBV), hepatitis C virus (HCV) and the human immunodeficiency virus (HIV) constitute well-recognized occupational risks for healthcare workers (HCWs). Avoiding occupational blood exposure by the adherence to principles of standard precautions through the use of appropriate work practices and personal protective equipment is a cornerstone for preventing transmission of these blood-borne pathogens (BBP) in the health-care setting.

In general, the risk of viral transmission after a percutaneous injury is highest for HBV, followed by HCV and HIV.

Occupational exposure is serious and every effort should be taken to prevent its occurrence. However, accidents may still happen and if so, risk assessment and counseling constitutes the basis of post exposure management. Appropriate post exposure prophylaxis (PEP) should be provided using a case-by-case evaluation approach.

Each healthcare institution should have personnel responsible for the Sharps Prevention Program

Types of exposure

1. Percutaneous Injury

- Puncture or laceration of the skin that penetrates into or below the dermis.
- For the purposes of this protocol, a percutaneous exposure to blood/body fluids which has one or more of the following factors present will be defined as a more severe exposure .
 - Deep percutaneous injury

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 35 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



- Visible blood present on the device associated with the exposure
- Exposure from a procedure which involved a needle placed directly into the Source's vein or artery
- Large-bore hollow needle
- A percutaneous exposure which has none of the above characteristics will be defined as a less severe exposure (e.g., superficial injury, no visible blood present on device associated with the exposure, procedure from which exposure resulted did not involve a needle being placed directly into the Source's vein or artery, solid needle)

2. Mucous Membrane and Non-intact Skin Exposures

- a. Mucous Membrane Exposure: When blood/body fluids come into contact with mucous membranes (e.g., eyes, oral cavity)
- b. Non-intact Skin Exposure: When blood/body fluids come into contact with an open wound or exposed skin that is chapped, abraded or non-intact because of dermatitis

A larger volume of blood/body fluid is associated with increased transmission risk for mucous membrane and non-intact skin exposures. For the purposes of this protocol, a mucous membrane or non-intact skin exposure involving a major splash of blood/body fluids will be defined as a large volume exposure. Exposures involving lesser amounts (e.g., only a few drops of fluid) will be defined as a small volume exposure.

3. **Human Bites:** Human bites may occur in both occupational and non-occupational settings. The person bitten has a potential percutaneous exposure and the person who was the biter has a potential mucous membrane exposure. Therefore, an individual who bites may be both the Source and Exposed in bite incidents.

- a. As HBV is present in saliva at concentrations 1,000 to 10,000 times less than in blood, for the purposes of post-exposure prophylaxis, generally only exposures to saliva containing visible blood would be considered for HBV PEP (such as deep bites associated with bleeding in the mouth of the biter)

4. Exposures to Blood/Body Fluids Obtained Through Cuts, Nosebleeds, Physical Assaults, Sports Injuries

5. **Consensual Sex** (Serodiscordant Partners) HBV and/or HIV PEP should be considered for the following unprotected (e.g., condom breakage) sexual exposures where the Source is known to be positive for the respective viruses

For example

The route of transmission for hepatitis B virus is through body substances such as blood and blood products, saliva, cerebrospinal fluid, peritoneal, pleural, pericardial and synovial fluid, amniotic fluid, semen and vaginal secretions and any other body fluid containing blood.

The risk of a health care worker acquiring HIV after a needle stick or other "sharps" injury is less than 0.5%.

Risk reduction must be undertaken for all blood borne pathogens, including:

- adherence to standard precautions using personal protective equipment
- appropriate use of safety devices and a needle disposal system to limit sharps exposure.
- Training for health care workers in safe sharps practice should be ongoing.
- Information on preventive measures must be provided to all staff with potential exposure to blood and blood products.
- Policies which are in keeping with the local and national guidelines must include
 - screening of patients
 - disposal of sharps and wastes
 - protective clothing
 - managing inoculation accidents
 - sterilization and disinfection.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 36 of 139
---------------------------	----------------------	---------------------------------------	-----------------------	----------------



- Hospital policy must include measures to obtain serological testing of source patients promptly where necessary, usually with the patient's informed consent.
- Post exposure prophylaxis should be started as per local or national guidelines.
- In case of hepatitis B, immunization is the best way of preventing transmission to health care staff.
 - All HCWs at risk must be vaccinated.
 - Staff infected with blood-borne pathogens may transmit these infections to patients and require careful evaluation with respect to their duties. This status should not be used as cause for discrimination
- If a staff member has been exposed to **Tuberculosis** (TB) they should report to the Infection Control Practitioner or the Staff Health Nurse depending on the hospital protocol for health care worker exposures.
- Health care workers in close respiratory contact with cases such as **meningococcal meningitis** should receive chemoprophylaxis with ciprofloxacin or an effective alternative agent. Close respiratory contact with the patient includes mouth-to-mouth contact, sharing of drink containers or cigarettes

Sharp injuries Needle

- stick injuries are the most common of sharps injuries, although other contaminated sharp instruments may also cause injuries.
- The majority of reported NIs involved hollow-bore needles (55-62%), and recapping was the most common behavior associated with NI. Overall, more than half of percutaneous injuries involving hollow-bore needles were potentially preventable through safer work practices or technologies.
- HCWs should prevent skin penetrating injuries by wearing appropriate clothing, shoes and personal protective equipment (PPE) where required. As a break in the skin can allow direct contact with blood and body substances these should be protected by keeping open wounds covered e.g. with a waterproof dressing or with appropriate clothing.
- Skin penetrating injuries can introduce infectious agents directly into the blood stream, e.g. tetanus and blood borne viruses such as hepatitis B, hepatitis C and HIV. It is very important that skin penetrating injuries are minimized e.g. through safe handling and disposal of sharps
- All health care workers with potential exposure should be vaccinated.
 - For other personnel, the risk of hepatitis B, hepatitis C and HIV infection should be assessed and appropriate immunization or chemo prophylactic steps taken.
 - Immediate treatment of such injuries should encourage
 - washing thoroughly with running water and an antiseptic solution.
 - Consult the infection control team for further advice.
- An incident reporting system should be in place. It should not be seen as punitive/disciplinary; active support by managers should encourage prompt and accurate reporting.

Exposures for which PEP is indicated

- Break in the skin by a sharp object (including hollow-bore, solid-bore, and cutting needles or broken glassware) that is contaminated with blood, visibly bloody fluid, or other potentially infectious material, or sharp objects had been in the source patient's blood vessel.
- Bite from a patient with visible bleeding (in the mouth) and which causes bleeding in the exposed worker.
- Splash of blood, visibly bloody fluid, or other potentially infectious material to a mucosal surface (mouth, nose, or eyes).

Remember, Health care workers should have immediate access to post exposure prophylaxis (PEP) , 24 hours a day, 7 days a week to be freely dispensed by any hospital (private or public), regardless of

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 37 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



the location or type of work they do. The minimum care following potential exposure to HIV should be risk assessment and, if deemed necessary, the first dose of PEP medication

General procedures

First Aid – when an exposure incident occurs, implement first aid

1. Following any exposure, the wound should be washed immediately and thoroughly with soap and water, flush the eyes with running water immediately following a bodily fluid splash. Alcohol, hydrogen peroxide, Betadine or other chemical cleansers are best avoided. Wound should not be squeezed or sucked.
2. For mucosal contact e.g. spillage into the conjunctivae, the exposed area should be immediately flushed with plenty of clean running water.
3. The exposed HCW is responsible for reporting the exposure incident to his/her supervisor and should then seek immediate medical advice for proper wound care and post-exposure management.
 - The following information should be recorded in the exposed worker's confidential medical record:
 - i. details about the source patient (e.g. name, NRIC No, diagnosis and any relevant information)
 - ii. date, time and place of the exposure
 - iii. details of the procedure being performed
 - iv. use of protective equipment at the time of the exposure
 - v. the type, severity, and amount of fluid to which the worker was exposed
4. The health care worker should be tested for HIV antibody, HCV, HBV antigen and antibody
5. The source patient's blood (if available) should be tested for HIV, HCV & HBV

Reporting

- All institutions should have a mechanism in place for reporting and managing of sharp injuries and mucosal exposure in the occupational setting. HCWs must know the reporting process to facilitate quick and smooth flow so as to allow the attending physician to evaluate the risk of exposure and provide prompt appropriate post exposure treatment
- In addition, a surveillance system of exposure events should be available to avoid similar incidents from occurring in the future.

Counseling

Until the risk of infection is ruled out, advice should be given to the exposed staff to refrain from donating blood, plasma, organs, tissue or semen. The use of condom during sexual intercourse should also be advised. A place for psycho-social support is clearly indicated

In general

Healthcare workers should practice the following:

- Follow safe work practices at all times
- Be familiar with employer's written departmental policies
- Know the potential health and safety hazards of the job and protective measures by participating in appropriate occupational health and safety training programs
- Use personal protective equipment (PPE) as trained and report any changes in personal medical condition that would require a change in status as to wearing PPE
- Know how to report unsafe working conditions
- Report any work-related injury or illness to supervisor
- Participate in accident and injury investigations
- Know what to do in an emergency – Participate in health and safety committees (when available) can be an important way to improve conditions on the job such as: – Provide a forum for employees and management to work together to solve health and safety problems

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 38 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



- Help prevent injury and illness on the job i.e. conduct regular walk-a-round inspections to identify potential health and safety hazards
- Increase awareness of health and safety issues among employees, supervisors, and managers i.e. analyze injury data, accident reports and report trends
- Develop strategies to make the work environment safe and healthy

Self-Check 5	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

- Blood is the most infectious body fluid for the transmission of (2 point each)
 - HIV
 - HBV
 - HCV
 - ALL**
- Puncture or laceration of the skin that penetrates into or below the dermis is(2 point each)
 - Mucous Membrane exposure
 - Human bites
 - Non-intact Skin Exposures
 - Percutaneous Injury**
- Risk reduction must be undertaken for all blood borne pathogens, including(2 point each)
 - Post exposure prophylaxis should be started
 - adherence to standard precautions using personal protective equipment
 - appropriate use of safety devices and a needle disposal system
 - ALL**
- In general healthcare workers should practice (2 point each)
 - Follow safe work practices at all times
 - Be familiar with employer’s written departmental policies
 - Report any work-related injury or illness to supervisor
 - Participate in accident and injury investigations
 - ALL**

Note: Satisfactory rating - 09 points Unsatisfactory - below 09 points
 You can ask you teacher for the copy of the correct answers.

<p>Information Sheet 6</p>	<p><i>Appropriate signs are placed when and where appropriate</i></p>
-----------------------------------	---

Appropriate signs are placed when and where appropriate

DANGER SIGN

- Used where an immediate hazard exists



CAUTION SIGN

- Used to warn against potential hazards or to caution against unsafe practices



SAFETY INSTRUCTION SIGNS

- Used where there is a need for general instruction and suggestions relative to safety measures



ACCIDENT PREVENTION TAGS

ACCIDENT PREVENTION TAGS

- Used for temporary use only, to specify precautions and safety needs



The Official Presentation Material from the Ministry of Education
CORE PROCESS 2. OUTCOME-BASED TRAINING
Module 4. Managing Training Facilities
23 May, 2011

ecbp

DANGER TAG

- Placed only where an immediate hazard exists



CAUTION TAG

- Used to warn against potential hazards or to caution against unsafe practices



OUT OF ORDER TAG

- Used to identify pieces of or equipment that are out of order



Flammable chemicals



Corrosive chemicals



Explosive chemicals




Summary of universal hazardous symbols



60

Summary of ...



61



Self-Check 5	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.


7. This picture is a sign for

- A. Biohazard
- B. Cytotoxic
- C. Biological item
- D. Waste



8. This picture is a sign for

- A. Irritant
- B. Cytotoxic
- C. Biological item
- D. Waste



9. This picture is a sign for

- A. Toxic
- B. Cytotoxic
- C. Biological item
- D. Waste





Information Sheet 7

remove **Spills** in accordance with the policies and procedures of the organization

2.4. Spills removal

Basic Principles:

- Spills of blood or body substances are to be dealt with as soon as possible. Isolate or restrict access to the area if there is an unavoidable delay.
- Standard precautions apply – assume all blood and body substances are potentially infectious and cover cuts, maintain hand hygiene and use appropriate PPE.
- Cover the spill, where applicable, to prevent the generation of splashes and aerosols from the spilled substance - o e.g. granular formulation such as vomit control o use a scraper and pan to remove the absorbed material
- after removing the bulk of the spill, clean the area thoroughly, rinse and dry.
- clean non-disposable cleaning equipment thoroughly after use, rinse and store dry.

Small Spills: e.g. spots or drops of blood and other small spills up to 10cm diameter. - wipe the area immediately with paper toweling - clean with warm water and detergent followed by rinsing - dry the area (as wet areas attract contaminants) - a sanitiser (e.g. alcohol wipe) can be used on the area after cleaning.

Large Spills: e.g. greater than 10 cm diameter.

Wet area – e.g. bathroom with a floor drain –

- wash carefully into the sewerage system using copious amounts of water, taking care to avoid splashes - clean the area with mop and bucket of warm water and detergent
- clean the bucket and mop thoroughly after use using warm soapy water and store dry.
- Carpet - contain and clean with warm water and detergent - do not use disinfectant.

Equipment

- Equipment (mop, bucket and cleaning agents) is to be readily available in a location known to all. Prepare for a range of likely occurrences at your location considering:
 - the nature of the spill (e.g. sputum, vomit, faeces, urine, blood or laboratory culture)
 - the germs most likely to be involved in these different types of spills (e.g. gastrointestinal germs associated with spills of vomit and diarrhoea)
 - the size of the spill
 - the type of surface (e.g. carpet or impervious flooring)
 - the location e.g. whether the spill occurs in a contained area such as a toilet cubicle or in a high traffic area such as a hallway or while in a public place such as on an excursion.

A portable ‘spills kit’ can be made up to manage likely spills for the area/activity e.g.

- a large (10 L) reusable plastic container or bucket with fitted lid, containing the following items
 - leak proof bags and containers for disposal of waste material
 - roll(s) of paper towel to contain and cover a spill
 - a designated, sturdy scraper and pan for spills (similar to a ‘pooper scooper’/dust pan)
 - sachets of a granular formulation containing 10,000 ppm available chlorine or equivalent (each sachet should contain sufficient granules to cover a 10-cm diameter spill) e.g. vomit control - disposable latex, vinyl or nitrile gloves suitable for cleaning - eye protection (disposable or reusable)
 - a plastic apron

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 45 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



- a respiratory protective device such as a disposable P2 respirator (for protection against inhalation of powder from the disinfectant granules, or aerosols, which may be generated from high-risk spills during the cleaning process).

Self-Check 6	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. Spills of blood or body substances are to be dealt with as soon as possible **True** False(2 point each)
2. **Small Spills** wipe the area immediately **True** False(2 point each)
3. Equipment (mop, bucket and cleaning agents) is to be readily available in a location known to all **True** False(2 point each)

**Operation Sheet 3****Prepare tools and equipment for identification and measuring Muda.**

1. Discuss and plan to prepare tools and equipment for Muda identification.
2. Prepare tools and equipment for Muda identification.



LAP Test	Practical Demonstration
-----------------	--------------------------------

Name: _____ Date: _____

Time started: _____ Time finished: _____

Instructions: Given necessary templates, workshop, tools and materials you are required to perform the following tasks.

Task 1: Identify and prepare tools and equipment for measuring and identification of Muda.



Instruction Sheet 3

LG08: Maintain personal hygiene

This learning guide is developed to provide you the necessary information regarding the following content coverage and topics –

Identify and respond to infection risks

- washing hands before and after contact client and any activity
- Hand washing procedures
- Implementation of *hand care* procedures
- covering Cuts and abrasions with water-proof dressings

This guide will also assist you to attain the learning outcome stated in the cover page. Specifically, upon completion of this Learning Guide, you will be able to –

- maintain **Hand hygiene** by washing hands before and after client contact and/or after any activity likely to cause contamination
- follow Hand washing procedure
- implement **Hand care** procedure
- covered Cuts and abrasion with water-proof dressings and changed as necessary

Learning Instructions:

1. Read the specific objectives of this Learning Guide.
2. Follow the instructions described in number **3 to 16**.
3. Read the information written in the “Information Sheets 1”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
4. Accomplish the “Self-check 1” **in page 8**.
5. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 1).
6. If you earned a satisfactory evaluation proceed to “Information Sheet 2”. However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 1.
7. Submit your accomplished Self-check. This will form part of your training portfolio.
8. Read the information written in the “Information Sheet 2”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
9. Accomplish the “Self-check 2” **in page 16**.
10. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 2).
11. Read the information written in the “Information Sheets 4”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 49 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



12. Accomplish the “Self-check 3” in page 19.
13. Ask your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 3).
14. If you earned a satisfactory evaluation proceed to “Operation Sheet 1” in page 21; However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 3.
15. Read the “Operation Sheet 1 and try to understand the procedures discussed.
16. Do the “LAP test” in page 22 (if you are ready). Request your trainer to evaluate your performance and outputs. Your trainer will give you feedback and the evaluation will be either satisfactory or unsatisfactory. If unsatisfactory, your trainer shall advice you on additional work. But if satisfactory you can proceed to Learning Guide #2.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 50 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Information Sheet-1	maintain Hand hygiene by washing hands before and after client contact
----------------------------	---

3.1. Hand hygiene

Why Hand Hygiene Is Important

Hand hygiene is a general term referring to any action of hand cleansing. It includes care of the hands, nails, and skin.

Proper hand hygiene is a key component in minimizing the spread of disease and in maintaining an infection-free environment. Hand hygiene significantly reduces the number of disease-causing microorganisms on hands and arms and can minimize cross-contamination (e.g., from health worker to patient).

It is the most important way to reduce the spread of infections in the health care setting. Hand hygiene practices such as hand washing and surgical hand scrubbing are intended to prevent hand-borne infections by removing dirt and debris and inhibiting or killing microorganisms on skin. This includes not only most of the organisms acquired through contact with patients and the environment, but also some of the permanent ones that live in the deeper layers of the skin.

Studies indicated that failure to perform appropriate hand hygiene is considered to be the leading cause of health care-associated infections and the spread of multidrug-resistant microorganisms and has been recognized as a significant contributor to outbreaks (Boyce and Pittel 2002).

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 51 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



self-Check 1	Written Test
---------------------	---------------------

Directions: Answer all the questions listed below. Use the Answer sheet provided in the next page:

10. From the various hand hygiene practices available, which one of the following remains the most common and the most important when hands are visibly soiled
- A. The use of soap and water
 - B. The use of medicated soap and water
 - C. Boiling water and salt
 - D. Mixture of alcohol, soap and water(10 points)
11. Which of the following is TRUE about Hand hygiene
- A. any action of hand cleansing
 - B. key component in minimizing the spread of disease
 - C. reduces the number of disease-causing microorganisms
 - D. ALL
 - E.

Note: Satisfactory rating - 16 points Unsatisfactory - below 16 points
 You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

**Hand washing procedures**

Hand hygiene can be accomplished by:

- **Hand washing**
- **Hand antiseptics**
- **Antiseptic hand rub**
- **Surgical hand scrub**

From the various hand hygiene practices available, the use of soap and water remains the most common and the most important when hands are visibly soiled. For hand hygiene in the absence of dirt or debris, however, alternatives such as antiseptic hand rubs, which are rapid acting, inexpensive, and easy to make, are gaining acceptance, especially where access to sinks and clean water is limited

The decision of which type of hand hygiene practice to use depends on:

- *Intensity of contact with patient and/or blood and body fluids*
- *The likelihood of microbial transmission*
- *Patient's susceptibility to infection*
- *Procedure being performed*



Self-Check 2	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. *Hand hygiene can be accomplished by:*

- A. Hand washing**
- B. Hand antiseptics**
- C. Antiseptic hand rub**
- D. Surgical hand scrub** (4 points)
- E. ALL**
- F.**

2. *The decision of which type of hand hygiene practice to use depends on:*

- A. Intensity of contact with patient and/or blood and body fluids*
- B. The likelihood of microbial transmission*
- C. Patient's susceptibility to infection*
- D. Procedure being performed*
- E. ALL**
- F.**

Note: Satisfactory rating - 12 points Unsatisfactory – below 12 points
 You can ask your trainer for the copy of the correct answers.

1. _____

2. _____



Information Sheet 3	<i>Implementing Hand care procedures</i>
----------------------------	---

Implementing Hand care procedures

Hand Washing

Healthcare-associated infections (HCAI) are infections that are acquired in healthcare facilities or as a result of healthcare interventions and are a major problem for patient safety. Hand hygiene is one of the most effective means of preventing HCAs.

The purpose of hand washing with plain soap and water is to mechanically remove soil and debris from skin and reduce the number of transient microorganisms. Hand washing with plain soap and clean water is as effective as washing with antimicrobial soaps (Pereira, Lee, and Wade 1997). But if the tap water is contaminated, hand washing with plain soap is only effective in removing dirt and debris. If tap water is contaminated, use water that has been boiled for 10 minutes and filtered to remove particulate matter (if necessary), or use chlorinated water.

When Do We Wash Our Hands?

- Immediately after arriving and leaving work (the health facility)
- Before and after examining (coming in direct contact with) a client/patient
- After touching contaminated instruments or items
- After exposure to mucous membranes, blood, body fluids, secretions, or excretions
- Before putting on gloves and after removing them
- Whenever our hands become visibly soiled
- After blowing nose or covering a sneeze
- Before eating or serving food
- After visiting the toilet

The 'WHO five moments for hand hygiene

- 1. before touching a patient;**
- 2. before clean/aseptic procedures;**
- 3. after bodily fluid exposure/risk**
- 4. after touching a patient; and**
- 5. after touching patient surroundings.**

Note: Hands should be washed with soap and clean water (or an antiseptic hand rub) after removing gloves because the gloves may have tiny holes or tears, and bacteria can rapidly multiply on gloved hands due to the moist and warm environment within the glove (CDC 1989; Korniewicz et al. 1990).

Steps for Routine Hand Washing

- Thoroughly wet hands with water.
- Apply plain soap (antiseptic agent is not necessary).
- Vigorously rub all areas of hands and fingers for 10 to 15 seconds, paying close attention to areas under fingernails and between fingers.
- Rinse hands thoroughly with clean water.
- Dry hands with personal dry clean towel or paper towel.
- Use a paper towel or a single-use towel after drying hands to turn off the water (faucet handles are contaminated).

When drying hands, using common towels should be avoided. Shared towels may harbor microorganisms and contaminate hands even after proper hand washing.

Note:

- If bar soap is used, provide small bars and soap racks that drain.
- Use running water and avoid dipping hands into a basin containing standing water.
- If liquid soap is being used, do not add soap to a partially empty liquid soap dispenser. This practice of “topping off” dispensers may lead to bacterial contamination of the soap. Liquid soap dispensers should be thoroughly washed and dried before refilling.
- A bucket with a tap or a bucket with a pitcher or jug can be used if running water is not available.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 55 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



- Used water should be collected in a basin and discarded in a latrine if a drain is not available.

Hand Antisepsis and Antiseptic Hand Rub

Hand Antisepsis

The purpose of hand antisepsis is to remove soil and debris and reduce both transient and resident flora on the hands. The technique for hand antisepsis is similar to hand washing except that it involves use of soap containing an antimicrobial agent (often chlorhexidine, iodophors, or triclosan) instead of plain soap or detergent. Medicum, Life Boy, and Dettol are some of the commonly found soaps with antimicrobial agents.

Hand antisepsis should be done before:

- Examining or caring for highly susceptible patients (e.g., premature infants, elderly patients, patients with advanced AIDS)
- Performing an invasive procedure such as placement of an intravascular device
- Leaving the room of patients on contact precautions (e.g., with hepatitis A or E) or who have drug-resistant infections

Antiseptic Hand Rub

Hand rub product is more effective in killing transient and resident flora than plain or medicated soap and water. Antiseptic hand rub is quicker and easier to use and gives a greater initial reduction in hand flora (Girou et al. 2002). Hand rubs also contain a small amount of an emollient such as glycerin, propylene glycol, or sorbitol that protects and softens skin. It is also less irritating to skin than medicated soaps. But, if hands are visibly soiled, hand washing with water and a hand-washing agent should be done first.

Alcohol-based hand rubs provide several advantages compared with hand washing with soap and water because they:

- Require less time
- Act faster
- Are more accessible than sinks
- Are more effective for standard hand washing than soap
- Can provide improved skin condition

A nonirritating, antiseptic hand rub can be made by adding glycerin, propylene glycol, or sorbitol to alcohol (2 mL in 100 mL of 60 to 90 percent ethyl or isopropyl alcohol solution) (Larson 1999).

The technique for performing antiseptic hand rub is as follows:

- Apply enough (5 mL) alcohol-based hand rub to cover the entire surface of hands and fingers.
- Rub the solution vigorously into hands, especially between the fingers and under the nails until dry (15 to 30 seconds).
- Do not rinse hands after applying hand rub.

There are 2 situations where alcohol hand rub alone is not sufficient:


- After contact with a patient with known or suspected diarrhea (e.g. Clostridium Difficile or Norovirus.)
- Where hands are visibly soiled.

In these instances hand wash with antiseptic soap or plain soap followed by use of an alcohol rub is recommended


Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 56 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------

Figure 3.1. Hand-washing techniques with soap and water

Hand washing Technique with Soap and Water



Hand Hygiene Technique with Alcohol-Based Formulation





2. Thoroughly wash hands, especially between fingers, and forearms up to the elbows with soap and water. (If a brush is used, it should be cleaned and either sterilized or high-level disinfected before reuse or shared with others. Sponges, if used, should be discarded.)
3. Clean nails with a nail cleaner.
4. Rinse hands and forearms thoroughly with clean, running water.
5. Apply an antiseptic agent (e.g., 2 to 4 percent chlorhexidine gluconate [CHG]) to all surfaces of hands and forearms to the elbows and rub hands and forearms vigorously for at least two minutes.
6. Rinse hands and arms thoroughly, holding hands higher than the elbows (if tap water is contaminated, use boiled and cooled water or chlorinated water and filter if necessary).
7. Keep hands up and away from the body, do not touch any surface or articles, and dry the hands and forearms with a sterile towel.
8. Put sterile surgical gloves on both hands.

Applying an antiseptic minimizes the number of microorganisms on hands under the gloves and minimizes growth of flora during surgery. Skin damage caused by allergic reactions provides an ideal place for microorganisms to multiply and should be avoided. Personnel with allergies to antiseptics may use plain soap followed by applying the waterless, alcohol- based hand rub.

Other Issues and Considerations Related to Hand Hygiene

Gloves:

- Wearing gloves does not replace the need for hand hygiene.

Hand lotions and hand creams:

- To minimize contact dermatitis related to frequent hand washing (more than 30 times per shift) due to the use of harsh detergents and frequent exposures to antiseptic agents, health care workers may use hand lotions, creams, and moisturizing skin care products. Such products should be water based and without fragrance. Oil-based barrier products, such as those containing petroleum jelly (Vaseline or lanolin), should not be used because they damage latex rubber gloves.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 58 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Self-Check 3	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. When Do We Wash Our Hands?

- A. Immediately after arriving and leaving work
- B. Before and after examining a client/patient
- C. After touching contaminated instruments or items
- D. After exposure to mucous membranes, blood, body fluids, secretions
- E. ALL**

2. In which two situations that alcoholhand rub alone is not sufficient

- A. When hands are visibly soiled and after contact with a patient with known cases**
- B. When hands are not visibly soiled and before contact with a patient with known cases
- C. When hands are visibly soiled and before contact with a patient with known cases
- D. None of the above

3. Alcohol-based hand rubs provide several advantages compared with hand washing with soap and water because they:

- A. Require less time
- B. Act faster
- C. Are more accessible than sinks
- D. Are more effective for standard hand washing than soap
- E. Can provide improved skin condition
- F. ALL**

Note: Satisfactory rating - 09 points Unsatisfactory - below 09 points

You can ask you teacher for the copy of the correct answers.

Score = _____
Rating: _____

Name: _____

Date: _____



Cover Cuts and abrasions

Lesions and skin breaks:

- Cuticles, hands, and forearms should be free from lesions (dermatitis or eczema) and skin breaks. Cuts and abrasions should be covered with waterproof dressings.

Fingernails and artificial nails:

- Long nails may serve as a reservoir for microorganisms, and long nails, either natural or artificial, tend to puncture gloves more easily. As a result, it is recommended that nails be kept moderately short and be less than 0.5 cm long beyond the fingertip. The use of artificial nails by health workers should be restricted (WHO 2009a).

Nail polish:

- Dark-colored nail polish may prevent dirt and debris under fingernails from being seen and removed. Although there is no restriction to wearing nail polish, it is suggested that surgical team members and staff working in specialty areas wear freshly applied, clear nail polish. Chipped nail polish supports the growth of larger numbers of organisms on fingernails compared to freshly polished or natural nails.

Jewelry:

- Although several studies have shown that skin under rings is more heavily colonized than comparable areas of skin on fingers without rings, at the present time, it is not known whether wearing rings results in greater transmission of pathogens. It is suggested that surgical team members not wear rings because it may be more difficult for them to put on surgical gloves without tearing them

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 60 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Self-Check 4	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. Cuticles, hands, and forearms should be free from lesions (dermatitis or eczema) and skin breaks (True False). (2 points).
2. Long nails may serve as a reservoir for microorganisms (True False).
3. Dark-colored nail polish may prevent dirt and debris under fingernails (True False).
4. Skin under rings is more heavily colonized than comparable areas of skin on fingers without rings (True False).

Note: Satisfactory rating - 09 points Unsatisfactory - below 09 points

You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Short Answer Questions

2. _____



Steps in correct hand washing technique:

1. Wet hands with warm water.
2. Apply antimicrobial soap.
3. Rub to form a lather, create friction, and loosen debris.
4. Thoroughly clean between fingers, including thumbs, under fingernails and rings, and up to the wrist, for at least 15 seconds.
5. Rinse hands in a downward position.
6. Dry with a paper towel.
7. Turn off faucets with a clean paper towel to prevent recontamination



LAP Test	Follow hand washing procedures	Practical Demonstration
-----------------	---------------------------------------	--------------------------------

Name: _____ Date: _____

Time started: _____ Time finished: _____

Instructions: Given necessary templates, workshop, tools and materials you are required to perform the following tasks.

Task 1: Perform hand washing procedures



The steps in surgical hand scrub include the following:

1. Remove all rings, watches, and bracelets.
2. Thoroughly wash hands, especially between fingers, and forearms up to the elbows with soap and water. (If a brush is used, it should be cleaned and either sterilized or high-level disinfected before reuse or shared with others. Sponges, if used, should be discarded.)
3. Clean nails with a nail cleaner.
4. Rinse hands and forearms thoroughly with clean, running water.
5. Apply an antiseptic agent (e.g., 2 to 4 percent chlorhexidine gluconate [CHG]) to all surfaces of hands and forearms to the elbows and rub hands and forearms vigorously for at least two minutes.
6. Rinse hands and arms thoroughly, holding hands higher than the elbows (if tap water is contaminated, use boiled and cooled water or chlorinated water and filter if necessary).
7. Keep hands up and away from the body, do not touch any surface or articles, and dry the hands and forearms with a sterile towel.
8. Put sterile surgical gloves on both hands.



LAP Test	Practical Demonstration
-----------------	--------------------------------

Instructions: Given necessary templates, tools and materials you are required to perform the following tasks within 5min.

Task 1: perform surgical hand rub



Instruction Sheet

LG09: Use PPE

This learning guide is developed to provide you the necessary information regarding the following **content coverage and topics** –

Identify and respond to infection risks

- Using of **PPE**
- Changing of PPE daily

This guide will also assist you to attain the learning outcome stated in the cover page. Specifically, upon completion of this Learning Guide, you **will be able to** –

- Personal **PPE** that complies with standards, and is appropriate for the intended use are worn
- Protective clothing and gowns/aprons are changed daily, more frequently if soiled and where appropriate, after each client contact

Learning Instructions:

1. Read the specific objectives of this Learning Guide.
2. Follow the instructions described in number **3 to 16**.
3. Read the information written in the “Information Sheets 1”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
4. Accomplish the “Self-check 1” **in page 8**.
5. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 1).
6. If you earned a satisfactory evaluation proceed to “Information Sheet 2”. However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 1.
7. Submit your accomplished Self-check. This will form part of your training portfolio.
8. Read the information written in the “Information Sheet 2”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
9. Accomplish the “Self-check 2” **in page 16**.
10. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 2).
11. Read the information written in the “Information Sheets **2**”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
12. Accomplish the “Self-check 3” in page 19.
13. Ask your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 3).

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 66 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



14. If you earned a satisfactory evaluation proceed to “Operation Sheet 1” in **page 21**; However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 3.
15. Read the “Operation Sheet 1 and try to understand the procedures discussed.
16. Do the “LAP test” in page 22 (if you are ready). Request your trainer to evaluate your performance and outputs. Your trainer will give you feedback and the evaluation will be either satisfactory or unsatisfactory. If unsatisfactory, your trainer shall advice you on additional work. But if satisfactory you can proceed to Learning Guide #2.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 67 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Information Sheet-1	Personal Protective Equipment
----------------------------	--------------------------------------

Personal Protective Equipment

Personal protective equipment (PPE) refers to a range of barriers and respirators used alone or in combination to protect mucous membranes, airways, skin, and clothing from contact with infectious agents

Protective barriers and clothing are now commonly referred to as PPE. PPE includes gloves, masks/respirators, eyewear (face shields, goggles, or glasses), caps, gowns, aprons, and other items. The basic principle behind wearing PPE is to provide a physical barrier/protection for health care providers and patients/clients from microorganisms.

The most effective barriers are made of treated fabrics or synthetic materials that do not allow water or other liquids (blood or body fluids) to penetrate them. However, these fluid-resistant materials are not widely available because they are expensive. In many countries, caps, masks, gowns, and drapes are made of cloth or paper. Lightweight cotton cloth (with a thread count of 140/inch²) is the material most commonly used for surgical clothing (masks, caps, and gowns) and drapes in many countries. Unfortunately, lightweight cotton does not provide an effective barrier because moisture can pass through it easily, allowing contamination.

Table 4.1. Types of personal protective equipment

Type of Personal Protective Equipment	Must Be Used For:	Primarily Protects:
Caps, gowns/scrub suits, masks, aprons, drapes	Invasive procedures where tissue beneath the skin is exposed	Service provider and client
Closed boots or shoes (open sandals are not acceptable)	Situations involving sharp instruments or when contact with blood and/or body fluids is likely	Service provider
Goggles or glasses, masks, apron or mackintosh	Situations where splashing of blood, body fluids, secretions, or excretions is likely	Service provider
Apron or Mackintosh	Situations where splashing or spillage of blood, body fluids, secretions, or excretions is likely	Service provider
Masks	Situation that call for airborne or droplet transmission precautions	Service providers
Sterile drapes	Major or minor surgical procedures	Client

and hair cover
 all staff body assistant cover use the

Respirators

Particulate respirators are specialized types of masks that are worn by health care personnel to protect them from inhalation exposure to airborne infectious agents that are less than 5 µm in size. These include infectious droplet nuclei from patients with *Mycobacterium tuberculosis* and dust particles that contain infectious particles, such as spores of environmental fungi (e.g., *Aspergillus* spp.). The N95 disposable, particulate, air-purifying respirator is the type used most commonly by health care personnel. (For safe donning and removing of respirator, refer to page 60 of *Infection Prevention and Patient Safety Reference Manual for Service Providers and Managers in Health Care Facilities of Ethiopia*).

Eyewear



- Eyewear protects staff in the event of an accidental splash of blood or other body fluid by covering the eyes. Eyewear includes clear plastic goggles, safety goggles, and faces shields.

Scrub suits or cover gowns

- Scrub suits are worn over, or instead of, street clothes. The main use of cover gowns is to protect the health care workers' clothing.

Surgical gowns

- Surgical gowns are intended to protect patients from microorganisms present on the abdomen and arms of the health care staff during surgery. Lightweight cloth gowns, generally available in Ethiopia, however, offer little protection. Under the circumstances, either wear a plastic apron before putting on the surgical gowns or, if large spills occur, take a shower or bathe as soon as possible after completing the surgery or the procedure. When surgical gowns are worn, sleeves should either taper gently toward the wrists or end with elastic or ties around the wrists. (Large, droopy sleeves invite accidental contamination.) In addition, the cuffs of the surgical gloves should completely cover the end of the sleeves.

Mackintosh or plastic apron

- Plastic aprons are used to protect clothing or surfaces from contamination. Aprons made of rubber or plastic provide a waterproof barrier along the front of the health care worker's body and should also be worn during cleaning and procedures where there is a likelihood of splashes or spillage of blood, body fluids, secretions, or excretions (e.g., when conducting deliveries).

Footwear

- Footwear is worn to protect feet from injury by sharp or heavy items or fluids that may accidentally fall or drip on them. For this reason, sandals, "thongs," or shoes made of soft materials are not acceptable.

Drapes

- Drapes are used to create an operative field around an incision, wrap instruments and other items for sterilization, cover tables in the operating room (OR), and keep clients warm during surgical procedures.
- There are four types of drapes:
 - **Towel drapes** (used for drying hands, squaring off the operative site, and wrapping small items)
 - **Drapes or lap sheets** (used for covering the patient)
 - **Site drapes** (used for minor surgical procedures and have a circular opening)
 - **Pack wrapper drapes** (large drapes that become a table cover when the sterile instrument pack is opened)

Using drapes for a surgical procedure:

All drapes should be applied around a completely dry, wide area of the skin around the site of incision to reduce risk of contamination.

If sterile drapes are used, sterile surgical gloves should be worn when placing the drapes (when putting drapes in place, care must be taken not to touch the patient's body with gloved hands).

- Drapes should be handled as little as possible and should never be shaken or flapped. Always hold drapes above the area to be draped, and discard the drape if it falls below this area.

Remember:

- Once a sterile drape touches the patient's skin, it is no longer sterile.

Sterile cloth drapes do not replace good aseptic technique

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 69 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Self-Check 1	Personal Protective Equipment
---------------------	--------------------------------------

Directions: Answer all the questions listed below. Use the Answer sheet provided in the next page:

12. _____ refers to a range of barriers and respirators used alone or in combination to protect mucous membranes, airways, skin, and clothing from contact with infectious agents
- A. Hand hygiene
 - B. Wearing gloves
 - C. Personal protective equipment
 - D. Massive treatment
13. To contain moisture droplets expelled as health workers speak, cough, or sneeze, which of the following PPE is recommended
- A. Apron
 - B. Scissors
 - C. Pickup forceps
 - D. Masks
14. When performing invasive medical or surgical procedures, which type of glove is used
- A. Surgical Glove
 - B. Clean glove
 - C. Examination glove
 - D. Utility glove

Note: Satisfactory rating - 16 points Unsatisfactory - below 16 points
 You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____



Changing of PPE daily

4.2 Protective clothing and gowns/aprons are changed**Gloves**

Hand hygiene, coupled with the use of protective gloves, is a key component in minimizing the spread of disease and maintaining an infection-free environment.

Health care workers wear gloves for the following three reasons:

1. To reduce the risk of staff acquiring bacterial infections from patients
2. To prevent staff from transmitting their skin flora to patients
3. To reduce contamination of the hands of staff by microorganisms that can be transmitted from one patient to another (cross-contamination)

Types of Gloves

Three types of gloves are used in health care facilities:

1. **Surgical gloves** should be used when performing invasive medical or surgical procedures. The best surgical gloves are made of latex rubber, because of rubber's natural elasticity, sensitivity, and durability. In addition, it provides a comfortable fit. Current standards in Ethiopia recommend that high-level disinfected surgical gloves are the only acceptable alternative if sterile surgical gloves are not available, when performing surgical or invasive procedures (FMOH, *Infection Prevention and Patient Safety Reference Manual for Service Providers and Managers in Health Care Facilities of Ethiopia*, February 2011).
2. **Clean examination gloves** provide protection to health care workers when performing many of their routine duties. These can be used when there is contact with mucous membranes and nonintact skin (e.g., performing medical examinations and procedures such as pelvic examinations).
3. **Utility or heavy-duty household gloves** should be worn for processing instruments, equipment, and other items; for handling and disposing of contaminated waste; and when cleaning contaminated surfaces. Double gloving using either new examination gloves or reprocessed surgical gloves provides some protection in case utility gloves are not available.

When to Wear Gloves

Depending on the situation, surgical gloves, clean examination gloves, or utility gloves should be worn by all staff when:

- There is reasonable chance of hands coming in contact with blood or other body fluids, mucous membranes, or nonintact skin.
- They perform invasive medical procedures (e.g., inserting vascular devices such as peripheral venous lines).
- They handle contaminated waste items or touch contaminated surfaces.



Note:

- When using latex rubber gloves, do not use hand creams or lotions that contain mineral oil, petroleum jelly (Vaseline), or lanolin to protect your hands because they may cause the gloves to break down within minutes.
- A separate pair of gloves must be used for each client to avoid cross-contamination or when moving from one site to another site on the same patient (i.e., from respiratory care to a dressing change).
- It is preferable to use new and single-use (disposable) gloves.

Removing and discarding or reprocessing gloves:


- If gloves are to be discarded, briefly immerse them in 0.5 percent chlorine solution, remove, and dispose in a container for contaminated waste.
- If gloves are to be reprocessed and reused, immerse them in a 0.5 percent chlorine solution briefly, remove gloves by inverting them, and then soak the gloves in the 0.5 percent chlorine solution for 10 minutes before cleaning and processing them

Figure 4.3. How to remove examination gloves



1. Perform hand hygiene before an "aseptic procedure" by handrubbing or hand washing.
2. Check the package for integrity. Open the first non-sterile packaging by peeling it completely off the heat seal to expose the second sterile wrapper, but without touching it.
3. Place the second sterile package on a clean, dry surface without touching the surface. Open the package and fold it towards the bottom so as to unfold the paper and keep it open.
4. Using the thumb and index finger of one hand, carefully grasp the folded cuff edge of the glove.
5. Slip the other hand into the glove in a single movement, keeping the folded cuff at the wrist level.
- 6-7. Pick up the second glove by sliding the fingers of the gloved hand underneath the cuff of the glove.
- 8-10. In a single movement, slip the second glove on to the ungloved hand while avoiding any contact/resting of the gloved hand on surfaces other than the glove to be donned (contact/resting constitutes a lack of asepsis and requires a change of glove).
11. If necessary, after donning both gloves, adjust the fingers and interdigital spaces until the gloves fit comfortably.
- 12-13. Unfold the cuff of the first gloved hand by gently slipping the fingers of the other hand inside the fold, making sure to avoid any contact with a surface other than the outer surface of the glove (lack of asepsis requiring a change of gloves).
14. The hands are gloved and must touch exclusively sterile devices or the previously-disinfected patient's body area.

Figure 4.5. How to remove sterile gloves



- 15-17. Remove the first glove by peeling it back with the fingers of the opposite hand. Remove the glove by rolling it inside out to the second finger joints (do not remove completely).
18. Remove the other glove by turning its outer edge on the fingers of the partially ungloved hand.
19. Remove the glove by turning it inside out entirely to ensure that the skin of the health-care worker is always and exclusively in contact with the inner surface of the glove.
20. Discard gloves.
21. Perform hand hygiene after glove removal according to the recommended indication.

NB: Donning surgical sterile gloves at the time of a surgical intervention follows the same sequences except that:

- it is preceded by a surgical hand preparation;
- donning gloves is performed after putting on the sterile surgical gown;
- the opening of the first packaging (non-sterile) is done by an assistant;
- the second packaging (sterile) is placed on a sterile surface other than that used for the intervention;
- gloves should cover the wrists of the sterile gown.

sticks or other injuries occur, it may decrease the risk of blood to hand contact.

Double gloving can be used during the following:

- Procedures that involve coming in contact with large amounts of blood or other body fluids (e.g., vaginal deliveries and cesarean sections)
- Orthopedic procedures in which sharp bone fragments, wire sutures, and other sharps are likely to be encountered
- Surgical procedures lasting more than 30 minutes

Some Dos and Don'ts Regarding Gloves

Do

- Wear the correct size gloves, particularly for surgical gloves. A poorly fitting glove can limit your ability to perform the task and may get damaged easily.
- Change surgical gloves periodically (every 45 minutes) during long cases because the protective effect of latex gloves decreases with time and unapparent tears may occur.
- Keep fingernails trimmed moderately short (less than 3 mm beyond the fingertip) to reduce the risk of tears.
- Pull gloves up over cuffs of gown (if worn) to protect the wrists.
- Use water-soluble hand lotions and moisturizers often to prevent hands from drying and cracking due to frequent hand washing and gloving.

Don't

- Use oil-based hand lotions or creams because they will damage latex surgical and examination gloves.
- Use latex gloves if you or the patient has an allergy to latex.



- Store gloves in areas where there are extremes of temperature (e.g., direct sunlight; near the heater, air conditioner, ultraviolet light, or x-ray machine). These conditions may damage the gloves (cause breakdown of the material they are made of), thus reducing their effectiveness as a barrier.
- Reprocess gloves that are cracked or have detectable holes/tears (FMOH, *Infection Prevention and Patient Safety Reference Manual for Service Providers and Managers in Health Care Facilities of Ethiopia*, February 2011).
- Reprocess examination gloves for reuse (FMOH, *Infection Prevention and Patient Safety Reference Manual for Service Providers and Managers in Health Care Facilities of Ethiopia*, February 2011).

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 74 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Self-Check 2	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. Health care workers wear gloves for the following three reasons: (4 points)
 - A. To reduce the risk of staff acquiring bacterial infections from patients
 - B. To prevent staff from transmitting their skin flora to patients
 - C. To reduce contamination of the hands of staff
 - D. ALL

2.provide protection to health care workers when performing many of their routine duties
 - A. Surgical gloves
 - B. Clean examination gloves
 - C. heavy-duty household gloves
 - D. ALL

3. When to Wear Gloves
 - A. hands coming in contact with blood or other body fluids
 - B. perform invasive medical procedures
 - C. handle contaminated waste items or touch contaminated surfaces
 - D. ALL

4. Double gloving can be used during the following: (4 points)
 - A. Orthopedic procedures
 - B. Surgical procedures lasting more than 30 minutes
 - C. contact with large amounts of blood or other body fluids
 - D. ALL

Note: Satisfactory rating - 12 points

Unsatisfactory – below 12 points

You can ask your trainer for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

**Steps for donning examination glove**

1. Take out glove from its original box
2. Touch only restricted surface of the glove corresponding to the wrist (at the top edge of the cuff)
3. Don the final glove
4. Take the second glove with the bare hand and touch only restricted surface
5. To avoid touching the skin of the forearm with the gloved hand turn.
6. Once gloved hands should not touch anything else that is not designed by indications and conditions for glove use.

1.2. Steps for donning surgical glove

1. Perform hand hygiene before an “aseptic procedures “by hand rubbing or hand is washing.
2. Check the package for integrity open the first non sterile package by peeling it completely off the heat seal to expose the second sterile wrapper but without touching it.
3. Place the second sterile package on a clean dry surface without touching the surface open the package and fold it toward the bottom so as to unfold the paper and keep it open.
4. Using the thumb and index finger of the hand carefully grasp the folded cuff edge of the gloves.
5. Slip the other hand in to the glove in to a single movement keeping the folded cuff at the wrist level.
6. Pick up the second glove by sliding the fingers of the gloved hand under neath the cuff of the glove.
7. In a single movement slip the second glove on to the un gloved hand while avoiding any contact/resting On the gloved hand o surface other than the glove to be donned.
8. If necessary after donning both gloves adjust the fingers and inter digital spaces until the glove fit comfortable.
9. Unfold the cuff of the first gloved hand by gently slipping the fingers of the other hand inside the fold making sure to avoid any contact with a surface other than the outer surface of the gloves (lack of asepsis requiring a change of glove.)
10. The hand is gloved and must touch exclusively sterile devices or the previously disinfected patient body area.

1.3. Steps for removing glove

1. Pinch one glove at the wrist level to remove it without touching the skin of the forearm and peel away from the hand thus allowing the glove to turn inside out.
2. Hold the removed glove in the gloved hand and slide the fingers of the un gloved hand inside b/n the gloved and the wrist remove the second glove by rolling it down the hand and fold into the first glove.
3. Discard the removed gloves



LAP Test	Practical Demonstration
-----------------	--------------------------------

Name: _____ Date: _____

Time started: _____ Time finished: _____

Instructions: Given necessary templates, workshop, tools and materials you are required to perform the following tasks.

Task 1: perform donning examination glove.

Task 2: perform donning surgical glove.

Task 3: Perform removing glove.



Instruction Sheet **LG10: Limit contamination**

This learning guide is developed to provide you the necessary information regarding the following **content coverage and topics** –

Limit contamination

- Demarcating of Clean and contaminated zones
- Storage of records, materials and medicaments to a clean zone
- well-designating Contaminated instruments and equipment to contaminated zone

This guide will also assist you to attain the learning outcome stated in the cover page. Specifically, upon completion of this Learning Guide, you **will be able to** –

- demarcate and maintain clean and contaminate zones in all aspects of health care work
- confine records, materials and medicaments are to a well-designated clean zone
- Confined contaminate instruments and equipment to a well-designated contaminated zone

Learning Instructions:

1. Read the specific objectives of this Learning Guide.
2. Follow the instructions described in number **3 to 16**.
3. Read the information written in the “Information Sheets 1”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
4. Accomplish the “Self-check 1” **in page 8**.
5. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 1).
6. If you earned a satisfactory evaluation proceed to “Information Sheet 2”. However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 1.
7. Submit your accomplished Self-check. This will form part of your training portfolio.
8. Read the information written in the “Information Sheet 2”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
9. Accomplish the “Self-check 2” **in page 16**.
10. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 2).
11. Read the information written in the “Information Sheets **3**”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
12. Accomplish the “Self-check 3” in page 19.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 78 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



13. Ask your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 3).
14. If you earned a satisfactory evaluation proceed to “Operation Sheet 1” in page 21; However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 3.
15. Read the “Operation Sheet 1 and try to understand the procedures discussed.
16. Do the “LAP test” in page 22 (if you are ready). Request your trainer to evaluate your performance and outputs. Your trainer will give you feedback and the evaluation will be either satisfactory or unsatisfactory. If unsatisfactory, your trainer shall advice you on additional work. But if satisfactory you can proceed to Learning Guide #2.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 79 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Demarcating Clean and contaminated zones

When something has been exposed to infectious agents it is considered to have been contaminated. Ways to limit contamination in the healthcare setting may include:

- cleaning surfaces
- protecting all materials, equipment and instruments from contaminants
- maintaining sterile objects
- being aware of the guidelines for single-use objects.

An important strategy to limit contamination in healthcare settings is to set and maintain:

- clean zones
- treatment zones
- contaminated zones.

Maintaining these three zones helps reduce the risk of contamination. it is the responsibility of every person working in the healthcare environment to:

- understand where the clean, treatment and contaminated zones are
- do what is required to maintain them.

Zones

Maintaining clean, treatment and contaminated zones helps reduce the risk of contamination and makes it easier to remember that anything entering a clean zone must first be decontaminated.

The clean Zone

Clean areas include those surfaces and drawers where clean, disinfected or sterilized instruments are stored and never come in contact with contaminated instruments or equipment

The clean zones are areas where non-contaminated items are kept. Example of these items and zones include:

- sterile instruments
- clean linen
- medical records
- kitchen preparation areas
- supply stores.

Before entering a clean zone it is important to remove contaminated gloves and other PPE and perform hand hygiene (decontamination).

The treatment Zone

The treatment zone is where items are currently being used by the client or healthcare worker.

For example, the client's bedside is a treatment zone as it has been exposed to microorganisms.

Contaminated zone

Contaminated zones are for objects and waste that is waiting for decontamination sterilization or disposal

The contaminated zone boundaries should be clearly defined, because this has implications for surface management and for the placement of equipment.

Instruments placed into the contaminated zone for a treatment session but not used during the session must be regarded as contaminated. For this reason, all bulk supplies such as opened boxes of gloves, cotton rolls or gauze must be stored outside the contaminated zone and protected from contamination from splashes and aerosols.

An example of this zone is a linen skip, in which used linen is stored while awaiting decontamination in the laundry.

If there is any possibility that any item **may** have been contaminated, it should be treated as if it **has** been contaminated.



Self-Check 1	Written Test
---------------------	---------------------

Directions: Answer all the questions listed below. Use the Answer sheet provided in the next page:

1. Ways to limit contamination in the healthcare setting may include:
 - A. cleaning surfaces
 - B. protecting all materials, equipment and instruments from contaminants
 - C. maintaining sterile objects
 - D. being aware of the guidelines for single-use objects.
 - E. ALL**
2. An important strategy to limit contamination in healthcare settings is to set and maintain:
 - A. clean zones
 - B. treatment zones
 - C. contaminated zones
 - D. ALL**
3. Ways to limit contamination in the healthcare include all of the following, except
 - a. Cleaning surfaces
 - b. Protecting all materials from contaminants
 - c. Maintaining sterile objects
 - d. None of the above**
4. An important strategy to limit contamination in healthcare settings is to set and maintain
 - a. Demarcation among different zones**
 - b. Effective treatment
 - c. Taking prophylaxis
 - d. Isolation of patients
5. Areas where non-contaminated items are kept is
 - a. Treatment Zone
 - b. Contaminated zone
 - c. Clean zone**
 - d. Patient zone
- 6.

Note: Satisfactory rating - 10 points Unsatisfactory - below 5 points
 You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____



Information Sheet 2

Storage of records, materials and medicaments to a clean zone

Storage of records, materials and medicaments to a clean zone

Records, materials and medicaments

All records that are kept and archived should be kept in a clean zone. All packaging materials and medicaments should be kept in a clean zone. These items should not be stored in a contaminated zone or a risk of cross contamination will occur.

Contaminated instruments and equipment

When items are received into the sterilizing facility, all reusable items that have been used or unused during patient treatment need to be cleaned in a physically separate area to prevent possible contamination of processed items



Self-Check 2	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. All records that are kept and archived should be kept in
 - A. a clean zone
 - B. Treatment Zone
 - C. Patient zone
 - D. Contaminated zone
 - E.

2. All packaging materials and medicaments should be kept in (4 points)
 - A. a clean zone
 - B. Treatment Zone
 - C. Patient zone
 - D. Contaminated zone

3. Which zone is used for objects and waste that is waiting for decontamination sterilization or disposal
 - A. Treatment Zone
 - B. Contaminated zone
 - C. Clean zone
 - D. Patient zone

Note: Satisfactory rating - 12 points Unsatisfactory – below 12 points
 You can ask your trainer for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Information Sheet 3	well-designating Contaminated instruments and equipment to contaminated zone
----------------------------	--

well-designating Contaminated instruments and equipment to contaminated zone

Workflow

To maintain the separation of clean and contaminated zones, workflow should be from the cleanest to the most contaminated areas – this ensures that there is never any movement of contaminated items into clean zones.

The cleanest to most contaminated approach also applies to cleaning surfaces, when you should start at the cleanest area and work out toward the most contaminated area. Make sure that cleaning equipment is correctly decontaminated and stored after use, to prevent it becoming another source of contamination.

Objects that have moved from the clean to the contaminated zone may only return when they have been cleaned, decontaminated, or sterilized as required. Take them from the contaminated zone, decontaminate them, and then place them in the clean zone. Be careful not to place a newly decontaminated item back in a contaminated zone, such as on a contaminated bench.

If there is a possibility that something may possibly have been contaminated, it should be treated as if it has definitely been contaminated.


Many healthcare facilities use dedicated trolleys and containers for holding and moving contaminated items and clean or sterile items, and you must ensure that the correct trolleys and containers are used. If a single trolley is used for both, it must be thoroughly cleaned before use for clean items. Clean and contaminated items can never be placed on the trolley together.

There are two basic rules to follow to limit contamination.

- Maintain clean zones and contaminated zones within the workplace.
- All movement of instruments and equipment must be from clean to contaminated.

The aim of barrier nursing is to protect the HCW but also the community from transmission of Infectious disease. Proper barrier management is the cornerstone in containing the spread of Infectious disease. in healthcare settings.

Figure . Zones and colour code



Different zones	Activities	Specification
Red zone • Patient treatment area	<ul style="list-style-type: none"> • Patient treatment area • Point of care; diagnostics • After visible contamination, cleaning and disinfection of HCW 	<ul style="list-style-type: none"> • Monitored by direct or video assisted observation
Dark yellow zone	<ul style="list-style-type: none"> • First re-entry step for staff exiting the red zone • Assisted disinfection and doffing for exit HCW • Potentially contaminating processes, such as cleaning and disinfection of boots and waste bags • Preparing waste for further processing, such as packaging waste bags in containers with non-removable clip-on lids 	<ul style="list-style-type: none"> • Critical zone for prevention and control of secondary contamination. • Abundant space required for unrestricted assisted doffing process of two HCW in PPE • Additional dedicated cleaning and disinfection areas
Light yellow zone	<ul style="list-style-type: none"> • Storage of waste • Hand disinfection for HCW before stepping into the green zone 	<ul style="list-style-type: none"> • Additional generously dimensioned waste storage areas. • Supervisor (barrier nursing guardian)
Green Zone	<ul style="list-style-type: none"> • Second step re-entry of staff from light yellow zone • Complete assisted donning for entry HCW • Briefing and de-briefing of staff • Staff coordination and supervision of activities • Inbound and outbound communications 	<ul style="list-style-type: none"> • Strict access control • Space for clean supplies • Zones for staff recreation

Functionally, the yellow zone needs to be understood as the decisive area, in which secondary contamination is prevented and controlled: Here a contaminated HCW exiting from the red zone is brought in to clean conditions, which enable him or her to safely re-enter the green zone. In addition, any material coming from the red zone, such as waste bags, re-usable PPE items, patient samples needed to be processed outside of the isolation unit are first cleaned and disinfected in the yellow zone. A particular function lies in the temporary storage of considerable amounts of waste produced every day in the nursing of a patient with an Infectious disease.

Practical hints

- Different zones need to be clearly marked.
- Prevention and control of secondary contamination happens in the yellow zone.
- The yellow zone has a virtual gradient from 'high potential for contamination' (dark yellow) to 'low potential of contamination' (light yellow).
- Instructions for staff should be displayed at the entry of the isolation area.
- If there is a cross-contamination incident outside the red zone (e.g. patient leaving the red zone), the contaminated area has also to be considered as a red zone. New yellow and green zones need to be established around the new red zone. The zones can be put back into normal function by room disinfection, once the patient has been dismissed.
- Donning and doffing areas must be separated and visually marked. The donning area is in the green zone.
- The doffing area must be in the dark yellow zone, but has to be clearly separated from the light yellow zone.

**Self-Check 3****Written Test**

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page:

1. Patient treatment zone
 - A. Red zone
 - B. Dark yellow zone
 - C. Green Zone
 - D. NONE
 2. Critical zone for prevention and control of secondary contamination.
 - A. Red zone
 - B. Dark yellow zone
 - C. Green Zone
 - D. NONE
 3. Zones for staff recreation
 - A. Red zone
 - B. Dark yellow zone
 - C. Green Zone
 - D. NONE
- A.

Note: Satisfactory rating - 09 points

Unsatisfactory - below 09 points

You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____

Rating: _____

Name: _____

Date: _____

**Operation Sheet 3****Prepare tools and equipment for identification and measuring Muda.**

3. Discuss and plan to prepare tools and equipment for Muda identification.
4. Prepare tools and equipment for Muda identification.



LAP Test	Practical Demonstration
-----------------	--------------------------------

Name: _____ Date: _____

Time started: _____ Time finished: _____

Instructions: Given necessary templates, workshop, tools and materials you are required to perform the following tasks.

Task 1: Identify and prepare tools and equipment for measuring and identification of Muda.



Instruction Sheet	LG11: Handle, package, label, store, transport and dispose of clinical and other waste
--------------------------	---

This learning guide is developed to provide you the necessary information regarding the following **content coverage and topics** –

Handle, package, label, store, transport and dispose of clinical and other waste

- wearing appropriate personal protective equipment when handling waste
- separating and disposing waste into color coded waste containers
- storage of Clinical or related waste into accessible area
- waste management to minimize potential risk
- safely waste disposal in accordance with policies

This guide will also assist you to attain the learning outcome stated in the cover page. Specifically, upon completion of this Learning Guide, you **will be able to** –

- worn appropriate personal protective clothing and equipment in accordance with OHS policies and procedures when handling waste
- separate and identify waste at the point where it has been generated and dispose of into waste containers that are color coded
- Clinical or related waste is stored in an area that is accessible only to authorized persons
- handle, package, label, store, transport and disposed of appropriately waste to minimize potential for contact with the waste and to reduce the risk to the environment from accidental release
- dispose of waste safely in accordance with policies and procedures of the organization and legislative requirements

Learning Instructions:

1. Read the specific objectives of this Learning Guide.
2. Follow the instructions described in number **3 to 16**.
3. Read the information written in the “Information Sheets 1”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
4. Accomplish the “Self-check 1” **in page 8**.
5. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 1).
6. If you earned a satisfactory evaluation proceed to “Information Sheet 2”. However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 1.
7. Submit your accomplished Self-check. This will form part of your training portfolio.
8. Read the information written in the “Information Sheet 2”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 89 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



9. Accomplish the “Self-check 2” in page 16.
10. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 2).
11. Read the information written in the “Information Sheets 5”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
12. Accomplish the “Self-check 3” in page 19.
13. Ask your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 3).
14. If you earned a satisfactory evaluation proceed to “Operation Sheet 1” in page 21; However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 3.
15. Read the “Operation Sheet 1 and try to understand the procedures discussed.
16. Do the “LAP test” in page 22 (if you are ready). Request your trainer to evaluate your performance and outputs. Your trainer will give you feedback and the evaluation will be either satisfactory or unsatisfactory. If unsatisfactory, your trainer shall advice you on additional work. But if satisfactory you can proceed to Learning Guide #2.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 90 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Information Sheet-1	wearing appropriate personal protective equipment when handling waste
----------------------------	---

1. Wearing appropriate personal protective equipment

1.1. Health care wastes

Health care waste is a byproduct of health care that includes potential risk and nonrisk wastes. It includes all the waste generated by health care establishments, research facilities, and laboratories. Health care waste can be classified as high-risk and low-risk wastes depending on the level of the risk they pose to the health provider, patient, and community.

Appropriate handling and disposal of potentially infectious waste helps to prevent the spread of infection, illness and disease

High-Risk Waste

High-risk waste includes the following:

- **Infectious waste:**
 - Blood, blood products, and other bodily fluids or items contaminated with similar fluids
 - Cultures and stocks of infectious agents from the laboratory and items contaminated with such agents
 - Isolation waste from highly infectious patients (including food residue)
 - Discarded live and attenuated vaccines
 - Waste, bedding, bandages, surgical dressings, and other contaminated material infected with human pathogens
- **Anatomical waste:**
 - Human tissues, body parts, and fetuses
 - Biopsies, carcasses, organs, and tissues infected with human pathogens
- **Sharps waste (used or unused):**
 - Syringes, needles, scalpel blades, suture needles, razors, and intravenous set needles
- **Chemical waste:**
 - Formaldehyde, photographic chemicals, solvents, organic and inorganic chemicals
- **Pharmaceutical waste:**
 - Outdated medications and residuals of drugs used in chemotherapy
 - Items contaminated by or containing pharmaceutical bottles/boxes
- **Radioactive waste:**
 - Contamination with radioactive isotopes
- **Genotoxic waste:**
 - Cytostatic drugs
 - Vomit, urine, or feces from patients treated with cytotoxic drugs, chemicals, and radioactive material
- **Pressurized containers:**
 - Explosion of cylinders containing gases or aerosols
- **Waste with high content of heavy metals:**
 - Batteries, broken thermometers, blood pressure gauges, etc.

Low-Risk Waste

Noninfectious waste:

Commercial waste is non-contaminated waste and poses no infectious risk to persons who handle it. Examples include paper, trash, boxes, bottles, plastic containers, leftover foods, and food products

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 91 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Not all health care waste is risky; most waste (80 to 85 percent) generated from health care facilities is believed to be noninfectious and non-risky (FMOH, *Infection Prevention and Patient Safety Reference Manual for Service Providers and Managers in Health Care Facilities of Ethiopia*, February 2011).

The proportion of waste generated from health care is as follows:

- Noninfectious waste: 80 percent
- Pathological waste and infectious waste: 15 percent
- Sharps waste: 1 percent
- Chemical or pharmaceutical waste: 3 percent
- Pressurized cylinders, thermometers: less than 1 percent

Risks of Health Care Waste

Inadequate and inappropriate handling of health care waste may have serious public health consequences and a significant impact on the environment. Injuries, transmission of infections, environmental pollution, fire hazards, and public nuisance (offensive smells, unsightly debris, etc.) are the major risks and hazards of poorly managed health care waste.

Improper health care waste management can expose health workers, patients, and the community to the risk of being exposed and potentially infected by blood-borne pathogens. Studies revealed that 33 percent of HBV and 42 percent of HCV infections occur due to direct or indirect exposure to infectious waste (WHO 2005a).

Improper health care waste management can also expose people to gastro enteric and respiratory infections.

In addition to health risks from direct contact, health care waste can impact human health by contaminating bodies of water and polluting the air. Emission of persistent organic pollutants/toxic gases like dioxins, furans, and polychlorinated biphenyls is dangerous to human health.

1.2. personal protective clothing and equipment during waste handling

Personal protective clothing and equipment during waste handling includes

- **Disposable plastic aprons and Gowns**
 - Aprons should be put on at the beginning of the activity
- **Full-body gowns**
 - Use where there is a risk of extensive splashing of blood, bodily fluids, secretions or excretions on to the skin or clothing of a HCW
- **Gloves including rubber gloves**
- **Face masks/respirators**
 - The mucous membrane of the mouth, nose and eyes are portals of entry for infectious agents as other skin surfaces if not intact.
- **Eye protect ion/face visors/protective spectacles**
 - Goggles, visors or protective spectacles must be worn to protect the eye
- **Forearm protection**
 - Forearm protection should be available for use in areas that there is a risk of injury. It should be used in conjunction with a detailed plan of care to minimize injuries from scratches and bites
- **Foot protection**
 - To protect feet and legs from falling objects, moving machinery, sharp objects, hot materials, chemicals, or slippery surfaces, employees should wear closed-toed shoes, boots, foot guards, leggings, or safety shoes as appropria

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 92 of 139
---------------------------	----------------------	---------------------------------------	-----------------------	----------------



Self-Check 1	Written Test
---------------------	---------------------

Directions: Answer all the questions listed below. Use the Answer sheet provided in the next page:

1. High-risk waste includes the following
 - A. Anatomical waste
 - B. Pharmaceutical waste
 - C. Radioactive waste
 - D. ALL**

2. Waste that are arising from Cytostatic drugs are
 - a. Drug waste
 - b. Genotoxic waste**
 - c. Pharmacological waste
 - d. Low risk waste

3. Batteries, broken thermometers, blood pressure gauges are
 - a. Medical equipment waste
 - b. Genotoxic waste
 - c. Pharmacological waste
 - d. Waste with high content of heavy metals**

4. Most waste (80 to 85 percent) generated from health care facilities is believed to be
 - a. Infectious
 - b. Risky
 - c. Noninfectious and non-risky**
 - d. Blood borne

Note: Satisfactory rating - 16 points Unsatisfactory - below 16 points
 You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Answer

No	
1	D
2	B
3	D
4	C



Information Sheet 2	waste separation
----------------------------	-------------------------

2. waste separation

Contaminated and non-contaminated wastes should be separated at origin, to reduce the volume of contaminated waste and minimize the cost to the institution for more expensive procedures required for managing and disposing of contaminated waste properly.

- When possible, use separate containers for combustible and non-combustible waste.
- Never sort through contaminated wastes (e.g., do not try to separate non-contaminated waste from contaminated wastes, or combustible from non-combustible, after they have been combined).

STEPS OF WASTE MANAGEMENT

- Segregation
- Collection
- Transportation
- Disposal

Self-Check 2	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page:

:

1. Contaminated and non-contaminated wastes should be separated at

- A. At Origin
- B. At the final site
- C. At the site of disposal
- D. None of the above

Note: Satisfactory rating - 16 points Unsatisfactory - below 16 points

You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____

Rating: _____

Name: _____

Date: _____



Self-Check 3	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

1. What is OHS represents for (2 point)
2. What is OSH represents for? (1 point)
3. What does WHS represent for? (1 point)
4. What are the goals of OHS? (2 points)
5. List at least two examples of OHS requirements in your work areas. (10 points)
6. List at least four workplace hazards? (4 points)

Note: Satisfactory rating - 12 points Unsatisfactory – below 12 points
 You can ask your trainer for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____



Information Sheet 4

waste management to minimize potential risk

Waste Minimization/Containment

Waste minimization is the first and best way to reduce health care waste quantities and costs, environmental impact, and exposure to health care workers, patients, and communities. Effective waste minimization practices require that the purchases of all materials and supplies be made with waste reduction in mind.

Segregation

Waste segregation is separating waste by type at the place where it is generated. Waste should immediately be separated by the person generating the waste, according to its type, and placed in a bin with an appropriate colored bin liner or into a sharps container. Waste handlers should never sort through waste after it has been placed in the bin. The color-coding system aims at ensuring immediate and nonequivocal identification and segregation of the hazards associated with the type of health care waste that is handled or treated. It is very important that both providers and waste handlers understand the color-coding system and handle waste accordingly. The following table shows the recommended color coding for categories of health care waste.

Category	Example	Color of Bin Liner
Noninfectious	Paper, packing materials, plastic bottles, food, cartons	Black
Infectious	Gloves, dressings, blood, body fluids, used specimen containers	Yellow with biohazard symbol
Highly infectious	Anatomical waste, pathological waste	Red with biohazard symbol
Chemical	Formaldehyde, pathological chemicals, solvents, organic and inorganic chemicals	Brown
Radioactive	Any solid, liquid, or pathological waste contaminated with radioactive isotopes of any kind	Yellow with radioactive label
Sharps	Needles, needles from intravenous sets, scalpels, blades, lancets, broken glass, syringes, and needles	Yellow box marked "SHARPS" with biohazard symbol

Waste handling refers to activities involving the handling of waste at the health care facility as well as its collection and storage.

Waste collection is the process of removing waste bags from the service point and taking to storage or disposal area. It also includes quantifying waste by volume, labeling as to its source, and recording.

Waste storing is the process of placing waste in a secure place until it can be disposed of. The ideal storage area should be designated (for waste only), secure (only authorized persons should have access), and kept clean, dry, and pest free. The designated central storage facility should be located within the premises of the health facility, close to the treatment unit but away from food storage or food preparation areas. Health care waste should be stored for no longer than two to three days, depending on weather conditions. However, in the case of safety boxes, the filled box can be stored in a locked room for up to one week at lower-level health care facilities where there is no incinerator. Organic waste should be disposed of daily. Segregation must be maintained throughout until final disposal.



Waste Transportation

Waste transportation is movement of waste from one place to another. Waste transportation can be either on-site or off-site and should also maintain waste segregation.

- **On-site:** Moving waste from one point to another within the health care facility
- **Off-site:** Transporting waste outside the health facility

Treatment and Disposal

Health care waste is treated to render it nonhazardous. Noninfectious waste does not need to be treated. Disposal is a process of eliminating health care waste without posing any risk to health facility workers or the general public.

The following are health care waste treatment and disposal options at the health care facility level: **Sharps waste:**

- Incineration using properly built and maintained medium- or high-temperature incinerator on-site
- Transport to off-site incinerators, if there is centralized treatment service
- On-site burial in a protected pit

Infectious waste:

- On-site burial in a protected pit
- On-site incineration in a medium- or high-temperature incinerator provided that the incinerator is standard and capable of destroying such wastes
- Transport to off-site treatment/disposal site, if the service is available

Non-risk waste:

- Collection by municipal truck or other private trash collection firm for landfill disposal

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 97 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



Self-Check 4	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page:

2. According to the color coding system, infectious wastes such as gloves, dressing, blood, body fluids and used specimen containers are put in
 - A. Yellow color container
 - B. Red color container
 - C. Blue color container
 - D. Dark color container
3. According to the color coding system, Highly infectious wastes such as anatomical and pathological waste are put in
 - A. Yellow color container
 - B. Red color container
 - C. Blue color container
 - D. Dark color container
4. Process of placing waste in a secure place until it can be disposed of is
 - A. Waste storing
 - B. Waste handling
 - C. Waste collection
 - D. None of the above
5. Moving waste from one point to another within the health care facility is
 - A. On site
 - B. Off site
 - C. Far site
 - D. Near site

Note: Satisfactory rating - 12 points Unsatisfactory – below 12 points
 You can ask your trainer for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____



Health care wastes disposal methods

Recommended Disposal Methods of Health Care Waste

1. Burial in a protected pit:

- Waste is placed into a pit (1 to 2 meters wide, 2 to 5 meters deep, and at least 1.5 meters above the water table) and covered with soil.
- Protected burial pits are an acceptable, and perhaps the most appropriate, disposal option for infectious wastes in rural health care facilities.
- Pits should be at least 50 meters away from any water source, be fenced to restrict unauthorized access, and be located away from public areas.
- Keep waste covered with a 10- to 30-cm layer of soil every time waste is added to the pit.
- Only contaminated and hazardous waste needs to be buried.
- Expired vaccines and drugs should be encapsulated and buried. Place the expired drugs and vaccines in a hard container such as a metal drum. When the container is mostly full (3/4) add a mixture of 1 part cement, 1 part lime, 4 parts sand, and 1/3 to 1/2 part water. Lime works as a disinfectant, and it also helps the cement flow into empty spaces to completely surround the waste. Seal the container and bury it in a pit. Expired vaccines and drugs should not be burned unless there is appropriate incineration technology.
- When the level of waste reaches to within 30 to 50 cm to the surface of the ground, completely fill the pit in with soil and dig another pit.

Advantages: Simple and inexpensive

Disadvantages: Can handle only a small volume of waste in areas where there is shortage of space and presents a danger to the community if not buried or covered properly

2. Burn

- Waste is placed into a pit and burned on a regular basis (at least once a week, according to volume of waste and size of pit). Waste must be burned thoroughly, and ashes must be covered with soil.
- Pits should be dug 1 to 2 meters wide and to a depth of 2 to 5 meters, but also at least 1.5 meters above water table
- The pit should be fenced off to restrict unauthorized access. The burn pit must be located away from public areas, and smoke from burning waste must not affect the surrounding area.
- Open burning (outside of a pit, on the ground) should not be practiced.
- Medical waste may not burn easily, especially if it is wet. Add kerosene to make the fire hot enough to burn all wastes.

3. Medium- or high-temperature incineration:

- Incineration is medium- or high-temperature burning. It reduces the volume of the waste and, if high enough temperatures are reached, eliminates pathogens.
- Proper incineration produces fewer pollutants than open-air burning and is preferred if a good-quality incinerator is available with a well-trained operator. Proper incineration includes:
 - Clear operation procedures
 - ✓ Trained operator
 - ✓ Reliable segregation system

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 99 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	----------------



- ✓ Reliable transport system
- ✓ Ash pit
- ✓ Maintenance performed on schedule
- ✓ Adequate supply of fuel
- ✓ Secured incinerator

When using incinerator:

- Keep incinerator clean. Remove ash from ash chamber and grate and dispose the ash into ash pit. Do not store waste in incinerator.
- Some incinerators need to be preheated by burning general or nonmedical waste (e.g., paper) until the incinerator reaches the recommended temperature for incinerating health care waste (800°C in the burning chamber).
 - Polyvinyl chloride plastics (like blood bags and intravenous lines), large amounts of reactive chemical waste, silver salts and photographic or radiographic waste (x-ray materials), waste with high mercury (such as broken mercury thermometers) or cadmium content, batteries, aerosol cans or pressurized gas containers, and glass vials must never be incinerated. **N.B. Syringes are not made of polyvinyl chloride plastic and are therefore safe to burn.**

Advantages: Treats and greatly reduces waste volume

Disadvantages: Overfilling the combustion chamber and wastes with high moisture content can produce smoke, and also may produce emission and hazardous ash that contain dioxins and metals. It may require pollution control equipment to meet local environmental regulation.

In general, medical wastes require careful disposal and containment before collection and consolidation for treatment. These measures are designed to protect the workers who generate medical wastes and who manage the wastes from point of generation to disposal.

- A single, leak-resistant biohazard bag is usually adequate for containment of regulated medical wastes, provided the bag is sturdy and the waste can be discarded without contaminating the bag's exterior.
- All bags should be securely closed for disposal.
- Puncture-resistant containers located at the point of use (e.g., sharps containers) are used as containment for discarded slides or tubes with small amounts of blood, scalpel blades, needles and syringes, and unused sterile sharps.
- To prevent needle stick injuries, needles and other contaminated sharps should not be recapped, purposefully bent, or broken by hand.
- Infected materials should be put in impermeable bags or hard plastic containers. These bags and containers need to be clearly labelled as highly infectious waste. Considerations on waste management:
 - ✓ Solid non-sharp waste should be placed in impermeable, clearly labelled bags to be discarded following applicable environmental regulations for the disposal and inactivation of infectious medical waste. Temporary disposal sites should be located as close as possible to the patient care area.
 - ✓ Sharp, pointed objects (e.g. open vials, needles) should be placed in hard plastic containers and labelled clearly.
 - ✓ Liquid waste (e.g. vomit, urine and diarrheal fluids) may only be disposed in the sanitary sewer if the pathogen in question would allow such procedure. Alternatively all bodily fluids need to be collected in tissues/diapers and then been disposed with other waste.



Self-Check 4	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page:

1. the most appropriate, disposal option for infectious wastes in rural health care facilities is
 - A. Burial in a protected pit
 - B. Burn
 - C. Medium- or high-temperature incineration
 - D. ALL
2. Which of the following **is True** about incinerator
 - A. Keep incinerator clean
 - B. Remove ash from ash chamber
 - C. Do not store waste in incinerator
 - D. ALL

Note: Satisfactory rating - 09 points Unsatisfactory - below 09 points
 You can ask your teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____



Operation Sheet 3

Prepare tools and equipment for identification and measuring Muda.

1. Discuss and plan to prepare tools and equipment for Muda identification.
2. Prepare tools and equipment for Muda identification.



LAP Test	Practical Demonstration
-----------------	--------------------------------

Name: _____ Date: _____

Time started: _____ Time finished: _____

Instructions: Given necessary templates, workshop, tools and materials you are required to perform the following tasks.

Task 1: Identify and prepare tools and equipment for measuring and identification of Muda.



Instruction Sheet **LG12: Clean environmental surfaces**

This learning guide is developed to provide you the necessary information regarding the following **content coverage and topics** –

Clean environmental surfaces

- wearing PPE during cleaning procedures
- removal of dust, dirt and physical debris from work surfaces
- cleaning work surfaces before and after any activity
- decontamination of equipments
- drying work surfaces before and after use
- replacing surface covers
- maintaining and storing Cleaning equipments

This guide will also assist you to attain the learning outcome stated in the cover page. Specifically, upon completion of this Learning Guide, you **will be able to** –

- worn personal protective clothing and equipment during cleaning procedures
- remove all dust, dirt and physical debris from work surfaces
- clean all work surfaces with a neutral detergent and warm water solution before and after each session or when visibly soiled
- decontaminate equipment requiring special processing in accordance with quality management systems to ensure full compliance with cleaning, disinfection and sterilization protocols
- all work surfaces are dried before and after use
- replace surface cover where applicable
- maintain and store cleaning equipment

Learning Instructions:

1. Read the specific objectives of this Learning Guide.
2. Follow the instructions described in number **3 to 16**.
3. Read the information written in the “Information Sheets 1”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
4. Accomplish the “Self-check 1” **in page 8**.
5. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 1).
6. If you earned a satisfactory evaluation proceed to “Information Sheet 2”. However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 1.
7. Submit your accomplished Self-check. This will form part of your training portfolio.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 104 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



8. Read the information written in the “Information Sheet 2”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
9. Accomplish the “Self-check 2” in page 16.
10. Ask from your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 2).
11. Read the information written in the “Information Sheets 7”. Try to understand what are being discussed. Ask your trainer for assistance if you have hard time understanding them.
12. Accomplish the “Self-check 3” in page 19.
13. Ask your trainer the key to correction (key answers) or you can request your trainer to correct your work. (You are to get the key answer only after you finished answering the Self-check 3).
14. If you earned a satisfactory evaluation proceed to “Operation Sheet 1” in page 21; However, if your rating is unsatisfactory, see your trainer for further instructions or go back to Information sheet 3.
15. Read the “Operation Sheet 1 and try to understand the procedures discussed.
16. Do the “LAP test” in page 22 (if you are ready). Request your trainer to evaluate your performance and outputs. Your trainer will give you feedback and the evaluation will be either satisfactory or unsatisfactory. If unsatisfactory, your trainer shall advice you on additional work. But if satisfactory you can proceed to Learning Guide #2.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 105 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



Information Sheet-1	wearing PPE during cleaning procedures
----------------------------	--

- wearing PPE during cleaning procedures

7.1 Personal protective clothing and equipment during *cleaning procedures*

Wear **gloves** while cleaning instruments and equipment. (Thick household or utility gloves work well.) If torn or damaged, they should be discarded; otherwise they should be cleaned and left to dry at the end of the day for use the following day.

- Even when wearing heavy-duty utility gloves, care should be taken to prevent needle sticks or cuts when washing sharps.

Wear protective eyewear (**plastic visors, face shields, goggles or glasses**) and a plastic apron, if available, while cleaning instruments and equipment to minimize the risk of splashing contaminated fluids into the eyes and onto the body.

- In most cleaning situations, gloves and aprons are sufficient.
- If there is risk of splashing waste into the face or eyes, face shields should also be used.
- PPE can only work if it is intact and undamaged, is fit for the purpose and is worn correctly
- Make sure the PPE fits you securely.
- If you notice any damaged PPE, notify your supervisor immediately
- Be familiar with the type of clothing you are required to wear for the purpose; for example, should the apron be waterproof



Self-Check 1	Written Test
---------------------	---------------------

Directions: Answer all the questions listed below. Use the Answer sheet provided in the next page:

1. In most cleaning situations, gloves and aprons are sufficient. **True** | False
2. If there is risk of splashing waste into the face or eyes, face shields should also be used. **True** | False
3. PPE can only work if it is intact and undamaged, is fit for the purpose and is worn correctly **True** | False
4. In most cleaning situations make sure the PPE fits you securely. **True** | False
5. Be familiar with the type of clothing you are required to wear for the purpose; for example, should the apron be waterproof. **True** | False

Note: Satisfactory rating - 16 points Unsatisfactory - below 16 points
 You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____



7.2 Removing dust, dirt and physical debris

Dust, soil and microbes on surfaces can all transmit infection. Cleaning removes foreign material and reduces the numbers of infectious agents and thus makes transmission of infectious organisms less likely

Floors

- Vacuum carpets and clean hard floors daily. Sweeping with an ordinary broom releases dust and bacteria into the air and so is not suitable.
- Wash buckets and mops after use with detergent and water and store them dry

Walls and fittings

- Generally, walls, blinds and curtains need less attention than floors unless they are in areas where soiling often happens (particularly if soiling is with blood or other body fluids). They should all be cleaned regularly and when visibly soiled. Curtains need changing regularly.



Self-Check 2	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

- _____ is the removes foreign material and reduces the numbers of infectious agents and thus makes transmission of infectious organisms less likely
 - Disposing
 - Disinfecting
 - Sterilization
 - Cleaning**

Note: Satisfactory rating - 12 points Unsatisfactory – below 12 points
 You can ask your trainer for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____



Information Sheet 3

cleaning work surfaces before and after any activity

7.3 Cleaning

Cleaning is the process that physically removes soiling, including large numbers of microorganisms and the organic material on which they grow. This is usually carried out using neutral detergent and warm water. Or **Cleaning** is the removal of visible soil, organic, and inorganic contamination from a device or surface, using either the physical action of scrubbing with a surfactant or detergent and water or an energy based process with appropriate chemical agents.

Routine environmental cleaning is required to minimize the number of microorganisms in the environment. A risk assessment must be undertaken to determine the method of decontamination required. Items not in close contact with patients or their immediate surroundings, e.g., floors, walls, ceiling, sinks and drains, present a minimal risk to patients and do not usually require disinfection. In most cases cleaning with a neutral detergent and warm water will be sufficient.

Disinfection: describes a process that eliminates **many** or all pathogenic microorganisms, except **bacterial** spores, on inanimate objects

Sterilization is a process that destroys or eliminates all forms of microbial life, including bacterial spores

Routine cleaning is the standard, everyday procedure for cleaning of clinical areas, including mopping of floors, damp dusting of surfaces with detergent, etc.

Terminal cleaning is performed when a patient with a transmissible illness is discharged (usually for isolation rooms), e.g. MRSA and other drug-resistant bacteria, tuberculosis, Clostridium difficile. The terminal cleaning process requires:

- Removal and discarding of all unused consumables and personal protective equipment (PPE) from the room
- Removal and laundering of all linen
- Removal and safe disposal of all waste
- Washing of all surfaces with detergent (including walls to a height of 2 metres)
- Wiping of all surfaces with an appropriate disinfectant (including bed frame, mattress and pillows). The IPC practitioner should be asked to advise on an appropriate disinfectant (usually alcohol-based or chlorine-based disinfectants at an appropriate strength or dilution). Remember that chlorine can be corrosive (causing damage to metal surfaces).
- Allowing all surfaces to dry before admission of a new patient.

Cleaning Methods

Dry Methods

- Dust attractant mop
- Dry dusting is not recommended as it may disperse dust and micro-organisms
- Sweeping brushes must not be used in clinical areas as they disperse dust and micro-organisms

Wet Methods

- Scrubbing
- Mopping
- Damp dusting

Once an item is washed it also needs to be rinsed and usually dried. Thorough rinsing with **clean** water removes any soap residue that can interfere with sterilization or HLD.3 After rinsing, items should be dried, especially if they will be sterilized or high-level disinfected using chemical disinfectants. Water remaining on the items (e.g., surgical instruments) dilutes the solution and may cause the process to fail

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 110 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



7.3.1 Purposes of cleaning

Cleaning in healthcare serves the dual functions of providing surface cleanliness and infection prevention and control

Cleaning holds special importance for hospitals and other healthcare facilities. The healthcare industry represents a significant population for health studies on cleaners because of the intensive and frequent cleaning with a wide range of cleaning and disinfecting products.

While the aesthetic benefits of cleaning are necessary for attracting and retaining patients, cleaning and disinfection play an essential role in healthcare settings by preventing healthcare-associated infections (HAIs)

Cleaning is important because:

- It is an effective way to reduce the number of microorganisms, especially endospores that cause tetanus, on soiled instruments and equipment.
- Neither sterilization nor high-level disinfection is effective without prior

A thorough washing with soap and clean water also physically removes organic material such as blood and body fluids.² This is important because dried organic material can entrap microorganisms, including endospores, in a residue that protects them against sterilization or disinfection. Organic matter also can partially inactivate some high-level disinfectants, rendering them less effective (AORN 1992; Rutala et al 1998).

Use of soap is important for effective cleaning because water alone will not remove protein, oils and grease (Nyström 1981). The use of hand (bar) or powdered soap is discouraged because the fatty acids in bar soap react with the minerals in hard water leaving a residue or scum (insoluble calcium salt), which is difficult to remove. Using liquid soap, if available, is preferable because it mixes more easily with water than bar or powdered soaps. In addition, liquid soap breaks up and dissolves or suspends grease, oil and other foreign matter in solution so that they can be removed more easily by the cleaning process.

7.3.2 Maintaining a clean work place

The importance of a clean workplace

The workplace environment influences employees' productivity, performance and well-being. No matter the industry, maintaining a clean workplace may help keep staff members safe, healthy and efficient. However, busy production schedules and increasing workloads may cause standards to dip.

Essential to safety

- When employees work in a messy environment, they may not notice all hazards, which increases the risk of an accident. According to the Occupational Safety and Health Administration (OSHA), an occupational hazard is anything in the workplace that may cause harm.

An occupational hazard is commonly caused by neglect on the part of the employer or a lack of awareness by workers. When the office or worksite isn't clean, it may increase the chance that a hazard will go unnoticed by a supervisor and staff members.

Employers may want to remember to keep the workplace free of debris and remind workers to put all equipment, such as personal protective equipment (PPE), in designated places to prevent an accident.

Crucial to health

- Flu season is rapidly approaching and workplaces may see an increase in the number of employees using sick days if they become ill. According to Kimberly-Clark Professional, germs can spread quickly through the workplace if supervisors and employees don't adequately sanitize their hands and their workspaces. Commonly used spaces, such as break rooms, can be hot spots for germs to accumulate.

Cleanliness is crucial in healthcare settings

- Controlling the spread of infections or viruses is vitally important in many different locations and settings – such as schools, leisure centres and the workplace – but it is even more crucial for healthcare

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 111 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



providers. People visiting or receiving treatment in these environments are already vulnerable to the spread of infection, so making sure that effective cleaning regimes are in place for waiting rooms, corridors, reception areas and wards is key. A clean and welcoming environment is also important from an aesthetic point of view, engendering feelings of well-being and trust in people who may be anxious or unwell.

The steps for cleaning are:

- Wear gloves while cleaning instruments and equipment. (Thick household or utility gloves work well.)
- Wear protective eyewear (plastic visors, face shields, goggles, or glasses), protective shoes, and a plastic apron, if available, while cleaning instruments and equipment to minimize the risk of splashing contaminated fluids into the eyes and onto the body.
- Using a soft brush or old toothbrush, detergent, and water, scrub instruments and other items vigorously to completely remove all blood, other bodily fluids, tissue, and other foreign matter. Hold items under the surface of the water while scrubbing and cleaning to avoid splashing. Disassemble instruments and other items with multiple parts, and be sure to brush in the grooves, teeth, and joints of items where organic materials can collect and stick.
- Rinse items thoroughly with clean water to remove all detergent. Any detergent left on the items can reduce the effectiveness of further chemical processing.
- Allow items to air dry (or dry them with a clean towel).

The guidance goes on to say that the arrangements for cleaning should include:

- Clear definition of specific roles and responsibilities for cleaning
- Clear, agreed and available cleaning routines
- Sufficient resources dedicated to keeping the environment clean and fit for purpose
- Consultation with Infection Control Teams (ICTs) or equivalent local expertise on cleaning protocols when internal or external contracts are being prepared
- Details of how staff can request additional cleaning, both urgently and routinely

6 Reasons Why a Clean Workplace Means a Safe Workplace

A clean workplace means more than just having a sparkling, fresh building. A clean workplace also ensures the safety and health of employees and visitors. In 2012 alone, nearly 3 million nonfatal workplace injuries and illnesses were reported by private industry employers. Workplace injuries can be prevented by taking action to ensure a clean, safe work environment.

Here are six reasons why a clean workplace also means a safe workplace:

- 1. Clean, dry floors to prevent slips and falls.**
 - Maintaining clean, dry floors is essential for the prevention of slips and falls in the workplace.
- 2. Disinfectants prevent the spread of germs and illness, including the flu.**
 - Germs can easily spread throughout a workplace, particularly during flu season—but disinfecting surfaces and objects stop germs in their tracks.
- 3. Proper air filtration lowers employee exposure to hazardous substances.**
 - Dusts and vapors are hazardous substances that can create an unsafe environment for employees. Building ventilation is one important factor in reducing airborne transmission of respiratory infections and maintaining the health and productivity of workers. Regularly cleaning prevents them from becoming saturated, which could lead to potential microbial growth and odor concerns.
- 4. Clean light fixtures improve lighting efficiency.**
 - Dirty light fixtures can reduce essential light levels, making it difficult and unsafe for employees to complete their daily tasks. Clean light fixtures significantly improve lighting efficiency in the workplace.
- 5. Green cleaning products are safer for both your employees and the environment.**
 - Not all cleaning products are the same, and some are held to a higher standard than others. Products with third-party certifications, such as Green Seal or GREENGUARD, must meet



specific standards and guidelines, which ensures they're safer for both people and the environment.

6. Proper disposal of waste and recyclable materials keeps work areas clutter-free.

- Allowing trash to pile up not only produces clutter, but it also presents a breeding ground for pests that pose a threat to your work environment. Placing “no-touch” wastebaskets in key locations throughout your facility ensures materials are disposed of and reduces the spread of germs. Recycling materials using clearly labeled waste receptacles also makes for a more sustainable environment.

7.3.3 General Principles of Cleaning

1. Ensure any electrical equipment to be cleaned is unplugged before commencing the decontamination cleaning. (Ensure cold chain for vaccine storage is maintained if fridge is unplugged).
2. Refer to the health and safety data sheets before using any product.
3. Display warning signs in the area, ensuring all signs are visible.
4. Wear suitable personal protective equipment. For most tasks gloves and aprons are sufficient. If there is the potential of a splash/irritation to the conjunctiva or mucous membranes face protection such as a mask and visor or goggles must be worn.
5. Prepare a fresh solution of cleaning agent or disinfectant as required.
6. Do not over-wet the floor.
7. Clean using lightly moistened clean disposable cloths (color coded as per National Hospital Office, below). The number of cloths required for cleaning per room depends on the level of activity being carried out e.g. one for patient examination area, one for desk area, one for washbasin to prevent transfer of microorganism from one area to another.
8. Particular attention should be given to frequently touched surfaces and those most likely to be contaminated with blood or body fluids e.g., toilet seats/flush handles, sinks/taps and doorknobs.
9. Sodium hypochlorite may damage metal surfaces. Remove disinfectant solution with clean water and dry with a disposable paper towel.
10. Dry equipment with paper towel.
11. Change gloves and perform hand hygiene between different tasks.
12. Dispose of solution in the toilet or designated household sink. Do not discard into wash hand basins
13. Dispose of cleaning cloth.
14. All chemicals must be used in accordance with manufacturer's instructions to ensure correct dilution and usage. These should be stored in a locked cupboard/room safely away from public access.
15. All equipment should be left clean, dry and tidy in storage area after use.
16. Do not use abrasive cleaners (e.g., Vim® or Comet®) or steel wool because these products can scratch or pit metal or stainless steel. These scratches then become a nesting place for microorganisms, making cleaning more difficult, as well as increasing the chance of corrosion (rusting).

7.4 Cleaning agents

All work surfaces must be cleaned correctly both before and after each session and when visibly soiled. There are different procedures depending on the level of risk; that is, the probability and the consequences of contamination. The following outlines some of the cleaning products that are commonly used in cleaning different surfaces in different areas.

Detergents

- When dealing with non-critical areas, detergent and warm water is all that is needed to clean the work area.

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 113 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



- Typically, detergents with a neutral pH are used. This means that they are neither alkaline nor acidic. Both alkaline and acidic substances can damage the skin and eyes and will also reduce the life of equipment and furnishings.
- Detergents with a neutral pH are far gentler and they clean most work areas effectively.
- Always complete any documentation necessary to indicate that cleaning has been carried out

Sanitisers and disinfectants

- Stronger cleaning products can be used when appropriate. These vary in strength:
 - **Low-level** disinfectants are effective in killing some bacteria and viruses; these can be used in most domestic settings.
 - **Medium-level** disinfectants are effective at killing a range of bacteria and viruses; these should be used if clients have tuberculosis or where there is a high likelihood that tuberculoid-causing pathogens are present.

High-level disinfectants are effective in killing all pathogens except for bacterial spores; these disinfectants should be used when cleaning medical equipment

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 114 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



Self-Check 3	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

6. _____ is the removal of visible soil, organic, and inorganic contamination from a device or surface.
- A. Disposing
 - B. Disinfecting
 - C. Sterilization
 - D. Cleaning**
7. **An occupational hazard** is commonly caused by
- A. neglect on the part of the employer
 - B. lack of awareness by workers
 - C. When the office or worksite isn't clean
 - D. ALL**
 - E.

Note: Satisfactory rating - 09 points Unsatisfactory - below 09 points
 You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____



a. Instruments processing

One area of infection prevention and control (IPC) in health facilities is the proper processing of instruments and other items that have come in contact with patients’ bodily fluids. It is critical in reducing the transmission of infections during clinical procedures and patient care.

Every item, whether it is soiled metal instruments or a pair of surgical gloves, requires special handling and processing in order to minimize the risk of accidental injury or blood or body fluid exposure to cleaning and housekeeping staff; provide a high-quality end product (i.e., sterile or high-level disinfected instruments and other items).

The basic IPC practices recommended to reduce disease transmission from soiled instruments and other reusable items are decontamination, cleaning, and either sterilization or HLD.

After completing an operation or invasive medical procedure, and while still wearing gloves, the health worker should

- Dispose of contaminated objects (gauze or cotton and other waste items) in a plastic bag or leak-proof, covered container.
- Next, disposable sharps (e.g., scalpel blades and suture needles) should be placed in a sharps container.
- Finally, all instruments and reusable items such as surgical gloves and suction cannulas, whether or not they were used in the operation, should be decontaminated by soaking for 10 minutes in a disinfectant (e.g., 0.5 percent chlorine solution).

Following decontamination

- The instruments and reusable items should be thoroughly cleaned with soap and water, completely rinsed, and dried.
- The surgical instruments and those items that come in contact with the bloodstream or that normally touch sterile tissue beneath the skin (critical items) should be sterilized to destroy all microorganisms including bacterial endospores.
 - When sterilization is not feasible or equipment is not available, however, HLD by boiling, steaming, or soaking in a chemical disinfectant is the only acceptable alternative.
- Instruments and other items that touch only mucous membranes or broken skin (semi-critical items), however, only need to be high-level disinfected

Important Definitions

Decontamination: This is a process that makes inanimate objects safer to be handled by staff before cleaning (i.e., inactivates HBV, HCV, and HIV, and reduces the number of other microorganisms but does not eliminate them).

Cleaning: A process that physically removes all visible dust, soil, blood, or other body fluids from inanimate objects as well as removing sufficient numbers of microorganisms to reduce risks for those who touch the skin or handle the object. It consists of thoroughly washing with soap or detergent and water, rinsing with clean water, and drying.

HLD: A process that eliminates all microorganisms except some bacterial endospores from inanimate objects by boiling, steaming, or using chemical disinfectants.

Sterilization: A process that eliminates all microorganisms (bacteria, viruses, fungi, and parasites) including bacterial endospores from inanimate objects by high-pressure steam (autoclave), dry heat (oven), chemical sterilants, or radiation

Chemical Disinfectants

Disinfectants are chemicals that destroy or inactivate microorganisms on inanimate objects, such as instruments and surfaces. Disinfectants are not meant to be used on the skin or mucous membranes. Many disinfectants are used alone or in combination (e.g., hydrogen peroxide and peracetic acid) in the health care setting. In most

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 116 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



instances, a given product is designed for a specific purpose and is to be used in a certain manner. Therefore, users should read labels carefully to ensure the correct product is selected for the intended use and applied efficiently.

Decontamination


In instrument processing, decontamination is the first step in handling used instruments and gloves. Decontamination inactivates HBV, HCV, and HIV and reduces the number of microorganisms. It is one of the most highly effective IP measures that can minimize the risk of transmission of these viruses to health care workers, especially cleaning and housekeeping staff, when they handle soiled medical instruments, surgical gloves, or other items. The objective of decontamination is to protect individuals who handle surgical instruments and other items that have been in contact with blood or bodily fluids from serious diseases.

Before cleaning, all soiled surgical instruments, surgical gloves, and other items should be first decontaminated by placing them in a 0.5 percent chlorine solution for 10 minutes. Because of the potentially high load of microorganisms and/or other organic material (blood or other bodily fluids) on soiled items, using a 0.5 percent solution for decontamination

Decontaminating Used Instruments and Other Items

1. Place all instruments in a 0.5 percent chlorine solution for 10 minutes immediately after completing the procedure
2. Decontaminate any surfaces contaminated during the procedure by wiping them with a cloth soaked in a 0.5 percent chlorine solution.
3. Immerse gloved hands in a 0.5 percent chlorine solution.
4. Remove gloves by turning inside out. If disposing of gloves, place them in a leak-proof container or heavy-duty plastic container.
5. If reusing gloves, soak in a 0.5 percent chlorine solution for 10 minutes for decontamination.
6. Remove instruments from the 0.5 percent chlorine solution after 10 minutes and immediately rinse them with cool water to remove residual chlorine before being thoroughly cleaned.
7. Two buckets can be used in the procedure areas or operating rooms, one filled with 0.5 percent chlorine solution and one with water, so instruments can be placed in the water after 10 minutes to help prevent corrosion.
 - a. Use a plastic container for decontamination to help prevent:
 - i. Dulling of sharps (e.g., scissors) due to contact with metal containers
 - ii. Rusting of instruments

Key steps in processing instruments and other items



Cleaning

After decontamination, cleaning is the next step in instrument processing. Cleaning is a process of physically removing infectious agents and other organic matter on which they live and thrive. This process does not necessarily destroy infectious agents. This is important because dried organic material can entrap microorganisms, including endospores, in a residue that protects them against sterilization or disinfection. Cleaning is an essential prerequisite to ensure effective disinfection or sterilization. Neither sterilization nor HLD could be effective without prior cleaning

Recall the steps of cleaning from chapter six

Sterilization and Storage

Sterilization is the destruction of all microorganisms, including bacterial endospores. Sterilization in health facilities can be achieved by high-pressure steam (autoclaves), dry heat (oven), chemical sterilants (glutaraldehyde or formaldehyde solutions), or physical agents (radiation).

To be effective, sterilization requires time, contact, temperature, and, with steam sterilization, high pressure. Because sterilization is a process, not a single event, all components must be carried out correctly for sterilization to occur. The effectiveness of any method of sterilization is also dependent on the following four factors:

1. The type of microorganism present.
2. The number of microorganisms present.
3. The amount and type of organic material that protects the microorganisms. Blood or tissue remaining on poorly cleaned instruments acts as a shield to microorganisms during the sterilization process.
4. The number of cracks and crevices on an instrument that might harbor microorganisms.

Methods of Heat Sterilization

The most common and readily available sterilization methods are:

1. High-pressure steam sterilization (autoclaves)



2. Dry heat sterilization (oven)
3. Chemical sterilization

High-Pressure Steam Sterilization (Autoclaves)

High-pressure steam sterilization (autoclave) is an effective method of sterilization but is the most difficult to do correctly.

The temperature, pressure, and time combinations for steam sterilization are as follows:

- At a temperature of 121°C (250°F), pressure of 106 kPa (15 lb/in²) for 20 minutes for unwrapped items and 30 minutes for wrapped items.
- At a higher temperature of 132°C (270°F), pressure of 30 lb/in² for 15 minutes for wrapped items

Note: Pressure settings (kPa or lb/in²) may vary slightly depending on the sterilizer used. When possible, follow the manufacturer's recommendations.

The two reasons why steam sterilization is an effective sterilization are as follows:

- Saturated steam is an extremely effective carrier of thermal energy that makes it many times more effective in conveying the necessary energy to the items to be sterilized than dry air.
- Steam is an effective sterilant because any resistant, protective outer layer of the microorganisms can be softened by steam, allowing coagulation of the inner sensitive portion of the microorganisms.

Advantages:

- Most commonly used effective method of sterilization.
- Sterilization cycle time is shorter than with dry heat or chemical sterilants.

Limitations:

- Requires a continuous source of heat (wood fuel, kerosene, or electricity).
- Requires equipment (steam sterilizer) that must be expertly maintained to keep it in working condition.
- Requires strict adherence to time, temperature, and pressure settings.
- Difficult to produce dry packs because breaks in procedure are common (e.g., not allowing items to dry before removing, especially in hot, humid climates).
- Repeated sterilization cycles can cause pitting and dulling of cutting edges of instruments (i.e., scissors).
- Plastic items cannot withstand high temperatures.

Preparing Items for Steam Sterilization

All instruments and other items should be decontaminated and thoroughly cleaned and dried before being sterilized. All jointed instruments should be open (or in the unlocked position) and disassembled. Reusable cloth items should be laundered and dried prior to sterilization.

Packing and Wrapping

Wrapping items to be sterilized permits sterile items to be handled and stored without being contaminated.

Materials used for wrappers should:

- Allow air removal and steam penetration
- Act as a barrier to microorganisms and fluids
- Resist tears and punctures and be free of holes
- Be inexpensive

Dry Heat Sterilization (Oven)

Dry heat sterilization is caused by hot air that destroys microorganisms through oxidation that causes slow destruction of the microorganisms' protein. Initially heat is absorbed by the exterior surface of an item and then passed to the next layer. Eventually, the entire object reaches the temperature needed for sterilization. Dry heat sterilization can be achieved with a simple oven as long as a thermometer is used to verify the temperature inside the oven.

Advantages:

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 119 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



- Effective method, as dry heat by conduction reaches all surfaces of instruments, even of instruments that cannot be disassembled
- Protective of sharps or instruments with a cutting edge (fewer problems with dulling of cutting edges)
- Leaves no chemical residue
- Eliminates “wet pack” problems in humid climates

Limitations:

Plastic and rubber items cannot be dry heat sterilized because temperatures used (160–170°C) are too high for these materials.

- Dry heat penetrates materials slowly and unevenly.
- It requires oven and continuous source of electricity.

Instructions for Dry Heat Oven

To ensure correct operation, consult specific operating instructions supplied by the oven’s manufacturer.

- **Step 1:** Decontaminate, clean, and dry all instruments and other items to be sterilized.
- **Step 2:** If desired, wrap instruments in aluminum foil or place in a metal container with a tight-fitting, closed lid. Wrapping helps prevent recontamination prior to use. Hypodermic or suture needles should be placed in glass tubes with cotton stoppers.
 - **Note:** When using dry heat to sterilize instruments wrapped in cloth, be sure that the temperature does not exceed 170°C/340°F.
- **Step 3:** Place loose (unwrapped) instruments in metal containers or on trays in the oven and heat to desired temperature.
- **Step 4:** After the desired temperature is reached, begin timing
 - **Note:** Use dry heat only for items that can withstand a temperature of 170°C (340°F). Needles and other instruments with cutting edges should be sterilized at lower temperatures (160°C [320°F]), because higher temperatures can destroy the sharpness of cutting edges.
 - Depending on the temperature selected, the total cycle time (preheating, sterilization time, and cool down) will range from about 2.5 hours at 170°C to more than 8 hours at 121°C.
- **Step 5:** After cooling, remove packs and/or metal containers and store. Loose items should be removed with sterile forceps/pickups and used immediately or placed in a sterile container with a tight-fitting lid.

Chemical Sterilization

Some high-level disinfectants will kill endospores after prolonged (10- to 24-hour) exposure. If objects need to be sterilized, but using high-pressure steam or dry heat sterilization would damage them or equipment is not available (or operational), they can be chemically sterilized. Common disinfectants that can be used for chemical sterilization include glutaraldehydes and formaldehyde. Sterilization takes place by soaking for at least 10 hours in 2 to 4 percent glutaraldehyde solution or at least 24 hours in 8 percent formaldehyde. Glutaraldehydes, such as Cidex, are often in short supply and very expensive, but they are the only practical sterilants for some instruments, such as laparoscopes, that cannot be heated. Both glutaraldehydes and formaldehyde require special handling and leave a residue on treated instruments; therefore, rinsing with sterile water is essential if the item must be kept sterile.

Advantages:

- Glutaraldehydes and formaldehyde solutions are not readily inactivated by organic materials.
- Both can be used for items that will not tolerate heat sterilization such as laparoscopes.
- Formaldehyde solutions can be used for up to 14 days (replace sooner if cloudy); some glutaraldehydes can be used for up to 28 days.

Limitations:

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 120 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



- Glutaraldehydes and formaldehyde are chemicals that cause skin irritation; therefore, all equipment soaked in either solution must be thoroughly rinsed with sterile water after soaking.

Storage of Sterile Instruments and Items

All sterile items should be stored appropriately to protect them from dust, dirt, moisture, animals, and insects. The storage area should be located next to or connected to where sterilization occurs and in a separate enclosed area with limited access that is used only to store sterile and clean patient care supplies.

Instructions for Storing Sterile Items

1. Keep the storage area clean, dry, dust-free, and lint-free.
2. Control temperature and humidity (approximate temperature 24°C and relative humidity less than 70 percent) when possible.
3. Packs and containers with sterile (or high-level disinfected) items should be stored 20 to 25 cm off the floor, 45 to 50 cm from the ceiling, and 15 to 20 cm from an outside wall.
4. Do not use cardboard boxes for storage. Cardboard boxes shed dust and debris and may harbor insects.
5. Date and rotate the supplies (first in/first out). This process serves as a reminder, but does not guarantee sterility of the packs.
6. Distribute sterile and high-level disinfected items from this area.

High-Level Disinfection and Storage

When sterilization equipment is either not available or not suitable, HLD is the only acceptable alternative method for the final processing of instruments. The HLD process destroys all microorganisms (including vegetative bacteria, tuberculosis, yeasts, and viruses) except some bacterial endospores.

HLD can be achieved by:

- Boiling in water
- Steaming (moist heat)
- Soaking instruments in chemical disinfectants (chemical disinfection)

Boiling

Boiling in water is an effective, practical way to high-level disinfect instruments and other items. Although boiling instruments in water for 20 minutes will kill all vegetative forms of bacteria, viruses (including HBV, HCV, and HIV), yeasts, and fungi, boiling will not reliably kill all endospores.

Steps for High-Level Disinfection by Boiling

- **Step 1:** Decontaminate and clean all instruments and other items to be high-level disinfected
- **Step 2:** If possible, completely immerse items in water. Adjust the water level so that there is at least 2.5 cm (1 inch) of water above the instruments. In addition, make sure all bowls and containers to be boiled are full of water. For example, empty bowls that turn bottom side up and float to the surface contain air pockets.
- **Step 3:** Close lid over pan and bring water to a gentle, rolling boil. (Boiling too vigorously wastes fuel, rapidly evaporates the water, and may damage delicate [or sharp] instruments or other items.) Hence, a gentle rolling boil is sufficient and will prevent instruments or other items from being bounced around and possibly damaged by striking other instruments or the side walls of the boiling pot.
- **Step 4:** Boil all items for 20 minutes. Start timing when the water begins to boil.
- **Step 5:** After boiling for 20 minutes, remove objects with previously high-level disinfected forceps. Never leave boiled instruments in water that has stopped boiling. As

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 121 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



the water cools and steam condenses, air and dust particles are drawn down into the container and may contaminate those instruments.

- **Step 6:** Use instruments and other items immediately, or with high-level disinfected forceps or gloves, place objects in a high-level disinfected container with a tight-fitting cover. Once the instruments are dry, if any pooled water remains in the bottom of the container, remove the dry items and place them in another high-level disinfected container that is dry and can be tightly covered

Steaming

The best method for HLD of gloves and a useful method of HLD of cannulas used during MVA is to steam them in a steamer containing one to three tiers of gloves or cannulas.

Steps for High-Level Disinfection by Steaming

After instruments and other items have been decontaminated and thoroughly cleaned, they are ready for HLD by steaming.

- **Step 1:** Place instruments, plastic MVA cannulas, and other items in one of the steamer pans with holes in its bottom. To make removal from the pan easier, do not overfill the pan.
- **Step 2:** Repeat this process until up to three steamer pans have been filled. Stack the filled steamer pans on top of a bottom pan containing water for boiling. A second empty pan without holes should be placed on the counter next to the heat source.
 - **Remember:** Be sure there is sufficient water in the bottom pan for the entire 20 minutes of steaming.
- **Step 3:** Place a lid on the top pan and bring the water to a full rolling boil. (When water only simmers, very little steam is formed and the temperature may not get high enough to kill microorganisms.)
- **Step 4:** When steam begins to come out between the pans and the lid, start the timer or note the time on a clock and record the time in the HLD log.
- **Step 5:** Steam items for 20 minutes.
- **Step 6:** Remove the top steamer pan and put the lid on the pan that was below it (the pan now on top). Gently shake excess water from the pan just removed.
- **Step 7:** Put the pan that was just removed onto the empty pan (see Step 3). Repeat until all pans are restacked on this empty pan and the top pan is covered with the lid. (This step allows the items to cool and dry without becoming contaminated.)
- **Step 8:** Allow items to air dry in the steamer pans (one to two hours) before using.
- **Step 9:** Using a high-level disinfected forceps, transfer the dry items to a dry, high-level disinfected container with a tight-fitting cover. Instruments and other items can also be stored in the stacked and covered steamer pans as long as a bottom pan (no holes) is used.

Soaking Instruments in Chemical Disinfectants

Although a number of disinfectants are commercially available in most countries, four disinfectants—chlorine, glutaraldehydes, formaldehyde, and peroxide—are routinely used as high-level disinfectants. A high-level disinfectant should be selected for use based on the characteristics of the items to be disinfected, the physical area (i.e., is it well ventilated?), and the skills of personnel available to do the procedure.

HLD by chlorine solutions

- Prepare 0.1 percent chlorine solution using boiled water that has been filtered if the tap water is cloudy.
- Soak for 20 minutes.

Formaldehyde

- Use 8 percent formaldehyde, which is inexpensive and readily available.
- Soak for 20 minutes.

Hydrogen peroxide

Medical laboratory L- III	HLT MLT3 TTLM 0919v1	Author/Copyright: Federal TVET Agency	Version -1 Sept. 2019	Page 122 of 139
---------------------------	----------------------	---------------------------------------	--------------------------	--------------------



- Use a 6 percent solution. The 3 percent hydrogen peroxide solutions used as antiseptics should not be used as a disinfectant.
- Soak for 20 minutes.

Glutaraldehydes

- Use a 2 to 4 percent solution.
- Soak for 20 minutes at 25°C.

Key Steps in Chemical High-Level Disinfection

- Decontaminate instruments and other items that may have been contaminated with blood and bodily fluids, and thoroughly clean and dry them before placing them in the disinfectant solution.
- Completely immerse all items in the high-level disinfectant.
- Soak for 20 minutes.
- Remove items using high-level disinfected or sterile forceps or gloves.
- Rinse well with boiled and filtered (if necessary) water three times and air dry
- Use promptly or store in a dry, high-level disinfected, covered container.



Self-Check 4	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page.

8. A process that makes inanimate objects safer to be handled is
A. Decontamination
 B. Chlorination
 C. Boiling
 D. Incineration
9. Which of the following method of instrumental processing is used to eliminates all microorganisms including bacterial endospores from inanimate objects
 A. HLD
B. Sterilization
 C. Contamination
 D. Decontamination
10. The effectiveness of any method of sterilization is also dependent on the following four factors:
 A. The type of microorganism present.
 B. The number of microorganisms present.
 C. The amount and type of organic material that protects the microorganisms.
 D. The number of cracks and crevices on an instrument
E. ALL
11. _____ is an effective method of sterilization but is the most difficult to do correctly
A. High-pressure steam sterilization
 B. Dry Heat Sterilization
 C. Chemical sterilization
 D. Incineration
12. The first step in processing of instrument is
 A. HLD
 B. Sterilization
 C. Contamination
D. Decontamination

Note: Satisfactory rating - 09 points Unsatisfactory - below 09 points

You can ask you teacher for the copy of the correct answers.

Answer Sheet

Score = _____
Rating: _____

Name: _____

Date: _____

Storage of Sterile Instruments and Items

All sterile items should be stored appropriately to protect them from dust, dirt, moisture, animals, and insects. The storage area should be located next to or connected to where sterilization occurs and in a separate enclosed area with limited access that is used only to store sterile and clean patient care supplies.

Instructions for Storing Sterile Items

1. Keep the storage area clean, dry, dust-free, and lint-free.
2. Control temperature and humidity (approximate temperature 24°C and relative humidity less than 70 percent) when possible.
3. Packs and containers with sterile (or high-level disinfected) items should be stored 20 to 25 cm off the floor, 45 to 50 cm from the ceiling, and 15 to 20 cm from an outside wall.
4. Do not use cardboard boxes for storage. Cardboard boxes shed dust and debris and may harbor insects.
5. Date and rotate the supplies (first in/first out). This process serves as a reminder, but does not guarantee sterility of the packs.
6. Distribute sterile and high-level disinfected items from this area.

SHELF LIFE

- The shelf life of an item (how long items can be considered sterile) after sterilization is event-related. An item remains sterile until something causes the package or container to become contaminate—time elapsed since sterilization is not the determining factor.
- To make sure items remain sterile until you need them, prevent events that can contaminate sterile packs, & protect them by placing them in plastic covers (thick polyethylene bags). An event can be a tear or worn area in the wrapping, the package becoming wet or anything else that will enable microorganism to enter the package or container.
- Before using any sterile item, look at the package to make sure the wrapper is intact and the seal unbroken and is clean and dry (as well as having not water stains).
- If the quality of wrapping cloth is poor and plastic bags are not available, limiting the shelf life is a reasonable option to ensure the sterility of the instruments



Self-Check 7	Written Test
---------------------	---------------------

Instructions: Answer all the questions listed below. Illustrations may be necessary to aid some explanations/answers. Write your answers in the sheet provided in the next page

1. Keep the storage area clean, dry, dust-free, and lint-free.
2. Control temperature and humidity (approximate temperature 24°C and relative humidity less than 70 percent) when possible.
3. Do not use cardboard boxes for storage.

**Operation Sheet 1****Decontaminating Used Instruments and Other Items**

1. Place all instruments in a 0.5 percent chlorine solution for 10 minutes immediately after completing the procedure
2. Decontaminate any surfaces contaminated during the procedure by wiping them with a cloth soaked in a 0.5 percent chlorine solution.
3. Immerse gloved hands in a 0.5 percent chlorine solution.
4. Remove gloves by turning inside out. If disposing of gloves, place them in a leak-proof container or heavy-duty plastic container.
5. If reusing gloves, soak in a 0.5 percent chlorine solution for 10 minutes for decontamination.
6. Remove instruments from the 0.5 percent chlorine solution after 10 minutes and immediately rinse them with cool water to remove residual chlorine before being thoroughly cleaned.
7. Two buckets can be used in the procedure areas or operating rooms, one filled with 0.5 percent chlorine solution and one with water, so instruments can be placed in the water after 10 minutes to help prevent corrosion.
8. Use a plastic container for decontamination to help prevent:
 - i. Dulling of sharps (e.g., scissors) due to contact with metal containers
 - ii. Rusting of instruments



LAP Test	Practical Demonstration
-----------------	--------------------------------

**Instr
uctio
ns:**

Given necessary templates, tools and materials you are required to perform the following tasks within 5min.

Task 1: Perform Decontamination



Operation Sheet 2	Instructions for Dry Heat Oven
--------------------------	---------------------------------------

Step 1: Decontaminate, clean, and dry all instruments and other items to be sterilized.

Step 2: If desired, wrap instruments in aluminum foil or place in a metal container with a tight-fitting, closed lid. Wrapping helps prevent recontamination prior to use. Hypodermic or suture needles should be placed in glass tubes with cotton stoppers.

- **Note:** When using dry heat to sterilize instruments wrapped in cloth, be sure that the temperature does not exceed 170°C/340°F.

Step 3: Place loose (unwrapped) instruments in metal containers or on trays in the oven and heat to desired temperature.

Step 4: After the desired temperature is reached, begin timing

- **Note:** Use dry heat only for items that can withstand a temperature of 170°C (340°F). Needles and other instruments with cutting edges should be sterilized at lower temperatures (160°C [320°F]), because higher temperatures can destroy the sharpness of cutting edges.
- Depending on the temperature selected, the total cycle time (preheating, sterilization time, and cool down) will range from about 2.5 hours at 170°C to more than 8 hours at 121°C.

Step 5: After cooling, remove packs and/or metal containers and store. Loose items should be removed with sterile forceps/pickups and used immediately or placed in a sterile container with a tight-fitting lid



LAP Test	Practical Demonstration
-----------------	--------------------------------

Name: _____ Date: _____

Time started: _____ Time finished: _____

Instructions: Given necessary templates, tools and materials you are required to perform the following tasks within 5min.

Task 1: Perform dry heat sterilization



Steps for Storing Sterile Items

1. Keep the storage area clean, dry, dust-free, and lint-free.
2. Control temperature and humidity (approximate temperature 24°C and relative humidity less than 70 percent) when possible.
3. Packs and containers with sterile (or high-level disinfected) items should be stored 20 to 25 cm off the floor, 45 to 50 cm from the ceiling, and 15 to 20 cm from an outside wall.
4. Do not use cardboard boxes for storage. Cardboard boxes shed dust and debris and may harbor insects.
5. Date and rotate the supplies (first in/first out). This process serves as a reminder, but does not guarantee sterility of the packs.
6. Distribute sterile and high-level disinfected items from this area.



LAP Test	Practical Demonstration
-----------------	--------------------------------

Name: _____ Date: _____

Time started: _____ Time finished: _____

Instructions: Given necessary templates, tools and materials you are required to perform the following tasks within 5min.

Task 1: Perform storing of sterile items

**❖ Steps for High-Level Disinfection by Boiling**

1. Decontaminate and clean all instruments and other items to be high-level disinfected
2. If possible, completely immerse items in water.
 - ✓ Adjust the water level so that there is at least 2.5 cm (1 inch) of water above the instruments. In addition, make sure all bowls and containers to be boiled are full of water. For example, empty bowls that turn bottom side up and float to the surface contain air pockets.
3. Close lid over pan and bring water to a gentle, rolling boil. (Boiling too vigorously wastes fuel, rapidly evaporates the water, and may damage delicate [or sharp] instruments or other items.) Hence, a gentle rolling boil is sufficient and will prevent instruments or other items from being bounced around and possibly damaged by striking other instruments or the side walls of the boiling pot.
4. Boil all items for 20 minutes. Start timing when the water begins to boil.
5. After boiling for 20 minutes, remove objects with previously high-level disinfected forceps. Never leave boiled instruments in water that has stopped boiling.
 - ✓ As the water cools and steam condenses, air and dust particles are drawn down into the container and may contaminate those instruments.
 - ✓ Use instruments and other items immediately, or with high-level disinfected forceps or gloves, place objects in a high-level disinfected container with a tight-fitting cover.
 - ✓ Once the instruments are dry, if any pooled water remains in the bottom of the container, remove the dry items and place them in another high-level disinfected container that is dry and can be tightly covered



LAP Test	Practical Demonstration
----------	-------------------------

Instructions: Given necessary templates, tools and materials you are required to perform the following tasks within 5min.

Task 1: Perform **High-Level Disinfection by Boiling**

**Steps for High-Level Disinfection by Steaming.**

1. Place instruments, plastic MVA cannulas, and other items in one of the steamer pans with holes in its bottom. To make removal from the pan easier, do not overfill the pan.
2. Repeat this process until up to three steamer pans have been filled. Stack the filled steamer pans on top of a bottom pan containing water for boiling. A second empty pan without holes should be placed on the counter next to the heat source.
 - ✓ **Remember:** Be sure there is sufficient water in the bottom pan for the entire 20 minutes of steaming.
3. Place a lid on the top pan and bring the water to a full rolling boil. (When water only simmers, very little steam is formed and the temperature may not get high enough to kill microorganisms.)
4. When steam begins to come out between the pans and the lid, start the timer or note the time on a clock and record the time in the HLD log.
5. Steam items for 20 minutes.
6. Remove the top steamer pan and put the lid on the pan that was below it (the pan now on top). Gently shake excess water from the pan just removed.
7. Put the pan that was just removed onto the empty pan (see Step 3). Repeat until all pans are restacked on this empty pan and the top pan is covered with the lid. (This step allows the items to cool and dry without becoming contaminated.)
8. Allow items to air dry in the steamer pans (one to two hours) before using.
9. Using a high-level disinfected forceps, transfer the dry items to a dry, high-level disinfected container with a tight-fitting cover. Instruments and other items can also be stored in the stacked and covered steamer pans as long as a bottom pan (no holes) is used.



LAP Test	Practical Demonstration
----------	-------------------------

Instructions: Given necessary templates, tools and materials you are required to perform the following tasks within 5min.

Task 1: Perform **High-Level Disinfection by Steaming**



Steps in Disinfection High-Level Chemical

1. Decontaminate instruments and other items that may have been contaminated with blood and bodily fluids, and thoroughly clean and dry them before placing them in the disinfectant solution.
2. Completely immerse all items in the high-level disinfectant.
3. Soak for 20 minutes.
4. Remove items using high-level disinfected or sterile forceps or gloves.
5. Rinse well with boiled and filtered (if necessary) water three times and air dry
6. Use promptly or store in a dry, high-level disinfected, covered container



LAP Test	Practical Demonstration
----------	-------------------------

Instru
ctions

: Given necessary templates, tools and materials you are required to perform the following tasks within 5min.

Task 1: Perform **Disinfecting high-level chemical**

List of Reference Materials

1. Federal Ministry of Health Ethiopia, April 2012, Infection Prevention and Patient Safety, Addis Ababa, Ethiopia: Federal Ministry of Health
2. Federal Ministry of Health, Ethiopia. 2004. *Infection Prevention Guidelines for Health Care Facilities in Ethiopia*. Addis Ababa, Ethiopia: Federal Ministry of Health.
3. Linda, Tietjen, Débora, Bossemeyer Noel McIntosh JHPIEGO, USIAD 2003 Guidelines for Healthcare Facilities with Limited Resources, , Johns Hopkins University,
4. WHO, , 2004, Practical Guidelines for Infection Control in Health Care Facilities World Health Organization Regional Office for Western Pacific, Manila Regional Office for South-East Asia, New Delhi
5. Helen Lemass , Niamh McDonnell , Dr. Nuala O'Connor , Dr. Sheila Rochford HCAI/AMR 2013, "INFECTION PREVENTION AND CONTROL FOR PRIMARY CARE IN IRELAND" ,
6. AG, Australian Guidelines for the Prevention and Control of Infection in Healthcare (2010)

Prepared By							
No	Name	Educational Back grund	LEVEL	Region	College	Email	Phaone Number
1	Kalicha Boru	Laboratory	B	oromia	Nagelle HSC	boru9683@gmail.com	0912493885
2	Furo Beshir	Laboratory	A	Harari	Harar HSC	nebi.furo@gmail.com	0911739970
3	Motuma Chali	Laboratory	B	oromia	Nekemte HSC	lammiifcaalii@gmail.com	0938456753
4	Abdirahman Mahad	Laboratory	A	Somali	Jigjiga HSC	abdirahman7584@gmail.com	0911044715
5	Adisu Tesfaye	Laboratory	B	Somali	Jigjiga HSC	adistesfaye21@gmail.com	0931747320
6	Kebebe Tadesse	Laboratory	B	BGRS	Pawi HSC	no	0926841290
7	Tagel Getachew	Laboratory	A	Harari	Harar HSC	tagegetachew@gmail.com	0915746748